Trafficking dynamics of VEGFR1, VEGFR2, and NRP1 in human endothelial cells

Sarvenaz Sarabipour[1], Karina Kinghorn[2], Kaitlyn M Quigley[2], Anita Kovacs-Kasa[3], Brian H Annex[3], Victoria L Bautch[2,4], Feilim Mac Gabhann[1]

[1]Institute for Computational Medicine and Department of Biomedical Engineering, Johns Hopkins University, Baltimore, Maryland, United States

[2]Department of Biology, University of North Carolina, Chapel Hill, North Carolina, United States

[3]Vascular Biology Center and Department of Medicine, Medical College of Georgia at Augusta University

[4]McAllister Heart Institute, University of North Carolina, Chapel Hill, North Carolina, United States

Abstract

The vascular endothelial growth factor (VEGF) family of cytokines are key drivers of blood vessel growth and remodeling. These ligands act via multiple VEGF receptors (VEGFR) and co-receptors such as Neuropilin (NRP) expressed on endothelial cells. These membrane-associated receptors are not solely expressed on the cell surface, they move between the surface and intracellular locations, where they can function differently. The location of the receptor alters its ability to 'see' (access and bind to) its ligands, which regulates receptor activation; location also alters receptor exposure to subcellularly localized phosphatases, which regulates its deactivation. Thus, receptors in different subcellular locations initiate different signaling, both in terms of quantity and quality. Similarly, the local levels of co-expression of other receptors alters competition for ligands. Subcellular localization is controlled by intracellular trafficking processes, which thus control VEGFR activity; therefore, to understand VEGFR activity, we must understand receptor trafficking. Here, for the first time, we simultaneously quantify the trafficking of VEGFR1, VEGFR2, and NRP1 on the same cells - specifically human umbilical vein endothelial cells (HUVECs). We build a computational model describing the expression, interaction, and trafficking of these receptors, and use it to simulate cell culture experiments. We use new quantitative experimental data to parameterize the model, which then provides mechanistic insight into the trafficking and localization of this receptor network. We show that VEGFR2 and NRP1 trafficking is not the same on HUVECs as on non-human ECs; and we show that VEGFR1 trafficking is not the same as VEGFR2 trafficking, but rather is faster and weighted to intracellular expression, due to faster internalization and lower recycling, resulting in increased overall degradation. As a consequence, the VEGF receptors are not evenly distributed between the cell surface and intracellular locations, with a very low percentage of VEGFR1 being on the cell surface. Our findings have implications both for the sensing of extracellular ligands and for the composition of signaling complexes at the cell surface versus inside the cell.

Introduction

Vascular endothelial cells that line blood vessels undergo sprouting angiogenesis, an essential process for tissue maintenance and remodeling. While successes have been achieved in the inhibition of angiogenesis in cancer and retinopathy, over a dozen human clinical trials of VEGF delivery in patients with peripheral artery disease (PAD) and coronary artery disease (CAD) who suffer from vascular insufficiencies have failed to produce clinical gain [1–3]. Thus, there is a need to increase blood flow via promotion of functional collateral vascularization. The inability to successfully bridge treatment from animals to humans demonstrates that our understanding of the complexities of how the VEGF receptor system controls angiogenesis is far from complete.

Vascular endothelial growth factor receptors (VEGFRs) are members of the receptor tyrosine kinase (RTK) superfamily and key targets for pro- and anti-angiogenic therapies [4]. The receptors bind to ligands to mediate signaling that controls endothelial cell behaviors during angiogenesis [4,7]; ligand binding induces a structural change in the receptor that increases phosphorylation [5]. The receptors can also interact in the absence of ligands [5,6]. VEGFR1 and VEGFR2 both bind common isoforms of VEGF-A; VEGFR1 also binds to isoforms of placental growth factor (PlGF) and VEGF-B [8]. In the body, different tissues (and tissues in different states) express different levels of VEGF/PlGF family isoforms, and the outcome of VEGFR signaling results from the competition of ligands for the receptors and the competition of receptors for the ligands. Select isoforms of VEGF and PlGF also interact with co-receptors expressed on endothelial cells, including heparan sulfate proteoglycans (HSPGs) and Neuropilin-1 (NRP1); these isoform-selective co-receptors play an important role in how the cell 'sees' and interprets the extracellular cytokine milieu. There are also direct (receptor-receptor) and indirect (receptor-ligand-receptor) associations between NRP1, a transmembrane glycoprotein, and the VEGFRs.

Of particular note among the VEGFRs, VEGFR2 has been more extensively studied, while VEGFR1, despite being a key regulator of vascular development [7,9], has been less well understood [4,8]. VEGFR1 is alternatively spliced into a membrane-integral form (mVEGFR1 or

mFlt1), which we focus on here, and a soluble form (sVEGFR1 or sFlt) [10]. sVEGFR1 is not cell-

associated and lacks a kinase domain, rendering it unable to initiate intracellular signaling; it binds

VEGFs and PlGFs and thus regulates the amounts of these ligands available to membrane-

associated VEGFRs. In some situations, membrane-associated VEGFR1 (mFlt1) appears to

behave as a localized decoy receptor, titrating VEGF access to VEGFR2 via non-signaling

complex formation; for example, mVEGFR1 is involved in connection-site selection as vessels

anastomose [13]. However, there is also evidence that VEGFR1 induces signaling pathways in

its own right, on endothelial cells [11] or other cell types [12]. For example, VEGFR1 can signal

through a non-canonical STAT3 pathway (specifically, phosphorylation of STAT3 independent of

JAK-STAT) to promote angiogenesis and perfusion recovery in mice and humans; this is typically

blocked by an alternately spliced VEGF isoform, VEGFA-165b, and suppression of VEGFA-165b

allows VEGF-A165a to bind VEGFR1, VEGFR1's cytoplasmic tyrosine Y1333 to be

phosphorylated, and STAT3 to be activated. [11]. In addition, recent experiments using

endothelium-specific VEGFR deletions in mice showed that VEGFR1 deletion impacts vascular

density even on a VEGFR2-deleted background, suggesting that VEGFR1 functions by signaling

in its own right, not only as a decoy for VEGFR2 [14]. The stability and trafficking of VEGFR1

alters its local availability (receptor density) and can therefore affect VEGFR1's decoy and

signaling functions.

While the trafficking pathways of VEGFR2 are well defined, much less is understood or

quantified about the trafficking of VEGFRs and NRP1. And yet, trafficking of these receptors is

responsible for multiple points of control in the VEGF pathway. Trafficking controls the quantity of

receptors available on the cell surface to bind to extracellular ligands in the first place; it controls

the stability and turnover of these receptors, and the balance between internal and external

expression. Trafficking also controls the movement of receptors post-ligation and activation [15].

As we have shown previously for VEGFR2, this trafficking exposes receptors to different

phosphatases, which selectively deactivate specific tyrosines and therefore specific downstream

pathways [16]. When receptors are internalized, they have two possible fates: degradation or

recycling back to the surface. VEGFR2 and NRP1 are recycled via a Rab4a-dependent route

(sometimes referred to as 'fast recycling') or a Rab11a-dependent route ('slow recycling') [2–4].

These two Rab-GTPases also control blood vessel formation and animal viability [17,18]. Both

unliganded and activated VEGFR2 are recycled [19,20]; however, how VEGFR1 internalization

is balanced with recycling and degradation has not previously been studied. NRP1 is also

trafficked and can impact VEGFR2 trafficking [16], but it is unclear how NRP1 affects VEGFR1

trafficking.

The VEGF-VEGFR system is targeted in many diseases, including ischemic diseases

such as peripheral artery disease, in an effort to modulate the vascular density and alter the extent

of perfusion of tissues. To design more effective therapies that target this system, a

comprehensive mechanistic understanding of how the VEGF-VEGFR system works is required.

Clearly, receptor trafficking plays an important role in this but has not yet been quantitatively

elucidated. For VEGFR2 and NRP1, trafficking has been studied *in vitro* [21–23] and *in silico*

[16,24,25], however previous computational models were based on measurements of receptors

in non-human cells, and/or in cells generated using overexpression of receptors or of trafficking

components. Thus, these previous models did not have detailed experimental measurements of

endogenous VEGFR1, VEGFR2, and NRP1 localization in human endothelial cells. Here, we

present such measurements and modeling for the first time; indeed, VEGFR1 trafficking is

investigated here for the first time in any cell type. To our knowledge, this is the first set of

VEGFR1, VEGFR2, and NRP1 trafficking experiments on the same human cell type, and the first

computational model to include detailed trafficking of VEGFR1, VEGFR2, and NRP1 explicitly.

Materials and Methods

Computational Model Construction

We developed a molecularly-detailed mechanistic computational model of membrane-integral

VEGF receptors, VEGFR1, VEGFR2, and Neuropilin-1, trafficking in human endothelial cells, and

parameterized it using experimental data from human endothelial cell culture. The model uses

coupled, nonlinear, deterministic ordinary differential equations to describe key biophysical and

biochemical reactions of receptor-receptor dimerization, and trafficking processes for receptors

(Figure 1). Binding and coupling reactions occur in every model compartment where associated

species are present. The model uses experimentally-derived surface receptor expression levels

and interaction kinetics for receptors expressed in HUVECs that have been previously measured

or derived (Supplementary Tables S1–S2) and incorporates geometry and initial concentrations

specific to the *in vitro* HUVEC cell culture experiments being simulated, including those previously

published and new experiments described here.

Receptor binding reactions and trafficking processes. In the model, we assume that VEGFR1

receptor monomers can undergo reversible homodimerization in the absence of ligands, as has

been shown for VEGFR2 [5,6]. The binding affinities and on- and off-rates of binding used here

are based on previously estimated reaction parameters (Supplementary Table S2). The model

includes five compartments: one extracellular (the fluid space outside the cell, i.e. culture media)

and four subcellular: cell surface; early endosomes (Rab4a-expressing and Rab5a-expressing

endosomes); recycling endosomes (Rab11a-expressing); and a degradation compartment

(representing lysosomes). Concentrations of components in each compartment are assumed to

be uniform (well mixed) but not constant. The model includes constant synthesis by the cell of

new VEGFR1, VEGFR2, and NRP1; the synthesis process inserts them into the cell surface

compartment. From there, movement of receptors between compartments occurs through first-

order transport processes; the trafficking processes are summarized in Figure 1. Surface

receptors are internalized to the early endosome compartment. From these endosomes,

receptors can either be degraded (via trafficking to lysosomes), recycled to the surface directly,

or trafficked to the Rab11a compartment for recycling; two recycling pathways are included

because previous evidence supports a role for both Rab4a and Rab11a endosomal recycling

pathways for VEGFR2 and NRP1 in endothelial cells [22]. The complete set of biochemical and

trafficking reactions can be found in the Supplementary tables S1-S4. The balance of synthesis,

degradation, and trafficking results in a steady-state, constant surface level of VEGFR1, VEGFR2,

and NRP1 populations in the absence of ligand. As part of the parameter optimization process,

this initial surface level of VEGFR1, VEGFR2, and NRP1 matches published experimental data

[26].

Model solution and outputs. The complete model contains 314 molecular species, including

separately tracking the levels of each receptor or receptor complex at each subcellular location

(plasma membrane, Rab4a/5a endosomes, Rab11a endosomes, lysosome). Code to simulate

the set of coupled ordinary differential equations that comprise this model were generated using

the rule-based BioNetGen software [27] and numerically solved using MATLAB. The model

outputs are the concentrations of each molecule or molecular complex over time. To facilitate

direct comparison to experimental data, the output concentrations of specific molecules or

complexes are combined into aggregated quantities of interest; for example, the total VEGFR1

levels on the cell surface is the sum of VEGFR1 present in multiple different complexes.

Depending on the experimental measurement, e.g. levels of proteins from whole cell lysates, or

only surface proteins isolated via biotin labeling, the appropriate molecular complexes from the

appropriate subcellular compartments in the model are aggregated to better compare the

simulated results to the experimental results.

Model parameter optimization. For HUVECs, we estimated the values of 15 receptor trafficking

parameters and 3 receptor production rate parameters in the absence of exogenous ligands.

These parameter estimates are based on optimization (fitting) to the results of new receptor

expression measurements and surface/internal expression percentages, described below,

including perturbations to receptor trafficking, that provide the most detailed and complete

quantitative data yet on VEGFR1, VEGFR2, and NRP1 trafficking from a single consistent set of

human endothelial cell experiments. Previously-published measurements of endogenous cell

surface VEGFR1, VEGFR2, and NRP1 (1,800, 4,900, and 68,000 receptors per cell respectively

for HUVECs) were also part of the optimization data [26]. When comparing simulation results to

experimental results, the simulation initial conditions, geometry, and parameter perturbations are

faithful to the experimental protocol for that experiment; and the experimental measurements

(such as surface and whole-cell receptor expression) are directly compared to the relevant

aggregated model outputs. As the experimental data are primarily western blots, we compare

normalized values of the experimental results to normalized values of the simulation results. For

optimization, we used a nonlinear least-squares optimization approach with the Levenberg-

Marquardt algorithm in MATLAB. As is common for biological systems with many components

and measurements, the experimental results available do not uniquely determine a single best fit

parameter set. Therefore, we initiated many separate optimization runs with different randomly

generated initial values, and bounded the parameters with a large range (five orders of magnitude:

$[10^{-6}, 10^{-1}]$ s^{-1}). Some parameters exhibit more uncertainty than others, but all parameters were

constrained to some degree (Figure 4).

Sensitivity Analysis. We performed univariate local sensitivity analysis to identify parameters

that most strongly affect model outputs. Key parameters were varied and the change in each

selected output calculated. We calculate the relative sensitivity metric as % change in output

divided by % change in parameter. Model outputs selected included levels of expression of the

receptor in the whole-cell and in subcellular locations (Figure 6).

Experimental Methods

Cell culture. Human umbilical vein endothelial cells (HUVEC) (see Supplementary Table S5 for

manufacturers, catalog numbers, and other details of all reagents used) were cultured at 37 °C in

EBM-2 medium supplemented with a bullet kit (EGM-2) and used at passages 3–5. For each

experiment, cells were used when they became confluent (2-3 days post-plating).

Perturbations. Several treatments were used to inhibit specific processes in the receptor

trafficking system (Figure 1). To inhibit new protein synthesis, following serum starvation HUVEC

were treated with 50 µg/mL cycloheximide (CHX). To inhibit lysosomal degradation, serum-

starved (overnight) HUVEC were treated with 10 $\mu g.mL^{-1}$ of chloroquine (CHQ). To deplete

expression of recycling-associated proteins Rab4a and Rab11a in HUVEC, approximately 1×10^{6}

cells were suspended in 100 ml nucleofector solution before adding 50–150 pmol of siRNA

(Silencer Select Locked Nucleic Acids) and using the A-034 program of the Amaxa Nucleofector

system for electroporation. Cells were transfected with control (non-targeting) siRNA duplex and

siRNA duplex targeting Rab4a and/or Rab11a. After 48 hr, the cells were serum-starved

(OptiMem + 0.1% FBS) for 18 hr, followed by CHX or CHQ addition.

Whole cell protein isolation. For selected experiments, total protein expression of VEGFR1,

VEGFR2, and NRP1 from whole cell lysates was measured. At appropriate time points, RIPA

buffer (50 mM Tris HCl, 150 mM NaCl, 1.0% (v/v) NP-40, 0.5% (w/v) Sodium Deoxycholate, 1.0

mM EDTA, 0.1% (w/v) SDS and 0.01% (w/v) sodium azide) and protease/phosphatase inhibitor

(Cell Signaling) at a pH of 7.4 was added to the plates, lysates were collected with cell scrapers,

subsequently boiled for 5 min at 95 °C, and used for immunoblot analysis as described below.

Experiments were repeated three times.

Internal and cell surface protein isolation. For other experiments, the protein expression of

receptors on the cell surface and expression inside the cell were separated and measured.

Biotinylation and isolation of cell surface proteins was performed according to the manufacturer's

protocol (Pierce™ Cell Surface Biotinylation and Protein Isolation Kit, cat#A44390). Briefly,

serum-starved HUVEC were washed twice in 6 ml of ice-cold PBS (Sigma-Aldrich) to stop

internalization. Surface VEGFR1, VEGFR2, and NRP1 were labeled with 0.25 mg.ml^{-1} of the

membrane-impermeant biotinylation reagent EZ-link Sulfo-NHS-SS-Biotin (ThermoFisher

Scientific) at 4°C for 30 min with constant rocking. The unreacted biotinylation reagent was

quenched by incubation with 50 mM glycine. HUVEC were washed twice with ice-cold PBS and

removed from the plate by gently scraping. After centrifugation at 800g for 5 min at 4°C, cells

were lysed using the provided lysis buffer supplemented with protease/phosphatase inhibitors

(Cell Signaling Technologies). A percentage of the total lysate was reserved for total protein

analysis, and the rest was put over a NeutrAvidin Agarose slurry-containing spin column

(ThermoFisher) at 4°C for 1 hr. Flow-through was collected in a given volume (same for all

samples), and the column was then eluted with 1 M dithiothreitol (DTT), and the eluate was

resuspended in the same volume as the flow-through, to facilitate determination of

surface:internal ratios. Experiments were repeated three times.

Immunoblots. Immunoblots were performed as previously described [18]. Briefly, whole cell

lysates, or cell surface and internal fractions, were collected as described above. For whole cell

lysates, approximately 10μg of protein was separated by SDS-PAGE on 10% Tris-Acetate gels

and transferred onto PVDF membranes. For biotin labeling experiments, equal volumes of the

flow-through (~10μg) and eluate were loaded to facilitate determination of surface:internal ratios.

Membranes were blocked for 1 hr in OneBlock (Prometheus) and incubated at 4°C with primary

antibodies (Supplementary Table S5) in OneBlock overnight. Membranes were washed 3X in

PBST (PBS 0.1% Tween-20) before adding HRP-conjugated secondary antibodies for 1 hr at RT.

Secondary antibody was removed and membranes washed four times in PBST before addition of

Luminata Forte (Millipore). See Supplementary Table S5 for details of antibodies, inhibitors and

siRNAs used for this study. BioRad software was used to image the western blots, and ImageJ

software was used to isolate and quantify the specific bands produced by the antibody. Loading

control (typically alpha tubulin) was used to normalize amount of total protein present.

Results

VEGF receptors exhibit different stability and turnover on HUVECs

Under normoxic culture conditions, we measured the whole-cell expression levels of VEGFR1,

VEGFR2, and NRP1 proteins in serum-starved primary HUVEC in the absence of exogenous

ligands. We compared untreated cells to cells treated with 50 ug.mL^{-1} cycloheximide (CHX) to

inhibit new protein synthesis and reveal the dynamics of trafficking and degradation not balanced

by new protein synthesis. Over time, NRP1 did not exhibit significant turnover, but rather was

stable throughout the experiment (Figure 2A). In contrast, whole-cell levels of VEGFR1 and

VEGFR2 decreased quickly, with half-lives of approximately 45 minutes for VEGFR1 and 75-90

minutes for VEGFR2; this was consistent with previous measurements of VEGFR2 loss following

CHX in other HUVEC studies [19]. Although VEGFR1 was suggested to be more stable than

VEGFR2 in the absence of ligand [18,29], here we carefully validated a different VEGFR1

antibody to ensure that it specifically identifies membrane-integral VEGFR1 (Supplementary

Figure S1).

VEGF receptors exhibit different localization patterns (cell surface vs intracellular)

Whole cell expression of receptors gives a partial view of receptor dynamics; to further refine

receptor localization we measured the cell-surface and intracellular distribution of VEGFR1,

VEGFR2, and NRP1 via biotin labeling of cell surface proteins on serum-starved HUVECs. Absent

additional perturbation, the percentage of each receptor expressed on the surface estimated from

these results (Figure 2B) is 10%, 51%, and 74% for VEGFR1, VEGFR2, and NRP1, respectively.

This is consistent with previous estimates of ~55-60% for VEGFR2 surface localization [30]; in

most endothelial cell types, NRP1 is present at cell surface densities higher than those of

VEGFR2 [26], and here we show that this difference is likely partly due to NRP1 being biased

towards surface expression. A previous estimate of ~25% VEGFR1 on the surface may have

been a result of overcounting using an antibody that is not specific for membrane-integral

VEGFR1 [29]. The absolute levels of VEGFR1 on the cell surface have been measured previously

[26] as being slightly lower or slightly higher than VEGFR2 levels, depending on cell type; however

the new observation that little of the total VEGFR1 is on the plasma membrane (with most of the

receptor being intracellular) has significant implications for how local receptor compositions differ.

Recycling pathway perturbations

Early endosomes sort internalized cargo for recycling or degradation, and proteins typically

recycle from endosomes to the cell surface in a Rab4a-dependent or Rab11a-dependent manner.

We disrupted these recycling pathways via siRNA-mediated knockdown of Rab4a and Rab11a in

HUVEC. Whole cell expression of VEGFR1, VEGFR2, and NRP1 was measured following siRNA-

Rab4a, siRNA-Rab11a, or siRNA-Rab4a/siRNA-Rab11a treatment; this was done in the absence

and presence of 50 ug.mL^{-1} CHX to assess changes to stability/turnover. The levels of VEGFR1,

VEGFR2, and NRP1 were essentially unaffected by Rab4a knockdown and Rab11a knockdown

experiments, and receptor stability in response to CHX was similarly unaffected (Figure 2C-D).

Previous estimates of receptor trafficking do not match HUVEC data

The goal of generating a computational model of VEGF receptor trafficking was to understand

and quantify the mechanistic basis for the observed experimental data. We previously estimated

trafficking parameters for VEGFR2 and NRP1 [16,25], however those parameters were based on

data from porcine aortic endothelial cells (PAECs) into which human receptors were transfected

and overexpressed [22]. We also previously used those PAEC-derived parameters to simulate

endothelial cells in human *in vivo* models [25]. To check whether those parameters are consistent

with the new HUVEC data, we used them in our trafficking model and simulated the HUVEC

experiments described above. The simulated surface vs internal levels of VEGFR2 and NRP1

appear to match well - perhaps these are similar on PAECs and HUVEC. However, using similar

trafficking parameters for VEGFR1 and VEGFR2 (this assumption was used in previous models

[25]) was inaccurate in HUVEC, since the simulated surface levels of VEGFR1 with that

assumption were higher than measured here. In addition, the stability of VEGFR1 and VEGFR2

in the simulations following CHX-induced protein synthesis shutdown was higher than measured.

These discrepancies suggest that the trafficking model parameters from the PAEC study are not

applicable to HUVEC (Figure 3A-B). This might be due to the species difference, tissue of origin

differences, or to differences in levels of receptor expression.

New estimates of VEGF receptor trafficking in HUVEC

Since the previous estimates for trafficking parameters did not fit the HUVEC data well, we

performed a new optimization of all model trafficking parameters for VEGFR1, VEGFR2, and

NRP1, using the experimental data from HUVEC (Figure 2, 5A). We performed the optimization

over a hundred times, and found that while there is not a single unique solution to the 18-

parameter optimization, most of the parameters are well constrained (Figure 4), particularly

receptor production, internalization, and degradation. Bearing in mind that the parameter

distributions shown are the aggregation of individual optimization parameter sets, we took the

geometric mean of each to create an estimated parameter set to test against the observed data.

Simulations using these parameters show good reproduction of the HUVEC data (Figure 5A-C).

For example, the receptor stability/turnover dynamics following CHX are reproduced for all three

receptors (Figure 5F-I); the percentage of each receptor that is present on the surface is matched

(Figure 5F-I); and the response to Rab4a and Rab11a knockdowns is similarly consistent (Figure

5F-I). Exploration of the parameter estimates highlights the mechanistic differences across the

receptors: VEGFR1 has the highest internalization rate constant, while VEGFR2 and NRP1 have

similar values; NRP has the slowest degradation rate constant, while VEGFR1 and VEGFR2 are

similar; NRP1 also has the highest recycling rates, with VEGFR1 and VEGFR2 being close to

each other (VEGFR1 being perhaps slightly faster via the Rab11a pathway) (Figure 4). In addition,

the production rates of the three receptors are comparable, suggesting that the observed receptor

levels are primarily controlled by trafficking and degradation.

Sensitivity analysis

We performed local sensitivity analysis to identify which of the trafficking parameters most

strongly affect key model outputs. Model outputs observed here are the same outputs that we

compare to experimental data - whole cell receptor expression, surface expression, intracellular

expression, and the percentage of each receptor on the cell surface. Trafficking parameters were

increased by 5%, and the percent change in each output was calculated (Figure 6). Increased

production of VEGFR1 or VEGFR2 increased receptor levels evenly across the cell; however,

production of NRP1 followed a more nonlinear path, with increased NRP1 production rates

causing large (hyperlinear) increases in NRP1 levels. NRP1 levels are also highly dependent on

VEGFR1 but not VEGFR2 production; this is likely due to formation of VEGFR1-NRP1 complexes

[31]; these complexes have a faster internalization and degradation rate and therefore deplete

NRP1 overall. All receptors are sensitive to changes in the degradation rate constant, although

NRP1 is less sensitive, presumably because it is already very low and increased degradation also

leads to a slight bias towards surface expression. Internalization has a stronger effect on surface

receptor levels (decreasing them) than it does on the internal levels, which are controlled primarily

by the balance of production and degradation. As a result of VEGFR1 being primarily intracellular

while VEGFR2 is more equally distributed between the surface and intracellular locations,

increased internalization has a much larger effect on total VEGFR2 cell levels than those of

VEGFR1. Finally, the predominance of Rab4-pathway recycling for VEGFR2 can be seen in the

sensitivity analysis results, along with evidence that the pathways are redundant and neither

receptor uses one exclusively.

Overall transport rates of VEGFR1, VEGFR2 and NRP1

We calculated the net rate of receptor movement in surface and endosomal compartments (Figure

7) to examine the contributions of each trafficking process to the steady-state distribution. At

steady-state, the model predicts a concentration (receptor density) of VEGFR1, VEGFR2, and

NRP1 complexes in the plasma membrane and in each of the endosomal compartments (Figure

1). The parameters calculated for the model are rate constants, and the overall rate of transport

for each process is the rate constant multiplied by the relevant concentration. A rate constant

could be high, but if the corresponding concentration is low, then the rate of movement will be

low; note that VEGFR1, VEGFR2 and NRP1 are not uniformly distributed across cellular

compartments - for HUVEC, cell surface levels of these receptors used in our model and based

on experimental data [26] are 1,800 VEGFR1, 4,900 VEGFR2, and 68,000 NRP1 per cell. Inside

the cell, the balance is different: 16,200 VEGFR1, 4,010 VEGFR2, and 19,180 NRP1 per cell

(Figure 7A). At steady state, the transport rates in and out of each compartment (subcellular

location) are balanced, so that 'net' rates of receptor movement are close to zero. For the Rab11-

recycling pathway endosomes, the receptors coming in from Rab4a are balanced by receptors

leaving via recycling to the surface (Figure 7C). For Rab4 early endosomes, receptors due to

internalization from the surface are balanced by the sum of receptors being degraded or recycled

via both pathways. For the cell surface, receptors out due to internalization are balanced by the

sum of receptors in via new synthesis and two recycling pathways. The internalization rate of

VEGFR1 was found to be 4.0×10^{-2} s^{-1}, for a half-life (residence time on the surface) of $\ln(2)/k_{int}$

= 27 seconds. For VEGFR2, half-life (residence time on the surface) is 4.5 minutes, and NRP1,

6 minutes. These are short compared to the overall half-life of the receptors (Figure 2A), because

once internalized, most of the receptors are recycled rather than degraded, emphasizing the

dynamic nature of receptor trafficking and cycling between the surface and internal receptor pools

(even for the highly stable NRP1). Any perturbations - for example, addition of CHX, CHQ, or

depletion via siRNA - in addition to its effect on the appropriate rate constant, would result in

changes to the local receptor concentrations, changes to the related overall rates, and (at least

transiently) changes to the net rates of movement (which is what enables the local receptor

concentration changes).

We can directly compare rate constants for processes that act on the same set of receptors (for

example, the processes that lead from a certain subcellular location). This is most obvious when

looking at the early endosomes, which function as the main crossroads in the model (Figure 1,

Figure 7B marked 'Rab4a'). First, comparing the two major fates of endosomal receptors

(recycling and degradation), while the combined recycling rate constants via the two pathways is

greater than degradation rate for each of the three receptors, the ratios are very different: 19:1,

11:1 and over 7000:1 for VEGFR1, VEGFR2, and NRP1 respectively (Figure 7B, left). This means

that for every VEGFR1 degraded, 19 are recycled; or viewed another way, there is an

internalization-recycling loop for which a VEGFR1 makes on average 19 in-out loops before

internalization; VEGFR2 receptors undergo fewer and NRP1 very many more internalization-

recycling loops, showing that despite the stability (lifetime) of receptors on the cell overall (Figure

2A), there is also frequent movement between the surface and intracellular locations; the

receptors are more likely to be recycled than degraded while in the intracellular locations. Within

recycling, the receptors are biased differently between the Rab4a- and Rab11a-dependent

pathways (Figure 7B, middle). VEGFR2 preferentially uses Rab4a, while NRP1 and VEGFR1 use

the Rab4a and Rab11a pathways more equally; none of the receptors uses one pathway

exclusively, which is in keeping with previously observed VEGFR2 and NRP1 observations

[16,22] and with the redundancy shown by the knockdown experiments here. Using a similar

analysis, we can see the mechanistic basis for the differently-skewed surface vs internal levels of

the receptors (Figure 7B, right): the rate constant for internalization is significantly larger than that

of recycling for VEGFR1, but the inverse for NRP1. VEGFR2 is much more balanced, reflecting

the almost-even distribution of VEGFR2 between the surface and intracellular locations.

Chloroquine (CHQ) and VEGFR1 degradation. We previously showed that loss of new protein

synthesis resulted in a decrease in VEGFR1 levels over time as the receptor is degraded. The

consequence of inhibited degradation, if production is unchanged, should be an increase in

receptor levels. We treated cells with Chloroquine (CHQ), an inhibitor of lysosomal degradation,

and indeed see VEGFR1 accumulation (Figure 8). Using computational simulations that recreate

the conditions of the CHQ experiments, we estimate that this observed increase in VEGFR1 levels

is consistent with CHQ treatment inhibiting approximately 42% of VEGFR1 degradation; if

VEGFR1 degradation were completely inhibited, the predicted accumulation of VEGFR1 would

be even higher. This suggests that at least half of VEGFR1 loss is through non-lysosomal

degradation pathways, i.e. is not inhibited by CHQ; alternatively, the accumulation of VEGFR1

may cause a downregulation of VEGFR1 synthesis to compensate. VEGFR2 and NRP1 levels

did not increase, again suggesting non-lysosomal degradation pathways or adaptation of

production to prevent receptor build up.

Discussion

The intracellular trafficking of receptors is central to the sensing of ligands and the specificity of

downstream signaling. Trafficking that differs between receptors, or between cells, or becomes

altered within the same cell type based on condition, results in different surface and total receptor

levels, affecting availability to ligands, and exposure to different regulators, such as

phosphatases, that can result in different signaling. Specifically, for the VEGF system, with

multiple receptors (VEGFR1, VEGFR2, and NRP1) competing and cooperating, the local

expression of each of the receptors influences the ligand binding and activation of the others; they

form a system. The intracellular location of VEGFR2, in particular, influences the downstream

signaling pathways and cellular behaviors that are set in motion, and this impacts tissue-level

properties such as vascular network architecture and function [16]. In this study, we quantified

VEGF receptor trafficking rates and distribution in HUVEC using a mechanistic computational

model fitted to detailed new experimental data.

While there are some previous estimates of VEGF receptor trafficking, this is the first

quantitative rate analysis of VEGFR trafficking in human endothelial cells. It is of particular

importance that VEGFR1, VEGFR2, and NRP1 were measured in the same cell type and at the

same time, because of the interdependence of these receptors. We compared these new results

to VEGFR2 and NRP1 parameters obtained for a related computational model [16] using

experimental data from porcine aortic endothelial cells (PAECs) [22] and found that values were

not consistent between the studies. There are multiple reasons why VEGFR trafficking may be

different in the PAEC study versus this HUVEC study, including different species of the cells and

different tissue origins of the cells. Moreover, human VEGFR2 and NRP1 were overexpressed in

the PAECs, and the PAEC study also used data from overexpression of tagged Rab proteins,

which can impact Rab biogenesis and alter endosomal trafficking [33]. The rate constants of

HUVEC trafficking of VEGFRs quantified here are in general higher (faster) than those identified

from PAECs, with the exception of NRP1 degradation (Figure 3A, 5A). The fast kinetics of

endogenous proteins indicate a highly active system, with frequent movement between

intracellular compartments, indicative of cells ready to respond to extracellular stimulation.

Given the differences between previous trafficking parameters and the parameters

described here, it may be important in the future to quantify trafficking rates by species and cell

type, including across human endothelial cells of different tissue origin [26], or cells under different

pathological conditions such as hypoxia. For example, endothelial cells in different tissues

express different amounts of VEGF receptors [26,31,32]. In addition, not only does receptor

trafficking determine the availability of receptors locally to interact with ligands and with each

other, but also the binding of ligands and activation of receptors can influence trafficking.

A key observation from this study is that trafficking of the three VEGF receptors studied is

quite different. NRP1 is highly stable, with very low turnover as evidenced by consistent

expression after cessation of new protein synthesis. Although it is stable on the cell, the

mechanistic rate constants suggest that NRP1 is, like the other receptors, actively and rapidly

exchanged between the surface and endosomes. VEGFR1 and VEGFR2 are much less stable,

and their levels decline rapidly when new protein synthesis is blocked. This propensity for high

production and high degradation may be relevant to sensitivity to microenvironmental ligand

levels, and to downregulation of or adaptation to those stimuli. The trafficking of VEGFR1 and

VEGFR2 are also different, perhaps most obviously in the resulting differences in receptor levels

on the cell surface and internally. Most VEGFR1 appears to be intracellular, while VEGFR2 is

balanced between the two locations. In terms of absolute numbers of receptors, this means that

VEGFR2 is in excess on the cell surface, but VEGFR1 is in excess intracellularly. These different

subcellular regions (further distinguished by a bias towards surface expression for NRP1) are

therefore very different domains for ligand binding and downstream signaling, and the implications

of these local levels will need further study.

VEGFR1 was a particular focus of our study due to it being understudied compared to

VEGFR2. Quantifying VEGFR1 trafficking in the same experiments as VEGFR2 and NRP1 is

important, as these receptors are commonly co-expressed on endothelial cells and can interact.

VEGFR1 and NRP1, in particular, can interact directly in the absence of ligands (unlike VEGFR2

and NRP1) [32], and this VEGFR1-NRP1 complex forms an important part of the computational

model. A key assumption made here was that this complex trafficked at rates similar to VEGFR1;

identifying an independent set of rate constants for this complex was not possible while measuring

the receptors independently. We also tested an alternate assumption that the rates for VEGFR1-

NRP1 would be similar to those for NRP1, however this did not provide a good fit to the

experimental data. This VEGFR1-NRP1 complex has important consequences, including

exhibiting ligand-binding specificity that impacts not just VEGFR1 binding but also ligand

availability to VEGFR2. The presence of the complex also partly explains the low degradation rate

constant of NRP1 in the model, in that the newly produced NRP1 in the simulations is not

balanced at steady state by NRP1 degradation but rather by the combined degradation of NRP1

and VEGFR1-NRP1 , and this is mostly through degradation of the VEGFR1-NRP1 complex.

When new protein synthesis is stopped, VEGFR1 decreases, VEGFR1-NRP1 levels decrease,

and overall NRP1 protein levels remain stable as NRP1 alone is degraded slowly. Notably, when

VEGFR1 or NRP1 production rates are increased (as in our sensitivity analysis, Figure 6), the

effects on NRP1 levels are very strong, as the ratio of NRP1:VEGFR1-NRP1 changes and thus

the aggregate degradation of NRP1 changes.

Combining comprehensive quantitative experimental data with mechanistic computational

models has enabled us to identify which subcellular pathways are most influential for trafficking

of each of the VEGF receptors, and to gain insight into how the VEGF receptor system is in quite

a different state (different relative levels of the receptors) in different parts of the cell. This local

regulation and the local conditions likely play an important role in response to ligand stimulation,

and can potentially be harnessed to enhance therapeutic targeting of the VEGF system.

Author contributions

Author contributions

SS and FMG designed and wrote the code, performed data visualization, and wrote the original

draft of the manuscript. KK, KMQ, and VLB planned and KK and KMQ performed the experiments

and experimental data analysis. FMG, VLB, and BHA acquired resources and funding. SS, FMG,

KMQ, KK, VLB, AKK, and BHA performed conceptualization, formal analysis, investigation,

methodology, validation, and wrote, reviewed and edited the manuscript.

Conflicts of interest

The authors declare no conflict of interest.

Acknowledgements

This work was supported by NIH grant R01-GM129074.

Supplementary File

Contains five Tables and one Figure.

516
517

518 **FIGURES**
519

520
521
522
Figure 1. Schematic of molecular interactions of the computational model. Cellular
biophysical and biochemical reactions between VEGFR1, VEGFR2 and NRP1 receptors. The
receptors can homodimerize (at a lower level than that induced by ligands), and unlike VEGFR2,
VEGFR1 can form a complex with NRP1 in the absence of ligands [32]. During trafficking, surface
protein complexes (monomeric, dimeric, or higher order) can be internalized (rate constants
denoted k_{int}). Early endosomal ("Rab4a/5a") receptors can be degraded (rate constant k_{deg}),
recycled (rate constant k_{rec4}), or transferred to the Rab11a compartment (rate constant k_{4to11})
which leads to an additional recycling pathway (rate constant k_{rec11}). New surface receptors
(monomers) are produced at rate k_{prod}. Each of the rate constant values can be different for the
different receptors. Reaction rates and species concentrations are detailed in Supplementary
Tables S1-S4.
534
535

536
537
538
Figure 2. Experimental measures of VEGFR1, VEGFR2, and NRP1: whole-cell and surface
levels in HUVEC. Western blot of total HUVEC lysates treated as indicated for 1-24 hours with
50 µg/ml cycloheximide (CHX) and stained for VEGFR1, VEGFR2 and NRP1. Representative of
n=3 replicates. **B,** Western blot of biotin labeling assay to measure surface and internal VEGFR1,
VEGFR2, and NRP1 levels in HUVEC. Representative of n=3 replicates. **C,** Western blot of
VEGFR receptor levels following Rab4a or Rab11a siRNA treatment shows similar receptor
stability as in response to CHX alone. Additional measurements at 60, 90, 120 and 240 min
showed similar results (data not shown). Representative of n=3 replicates. **D,** Western blot
showing the effect on VEGFR1 levels of depletion of both Rab4a and Rab11a.
548

A

Previously estimated parameters (2015-2017)				
Parameter	Description	VEGFR1 (s^{-1})	VEGFR2 (s^{-1})	NRP1 (s^{-1})
kint	Internalization	2.6×10^{-3}	2.6×10^{-3}	2.6×10^{-3}
kdeg	Degradation	8.6×10^{-6}	8.6×10^{-6}	3.8×10^{-4}
krec4	Early recycling	3.8×10^{-3}	3.8×10^{-3}	3.8×10^{-5}
k4to11	Pre-recycling	1.0×10^{-5}	1.0×10^{-5}	1.9×10^{-2}
krec11	Late recycling	1.4×10^{-4}	1.4×10^{-4}	1.4×10^{-2}
kprod	production rate	1.0×10^{-4}	2.8×10^{-4}	3.5×10^{-3}

B

C

Figure 3. Previous estimates of VEGFR trafficking parameters are not consistent with HUVEC observations. A, VEGFR2 and NRP1 trafficking parameters as previously estimated using a computational model and published experimental data from porcine aortic endothelial cells overexpressing human VEGFR2 and NRP1 [16]; and VEGFR1 parameters as then assumed

based on VEGFR2 parameters for computational models incorporating all three receptors [25].
B, Model-predicted levels ("Sim") of VEGFR1, VEGFR2 and NRP1 at steady state on the cell
surface and in intracellular endosomes (Rab4a/5a and Rab11a) using the 2015-2017 trafficking
parameters based mainly on PAEC data. Experimental data comparisons ("Exp") for surface
receptor levels are for HUVECs from this study. **C,** Simulations (lines) and experimental data
(dots) for whole-cell VEGFR1, VEGFR2, and NRP1 over time following CHX treatment. The
schematic shows how CHX inhibition of protein synthesis is represented in the model.
561

Figure 4. Trafficking parameter estimates for VEGFR1, VEGFR2, and NRP1 based on optimization to experimental data from HUVEC. While there is not one single unique optimization solution, the optimized parameter sets are well constrained, for most the majority of estimates are constrained within an order of magnitude. For each parameter, we used the median of the 76 optimizations to create a standard parameter set for use through the rest of the manuscript (see Supplementary Table S4 and Figure 5A).

A

Updated trafficking parameters (2022)			
Parameter	Description	VEGFR1 (s^{-1})	VEGFR2 (s^{-1})	NRP1 (s^{-1})
kint	Internalization	4.1×10^{-2}	3.6×10^{-3}	2.5×10^{-3}
kdeg	Degradation	2.5×10^{-4}	3.2×10^{-4}	5.5×10^{-6}
krec4	Early recycling	2.5×10^{-3}	2.6×10^{-3}	1.9×10^{-2}
k4to11	Pre-recycling	2.2×10^{-3}	7.6×10^{-4}	1.9×10^{-2}
krec11	Late recycling	5.0×10^{-2}	1.2×10^{-2}	2.1×10^{-2}
kprod	production rate	3.8×10^{-3}	1.5×10^{-3}	3.7×10^{-3}

Figure 5. Computational model predictions and model validation with trafficking parameters. A, Updated trafficking parameters for VEGFR1, VEGFR2, and NRP1 based on geometric means of optimized parameter distributions. **B,** Surface, Rab4a/5a and Rab11a distributions of unligated VEGFR1, VEGFR2 and NRP1 at steady state with the 2022 trafficking parameters. Simulations ("Sim") and experimental data ("Exp") of surface VEGFR1, VEGFR2 and NRP1 levels. **C,** Simulations (lines) and experimental data (dots) for whole-cell VEGFR1, VEGFR2, and NRP1 over time following CHX treatment. **D-F,** Each panel shows the response to CHX at time zero following no knockdown, Rab4a knockdown, Rab11a knockdown, and double knockdown, for total (whole cell) (D, F) and surface (E) VEGFR1, VEGFR2, and NRP1. The schematics show how the Rab knockdown experiments are represented in the model.

582
583

Figure 6. Sensitivity of model outputs to VEGFR1, VEGFR2 and NRP1 trafficking parameters. Local sensitivity analysis was performed by examining the sensitivity of model outputs to small changes in each of the 15 receptor trafficking and 3 receptor production parameters. Sensitivity values are the ratio of percent change in key model outputs (x-axis) to percent change in the parameter values (y-axis). The gray values represent higher than linear sensitivity that negatively (production of VEGFR1) or positively (production of NRP1) affect NRP1 levels due to the nonlinearity of VEGFR1-NRP1 interactions.

Figure 7. Transport rate analysis on differential transport of VEGFR1, VEGFR2 and NRP1.
A, Receptor expression levels on the cell surface and intracellularly in our model without
perturbation. The surface levels are as previously measured experimentally [26] and the internal
levels are calculated here based on the surface levels and the percentage of each receptor on
the surface based on the biotin labeling experiments (Figure 2). **B,** Mechanistic insights from
trafficking parameters, informed by key ratios of rate constants. **C,** The overall transport rates
(rate constant multiplied by concentration) for VEGFR1, VEGFR2 and NRP1 in each subcellular
location. A rate constant could be high, but if the corresponding concentration is low (and the
receptors are not uniformly distributed across cellular compartments), then the rate of movement
will be low. At steady state, these overall rates in and out are balanced, so the 'net' rates are close
to zero. For Rab11a: receptors arrive from Rab4a balances out by recycling, for Rab4a: receptors
are internalized to Rab4a balances out the sum of degraded, recycling, and to Rab4a, for surface:
receptor internalization balances the sum of new synthesis and recycling from Rab4a and
Rab11a.

617 **A**

624 **Figure 8. Impact of inhibiting lysosomal degradation on whole cell and surface VEGFR1,**
625 **VEGFR2, NRP1 trafficking.** Chloroquine (CHQ) treatment inhibits lysosomal degradation. **A,**
626 Western blot of total (whole cell) VEGFR1, VEGFR2, NRP1 levels after indicated times of CHQ
627 treatment. **B,** Comparison of simulation result (line) and quantification of Western blot (dot) for
628 whole cell VEGFR1 levels after indicated times of CHQ treatment.

References

1. Annex BH. Therapeutic angiogenesis for critical limb ischaemia. Nat Rev Cardiol. 2013;10: 387–396.

2. Annex BH, Cooke JP. New directions in therapeutic angiogenesis and arteriogenesis in peripheral arterial disease. Circ Res. 2021;128: 1944–1957.

3. Mac Gabhann F, Annex BH, Popel AS. Gene therapy from the perspective of systems biology. Curr Opin Mol Ther. 2010;12: 570.

4. Simons M, Gordon E, Claesson-Welsh L. Mechanisms and regulation of endothelial VEGF receptor signalling. Nat Rev Mol Cell Biol. 2016;17: 611–625. doi:https://doi.org/10.1038/nrm.2016.87

5. Sarabipour S, Ballmer-Hofer K, Hristova K. VEGFR-2 conformational switch in response to ligand binding. eLife. 2016;5: e13876. doi:http://doi.org/10.7554/eLife.13876

6. da Rocha-Azevedo B, Lee S, Dasgupta A, Vega AR, de Oliveira LR, Kim T, et al. Heterogeneity in VEGF Receptor-2 Mobility and Organization on the Endothelial Cell Surface Leads to Diverse Models of Activation by VEGF. Cell Rep. 2020;32: 108187.

7. Chappell JC, Bautch VL. Vascular development: genetic mechanisms and links to vascular disease. Curr Top Dev Biol. 2010;90: 43–72.

8. Markovic-Mueller S, Stuttfeld E, Asthana M, Weinert T, Bliven S, Goldie KN, et al. Structure of the Full-length VEGFR-1 Extracellular Domain in Complex with VEGF-A. Structure. 2017;25: 341–352. doi:https://doi.org/10.1016/j.str.2016.12.012

9. Bautch VL. VEGF-directed blood vessel patterning: from cells to organism. Cold Spring Harb Perspect Med. 2012;2: a006452.

10. Kendall R, Thomas K. Inhibition of vascular endothelial cell growth factor activity by an endogenously encoded soluble receptor. Proc Natl Acad Sci U S A. 1993;90: 10705–1070. doi:https://doi.org/10.1073/pnas.90.22.10705

11. Ganta VC, Choi M, Kutateladze A, Annex BH. VEGF165b Modulates Endothelial VEGFR1–STAT3 Signaling Pathway and Angiogenesis in Human and Experimental Peripheral Arterial Disease. Circ Res. 2016;120: 282–295. doi:https://doi.org/10.1161/CIRCRESAHA.116.309516

12. Ganta VC, Choi M, Farber CR, Annex BH. Antiangiogenic VEGF165b Regulates Macrophage Polarization via S100A8/S100A9 in Peripheral Artery Disease. Circulation. 2018;139: 226–242. doi:https://doi.org/10.1161/CIRCULATIONAHA.118.034165

13. Nesmith JE, Chappell JC, Cluceru JG, Bautch VL. Blood vessel anastomosis is spatially regulated by Flt1 during angiogenesis. Development. 2017;144: 889–896. doi:http://doi.org/10.1242/dev.145672

14. Karaman S, Paavonsalo S, Heinolainen K, Lackman MH, Ranta A, Hemanthakumar KA, et al. Interplay of vascular endothelial growth factor receptors in organ-specific vessel maintenance. J Exp Med. 2022;219: e20210565.

15. Sadowski L, Pilecka I, Miaczynska M. Signaling from endosomes: Location makes a difference. Exp Cell Res. 2009;315: 1601–1609. doi:https://doi.org/10.1016/j.yexcr.2008.09.021

16. Wendel Clegg L, Mac Gabhann F. Site-Specific Phosphorylation of VEGFR2 Is Mediated by Receptor Trafficking: Insights from a Computational Model. PLOS Comput Biol. 2015;11: e1004158. doi:https://doi.org/10.1371/journal.pcbi.1004158

17. Smith GA, Fearnley GW, Tomlinson DC, Harrison MA, Ponnambalam S. The cellular response to vascular endothelial growth factors requires co-ordinated signal transduction, trafficking and proteolysis. Biosci Rep. 2015;35: e00253.

18. Boucher JM, Clark RP, Chong DC, Citrin KM, Wylie LA, Bautch VL. Dynamic alterations in decoy VEGF receptor-1 stability regulate angiogenesis. Nat Commun. 2017;8: 15699. doi:https://doi.org/10.1038/ncomms15699

19. Jopling HM, Howell GJ, Gamper N, Ponnambalam S. The VEGFR2 receptor tyrosine kinase
undergoes constitutive endosome-to-plasma membrane recycling. Biochem Biophys Res
Commun. 2011;410: 170–176.
20. Gampel A, Moss L, Jones MC, Brunton V, Norman JC, Mellor H. VEGF regulates the
mobilization of VEGFR2/KDR from an intracellular endothelial storage compartment. Blood.
2006;108: 2624–2631. doi:https://doi.org/10.1182/blood-2005-12-007484
21. Bruns AF, Herbert SP, Odell AF, Jopling HM, Hooper NM, Zachary IC, et al. Ligand-
stimulated VEGFR2 signaling is regulated by co-ordinated trafficking and proteolysis.
Traffic. 2010;11: 161–174.
22. Ballmer-Hofer K, Andersson AE, Ratcliffe LE, Berger P. Neuropilin-1 promotes VEGFR-2
trafficking through Rab11 vesicles thereby specifying signal output. Blood. 2011;118: 816–
826. doi:https://doi.org/10.1182/blood-2011-01-328773
23. Jopling HM, Odell AF, Pellet-Many C, Latham AM, Frankel P, Sivaprasadarao A, et al.
Endosome-to-plasma membrane recycling of VEGFR2 receptor tyrosine kinase regulates
endothelial function and blood vessel formation. Cells. 2014;3: 363–385.
24. Mac Gabhann F, Popel AS. Model of competitive binding of vascular endothelial growth
factor and placental growth factor to VEGF receptors on endothelial cells. Am J Physiol-
Heart Circ Physiol. 2004;286: H153–H164.
25. Wendel Clegg L, Mac Gabhann F. A computational analysis of in vivo VEGFR activation by
multiple co-expressed ligands. PLOS Comput Biol. 2017;13: e1005445.
doi:https://doi.org/10.1371/journal.pcbi.1005445
26. Imoukhuede PI, Popel AS. Quantification and cell-to-cell variation of vascular endothelial
growth factor receptors. Exp Cell Res. 2011;317: 955–965.
doi:https://doi.org/10.1016/j.yexcr.2010.12.014
27. Harris LA, Hogg JS, Tapia J-J, Sekar JAP, Gupta S, Korsunsky I, et al. BioNetGen 2.2:
advances in rule-based modeling. Bioinformatics. 2016;32: 3366–3368.
doi:https://doi.org/10.1093/bioinformatics/btw469
28. Calera MR, Venkatakrishnan A, Kazlauskas A. VE-cadherin increases the half-life of VEGF
receptor 2. Exp Cell Res. 2004;300: 248–56. doi:https://doi.org/10.1016/j.yexcr.2004.07.007
29. Mittar S, Ulyatt C, Howell GJ, Bruns AF, Zachary I, Walker JH, et al. VEGFR1 receptor
tyrosine kinase localization to the Golgi apparatus is calcium-dependent. Exp Cell Res.
2009;315: 877–889.
30. Napione L, Pavan S, Veglio A, Picco A, Boffetta G, Celani A, et al. Unraveling the influence
of endothelial cell density on VEGF-A signaling. Blood. 2012;119: 5599–5607.
doi:https://doi.org/10.1182/blood-2011-11-390666
31. Wu FTH, Stefanini MO, Mac Gabhann F, Popel AS. A Compartment Model of VEGF
Distribution in Humans in the Presence of Soluble VEGF Receptor-1 Acting as a Ligand
Trap. PLoS ONE. 2009;4: e5108. doi:https://doi.org/10.1371/journal.pone.0005108
32. Fuh G, Garcia KC, de Vos AM. The interaction of neuropilin-1 with vascular endothelial
growth factor and its receptor flt-1. J Biol Chem. 2000;275: 26690–26695.
33. Baumdick M, Brüggemann Y, Schmick M, Xouri G, Sabet O, Davis L, Chin JW, Bastiaens
PIH. EGF-dependent re-routing of vesicular recycling switches spontaneous
phosphorylation suppression to EGFR signaling. eLife. 2015;4: e12223.
https://doi.org/10.7554/eLife.12223