TITLE

Optogenetic Miro cleavage reveals direct consequences of real-time loss of function in

Drosophila.

Francesca Mattedi[1], Ethlyn Lloyd-Morris[1], Frank Hirth[1], and Alessio Vagnoni[1*].

[1]Department of Basic and Clinical Neurosciences, Maurice Wohl Clinical Neuroscience

Institute, Institute of Psychiatry, Psychology and Neuroscience, King's College London, 5

Cutcombe Road, London, SE5 9RX, UK

*Corresponding author

E-mail: alessio.vagnoni@kcl.ac.uk

ABSTRACT

Miro GTPases control mitochondrial morphology, calcium homeostasis and regulate

mitochondrial distribution by mediating their attachment to the kinesin and dynein motor

complex. It is not clear, however, how Miro proteins spatially and temporally integrate

their function as acute disruption of protein function has not been performed. To address

this issue, we have developed an optogenetic loss of function 'Split-Miro' allele for precise

control of Miro-dependent mitochondrial functions in *Drosophila*. Rapid optogenetic

cleavage of Split-Miro leads to a striking rearrangement of the mitochondrial network,

which is mediated by mitochondrial interaction with the microtubules. Unexpectedly, this

treatment did not impact the ability of mitochondria to buffer calcium or their association

with the endoplasmic reticulum. While Split-Miro overexpression is sufficient to augment

mitochondrial motility, sustained photocleavage shows Split-Miro is surprisingly

dispensable to maintain elevated mitochondrial processivity. In adult fly neurons *in vivo*,

Split-Miro photocleavage affects both mitochondrial trafficking and neuronal activity.

Furthermore, functional replacement of endogenous Miro with Split-Miro identifies its

essential role in the regulation of locomotor activity in adult flies, demonstrating the

feasibility of tuning animal behaviour by real-time loss of protein function.

INTRODUCTION

Methods to observe loss of function (LoF) phenotypes are used to study many biological processes. Although important tools for elucidating gene function, disruption of genes by genomic mutations or RNA interference often does not have the spatio-temporal resolution to capture the direct cellular and organismal consequences of LoF and to report specifically on a protein's primary function. The observed 'end-point' phenotypes might thus be the result of compensatory mechanisms to loss of protein, and gene pleiotropy means that, even in cell culture models, it is often difficult to dissect the causality of the observed phenotypes.

This issue is especially problematic when studying complex cellular processes such as mitochondrial dynamics, and it is particularly evident for Miro proteins, mitochondrial Rho GTPases that influence the motility, morphology and physiology of mitochondria (1–3). In *Drosophila*, homozygous mutants of *miro* are developmentally lethal (4) while knockout of *Miro1* in mice leads to perinatal lethality (5,6). Chronic loss of Miro is detrimental for mitochondrial transport in *Drosophila* and mammalian neurons (7–9), where it leads to alteration of synaptic strength (4,10) whereas disruption of Miro in *Drosophila* additionally impairs mitochondrial calcium homeostasis (11,12). Reducing Miro abundance has also profound effects on mitochondrial morphology with fragmented mitochondria observed in yeast (13), *Drosophila* larval motor neurons (7) and mouse embryonic fibroblasts (6). It is not yet clear, however, to what extent these phenotypes are directly consequences of Miro disruption and whether they might arise independently. Any attempt to dissect the causality of cellular and organismal phenotypes after Miro manipulation is challenging as the current tools do not allow acute disruption of function.

Here we present the generation of 'Split-Miro', a photocleavable variant of *Drosophila* Miro, to achieve rapid and controlled loss of protein function. Mitochondrial network remodelling is a rapid response to Split-Miro photocleavage in *Drosophila* S2R+ cells. This effect, mediated by loss of anchorage to the microtubule network, impacts on mitochondrial distribution into cell processes where the use of Split-Miro shows that this protein is sufficient to increase mitochondrial motility but dispensable for the maintenance of elevated mitochondrial velocities. This effect is mirrored in adult fly neurons *in vivo*, where Split-Miro affects the motility, but not the processivity, of the mitochondria. Unexpectedly, we show that Split-Miro photocleavage neither directly impacts mitochondrial calcium homeostasis nor the association between the mitochondria and the endoplasmic reticulum in S2R+ cells. Finally, we demonstrate that Split-Miro modulates neuronal activity in adult flies and, by rescuing the lethality associated with classical Miro LoF mutations via pan-neuronal expression, we provide proof of concept that Split-Miro affords control of fly locomotor activity through exposure to blue light.

RESULTS

Design of a photocleavable Miro variant in Drosophila.

To gain real-time spatio-temporal control of Miro LoF, we created a Miro variant that contains the LOV2-Zdk1 protein pair (Fig 1A-B) that undergoes light-induced dissociation upon exposure to blue light (14–16) and so is predicted to achieve rapid and reversible Miro LoF through protein photocleavage (Fig 1B).

We fused Zdk1 N-terminally to the mitochondrial targeting transmembrane domain of

Miro, with LOV2 fused C-terminally to the rest of the protein (Fig 1A-B), and expressed

this optogenetic variant of Miro (herein called 'Split-Miro') as two components in

Drosophila S2R+ cells (Fig 1B-C and S1A Fig). In absence of blue light, LOV2 interacts

with Zdk1 thus reconstituting Split-Miro at the mitochondria (Fig 1C). However, upon blue

light exposure, the LOV2 Jα helix undergoes a conformational change that prevents Zdk1

from binding (14,15), resulting in the photocleavage of Split-Miro and the complete and

rapid release of the N-terminal moiety into the cytoplasm (Fig 1D-F, Movie 1).

Quantifications of the kinetics of release after blue light exposure showed release half-

life of 1.6 ± 0.2 seconds (Fig 1F) and full Split-Miro reconstitution within 3 minutes (Fig

1D-E, G) after removal of blue light, in line with what was previously reported for the

LOV2-Zdk1 association in mammalian cells (14,17). Tagging Split-Miro only at the N-

terminus did not significantly change the kinetic of photocleavage (S1B-G Fig, Movie 2)

thus showing the Split-Miro design supports a versatile tagging strategy to facilitate

fluorophore multiplexing.

Figure 1. Split-Miro is a photocleavable version of *Drosophila* Miro. A) Schematic of

Split-Miro functional domains. **B)** Schematic representation of reconstituted and

photocleaved mitochondrial-bound Split-Miro in its dark and lit state, respectively. Split-

MiroN and Split-MiroC moieties can be followed independently by tagging them with

different combinations of fluorescent proteins (e.g., EGFP/mCherry). **C)** Localisation of

Split-Miro in S2R+ cells co-transfected with mCherry-Split-MiroN (magenta) harbouring

the T406A, T407A mutations in the N-terminus of LOV2 (15,18) and EGFP-tagged Split-

MiroC, before and after a 570-ms pulse of blue light. Before irradiation, the mCherry and

EGFP signals co-localise while mCherry-Split-MiroN is fully released into the cytoplasm

(indicated by the homogenous magenta colour) immediately after irradiation (16 s), and

it fully reconstitutes after 3 minutes. Cartoon depicts photocleavage and reconstitution of

mCherry-Split-Miro-EGFP at different time points. s, seconds. **D)** Quantification of

mCherry-Split-MiroN fluorescence at the mitochondria and in the cytoplasm before

irradiation, at the point of maximum release and recovery indicates the N-terminus moiety

is completely released into the cytoplasm (mean ratio=1 at release) and fully reconstituted

afterwards. **E)** Quantification of EGFP-Split-MiroC fluorescence indicates the localisation

of the C-terminus moiety does not change during the experiment. **F-G)** Quantification of

mCherry-Split-MiroN half-time release (F) and recovery (G) after photocleavage, relative

to C. In (G), left panel shows levels of cytosolic Split-Miro N-terminus quantified after the

maximum release is reached; right panel: recovery half-time. Data are shown as mean ±

SEM. Solid line in (G) is exponential curve fit. Circles, number of cells, from three

independent experiments. Scale bars: 10 µm. **** $p < 0.0001$.

Overexpression of Split-Miro increases the proportion of motile mitochondria and

their processivity in cell processes.

Drosophila S2R+ cells are often used for intracellular trafficking experiments (19,20) and

can be induced to extend processes (Fig 2A) with a stereotypical plus-end out microtubule

array (20,21) that display Miro-dependent long-range, bidirectional mitochondrial motility

(Fig 2B-D and S2A-C Fig). With the aim of studying whether mitochondrial motility can be

manipulated via Split-Miro, we first transfected S2R+ cells with either Split-Miro or wild-

type Miro (wt-Miro) N-terminally tagged with mCherry. We initially exploited the mCherry

tag to follow mitochondria with a 561-nm laser line, which does not photocleave Split-Miro

(Fig 2E-G, 561-nm laser).

Miro links kinesin and dynein motors to mitochondria via milton (TRAK1/2 in mammals)

(8,9,22–25) (S2D-E Fig) and so overexpression of Miro is predicted to favour the

recruitment of motor proteins on mitochondria for processive transport. Consistent with

this idea, mitochondria spent more time on long runs, paused less and engaged less

frequently in short runs after either Miro or Split-Miro overexpression compared to controls

(Fig 2F, 561-nm laser, and S2F-G Fig). Thus, both Miro isoforms turn the motility of

mitochondria from predominantly bidirectional with frequent reversals to markedly more

processive. Further supporting this notion, both Miro isoforms caused a higher proportion

of mitochondria in cells processes to be motile (Fig 2G, 561-nm laser) and their duty cycle

increased in both the anterograde and retrograde directions (S2H Fig), suggesting that

Miro participates in the activation of transport complexes for bidirectional mitochondrial

transport. Interestingly, while there was no difference in the retrograde run velocities when

Miro or Split-Miro were overexpressed, mitochondria traveling in the anterograde

direction moved at nearly double the speed when compared to control (Fig 2H, 561-nm

laser). This observation might reflect a preference for Miro to recruit milton-kinesin

complexes, consistent with what was observed in larval segmental neurons after Miro

overexpression (26). Collectively, these experiments shed light on Miro's role in regulating

mitochondrial motility in S2R+ cells and demonstrate that, in absence of blue light

exposure, Split-Miro and Miro are functionally equivalent.

Figure 2. Miro regulates mitochondrial motility in the processes of S2R+ cells and

Split-Miro photocleavage reverts the effect of Split-Miro overexpression on

mitochondrial motility but not velocity. A) Example of S2R+ cell treated with

cytochalasin D to induce the formation of microtubule-rich processes. The microtubules

and the mitochondria are stained with Tubulin Tracker and with MitoTracker Green

(MitoTracker), respectively. White arrows indicate examples of cellular processes

containing mitochondria. Scale bar: 10 µm. **B)** Representative kymographs of

mitochondrial transport in the cellular processes of S2R+ cells treated with control (upper

panel) or Miro dsRNA (bottom panel). Yellow highlights indicate examples of short (< 2

µm) and long runs (≥ 2 µm). Scale bars: 1 µm (distance) and 5 s (time). **C)** Duty cycle

analysis describes the average time mitochondria spend on long runs, short runs or

pausing. For each parameter, all mitochondrial values from each cell were averaged and

compared between control and Miro dsRNA condition. **D)** Percentage of motile

mitochondria in cellular processes. Number of mitochondria analysed are in brackets,

from 29 (Ctrl dsRNA) and 36 (Miro dsRNA) cells, from two independent experiments.

Data are shown as mean ± SEM, multiple Student's t-tests (C) and Mann-Whitney test

(D). This analysis suggests Miro is necessary to drive mitochondrial motility in *Drosophila*

S2R+ cells and this is mainly due to modulation of long-range transport, while short-range

motility appears to be largely independent of Miro. **E)** Representative kymographs

showing mitochondrial transport in S2R+ cellular processes before and during exposure

to blue light. Control cells were co-transfected with mCherry-tagged wt-Miro, which cannot

be photocleaved, and EGFP targeted to the mitochondria via the Zdk1-MiroC anchor

(EGFP-mito). Split-Miro was tagged with both EGFP and mCherry to independently follow

the C-terminal and N-terminal half, respectively. Mitochondrial transport was first imaged

with a 561-nm laser, to capture the mCherry signal, and then with a 488-nm laser, to

capture the EGFP signal while photocleaving Split-Miro (Movie 4-5). Cartoon depicts

schematic of the transfected constructs. Scale bars: 5 µm (distance) and 30 s (time). **F)**

Duty cycles analysis and **G)** percentage of motile mitochondria in cellular processes. In

F-G, mitochondrial transport was first analysed following the mCherry tag in cells

expressing either mCherry-tagged wt-Miro or Split-Miro (561-nm laser). In a separate

experiment (488-nm laser), mitochondrial transport was quantified following the EGFP

tag at the beginning (1st minute) and at the end (7th minute) of the time-lapse imaging with

488-nm blue light, as shown in panel E. In F, for each parameter, all mitochondrial values

from each cell were averaged and compared to the control condition (561-nm laser) or to

the first minute of blue light exposure (488-nm laser). Data are shown as mean ± SEM,

from three independent experiments. Statistical significance was evaluated by one-way

ANOVA followed by Tukey's post-hoc test (561-nm laser) or multiple paired t-tests (488-

nm laser) in (F) and by Kruskal-Wallis test followed by Dunn's post-hoc test (561-nm

laser) or by Mann-Whitney tests (488-nm laser) in (G). Number of mitochondria analysed

are in brackets, from 16 (Control), 15 (wt-Miro) and 15 (Split-Miro) cells under 561-nm

laser, and 11 (wt-Miro) and 17 (Split-Miro) cells under 488-nm laser. **H)** Distribution of

anterograde and retrograde long run velocity after wt-Miro and Split-Miro overexpression

(561-nm laser) and after irradiation with blue light to photocleave Split-Miro (488-nm

laser). Solid lines are fitted curves. Statistical significance was calculated with a Kruskal-

Wallis test followed by Dunn's post-hoc test (561-nm laser) and a Mann-Whitney test

(488-nm laser). N = number of mitochondrial runs. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$,

**** $p < 0.0001$. **I)** Model for Split-Miro-mediated regulation of mitochondrial motility. The

Split-Miro trafficking data are consistent with a model in which two different pools of

mitochondria co-exist in S2R+ cells. After overexpression, Split-Miro recruits the motor

complexes on the mitochondria and activates transport. On one mitochondrial pool

(*mitochondrial pool 1*), Split-Miro links directly or indirectly (for example, via Milton/Trak,

not depicted) to the motor proteins. On another subset of mitochondria (*mitochondrial*

pool 2), the motor complexes are instead stabilised on the organelle by an unknown factor

(question mark) following Miro-dependent recruitment. After Split-Miro photocleavage,

mitochondrial pool 1 can reverse to control levels of processivity or become stationary

(Fig 2F-G). However, the processivity (e.g., velocity and run length) of the *mitochondrial*

pool 2 is not affected as the recruited motors are not directly linked to the mitochondria

via Miro (Fig 2H and S2J-K Fig). This model is consistent with previous reports describing

Miro-independent mitochondrial motility (6,24). Split-Miro-dependent recruitment of motor

complexes on an interconnected mitochondrial network, such as the one found in the

perinuclear area of the cell soma, could cause significant tension on the network.

Releasing the Split-Miro anchor by photocleavage, even if only on a subset of the

mitochondria, would be sufficient to release the tension and cause mitochondrial network

retraction, as shown in Fig 3. Not depicted, dynein motor complex.

Optogenetic cleavage of Split-Miro reduces the proportion, but not the

processivity, of motile mitochondria in the cell processes.

To test whether Split-Miro photocleavage reverses the observed Miro gain-of-function

effects on mitochondrial transport, we analysed mitochondrial motility under blue light in

wt-Miro and Split-Miro transfected cells. Time-lapse imaging with the 488-nm laser line

ensured Split-Miro was not reconstituted while recording mitochondrial motility via the

EGFP-tag (Fig 2E, green). wt-Miro, which cannot be photocleaved, was used as a control.

We found that while there was no detectable difference in the motility after imaging for

one minute (S2I Fig), sustained Split-Miro photocleavage (7 minutes) reduced the time

mitochondria spent on long runs and the proportion of organelles in the processes that

were motile to the levels observed prior to overexpression, while there was no such effect

in wt-Miro controls (Fig 2E-G, 488-nm laser). Cleaving Split-Miro from mitochondria did

not, however, have any major effects on the velocity and run length of the moving

organelles, which remained elevated and did not return to control levels (Fig 2H, 488-nm

laser, and S2J Fig). This was also the case when endogenous Miro was depleted in the

Split-Miro condition by RNAi (S2K Fig). Thus, using a strategy combining upregulation

(via overexpression) and downregulation (via Split-Miro photocleavage), these results

indicate that the proportion and processivity of motile mitochondria are controlled by

separate Miro-dependent and Miro-independent mechanisms (Fig 2I).

Split-Miro photocleavage triggers a rapid collapse of the mitochondrial network

which is rescued by mitochondrial anchoring to the microtubules.

We next examined the effect of Split-Miro on the integrity of the mitochondrial network in

the cell soma. Strikingly, exposing Split-Miro-transfected S2R+ cells to blue light triggered

a rapid (< 3 min) and dramatic remodelling of the entire mitochondrial network which

progressively collapsed towards the centre of the cell (Fig 3A, Movie 3). Mitochondria

shortened along their long axis taking up a rounder shape (Fig 3B), which was associated

with a strong reduction in the number of branches (Fig 3C) and reduction of the total area

covered by mitochondria (S3A Fig). Again, the presence of endogenous Miro was

dispensable for Split-Miro functionality as reducing Miro levels by RNAi did not affect the

mitochondrial phenotype in Split-Miro-transfected cells (Fig 3D). This phenotype is not

dependent on adverse effects caused by increased cytoplasmic concentration of Split-

MiroN, as overexpression of the Miro N-terminal moiety alone, which diffuses throughout

the cytoplasm, does not cause any gross mitochondrial morphological aberration (S3B

Fig and (24,25)). Of note, neither the motility and distribution of peroxisomes in the cell

processes (S3C-D Fig) nor the overall organisation of the microtubule network (S3E Fig)

were affected by this rapid cellular-scale change. Nevertheless, we reasoned that

mitochondrial network collapse might impede organelle delivery towards the periphery.

The number of mitochondria in the cell processes showed a progressive decline in

mitochondrial content in cells expressing Split-Miro under blue light, while no effect was

observed in the presence of wt-Miro (S3F-G Fig). We conclude that the rapid

mitochondrial shape transition with associated loss of network integrity, clearly detectable

within 3 minutes under blue light, strongly contributes to the progressive depletion of

mitochondria from the cell processes.

The Miro-milton-motor complex provides a link for the attachment of mitochondria onto

the microtubules (27,28). To test the hypothesis that loss of microtubule tethering is

responsible for the collapse of the mitochondrial network, we set out to induce

mitochondrial tethering to the microtubules in a Miro-independent manner. In mammals,

syntaphilin (SNPH) anchors mitochondria onto the microtubules (29), although a

Drosophila homologue has not yet been found. Thus, we co-expressed EGFP-tagged

human SNPH with Split-Miro in S2R+ cells stained with MitoTracker. SNPH signal in

Drosophila cells overlaps with mitochondria and, as observed in mammalian neurons,

SNPH puncta often localise at mitochondrial ends and are associated with strong

reduction in mitochondrial dynamics (Fig 3E-E' and S3H-I Fig). While in cells devoid of

SNPH the mitochondrial network retracts after Split-Miro photocleavage, the presence of

SNPH prevents this phenotype (Fig 3E-G). These observations support the notion that

loss of mitochondrial anchoring on microtubules is responsible for the rapid mitochondrial

network collapse when Split-Miro is cleaved.

Figure 3. Split-Miro-dependent changes of mitochondrial morphology and

connectivity are rescued by SNPH expression. A) Representative images of

mitochondria in *Drosophila* S2R+ cells at the beginning (0 min) and after 3 and 7 minutes

of exposure to blue light, which leads to Split-Miro, but not wt-Miro, photocleavage.

Control cells (top panels) were co-transfected with mCherry-tagged wt-Miro (magenta)

and EGFP targeted to the mitochondria via the Zdk1-MiroC anchor (EGFP-mito, grey).

Split-Miro (bottom panels) was tagged with both EGFP and mCherry to independently

follow the C-terminal and N-terminal half, respectively. The mCherry is shown at the

beginning and end of the imaging period to confirm retention on and release from the

mitochondria in wt-Miro and Split-Miro transfected cells, respectively, under blue light.

Scale bar: 10µm. **B)** Quantification of mitochondrial aspect ratio (AR) and **C)** of the

number of mitochondrial branches within the network, relative to A. **D)** Quantifications of

the mitochondrial collapse phenotype after 7 minutes of time-lapse imaging with 488-nm

blue light in S2R+ cells overexpressing wt-Miro and Split-Miro with or without a Miro

dsRNA construct (Miro^{RNAi}). Number of cells: wt-Miro = 15, Split-Miro = 15, Split-Miro +

Miro^{RNAi} = 10, Fisher's exact test. **E)** Representative images of cells expressing mCherry-

Split-Miro (Split-Miro, yellow) with either an empty vector (top panels) or with EGFP-

SNPH (SNPH, bottom panels, grey) and stained with MitoTracker DeepRed (MTDR,

magenta). White arrows show examples of mitochondria that have retracted after Split-

Miro photocleavage. Diffuse cytoplasmic yellow signal indicates release of Split-Miro N-

terminus from the mitochondria. Scale bar: 10 µm. **E')** Magnified inset shows examples

of stable SNPH-positive mitochondria (white arrowheads) and dynamic mitochondrial

membranes devoid of SNPH (magenta arrowheads). Scale bar: 2 µm. Not shown, Split-

Miro. **F)** Quantification of mitochondrial AR and **G)** number of mitochondrial branches at

the time points indicated, relative to E. Circles represent the average AR calculated from

single mitochondria within the same cell (B, F) and the average number of branches per

cell normalised to the average group value (Split-Miro, Split-Miro + SNPH) at timepoint 0

(C, G). Comparison across timepoints was performed by repeated-measures one-way

ANOVA followed by Tukey's post-hoc test (B-C, F) and Friedman test followed by Dunn's

post-hoc test (G), from three independent experiments. Data are reported as mean ±

SEM. * p < 0.05, ** p < 0.01, **** p < 0.0001.

Split-Miro photocleavage does not affect mitochondrial calcium buffering or

mitochondria-ER association.

Mitochondria buffer calcium to help maintain cellular homeostasis and loss of Miro

reduces calcium levels in the mitochondria of the *Drosophila* brain (11,12,30). However,

the mechanisms underlying decreased calcium uptake when Miro is disrupted are not

understood. In S2R+ cells, Split-Miro photocleavage did not induce any changes in the

fluorescent intensity of the mitochondrial calcium $[Ca^{2+}]_m$ indicator mito-GCaMP6f when

compared to control cells (Fig 4A-B), indicating that the steady-state level of $[Ca^{2+}]_m$ is

not affected by this manipulation. Optogenetic inactivation of Split-Miro also did not affect

$[Ca^{2+}]_m$ uptake when S2R+ cells were challenged with ionomycin, an ionophore that

causes a sharp increase in cytosolic calcium (27,31) (Fig 4C-E). This result indicates that

the mitochondrial morphological changes induced by Split-Miro are not sufficient to alter

$[Ca^{2+}]_m$ homeostasis. Likewise, overexpression of SNPH, which rescues the Split-Miro-

induced mitochondrial collapse, did not have any effect on $[Ca^{2+}]_m$ uptake when Split-Miro

was cleaved (Fig 4C-E).

It has been shown that a subset of Miro is found at the mitochondria-ER interface to

regulate the contacts between these two organelles (12,32,33) and the mitochondria-ER

contacts sites (MERCS) are known to mediate Ca^{2+} exchange between the two organelles

(34). However, we did not detect any significant difference in the number of MERCS

visualised using a split-GFP assay (35,36) after Split-Miro photocleavage (Fig 4F-H).

Furthermore, the morphological changes displayed by mitochondria after Split-Miro

photocleavage were not associated with detectable changes in their membrane potential

(S4 Fig), implying that mitochondria do not become dysfunctional during this rapid

morphological transition. Together these findings suggest that acutely modulating

mitochondrial network integrity via Split-Miro/SNPH or blocking mitochondrial motility via

SNPH (S3F-G Fig) are not sufficient to perturb $[Ca^{2+}]_m$ homeostasis in this context.

Figure 4. Split-Miro photocleavage does not affect $[Ca^{2+}]_m$ abundance. A)

Representative image of an S2R+ cell showing mitochondrial targeting of mito-GCaMP6f

(cyan) in the perinuclear region (white arrows) and in single mitochondria (arrowheads).

Yellow is mCherry-tagged wt-Miro (wt-Miro), magenta is MitoTracker DeepRed (MTDR).

Scale bar: 5 μm. **B)** Cells transfected with mito-GCaMP6f and either mCherry-wt-Miro

(wt-Miro) or mCherry-Split-Miro (Split-Miro) were stained with MTDR and the ratio of mito-

GCaMP6f/MTDR signal intensity was analysed at the beginning (1st minute) and at the

end (7th minute) of the time-lapse imaging under blue light. Number of cells: wt-Miro = 12,

Split-Miro = 16; from three independent experiments. Data are shown as mean ± SEM.

Repeated measures one-way ANOVA followed by Tukey's post-hoc test did not show any

significant difference between groups, indicating the basal $[Ca^{2+}]_m$ does not significantly

change after Split-Miro photocleavage. **C)** Expression of mCherry-Split-Miro (Split-Miro)

or EBFP-SNPH (SNPH) does not significantly alter $[Ca^{2+}]_m$ uptake in cells challenged with

ionomycin, compared to control conditions. Traces indicate the average mito-GCaMP6f

fluorescence intensity values (circles) at individual time point before and after cell

exposure to ionomycin (arrow). N = number of cells, from five independent experiments.

D-E) Normalised response peak and time to reach the peak, respectively, relative to the

data shown in C. Circles, number of cells. Kruskal-Wallis test followed by Dunn's multiple

comparisons showed no difference between conditions. **F)** Wild-type Miro or Split-Miro

(magenta) were co-expressed with the ER-mito::SPLICS probe (cyan) in S2R+ cells. The

SPLICS probe displays a typical punctuated stain in these cells, as previously observed

in mammalian and *Drosophila* cells (34,36). **G-H)** Quantification of the SPLICS puncta

and statistical analysis by Wilcoxon test showed no significant difference between the

beginning (1st minute) and the end (7th minute) of the time-lapse imaging under blue light

in wild-type and Split-Miro-transfected cells. Data are presented as % change of SPLICS

puncta at 7th minute compared to 1st minute. Number of contacts analysed are in brackets

from 5 (wt-Miro) and 9 (Split-Miro) cells from two independent experiments.

In vivo regulation of mitochondrial motility by Split-Miro.

To gain insight into the *in vivo* roles of Split-Miro in regulating mitochondrial dynamics, we

expressed *UAS-mCherry-MiroN-LOV2* and *UAS-EGFP-Zdk1-MiroC* (herein '*UAS-Split-*

Miro') with the *Appl-Gal4* driver and imaged mitochondrial motility in the axons of the adult

wing neurons *in situ* (37). Split-Miro is reconstituted in adult fly neurons and can be

photocleaved efficiently through exposure to blue light (S5A-D Fig). Overexpression of

Miro in *Drosophila* neurons *in vivo* has been shown to be detrimental for mitochondrial

motility and overall neuronal viability (7,20). However, expressing Split-Miro and the wt-

Miro control in a *miro^+/-^* background is permissive towards mitochondrial motility (Fig 5A-

B and S5E-F Fig, Movie 6). Tracking of mitochondrial movements revealed that, while

exposing the *UAS-wt-Miro^+^* neurons to pulses of blue laser light did not cause any change

in the number of motile mitochondria in 2-d-old flies (Fig 5A,C, Movie 6), *UAS-Split-Miro^+^*

neurons showed a strong decline after 5 minutes (Fig 5B,D). Mitochondrial velocity and

run length, however, were not affected by this manipulation (Fig 5E-F), similarly to what

observed in S2R+ cells after Split-Miro photocleavage. Interestingly, 7-d old *UAS-Split-*

*Miro^+^* neurons did not show any changes in mitochondrial motility under the same

experimental conditions (S5E-H Fig). This suggests that in older flies, when the rates of

mitochondrial transport are known to decline (20,38), alternative mechanisms might

become predominant towards controlling mitochondrial trafficking.

Figure 5. Split-Miro-dependent control of mitochondrial motility *in vivo.* A-B) Stills

from movies of GFP-labelled mitochondria in wing neuronal axons expressing *UAS-wt-*

Miro (A) or *UAS-Split-Miro* (B) in *miro$^{Sd32/+}$* background during the first minute (top panels)

and fifth minute (bottom panels) of blue light exposure. Traces of transported

mitochondria in corresponding movies are overlayed onto the images. **C-D)** Number of

motile mitochondria captured in a 50-μm axonal tract in wing neurons expressing *UAS-*

wt-Miro (C) or *UAS-Split-Miro* (D). Bar charts show the average mitochondrial content at

each time point. Filled circles represent the number of mitochondria within each axonal

bundle at minute 1 and 5. Data were analysed by paired Student's t-test. Number of wings

analysed: *UAS-wt-Miro* = 8, *UAS-Split-Miro* = 8, from two independent experiments. **E-F)**

Anterograde velocity (E) and run length (F) of axonal mitochondria in wing neurons

expressing *UAS-wt-Miro* or *UAS-Split-Miro*, during the first and fifth minute of blue light

exposure, relative to C-D. Due to the overall lower number of bidirectional and retrograde-

moving mitochondria, a meaningful statistical analysis of their velocity and run length is

not possible. Circles represent tracked mitochondria. Data mean ± SEM, Mann-Whitney

test. *** $p < 0.001$.

Split-Miro controls neuronal activity in vivo.

Having observed a strong reduction in mitochondrial trafficking in 2d-old Split-Miro flies,

we next asked whether rapid Split-Miro photocleavage could affect wider neuronal

physiology. Reduced mitochondrial motility has been associated with alterations in

neuronal activity (4,10,39,40). Thus, we decided to measure the activity of adult wing

neurons by recording spontaneous Ca^{2+} transient with the GCaMP5 Ca^{2+} indicator (Fig

6A-B). As done for the trafficking experiment, we expressed Split-Miro and the wt-Miro

control in a *miro$^{+/-}$* background and recorded basal neuronal responses under exposure

to blue laser light. We found that *UAS-Split-Miro$^+$* neurons displayed a milder and slower

response compared to *UAS-wt-Miro$^+$* control neurons (Fig 6C-E). Interestingly, the

dampened neuronal firing was offset by an enlarged area of active neurons (Fig 6F),

suggesting that Split-Miro photocleavage induced distributed network performance which

characterises neural ensembles and manifolds (41,42).

Figure 6. Basal neuronal activity is affected by Split-Miro. A-B) Examples of

spontaneous bouts of neuronal activity, indicated by transient fluorescence increase of

the GCaMP5 Ca^{2+} indicator in the cell body (A) and axonal bundle (B) of adult wing

neurons. White circle, neuronal soma (A). In B, the arrows indicate a single axon within

the axonal bundle. Scale bars: 5 μm. **C)** Pan-neuronal expression of *UAS-Split-Miro*

significantly reduces the amplitude of Ca^{2+} transients, compared to *UAS-wt-Miro*. Traces

indicate the average GCaMP5 fluorescence intensity at individual time points in neuronal

axons. N = number of bouts of activity from 16 wings (*UAS-wt-Miro* = 8; *UAS-Split-Miro* =

8) and two independent experiments. **D-E)** Peak of GCaMP fluorescence expressed as

fluorescence fold increase from baseline (D) and time to reach the peak (E). Circles,

combined bouts of activity from neuronal soma and axons, relative to A-C. Data are mean

± SEM, Mann Whitney test. **F)** Quantification of the number of active neuronal regions

after 5 minutes of time-lapse imaging with 488-nm blue light in *UAS-wt-Miro* and *UAS-*

Split-Miro flies. The number of neuronal regions (cell bodies and axons) that respond

were counted in 16 (*UAS-wt-Miro*) and 15 (*UAS-Split-Miro*) fields of view, Fisher's exact

test. * p < 0.05, *** p < 0.001, **** p < 0.0001. Refer to S3 Table for full genotypes.

Inducible hyperactivity of Split-Miro flies is exacerbated by age.

Genetic mutations and RNAi have shown that Miro is critical for mitochondrial functionality in the nervous system (5,7,8). Homozygous *miro* gene loss of function alleles are lethal, thus precluding a comprehensive analysis of Miro function in adult animals. Conditional loss of Miro1 in mouse neurons causes severe movement defects within 30 days post-natal (5) and both increased and reduced Miro abundance in the *Drosophila* nervous system can rescue fly climbing activity in models of neurodegeneration (43,44). These findings suggest that real-time Miro disruption in the adult nervous system could be exploited to manipulate animal behaviour.

Having demonstrated that Split-Miro affects mitochondrial functionality in S2R+ cells and neuronal physiology *in vivo,* we turned to behavioural genetics to assess Split-Miro versatility for the study of organismal phenotypes. Pan-neuronal expression of *UAS-mCherry-MiroN-LOV2* and *UAS-Zdk1-MiroC* ('UAS-Split-Miro') with the *Appl-Gal4* driver rescued the lethality associated with classic null mutations of the Miro gene (*miro$^{Sd32/B682}$*) (Fig 7A and S5A Fig). Adult flies were assayed for their motor behaviour at day 2 and day 8 before, during and after blue-light exposure using the 'Opto-DART' system (Fig 7B). *miro$^{Sd32/B682}$*, *UAS-Split-Miro* flies at 2 days of age showed a modest increase in locomotor activity after an hour of exposure to blue light, which was sustained also after the exposure to light ceased (Fig 7C-E). These observations suggest that the effects of Split-Mito photocleavage were not rapidly reversed after removal of blue-light.

Strikingly, 8 days-old *miro$^{Sd32/B682}$*, *UAS-Split-Miro* flies displayed a rapid and sustained increase in motor activity when exposed to blue light compared to control, while no

difference was observed in their baseline activity prior to exposure to blue light (Fig 7F-

H). While the fly speed returned to control level as the blue light was turned off, the action

initiation revealed a prolonged effect on locomotor activity, similar to what observed in

younger flies (Fig 7F-H). Remarkably, Split-Miro-dependent effects on fly activity and

action initiation could be reversed by co-expressing human SNPH (*UAS-SNPH*) with

Appl-Gal4 (Fig 7I-K). Thus, SNPH can also suppress the effect of Miro inactivation in the

context of animal behaviour. Together, these results show that acute loss of Miro function

leads to fly hyperactivity, and this phenotype is exacerbated by ageing and suppressed

by expression of SNPH. Collectively, these experiments reveal that Split-Miro is effective

in an adult animal and uncover a previously unknown role of Miro in the regulation of

Drosophila locomotor behaviour.

Figure 7. Optogenetic control of *Drosophila* locomotor behaviour in Split-Miro flies.

A) Adult *miro$^{Sd32/B682}$* flies eclosed after *Appl-Gal4*-driven expression of *UAS-wt-Miro*,

UAS-Split-Miro or *UAS-Split-Miro + UAS-EGFP-SNPH* (*UAS-SNPH*) were counted from

6, 5, and 3 independent crosses, respectively. Data are reported as percentage of flies

expected from mendelian ratios. Data are mean ± SEM, one-way ANOVA with Tukey's

post-hoc test showed no statistical difference. **B)** Schematic of the 'Opto-DART'

behavioural setup consisting of a custom-made optogenetic enclosure equipped with

LEDs for blue light stimulation and a camera to record fly activity. Flies are transferred to

a 1D platform for automated recording of activity using the DART system[45,46]. DAC:

Digital-to-Analog Converter for multi-platform integration. **C-H)** Overall activity, action

initiation, and average speed of 2-day old flies (C-E) and 8-day old flies (F-H) expressing

UAS-wt-Miro or UAS-Split-Miro, before, during and after blue light exposure (shaded blue

rectangles). In C and F, the average activity of flies expressing UAS-wt-Miro vs UAS-

Split-Miro is: *before light exposure*, 3.78% ± 0.94 vs 2.34% ± 0.72 (p = 0.11)(C) and 5.62%

± 1.02 vs 8.7% ±1.81 (p = 0.14)(F); *under blue light*, 9.52% ± 0.96 vs 7.32% ± 0.81 (p =

0.13)(C) and 8.45% ± 0.82 vs 21.87% ± 2.07**** (F); *after blue light exposure*, 3.36% ±

1.18 vs 7.85% ± 1.96 (p = 0.13)(C) and 3.23% ± 1.1 vs 13.19% ± 2.39** (F). **I-K)** Overall

activity, action initiation, and average speed of 8-day old flies expressing *UAS-Split-Miro*

or *UAS-Split-Miro + UAS-SNPH*, before, during and after blue light exposure (shaded

blue rectangles). In I, the overall activity of flies expressing *UAS-Split-Miro* vs *UAS-Split-*

Miro + UAS-SNPH is: 6.09% ± 0.89 vs 0.95% ± 0.35**** before light exposure; 18.62% ±

1.4 vs 3.55% ± 0.75**** under blue light; 9.9% ± 1.4 vs 3.91% ± 1.25* after blue light

exposure. In C-K, n = number of flies. Values are means ± SEM, Mann-Whitney test

(C,F,I) and multiple unpaired t-test (D-E, G-H, J-K) with Holm-Sidak correction for multiple

comparisons. * p < 0.05, ** p < 0.01, **** p<0.0001.

DISCUSSION

Using optogenetics to implement a real time LoF paradigm by targeting Miro, we show

that collapse of the mitochondrial network is an immediate response to Miro

photocleavage in S2R+ cells which temporally precedes the defects observed in

mitochondrial trafficking. We found that Miro overexpression increases the proportion and

the processivity of mitochondria transported in the processes of S2R+ cells. Surprisingly,

although sustained Split-Miro photocleavage reverted the proportion of transported

mitochondria to control levels, the velocities and run lengths of the motile organelles were

largely unaffected. Interestingly, Split-Miro photocleavage decreases the number, but not

the velocities and run length, of motile mitochondria in adult fly neurons *in vivo*,

suggesting that a potentially similar mechanism might account for the regulation of

mitochondrial motility in the processes of S2R+ cells and in adult neurons. This scenario

is consistent with an essential role for Miro in the recruitment of transport complexes for

activation of bidirectional transport, likely by recruiting (47,48) or directly activating (49)

molecular motor complexes. However, we hypothesise that once motors have been

recruited onto mitochondria, they may link to the organelles via additional factors in a

Miro-independent way, at least on a proportion of mitochondria (Fig 2I). Future studies

should focus on discovering these factors, for example by testing if the functional

homologs of metaxins (50) in *Drosophila* could fulfil this role.

We previously showed that mitochondrial motility declines with age in the axons of

Drosophila neurons (20,38), which can be partly rescued by boosting the cAMP/PKA

signalling pathways and the abundance of the kinesin-1 motor protein (38). However, the

mechanisms underlying transport decline are still not clear. In the current study, we

observed that Split-Miro photocleavage decreases mitochondrial motility in 2d-old flies

while there was no effect in 7d-old flies. This result is intriguing as it suggests that, while

important to maintain transport in young flies, Miro is largely dispensable to sustain the

less abundant transport typically observed at later stages (20) implying that Miro-

independent mechanisms may become predominant.

Rapid retraction of the mitochondrial network in the cell soma after Split-Miro

photocleavage is conceivably a consequence of releasing membrane tension that

accumulates under stretch, reminiscent of the recoil of daughter mitochondria after fission

(51). Increased mitochondrial tension following Miro overexpression is consistent with the

idea that more motors are recruited and pull onto the mitochondrial membrane via

cytoskeletal interaction (52–55), which would likely contribute to the build-up of

mechanical energy onto an interconnected network. According to this view, releasing the

link between mitochondria and the microtubules then triggers the rapid collapse of the

network. It would be interesting in future investigations to establish whether the phenotype

that we observe could be regarded as a 'mitoquake', i.e. rapid mitochondrial network

disruption with associated release of mechanical energy, similar to the sudden

cytoskeletal rearrangements ('cytoquakes') that were proposed to underpin mechanical

adaptivity during cellular dynamic processes (56).

We found that using SNPH to tether mitochondria onto the microtubule network prevented

Split-Miro-induced mitochondrial reorganisation, indicating that Miro stabilises the

mitochondrial network by providing an anchor to the cytoskeleton. It is possible, however,

that Miro may stabilise the mitochondrial network by simultaneously bridging

mitochondria to different cellular structures. In this regard, we did not find that the actin

cytoskeleton plays a significant role in S2R+ cells (S6 Fig), although it is conceivable that

the actin network contributes to mitochondrial stability via Miro-Myosin interactions in

other cell types (6,57–59). We showed that Split-Miro photocleavage does not affect

mitochondria-ER contacts using a split-GFP reporter, suggesting that the associations

mitochondria establish with the ER do not significantly contribute to the stability of the

mitochondrial network. Testing whether Miro-dependent mitochondrial interaction with

other cellular structures is necessary to maintain mitochondrial network stability is a goal

for future studies.

Decreasing the abundance of Miro by RNAi reduced $[Ca^{2+}]_m$ levels in the neurons of

Drosophila brain (11,12) and mutating the Ca^{2+}-binding EF domains of Miro reduced

$[Ca^{2+}]_m$ uptake in mouse hippocampal neurons (60), although Miro1-KO and Miro-EF

mutant MEFs did not show any disturbances in $[Ca^{2+}]_m$ homoeostasis (5,27). In S2R+

cells, shedding Split-Miro functional domains (including the Ca^{2+}-binding EF-hands

motifs) did not have any effect on $[Ca^{2+}]_m$ uptake. We hypothesise that impaired $[Ca^{2+}]_m$

homeostasis shown with classical Miro LoF approaches (i.e., RNAi, knock-out) may be a

secondary effect of Miro LoF, potentially a consequence of sustained morphological and

transport defects of mitochondria. However, because Miro was shown to interact with

MCU and the Sam/MICOS complexes (31,32), presumably via direct interaction with the

Miro transmembrane domain, we cannot exclude that the short mitochondrial targeting

sequence might still mediate $[Ca^{2+}]_m$ homeostasis. Our finding that MERCS were not

affected by Split-Miro photocleavage supports the idea that $[Ca^{2+}]_m$ uptake, known to be

regulated by MERCS, is not a primary role of Miro in this context.

Overexpressing SNPH to rescue Split-Miro-dependent mitochondrial network retraction

also did not affect $[Ca^{2+}]_m$ uptake. Because SNPH also locks mitochondria into a

stationary state with little network dynamic, these results raise the intriguing possibility

that mitochondrial movements are not critical to maintain $[Ca^{2+}]_m$ homeostasis, as long as

mitochondria maintain their functionality.

Elegant methods for light-induced repositioning of trafficked vesicles and mitochondria

have been developed which are based on the recruitment of truncated forms of motor

proteins to overpower the endogenous transport machinery and so to achieve controlled

redistribution of cellular cargoes (61–64). Engineering the LOV2-Zdk1 domain into

subunits of motor and adaptor proteins could offer a complementary strategy for studying

intracellular trafficking when a real-time LoF approach is preferred. Because protein

photocleavage is reversible, the LOV2-Zdk1 methodology also offers significant

advantages over existing methods based on the rapid, non-reversible, degradation of a

target protein by the proteasome, such as the degron (65) or the TRIM-away systems

(66).

Opsin-based optogenetic approaches to activate/repress specific neurons and study

associated animal behaviour have been extensively used in live animals (67). A LOV2-

controlled CaMKII inhibitor was used to impair memory formation in live mice after blue-

light stimulation for 1 hour (68). By creating Split-Miro flies, we combine optogenetics with

Drosophila behaviour and neuronal specificity to perform LoF experiments in adult

animals in real time. Although the locomotor behaviour of Split-Miro flies is

indistinguishable from the wild-type counterpart before exposure to blue light, their activity

is enhanced under blue light and this hyperactivity becomes more pronounced with age.

The mechanisms underlying the exacerbated hyperactivity phenotype in older flies are

unexplained. It is conceivable that the alteration of neuronal activity observed in young

Split-Miro flies is linked to an acute imbalance of synaptic transmission and leads to

augmented locomotor activity. We speculate that, in older flies, a potentially compromised

cellular state might contribute to amplify this response. In line with this notion, ageing has

been associated with increased activity of excitatory neurons in *C. elegans*, flies and mice

(69–72). We are mindful that we performed these experiments in the wing neurons of the

flies and, although these cells can relay signals to affect motor phenotypes (73,74), we

are cautious not to generalise our findings to all neurons in the fly.

The observed reduction in mitochondrial axonal transport induced by Split-Miro may also

be a contributing factor towards enhanced fly activity, at least in young flies, by causing

an imbalance in neuronal activity. It is known that reducing mitochondrial number

positively correlates with activity-dependent vesicular release at the presynapses of

hippocampal and cortical neurons (10,39) and with miniature excitatory junction potentials

at the *Drosophila* NMJ (4). In this view, the remarkable rescue of Split-Miro-induced

hyperactivity in older animals by SNPH overexpression could conceivably occur via

further reduction of the remaining mitochondrial transport in older neurons or via a direct

effect on synaptic mitochondria (40). Overall, our data point to an important role of

neuronal mitochondrial mobility for animal behaviour and suggest that Miro could play a

crucial role in preventing hyperexcitation in the ageing nervous system with potential

ramification in the context of neurodegeneration.

MATERIALS AND METHODS

Generation of plasmid DNA.

The new constructs produced in this study are reported in S1 Table and were generated

either through restriction enzymes mediated cloning or NEBuilder HiFi DNA Assembly

(NEB) using the primers listed in S2 Table. The plasmid inserts were amplified by PCR

using the Q5® Hot-Start High-Fidelity 2X Master Mix (NEB). Site-directed mutagenesis

was performed using the Q5® Site-directed mutagenesis kit (NEB) following the

manufacturer's instructions. The fidelity of all assembled constructs was verified by

Sanger sequencing (Eurofins Genomics).

Isolation of Drosophila Miro cDNA.

Five Oregon-R flies were anesthetised on dry ice, rapidly grinded to powder with a plastic

pestle in an Eppendorf tube and the total RNA extracted with the RNeasy Micro kit

(Qiagen) following the manufacturer's instructions. Reverse transcription was performed

with the iScript Select cDNA Synthesis kit (BioRad), using 1 µl of total RNA and Oligo dT

primers. *Drosophila* Miro is encoded by a single gene, located on the third chromosome,

from which potentially four different transcripts are produced (FlyBase reference:

FBgn0039140). The coding sequence of the longest Miro-RE/Miro-RF isoform was

amplified from the total *Drosophila* cDNA with the primer pair #1 and #2 (S2 Table) and

was used as a template to engineer wild type Miro and the Split-Miro variants employed

in this study.

Cell culture, transfection and RNAi.

Drosophila S2R+ cells were obtained from the Drosophila Genomics Research Centre

(DGRC, Indiana University) and cultured in Schneider's insect medium (Gibco) at 25°C.

Cells were transfected using FuGene HD transfection reagent (Promega) using a

DNA:FuGene ratio of 1:3 following the manufacturer's instructions.

For RNAi experiments, 532 bp (Miro) and 599 bp (Control) dsRNA molecules were

transcribed with the MEGAScript RNAi kit (Thermo Scientific) from templates generated

by PCR of sequences within the Miro 3'-UTR (targeting only endogenous, and not

overexpressed, Miro and Split-Miro) and the pT2-DsRed-UAS plasmid backbone,

respectively. Primers used are listed in S2 Table. Cells were treated with 15 µg/ml dsRNA

for a total of 6 days, replacing the dsRNA every 24 h in fresh media, prior to imaging.

Drosophila* husbandry and generation of transgenic flies*.

The fly strains used in this study are listed in S3 Table. Flies were maintained on 'Iberian'

food [70 mg/ml yeast (Brewer's yeast, MP Biomedicals, 903312), 55 mg/ml glucose

(VWR, 10117HV), 7.7 mg/ml agar (SLS, FLY1020), 35 mg/ml organic plain white flour

(Doves Farm, UK), 1.2 mg/ml Nipagin (Sigma, H3647), 0.4% propionic acid (Sigma-

Aldrich, P5561] at 25°C and 60% humidity with a 12-h-light-12-h-dark cycle.

The transgenic fly lines generated in this study were obtained by phiC31-mediated

transgenesis to integrate the relevant constructs into either the attP40 (25C6) or attP2

(68A4) landing sites following embryo injection.

Immunoprecipitation.

Cells were transfected in 10-cm dishes, washed with PBS and incubated with 900 μl IP

lysis buffer (50 mM Trizma pH 7.4, 150 mM NaCl, 5 mM EDTA, 0.5% Triton X-100, 1x

PhosSTOP^{TM}, 1x cOmplete EDTA-free protease inhibitor cocktail) on ice for 20 minutes.

The cell lysate was homogenised through a 23G syringe needle and centrifuged at 21'000

g for 30 min at 4°C. 25 μl of GFP-Trap beads (Chromotek) were equilibrated by washing

3x times in IP lysis buffer prior to the incubation with 1 mg of cleared lysate at 4°C

overnight. After washing the beads 3x times with IP buffer, the immunoprecipitated

material was eluted in 20 μl of 2X NuPAGE sample buffer supplemented with 40 mM DTT

for 10 minutes at 95°C.

Western blotting.

Samples were loaded on a NuPAGE 4-12% Bis-Tris protein gel and transferred onto an

Immobilion PVDF membrane (Merk-Millipore) for 1 h at 35 V in a wet transfer apparatus

buffered with 12.5 mM Trizma-Base (Sigma), 96 mM glycine (Sigma), 10% methanol

(Fisher Chemicals). The membrane was incubated in blocking solution [5% milk in PBST

(PBS (pH 7.4) with 0.1% Tween-20 (VWR))] for 2 h and probed overnight with the

following primary antibodies: anti-βtubulin (1:100, Developmental Studies Hybridoma

Bank, DSHB #7, AB_2315513), anti-dMiro (75) (1:50000), anti-Milton (76) (1:1000,

monoclonal antibody 2A108), anti-GFP (1:1000, Chromotek, PABG1). Membranes were

incubated for 1 h with either an IRDye secondary antibody (for LI-COR Odyssey imaging)

or with horseradish peroxidase (HRP)-conjugated secondary antibodies followed by a 3-

min incubation with a chemiluminescent substrate (GE Healthcare) for ChemiDoc

imaging. Secondary HRP-conjugated antibodies: anti-rabbit (1:5000, NIF824), anti-

mouse (1:5000, NIF825), anti-guinea pig (1: 5000, SeraCare 5220-0366). LI-COR

secondary antibody: IR Dye 800CW goat anti-mouse IgG (1:10000).

Live cell imaging of cargo transport and photostimulation.

S2R+ cells were seeded in 4-well Nunc Lab-Tek chambered coverglass (Thermo

Scientific) coated with 0.5 mg/ml Concanavalin A (Sigma). To induce the formation of

cellular processes, cells were treated with 1 μM cytochalasin D (Sigma) for 3-4 h before

imaging and throughout the imaging period. Before imaging, cells were stained with the

following dyes, depending on the specific experimental setting: 200 nM MitoTracker

Green FM, 200 nM MitoTracker DeepRed FM (Thermo Scientific), 100 nM MitoView 405

(Biotium), 1X TubulinTracker DeepRed (Thermo Scientific), 1X ActinTracker (CellMask

Actin Tracker Stain, Invitrogen). The MitoView 405 dye was added to the cells 15 min

prior imaging and kept throughout. All the other dyes were washed off prior to imaging,

after 30 minutes of incubation. Live cell imaging was performed using a Nikon A1RHD

inverted confocal microscope, equipped with a photo-stimulation unit, GaAsP and multi-

alkali PMT detectors, 405-nm, 488-nm, 561-nm and 640-nm laser lines and a Nikon

60X/1.4NA Plan Apochromatic oil immersion objective, unless specified otherwise.

Experiments were carried out at a constant 25°C temperature in an Okolab incubation

system and time series were digitally captured with the Nikon NIS-Elements software at

1 frame per seconds (fps), unless specified otherwise.

Sustained Split-Miro photocleavage in S2R+ cells was achieved with a 0.3% (0.017 mW)

488-nm laser, scan zoom of 4, at a framerate of 1 fps for the entire duration of the

experiment. For dual colour imaging during sustained Split-Miro photocleavage, a frame

every 1 min was acquired with either the 405-nm, 561 or 640-nm laser line. Transient

whole-cell photocleavage of Split-Miro with untagged C-terminus was achieved with a

stimulation step of 570 ms using a 4% 488-mn laser at a scan speed of 4 fps and 2

scanning iterations. In these experiments, a Nikon 40X/1.15NA Apochromatic water

immersion objective with a scan zoom of 4 was used. The stimulation area was set to

492.7 μm^2, sufficient to cover the whole cell area. Time series before and after stimulation

were acquired at 0.5 fps. To achieve transient photocleavage of double-tagged Split-Miro,

cells were imaged with both a 488-nm (0.8%, scan time 0.6 s/frame) and a 561-mn (1%,

scan time 0.6 s/frame) laser for 8 s with a frame rate of 0.5 fps, scan zoom of 4. Recovery

was monitored by imaging with a 561-nm laser with a frame rate of 0.5 s for up to 3 min.

To combine optogenetic stimulation with three-colour imaging and fast acquisition (as in

Fig 3), the same settings were maintained, and the 640-nm laser power was set to 0.8%.

Imaging the neurons of the adult Drosophila wing was performed as reported in Vagnoni

et al. (37,38). For rapid blue-light photostimulation in this tissue, a 5% (0.184 mW) 488-

nm laser, scan zoom of 4, was used with a scan speed of 4 fps and 4 scanning iterations.

The total duration of the stimulation step was 1.14 s and the stimulation area was set to

cover a 46.23 μm^2 region of the neuronal axons.

Mitochondrial trafficking in adult wing neurons was recorded with a frame rate of 0.5

frames/s for 5 minutes and quantified from the bundle of marginal axons along the L1

vein, essentially as we described previously (37,38). The *in vivo* trafficking experiments

were performed using a Nikon spinning disk system with a CSU-X1 scanning head

(Yokogawa) and an inverted microscope stand (Eclipse Ti-E (Nikon)) equipped with a 60

x CFI Apo oil immersion objective (1.4 NA) and an EM-CCD camera (Du 897 iXon Ultra

(Andor)). Imaging of mitochondrial motility and concomitant Split-Miro photocleavage

were achieved using a 15% 488-nm laser light and 400-ms light exposure.

Analysis of mitochondrial and peroxisomal transport in S2R+ cells.

Organelle motility was quantified in 2 – 6 processes for each cell. Only processes which

remained in the same focal plane throughout acquisition and had no or minimal crossing

with other processes, were selected. Each process was straightened using the

'Straighten' function in Fiji/ImageJ and kymographs produced and analysed with the

'Velocity Measurement Tool' (https://dev.mri.cnrs.fr/projects/imagej-

macros/wiki/Velocity_Measurement_Tool) in Fiji/ImageJ.

'Run' was defined as a continuous movement which lasts for at least 3 seconds without

any change in direction or velocity. Organelles were defined as 'motile' if engaging in at

least one run ≥ 2 µm. A 'pause' was defined as a period of at least 3 seconds during

which the mitochondrion does not change its position. The duty cycle describes the motile

behaviour of the organelles as percentage of time spent on long runs (≥ 2 µm), short runs

(< 2 µm) and pausing, or in the anterograde and retrograde direction.

Analysis of mitochondrial morphology and mitochondria-ER contacts.

Mitochondrial morphology in the cell soma was analysed from cells showing a clearly

interconnected network before photocleavage, using Fiji/ImageJ and a pipeline adapted

from Chaudhry *et al.* (77). Briefly, the 'Adaptive Threshold' command was used to create

binary images which were individually checked for accuracy in marking the mitochondrial

network. The 'Analyse particles' command was used to measure mitochondrial size, with

a cut-off set at 0.09 μm^2. A skeleton of the mitochondrial network within each cell was

obtained and analysed with the 'Skeletonize' and 'Analyze skeleton (2D/3D)' functions,

respectively. The same analysis pipeline was used to quantifying mitochondria-ER

contacts by creating a mask of the SPLICS signal and counting the GFP puncta

throughout the cell.

Mitochondrial mass was defined as the total area covered by mitochondria within the cell

and normalised to the first time point. The mitochondrial aspect ratio (AR) was measured

in single mitochondria smaller than 6 μm^2 and was defined as the ratio between the

longest and the shortest axis of the organelle. For the analysis of the mitochondrial

network branching, only branches emanating from the main mitochondrial skeleton were

considered.

Analysis of Split-Miro kinetics.

The kinetics of the cytoplasmic release of mCherry-tagged Split-Miro N-terminus after

photocleavage and the reconstitution at the mitochondria was analysed by measuring the

cytosolic fluorescence intensity of the mCherry before and after photocleavage. For cell

culture experiments with single-tagged Split-Miro, images were analysed with the General

Analysis tool of the Nikon NIS-Elements Software by defining an ROI covering the cellular

cytosol (with no mitochondria) and measuring the fluorescence intensity at each time

point. For cell culture experiments with double-tagged Split-Miro and for neuronal *in vivo*

experiments, images were analysed with Fiji/ImageJ by measuring the fluorescence

intensity of a cytosolic ROI devoid of mitochondria (area is 0.451-μm^2 and 0.126-μm^2,

respectively). For each cell or fly wing, the fluorescence intensity over time was

normalised to a scale from 0 to 1, where 0 corresponds to the minimum value measured

before Split-Miro N-terminus release and 1 to the maximum value measured after

779 photocleavage. Mitochondria/cytosolic ratio of both mCherry-tagged Split-MiroN and

780 EGFP-tagged Split-MiroC in S2R+ cells were calculated by measuring the fluorescence

781 intensity at different time points from two 0.451-μm^2 ROIs located within the mitochondrial

782 network and within a proximal area of the cytosol devoid of mitochondria. An mCherry

783 ratio of 1 indicates the signal is homogeneous throughout the cytoplasm and thus

784 mCherry-Split-MiroN has reached complete release.

786 ***Live cell calcium imaging in S2R+ cells.***

787 S2R+ cells were imaged with a Nikon Ti-E inverted epifluorescence microscope equipped

788 with a Nikon 60X/1.4NA Plan Apochromatic oil immersion objective, a mercury lamp

789 (Nikon Intensilight C-HGFI) for illumination and a Dual Andor Neo sCMOS camera for

790 detection. Cells were plated on a Concanavalin A-coated 18-mm coverslip and imaged in

791 a Ludin imaging chamber (Type 1, Life Imaging Services) in S2 media. Cells were imaged

792 at 1 fps for 3 minutes under blue-light which was sufficient to induce Split-Miro-dependent

793 mitochondrial network retraction, prior to ionomycin stimulation. A peristaltic pump system

794 (Ismatec) was used to perfuse the imaging chamber with 2.5 μM ionomycin

795 (ThermoFisher Scientific) until all the media in the chamber was replaced. Additional 5-

796 minutes imaging at 1 fps was performed to record the dynamic mito-GCaMP6 signal. To

797 verify Split-Miro cleavage, a frame with green light (to capture mCherry-Split-MiroN) was

798 acquired at the beginning and the end of each acquisition period.

799 To quantify mitochondrial calcium, a region of interest (ROI) was drawn to outline single

800 cells and the fluorescence intensity of mito-GCaMP6f signal measured using the Nikon

801 NIS-Elements software. Response curves were aligned at the base of the response peak

802 and, for each cell, the fluorescence intensity at every time point was normalised to the

803 value of the first frame. Only cells that responded at least 2-fold to ionomycin in the first

100 s were included in the quantification. mito-GCaMP6f 'peak' was defined as the

maximum fold-change value reached during imaging. The time to reach the peak was

calculated as the Δt between the 'peak' timepoint and the time of first response to

stimulation.

Live cell calcium imaging in adult neurons.

Calcium transients in the wing marginal neurons were acquired using the same spinning

disk imaging system described to capture mitochondrial trafficking. Image series were

recorded for 5 minutes using a 10% 488-nm laser light with an acquisition time of 2

frames/s and exposure time of 200 ms. Image series from 1-2 fields of view were acquired

from the same areas of each wing (i.e., mid-margin to the L1–L2 vein intersection). To

quantify the GCaMP fluorescence during spontaneous calcium transients, an ROI was

drawn around each responding neuronal soma and axons. After background subtraction

in Fiji/ImageJ (rolling ball radius = 50 px), the ROI mean intensity over time was calculated

using the Plot Z-axis profile command. GCaMP fluorescence peak and time to reach the

peak were calculated as done for calcium imaging in S2R+ cells.

Behavioural assay.

Flies were anesthetised with CO_2 on the day of eclosion, transferred to new food and

aged for either 2 or 8 days prior to the assay. On the day of the assay, flies were briefly

anesthetised on ice and individually loaded into 65-mm glass-tubes on custom-made

platforms. Fly activity was recorded in an optogenetic enclosure (BFK Lab Ltd) equipped

with LEDs for 444-nm blue light stimulation and a webcam for video recording. The

position of each fly was automatically tracked and the quantification of the fly locomotor

activity performed using the *Drosophila* ARousal Tracking (DART) system (45,46). Flies

were left to acclimatise for 20 min before starting the experiment. Fly activity was recorded

continuously for 30 minutes before the blue LED light was switched on for 2 hours (power

at the platform: 5 mW/cm^2). After the blue light was switched off, fly movements were

recorded for an additional hour, for a combined experimental time of 3 hours, 30 minutes.

Flies were defined as 'active' if displaying bouts of activity of at least 2 mm/s. The active

average speed refers to the speed of fly movement when active. The action initiation is

defined as the number of times a bout of activity is started per second. The % activity

represents the percentage of time the fly spends moving.

Statistical analysis and image preparation.

Data were analysed with Microsoft Excel and GraphPad Prism 9. Statistical tests and

number of replicates are reported in the figure legends. Images were assembled using

Fiji/ImageJ. For presentation purposes, 'Subtract Background' (rolling ball radius = 20

pixels), and either the 'Despeckle' or 'Smooth' filters were used to reduce salt and pepper

noise. All images in the same experimental series were processed in the same manner.

ACKNOWLEDGMENTS

We thank Tito Calì, Manolis Fanto, Gohta Goshima, Marc-David Ruepp, Tom Schwarz,

Zu-Hang Sheng, Konrad Zinsmaier and Alex Whitworth for sharing reagents, the Fly

Facility of the Department of Genetics, University of Cambridge for help with *Drosophila*

embryo injections, the BFK Lab Ltd for assistance with the behavioural assays, the Wohl

Cellular Imaging Centre at King's College London for help with light microscopy, and the

Bloomington Drosophila Stock Center for fly stocks. We thank members of the Vagnoni

REFERENCE LIST

1.   Eberhardt EL, Ludlam A V., Tan Z, Cianfrocco MA. Miro: A molecular switch at the

center of mitochondrial regulation. Protein Sci. 2020;29(6):1269–1284.

2.   Reis K, Fransson Å, Aspenström P. The Miro GTPases: At the heart of the

mitochondrial transport machinery. FEBS Lett. 2009;583(9):1391–1398.

3.   Lee KS, Lu B. The myriad roles of Miro in the nervous system: axonal transport of

mitochondria and beyond. Front Cell Neurosci. 2014;8:330.

4.   Guo X, Macleod GT, Wellington A, Hu F, Panchumarthi S, Schoenfield M, et al. The

GTPase dMiro is required for axonal transport of mitochondria to Drosophila

synapses. Neuron. 2005;47(3):379–393.

5.   Nguyen TT, Oh SS, Weaver D, Lewandowska A, Maxfield D, Schuler MH, et al.

Loss of Miro1-directed mitochondrial movement results in a novel murine model for

neuron disease. Proc Natl Acad Sci U S A. 2014;111(35): E3631-3640.

6.   López-Doménech G, Covill-Cooke C, Ivankovic D, Halff EF, Sheehan DF, Norkett

R, et al. Miro proteins coordinate microtubule- and actin-dependent mitochondrial

transport and distribution. EMBO J. 2018;37(3):321–336.

7.   Russo GJ, Louie K, Wellington A, Macleod GT, Hu F, Panchumarthi S, et al.

Drosophila Miro is required for both anterograde and retrograde axonal

mitochondrial transport. J Neurosci. 2009;29(17):5443–5455.

8.   MacAskill AF, Rinholm JE, Twelvetrees AE, Arancibia-Carcamo IL, Muir J,

Fransson A, et al. Miro1 Is a Calcium Sensor for Glutamate Receptor-Dependent

Localization of Mitochondria at Synapses. Neuron. 2009;61(4):541–555.

9.   Wang X, Schwarz TL. The Mechanism of Ca^{2+}-Dependent Regulation of Kinesin-

Mediated Mitochondrial Motility. Cell. 2009;136(1):163–174.

10.   Vaccaro V, Devine MJ, Higgs NF, Kittler JT. Miro1-dependent mitochondrial

positioning drives the rescaling of presynaptic Ca^{2+} signals during homeostatic

plasticity. EMBO Rep. 2017;18(2):231–240.

11.   Li L, Conradson DM, Bharat V, Kim MJ, Hsieh CH, Minhas PS, et al. A mitochondrial

membrane-bridging machinery mediates signal transduction of intramitochondrial

oxidation. Nat Metab. 2021;3(9):1242–1258.

12.   Lee S, Lee KS, Huh S, Liu S, Lee DY, Hong SH, et al. Polo Kinase Phosphorylates

Miro to Control ER-Mitochondria Contact Sites and Mitochondrial Ca^{2+}

Homeostasis in Neural Stem Cell Development. Dev Cell. 2016;37(2):174–189.

13.   Frederick RL, McCaffery JM, Cunningham KW, Okamoto K, Shaw JM. Yeast Miro

GTPase, Gem1p, regulates mitochondrial morphology via a novel pathway. J Cell

Biol. 2004;167(1):87–98.

14.   Wang H, Vilela M, Winkler A, Tarnawski M, Schlichting I, Yumerefendi H, et al.

LOVTRAP: an optogenetic system for photoinduced protein dissociation. Nat

Methods. 2016;13(9):755–758.

15.   Van Haren J, Charafeddine RA, Ettinger A, Wang H, Hahn KM, Wittmann T. Local

control of intracellular microtubule dynamics by EB1 photodissociation. Nat Cell

Biol. 2018;20(3):252–261.

16.   Dema A, van Haren J, Wittmann T. Optogenetic EB1 inactivation shortens

metaphase spindles by disrupting cortical force-producing interactions with astral

microtubules. Curr Biol. 2022;32(5):1197-1205.e4.

17.   van Haren J, Adachi LS, Wittmann T. Optogenetic Control of Microtubule

Dynamics. Methods Mol Biol. 2020;2101:211–234.

18.   Strickland D, Lin Y, Wagner E, Hope CM, Zayner J, Antoniou C, et al. TULIPs:
Tunable, light-controlled interacting protein tags for cell biology. Nat Methods.
2012;9(4):379–384.

19.   Ally S, Larson AG, Barlan K, Rice SE, Gelfand VI. Opposite-polarity motors activate
one another to trigger cargo transport in live cells. J Cell Biol. 2009 Dec
28;187(7):1071–1082.

20.   Vagnoni A, Hoffmann PC, Bullock SL. Reducing Lissencephaly-1 levels augments
mitochondrial transport and has a protective effect in adult Drosophila neurons. J
Cell Sci. 2016;129(1):178–190.

21.   Jolly AL, Luan CH, Dusel BE, Dunne SF, Winding M, Dixit VJ, et al. A Genome-
wide RNAi Screen for Microtubule Bundle Formation and Lysosome Motility
Regulation in Drosophila S2 Cells. Cell Rep. 2016;14(3):611–620.

22.   van Spronsen M, Mikhaylova M, Lipka J, Schlager MA, van den Heuvel DJ, Kuijpers
M, et al. TRAK/Milton motor-adaptor proteins steer mitochondrial trafficking to
axons and dendrites. Neuron. 2013;77(3):485–502.

23.   Brickley K, Stephenson FA. Trafficking kinesin protein (TRAK)-mediated transport
of mitochondria in axons of hippocampal neurons. J Biol Chem.
2011;286(20):18079–18092.

24.   Glater EE, Megeath LJ, Stowers RS, Schwarz TL. Axonal transport of mitochondria
requires milton to recruit kinesin heavy chain and is light chain independent. J Cell
Biol. 2006;173(4):545–557.

25.   Fransson Å, Ruusala A, Aspenström P. The atypical Rho GTPases Miro-1 and
Miro-2 have essential roles in mitochondrial trafficking. Biochem Biophys Res
Commun. 2006;344(2):500–510.

26. Liu S, Sawada T, Lee S, Yu W, Silverio G, Alapatt P, et al. Parkinson's disease-
associated kinase PINK1 regulates Miro protein level and axonal transport of
mitochondria. PLoS Genet. 2012;8(3):e1002537.

27. Nemani N, Carvalho E, Tomar D, Dong Z, Ketschek A, Breves SL, et al. MIRO-1
Determines Mitochondrial Shape Transition upon GPCR Activation and Ca^{2+}
Stress. Cell Rep. 2018;23(4):1005–1019.

28. Chung JYM, Steen JA, Schwarz TL. Phosphorylation-Induced Motor Shedding Is
Required at Mitosis for Proper Distribution and Passive Inheritance of Mitochondria.
Cell Rep. 2016;16(8):2142–2155.

29. Kang J-S, Tian J-H, Pan P-Y, Zald P, Li C, Deng C, et al. Docking of axonal
mitochondria by syntaphilin controls their mobility and affects short-term facilitation.
Cell. 2008;132(1):137–148.

30. Lee K-S, Huh S, Lee S, Wu Z, Kim A-K, Kang H-Y, et al. Altered ER-mitochondria
contact impacts mitochondria calcium homeostasis and contributes to
neurodegeneration in vivo in disease models. Proc Natl Acad Sci U S A.
2018;115(38):E8844–8853.

31. Niescier RF, Hong K, Park D, Min KT. MCU Interacts with Miro1 to Modulate
Mitochondrial Functions in Neurons. J Neurosci. 2018;38(20):4666–4677.

32. Modi S, López-Doménech G, Halff EF, Covill-Cooke C, Ivankovic D, Melandri D, et
al. Miro clusters regulate ER-mitochondria contact sites and link cristae
organization to the mitochondrial transport machinery. Nat Commun.
2019;10(1):4399.

33. Guillén-Samander A, Leonzino M, Hanna MG, Tang N, Shen H, De Camilli P.
VPS13D bridges the ER to mitochondria and peroxisomes via Miro. J Cell Biol.
2021;220(5):e202010004.

34.   Hewitt VL, Miller-Fleming L, Andreazza S, Mattedi F, Prudent J, Polleux F, et al.
954   Decreasing pdzd8-mediated mitochondrial-ER contacts in neurons improves
955   fitness by increasing mitophagy. Life Sci Alliance. 2022; 13;5(11):e202201531.

956   35.   Cieri D, Vicario M, Giacomello M, Vallese F, Filadi R, Wagner T, et al. SPLICS: a
957   split green fluorescent protein-based contact site sensor for narrow and wide
958   heterotypic organelle juxtaposition. Cell Death Differ. 2018; 25(6):1131-1145.

959   36.   Granat L, Ranson D, Hamer E, Chakrabarty RP, Mattedi F, Fort-Aznar L, et al.
960   Yeast NDI1 reconfigures neuronal metabolism and prevents the unfolded protein
961   response in mitochondrial complex I deficiency. bioRxiv. 2022; doi:
962   10.1101/2022.09.22.509020v2.

963   37.   Vagnoni A, Bullock SL. A simple method for imaging axonal transport in aging
964   neurons using the adult Drosophila wing. Nat Protoc. 2016;11:1711–1723.

965   38.   Vagnoni A, Bullock SL. A cAMP/PKA/Kinesin-1 Axis Promotes the Axonal
966   Transport of Mitochondria in Aging Drosophila Neurons. Curr Biol. 2018;28:1265–
967   1272.

968   39.   Kwon SK, Sando R, Lewis TL, Hirabayashi Y, Maximov A, Polleux F. LKB1
969   Regulates Mitochondria-Dependent Presynaptic Calcium Clearance and
970   Neurotransmitter Release Properties at Excitatory Synapses along Cortical Axons.
971   PLOS Biol. 2016;14(7):e1002516.

972   40.   Sun T, Qiao H, Pan PY, Chen Y, Sheng ZH. Motile axonal mitochondria contribute
973   to the variability of presynaptic strength. Cell Rep. 2013 Aug 15;4(3):413–419.

974   41.   Aery Jones EA, Giocomo LM. Neural ensembles in navigation: From single cells to
975   population codes. Curr Opin Neurobiol. 2023;78:102665.

976   42.   Langdon C, Genkin M, Engel TA. A unifying perspective on neural manifolds and
977   circuits for cognition. Nat Rev Neurosci 2023;13;1–15.

43. Panchal K, Tiwari AK. Miro, a Rho GTPase genetically interacts with Alzheimer's
disease-associated genes (Tau, Aβ 42 and Appl) in Drosophila melanogaster. Biol
Open. 2020;9(9):bio049569.

44. Shaltouki A, Hsieh CH, Kim MJ, Wang X. Alpha-synuclein delays mitophagy and
targeting Miro rescues neuron loss in Parkinson's models. Acta Neuropathol.
2018;136(4):607–620.

45. Faville R, Kottler B, Goodhill GJ, Shaw PJ, van Swinderen B. How deeply does
your mutant sleep? Probing arousal to better understand sleep defects in
Drosophila. Sci Rep. 2015;5:8454.

46. Kottler B, Faville R, Bridi JC, Hirth F. Inverse Control of Turning Behavior by
Dopamine D1 Receptor Signaling in Columnar and Ring Neurons of the Central
Complex in Drosophila. Curr Biol. 2019;29(4):567-577.e6.

47. Fenton AR, Jongens TA, Holzbaur ELF. Mitochondrial adaptor TRAK2 activates
and functionally links opposing kinesin and dynein motors. Nat Commun.
2021;12(1):4578.

48. Henrichs V, Grycova L, Barinka C, Nahacka Z, Neuzil J, Diez S, et al. Mitochondria-
adaptor TRAK1 promotes kinesin-1 driven transport in crowded environments. Nat
Commun. 2020;11(1):3123.

49. Canty JT, Hensley A, Yildiz A. TRAK adaptors coordinate the recruitment and
activation of dynein and kinesin to control mitochondrial transport. bioRxiv. 2021;
doi:10.1101/2021.07.30.454553v1.

50. Zhao Y, Song E, Wang W, Hsieh CH, Wang X, Feng W, et al. Metaxins are core
components of mitochondrial transport adaptor complexes. Nat Commun.
2021;12(1):83.

51. Mahecic D, Carlini L, Kleele T, Colom A, Goujon A, Matile S, et al. Mitochondrial

membrane tension governs fission. Cell Rep. 2021;35(2):108947.

52.   Helle SCJ, Feng Q, Aebersold MJ, Hirt L, Grüter RR, Vahid A, et al. Mechanical

force induces mitochondrial fission. Elife. 2017;6:e30292.

53.   Song Y, Huang P, Liu X, Zhao Z, Wang Y, Cui B, et al. Light-inducible deformation

of mitochondria in live cells. Cell Chem Biol. 2022;29(1):109-119.e3.

54.   König T, Nolte H, Aaltonen MJ, Tatsuta T, Krols M, Stroh T, et al. MIROs and DRP1

drive mitochondrial-derived vesicle biogenesis and promote quality control. Nat Cell

Biol 2021;23(12):1271–1286.

55.   Qin J, Guo Y, Xue B, Shi P, Chen Y, Su QP, et al. ER-mitochondria contacts

promote mtDNA nucleoids active transportation via mitochondrial dynamic

tubulation. Nat Commun. 2020;11(1):4471.

56.   Floyd C, Levine H, Jarzynski C, Papoian GA. Understanding cytoskeletal

avalanches using mechanical stability analysis. Proc Natl Acad Sci U S A.

2021;118(41):e2110239118.

57.   Oeding SJ, Majstrowicz K, Hu XP, Schwarz V, Freitag A, Honnert U, et al.

Identification of Miro1 and Miro2 as mitochondrial receptors for myosin XIX. J Cell

Sci. 2018;131(17):jcs219469.

58.   Bocanegra JL, Fujita BM, Melton NR, Cowan JM, Schinski EL, Tamir TY, et al. The

MyMOMA domain of MYO19 encodes for distinct Miro-dependent and Miro-

independent mechanisms of interaction with mitochondrial membranes.

Cytoskeleton (Hoboken). 2020;77(3–4):149–166.

59.   Sato O, Sakai T, Choo YY, Ikebe R, Watanabe TM, Ikebe M. Mitochondria-

associated myosin 19 processively transports mitochondria on actin tracks in living

cells. J Biol Chem. 2022;298(5):101883.

60.   Chang KT, Niescier RF, Min K-T. Mitochondrial matrix Ca^{2+} as an intrinsic signal

regulating mitochondrial motility in axons. Proc Natl Acad Sci U S A.
2011;108(37):15456–15461.

61.   van Bergeijk P, Adrian M, Hoogenraad CC, Kapitein LC. Optogenetic control of
organelle transport and positioning. Nature. 2015;518(7537):111–114.

62.   Ballister ER, Ayloo S, Chenoweth DM, Lampson MA, Holzbaur ELF. Optogenetic
control of organelle transport using a photocaged chemical inducer of dimerization.
Curr Biol. 2015;25(10):R407–408.

63.   Gutnick A, Banghart MR, West ER, Schwarz TL. The light-sensitive dimerizer
zapalog reveals distinct modes of immobilization for axonal mitochondria. Nat Cell
Biol. 2019; 21:768-777.

64.   Harterink M, Van Bergeijk P, Allier C, De Haan B, Van Den Heuvel S, Hoogenraad
CC, et al. Light-controlled intracellular transport in Caenorhabditis elegans. Curr
Biol. 2016;26(4):R153–154.

65.   Yesbolatova A, Saito Y, Kitamoto N, Makino-Itou H, Ajima R, Nakano R, et al. The
auxin-inducible degron 2 technology provides sharp degradation control in yeast,
mammalian cells, and mice. Nat Commun. 2020;11(1):1–13.

66.   Clift D, McEwan WA, Labzin LI, Konieczny V, Mogessie B, James LC, et al. A
Method for the Acute and Rapid Degradation of Endogenous Proteins. Cell.
2017;171(7):1692-1706.e18.

67.   Deubner J, Coulon P, Diester I. Optogenetic approaches to study the mammalian
brain. Curr Opin Struct Biol. 2019 Aug 1;57:157–163.

68.   Murakoshi H, Shin ME, Parra-Bueno P, Szatmari EM, Shibata ACE, Yasuda R.
Kinetics of endogenous CaMKII required for synaptic plasticity revealed by
optogenetic kinase inhibitor. Neuron. 2017;94(1):37-47.e5.

69.   Zullo JM, Drake D, Aron L, O'Hern P, Dhamne SC, Davidsohn N, et al. Regulation

of lifespan by neural excitation and REST. Nature 2019;574(7778):359–364.

70. Simkin D, Hattori S, Ybarra N, Musia TF, Buss EW, Richter H, et al. Aging-Related

Hyperexcitability in CA3 Pyramidal Neurons Is Mediated by Enhanced A-Type K+

Channel Function and Expression. J Neurosci. 2015;35(38):13206-13218.

71. Haberman RP, Koh MT, Gallagher M. Heightened cortical excitability in aged

rodents with memory impairment. Neurobiol Aging. 2017;54:144–151.

72. Wong CO, Karagas NE, Jung J, Wang Q, Rousseau MA, Chao Y, et al. Regulation

of longevity by depolarization-induced activation of PLC-β-IP3R signaling in

neurons. Proc Natl Acad Sci U S A. 2021;118(16):e2004253118.

73. Raad H, Ferveur J-F, Ledger N, Capovilla M, Robichon A. Functional gustatory role

of chemoreceptors in Drosophila wings. Cell Rep. 2016;15(7):1442–1454.

74. Zhang N, Simpson JH. A pair of commissural command neurons induces

Drosophila wing grooming. iScience. 2022;25(2):103792.

75. Babic M, Russo GJ, Wellington AJ, Sangston RM, Gonzalez M, Zinsmaier KE.

Miro's N-terminal GTPase domain is required for transport of mitochondria into

axons and dendrites. J Neurosci. 2015;35(14):5754–5771.

76. Stowers RS, Megeath LJ, Górska-Andrzejak J, Meinertzhagen IA, Schwarz TL.

Axonal transport of mitochondria to synapses depends on milton, a novel

Drosophila protein. Neuron. 2002;36(6):1063–1077.

77. Chaudhry A, Shi R, Luciani DS. A pipeline for multidimensional confocal analysis

of mitochondrial morphology, function, and dynamics in pancreatic β-cells. Am J

Physiol Endocrinol Metab. 2020;318(2):E87–101.

1075

1076 **SUPPORTING INFORMATION**

1077

1078 **Data S1.** Excel spreadsheet containing the numerical data and details of statistical

1079 analysis for Figure panels 1D, 1E, 1F-G, 2C-D, 2F, 2G, 2H, 3B-D, 3F-G, 4B, 4C, 4D-E,

1080 4G-H, 5C-D, 5E-F, 6C, 6D-F, 7A, 7C, 7D-E, 7F, 7G-H, 7I, 7J-K, S1C-D, S1F-G, S2B-C,

1081 S2G, S2H, S2I, S2J, S2K, S3A, S3C, S3D, S3F, S3G,I, S4B, S5C-D, S5E-F, S5G-H.

1082

S1 Fig. Split-Miro reconstitution and kinetics of Split-Miro harbouring wild-type and

mutant LOV2 proteins. A) EGFP-Split-Miro-mCherry is reconstituted and localises at

the mitochondria in the absence of sustained blue-light irradiation. S2R+ cells were co-

transfected with mCherry-tagged Split-MiroC (magenta) and EGFP-tagged Split-MiroN

(green), mitochondria are stained with MitoTracker Deep Red (MTDR, cyan). Note the

EGFP/mCherry tags are appended at different termini compared to the mCherry-Split-

Miro-EGFP shown in Fig 1. **B, E)** Localisation of mCherry-Split-MiroN before and after a

570-ms pulse of blue light. S2R+ cells are co-transfected with untagged Split-MiroC and

mCherry-Split-MiroN containing the T406A, T407A mutations in the N-terminus of LOV2

(LAAA, in B) or mCherry-Split-MiroN containing the wild-type sequence (LATT, in E).

Before irradiation, Split-Miro is reconstituted at the mitochondria, indicated by the co-

localisation of mCherry-Split-MiroN (grey) and the MitoTracker Deep Red (MTDR)

staining. Immediately after irradiation (18 s), Split-MiroN is released into the cytoplasm

(indicated by a more homogenous grey colour), and it fully reconstitutes within 2 minutes.

C-D) Quantification of mCherry-Split-MiroN half-time release (C) and recovery (D) after

photocleavage, relative to B. In (D), left panel shows levels of cytosolic Split-Miro N-

terminus quantified after the maximum release is reached; right panel: recovery half-time.

Solid line in (D) is exponential curve fit. In **B** and **E**, the cartoon depicts photocleavage

and reconstitution of mCherry-tagged Split-Miro. s, seconds. Scale bar: 10 μm. **F-G)**

Comparison of mCherry-Split-MiroN release (F) and recovery (G) half-time after

photocleavage shows no significant difference between the two LOV2 variants (unpaired

Student's t-test). The LATT (wild-type) to LAAA mutation in the N-terminus of LOV2 has

been reported to have a stabilising effect on the Jα helix in cultured cells at 37°C (17,18).

We did not find any noticeable difference in the steady-state reconstitution efficiency of

the two variants in S2R+ cells cultured at 25°C. Data are shown as mean ± SEM. Circles,

number of cells, from two independent experiments. Scale bar: 10 μm (A) and 5 μm (B,

E).

1110

S2 Fig. Insight into the regulation of mitochondrial motility in the processes of

S2R+ cells by endogenous Miro and Split-Miro. A) Representative western blots of

Miro from total lysates of control and Miro RNAi-treated S2R+ cells. **B)** Duty cycle analysis

describes the average time mitochondria spend moving anterogradely, retrogradely or

pausing. For each parameter, all mitochondrial values from each cell were averaged and

compared between control and Miro dsRNA condition using a multiple Student's t-tests.

Number of mitochondria analysed are in brackets from 29 (Ctrl dsRNA) and 36 (Miro

dsRNA) cells, respectively, from two independent experiments. **C)** Run velocities of short

and long runs in control dsRNA-treated S2R+ cells showing that Miro-dependent long

runs are significantly more processive than the short, Miro-independent runs (Fig 2C).

Number of runs analysed are in brackets, from two independent experiments. Mann-

Whitney test. **D)** Split-Miro interacts with Milton in S2R+ cells. Cells were transfected with

Split-Miro or Control (as shown in E) and the total cell lysates immunoprecipitated using

GFP-beads to pull down EGFP-tagged Split-Miro C-terminus. Immunoprecipitates were

blotted and probed with anti-GFP antibody (to detect Split-Miro C-terminus), anti-Miro

antibody (to detect Split-Miro N-terminus) and an anti-Milton antibody. Inputs are total

lysates (25 μg protein). **E)** Cartoon showing Split-Miro and Control constructs with the

GFP and Miro antibodies used for immunoprecipitation and western blotting in (D). **F)**

Representative kymographs of mitochondrial transport in the processes of S2R+ cells

transfected with mCherry-tagged Zdk1-MiroC (Control), mCherry-Miro (wt-Miro) and

mCherry-Split-Miro (Split-Miro). Scale bars: 2 μm (distance) and 5 s (time). **G)** Distribution

of mitochondria run lengths in the processes of S2R+ cells, transfected with control, wt-

Miro and Split-Miro as shown in F. N = number of mitochondrial runs. One-way ANOVA

with Tukey's post-hoc test. **H)** Duty cycle analysis describing the average time

mitochondria spend moving anterogradely, retrogradely or pausing in control, wt-Miro and

Split-Miro-transfected cells, relative to F. For each parameter, all mitochondrial values per

cell were averaged and compared by one-way ANOVA followed by Tukey's post-hoc test.

Number of mitochondria are in brackets from 16 (control), 15 (wt-Miro), and 15 (Split-

Miro) cells, respectively, from three independent experiments. **I)** S2R+ cells transfected

with wt-Miro and Split-Miro were imaged for 1 minute with a 561-nm laser, to capture the

mCherry signal, followed by 1-minute imaging with 488-nm blue light, to capture the

EGFP signal after Split-Miro photocleavage (relative to Fig 2E). Number of mitochondria

are in brackets from 11 (wt-Miro) and 17 (Split-Miro) cells, from three independent

experiments. Data are shown as mean ± SEM. Kolmogorov-Smirnov test showed no

statistical difference between groups. **J)** Distribution of mitochondrial run lengths in the

anterograde direction in the processes of S2R+ cells, during the 1st and 7th minute of

time-lapse imaging with blue light in cells transfected with wt-Miro or Split-Miro. N =

number of runs. Mann-Whitney test showed no statistical difference between groups. **K)**

Run velocities for long processive anterograde and retrograde runs in S2R+ cells

transfected with Split-Miro and Miro dsRNA (which targets endogenous Miro) and imaged

by time-lapse for seven minutes under blue light. Circles, number of runs, from two

independent experiments. Mann-Whitney test shows no difference between first and

seventh minutes of imaging, with the velocities remaining high compared to non-transfected condition (e.g., Fig 2H, S2C Fig). This result shows that the velocities of the processive mitochondria, augmented as a consequence of Split-Miro overexpression, remain elevated even after reduction of endogenous Miro, suggesting Miro is not necessary for maintaining mitochondrial velocities. * $p < 0.05$, ** $p < 0.01$, **** $p < 0.0001$.

S3 Fig. Extended characterisation of the effects of Split-Miro photocleavage and SNPH overexpression in S2R+ cells. A) Quantification of the total area covered by the mitochondria within the cell. Each measurement was normalised to the average group value (wt-Miro, Split-Miro) at timepoint 0. Comparison across timepoints was performed by repeated-measures one-way ANOVA followed by Tukey's post-hoc test. Data are reported as mean ± SEM. Circle, number of cells, from three independent experiments. **B)** S2 cells are transfected with EGFP-tagged Split-Miro N-terminus (green) and stained with MitoTracker DeepRed (MTDR, cyan). Middle panel: white and magenta stars indicate untransfected and transfected cells, respectively. Scale bar: 10 µm. **C)** Duty cycle analysis describes the average time peroxisomes spend on long runs, short runs or pausing. For each parameter, all peroxisomal values from each cell were averaged and compared between time points. Statistical significance was evaluated by multiple Mann-Whitney tests. **D)** Bar chart shows the average peroxisomal content at minute 1 and 7 of time-lapse imaging with blue light. Circles represent the number of peroxisomes within each process. Statistical significance was evaluated by Wilcoxon test. In C-D), number of processes and cells: wt-Miro = 24, 11, Split- Miro = 39, 16, from three independent experiments. There is no significant difference in the motility and number of peroxisomes in each process between timepoints. **E)** Representative images showing S2R+ cells transfected with Split-Miro and imaged by time-lapse with blue light for seven minutes.

Exposure to blue-light (to induce Split-Miro photocleavage) leads to altered mitochondrial

morphology and distribution, without noticeable disorganisation of the microtubule

network as detected by the Tubulin Tracker (magenta). Scale bar: 10 µm. Not shown,

mCherry-tagged Split-Miro N-terminus. **F)** Bar chart shows the average mitochondrial

content at minute 1, 3 and 7 of time-lapse imaging with blue light in the processes of

S2R+ transfected with either wt-Miro or Split-Miro. Circles represent the number of

mitochondria within each process at minute 1 and 7. Data were analysed by Friedman

test with Dunn's multiple comparison test. Number of processes: wt-Miro = 39, Split-Miro

= 50 from 11 and 17 cells, respectively, from three independent experiments. **G)** Length

of cellular processes imaged for 7 minutes under time-lapse exposure to 488-nm blue

light to achieve Split-Miro photocleavage. Circles, number of the processes. Number of

cells: wt-Miro = 11, Split-Miro = 17, from three independent experiments. Statistical

significance was evaluated by Mann-Whitney test. **H)** Representative kymographs from

processes of cells transfected with an empty vector (Control) or EGFP-SNPH (SNPH).

Mitochondria were stained with MitoTracker Deep Red (MitoTracker). Scale bars: 2 µm

(distance) and 10 s (time). **I)** Percentage of motile mitochondria in cellular processes of

control or SNPH-expressing cells. Number of processes: control = 9, SNPH = 9, from

eight cells, from two independent experiments. Data are shown as mean ± SEM and were

analysed by unpaired Student's t-test. *p < 0.05, **** p < 0.0001.

1197

S4 Fig. Split-Miro photocleavage does not affect mitochondrial membrane potential

in S2R+ cells. A) Representative images of cells co-transfected either with wild-type Miro

(mCherry-wt-Miro, EGFP-mito) or Split-Miro (mCherry-Split-Miro-EGFP). EGFP signal is

used to mark the mitochondria over time; mCherry, not shown. MitoView 405 was used

to monitor the mitochondrial membrane potential during the imaging period. Scale bar: 10

µm. **B)** Ratio of MitoView and EGFP mitochondrial fluorescence intensity at the time

points indicated. Number of cells: wt-Miro = 7, Split-Miro = 7, from two independent

experiments. Data are shown as mean ± SEM. Each group was analysed by repeated-

measures one-way ANOVA followed by Dunnett's post-hoc test, with each time point

compared to the t = 0 min.

1208

S5 Fig. Extended characterisation of Split-Miro *in vivo*.

A) Western blots of lysates from male fly heads of the reported genotypes confirms the

expression of *UAS-wt-Miro* and *UAS-Split-Miro*, in either Miro heterozygous ($miro^{Sd32/+}$)

and null backgrounds ($miro^{Sd32/B682}$), using the Appl-Gal4 driver. *UAS-wt-Miro* is

expressed from attP2, the *UAS-MiroN* and *UAS-MiroC* (to reconstitute Split-Miro) were

expressed from attP40 and attP2, respectively. The higher molecular weight of wt-Miro

and Split-Miro compared to endogenous Miro is consistent with the presence of the

mCherry tag (wt-Miro) and mCherry/LOV2 tags (Split-Miro). Please note the lower

expression of Split-Miro compared to wt-Miro does not significantly affect the proportion

of rescued flies, as shown in Fig 7A. The top membrane was blotted with an anti-Miro

antibody recognising an N-terminal epitope (see also S2E Fig). **B)** Representative images

of axons in the L3 vein of the adult live fly wing. For co-localisation experiments, flies

express mCherry-tagged wt-Miro (attP40) and EGFP-tagged Miro C-terminus (attP2).

Split-Miro flies express the N-terminus and C-terminus halves of Split-Miro from attP40

and attP2, respectively. All constructs were expressed under the control of the Appl-Gal4

driver in a $miro^{+/+}$ background. Arrows highlight examples of co-localised signal. Scale

bar: 10 µm. **C)** Quantification of the kinetics of Split-Miro recovery after photocleavage in

the neurons of the adult fly wing *in vivo*. UAS-mCherry-Split-MiroN and UAS-MiroC were

both expressed from attP40, using the Appl-Gal4 driver. The first time point corresponds

to maximum levels of mCherry-Split-MiroN release. Data are shown as mean ± SEM. Red

solid line, exponential curve fit (n = 6 wings, from 3 flies). **D)** There is no difference in the

half-life recovery of Split-Miro after photobleaching in S2R+ cells (*in vitro*) and in the

neurons of the adult fly wing (*in vivo*). Data are shown as mean ± SEM and were analysed

by Mann-Whitney test. Circles, number of cells (*in vitro*) and wings (*in vivo*) analysed. **E-**

F) Number of motile mitochondria captured in a 50-μm axonal tract in 7d-old wing neurons

expressing wt-Miro (E) or Split-Miro (F). Bar charts show the average mitochondrial

content at each time point. Filled circles represent the number of mitochondria within each

axonal bundle at minute 1 and 5. Data were analysed by paired Student's t-test. Number

of wings analysed: wt-Miro = 5, Split-Miro = 6, from two independent experiments. **G-H)**

Anterograde velocity (G) and run length (H) of axonal mitochondria in wing neurons

expressing wt-Miro or Split-Miro, during the first and fifth minute of blue light exposure,

relative to E-F. Circles represent tracked mitochondria. Data were analysed by Student's

t-test. Due to the overall lower number of bidirectional and retrograde-moving

mitochondria, a meaningful statistical analysis of their velocity and run length is not

possible.

1244

S6 Fig. The role of actin in the Split-Miro-dependent retraction of the mitochondrial

network in S2R+ cells.

A) S2R+ cell transfected with mCherry/EGFP-tagged Split-Miro and not treated with

cytochalasin D (to maintain an intact actin network) were imaged by time-lapse with a

488-nm laser for 7 minutes. The presence of an intact actin network does not prevent

mitochondrial network collapse. Scale bar: 10 µm. Cartoon depicts the reconstitution state

of Split-Miro at 0 and 7 minutes (min) under blue light. **B)** Representative images of S2R+

cells treated with or without cytochalasin D (cytoD) for 4 hours before imaging. Before

imaging, cells were stained with MitoTracker Green (MitoTracker) and ActinTracker

DeepRed (ActinTracker) to visualise the mitochondria and the actin network, respectively.

Note the depolymerisation of the actin network after cytoD treatment, which is required

for process extension in S2R+ cells. Scale bar: 5 µm.

1257

1258   **S1 Table.** Plasmids used in this study.

1259

S2 Table. Primers used in this study. For NEBuilder HiFi DNA Assembly, bold fonts

indicate primer overlap with the amplified gene product, underline indicated overlap with

the adjacent insert or plasmid sequence. For site-directed mutagenesis, bold fonts

indicate mutated nucleotides. For dsRNA template production, underline indicates T7

promoter sequence which precedes a sequence overlapping with either Miro 3'-UTR

(dsRNA) or a non-targeting sequence within the pT2-DsRed-UAS plasmid backbone

(Control).

1267

1268   **S3 Table.** Fly lines used in this study and genotypes analysed in the behavioural assay.

1269

Movie 1. Representative time-lapse movie of S2R+ cells transfected with mCherry-Split-

MiroN (magenta) and EGFP-tagged Split-MiroC (green). mCherry-Split-Miro-EGFP is

reconstituted at the mitochondria and, upon irradiation with 488-nm blue light, mCherry-

Split-MiroN (but not EGFP-tagged Split-MiroC) is released into the cytoplasm and fully

reconstitutes within 3 minutes. Note mCherry-Split-MiroN release is already evident at 2

seconds, followed by complete diffusion throughout the cytoplasm.

1276

1277   **Movie 2.** Representative time-lapse movie of S2R+ cells transfected with mCherry-Split-

1278   MiroN and untagged Split-MiroC, before and after a 570-ms pulse of blue light. Upon

1279   irradiation with 488-nm blue light, mCherry-Split-MiroN (grey) is released from the

1280   mitochondria into the cytoplasm and fully reconstitutes within 120 s.

1281

1282   **Movie 3.** Representative time-lapse movie of S2R+ cells transfected with mCherry-Split-

1283   MiroN and EGFP-tagged Split-MiroC and imaged with the 488-nm laser to capture EGFP-

1284   targeted mitochondria (grey) while photocleaving Split-Miro. Note the strong

1285   mitochondrial morphological changes during the period of blue-light imaging. Not shown,

1286   mCherry-Split-MiroN.

1287

Movie 4. Representative time-lapse movie of S2R+ cells transfected with mCherry-Split-

MiroN and EGFP-tagged Split-MiroC and imaged with the 561-nm laser for 1 minute (to

capture Split-Miro via mCherry, red) before switching to the 488-nm laser for 7 minutes

to capture EGFP-targeted mitochondria (green) while photocleaving Split-Miro. Note the

strong mitochondrial morphological changes and reduced motility during the period of

blue-light imaging.

1294

Movie 5. Representative time-lapse movie of S2R+ cells transfected with mCherry-wt-

Miro and EGFP-tagged Split-MiroC and imaged with the 561-nm laser for 1 minute (to

capture wt-Miro via mCherry, red) before switching to the 488-nm laser for 7 minutes to

capture EGFP-targeted mitochondria (green). Note that, contrarily to Split-Miro

transfected cells (Movie 4), there are no mitochondrial morphological changes or reduced

motility during the period of blue-light imaging.

1301

1302 **Movie 6.** Representative time-lapse movie of mitochondria motility in a 30-μm axonal tract

1303 of adult wing neurons expressing either wt-Miro (top movie) or Split-Miro (bottom movie)

1304 in a *miro$^{+/-}$* background. Mitochondrial motility is shown after 1 and 5 minutes of blue-light

1305 exposure. Each movie corresponds to 30 seconds of real time imaging.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

Appl-GAL4>UAS-wt-Miro, UAS-mito::GFP, Miro^{Sd32}

1st minute

2.1
1.3
1.1
1.2

5th minute

1.1
1.2
3.1
1.3

Retrograde   Bidirectional   Anterograde

B

Appl-GAL4>UAS-Split-Miro, Miro^{Sd32}

1st minute

2.3
1.6
1.4
1.5 1.2
2.1 1.1
2.2
1.3
2.4

5th minute

1.2
1.1
2.1
1.3

Retrograde   Anterograde

C

2 days

motile mitochondria/50 μm

UAS-wt-Miro 1st   UAS-wt-Miro 5th

D

2 days

motile mitochondria/50 μm

***

UAS-Split-Miro 1st   UAS-Split-Miro 5th

E   Anterograde velocity

μm/s

UAS-wt-Miro 1st   UAS-wt-Miro 5th

μm/s

UAS-Split-Miro 1st   UAS-Split-Miro 5th

F   Anterograde run length

μm

UAS-wt-Miro 1st   UAS-wt-Miro 5th

μm

UAS-Split-Miro 1st   UAS-Split-Miro 5th

Figure 6

A *Appl-GAL4>UAS-GCaMP5*

GCaMP   32 s   46 s   75 s

B *Appl-GAL4>UAS-GCaMP5*

GCaMP   454 s   460 s   466 s

C *Appl-GAL4>UAS-GCaMP5*

Normalised GCaMP5 fluorescence (F/F_0)

time (s)

○ *UAS-wt-Miro* (n = 34)
■ *UAS-Split-Miro* (n = 35)

D *UAS-GCaMP5*

Peak fold increase (F/F_0)

****

UAS-wt-Miro   *UAS-Split-Miro*

Appl-GAL4

E *UAS-GCaMP5*

time to peak (s)

***

UAS-wt-Miro   *UAS-Split-Miro*

Appl-GAL4

F *UAS-GCaMP5*

fields of view

□ No response
■ Response

*

UAS-wt-Miro   *UAS-Split-Miro*

Appl-GAL4

Figure 7