Spatially defined multicellular functional units in colorectal cancer revealed from single cell and spatial transcriptomics

Inbal Avraham-Davidi[1,20]*, Simon Mages[1,2,20], Johanna Klughammer[1,2,20], Noa Moriel[3], Shinya Imada[4], Matan Hofree[1,] Evan Murray[5], Jonathan Chen[5,6,7], Karin Pelka[5,6,8,9], Arnav Mehta[5,6,10], Genevieve M. Boland[5,11], Toni Delorey[1], Leah Caplan[1], Danielle Dionne[1], Robert Strasser[2], Jana Lalakova[12], Anezka Niesnerova[12], Hao Xu[12], Morgane Rouault[12], Itay Tirosh[13], Nir Hacohen[5,6], Fei Chen[5,14], Omer Yilmaz[4,15], Jatin Roper[4,16,*], Orit Rozenblatt-Rosen[1,17,*], Mor Nitzan[3,18,19*], Aviv Regev[1,4,17*]

[1]Klarman Cell Observatory, Broad Institute of MIT and Harvard, Cambridge, MA, USA

[2]Gene Center and Department of Biochemistry, Ludwig-Maximilians-Universität München, Munich, Germany

[3]School of Computer Science and Engineering, The Hebrew University of Jerusalem, Jerusalem, Israel

[4]Departmen of Biology at MIT, Koch Institute for Integrative Cancer Research at MIT, Cambridge, MA, USA

[5]Broad Institute of Massachusetts Institute of Technology and Harvard, Cambridge, MA, USA

[6]Massachusetts General Hospital (MGH) Cancer Center, Harvard Medical School (HMS), Boston, MA, USA

[7]Department of Pathology, MGH, Boston, MA, USA

[8]Current address: Gladstone-UCSF Institute of Genomic Immunology, Gladstone Institutes, CA, USA

[9]Current address: Department of Microbiology and Immunology, UCSF, CA, USA

[10]Dana Farber Cancer Institute, Boston, MA, USA

[11]Department of Surgery, MGH, Boston, MA, USA

[12]10xGenomics, Solnavägen 3H, Stockholm, Sweden

[13]Department of Molecular Cell Biology, Weizmann Institute of Science, Rehovot, Israel

[14]Harvard Stem Cell and Regenerative Biology, Cambridge MA, USA

[15]Department of Pathology, Massachusetts General Hospital, Boston, MA, USA

[16]Current address: Department of Pharmacology and Cancer Biology and Department of Medicine,

Division of Gastroenterology, Duke University, Durham, NC, USA

[17]Current address: Genentech, 1 DNA Way, South San Francisco, CA, USA

[18]Racah Institute of Physics and Faculty of Medicine, The Hebrew University of Jerusalem,

Jerusalem, Israel

[19]Faculty of Medicine, The Hebrew University of Jerusalem, Jerusalem, Israel

[20]These authors contributed equally

*Correspondence: inbalavr@gmail.com (I.A.-D.), jatin.roper@duke.edu (J.R.),

orit.r.rosen@gmail.com (O.R.-R.), mor.nitzan@mail.huji.ac.il (M.N.), aviv.regev.sc@gmail.com

(A.R.)

Abstract

While advances in single cell genomics have helped to chart the cellular components of tumor

ecosystems, it has been more challenging to characterize their specific spatial organization and

functional interactions. Here, we combine single cell RNA-seq, spatial transcriptomics by Slide-

seq, and *in situ* multiplex RNA analysis, to create a detailed spatial map of healthy and dysplastic

colon cellular ecosystems and their association with disease progression. We profiled inducible

genetic CRC mouse models that recapitulate key features of human CRC, assigned cell types and

epithelial expression programs to spatial tissue locations in tumors, and computationally used them

to identify the regional features spanning different cells in the same spatial niche. We find that

tumors were organized in cellular neighborhoods, each with a distinct composition of cell

subtypes, expression programs, and local cellular interactions. Comparing to scRNA-seq and

Slide-seq data from human CRC, we find that both cell composition and layout features were

conserved between the species, with mouse neighborhoods correlating with malignancy and

clinical outcome in human patient tumors, highlighting the relevance of our findings to human

disease. Our work offers a comprehensive framework that is applicable across various tissues,

tumors, and disease conditions, with tools for the extrapolation of findings from experimental

mouse models to human diseases.

INTRODUCTION

The spatial organization of diverse cells in the tumor ecosystem impacts and drives interactions between malignant cells and neighboring immune and stromal cells, either promoting or suppressing tumor growth[1]. Recent studies have shown that systematic understanding of the spatial organization of tumors can shed light on disease progression and response to therapy, with specific features correlated with tumor subtypes[2–4], cancer prognosis[5–7], or response to treatment[8,9].

However, genome-scale, high-resolution dissection of the spatial organization of tumors and its functional implications remains challenging, largely due to technical limitations. Methods such as fluorescent *in situ* hybridization (FISH) and immunohistochemistry can only measure a handful of pre-selected transcripts or proteins, whereas single cell RNA-seq (scRNA-seq) does not directly capture spatial relations. Recent advances in spatial genomics and proteomics allow multiplexed or genome-scale measurements *in situ*[10–17], but with a trade-off between genomic scale and spatial resolution[18]. As a result, data from different experimental methods need to be integrated for a comprehensive view of the tissue biology. Many analytical tools have been developed to integrate some crucial aspects of the data[19–23], but it can be challenging to deploy them and distill answers to specific disease biology questions. This leaves open many fundamental questions about tissue organization and collective function, including whether there are canonical functional units in tumors, what may be their organization in the tumor landscape, and what role does each play in tumor progression.

A case in point is colorectal cancer (CRC), where initial lesions (adenomatous polyps) progress

over time to carcinoma and eventually to metastatic disease. While the mutations that drive this

process were extensively studied[24–27], and the cellular ecosystem of CRC has now been deeply

charted[2,28,29], the spatial landscape is less well-characterized. In a recent study of human CRC[2], we

statistically associated cell profiles across tumors and showed that they map to different cellular

communities, reside in different locations in the tumor and reflect different tumor subtypes.

However, absent genome wide *in situ* measurements, these statistical inferences do not yet reflect

the full spatial organization of the tumor. Moreover, it is important to relate such patterns to those

in animal models used in mechanistic studies and as pre-clinical models, to understand the relation

between lab models and human tumors.

Here, we deciphered the spatial and cellular organization of colorectal cancer (CRC) by combining

scRNA-seq, spatial transcriptomics by Slide-seq, and *in situ* RNA multiplex analysis, using a novel

computational framework. We first profiled two inducible genetic mouse models of colorectal

cancer that recapitulate key features of human CRC, before and after tumor initiation. We

integrated the spatial and cell profiles to create a spatial cell map of the tumor landscape, revealing

dysplasia-specific cellular layout and potential physical interactions. We found that the tumor

landscape is organized in cell neighborhoods, each with distinct epithelial, immune, and stromal

cell compositions, and governed by different gene programs. Three of the cell neighborhoods are

associated with tumor progression, each activating different biological pathways but all active

simultaneously albeit in different parts of the tumor. We devised a computational framework,

based on the TACCO[21] method, extending it to compare single cell and spatial features of tumors

between species and applied it to scRNA-seq data from human CRC. Multiple features were

conserved between tumors in the mouse model and the human patients, and the mouse cellular

neighborhoods correlated with malignancy and clinical outcome (progression-free intervals (PFI)

and overall survival (OS)) in human patient tumors. Our work provides a general approach that

can be applied to other tissues, tumors and disease conditions.

RESULTS

A cell atlas of genetic models of colorectal cancer

To chart the cell ecosystem of CRC and how it changes during tumor progression, we studied two

genetic mouse models of CRC, one with inactivation of *Apc* and another in which *Apc* inactivation

is accompanied by an oncogenic $Kras^{G12D/+}$ mutation and inactivation of $Trp53^{30–32}$ (**Figure 1A**).

In the AV model, $Apc^{fl/fl}Villin^{creERT2}$ mice are injected with 4-hydroxytamoxifen to the submucosal

layer of the colon, inducing the deletion of *Apc* specifically in epithelial cells within the injection

site[30]. In the AKPV model ($Apc^{fl/fl}$; LSL-$Kras^{G12D}$; $Trp53^{fl/fl}$;$Rosa^{26LSL\text{-}tdTomato/+}$; $Villin^{CreERT2}$ mice,

Methods), 4-hydroxytamoxifen injection also induces an oncogenic $Kras^{G12D/+}$ mutation and then

inactivation of *Trp53*. In both cases, 4-hydroxytamoxifen injection leads to the formation of local

lesions that resemble human dysplastic lesions[30].

We first generated a single-cell atlas of the models consisting of 48,115 high quality scRNA-seq

profiles from normal colon, AV (3 weeks after 4-hydroxytamoxifen induction) and AKPV (3 and

9 weeks after induction) tissues. We captured a diverse cell census (**Figure 1B,C**, **Methods**), with

35 clusters annotated *post hoc* by the expression of known marker genes (**Figure 1B**, **Figure**

S1A,B, **Methods**), across epithelial, immune and stromal cell compartments (**Figure S1C**). For

validation purposes, we also generated multiplex *in situ* RNA profiles (with the Cartana[33] method)

in 6 sections each from normal and AV conditions, using a panel of 66-180 marker genes chosen

to best represent the cell types and programs found in the tissues (**Figure S1D, Methods**).

Tumorigenic genotypes cause shifts in the composition of epithelial cell populations and their

microenvironment in AV and AKPV lesions

Dysplastic lesions exhibited shifts in proportions of immune and stromal cells, including changes

in subsets pre-existing in normal tissue, as well as infiltration of new cell subsets (**Figure 1C,D**

and **Figure S1D,E,F**). This resulted in both increase in cells of existing populations (*e.g.*, some T

cell subsets (TNK05 (GdT/Il17+), TNK06 (Treg)) and emergence of new dysplasia-associated

cells (*e.g.*, granulocytes (Gran01, Gran02) and monocytes (Mono02, Mono03)) mirroring

observations in human CRC[2], breast cancer[34], and non-small cell lung cancer[35] (**Figure 1C,D** and

Figure S1D,E and **Figure S2 A-C**). We validated these patterns using multiplex *in situ* RNA

analysis (**Figure 1E**, **Figure S1F-H** and **Supplementary item 1 and 2**). Infiltration is likely to

underlie many of these changes as many of the increasing cell subsets (granulocytes, monocytes,

mast cells) expressed genes, such as Sell and Ccr2, indicating tissue recruitment, and as the cells

dramatically increase in proportion despite negligible signals of proliferation programs.

Two of four monocyte subsets, Mono02 and Mono03, were unique dysplasia-associated cells

(**Figure S2D-F**) and were respectively enriched for general inflammatory response genes

(FDR=5.7 10^{-30}, two-sided Fisher's exact test in GO term enrichment) and interferon beta and

gamma response genes (FDR=3.5 10^{-11}, 1.0 10^{-13}). T cell subsets showed the expected diversity

across nine subsets (**Figure S2G-I**)[36], with a significant decrease (out of all T cells) in TNK01

(GdT/Cd8) in the dysplastic microenvironment and an increase in TNK05 (GdT/Il17+) (**Figure**

S2J). This is consistent with the T cell composition in tumors from mismatch repair proficient

(MMRp) CRC patients (**Figure S2K**). RNA velocity analysis[37] of T cells from normal and AV

tissue (**Figure S3A**) showed a change in inferred cellular relationships with TNK03 (naïve T) and

TNK02 (Th1/Th17) preceding TNK06 (Treg), consistent with the promotion of an

immunosuppressive microenvironment, and TNK08 (proliferating T) also preceding TNK02

(Th1/Th17), TNK04 (GdT/Cd8) and TNK05 (GdT/Il17+) populations.

Within the five subsets of stromal cells (including vascular endothelial and lymphatic endothelial

cells and three fibroblast subsets, **Figure S2L-N** and **Methods**), Endo01 (vascular) were enriched

in dysplastic lesions compared to normal colon (FDR=1.8×10^{-3}, two-sided Welch's t test on CLR

transformed compositions; **Figure S2O**). Angiogenesis-related pathways, such as angiogenesis

(FDR=1.0×10^{-11}) and positive regulation of angiogenesis (FDR=3.4×10^{-4}) as well glycolytic process

(FDR=1.4×10^{-4}) were enriched in Endo01 (vascular) from lesions compared to normal colon (**Table**

S1). This is in line with vascular adaptation to the tumor's growing needs for nutrients and

oxygen[38] and with the increased expression of the vascular growth factor *Vegf-A* in both monocytes

and macrophages (**Figure S2P**).

Cell-intrinsic expression shifts in different sub-lineages in the dysplastic epithelium

Epithelial cells showed dramatic cell-intrinsic changes between normal tissues and either AV or

AKPV lesions, such that the cell profiles of dysplastic epithelial cells in both models were highly

distinct from normal epithelial cells (and similar to each other) (**Figure 1C**, **Figure S4A** and

Methods). Epithelial cell profiles from normal mice (41% of cells) separated from most of those

from AV and AKPV models (59% of cells) (**Figure 2A** and **Figure S4A,B**), suggesting a common

shift in all dysplastic cells from the normal state. Notably, 11% of epithelial cells from AV/AKPV

mice were classified as non-dysplastic healthy cells, indicating that normal, non-dysplastic, cells

may be present in or adjacent to the lesion microenvironment (**Figure 2B** and **Figure S4A**),

although we cannot firmly rule out adjacent tissue contamination. We annotated two cell clusters

– Epi01 (dysplastic stem-like) and Epi05 (dysplastic secretory-like) – as dysplastic, due to their

virtually exclusive presence in AV and AKPV models and because they expressed high levels of

Apc target genes (*e.g.*, *Axin2, Ascl2, Myc, Ccnd1, Lgr5*) and were enriched in tdTomato[+] cells from

AKPVT mice (**Figure 2C** and **Figure S4C,D**) while also spanning the enterocyte to secretory

continuum with healthy cells (**Figure S4B**, PC2).

Interestingly, Epi05 (dysplastic secretory-like) had distinguishing markers (*e.g.*, *Ccl9, Mmp7,*

Ifitm3) from their counterparts in normal tissue, Epi04 (secretory) (**Figure 2A-C** and **Figure**

S4A,C,E). *Mmp7* and *Ifitm3* are known to promote metastasis in human CRC[39,40], and *Ccl9*

expression by epithelial cells promotes tumor invasion through recruitment of Ccr1[+] myeloid cells

to the tumor's invasive front in a mouse model of CRC[41]. Notably, *Ccr1* is expressed by newly

recruited monocytes, macrophages and granulocytes in our model, suggesting a potential

mechanism for tumor infiltration and invasion (**Figure S4E**). Thus, dysplastic secretory epithelial

cells may perform additional functions that differ from those of their healthy counterparts.

RNA velocity[37] analysis of the epithelial cell compartment predicted that in the dysplastic

epithelium (**Figure S3B, right**) a proliferating stem-cell like dysplastic subset is a direct source to

both a massively expanded and heterogeneous non-proliferating population of dysplastic stem-like

cells (expressing WNT signaling and angiogenesis programs) and to dysplastic cells of different

'differentiation states' (MHCII expressing stem/progenitors leading to enterocytes and secretory-

like cells) (**Figure S3B,C**). Conversely, in the normal epithelium (**Figure S3B**, left), the 'root' is

placed in a much smaller population of proliferating intestinal stem cells, and the stem cell

compartment is overall much more modest. The proliferation and differentiation path of previously

identified normal epithelial cells remains intact (*i.e.*, reminiscent of the one in normal samples)

even in the dysplastic lesions (**Figure S3B**). While our RNA velocity analysis provides insights

into potential cellular trajectories, further experimental validation is required to confirm these

findings.

Expression programs for stem-like functions, Wnt signaling, angiogenesis and inflammation

are activated in dysplastic epithelial cells

Both normal and dysplastic epithelial cells varied along a continuum, as expected and previously

observed in the ongoing differentiation in the colon epithelium[2,42–44] and our RNA Velocity

analysis (**Figure S3**). Using non-negative matrix factorization (iNMF from LIGER[45], **Methods**),

we recovered 20 expression programs spanning the different epithelial functions, and annotated

them by Gene Ontology terms enriched in their top 100 weighted genes (**Figure 2D,E**, **Figure**

S3C, S4F-L, Methods).

The programs enriched in different dysplastic cells highlighted key processes that play a role in

tumor promotion, including stem cell programs, Wnt signaling, angiogenesis, and inflammation

and innate immunity, including interferon alpha, beta and gamma pathways (**Figure 2D,E, Figure**

S3C). In particular, the stem cell program (#16) detected in some cells across all conditions, was

enriched in dysplastic samples (FDR=5.6 10^{-10}, two-sided Welch's t-test on CLR transformed

compositions), reminiscent of a recently described population in human[28,29]. Comparing cells from

dysplastic and normal samples that express the stem cell program, the dysplastic cells had distinct

expression profiles with induction of negative regulators of Wnt signaling (FDR=4.5 10^{-5}, two-

sided Fisher's exact test in GO term enrichment, *e.g.*, *Notum*, Wnt inhibitory factor 1 (*Wif1*) and

Nkd1) and genes that are related to cellular response to interferon-gamma (FDR=1.7 10^{-6}, *e.g.*,

Ccl9, Ccl6) and immune system process (FDR=6.4 10^{-4}, *e.g.*, Ifitm1 and Ifitm3) (**Figure 2E,F**).

This is consistent with recent studies showing that *Apc*-mutant stem cells secrete negative

regulators of Wnt signaling to induce the differentiation of the WT stem cells in their proximity,

thereby outcompeting them and promoting tumor formation[46,47]. Thus, stem cells from dysplastic

lesions may have non-canonical function and regulation. In addition, the programs for Wnt

signaling (expressing both positive and negative regulators; #4, FDR=2.8 10^{-6}), angiogenesis (#14,

FDR=1.2 10^{-9}), inflammatory response (#6, FDR=1.4 10^{-6}), and innate immune response and

interferon response (#7, FDR=1.2 10^{-2}) were all predominantly expressed or enriched in

AV/AKPV epithelium (all with two-sided Welch's t-test on CLR transformed compositions,

Figure 2E and **Figure S4I-M**). These results are consistent with the known role of the Wnt

signaling pathway in CRC, and of angiogenesis, response to hypoxia and inflammation in tumor

progression[48–50].

Malignant-like tissue programs and composition are conserved between mouse and human

tumors

To evaluate the relevance of our findings to human colorectal cancer, we compared them to a

scRNA-seq atlas we recently generated from tumor and adjacent normal tissue from 62 patients

with either MMRp or MMRd CRC[2]. We compared mouse and human tumors in terms of their

epithelial expression programs, cellular composition, and cell associations in multicellular hubs[2].

While our mouse model should more closely resemble MMRp tumors, we compared to both

classes separately and together to identify any shared features.

To assess the similarity between mouse and human programs we controlled for overall cross-

species and batch differences by normalizing program-specific expression profiles with species-

specific background profiles (**Methods**). The Pearson correlation coefficients of these normalized

scores between the human and mouse programs indicate some overlap in the programs defined on

mouse and human data (**Figure S5A**). Epithelial cells from human and mouse tumors expressed

many programs highly correlated between the species (**Figure S5B**, **Methods**), including for cell

cycle, inflammation, epithelial secretory, angiogenesis, Wnt signaling, and normal colon

functions.

Co-variation in cell proportions across samples (by scRNA-seq) was also conserved between

human and mouse tumors, suggesting broad conservation of tumor composition. For example, in

both species the proportion of endothelial cells and fibroblasts correlated across samples, as did T

and B cell proportions in human tumors and mouse dysplastic lesions (**Figure S5C**). Moreover,

when we transferred epithelial program annotations from mouse to human scRNA-seq and

calculated their co-variation across samples in each species, programs 11 (proliferation), 14

(angiogenesis) and 16 (stem cells), co-varied both across dysplastic mouse tumors and across

human MMRp and MMRd tumors (**Figure S5D** and **Methods**), suggesting a conserved dysplastic

tissue architecture.

Integrated spatial and single-cell atlas of mouse CRC tumors

To comprehensively decipher the distribution of cells and programs in the tumor spatial niche, we

next used Slide-seqV2[15] for genome-wide spatial RNA-seq at 10 μm resolution. We sectioned and

profiled frozen tissues from four normal colon and four AV lesions using 10 Slide-seqV2 pucks

(**Methods**), recovering 221,936 high quality beads (**Figure S6A-C**, **Methods**). We then integrated

the single cell census and spatial profiles using TACCO, which allowed us to annotate each bead

with compositions of discrete cell types (from epithelial, immune and stromal compartments) and

to further annotate the epithelial fraction of each bead with a composition of epithelial program

activity (**Figure 1A** "annotation").

We first used TACCO to annotate every bead in the Slide-seq data with a composition of discrete

cell subtypes for every puck separately, using its matching single-cell reference (normal or disease;

Figure 1A, **Methods**). To this end, TACCO iteratively solved optimal transport problems to assign

cell subtypes to fractions of reads of the beads. TACCO relies on unbalanced optimal transport to

allow for shifts in the frequency of cell subtypes in the pucks *vs.* the single-cell dataset, while using

the reference cellular frequencies as prior knowledge (**Figure S6E,G**). TACCO's cell type

mapping recapitulated the muscularis layer in its expected tissue location based on the inferred

cellular composition pattern (**Figure 3A, 4A and S6D**). This illustrates that TACCO's

compositional annotations align well with biological patterns.

Next, we used TACCO to map the epithelial gene programs (defined above), focusing on transcript

counts that are inferred as derived from epithelial cells. TACCO partitioned the read count matrices

for each puck, assigning counts to epithelial cells based on the mapped per-bead cell subtype

annotations (from the first step) and the expression profiles associated with each subtype

(**Methods**). It then summed all epithelial contributions into an epithelial-only spatial count matrix,

followed by optimal transport to assign epithelial program contributions to individual beads, based

on epithelial cell-only read signals. As for cell type mapping, the proportional contribution of the

programs largely recapitulated their contributions in scRNA-seq (**Figure S6F,H**).

Altered and less ordered local cellular organization of dysplastic lesions

We assessed the local cellular architecture in terms of the preferential proximity of cells of certain

type or expressing particular epithelial programs, within a fixed-sized neighborhood, by adapting

an earlier method. We defined a z-score as significance of the observed neighborship relations

compared to the null for neighborhoods of 20, 40 or 60 μm diameter (**Figure 3B,C** and **Figure**

S7A,B). This z-score is defined with respect to a population of random cell type annotations

generated by random permutations of the cell type annotations between the beads, where in our

case we permute fractional cell type contributions.

Cell proximity preferences in the normal colon tissue are consistent with the expected morphology,

validating our approach (**Figure 3A,B**). Epithelial cells were organized such that the differentiated

Epi02 (Enterocytes) are excluded from the stem cell niche (**Figure 3B** and **Figure S7C**), and

endothelial cells and fibroblasts were also spatially co-located in a focused region (**Figure 3B**),

with T cells in their vicinity (**Figure 3B**). Our multiplex *in situ* RNA analysis validated the

exclusion of enterocytes from the stem cell niche, as also seen in Slide-seq data (**Figure S7C,D**).

While some normal tissue features are preserved in dysplastic samples, including co-location of

cells of the same lineage[5,10] (**Figure 3A-C**), there were notable changes, and more disorder. Cell

types were more randomly distributed in AV lesion *vs*. normal tissue, reflected in lower z-scores

(p=1.6 10^{-37}, one-sided Mann-Whitney U test; **Figure S7E**). At short distances, all epithelial cells

(normal and dysplastic) were preferentially located close to cells from the same subtype (**Figure**

3C) and even to cells with similar functions: epithelial cells expressing programs associated with

malignant-like function (*e.g.*, program 4 (Wnt signaling), 14 (angiogenesis) and 16 (stem cells))

resided close to each other and were spatially distant from cells expressing programs that are

related to normal epithelial functions (*e.g.*, program 5 (basolateral plasma membrane), 8 (apical

plasma membrane) and 10 (oxidation-reduction process)), supporting a model where tumor

progression is structured and compartmentalized (**Figure S7F**). Immune and stromal cells were

generally excluded from epithelial cell neighborhoods. Granulocytes aggregated together (self-

proximal) (**Figure 3C**) and were relatively close to endothelial cells and dysplasia-associated

monocytes (Mono02, Mono03), consistent with their recruitment from the blood through the

vessels (**Figure 3C** and **Figure S7G**). Our multiplex *in situ* RNA analysis validated the spatial

enrichment of monocytes and granulocytes near vessels (**Figure S7H**).

Epithelial regional analysis recovers canonical structures in normal colon

To detect distinctive tissue regions in tumors, which lack traditional tissue references, we identified

cellular neighborhoods with both similar epithelial program activity and a particular composition

of immune and stromal cells. Specifically, we first identified "epithelial program regions" as areas

of distinct epithelial program activity, and then found immune or stromal cells associated with

each region (**Figure 1A** "annotation"). Intuitively, we defined "regions" based on both the

similarity in epithelial expression program activity and proximity in space. To do this, after

assigning epithelial programs to epithelial beads, we clustered the beads based on a weighted sum

of spatial proximity and expression program similarity. This results in spatially contiguous

annotation of beads with distinct epithelial program activity, which, together with the immune and

stromal cells in their proximity, compose the "region". Specifically, using TACCO, we defined

epithelial program regions by Leiden clustering of the weighted sum of neighborship graphs for

spatial bead proximity and epithelial expression program similarity, such that transcriptionally

similar epithelial beads on different pucks can be connected (despite "infinite" spatial distance,

Methods). We then used this single framework for region annotation across all pucks (**Figure 4A**),

to determine the distinctive composition of additional cell types in the same set of spatial regions

(**Figure 4B-D** and **Figure S8A**).

In the normal colon, the regional analysis (**Figure 4A**, bottom) robustly recovered the expected

spatial organization of the healthy colon across five regions and their cellular composition and

sublayers (**Figure 1A**), from luminal/apical to basal. Four regions recovered by TACCO

corresponded to different layers of the mucosa (**Figure 4A** and **4E,F**): a luminal layer with reads

found beyond the cellular layer and likely representing cellular debris trapped in the mucus; three

apical layers expressing programs related to normal epithelial function (transmembrane transport,

oxidation-reduction process) with gradual transition from apical to basal features; and a basal-most

layer, enriched for the deep crypt, proliferation (G1/S,G2/M), MHCII and basolateral plasma

membrane programs, all common features of the deep crypt area. Finally, region 2, enriched with

fibroblasts, myofibroblasts and endothelial cells, and located in the most basal side of the tissue,

captured the submucosal and muscularis propria layers, which are predominantly comprised of

fibroblasts and muscle, respectively, alongside blood and lymphatic vessels, nerves and immune

cells. Overall, TACCO recovered the known organization of the colon, showing the power of our

unsupervised mapping approach and shedding light on expression programs that are required for

the maintenance of normal colon homeostasis.

Dysplastic lesions maintain some of the programs of the corresponding regions in healthy

tissue

AV lesions did not maintain the robust organization of normal tissues, and reflected the expected

histopathology of high grade dysplasia, when dysplastic cells are confined to the mucosal layer

and do not invade the submucosa[51] (**Figure 1A** and **4A**, top). Specifically, the submucosal and

muscularis propria layers from both normal and AV lesions were assigned to region 2 (**Figure**

4A).

Despite the altered morphology, some of the disrupted regions also expressed programs

characteristic of their normal healthy function, suggesting that tumor progression is spatially

structured and compartmentalized. For example, the region above the submucosa, captured as

region 1 in AV lesions and region 5 in normal colon (**Figure 4A**), had similar features in both AV

lesions and normal samples. Thus, although the overall spatial organization was disrupted in the

lesion, region 1 in AV lesions expressed programs that are reminiscent of the normal deep crypt

region 5, and was enriched for deep crypt cells and programs that are related to proliferation and

MHC II[43] (**Figure 4A,C**). These included proliferation programs 3 and 11 and both normal stem

cells (Epi03) and dysplastic secretory-like cells (Epi05), as well as dysplastic stem cells (Epi01,

though to a lesser extent than some other regions), so it may reflect one of the proliferative stem

cell (and dysplastic secretory-like) niches in AV models (**Figure 4B,C**). Other regions in the AV

lesions also contained some epithelial cells with normal profiles, expressing programs that should

allow them to maintain their capacity to perform normal tasks. For example, region 3 expressed

apical plasma membrane functions and region 10 was enriched with oxidation-reduction functions

(**Figure 4C**).

To learn about the spatial distribution of the dysplastic regions, we measured their distance from

region 2 (muscularis), which is a stable landmark in the lesions. Remnants of the layered structure

of the healthy tissue were still observed in the AV tissue, especially at relatively low distances

from the muscularis. For example, healthy region 5 – characteristically located at distances of

about 150-200µm from the muscularis – is replaced by dysplastic region 1, peaking at 200µm. All

malignant-like regions (6/8/11) were spatially associated at ~300-700µm from the muscularis

(**Figure S8C**), located ~100-400µm apart from each other (**Figure S8D**). We further validated this

result at the protein level, by staining for b-catenin, showing an (inactive) cytoplasmic localization

in the region adjacent to the muscularis, and mostly nuclear (active) localization in distal regions,

near the lumen (**Figure S8B**).

Three spatially and functionally distinct tumor regions enriched in AV lesions

Three regions – 6, 8, and 11 – had epithelial composition and programs that suggested advanced

malignant-like characteristics, each highlighting a potentially different mechanism for tumor

progression (**Figure 4G,H**). These three 'malignant-like regions', were enriched (*vs*. all other

regions) in stem cell, Wnt signaling and angiogenesis programs (16, #4 and #14; FDR=9.6 10^{-21},

8.1 10^{-10}, 3.7 10^{-10}, two-sided Welch's t-test on CLR transformed compositions) and depleted of

normal epithelial programs (#5, #8, and #10; FDR=5.5 10^{-12}, 1.6 10^{-12}, 4.2 10^{-10}, **Figure 4I**).

Furthermore, the malignant-like regions were enriched in immune cells, including monocytes-

macrophages (FDR<=1.5 10^{-5}; excluding Mac02 (Lyve1+)), T cell subsets TNK02 (Th1/Th17),

TNK05 (GdT/Il17+), TNK06 (Treg), TNK08 (proliferating T) (FDR=2.4 10^{-3}, 2.2 10^{-4}, 1.7 10^{-3},

9.6 10^{-4}; two-sided Welch's t-test on CLR transformed compositions), infiltrating granulocytes

(FDR<=9.7 10^{-3}), and mast cells (FDR=1.4 10^{-2}), suggesting an ongoing immune response

(**Figure 4J**). However, each one of the three regions had a different epithelial program

composition, suggesting that in each type of region there is a different dominant pathway/feature

that may drive tumor progression (**Figure 4C, Table S2**).

Region 6 was characterized by an inflammatory and angiogenic multicellular community, with

epithelial and immune cells expressing inflammatory programs, endothelial cells and monocytes

connected in a pro-angiogenic circuit, and pro-invasive genes expressed by both endothelial and

immune cells (**Figure 4B,C**). Specifically, region 6 was distinctly enriched for proliferation

(programs 3 and 11; FDR=2.2 10^{-14}, 1.2 10^{-15}, two-sided Welch's t-test on CLR transformed

compositions) and inflammatory epithelial programs (programs 6 and 7; FDR=9.0 10^{-7}, 2.0 10^{-11}

^{11}), and its non-epithelial compartment was correspondingly enriched for genes from

inflammatory pathways, including the response to TNF, IL-1 and IFN γ (FDR=3.1 10^{-4} , 2.8 10^{-3},

4.9 10^{-6}, two-sided Fisher's exact test in GO term enrichment), and chemotaxis of monocytes,

neutrophils, and lymphocytes (FDR=1.5 10^{-3}, 3.2 10^{-10}, 1.0 10^{-2}, **Table S2**), suggesting

recruitment of inflammatory cells from the circulation or other parts of the tissue. Region 6 was

also enriched for collagen binding genes and collagen-containing extracellular matrix (ECM)

genes (FDR=1.4 10^{-2}, 7.6 10^{-5}, two-sided Fisher's exact test in GO term enrichment, **Table S2**),

which are important for migration and invasiveness[52]. These include *Sparc*, expressed mainly by

endothelial cells and fibroblasts in our data, known to promote colorectal cancer invasion[53]; and

Ctss, a peptidase expressed by T cells and monocytes-macrophages that promotes CRC

neovascularization and tumor growth[54]. Finally, gene expression patterns in endothelial cells and

monocytes in region 6 suggested active angiogenesis through a multi-cellular feedback loop, with

enriched numbers of vascular and lymphatic endothelial cells expressing immune-attracting

chemokines (*Cxcl9*) and adhesion molecules (*e.g. Chd5, Mcam*), monocytes expressing

proangiogenic factors that induce proliferation of endothelial cells (*e.g.*, *Mmp12*), and monocytes

and macrophages expressing *Ctsd*, which increases tumorigenesis in CRC models[55] (**Figure S8E**).

Region 8 was enriched for deep crypt cells (program 13; FDR=1.7 10^{-14}, two-sided Welch's t-test

on CLR transformed compositions), reminiscent of the normal stem cell niche in normal colon, an

epithelial innate immune program (program 1; FDR=3.4 10^{-100}) expressed by secretory cells in AV

and AKPV lesions, and plasma and B cell activity. Unlike the canonical (normal) deep crypt region

(region 5), which is enriched for MHCII expression (program 18; FDR=8.6 10^{-28}), this region was

depleted for the program's expression (FDR=2.1 10^{-26}), which may indicate an earlier stem cell-

like state[43], or a decoupling of the cell cycle and MHCII programs (which are coupled in normal

ISC differentiation, and allow a cross talk with T cells to modulate T cell differentiation) (**Figure

4B,C**). The region's non-epithelial compartment was enriched for B cell activation and BCR

signaling genes (FDR=4.0 10^{-4}, 1.9 10^{-3}, two-sided Fisher's exact test in GO term enrichment,

Table S2). This may be related to B cell function in protection from lumen antigens[56] or to tertiary

lymphoid structures (TLS), which is correlated with clinical benefits in cancer patients[57]. Notably,

Epi05 (dysplastic secretory-like) enriched in Region 8 (**Figure 4B**) expressed higher levels of

inflammatory genes and immune chemokines (*e.g.*, *Ccl9, Ifitm3*) compared to normal counterparts,

Epi04 (secretory) (**Figure 2C** and **Figure S4C,E**), and may thus promote the formation of this

region. We validated the presence of TLS-like structures in association with deep crypt secretory

cells in AV lesions using multiplex RNA analysis (Cartana) (**Supplementary item 3**), showing

that the dominant population of B cells is accompanied by monocyte-macrophages and T cells

characteristic of TLSs, as well as the expression of *Reg4* and *Muc2*, deep crypt goblet/secretory

cell markers.

Region 11 was populated by cells expressing the Wnt signaling pathway program (4, FDR=9.3 10^{-}

^{13}, two-sided Welch's t-test on CLR transformed compositions), with several lines of evidence

supporting an active epithelial to mesenchymal transition (EMT) in this region. Epithelial cells in

region 11 were enriched for the expression of mesenchymal genes, including Vimentin[58] (*Vim*,

FDR=7.4 10^{-245}, one-sided Fisher's exact test), *Prox1*[59] (FDR=3.7 10^{-153}), and *Sox11*[60] (FDR=7.7

10^{-224}) (**Figure S8F**), as well as for EMT signatures from a mouse model of lung adenocarcinoma

[61] (FDR=1.5 10^{-142}, two-sided Mann-Whitney U test) and from human head and neck squamous

cell carcinoma tumors[62] (FDR=6.9 10^{-55}, two-sided Mann-Whitney U test). This is consistent with

the role of Wnt signaling in promoting EMT and a mesenchymal phenotype in CRC, breast cancer

and other epithelial tumors[63,64]. Region 11 non-epithelial cells also expressed genes encoding

MHC-I binding proteins (FDR=4.6 10^{-2}, two-sided Fisher's exact test in GO term enrichment,

Table S2) and actin cytoskeleton, filament and binding proteins (FDR=3.1 10^{-6}, 7.7 10^{-3}, 2.7 10^{-}

^{10}). Organization of the cytoskeleton affects migration, adherence, and interaction of lymphocytes

with antigen presenting cells[65]. Notably, region 11 also concentrated at a more distal part of the

tissue at ~900μm from the muscularis suggesting an outgrowth of the tissue towards the lumen

(**Figure S8C**).

Non-epithelial cells formed two cellular hubs in the malignant-like regions (6,8,11) (**Figure S8G**):

An endothelial-fibroblast hub, detected in all three regions, and an immune hub with B cells, TNK

cells, monocytes, and macrophages, which was prominent in inflammatory region 6, weaker (less

spatially correlated) in region 8 (but validated *in situ*), and not correlated in region 11. Thus,

activation of an immune response is reflected by close proximity between immune cells. We

further characterized the organization of the vascular niche using our multiplex *in situ* RNA data,

finding that while neighbors of the Pdgfrb-expressing pericytes are mainly other Pdgfrb-

expressing pericytes and endothelial cells, Pecam1-expressing endothelial cells appear self-

enriched next to themselves at cellular scale distances and close to Pdgfrb-expressing pericytes

for larger distances (**Figure S8H**).

Overall, three multicellular community regions were enriched in AV lesions: (**1**) inflammatory

epithelial regions with endothelial cells and monocytes expressing angiogenesis, inflammation and

invasion programs; (**2**) epithelial stem-like regions, associated with plasma and B cell activity; and

(**3**) regions with epithelial to mesenchymal transition (EMT) and Wnt signaling dysplastic cells.

Each region highlights different processes that modulate tumorigenesis or invasion, and the three

regions co-exist in the same tumor at different spatial locations.

Cell-cell interactions are rewired in AV lesions

To identify cell-cell signaling mechanisms that may underly these regional associations, we used

COMMOT[66], a computational framework that uses Optimal Transport to infer cell-cell

communication from receptor-ligand expression patterns in spatially resolved data. We used

COMMOT's bead-wise communication 'output' and devised a method to address p-value inflation

in statistical enrichment testing, using spatially-informed data aggregation (**Methods**).

We observed stronger and distinct ligand-receptor interactions in AV *vs*. normal samples,

reflecting the activated state in the dysplastic tissue (**Figure 4K, Supplementary item 4A**). In

particular, while interactions enriched in AV lesions involved immune, epithelial and stromal

signaling, those enriched in normal tissue involved neuropeptides, such as NPY and GCG

(glucagon). Moreover, malignant-like regions 6 and 11 as well as region 2 (muscularis) were

particularly enriched for active communication pathways (**Figure S8I** and **Supplementary item**

4B). This included the WNT signaling pathway, angiogenesis (VEGF, PDGF, FGF) and the OSM

pathway.

Similar spatial patterns in human and mouse tumors

The overall spatial distribution of cell types and epithelial profiles was conserved between mouse

and human tumors, when comparing to scRNA-seq[2,28,29,67–70]. We examined mouse-defined

regions in human tumors, using TACCO to map the expression profiles associated with the

epithelial, immune and stroma compartments in each of the TACCO-identified mouse regions to

scRNA-seq profiles from human CRC, and probabilistically annotated region-specific expression

profiles for each scRNA-seq profile from the human samples. This identified two main "meta

compartments", with epithelial, stromal and immune profiles from human MMRp and MMRd

tumors associated with regions 6, 8 and 11 that were enriched in AV lesions (as well as 0 and 2),

while those from normal human tissue were associated with normal regions (*e.g.*, 5, 10, 12)

(**Figure 5A** and **Figure S9A**, **Methods**).

Malignant-like regions are associated with tumor progression in human colorectal tumors

We next assessed if the regional epithelial programs that we spatially identified in mouse are

conserved in human. To this end, we constructed pseudo-bulk profiles from epithelial cells for our

mouse samples and for recently published human samples profiled along different stages of

malignant transformation, from normal tissue to polyp to CRC[2,28,29,67–70]. We scored each epithelial pseudo-bulk profile with the differentially expressed genes between the epithelial parts of the regions and computed the principal components of these scores across all human and mouse samples (**Methods**). The first principal component (PC1) captured features that are related to malignancy, with higher values for human tumors *vs*. polyps (**Figure 5B** and **Figure S9B**). In addition, malignant-like region (6/8/11) scores were higher in dysplastic *vs*. normal samples (**Figure 5C**). Thus, the spatial region profiles defined in mouse capture features that correlate with malignant transformation in human.

We further classified each full pseudo-bulk profile from the dysplastic human samples into one of the four groups in the CMS expression-based classification[71] (**Methods**) and compared the mouse region scores for each class of samples (**Figure S9C, Methods**). CMS2 classified samples were most closely related to our dysplastic mouse models: all of the dysplasia -associated regions were enriched in CMS2 tumors while most normal regions were depleted, relative to the other CMS classes.

Finally, we found that expression of the malignant like regions (6/8/11) in tumors was associated with clinical outcome. We scored each tumor based on genes that were differentially expressed between the full expression profile of malignant-like regions (6/8/11) and compared the progression-free interval (PFI) and overall survival (OS) for patients in TCGA whose RNA-seq profiles were in the top and bottom quartile of malignant-like region scores (**Methods**). High scores for malignant-like region 11 (EMT) were correlated with shorter PFI, while those for malignant-like region 6 (inflammation) correlate with longer PFI and longer OS, (**Figure 5D,E**).

These associations were driven primarily by MMRp tumors **(Figure S9D,E)**. This suggests that

region 11 is associated with pro-tumorigenic properties in human patients, while region 6 might

be associated with tumor controlling properties. This highlights the importance of multicellular

functional tissue modules in the CRC tumor ecosystem.

DISCUSSION

Here, we systematically charted the spatial organization of cellular expression in dysplastic tissue

of the colon, to help identify putative functional units in the tumor. We used TACCO[21] to integrate

scRNA-seq and Slide-seq data, not only by mapping cell types to their positions, but also

distinguishing different cell programs, the regions that they dominate, and their characteristic

microenvironments. This allowed us to overcome technical limitations, such as lack of spatial

context in scRNA-seq and sparse readout in Slide-seq, and to generate a high-resolution spatial

map of the dysplastic landscape transcending beyond the mapping of individual cells to spatial

positions. We used this map to show correlation with clinical outcome in human patient tumors.

Our scRNA-seq analysis revealed profound enrichment of a stem cell program in dysplastic

tissues. The profiles of dysplastic cells expressing this program are distinct from normal stem cells

and enriched with expression of negative regulators of the WNT signaling pathway and

inflammation, suggesting a non-canonical function. The abundance of these cells with stemness

potential across all our malignant-like regions, points to a dynamic population that can affect the

cells in its proximity, by secretion of negative regulators of the WNT signaling and inflammatory

function but may also adopt various functions depending on the environmental cues and dysplasia-

associated cells in its proximity. A similar population, designated "high-plasticity cell state", was

previously described in a mouse model of lung adenocarcinoma and in human patients, where it

was correlated with resistance to chemotherapy[61]. Whether these cells can be manipulated to take

on specific phenotypes or even to differentiate into normal-like enterocytes given the appropriate

signal from the microenvironment, remain as open questions.

Within the dysplastic lesions, alongside malignant-like regions, we found regions with normal

features (Regions 3,4,9,10), comparable to regions found in the normal colon, most likely

representing compartments driven by clones that were not affected by the genetic perturbation.

One of these regions, region 4, contained mainly goblet cells with normal expression profiles.

Whether this neighborhood represents normal cells that reside alongside malignant cells or a

cancer transition state, it may modify tumor progression, by recruiting immune cells or by

secreting factors that affect epithelial proliferation in adjacent regions. For example, region 4 in

dysplastic lesions is enriched with chemokine activity genes relative to region 4 in normal colon

suggesting a possible role in recruitment of immune cells to the dysplastic landscape. Further work

is required to understand the role of these regions (expressing normal features) in tumor

progression.

While the malignant-like regions were identified as discrete spatial entities, each with coordinated

features across epithelial, immune, and stromal cells, these regions are adjacent to each other.

Thus, they may still influence one another through signaling or by utilizing branches of the same

main vessels. For example, *Osm* is expressed by cells in region 6, whereas its receptor is expressed

on fibroblasts and endothelial cells enriched in region 11. *OSMR* was previously shown to be

expressed by inflammatory fibroblasts[44,72] and its activation in malignant cells promotes EMT in

breast cancer and pancreatic cancer[73,74] and a mesenchymal state in glioblastoma[75]. Future studies

can help determine if these regions are functionally inter-dependent and if they evolved from the

same clones and can inter-convert, or whether they developed independently.

Because animal models complement cell and tissue atlases of human colorectal cancer[2,28], by

allowing experimental manipulation for mechanistic studies[76,77], it is important to relate between

models and patients. By studying genetically engineered mouse models using high resolution

single cell spatial genomics we can help determine to what extent they recapitulate the cellular and

spatial organization of human disease, in the context of two distinct genetic states that represent

human CMS-2 lesions. To this end, we developed several approaches to allow cross species

comparison of tumors at the single cell and spatial level, despite the high level of both intra- and

inter-individual variation within each species. Comparing to human CRC, our analysis suggests

that the CRC landscape is organized in similar multicellular functional tissue modules between

human and mouse, and disease subtypes (*e.g.*, MMRp and MMRd). Future studies applying our

approaches to patient cohorts could help understand whether the expression of different tissue

modules may contribute to the partial response to immunotherapy reported for MMRd patients[78],

and to define specific tissue modules predictive of response to therapy. Notably, while our study

focused on the tumor landscape, its findings may be relevant for tissue response to other challenges

(*e.g.*, inflammation, fibrosis, wound healing), which involve activation of similar functional tissue

modules, a result of collective function of parenchymal, immune and stromal cells.

Taken together, our integrative approach facilitates spatial analysis with high resolution,

constructing regional neighborhoods and their spatial layout at both high cellular resolution and

genomic scale. Our work is an important step toward a systematic understanding of the

organization of dysplastic tissue with the potential to contribute to improved patient stratification

by the multicellular functional units in the tumor landscape.

Methods

Human subjects

The MGH Institutional Review Board approved protocols for tissue collection used for sequencing

(Protocol 02-240). Informed consent was obtained from all subjects prior to collection. Only

patients with primary treatment- naive colorectal cancer were included in this study.

Mice

Mice were housed in the animal facility at the Koch Institute for Integrative Cancer Research at

MIT. All animal studies described in this study were approved by the MIT Institutional Animal

Care and Use Committee (Protocol 1213-106-16). *Apc$^{fl/fl}$* mice[79] were obtained from NCI mouse

repository, *Kras$^{LSL-G12D/+}$* Ref.[80], *Rosa26$^{LSL-tdTomato}$* Ref.[81] and Trp53$^{fl/fl}$ Ref.[82] mice obtained from

Jackson, *Villin^{CreERT2}* Ref.[83] mice were a gift from Dr. Sylvie Robine. All mice were maintained

on C57BL/6J genetic background. Approximately equal numbers of male and female mice of 6–

10 weeks of age were used for all experiments. Where indicated, mice were injected to the

submucosal layer of the colon with 4-hydroxytamoxifen (EMD Millipore # 579002) dissolved in

ethanol at a concentration of 100 μM (for the mice that were kept for 3 weeks after injection) or

30 μM (for the mice that were kept for 9 weeks after injection). Tumors were resected at either 3

or 9 weeks after 4-hydroxytamoxifen injection. Colonoscopy and colonoscopy-guided injection

methods were previously described in detail[30,31].

Tissue processing for scRNA-seq

Single-cell suspensions from healthy colon or dysplastic lesions were processed using a modified

version of a previously published protocol[44]. Tissue samples were rinsed in 30ml of ice-cold PBS

(ThermoFisher 10010-049), chopped to small pieces and washed twice in 25 ml PBS, 5mM EDTA

(ThermoFisher AM9261), 1%FBS (ThermoFisher 10082-147). To prime tissue for enzymatic

digestion, samples were incubated for 10 minutes at 37°C, placed on ice for 10 minutes before

shaking vigorously 15 times followed by supernatant removal. Tissues were placed into a large

volume of ice-cold PBS to rinse prior to transferring to 5ml of enzymatic digestion mix (Base:

RPMI1640, 10 mM HEPES (ThermoFisher 15630-080), 2% FBS), freshly supplemented

immediately before use with 100 mg/mL of Liberase TM (Roche 5401127001) and 50 mg/mL of

DNase I (Roche 10104159001), and incubated at 37°C with 120 rpm rotation for 30 minutes. After

30 minutes, enzymatic dissociation was quenched by addition of 1ml of 100% FBS and 10mM

EDTA. Samples were then filtered through a 40 mM cell strainer into a new 50 mL conical tube

and rinsed with PBS to 30 mL total volume. Tubes were spun down at 400 g for 7 minutes, at 4°C.

Resulting cell pellets were resuspended in 1ml PBS, placed on ice and counted.

Cell hashing

Cell hashing was performed based on the published protocol[84] as summarized below. Dissociated

cells were resuspended in 1ml of Cell Hashing Staining Buffer (1× PBS with 2% BSA (New

England Biolabs, B9000S) and 0.02% Tween (Tween®-20 Solution, 10%, Teknova, VWR-

100216-360) and counted. 500,000 cells were resuspended in 100 µL of Cell Hashing Staining

Buffer and incubated for 30 minutes on ice, with 2 µL of the appropriate BioLegend TotalSeq™

Hashing antibody (a 1:50 dilution, using a total of 1 µg of antibody per cell suspension).

TotalSeq™-A anti-mouse Hashtag antibodies #1-8 (catalog numbers:155801, 155803, 155805,

155807, 155809, 155811, 155813, 155815) were used. Cells were washed three times with 0.5 mL

of Cell Hashing Staining Buffer and filtered through low-volume 40-µm cell strainers. All cell

suspensions were recounted to achieve a uniform concentration of 7,000 cells per microliter before

pooling for capture by 10x Chromium controller following the manufacturer protocol for the v2 or

v3 3' kit (10X Genomics, Pleasanton, CA).

Hashtag oligo (HTO) library preparation

Separation of hashtag oligo (HTO)-derived cDNAs (<180 bp) and mRNA-derived cDNAs

(>300 bp) was done after whole-transcriptome amplification by performing 0.6× SPRI bead

purification (Agencourt) on cDNA reactions as described in 10x Genomics protocol. Briefly,

supernatant from 0.6× SPRI purification contains the HTO fraction, which was subsequently

purified using 1.4 and 2× SPRI purifications per the manufacturer's protocol (Agencourt). HTOs

were eluted by resuspending SPRI beads in 15 µL TE. Purified HTO sequencing libraries were

then amplified by PCR (1µL clean HTO cDNA, 25µL 2X NEBNext Master Mix (NEB #M0541)),

10 µM SI-PCR and D701or D704 primers performed dial out PCR (98°C (10 sec), (98°C for 2 sec,

72°C for 15 sec) x 12/18 then 72°C for 1 min) for 12 and 18 cycles, and used the 18 cycles product

for sequencing. PCR reactions were purified using another 2× SPRI clean up and eluted in 15 µL

of 1× TE. HTO libraries were quantified by Qubit High sensitivity DNA assay (ThermoFisher)

and loaded onto a BioAnalyzer high sensitivity DNA chip (Agilent).

SI-PCR: AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGC*T*C

D701:

CAAGCAGAAGACGGCATACGAGATCGAGTAATGTGACTGGAGTTCAGACGTGTGC

D704 :

CAAGCAGAAGACGGCATACGAGATGGAATCTCGTGACTGGAGTTCAGACGTGTGC

Sequencing

Samples were sequenced using HiSeq X (Illumina). Hashing libraries were sequenced with spike-

ins of 2.5%.

Tissue processing for Slide-seq

Colons were flashed with cold PBS and a segment that includes the lesion and surrounding tissue

(or a respective healthy segment from normal mice) was dissected. Samples were then mounted in

cold OCT, flash frozen on dry ice covered with ETOH and long-term stored in -80°C.

Slide-seq

For mouse and human experiments, 10 μm sections were cut and the Slide-seq V2 protocol was

used as previously described[15]. For mouse experiments, four and six arrays were collected from

normal colons and AV lesions respectively. The muscularis was fit onto the array of both healthy

and dysplastic lesions to allow appropriate orientation.

Multiplex *in situ* RNA Analysis

Multiplex *in situ* RNA analysis was performed with Cartana[33] technology (a newer version is now

available as Xenium (10x Genomics)). In total we measured 3 samples with one section per sample

in each state (normal/AV) and probe set (V1/V2), with an additional replicate section for one of

the samples in normal V2.

Fresh Frozen OCT-embedded tissues from normal colon and AV lesions were cryosectioned as 10

μm sections and placed onto SuperFrost Plus glass slides (ThermoFisher) and further stored at -

80˚C before experiments. Samples were fixed (with 4% formaldehyde) for 5 minutes and

permeabilized for 5 minutes (with 0,1 mg/mL pepsin in 0.1M HCl (P7012 Sigma-Aldrich)) prior

to library preparation.

For library preparation, chimeric padlock probes (directly targeting RNA and containing an anchor

sequence as well as a gene-specific barcode) for a custom panel of 66 (V1) or 180 (V2) genes

(**Table S3**, see below) were hybridized overnight at 37°C, then ligated before the rolling circle

amplification was performed overnight at 30°C using the HS Library Preparation kit for

CARTANA technology and following manufacturer's instructions. All incubations were

performed in SecureSeal^{TM} chambers (Grace Biolabs). Note that prior to final library preparation,

optimal RNA integrity and assay conditions were assessed using Malat1 and Rplp0 housekeeping

genes only using the same protocol.

To prevent tissue sections detachment, an additional baking step of 30 minutes at 37°C was

performed before mounting. To quench autofluorescence background, TrueView (SP-8400

VectorLabs) was used for 1 minute at room temperature. For tissue sections mounting, Slow Fade

Antifade Mountant (Thermo Fisher) was used for optimal handling and imaging.

Quality control of library preparation was performed by applying anchor probes to detect

simultaneously all rolling circle amplification products from all genes in all panels. Anchor probes

are labeled probes with Cy5 fluorophore (excitation at 650 nm and emission at 670 nm).

All samples passed quality control and went through *in situ* barcode sequencing, imaging and data

processing. Briefly, adapter probes and sequencing pools (containing 4 different fluorescent labels:

Alexa Fluor® 488, Cy3, Cy5 and Alexa Fluor® 750) were hybridized to the padlock probes to

detect the gene-specific barcodes, through a sequence specific signal for each gene specific rolling

circle amplification product. This was followed by imaging and performed 6 times in a row to

allow for the decoding of all genes in the panel. To reduce lipofuscin autofluorescence, 1X

Lipofuscin Autofluorescence Quencher (Promocell) was applied for 30 seconds before

fluorescence labeling.

Raw data consisting of 20x or 40x images from 5 fluorescent channels (DAPI, Alexa Fluor® 488,

Cy3, Cy5 and Alexa Fluor® 750) were each taken as z-stack and flattened to 2D using maximum

intensity projection. After image processing and decoding, the results were summarized in a csv

file and gene plots were generated using MATLAB [85].

scRNA-seq pre-processing and quality control filtering

Count matrices for scRNA-seq were generated using the Cumulus feature barcoding workflow

v0.2.0[86] with CellRanger v3.1.0 and the mm10_v3.0.0 mouse genome reference. Cell profiles were

quality filtered by requiring between 1,000 and 50,000 counts, and between 500 and 7,000 genes,

less than 20% mitochondrial counts, and less than 10% hemoglobin counts. Cell profiles that did

not meet all these criteria were discarded. The top 5,000 highly variable genes were annotated on

the remaining cells after normalization to 10,000 counts and log1p transform using Scanpy's

"highly_variable_genes" function[87] and providing the chemistry (v2/v3) by hashing (True/False)

combination as batch-annotation. Putative doublets were removed using Scrublet[88] with default

parameters.

Selection of variable genes, dimensionality reduction and clustering

A preliminary clustering using the Leiden algorithm with resolution 1.0 was performed after normalization to 10,000 counts, log1p transform, correction for number of counts and percentage of mitochondrial genes, scaling with a max_value of 10, and generating a k-nearest neighbors (k-NN) graph with 15 neighbors on a PCA of the previously annotated 5000 highly variable genes with 50 components using Scanpy[87]. The single cell profiles were provisionally annotated with SingleR[89] cell-wise (i.e. without using clustering information) using the SingleR built-in MouseRNAseqData and an intestine specific dataset from Tabula Muris[90] (https://figshare.com/ndownloader/files/13092143). For further processing, the dataset was then split into the three compartments, epithelial, immune and stromal, using the provisional SingleR annotations.

For each compartment, the top 5,000 highly variable genes were annotated using Scanpy's "highly_variable_genes" function on cells normalized to 10,000 counts after log1p-transformation and providing the chemistry (v2/v3) by hashing (True/False) combination as batch-annotation.

Expression programs and batch correction

For the dataset of each compartment separately (generated as described above), an integrative NMF was performed (using a part of the LIGER[45] implementation) with k=20 and lambda=5 to identify 20 programs and their respective weights per cell. This iNMF factorization represents the single cell expression matrix as a weighted sum of profiles such that both the weights and programs contain only non-negative numbers, while allowing for and separating out batch-only contributions. The same approach was also used with a higher k (epithelial and immune: 200, stromal: 50) to yield a detailed and batch corrected decomposition of expression which was then

combined to obtain a count-like corrected expression matrix for the top 5,000 highly variable

genes. For each compartment separately, these batch-corrected data were normalized to 10,000

counts, log1p transformed, corrected for number of counts and percentage of mitochondrial genes

by linear regression, scaled with a max_value of 10, followed by a PCA of the previously annotated

5,000 highly variable genes. A k-nearest neighbors (k-NN) graph was constructed from the top 50

PCs, with k=15 neighbors using Scanpy, and clustered using a compartment-specific Leiden

resolution parameter (epithelial: 0.2, immune: 0.4, stromal: 0.1). This clustering was used as the

cluster level annotation of the mouse scRNA-seq data for the epithelial and stromal compartment.

Separately per compartment the data were annotated with SingleR using the cluster information.

The same per-compartment batch-corrected and preprocessed data from the Leiden clustering was

used to create UMAP embeddings with PAGA initialization using Scanpy.

To improve the clustering and annotation in the immune compartment and to filter out additional

doublets not detected by Scrublet, the immune data were separately filtered and clustered using

information from the compartment level clustering and annotation. To that end, myeloid and T/NK

cells were partitioned separately and further processed, and additional likely doublet cells were

labeled and removed by the following procedure:

1. Cells were labeled as doublets based on higher number of UMIs of marker genes for other

compartments than the 95th percentile observed in this immune partition (*i.e.*, Epcam and Cdh1

to remove immune-epithelial doublets and Cav1 and Kdr to remove immune-stromal doublets)

and other immune partitions (i.e., Cd3d, Cd3e, and Cd3g to remove myeloid-lymphoid

doublets from the myeloid cells). This type of filter criterion for lowly expressed genes ("larger

than some percentile" on integer counts) also allows to keep more than 95% of the cells if, for

example, all cells of this partition happened to have 0 UMIs of a particular marker gene.

2. Cells were labeled as doublets if they had inconsistent cell-wise and cluster-wise SingleR

annotations.

3. Cells were labeled as doublets if they had significantly (Benjamini-Hochberg FDR=0.05, one-

sided Fisher's exact test) more neighbors in the k-NN graph from the immune compartment

that were already marked as doublets.

4. All cells labeled as doublets were removed.

After filtering, the count matrices were batch corrected as above using the integrative NMF from

LIGER with k=20 and lambda=5, and clustered like above with group specific Leiden resolution

(myeloid: 0.2, TNK: 0.4). For myeloid and TNK cells, this clustering superseded the original

clustering. The integrative NMF result here was only used for updating the clustering and not for

generating an extra set of expression programs.

Note that cluster Epi06 shows a broad expression spectrum; while such pattern can best be

explained by remaining doublets, they were not called by Scrublet and could also not be

consistently removed by extra QC as was applied to the immune compartment. The interpretation

of Epi06 as containing doublets is supported by its overrepresentation in the annotation of the

Slide-seq data (Figure S6E), as both doublets and slide-seq beads represent compositions of

different cells. Still, non-doublet explanations (e.g. undifferentiated cells) cannot be ruled out.

Therefore, Epi06 cells are retained in the data but not interpreted biologically.

Marker selection for *in situ* RNA analysis (CARTANA)

CARTANA V1 markers were selected from genes differentially expressed between compartments,

cell types and clusters (for the immune and stromal compartments), and highly ranked genes for

programs (for the epithelial compartment) were first filtered by biological relevance and the

literature, obtaining 87 genes. To further reduce the set to the available panel size (66 genes), genes

were annotated into categories of potential redundancy. To choose between redundant genes, a

global objective function was optimized over the gene selection, looping over all potentially

redundant gene sets until convergence, exhaustively testing all choices within a gene set, and

accepting the best choice for this gene set in terms of the global objective function. The global

objective function was constructed as the mean 4-fold cross-validation scores (using

"GroupKFold" and "cross_val_score" from sklearn[91]) of multi-class logistic regression

classification (using "LogisticRegression" from sklearn) for discriminating cell classes and of

ridge regression of epithelial program weights (using "RidgeRegression" from sklearn) on the

scRNA-seq data subsampled to the expected sparsity of CARTANA data. The cell classifications

used were between the stromal, immune, and epithelial compartment, within the stromal

compartment between Endo01, Endo02, and fibroblasts, within the immune compartment between

myeloid and lymphoid lineage, within the lymphoid lineage between T cells and B cells, within

the myeloid lineage between granulocytes, mast cells, and all monocytes and macrophages

together, between monocytes and macrophages, and within monocytes between Mono01, Mono02,

Mono03, and Mono04. The programs used in ridge regression were programs 3, 4, 6, 7, 13, 14,

15, and 16. As the probes for Ly6c1 and Ly6c2 could not discriminate sufficiently between Ly6c1

and Ly6c2, we chose combined probes that measure both.

The CARTANA V2 panel, included 59 of the 66 genes in the V1 panel (the others had to be

removed for technical reasons), another 113 genes from the standard fixed gene panel for

CARTANA, and 8 selected genes from literature.

Analysis of CARTANA data

Each measured molecule was annotated with an originating cell type cluster label (using TACCO's

"tc.tl.annotate_single_molecules", with RCTD[19] as the core annotation method and parameters

bin_size=20, n_shifts=3, assume_valid_counts=True) separately for each sample. For this, genes

in the reference that would likely cross hybridize in the probe panel design were summed over

(Ly6c1 and Ly6c2). TLS-like regions were annotated by visual inspection of the cell type cluster

composition and morphology.

To assess cell type compositions of the full dataset, molecules with cell type cluster annotations

were binned into 10µm bins (using TACCO's "tc.utils.bin" and "tc.utils.hash" functions) and

cluster-level annotations were merged to cell-type level.

To assess the compositions of TLS-like regions, CARTANA v2 data were aggregated, conserving

the categorical TLS annotation (using TACCO's "tc.utils.bin" and "tc.tl.dataframe2anndata"

functions).

Comparison between experimental methods

To compare cell type composition between methods, CLR-transformed compositions of samples

(or of spatially split samples for the spatial methods, see subsection "Ligand-Receptor Analysis

and Spatially Informed Enrichment") were computed. Then, using a 100 bootstrapped means of

the compositions, mean and standard error of the mean were calculated once for the AV and once

for the normal samples. To compare gene expression, the mean difference of gene counts between

normal and AV samples and its standard error were calculated using all pairwise differences

between the bootstrapped normal and AV samples. For gene comparisons, the CLR-transformed

composition over all genes that were measured in all of scRNA-seq, Slide-seq, CARTANA V1,

and CARTANA V2 was used. Pearson correlation between mean compositions was calculated for

each pair of methods.

RNA-velocity analysis

Splicing aware count matrices for scRNA-seq were generated using CellRanger v6.1.2 and

velocyto v0.17.17[37] with the ensembl v108 mouse genome reference. Scvelo v0.2.5 was used to

infer velocity separately for the epithelial and TNK subsets (using the functions

"scv.pp.filter_and_normalize", "scv.pp.moments", "scv.tl.velocity" (with mode='stochastic'), and

"scv.tl.velocity_graph"). Scanpy and bbknn v1.5.1[92] were used to generate batch-corrected UMAP

embeddings for the two subsets for visualization with scvelo's

"scv.pl.velocity_embedding_stream" function.

Selection of human single cell data for the comparison of cell type and epithelial program

composition

ScRNA-seq data from Ref[2] was used as reference for human CRC. To avoid biases in cell type

compositions, only the subset of the data where "PROCESSING_TYPE = unsorted" were used.

Comparison of human and mouse samples by cell type composition

To compare human and mouse samples by composition of T/NK cell subsets, T/NK annotations

from mouse and human data[2] were matched by TACCO, using optimal transport (OT). First,

human expression data were mapped to mouse genes using MGI homology information

[subsection "Mapping of mouse and human orthologs"]. Then, human cell cluster annotations

('cl295v11SubFull') were mapped from the subset of human cells annotated as T/NK/ILC to the

subset of mouse cells annotated as T/NK using TACCOs "annotate" function with OT as core

method, basic platform normalization, entropy regularization parameter epsilon 0.005, marginal

relaxation parameter lambda of 0.1, and 4 iterations of bisectioning with a divisor of 3. Annotation

with maximum probability per cell was used as the unique cluster level annotation for mouse T/NK

cells. Annotations were aggregated per sample to yield a compositional annotation over the

identical cluster annotation categories (from the human dataset) for the T/NK subsets of human

and mouse samples. Annotation vectors were then processed using the sc.pp.neighbors and

sc.tl.umap functions from Scanpy[87] to yield a 2D sample embedding with respect to T/NK cell

composition. Using the coordinates in the UMAP in place of spatial coordinates, neighborship

enrichment z-scores were computed with TACCO's co_occurrence_matrix function with

max_distance=2 and n_permutation=100.

Slide-seq compositional annotation

Slide-seq data were annotated with scRNA-seq reference annotations. First, Slide-seq and scRNA-

seq data were filtered to retain only 15759 genes that were detected in both datasets and only beads

and cells that had at least 10 counts across all these common genes.

Next, TACCO[21] was used to perform compositional annotation of each bead, allowing the bead

counts to be explained by fractional contributions. In its basic application, TACCO finds an

"optimal" mapping between scRNA-seq annotation categories (*e.g.*, cell types) and beads by

solving a variant of an entropically regularized Optimal Transport (OT) problem in expression

space. In its iterative application, TACCO uses a bi-sectioning functionality iteratively, annotating

only fractions of the counts in each round, and reserving the remainder for the next round to

improve the sensitivity to sub-leading annotation contributions (that is, first capture a portion of

the counts for the "top" cell types, but preserving others for other, more minor, cell types).

For the compositional annotation of Slide-seq beads with the categorical cell clusters from the

single cell data, the "annotate" function of TACCO with OT was used as core annotation method

per Slide-seq puck with the subset of the single cell data with matching disease state, with basic

platform normalization, entropy regularization parameter epsilon 0.005, marginal relaxation

parameter lambda of 0.001, and 4 iterations of bisectioning with a divisor of 3.

For the compositional annotation of Slide-seq beads with the compositional epithelial programs,

the annotated beads were split using the "split_observations" function of TACCO on the cluster-

level annotation and then aggregated to compartment level using the "merge_observations"

function keeping only beads for a compartment with at least 50 counts assigned to that

compartment. The genes from the epithelial part were filtered to retain only those that were used

to define the epithelial programs in scRNA-seq, and then annotated using again the "annotate"

function with OT as core annotation method, basic platform normalization, entropy regularization

parameter epsilon 0.01, and a marginal relaxation parameter lambda of 0.001.

Slide-seq region annotation

Region annotation was done for all pucks (normal and AV pucks) in one step to get comparable region annotations across pucks, such that beads that have similar epithelial program activity and are spatially close are called as one region. Because cell type composition can change drastically from one bead to its neighbor at the length scales of Slide-seq data, there is a need to compromise between optimizing the two similarities. This is done with the "find_regions" function of TACCO, which performs a Leiden clustering on the weighted sum of connectivity matrices derived from epithelial program similarity and spatial proximity, using a position weight of 0.7, a Leiden resolution of 1.3, and 15 nearest neighbors per bead in position space and epithelial program space. To determine the neighbors in epithelial program space, the square-roots of the program weights were used for neighbor finding which effectively uses Bhattacharyya coefficients as overlap in epithelial program space instead of the Euclidean scalar products used for position space. These regions are defined by construction only on beads with a large enough epithelial contribution (see above) and are then extended to all beads by assigning unannotated beads the region from the nearest bead with region annotation.

Submucosal and muscularis propria layers are predominantly comprised of fibroblasts and muscle cells, respectively, alongside blood and lymphatic vessels, nerves and immune cells. Our algorithm depends on the epithelial expression component in beads. Since these layers do not contain epithelial cells[93], mapping likely relied either on "noisy" signal from non-epithelial cells or from the basal-most epithelial layer.

To determine region composition at a certain distance of a reference region, TACCO's "annotation_coordinate" function is used with max_distance=1000 and delta_distance=10.

Slide-seq quality filtering

948   For all downstream analyses, all beads with less than 100 reads were discarded.

Region- and cell type- characterizing genes in Slide-seq data

Genes to characterize regions on Slide-seq pucks irrespective of compartment composition were

found using Scanpy's rank_genes_groups function on the full bead expression profiles. To find

them separately for each compartment, the compartment-level split beads [sub-section "Slide-seq

annotation"] were used instead of the full beads. To compare gene expression between cell types

on Slide-seq pucks, cluster-level split beads [sub-section "Slide-seq annotation"] were aggregated

to cell type level.

EMT scoring

957   Malignant regions were scored for EMT signatures, using only counts attributed to the epithelial

958   compartments within these regions and only genes expressed on at least 3 beads. Bead profiles

959   were normalized to 10,000 counts, log1p transformed and scaled, and Scanpy's "sc.tl.score_genes"

960   function was used to score the top 50 genes in two EMT gene signatures[61,94].

Cell-type neighborships in Slide-seq data

To evaluate the local cell-type neighborship relations in the different disease states on the cluster

level, the clusters were filtered per disease state to contain only clusters which account for at least

1% of the UMIs in that state. Then neigbourhood-enrichment z-scores were calculated using

TACCO's "co_occurrence_matrix" function with max_distance=20 and n_permutation=10. To

evaluate the stability of the result, this is also repeated for (max_distance,

n_permutation)=(40,10),(60,10),(20,5), and (20,50). To get the significance of the overall change

in z-scores between the states, a one-sided Mann-Whitney-U test was performed on the values of

the upper triangular half of the matrix between the two disease states for (max_distance,

n_permutation)=(20,10).

A similar neighborship analysis was performed on the coarser cell-type level separately for the

three malignant regions 6, 8, and 11, using TACCO's "co_occurrence_matrix" function with

max_distance=20 and n_permutation=10.

Cell-type co-occurrence in Slide-seq data

Cell-type compositions relative to a spatial landmark, Region 2=muscularis, were evaluated using

TACCO's "annotation_coordinate" function with max_distance=1000 and delta_distance=10. To

reduce tissue structure bias from the muscularis, the distance dependency of cell-type frequency

relations was evaluated only for beads deep in the "epithelial domain", defined as follows. The

effective distance from stromal annotation was computed using TACCO's

"annotation_coordinate" function (with max_distance=100, delta_distance=10,

critical_neighbourhood_size=4.0) and only beads with a distance of at least 75μm were used. On

these remaining beads, TACCO's "co_occurrence" function was used (with delta_distance=20,

max_distance=1000) to compute cell types co-occurrence as a function of their distance.

Epithelial program neighborships in Slide-seq data

As for cell types above, neighborship relations were evaluated for epithelial programs in the AV

Slide-seq samples using TACCO's "co_occurrence_matrix" function with max_distance=20 and

n_permutation=10, after selecting only the programs which make up at least 1% of the UMIs in

the AV Slide-seq samples.

Ligand-Receptor analysis and spatially informed enrichment

For the ligand-receptor analysis on slide-seq data we used COMMOT v0.03[66]. COMMOT employs

optimal transport to construct sender and receiver side receptor-ligand interactions for every bead

in one run of COMMOT. After filtering to beads with at least 100 counts, we applied basic

preprocessing (normalization, log1p-transformation) and load the CellChat database as done in the

COMMOT tutorial (https://commot.readthedocs.io/en/latest/notebooks/Basic_usage.html). We

then follow the Slide-Seq v2 analysis from the COMMOT paper ("slideseqv2-mouse-

hippocampus/1-lr_signaling.ipynb" from https://doi.org/10.5281/zenodo.7272562) to filter the

database and to reconstruct the spatial communication network on ligand-receptor pair and

pathway level separately for every puck. In particular, we use the distance cutoff of 200μm for

inference of ligand-receptor interactions. The resulting bead-wise sender- and receiver

communication values were then used for enrichment analysis between disease states and between

spatial regions.

1004

Unlike the downstream Slide-Seq v2 analysis from the COMMOT paper ("slideseqv2-mouse-

hippocampus/2-downstream_analysis.ipynb" from https://doi.org/10.5281/zenodo.7272562), we

do not treat each bead on the puck as statistically independent observation in statistical tests, which

leads to unreliably small p-values. Instead, we split all pucks along their axis of greatest extent

(defined by the first principal component axis of the distribution of the spatial measurements) into

spatial patches discarding 400µm of boundary layer (twice the COMMOT distance cutoff, also co-

occurrences have decayed strongly at this distance) in between the patches to reduce the correlation

between the patches. This is done iteratively with the patches to get a set of weakly correlated

patches. On these patches we calculate the mean of the communication values and treat them as

statistically independent observations for statistical tests. We argue that multiple sufficiently

separated spatial patches of single spatial samples can be seen as multiple spatial samples using a

spatial method with a smaller measurement area, and therefore can be treated as replicates. Unlike

splitting into patches without removing a boundary layer, this procedure does not converge to the

case of treating each bead as an independent observation as the number of iterations rises as it

accounts for the spatial correlations between adjacent measurements. This gives a natural lower

limit of p-values reachable with p-values rising again if too many splits are performed as too much

data is lost to remove the correlations (**Supplementary item 4C,D**). We chose a number of

iterations of 2 as a compromise between having more patches and not discarding much data. For

the enrichment of the pathway communication values we used a two-sided Mann-Whitney U test

across the patches and cite Benjamini-Hochberg FDR values. For the enrichment testing in a given

group (e.g. an annotated spatial region) only those patches are used which have at least 100 beads

on the patch.

1027

1028   Unless otherwise noted, for consistency, an analogous enrichment procedure was also used for

1029   COMMOT-unrelated quantities, like cell types, even though their spatial footprint is not as large

1030   as COMMOT's (distance cutoff of 200µm). For cluster-level cell type enrichment analyses on

1031   Slide-seq data, only the top 3 contributing subtypes were considered, to better reflect the expected

1032   properties of Slide-seq data and better compare with categorically annotated scRNA-seq data.

Mapping of mouse and human orthologs

We applied TACCO's functions "setup_orthology_converter" and "run_orthology_converter" with option "use_synonyms=True" to map human to mouse genes using the ortholog mapping from Mouse Genome Informatics (http://www.informatics.jax.org/homology.shtml). Specifically we used http://www.informatics.jax.org/downloads/reports/HOM_MouseHumanSequence.rpt (downloaded April 26th, 2021) for the analyses comparing T/NK compositions of human and mouse scRNAseq data, analyses comparing epithelial program similarity in human and mouse scRNAseq data, analyses involving the annotation of human scRNA-seq data with mouse regions, and analyses comparing the cell type and epithelial program correlations cross samples between human and mouse scRNA-seq data, and http://www.informatics.jax.org/downloads/reports/HOM_AllOrganism.rpt (downloaded on August 8th, 2022) for the analyses involving the scoring of mouse regions in mouse and human scRNA-seq data and leading to the CMS classification.

GO term enrichment analysis

We used TACCO's functions "setup_goa_analysis" and "run_goa_analysis" to perform GO terms enrichment. As "gene_info_file" we used https://ftp.ncbi.nih.gov/gene/DATA/GENE_INFO/Mammalia/Mus_musculus.gene_info.gz, as "GO_obo_file" http://purl.obolibrary.org/obo/go/go-basic.obo, and as "gene2GO_file" https://ftp.ncbi.nih.gov/gene/DATA/gene2go.gz (all downloaded on August 10[th], 2022).

Comparison between mouse and human programs

To compare mouse and human epithelial expression programs[2], genes were mapped to mouse homologs using MGI homology information [subsection "Mapping of mouse and human orthologs"]. Mouse and human programs were then characterized by a single vector of mean expression per program in mouse gene space. Specifically, both mouse and human programs were defined such that their weighted sum approximates the expression profiles of the cells without any transformations. Programs and weights were normalized to sum to 1. To reduce batch-effects (including species-specific ones), a background expression profile was defined for each species dataset as the pseudo-bulk epithelial expression profile in the respective scRNAs-seq data. Program and background profiles were normalized to 10,000 counts and the log ratio of the normalized program and background expression vectors was used to define a vector for each species. Pearson correlation coefficients were calculated for each pair of program vectors (mouse vs. human).

Human expression program associations across mouse scRNA-seq

All sets of programs that were previously used to define human CRC tissue hubs[2] (epithelial, T/NK cells and myeloid cells) were mapped to mouse genes with MGI homology information [subsection "Mapping of mouse and human orthologs"] and then to mouse single cell data with TACCO, using TACCOs platform normalization to account for batch effects. The "annotate" function in TACCO was used with OT as the core annotation method, on the comparable subsets of cells from mouse and human single-cell datasets (*e.g.*, myeloid cells from mouse and human), with basic platform normalization, entropy regularization parameter epsilon 0.005, marginal relaxation parameter lambda of 0.1, and 4 iterations of bisectioning with a divisor of 3, and flat annotation prior distribution. The resulting probabilistic per-cell program annotations were aggregated to get

1077 probabilistic per-sample program annotations for all dysplastic mouse samples and CLR-

1078 transformed. For each pair of programs, the Pearson correlation coefficient was calculated on these

1079 transformed values.

1080

Annotating human scRNA-seq data with mouse-derived region information

Human scRNA-seq profiles[2] were mapped to mouse gene space using MGI homology information

[subsection "Mapping of mouse and human orthologs"]. Working in the same expression space,

the "annotate" function in TACCO with OT as core annotation method was used on the full human

scRNA-seq and mouse Slide-seq dataset with basic platform normalization, entropy regularization

parameter epsilon 0.005, marginal relaxation parameter lambda of 0.1, and 7 iterations of

bisectioning with a divisor of 3, and 10-fold sub-clustering of the region annotations. The region

transfer is done separately per compartment, with the Slide-seq compartment split as described

above and the human scRNA-seq data split using the cell type annotation of the data. For

validation, mapping was also performed with mouse scRNA-seq data, as well as mapping the

region information from the mouse pucks back to themselves.

1092

1093 To test for enrichments of region annotations across disease state, region composition was

1094 aggregated to sample-level (for Slide-seq to 4-way split pucks), CLR-transformed, and enrichment

1095 was calculated using a two-sided Welch's t-test. This was done for region annotation on human

1096 and mouse scRNA-seq data, and on the original and mapped region annotation on the mouse Slide-

1097 seq data.

1098

Cell-type associations across samples

To compare associations of cell types across samples in human and mouse scRNA-seq the

"clMidwayPr" cell type annotation in the human data[2] was aggregated to the same level as mouse

cell type annotation, and then aggregated per sample and CLR-transformed. Pearson correlation

coefficients were calculated for every cell type pair for different subsets of samples: all samples,

normal samples, dysplastic, and for human MMRd/MMRp samples.

1105

Epithelial program associations across human samples

To determine epithelial program associations across human samples, TACCO's "annotate"

function was used to annotate human epithelial scRNA-seq (after mapping to mouse orthologs

using MGI homology information [subsection "Mapping of mouse and human orthologs"]) with

mouse epithelial programs from mouse scRNA-seq data using OT as core method, basic platform

normalization, entropy regularization parameter epsilon 0.005, marginal relaxation parameter

lambda of 0.1, and 4 iterations of bisectioning with a divisor of 3. The remaining steps were

performed as for cell-type association (subsection "Cell-type associations across samples").

1114

Scoring epithelial mouse regions in mouse and human epithelial pseudo-bulk data

The published processed and filtered count matrices were used (where available) or instead raw

count matrices for single cell/nucleus RNA seq data from Pelka[2] (GEO accession number

GSE178341; downloaded on August 8th, 2022), Chen[28] (Synapse IDs syn27056096,

syn27056097, syn27056098, syn27056099; downloaded on August 5th, 2022), Khaliq[69] (GEO

accession number GSE200997; downloaded on August 5th, 2022), Becker[29] (GEO accession

number GSE201348; downloaded on August 5th, 2022), Zheng[68] (GEO accession number

GSE161277; downloaded on August 5th, 2022; excluding 'blood' samples), Che[67] (GEO accession

number GSE178318; downloaded on August 5th, 2022; only 'CRC' and 'LM' samples) and

Joanito[70] (Synapse IDs syn26844072, syn26844073, syn26844078, syn26844087, syn26844111;

downloaded on August 22nd, 2022; excluding the 'LymphNode' sample).

1126

To subset the human single cell data to epithelial cells, the epithelial annotation was used where

readily available[2,28]. For the remaining datasets[29,67–70], TACCOs tc.tl.annotate function was used

with default parameters to transfer the 'cl295v11SubShort' annotation from Pelka[2], from which a

compositional compartment annotation was constructed, and then a cell was assigned to the

epithelial compartment if it had more than 95% epithelial fraction.

To correct for batch effects between the different data sources, first batches were defined by

species times protokoll: 'mouse-10x3p', 'mouse-SlideSeq', 'human-10x3p'[2,67,68,70], 'human-

10x5p'[69,70], 'human-inDrop'[28], and 'human-snRNA'[29]. Then TACCO's "tc.pp.normalize_platform"

function was used to determine per gene batch normalization factors using only the normal samples

of one data source per batch (choosing the normal samples from Zheng for 'human-10x3p' and the

normal 5' samples from Joanito for 'human-10x5p'). The resulting factors are then used to rescale

the sample-by-gene count matrices for the full dataset per batch, i.e. including non-normal

samples. The normalization factors are calculated with respect to an (arbitrarily chosen) normal

reference dataset[68].

The epithelial mouse region score was defined as the mean of the CLR-transformed expression

values in the pseudo-bulk expression profile of the epithelial part of a dataset using the top 200

differentially expressed genes between all regions by a one-sided Fisher's exact test.

To account for species-specific biases (in-set *vs*. out-of-set prediction: the DEGs are calculated in

mouse), the scores per region across samples were zero-centered and scaled to unit variance across

all samples (including normal and non-normal samples and all batches) per species. A Principal

Components Analysis (PCA) of the region scores across all species, batches and samples was

conducted and the values for the first PC were compared between different conditions using a two-

sided Mann-Whitney U test with Benjamini-Hochberg FDR.

1150

Assessing the relationship between mouse regions and CMS tumor classification

We used the package 'CMSclassifier' (https://github.com/Sage-Bionetworks/CMSclassifier)

referred to in Ref.[71] to classify human pseudo-bulk CRC profiles from all samples which were

not normal or unaffected from the human studies above into CMS classes. We determine the

enrichment (Benjamini-Hochberg FDR, two-sided Welch's t test) of the same mouse region

scores in the CMS classes.

1157

Assessing the relation between mouse regions and clinical endpoints in human bulk RNA-

seq

Published RNA-seq data from the COAD and READ cohorts of TCGA PanCancerAtlas[95] were

used. Mouse region scores were defined as the mean of the log1p-transformed, zero-centered and

scaled expression values in the bulk expression profile using the top 200 differentially expressed

genes between the malignant mouse regions (6, 8, 11) by a one-sided Fisher's Exact test

(comparing each of the three regions to the other two). Scores were stratified into quartiles. PFI

and OS was compared between patients with tumors whose scores were in the lowest and highest

quartiles using the Logrank test as implemented in the lifelines package[96], followed by Benjamini-

Hochberg FDR.

1168

Compositional enrichment analyses

Enrichments on compositional data (cell type compositions, etc.) were evaluated with a two-sided Welch's t test on sample level using CLR-transformed compositions followed by Benjamini-Hochberg FDR. For the enrichment of tdTomato, counts and ALR-transformation were used instead with all non-tdTomato counts used as reference compartment. Enrichment analyses were performed using TACCO's "enrichments" function.

Code availability

The analysis code is available on GitHub (https://github.com/simonwm/mouseCRC).

Data availability

All data generated in this project is deposited on the Single Cell Portal and is available under the accession SCP1891 (https://singlecell.broadinstitute.org/single_cell/study/SCP1891). The raw scRNA-seq and Slide-seq data is also deposited on GEO under the GSE260801 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE260801). The public human scRNA-seq datasets can be found on GEO under accession numbers GSE178341, GSE200997, GSE201348, GSE161277, GSE178318 and on Synapse under the IDs syn27056096, syn27056097, syn27056098, syn27056099, syn26844072, syn26844073, syn26844078, syn26844087, syn26844111.

ACKNOWLEDGEMENTS

We thank N. Friedman, I. Benhar, N. Habib, M. Biton, K. Geiger-Schuller, J.C. Hütter, B. Dumitrascu, E. Baker and A. Greenwald for helpful discussions. We thank C. McCabe, O. Kuksenko and I. Barrera for technical assistance. We thank P. Yadollahpour, E. Dhaval, G. Smith-Rosario, S. Vickovic, D. Schapiro, S. Farhi, D. Abbondanza, A. Segerstolpe and T. Biancalani for their important contribution to this project. We thank L. Gaffney and A. Hupalowska for help with figure preparation. S.M. was supported by a DFG research fellowship (MA 9108/1-1), J.K. was supported by a HFSP long term fellowship (LT000452/2019-L), A.R. was a Howard Hughes Medical Institute (HHMI) Investigator when conducting this work. Work was supported by the Klarman Cell Observatory, a CEGS grant (5RM1HG006193-09) from the NHGRI, the NIH/NIAID (grants 1U24 CA180922, 1U19 MH114821, 1RC2 DK114784), the MIT Ludwig Center, the Manton Family Foundation, and HHMI (A.R.); Azrieli Foundation Early Career Faculty Fellowship, and an ISF Research Grant (1079/21) (M.N.), the Center for Interdisciplinary Data Science Research at the Hebrew University of Jerusalem (N.M. and M.N.), the Israeli Council for Higher Education Ph.D. fellowship (N.M.), SU2C Peggy Prescott Early Career Scientist Award PA-6146, SU2C Phillip A. Sharp Award SU2C-AACR-PS-32 and NIH/NCI R00CA259511 (K.P.), NIH/NCI R01 CA208756; Arthur, Sandra, and Sarah Irving Fund for Gastrointestinal Immuno-Oncology (N.H.), NIH/NCI R01CA257523, MIT Stem Cell Initiative (Foundation MIT) (O.Y.), and NIH R37CA259363, R21CA256414, R21DK125911, R41EB032693, R01CA254108, R01CA256530, and R01CA244359; DOD W81XWH-20-1-0203; and a Duke-NC State Translational Research Grant (J.R.). The authors gratefully acknowledge LMU Klinikum for providing computing resources on their Clinical Open Research Engine (CORE) and the

Bioinformatic Core Facility of the Biomedical Center Munich for providing computing resources

on their HPC system.

AUTHOR CONTRIBUTIONS

I.A-D. and A.R. conceived the study and designed experiments. I.A-D. J.R., S.Y., E.M., T.D., L.C.

and D.D. performed experiments, with guidance from O.Y. and A.R.. S.M., J.K., and N.M.

developed computational approaches and analyzed data with I.A-D., M.H., E.M. and R.S, and with

guidance from M.N. and A.R.. M.H., E.M., J.C., K.P., A.M. and G.M.B. generated and analyzed

human data, with guidance from N.H., I.T., N.H., F.C., O.Y., O.R-R, and A.R.. A.N. and H.X.

generated and processed the multiplex *in-situ* RNA data, with guidance from J.L. and M.R.. M.N

and A.R provided supervision and acquired funding. I.A-D., S.M., M.N. and A.R. wrote the

manuscript with input from all authors.

DECLARATION OF INTERESTS

A.M. has served a consultant/advisory role for Third Rock Ventures, Asher Biotherapeutics, Abata

Therapeutics, ManaT Bio, Flare Therapeutics, venBio Partners, BioNTech, Rheos Medicines and

Checkmate Pharmaceuticals, is currently a part-time Entrepreneur in Residence at Third Rock

Ventures, is an equity holder in ManaT Bio, Asher Biotherapeutics and Abata Therapeutics, and

has received research funding support from Bristol-Myers Squibb. G.M.B. has sponsored research

agreements with InterVenn Biosciences, Palleon Pharmaceuticals, Olink Proteomics, and Teiko

Bio. G.M.B. is a consultant for Ankyra Therapeutics and InterVenn Bio. G.M.B. has been on

scientific advisory boards for Merck, Iovance, Nektar Therapeutics, Instil Bio, and Ankyra

Therapeutics. G.M.B. holds equity in Ankyra Therapeutics. J.L. and M.R. hold equity in

10xGenomics. N.H. holds equity in BioNTech and is a founder of Related Sciences/DangerBio.

References

1241

1.   McAllister, S.S., and Weinberg, R.A. (2014). The tumour-induced systemic environment as
a critical regulator of cancer progression and metastasis. Nat. Cell Biol. *16*, 717–727.

2.   Pelka, K., Hofree, M., Chen, J.H., Sarkizova, S., Pirl, J.D., Jorgji, V., Bejnood, A., Dionne,
D., Ge, W.H., Xu, K.H., et al. (2021). Spatially organized multicellular immune hubs in
human colorectal cancer. Cell *184*, 4734-4752.e20.

3.   Wagner, J., Rapsomaniki, M.A., Chevrier, S., Anzeneder, T., Langwieder, C., Dykgers, A.,
Rees, M., Ramaswamy, A., Muenst, S., Soysal, S.D., et al. (2019). A Single-Cell Atlas of the
Tumor and Immune Ecosystem of Human Breast Cancer. Cell *177*, 1330-1345.e18.

4.   Hunter, M.V., Moncada, R., Weiss, J.M., Yanai, I., and White, R.M. (2021). Spatially
resolved transcriptomics reveals the architecture of the tumor-microenvironment interface.
Nat. Commun. *12*, 1–16.

5.   Keren, L., Bosse, M., Marquez, D., Angoshtari, R., Jain, S., Varma, S., Yang, S.-R., Kurian,
A., Van Valen, D., West, R., et al. (2018). A Structured Tumor-Immune Microenvironment
in Triple Negative Breast Cancer Revealed by Multiplexed Ion Beam Imaging. Cell *174*,
1373-1387.e19.

6.   Schürch, C.M., Bhate, S.S., Barlow, G.L., Phillips, D.J., Noti, L., Zlobec, I., Chu, P., Black,
S., Demeter, J., McIlwain, D.R., et al. (2020). Coordinated cellular neighborhoods orchestrate
antitumoral immunity at the colorectal cancer invasive front. Cell *182*, 1341-1359.e19.

7.   Jackson, H.W., Fischer, J.R., Zanotelli, V.R.T., Ali, H.R., Mechera, R., Soysal, S.D., Moch,
H., Muenst, S., Varga, Z., Weber, W.P., et al. (2020). The single-cell pathology landscape of
breast cancer. Nature *578*, 615–620.

8.   Jerby-Arnon, L., Shah, P., Cuoco, M.S., Rodman, C., Su, M.-J., Melms, J.C., Leeson, R.,
Kanodia, A., Mei, S., Lin, J.-R., et al. (2018). A Cancer Cell Program Promotes T Cell
Exclusion and Resistance to Checkpoint Blockade. Cell *175*, 984-997.e24.

9.   Grünwald, B.T., Devisme, A., Andrieux, G., Vyas, F., Aliar, K., McCloskey, C.W., Macklin,
A., Jang, G.H., Denroche, R., Romero, J.M., et al. (2021). Spatially confined sub-tumor
microenvironments in pancreatic cancer. Cell *184*, 5577-5592.e18.

10.   Goltsev, Y., Samusik, N., Kennedy-Darling, J., Bhate, S., Hale, M., Vazquez, G., Black, S.,
and Nolan, G.P. (2018). Deep Profiling of Mouse Splenic Architecture with CODEX
Multiplexed Imaging. Cell *174*, 968-981.e15.

11.   Angelo, M., Bendall, S.C., Finck, R., Hale, M.B., Hitzman, C., Borowsky, A.D., Levenson,
R.M., Lowe, J.B., Liu, S.D., Zhao, S., et al. (2014). Multiplexed ion beam imaging of human
breast tumors. Nat. Med. *20*, 436–442.

12.   Giesen, C., Wang, H.A.O., Schapiro, D., Zivanovic, N., Jacobs, A., Hattendorf, B., Schüffler,
P.J., Grolimund, D., Buhmann, J.M., Brandt, S., et al. (2014). Highly multiplexed imaging of
tumor tissues with subcellular resolution by mass cytometry. Nat. Methods *11*, 417–422.

13.   Ståhl, P.L., Salmén, F., Vickovic, S., Lundmark, A., Navarro, J.F., Magnusson, J.,
Giacomello, S., Asp, M., Westholm, J.O., Huss, M., et al. (2016). Visualization and analysis
of gene expression in tissue sections by spatial transcriptomics. Science *353*, 78–82.

14.   Rodriques, S.G., Stickels, R.R., Goeva, A., Martin, C.A., Murray, E., Vanderburg, C.R.,
Welch, J., Chen, L.M., Chen, F., and Macosko, E.Z. (2019). Slide-seq: A scalable technology
for measuring genome-wide expression at high spatial resolution. Science *363*, 1463–1467.

15.   Stickels, R.R., Murray, E., Kumar, P., Li, J., Marshall, J.L., Di Bella, D.J., Arlotta, P.,
Macosko, E.Z., and Chen, F. (2021). Highly sensitive spatial transcriptomics at near-cellular
resolution with Slide-seqV2. Nat. Biotechnol. *39*, 313–319.

16.   Marx, V. (2021). Publisher Correction: Method of the Year: spatially resolved
transcriptomics. Nat. Methods *18*, 219.

17.   Waylen, L.N., Nim, H.T., Martelotto, L.G., and Ramialison, M. (2020). From whole-mount
to single-cell spatial assessment of gene expression in 3D. Commun Biol *3*, 602.

18.   Palla, G., Fischer, D.S., Regev, A., and Theis, F.J. (2022). Spatial components of molecular
tissue biology. Nat. Biotechnol. https://doi.org/10.1038/s41587-021-01182-1.

19.   Cable, D.M., Murray, E., Zou, L.S., Goeva, A., Macosko, E.Z., Chen, F., and Irizarry, R.A.
(2021). Robust decomposition of cell type mixtures in spatial transcriptomics. Nat.
Biotechnol., 1–10.

20.   Kleshchevnikov, V., Shmatko, A., Dann, E., Aivazidis, A., King, H.W., Li, T., Elmentaite,
R., Lomakin, A., Kedlian, V., Gayoso, A., et al. (2022). Cell2location maps fine-grained cell
types in spatial transcriptomics. Nat. Biotechnol. *40*, 661–671.

21.   Mages, S., Moriel, N., Avraham-Davidi, I., Murray, E., Watter, J., Chen, F., Rozenblatt-
Rosen, O., Klughammer, J., Regev, A., and Nitzan, M. (2023). TACCO unifies annotation
transfer and decomposition of cell identities for single-cell and spatial omics. Nat.
Biotechnol., 1–9.

22.   Lopez, R., Li, B., Keren-Shaul, H., Boyeau, P., Kedmi, M., Pilzer, D., Jelinski, A., Yofe, I.,
David, E., Wagner, A., et al. (2022). DestVI identifies continuums of cell types in spatial
transcriptomics data. Nat. Biotechnol. *40*, 1360–1369.

23.   Biancalani, T., Scalia, G., Buffoni, L., Avasthi, R., Lu, Z., Sanger, A., Tokcan, N.,
Vanderburg, C.R., Segerstolpe, Å., Zhang, M., et al. (2021). Deep learning and alignment of
spatially resolved single-cell transcriptomes with Tangram. Nat. Methods *18*, 1352–1362.

24.   Cancer Genome Atlas Network (2012). Comprehensive molecular characterization of human
colon and rectal cancer. Nature *487*, 330–337.

25. Kwong, L.N., and Dove, W.F. (2009). APC and its modifiers in colon cancer. Adv. Exp. Med.
Biol. *656*, 85–106.

26. Fearon, E.R. (2011). Molecular genetics of colorectal cancer. Annu. Rev. Pathol. *6*, 479–507.

27. Fearon, E.R., and Vogelstein, B. (1990). A genetic model for colorectal tumorigenesis. Cell
61, 759–767.

28. Chen, B., Scurrah, C.R., McKinley, E.T., Simmons, A.J., Ramirez-Solano, M.A., Zhu, X.,
Markham, N.O., Heiser, C.N., Vega, P.N., Rolong, A., et al. (2021). Differential pre-
malignant programs and microenvironment chart distinct paths to malignancy in human
colorectal polyps. Cell *184*, 6262-6280.e26.

29. Becker, W.R., Nevins, S.A., Chen, D.C., Chiu, R., Horning, A.M., Guha, T.K., Laquindanum,
R., Mills, M., Chaib, H., Ladabaum, U., et al. (2022). Single-cell analyses define a continuum
of cell state and composition changes in the malignant transformation of polyps to colorectal
cancer. Nat. Genet., 1–11.

30. Roper, J., Tammela, T., Cetinbas, N.M., Akkad, A., Roghanian, A., Rickelt, S., Almeqdadi,
M., Wu, K., Oberli, M.A., Sánchez-Rivera, F.J., et al. (2017). In vivo genome editing and
organoid transplantation models of colorectal cancer and metastasis. Nat. Biotechnol. *35*,
569–576.

31. Roper, J., Tammela, T., Akkad, A., Almeqdadi, M., Santos, S.B., Jacks, T., and Yilmaz, Ö.H.
(2018). Colonoscopy-based colorectal cancer modeling in mice with CRISPR–Cas9 genome
editing and organoid transplantation. Nat. Protoc. *13*, 217–234.

32. Golovko, D., Kedrin, D., Yilmaz, Ö.H., and Roper, J. (2015). Colorectal cancer models for
novel drug discovery. Expert Opin. Drug Discov. *10*, 1217–1229.

33. Gyllborg, D., Langseth, C.M., Qian, X., Choi, E., Salas, S.M., Hilscher, M.M., Lein, E.S.,
and Nilsson, M. (2020). Hybridization-based in situ sequencing (HybISS) for spatially
resolved transcriptomics in human and mouse brain tissue. Nucleic Acids Res. *48*, e112.

34. Hagerling, C., Gonzalez, H., Salari, K., Wang, C.-Y., Lin, C., Robles, I., van Gogh, M.,
Dejmek, A., Jirström, K., and Werb, Z. (2019). Immune effector monocyte–neutrophil
cooperation induced by the primary tumor prevents metastatic progression of breast cancer.
Proc. Natl. Acad. Sci. U. S. A. *116*, 21704–21714.

35. Arenberg, D.A., Keane, M.P., DiGiovine, B., Kunkel, S.L., Strom, S.R., Burdick, M.D.,
Iannettoni, M.D., and Strieter, R.M. (2000). Macrophage infiltration in human non-small-cell
lung cancer: the role of CC chemokines. Cancer Immunol. Immunother. *49*, 63–70.

36. Smith, P.M., and Garrett, W.S. (2011). The gut microbiota and mucosal T cells. Front.
Microbiol. *2*, 111.

37. La Manno, G., Soldatov, R., Zeisel, A., Braun, E., Hochgerner, H., Petukhov, V.,
1346   Lidschreiber, K., Kastriti, M.E., Lönnerberg, P., Furlan, A., et al. (2018). RNA velocity of
1347   single cells. Nature *560*, 494–498.

1348   38. Ziyad, S., and Iruela-Arispe, M.L. (2011). Molecular mechanisms of tumor angiogenesis.
1349   Genes Cancer *2*, 1085–1096.

1350   39. Zeng, Z.-S., Shu, W.-P., Cohen, A.M., and Guillem, J.G. (2002). Matrix metalloproteinase-7
1351   expression in colorectal cancer liver metastases: evidence for involvement of MMP-7
1352   activation in human cancer metastases. Clin. Cancer Res. *8*, 144–148.

1353   40. Li, D., Peng, Z., Tang, H., Wei, P., Kong, X., Yan, D., Huang, F., Li, Q., Le, X., Li, Q., et al.
1354   (2011). KLF4-mediated negative regulation of IFITM3 expression plays a critical role in
1355   colon cancer pathogenesis. Clin. Cancer Res. *17*, 3558–3568.

1356   41. Kitamura, T., Kometani, K., Hashida, H., Matsunaga, A., Miyoshi, H., Hosogi, H., Aoki, M.,
1357   Oshima, M., Hattori, M., Takabayashi, A., et al. (2007). SMAD4-deficient intestinal tumors
1358   recruit CCR1+ myeloid cells that promote invasion. Nat. Genet. *39*, 467–475.

1359   42. Haber, A.L., Biton, M., Rogel, N., Herbst, R.H., Shekhar, K., Smillie, C., Burgin, G., Delorey,
1360   T.M., Howitt, M.R., Katz, Y., et al. (2017). A single-cell survey of the small intestinal
1361   epithelium. Nature *551*, 333–339.

1362   43. Biton, M., Haber, A.L., Rogel, N., Burgin, G., Beyaz, S., Schnell, A., Ashenberg, O., Su, C.-
1363   W., Smillie, C., Shekhar, K., et al. (2018). T Helper Cell Cytokines Modulate Intestinal Stem
1364   Cell Renewal and Differentiation. Cell *175*, 1307-1320.e22.

1365   44. Smillie, C.S., Biton, M., Ordovas-Montanes, J., Sullivan, K.M., Burgin, G., Graham, D.B.,
1366   Herbst, R.H., Rogel, N., Slyper, M., Waldman, J., et al. (2019). Intra- and Inter-cellular
1367   Rewiring of the Human Colon during Ulcerative Colitis. Cell *178*, 714-730.e22.

1368   45. Welch, J.D., Kozareva, V., Ferreira, A., Vanderburg, C., Martin, C., and Macosko, E.Z.
1369   (2019). Single-Cell Multi-omic Integration Compares and Contrasts Features of Brain Cell
1370   Identity. Cell *177*, 1873-1887.e17.

1371   46. van Neerven, S.M., de Groot, N.E., Nijman, L.E., Scicluna, B.P., van Driel, M.S., Lecca,
1372   M.C., Warmerdam, D.O., Kakkar, V., Moreno, L.F., Vieira Braga, F.A., et al. (2021). Apc-
1373   mutant cells act as supercompetitors in intestinal tumour initiation. Nature *594*, 436–441.

1374   47. Flanagan, D.J., Pentinmikko, N., Luopajärvi, K., Willis, N.J., Gilroy, K., Raven, A.P.,
1375   Mcgarry, L., Englund, J.I., Webb, A.T., Scharaw, S., et al. (2021). NOTUM from Apc-mutant
1376   cells biases clonal competition to initiate cancer. Nature *594*, 430–435.

1377   48. Clevers, H. (2006). Wnt/beta-catenin signaling in development and disease. Cell *127*, 469–
1378   480.

1379   49. Folkman, J. (2002). Role of angiogenesis in tumor growth and metastasis. Semin. Oncol. *29*,
1380   15–18.

50. Lasry, A., Zinger, A., and Ben-Neriah, Y. (2016). Inflammatory networks underlying
colorectal cancer. Nat. Immunol. *17*, 230–240.

51. Fleming, M., Ravula, S., Tatishchev, S.F., and Wang, H.L. (2012). Colorectal carcinoma:
Pathologic aspects. J. Gastrointest. Oncol. *3*, 153–173.

52. Winkler, J., Abisoye-Ogunniyan, A., Metcalf, K.J., and Werb, Z. (2020). Concepts of
extracellular matrix remodelling in tumour progression and metastasis. Nat. Commun. *11*, 1–
19.

53. Drev, D., Harpain, F., Beer, A., Stift, A., Gruber, E.S., Klimpfinger, M., Thalhammer, S.,
Reti, A., Kenner, L., Bergmann, M., et al. (2019). Impact of Fibroblast-Derived SPARC on
Invasiveness of Colorectal Cancer Cells. Cancers *11*.
https://doi.org/10.3390/cancers11101421.

54. Burden, R.E., Gormley, J.A., Jaquin, T.J., Small, D.M., Quinn, D.J., Hegarty, S.M., Ward,
C., Walker, B., Johnston, J.A., Olwill, S.A., et al. (2009). Antibody-mediated inhibition of
cathepsin S blocks colorectal tumor invasion and angiogenesis. Clin. Cancer Res. *15*, 6042–
6051.

55. Basu, S., Cheriyamundath, S., Gavert, N., Brabletz, T., Haase, G., and Ben-Ze'ev, A. (2019).
Increased expression of cathepsin D is required for L1-mediated colon cancer progression.
Oncotarget *10*, 5217–5228.

56. Spencer, J., and Sollid, L.M. (2016). The human intestinal B-cell response. Mucosal
Immunol. *9*, 1113–1124.

57. Sautès-Fridman, C., Petitprez, F., Calderaro, J., and Fridman, W.H. (2019). Tertiary lymphoid
structures in the era of cancer immunotherapy. Nat. Rev. Cancer *19*, 307–325.

58. Mendez, M.G., Kojima, S.-I., and Goldman, R.D. (2010). Vimentin induces changes in cell
shape, motility, and adhesion during the epithelial to mesenchymal transition. FASEB J. *24*,
1838–1851.

59. Lu, M.-H., Huang, C.-C., Pan, M.-R., Chen, H.-H., and Hung, W.-C. (2012). Prospero
homeobox 1 promotes epithelial-mesenchymal transition in colon cancer cells by inhibiting
E-cadherin via miR-9. Clin. Cancer Res. *18*, 6416–6425.

60. Oliemuller, E., Newman, R., Tsang, S.M., Foo, S., Muirhead, G., Noor, F., Haider, S.,
Aurrekoetxea-Rodríguez, I., Vivanco, M. dM, and Howard, B.A. (2020). SOX11 promotes
epithelial/mesenchymal hybrid state and alters tropism of invasive breast cancer cells. Elife
9, e58374.

61. Marjanovic, N.D., Hofree, M., Chan, J.E., Canner, D., Wu, K., Trakala, M., Hartmann, G.G.,
Smith, O.C., Kim, J.Y., Evans, K.V., et al. (2020). Emergence of a High-Plasticity Cell State
during Lung Cancer Evolution. Cancer Cell *38*, 229-246.e13.

62. Puram, S.V., Tirosh, I., Parikh, A.S., Patel, A.P., Yizhak, K., Gillespie, S., Rodman, C., Luo,
C.L., Mroz, E.A., Emerick, K.S., et al. (2017). Single-Cell Transcriptomic Analysis of
Primary and Metastatic Tumor Ecosystems in Head and Neck Cancer. Cell *171*, 1611-
1624.e24.

63. Schwab, R.H.M., Amin, N., Flanagan, D.J., Johanson, T.M., Phesse, T.J., and Vincan, E.
(2018). Wnt is necessary for mesenchymal to epithelial transition in colorectal cancer cells.
Dev. Dyn. *247*, 521–530.

64. DiMeo, T.A., Anderson, K., Phadke, P., Fan, C., Perou, C.M., Naber, S., and Kuperwasser,
C. (2009). A novel lung metastasis signature links Wnt signaling with cancer cell self-renewal
and epithelial-mesenchymal transition in basal-like breast cancer. Cancer Res. *69*, 5364–
5373.

65. Penninger, J.M., and Crabtree, G.R. (1999). The Actin Cytoskeleton and Lymphocyte
Activation. Cell *96*, 9–12.

66. Cang, Z., Zhao, Y., Almet, A.A., Stabell, A., Ramos, R., Plikus, M.V., Atwood, S.X., and
Nie, Q. (2023). Screening cell–cell communication in spatial transcriptomics via collective
optimal transport. Nat. Methods *20*, 218–228.

67. Che, L.-H., Liu, J.-W., Huo, J.-P., Luo, R., Xu, R.-M., He, C., Li, Y.-Q., Zhou, A.-J., Huang,
P., Chen, Y.-Y., et al. (2021). A single-cell atlas of liver metastases of colorectal cancer
reveals reprogramming of the tumor microenvironment in response to preoperative
chemotherapy. Cell Discovery *7*, 1–21.

68. Zheng, X., Song, J., Yu, C., Zhou, Z., Liu, X., Yu, J., Xu, G., Yang, J., He, X., Bai, X., et al.
(2022). Single-cell transcriptomic profiling unravels the adenoma-initiation role of protein
tyrosine kinases during colorectal tumorigenesis. Signal Transduction and Targeted Therapy
7, 1–14.

69. Khaliq, A.M., Erdogan, C., Kurt, Z., Turgut, S.S., Grunvald, M.W., Rand, T., Khare, S.,
Borgia, J.A., Hayden, D.M., Pappas, S.G., et al. (2022). Correction: Refining colorectal
cancer classification and clinical stratification through a single-cell atlas. Genome Biol. *23*,
156.

70. Joanito, I., Wirapati, P., Zhao, N., Nawaz, Z., Yeo, G., Lee, F., Eng, C.L.P., Macalinao, D.C.,
Kahraman, M., Srinivasan, H., et al. (2022). Single-cell and bulk transcriptome sequencing
identifies two epithelial tumor cell states and refines the consensus molecular classification
of colorectal cancer. Nat. Genet. *54*, 963–975.

71. Guinney, J., Dienstmann, R., Wang, X., de Reyniès, A., Schlicker, A., Soneson, C., Marisa,
L., Roepman, P., Nyamundanda, G., Angelino, P., et al. (2015). The consensus molecular
subtypes of colorectal cancer. Nat. Med. *21*, 1350–1356.

72. West, N.R., Hegazy, A.N., Owens, B.M.J., Bullers, S.J., Linggi, B., Buonocore, S., Coccia,
M., Görtz, D., This, S., Stockenhuber, K., et al. (2017). Oncostatin M drives intestinal

inflammation and predicts response to tumor necrosis factor-neutralizing therapy in patients
with inflammatory bowel disease. Nat. Med. *23*, 579–589.

73.   West, N.R., Murray, J.I., and Watson, P.H. (2013). Oncostatin-M promotes phenotypic
changes associated with mesenchymal and stem cell-like differentiation in breast cancer.
Oncogene *33*, 1485–1494.

74.   Smigiel, J.M., Parameswaran, N., and Jackson, M.W. (2017). Potent EMT and CSC
phenotypes are induced by oncostatin-M in pancreatic cancer. Mol. Cancer Res. *15*, 478–488.

75.   Hara, T., Chanoch-Myers, R., Mathewson, N.D., Myskiw, C., Atta, L., Bussema, L.,
Eichhorn, S.W., Greenwald, A.C., Kinker, G.S., Rodman, C., et al. (2021). Interactions
between cancer cells and immune cells drive transitions to mesenchymal-like states in
glioblastoma. Cancer Cell *39*, 779-792.e11.

76.   de Sousa e Melo, F., Kurtova, A.V., Harnoss, J.M., Kljavin, N., Hoeck, J.D., Hung, J.,
Anderson, J.E., Storm, E.E., Modrusan, Z., Koeppen, H., et al. (2017). A distinct role for
Lgr5+ stem cells in primary and metastatic colon cancer. Nature *543*, 676–680.

77.   Schepers, A.G., Snippert, H.J., Stange, D.E., van den Born, M., van Es, J.H., van de Wetering,
M., and Clevers, H. (2012). Lineage tracing reveals Lgr5+ stem cell activity in mouse
intestinal adenomas. Science *337*, 730–735.

78.   André, T., Shiu, K.-K., Kim, T.W., Jensen, B.V., Jensen, L.H., Punt, C., Smith, D., Garcia-
Carbonero, R., Benavides, M., Gibbs, P., et al. (2020). Pembrolizumab in Microsatellite-
Instability–High Advanced Colorectal Cancer. N. Engl. J. Med. *383*, 2207–2218.

79.   Kuraguchi, M., Wang, X.-P., Bronson, R.T., Rothenberg, R., Ohene-Baah, N.Y., Lund, J.J.,
Kucherlapati, M., Maas, R.L., and Kucherlapati, R. (2006). Adenomatous polyposis coli
(APC) is required for normal development of skin and thymus. PLoS Genet. *2*, e146.

80.   Johnson, L., Mercer, K., Greenbaum, D., Bronson, R.T., Crowley, D., Tuveson, D.A., and
Jacks, T. (2001). Somatic activation of the K-ras oncogene causes early onset lung cancer in
mice. Nature *410*, 1111–1116.

81.   Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., Ng, L.L.,
Palmiter, R.D., Hawrylycz, M.J., Jones, A.R., et al. (2009). A robust and high-throughput Cre
reporting and characterization system for the whole mouse brain. Nat. Neurosci. *13*, 133–140.

82.   Marino, S., Vooijs, M., van Der Gulden, H., Jonkers, J., and Berns, A. (2000). Induction of
medulloblastomas in p53-null mutant mice by somatic inactivation of Rb in the external
granular layer cells of the cerebellum. Genes Dev. *14*, 994–1004.

83.   el Marjou, F., Janssen, K.-P., Chang, B.H.-J., Li, M., Hindie, V., Chan, L., Louvard, D.,
Chambon, P., Metzger, D., and Robine, S. (2004). Tissue-specific and inducible Cre-
mediated recombination in the gut epithelium. Genesis *39*, 186–193.

84. Stoeckius, M., Zheng, S., Houck-Loomis, B., Hao, S., Yeung, B.Z., Mauck, W.M., 3rd,
Smibert, P., and Satija, R. (2018). Cell Hashing with barcoded antibodies enables
multiplexing and doublet detection for single cell genomics. Genome Biol. *19*, 224.

85. Qian, X., Harris, K.D., Hauling, T., Nicoloutsopoulos, D., Muñoz-Manchado, A.B., Skene,
N., Hjerling-Leffler, J., and Nilsson, M. (2019). Probabilistic cell typing enables fine mapping
of closely related cell types in situ. Nat. Methods *17*, 101–106.

86. Li, B., Gould, J., Yang, Y., Sarkizova, S., Tabaka, M., Ashenberg, O., Rosen, Y., Slyper, M.,
Kowalczyk, M.S., Villani, A.-C., et al. (2020). Cumulus provides cloud-based data analysis
for large-scale single-cell and single-nucleus RNA-seq. Nat. Methods *17*, 793–798.

87. Wolf, F.A., Angerer, P., and Theis, F.J. (2018). SCANPY: large-scale single-cell gene
expression data analysis. Genome Biol. *19*, 15.

88. Wolock, S.L., Lopez, R., and Klein, A.M. (2019). Scrublet: Computational Identification of
Cell Doublets in Single-Cell Transcriptomic Data. Cell Syst *8*, 281-291.e9.

89. Aran, D., Looney, A.P., Liu, L., Wu, E., Fong, V., Hsu, A., Chak, S., Naikawadi, R.P.,
Wolters, P.J., Abate, A.R., et al. (2019). Reference-based analysis of lung single-cell
sequencing reveals a transitional profibrotic macrophage. Nat. Immunol. *20*, 163–172.

90. Tabula Muris Consortium, Overall coordination, Logistical coordination, Organ collection
and processing, Library preparation and sequencing, Computational data analysis, Cell type
annotation, Writing group, Supplemental text writing group, and Principal investigators
(2018). Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris. Nature *562*,
367–372.

91. Pedregosa, F., Varoquaux, G., Gramfort, A., and Michel, V. Scikit-learn: Machine Learning
in Python. 12 (85): 2825–2830. : http://jmlr. org/papers/v12 ….

92. Polański, K., Young, M.D., Miao, Z., Meyer, K.B., Teichmann, S.A., and Park, J.-E. (2020).
BBKNN: fast batch alignment of single cell transcriptomes. Bioinformatics *36*, 964–965.

93. Rao, J.N., and Wang, J.-Y. (2010). Intestinal Architecture and Development (Morgan &
Claypool Life Sciences).

94. Puram, S.V., Parikh, A.S., and Tirosh, I. (2018). Single cell RNA-seq highlights a role for a
partial EMT in head and neck cancer. Mol Cell Oncol *5*, e1448244.

95. Liu, J., Lichtenberg, T., Hoadley, K.A., Poisson, L.M., Lazar, A.J., Cherniack, A.D.,
Kovatich, A.J., Benz, C.C., Levine, D.A., Lee, A.V., et al. (2018). An Integrated TCGA Pan-
Cancer Clinical Data Resource to Drive High-Quality Survival Outcome Analytics. Cell *173*,
400-416.e11.

96. Davidson-Pilon, C. (2019). lifelines: survival analysis in Python. The Journal of Open Source
Software *4*, 1317.

97. Fazilaty, H., Brügger, M.D., Valenta, T., Szczerba, B.M., Berkova, L., Doumpas, N.,
Hausmann, G., Scharl, M., and Basler, K. (2021). Tracing colonic embryonic transcriptional
profiles and their reactivation upon intestinal damage. Cell Rep. *36*, 109484.

1526

Figure 1. A single cell atlas of healthy colon and dysplastic lesions in mouse.

A. Study overview. **B.** Major cell subsets of healthy colon and dysplastic lesions. 2D embedding

of 48,115 single cell profiles colored by cluster (top, legend) or annotated cell type (bottom,

legend). **C,D.** Changes in cell composition in dysplastic tissues. C. 2D embedding of single cell

profiles, showing only the cells in each condition state, subsampled to equal numbers of cells per

condition state, colored by cluster (same legend as in B (top)). D. Proportion of cells (y axis) of

each cell type in each sample (x axis). **E.** multiplex RNA *in situ* analysis. Representative images

of Cartana analysis of normal colon (left) and AV lesions (right) colored by cell type assignment

(same as **Supplementary item 2A**).

1539

Figure 2. Composition and cell intrinsic expression program changes in dysplastic epithelial

cells.

A-C. Compositional changes in epithelial cells in dysplastic tissue. A. 2D embedding of epithelial

cell profiles colored by clusters (legend). Cluster Epi06: potential doublets (**Methods**). B.

Proportion of cells out of all epithelial cells (y axis) of each epithelial cell subset in each sample

(x axis). C. Fraction of expressing cells (dot size) and mean expression in expressing cells (dot

color) of marker genes (columns) for each cluster (rows). **D-E.** Use of epithelial cell programs

changes in dysplastic tissue. D. Weights (x axis) of each of the 20 top ranked genes (y axis) for

each program. E. Proportion of program weights summed over all epithelial cells (y axis) in each

sample (x axis). **F.** Stem cell program 16 is induced in epithelial cells in dysplastic tissue. Scaled

log-normalized expression (color bar) of the top 100 genes differentially expressed between cells

from normal colon and from dysplastic (AV and AKPV) across the 10,812 cells that accounted for

90% of program 16's expression across all epithelial cells (columns). Selected program genes are

marked.

1556

Figure 3. Altered cell type neighborship in CRC.

A. Cell type distributions *in situ*. Slide-seq pucks of dysplastic (top) and normal (bottom) tissue colored by TACCO assignment of cell labels (legend, light grey: low quality beads) (x and y axis: spatial coordinates in μm). **B,C.** Cell type neighborships in normal and dysplastic colon tissue. Short-range (up to 20μm) neighborhood enrichment (Z score, color bar) vs. a background of spatially random annotation assignments for each pair of cell annotations (rows, columns) in normal (A) and dysplastic (B) tissue.

1564

1565

Figure 4. **Three cellular neighborhoods associated with tumor progression.**

A. Spatial regions. Slide-seq pucks of AV (top) and normal (bottom) mouse colon colored by

TACCO regions (legend, light gray: low quality beads) (x and y axis are spatial coordinates in

μm). **B,C.** Enrichment and depletion of cell subsets and epithelial programs across different

regions. Significance (FDR, color bar, two-sided Welch's t-test on CLR-transformed

compositions) of enrichment (red) or depletion (blue) of specific cell subsets (rows, B) or epithelial

cell programs (rows, C) in the different regions defined by TACCO (columns) as well as all normal

("N (ref.)", leftmost column) and AV ("AV (ref.)", leftmost column) samples. **D.** TACCO defined

regions preferentially relate to normal or AV tissue. Significance (FDR, color bar, two-sided

Welch's t-test on CLR-transformed compositions) of enrichment (red) or depletion (blue) of each

TACCO defined region (rows) in normal ("N vs. rest") and AV ("AV vs. N") samples (columns).

E. TACCO reveals normal colon architecture. Left: Slide-seq puck of normal mouse colon colored

by TACCO region annotations (legend) (x and y axis: spatial coordinates (μm)). Right: Main

epithelial expression programs enriched in each region (FDR<6.3 10^{-4}, two-sided Welch's t test

on CLR-transformed compositions) except region 2 (muscularis), which is characterized by non-

epithelial (stromal) cell types. **F.** Expression signatures of cells in normal regions 3,5,10 and 12.

Scaled log-normalized expression of the top 20 differentially expressed genes (rows) for each bead

(columns) in the region. **G,H**. Malignant-like regions. G. Slide-seq pucks of two AV lesions

colored by TACCO annotations of malignant-like regions 6, 8 and 11. H. Scaled log-normalized

expression of the top 20 differentially expressed genes (rows) of each bead (left, columns) in the

region; or epithelial (middle left), immune (middle right) or stromal (right) fractions of beads

(columns) in regions 6, 8 and 11 in dysplastic lesions. **I,J.** Epithelial cell subsets and programs

associated with "malignant-like", "normal-like" and normal tissues. Significance (FDR, color bar,

two-sided Welch's t-test on CLR-transformed compositions) of enrichment (red) or depletion

(blue) of epithelial cell programs (I, rows) or epithelial, immune and stromal cell subsets (J, rows)

in different tissue types (columns) based on Slide-seq or scRNA-seq ("sc") samples. **K.** Inferred

interaction pathways. Enrichment (FDR) in AV *vs*. normal tissue of corresponding "sender" (x

axis) and "receiver" (y axis) (aggregated over ligand-receptor pairs) pathways (dots). Red/blue:

pathways significantly enriched in AV/normal samples and (light red: only for either sender or

receiver). The top 5 enriched pathways (in each direction) are labeled.

1596

Figure 5. Mouse tumor regions associated with tumor progression in human colorectal tumors.

A. Expression profiles characterizing mouse regions are recapitulated in human tumors. Significance (FDR, color bar, two-sided Welch's t-test on CLR-transformed compositions) of enrichment (red) or depletion (blue) of region-associated epithelial, immune or stromal profiles (rows) compared between normal, MMRp, or MMRd samples (columns). **B,C.** Mouse regions capture malignant features in human tumors. B. Top left: First (PC1, x axis) and second (PC2, y axis) principal components of mouse region scores of mouse and human epithelial pseudo-bulk

samples. Top right: PC1 loadings (x axis) of each mouse region score (y axis). Bottom: PC1 values

(box plots show mean, quartiles, and whiskers for the full data distribution except for outliers

outside 1.5 times the interquartile range (IQR)) for each type of mouse or human sample (x axis).

C. Significance (FDR, color bar, two-sided Welch's t-test) of enrichment (red) or depletion (blue)

of region-associated profile scores (rows) in normal and dysplastic samples (columns) in human

or mouse. **D,E.** Expression of malignant like regions 6 and 11 in tumors are associated with PFI

(D) and OS (E) in human patients. Kaplan-Meier PFI (e, n = 662[95]) or OS (f, n = 662[95]) analysis

of human bulk RNA-seq cohort stratified by malignant-like region profile scores.

Figure S1. **Marker genes, cell states and cell types in the healthy and dysplastic mouse**

colon atlas.

A. Cell-type expression signatures. Scaled log-normalized expression (color bar) of the top 20

differentially expressed genes (rows) in cells (columns) from each cell type. **B,C.** Distinct

condition states and compartments. 2D embedding of all single cell profiles (dots) colored by either

condition state (B) or compartment (C). **D.** Changes in cell composition in dysplastic lesions.

Significance (FDR, color bar, two-sided Welch's t-test on CLR-transformed compositions) of

enrichment (red) or depletion (blue) of each cell subset (rows) between samples from different

conditions (columns). **E.** Changes in cell composition between healthy and dysplastic tissue.

Proportion of cells (y axis) from each cell subset (x axis) out of all cells in each sample (x axis).

F-H. Multiplex *in situ* RNA profiles (Cartana) reproduce scRNA-seq and Slide-Seq findings. F.

Proportion of RNA molecules (y axis) attributed to each cell type in each sample (x axis) in two

Cartana experiments, with different gene panels. G. Change in cell proportions in AV vs. normal

samples (difference of CLR-transformed cell type fractions, x and y axes) for each cell type (dots,

color) based on scRNA-seq, Slide-Seq or Cartana (axis labels). Error bars: bootstrapped standard

error of the mean (normal n=8, 16, 12, 16 and AV n=11, 24, 12, 12 for scRNA-seq, Slide-seq,

Cartana V1, Cartana V2). H. Change in marker gene expression (dots) in AV vs. normal samples

(difference of CLR-transformed gene fractions, x and y axes) based on scRNA-seq, Slide-Seq or

Cartana (axis labels), for genes measured by all 3 methods. Genes with the 8 maximal and minimal

expression ranks across methods are labeled. Error bars: bootstrapped standard error of the mean

(normal n=8, 16, 12, 16 and AV n=11, 24, 12, 12 for scRNA-seq, Slide-seq, Cartana V1, Cartana

V2).

Figure S2. Compositional and cell intrinsic changes in stromal and immune cells.

A-C. Immune cell subsets and composition. A. 2D embedding of immune cell profiles colored by

clusters (as in B). B. Proportion of cells out of all immune cells (y axis) of each immune cell subset

in each sample (x axis). C. Scaled log-normalized expression (color bar) of the top 20 differentially

expressed genes (rows) in cells (columns) from each immune cell subset (color bar on top). **D-F**.

Monocyte and macrophage cell subsets and composition. D. 2D embedding of monocyte and

macrophage cell profiles colored by clusters (as in E). E. Proportion of cells out of all monocytes

and macrophages (y axis) of each monocyte and macrophage cell subset in each sample (x axis).

F. Scaled log-normalized expression (color bar) of the top 20 differentially expressed genes (rows)

in cells (columns) from each monocyte and macrophage cell subset (color bar on top). **G-I.** T/NK

cell subsets and composition. G. 2D embedding of T/NK cell profiles colored by clusters (as in

H). H. Proportion of cells out of all T/NK cells (y axis) of each T/NK cell subset in each sample

(x axis). I. Scaled log-normalized expression (color bar) of the top 20 differentially expressed

genes (rows) in cells (columns) from each T/NK cell subset (color bar on top). **J.** Enrichment of

IL17+ γδT cells (TNK05) and depletion of CD8+ γδT cells (TNK01) in dysplastic lesions.

Significance (FDR, color bar, two-sided Welch's t-test on CLR-transformed compositions) of

enrichment (red) or depletion (blue) of each T/NK cell subset (rows) between samples from

different conditions (columns). **K**. T/NK cell compositions is similar in human and mouse. Left:

2D embedding of T/NK cell composition profiles of human and mouse samples colored by sample

type (legend) (**STAR Methods**). Right: Similarity of T/NK cell composition (enrichment z-scores)

in the 2D embedding between each set of samples (rows, columns). **L-N.** Stromal cell subsets and

composition. L. 2D embedding of stromal cell profiles colored by clusters (legend). M. Proportion

of cells out of all stromal cells (y axis) of each stromal cell subset in each sample (x axis). N.

Scaled log-normalized expression (color bar) of the top 20 differentially expressed genes (rows)

in cells (columns) from each stromal cell subset (color bar on top). **O**. Enrichment of vascular

endothelial cells (Endo01) and depletion of myofibroblasts (Fibro02) in dysplastic lesions.

Significance (FDR, color bar, two-sided Welch's t test on CLR-transformed compositions) of

enrichment (red) or depletion (blue) of each stromal cell subset (rows) between samples from

different conditions (columns). **P.** Increased VEGFA expression in monocyte-macrophage

populations with dysplasia. Distribution of expression (y axis, log1p(counts)) of *VegfA* in

monocytes and macrophages from different conditions (x axis).

1670

Figure S3. Changes in T/NK and epithelial cell differentiation in dysplastic lesions.

A. T/NK cell differentiation. 2D embedding of T/NK cell profiles from normal tissues (left) and

dysplastic lesions (right), colored by cell subset. Streamlines: averaged and projected RNA

velocities. **B,C.** Proliferating stem-like cells give rise to an expanded stem-like compartment and

differentiated-like tumor cells in dysplastic lesions. 2D embedding of epithelial cell profiles from

normal tissues (B, left and C, top) and dysplastic lesions (B, right, and C, bottom), colored by cell

subset (B), or by expression of epithelial programs (C). Streamlines: averaged and projected RNA

velocities. Outlined dots (B): Proliferative cells (cells with more than 50% program weight in the

proliferation programs #3 and #11).

1680

Figure S4. Changes in cell composition and expression programs usage in dysplastic epithelium.

A-D. Changes in epithelial cell composition in dysplastic tissues. A. 2D embedding of all single cell epithelial profiles showing only the profiles (dots) of cells from each condition state, subsampled to equal numbers of cells per condition state, colored by cluster. B. PCA captures differentiation status and lineage of epithelial clusters. First (PC1, x axis) and second (PC2, y axis) principal components of the expression of epithelial clusters. C. Mean expression (dot color) and fraction of expressing cells (dot size) of epithelial cell marker genes[97] (columns) in each cluster (rows). D. Significance (FDR, color bar, two-sided Welch's t test on ALR-transformed compositions with all non-tdTomato counts used as reference compartment) of enrichment (red) or depletion (blue) of tdTomato expression in cells from AKPVT samples between every pair of epithelial clusters. **E.** Dysplastic secretory like (Epi05) and immune cells express tumor-related genes. Distribution of expression (y axis, log1p(counts)) of different marker genes in cells from each epithelial/immune cell cluster (x axis). **F-L**. Epithelial gene programs. 2D embedding of all epithelial cells colored by the weight of each program (color bar) and the expression of selected program genes (color bar). **M.** Epithelial program characteristics of normal and dysplastic colon. Heatmap significance (FDR, color bar, two-sided Welch's t test on CLR-transformed compositions) of enrichment (red) or depletion (blue) of each epithelial program (rows) between normal vs. dysplastic tissues (columns).

Figure S5. Conservation of cellular composition and expression programs between mouse

and human scRNA-seq data.

A,B. Conservation of general and subtype-specific human and mouse epithelial expression

programs. A. Pearson correlation coefficients (color) between program-specific expression

profiles (**Methods**) of human (all samples, left, MMRd, middle and MMRp, right)) (rows) and

mouse programs (columns). B. Mouse epithelial program enrichments in mouse and human tumor

and normal samples. Significance (FDR, color bar, two-sided Welch t-test on CLR-transformed

compositions) of enrichment (red) or depletion (blue) of mouse epithelial program expression

(rows) in sample classes from human or mouse (columns). **C,D.** Human-mouse conservation of

cell type and program associations. Pearson correlation coefficients (color) of the CLR-

transformed cell type (C) or epithelial program (D) compositions across samples in mouse (left)

or human (right) single cell data. In (D), data are hierarchically clustered for the "not normal"

mouse case (AV and AKPV) and this ordering is applied to all other panels.

1721

1722

Figure S6. **Spatial distributions of cells and programs across regions.**

A-C. Slide-seq quality controls. Slide-seq pucks of AV (top) and normal (bottom) mouse colon

colored by number of UMIs (A) or of genes (B) per bead. (x and y axis: spatial coordinates (μm)).

C. H&E staining of sections adjacent to those used for Slide-seq. **D.** Selected marker gene

expression. Slide-seq pucks from a normal (top) and AV (bottom) sample, colored by marker gene

detected per bead. **E,F.** Spatial mapping of cell types and programs yields comparable composition

to scRNA-seq. Distribution of the proportion (y axis) of contributions to each cell type (D) or

program (E; based on fractional annotations) in cells (for scRNA-seq; "sc") or beads (for Slide-

seq; "spatial"; based on fractional annotations) in samples from normal (N) or AV (AV) tissue.

G,H. Distinct cell types and programs associated with AV and normal colon. Significance (FDR,

color bar, two-sided Welch's t test on CLR-transformed compositions) of enrichment (red) or

depletion (blue) of cell types (F, rows) or epithelial programs (G, rows) in normal (N) or AV (AV)

tissues based on Slide-seq ("spatial") data or scRNA-seq ("sc").

Figure S7. Distinct cellular layout in normal and AV tissues.

A,B. Neighborship analysis is robust to the distance and number of randomizations. Neighborship enrichment (z-scores, color) *vs*. a background of spatially random annotation assignments for each pair of cell annotations (rows, columns) in normal (top) and AV (bottom) samples at varying distance (a, left: ≤40μm; right: ≤60 μm) or number of permutations (B, left: 5, right: 50). **C,D.** Reproducible spatial arrangement of epithelial cells in normal colon in Slide-seq and Cartana V2. Co-occurrence (y axis) of normal epithelial cell types (color) at different distances (x axis) from different central normal epithelial cell type (annotated on top) in Slide-Seq (C) and Cartana (D) data. **E.** Decreased spatial order of cell types in AV vs. normal tissue. Distribution of enrichment z-values (x axis) of cluster-cluster interactions (as shown in **Figure 3B,C**) in AV lesions (orange) and normal colon (blue). **F.** Epithelial program neighborships in AV tissue. Short-range (≤20μm) neighborship enrichment z-scores (color) *vs*. a background of spatially random annotation assignments for each pair of epithelial program annotations (rows, columns) in AV lesions. **G,H.** Reproducible spatial arrangement of dysplasia-associated immune cells relative to endothelial cells in Slide-seq and Cartana V2. Co-occurrence (y axis) of dysplasia-associated monocyte or granulocyte cells (color) with endothelial cells at different distances (x axis) in Slide-Seq (G) and Cartana (H) data.

Figure S8. Cellular neighborhoods.

A. Distinct characteristic region distributions in healthy and AV tissue. Proportion of UMIs assigned to each region (y axis) in each Slide-seq puck (x axis). **B.** β-catenin expression in AV lesions. Representative images (i) of β-catenin IHC on sections of AV lesions showing either nuclear (*) or cytoplasmic (**) staining pattern (x10 magnification), along with high magnification (x20, ii) of the region highlighted in (i), and H&E staining (iii) of a serial section showing a region proximal to the region in (i). (M-muscularis, L-lumen, dashed line: border between areas of nuclear and cytoplasmic staining of β-catenin). **C,D.** Spatial relations between the regions. Proportion of beads (y axis) of each region category (color code) at different distances (x axis) from region 2 (C, muscularis), 6 (D, left), 8 (D, middle) or 11 (D, right). **E,F.** Fraction of expressing cells (dot size) and mean expression per celltype (E, dot color) or per region (F, dot color) of marker genes (columns) across cell types in region 6 (E, rows) or regions 6, 8 and 11 (F, rows). **G.** Cell type neighborships in different malignant regions. Short-range (≤20μm) neighborship enrichment z-scores (color) *vs*. a background of spatially random annotation assignment for each pair of cell type annotations (rows, columns) in malignant-like regions 6, 8 and 11 within AV lesions. **H.** Spatial organization of endothelial cells and pericytes based on Cartana multiplex *in situ* RNA analysis. Co-occurrence (y axis) of Cdh5 (top) or Pecam1 (middle) -expressing (endothelial) cells or Pdgfrb (bottom) -expressing cells (pericytes) at different distances (x axis) from each central cell type (color code), based on expression of a specific gene (color bar). **I.** Cell-cell interaction pathways enriched in different regions. Enrichment (FDR) in regions 2 (muscularis, left), 6 (malignant-like, inflammation, middle) or 11 (malignant-like, EMT, right) of corresponding "sender" (x axis) and "receiver" (y axis) (aggregated over ligand-receptor pairs) pathways (dots). Red/blue: pathways significantly enriched/depleted in region (light red/blue: only

1784   enriched/depleted for either sender or receiver). The top 5 enriched pathways (in each direction)

1785   are labeled.

Figure S9. Transfer of mouse spatial region expression profiles to human patient data.

A. Expression profiles characterizing mouse regions in human and mouse samples. Significance

(FDR, color bar, two-sided Welch's t test on CLR-transformed compositions) of enrichment (red)

or depletion (blue) of mouse region-associated epithelial (left), immune (middle) or stromal

profiles (right) in pucks from normal colon and dysplastic lesions, in the same pucks but after

mapping the region annotation to itself (consistency check), in mouse single cell data after

mapping region annotation from mouse pucks, and in human single cell data after mapping region

annotation from mouse pucks. **B**. PC1 values (y axis; box plots show mean, quartiles, and whiskers

for the full data distribution except for outliers outside 1.5 times the interquartile range (IQR)) in

a PCA of region scores of mouse and human samples (x axis) sorted by malignant status and

colored by status (top, legend) or study (bottom, legend). **C**. Spatial regions profiles associated

with different CMS classes. Significance (Benjamini-Hochberg FDR, color bar, two-sided

Welch's t test) of enrichment (red) or depletion (blue) of each region profile (rows) in pseudo bulk

profiles of human tumor samples classified in each CMS class (columns). **D,E**. Expression of

malignant-like regions 6 and 11 in tumors is associated with PFI and OS in MMRd and MMRp

patients. Kaplan-Meier PFI (f, MMRd n = 189[95], MMRp n = 447[95]) or OS (g, MMRd n = 189[95],

MMRp n = 447[95]) analysis of human bulk RNA-seq cohort stratified by malignant-like region

profile scores.

1811

Supplementary item 1. Correspondence between scRNA-seq, Slide-seq, and Cartana data

A,B. Agreement in cell type proportions. Inferred cell type proportions (y and x axis; CLR-transformed cell type fractions) for each cell type (dot color) in normal (A) and AV (B) samples in scRNA-seq, Slide-Seq and Cartana data (axis labels). Pearson's r: top left corners. Error bars: bootstrapped standard error of the mean ((A) n=8, 16, 12, 16 and (B) n=11, 24, 12, 12 for scRNA-seq, Slide-seq, Cartana V1, Cartana V2). **C,D**. Agreement in gene expression levels. Measured expression (y and x axis; CLR-transformed gene fractions) for each gene (dot) measured by all three methods in normal (A) and AV (B) samples in scRNA-seq, Slide-Seq and Cartana data (axis labels). Pearson's r: top left corners. Error bars: bootstrapped standard error of the mean ((C) n=8, 16, 12, 16 and (D) n=11, 24, 12, 12 for scRNA-seq, Slide-seq, Cartana V1, Cartana V2). Genes with the 8 maximal and minimal expression ranks across methods are labeled.

1831

Supplementary item 2. Dysplasia-associated cells in the tumor microenvironment.

A,B. Representative images of multiplex RNA analysis in normal colon (left) and AV lesions

(right) colored by cell type assignment (A, the same is **Figure 1E**) and expression of marker genes

(B) for Mono02 (Arg1, Hilpda, Nos2, Cd274, Vegfa, Trem1; cyan), Mono03 (Osm, Ifi204 and

Thbs1; red), granulocytes (S100a8; yellow), as well as Sell (green), Ccr2 (magenta), Ptprc (blue)

and Cd14 (gray).

1838

Supplementary item 3. TLS-like structures in AV **lesions by Cartana multiplex** *in situ* **RNA**

profiles.

A. Proportion of RNA molecules (y axis) attributed to each cell type (top) or gene (bottom) in

TLS-like regions (x axis) in Cartana V2 multiplex *in situ* RNA experiment. Genes which constitute

at least 3% of all measured transcripts in any of the identified follicular structures are noted

(bottom). **B.** Representative images from TLS-like regions showing the distribution of cell type

(top row) and of cell type markers (bottom four rows; Cd14, Cd68 (monocytes, magenta), Cd3d,

Cd3e, Cd3g (T cells, yellow), and Cd79, Jchain, Igkc (B cells, dark cyan)).

Supplementary item 4. Ligand-receptor analysis.

A,B. Cell-cell interaction pathways enriched in normal and AV tissue and regions. Significance (FDR, color bar, two-sided Mann-Whitney U test) of enrichment (red) or depletion (blue) of expression of "sender" (left) and "receiver" (right) cell-cell communication pathways (rows) (aggregted from ligand-receptor pairs) in normal *vs*. AV samples (A, columns), or regions (B, columns). **C,D.** Split power analysis for cell-cell communication pathway analysis. Significance of enrichment ($\log_{10}$(FDR), y-axis) for receiver (solid lines) and sender (dashed line) for each of the top 5 enriched communication pathways (color) in each iteration for the spatial split of the Slide-Seq pucks into spatially disconnected and separated patches (x axis), for either normal vs. AV tissue enrichment (C) or for specific regions (D).