1 **A novel knock sideways approach for *Plasmodium* secreted proteins reveals a surprising**

2 **insensitivity to vacuolar depletion of PTEX component HSP101**

8 Manuel A Fierro[1], Tahir Hussain[1], Liam J Campin[2] and Josh R Beck[1,2*]

11 [1]Department of Biomedical Sciences, Iowa State University, Ames, IA, 50011, USA

12 [2]Roy J. Carver Department of Biochemistry, Biophysics and Molecular Biology, Iowa State

13 University, Ames, IA, 50011, USA

16 [*]Corresponding author: jrbeck@iastate.edu

Abstract

Obligate intracellular malaria parasites depend on protein secretion to penetrate, subvert, and exit their host cells. Despite this key reliance on the secretory pathway, none of the conditional mutagenesis tools available in *Plasmodium spp.* exploit the unique trafficking features of secreted proteins. Here we report knockER, a novel DiCre-mediated knock sideways approach that sequesters secreted proteins in the ER via conditional fusion of a KDEL ER-retrieval sequence to the target protein C-terminus. We applied knockER to a diverse set of reporters and endogenous *P. falciparum* and *P. berghei* proteins targeted to the rhoptries, dense granules, parasite vacuole and apicoplast and show conditional ER sequestration is not generally toxic, enabling loss-of-function studies. Taking advantage of the unique ability to redistribute secreted protein pools from their terminal destination to the ER, we employed knockER to study components of the PTEX translocon in an attempt to separate HSP101 vacuolar function in protein export from a recently proposed role in cargo recognition at the ER. Strikingly, while KDEL-fusion produced similar levels of ER retrieval for both HSP101 and the translocon adaptor PTEX150, parasite growth and effector export were completely unaffected by HSP101 retention while ER retrieval of PTEX150 produced a lethal export defect, indicating vacuolar HSP101 levels are uniquely maintained in excess. Intriguingly, redistribution of HSP101 to the ER did not further sensitize parasites to TetR-DOZI-mediated knockdown compared to parasites containing normal vacuolar levels of HSP101, suggesting PV-localized function does not fully account for HSP101 contribution to parasite fitness and consistent with an important role for HSP101 in the ER. Collectively, our work provides a novel tool for dissecting secreted protein function with sub-compartmental resolution that should be widely amenable to genetically tractable eukaryotes.

Introduction

Protein secretion is a fundamental biological process of all living cells. In eukaryotes this is accomplished by a dedicated secretory pathway that begins with import into the Endoplasmic Reticulum (ER) for initial folding and post-translational modification, followed by sorting in the Golgi and post-Golgi compartments. As an obligate intracellular parasite, secreted proteins play a critical role in the biology of the malaria parasite, *Plasmodium*. Through the secretory pathway, proteins are trafficked into both the plasma membrane and several organelles, including a relict plastid called the apicoplast that is critical for parasite metabolism (1). During schizogony, another subset of proteins is packaged into several distinct secretory organelles that are subsequently discharged to facilitate host cell attachment and penetration (2). Invagination of the host membrane by the parasites during invasion generates an intracellular microenvironment called the parasitophorous vacuole (PV) which forms the interface for host-parasite interactions (3). Resident PV proteins secreted into the vacuole perform critical functions at this interface, including nutrient/waste exchange and protein export (4). The later process enables parasites to deploy a battery of secreted effector proteins across the PV membrane (PVM) and into the host compartment, dramatically remodeling the erythrocyte to meet nutritional demands and avoid host defenses (5, 6). Remarkably, it is estimated that up to 10% of the parasite genome is devoted to encoding exported effectors, further highlighting the central importance of protein secretion to this parasitic lifestyle (7-11).

Despite their divergent position as early branching eukaryotes, *Plasmodium spp.* and other apicomplexan parasites have retained many of the canonical features of the secretory pathway (12). Like other eukaryotes, entry into the secretory pathway generally begins with co-translational import of proteins containing a signal peptide or transmembrane domain(s) into the ER via the Sec61 complex, followed by sorting from the Golgi to internal compartments, the PV or beyond into the host (13). ER-resident proteins that escape the ER via bulk flow in COPII vesicles are returned to the ER in COPI vesicles by ERD2 surveillance, which binds a C-terminal

retrieval sequence ("KDEL" and variants) in the *cis* Golgi and releases it in the ER in a pH-

dependent manner (14-16). Similar to other eukaryotes, ER retention of secreted reporters or

chimeric proteins can be achieved in *Plasmodium spp.* by a C-terminal KDEL or functional

variants (17-22).

In this present work, we developed and characterized a novel DiCre-based knock

sideways tool called knockER that enables conditional fusion of KDEL to the C-terminus of

secreted proteins, subjecting the target protein to ERD2 surveillance in the *cis* Golgi to cause its

retrieval to the ER. Using this approach, we show successful conditional retrieval of five

endogenous *Plasmodium* proteins (ClpB1, RON6, EXP2, PTEX150 and HSP101) that traffic

through the ER to diverse compartments including the apicoplast, rhoptries, dense granules, and

the PV. Importantly, conditional ER retrieval of both highly expressed reporters and endogenously

tagged proteins does not generally impact fitness, indicating knockER-mediated phenotypes are

not the indirect result of an integrated stress response, thus making this system suitable for protein

loss-of-function studies. Indeed, we show that knockER reproduces known loss-of-function

phenotypes previously established by other conditional knockdown strategies. We further apply it

to study loss of ClpB1 and RON6 function for the first time, revealing an essential apicoplast

function for the former and suggesting that RON6 is dispensable during the blood stage. Lastly,

we took advantage of the ability to redistribute secreted protein pools to interrogate a recently

proposed ER function for the PTEX translocon component HSP101 in addition to its role in the

PV (23). We reveal a surprising insensitivity to ER retrieval of HSP101 in contrast to the other

PTEX components, indicating HSP101 is maintained in excess in the PV. Interestingly, depletion

of HSP101 PV levels does not hypersensitize parasites to HSP101 knockdown, providing support

for an HSP101 function in the early secretory pathway in addition to its role in protein translocation

across the PVM in the assembled PTEX complex (24-26). These results establish knockER as a

novel tool to study secreted proteins and provide the first functional genetic evidence for an extra-

vacuolar HSP101 function, highlighting the unique ability of knockER to resolve function between

compartments for dually localized proteins.

Results

Conditional ER retrieval of a secreted reporter does not impact parasite fitness

To facilitate conditional ER retrieval of a protein of interest, we designed a strategy based

on the DiCre system where a gene of interest is C-terminally fused to a 3xFLAG tag and a stop

codon flanked by *loxP* sequences imbedded in small introns (27) which, upon addition of

rapamycin, undergo recombination via activation of the dimerizable Cre recombinase (27, 28).

Excision of the loxP-intervening sequence removes the 3xFLAG tag and stop codon and brings

into frame a 3xHA tag followed by the ER-retention sequence KDEL and a stop codon (Figure

1A). While the ER plays a crucial role in quality control and is sensitive to added stress (29, 30),

several previous studies have exogenously expressed various KDEL or SDEL fusion proteins in

P. falciparum and *P. berghei* without apparent impact on parasite fitness (21, 31-37). To directly

ascertain whether ER retrieval of a highly expressed secreted protein impacted parasite fitness,

we first appended the knockER system to an mNeonGreen (mNG) reporter and fused this to the

exp2 5' UTR and signal peptide, enabling expression from the strong *exp2* promoter and entry

into the ER followed by default secretion into the PV (38-40). This assembly was placed under

control of the anhydrotetracycline (aTc)-responsive TetR-DOZI-aptamers system to prevent

expression until aTc is present in the culture (Figure 1A) (41, 42). To combine the Bxb1 integrase

system with DiCre, we first introduced a DiCre expression cassette at the benign *pfs47* locus (27)

in the *P. falciparum* NF54[attB] strain (43) (Figure S1) and then integrated the mNG-knockER

reporter construct at the *attB* site in the *cg6* locus on chromosome 6 (43) in this NF54[attB:DiCre] line.

Upon treatment with rapamycin, parasites efficiently excised the knockER cassette (kER),

resulting in the expected tag switching from FLAG to HA (Figure 1B,C). Examination of untreated

parasites by live microscopy showed a peripheral mNG signal, consistent with secretion to the

PV (Figure 1D). In contrast, following treatment with rapamycin the mNG signal was dramatically

altered to an intracellular, perinuclear pattern that co-localized with ER Tracker, confirming

retention in the ER (Figure 1D). To determine the level of ER retention achieved by KDEL fusion,

we quantified the distribution of mNG fluorescence between the parasite periphery (PV) and

interior and observed substantial re-localization following KDEL fusion, indicating robust retrieval

by ERD2 (Figure 1E, mean internal mNG 42.54 ± 6.0% in DMSO vs 84.66 ± 6.9% in Rap).

Importantly, ER retention of mNG had no impact on parasite growth (Figure 1F) showing that

knockER can successfully retrieve a secreted reporter to the ER without impacting parasite

fitness.

knockER reveals Golgi-dependent trafficking and essential apicoplast functionality of

ClpB1

Next, we applied knockER to an endogenous gene for conditional loss-of-function

mutagenesis. We choose ClpB1, a poorly characterized, nuclear-encoded AAA+ class 1

Clp/HSP100 chaperone that is targeted to the apicoplast (44, 45). As apicoplast ablation can be

chemically rescued with isopentenyl pyrophosphate (IPP) (46), we reasoned that IPP

supplementation would provide a simple validation that any loss of parasite fitness is a direct

effect of sequestering ClpB1 away from its site of action and not an indirect result of ER stress.

ClpB1 contains both a signal peptide for ER entry and a transit peptide for apicoplast targeting

and localization of ClpB1 to this organelle was previously confirmed by both microscopy and

proteomic analyses (36, 45). Using CRISPR/Cas9 editing, we fused mNG to the C-terminus of

ClpB1 followed by the kER cassette (Figure 2A). As expected, IFA demonstrated co-localization

with the apicoplast marker ACP1, confirming ClpB1 localization to this organelle (Figure 2B).

Treatment with rapamycin resulted in efficient excision (Figure 2C) and switching to the 3xHA-

KDEL tag produced a shift in migration by Western blot consistent with an increase in molecular

weight of ~14 kDa in the rapamycin treated samples, indicating that the N-terminal transit peptide

(residues 24-152, 13.9 kDa) is no longer cleaved in the ClpB1-KDEL fusion (Figure 2D). Since

this maturation event takes place in the apicoplast (47), this suggested that ClpB1 was no longer

being trafficked to this organelle. Indeed, live microscopy showed a dramatic re-localization of

mNG from puncta to a perinuclear distribution that co-localized with ER Tracker, indicative of

successful ER retrieval (Figure 2E).

While its functional importance has not been directly tested, ClpB1 is expected to be

essential for blood-stage survival based on saturating insertional mutagenesis in *P. falciparum*

and pooled knockout screens in *P. berghei* (48, 49). Indeed, ER retention of ClpB1 was lethal and

resulted in delayed parasite death (Figure 2F), consistent with an essential role in the apicoplast

(50, 51). Moreover, supplementation with IPP fully rescued parasite growth, confirming the

observed phenotype was apicoplast-specific and not due to a disruption of ER homeostasis

following ClpB1 retention (Figure 2F). Rapamycin-treated parasites supplemented with IPP

showed dispersed, vesicular ACP1 signal consistent with plastid disruption, further indicating

ClpB1 is required for apicoplast maintenance (Figure 2G). Taken together, these results show for

the first time that ClpB1 is an indispensable component of the apicoplast machinery and illustrate

that knockER can be used for protein loss-of-function studies in *P. falciparum*.

knockER induces ER retrieval of rhoptry proteins

Having demonstrated that knockER is suitable to generate loss-of-function phenotypes by

protein mis-localization to the ER, we next tested this knock sideways approach on an expanded

set of endogenous soluble and membrane proteins that traffic through the secretory pathway to

diverse destinations, including secretory organelles and the PV/PVM. We began by attempting to

retrieve RON6, a soluble rhoptry-neck protein that is injected into the PV during invasion. Attempts

predating CRISPR technology to disrupt *Pfron6* or truncate its Cysteine-rich C-terminus were

unsuccessful (52). More recently, saturating transposon mutagenesis in *P. falciparum* recovered

a single insertion within an intron (48) expected to generate a truncation at residue 713 of 950,

leaving most of the protein intact (Figure 3A). Curiously, RON6 orthologs in rodent malaria

parasites lack the cysteine-rich region which accounts for the C-terminal ~23% of *P. falciparum*

RON6, roughly corresponding to the truncated region in the insertional mutant. As RON6 was not

targeted in the PlasmoGEM data set, its suspected essentiality remains uncertain (49). To directly

test the importance of RON6, we generated RON6-kER parasites, which underwent efficient

excision upon rapamycin treatment (Figure 3B). Rapamycin treatment resulted in re-localization

of RON6 from the rhoptries to the ER and produced a minor but reproducible growth defect (Figure

3C,D). These results suggest that RON6 is dispensable in the blood stage, although we cannot

exclude the possibility that small amounts of RON6-KDEL may escape ERD2 surveillance at

levels sufficient to mediate a critical function. Additionally, the minor fitness defect that results

from RON6-KDEL fusion could be the indirect result of modest ER stress induced by RON6

retention. Nonetheless, these findings clearly demonstrate that knockER efficiently retrieves

secretory proteins synthesized during the brief period of merozoite formation at the terminal phase

of intraerythrocytic development.

ER retention of EXP2 produces a lethal defect independent to loss of PVM transport

function

Having successfully retained several soluble proteins in the ER using knockER, we next

tested conditional retrieval of a membrane protein. Solute permeation across the PVM is mediated

by a nutrient-permeable channel (53, 54) whose identity was recently tied to EXP2, a PVM protein

broadly conserved among vacuole-dwelling apicomplexans (55, 56). EXP2 contains a signal

peptide followed by an N-terminal amphipathic helix and oligomerizes to form a heptameric pore

in the PVM (26, 57, 58). It is currently unknown how pore formation is specifically constrained to

the PVM to avoid perforating other membranes along the secretory pathway or the parasite

plasma membrane. While not structurally homologous to the EXP2 pore, bacterial α-helical pore

forming toxins are functionally analogous and provide a possible model for EXP2 pore formation.

These proteins are typically maintained in a monomeric state that shields a membrane-spanning

amphipathic helix, enabling soluble trafficking until contact with components of a target membrane

triggers conformational rearrangement, leading to membrane insertion and oligomerization (59).

If EXP2 traffics in a similar configuration, then its C-terminus will enter the secretory

pathway lumen and be sensitive to ER-retrieval by KDEL fusion. Indeed, we endogenously tagged

EXP2 with a version of the kER cassette lacking mNG and found that KDEL fusion disrupted

EXP2 trafficking to the PV and resulted in ER retention, confirming its C-terminus resides in the

secretory pathway lumen (Figure 4A-D and S2). Notably, ER retrieval of EXP2 caused rapid

parasite death at earlier time points than post-transcriptional conditional knockdown of EXP2 (55,

60) (Figure 4E). To test whether this was solely due to loss of EXP2 transport functions in the PV

and not an indirect result of retaining EXP2 in the ER, we generated parasites expressing a

second copy of EXP2-mNG-kER from the *attB* locus under the control of its own promoter, leaving

the endogenous EXP2 locus unaltered (Figure S2). Retrieval of this second copy of EXP2 to the

ER also resulted in parasite death, indicating that EXP2 retention in the ER is toxic independent

of compromised transport functions at the PVM (Figure 4F,G). In contrast, ER retrieval of a second

copy of EXP2 lacking the amphipathic helix rescued parasite death (Figure 4H,I and S2). As

expected, prior to KDEL fusion the ΔTM version of EXP2 localized to the PV and the digestive

vacuole (due to endocytosis of PV proteins along with RBC cytosol); surprisingly, however, some

cells also showed EXP2ΔTM-mNG export into the RBC cytoplasm (arrow, Figure 4H). The

requirement for the amphipathic helix in EXP2-kER toxicity suggests that increasing EXP2 dwell

time in the early secretory pathway may trigger an unfolded protein response or cause aberrant

EXP2 pore formation in the ER membrane. Thus, this may provide a new tool for studying EXP2

pore formation but also indicates knockER may not be appropriate for loss-of-function studies of

certain membrane proteins.

ER retention of PTEX150 but not HSP101 results in loss of PTEX function *in vitro*

In malaria parasites, the nutrient-permeable channel has been additionally functionalized

by the flange-like adaptor PTEX150 which docks the AAA+ chaperone HSP101 onto EXP2 to

form the _Plasmodium_ Translocon of Exported proteins (PTEX), transforming EXP2 into a protein-

conducting pore to translocate effector proteins across the PVM and into the erythrocyte (24-26,

55, 61). Interestingly, HSP101 shows a dual localization to both the PV and the ER as opposed

to EXP2 and PTEX150, which localize exclusively to the PV and can form a sub-complex

independent of HSP101 (23, 38, 62, 63). We verified this unique ER localization of HSP101 by

protease protection assays in parasite lines expressing fluorescently tagged versions of each

PTEX core component, confirming an internal, perinuclear pool of HSP101 but not EXP2 or

PTEX150 (Figure S3). This arrangement suggests that in addition to powering translocation in

the assembled PTEX complex, HSP101 may also function upstream of the vacuole, possibly

initiating cargo selection early in the secretory pathway (23). While several HSP101 conditional

knockdown (23, 25) or inactivation (24) strategies have demonstrated an essential role in protein

export, it is difficult to resolve distinct compartmental functions with these mutants given that they

impact the total protein pool and have no sub-compartmental resolution. In contrast, knockER

provides a strategy to resolve potential HSP101 functions across distinct compartments by

redistributing the target protein pool from the PV to the ER.

In an attempt to separate HSP101 ER function from its role in the PV, we generated a kER

fusion to the endogenous copy of HSP101. In parallel, we also generated a PTEX150-kER line,

reasoning that since HSP101 and PTEX150 have similar expression timing/levels and are

exclusively involved in protein export but not nutrient uptake (55), PTEX150 would provide a

control for knockER inactivation of PTEX activity independent of ER function. As expected, the

PTEX150-KDEL fusion was efficiently retrieved to the ER, producing a lethal growth defect and

accompanying block in protein export (Figure 5A-E), similar to _glmS_ ribozyme-mediated

knockdown of PTEX150 (25). Importantly, ER retention of a second copy of PTEX150-kER in

parasites where the endogenous PTEX150 locus was unmodified did not impact growth,

demonstrating that the phenotype observed upon ER retention of the endogenously tagged

protein is a direct result of functional disruption of PTEX150 (Figure S4A-D).

Prior to knockER induction, HSP101-mNG-3xFLAG showed a small but significantly more

internal distribution than PTEX150-mNG-3xFLAG, consistent with the unique HSP101 ER pool

(Figure S4E, 24 hrs post DMSO treatment: $39.41\pm11.00\%$ internal HSP101 vs $35.58\pm7.41\%$

internal PTEX150, P = 0.0024, unpaired t test). Unexpectedly, although the rapamycin-induced

HSP101-KDEL fusion increased ER localization similar to PTEX150 (Figure 5F-H), parasite

growth was completely unaffected and protein export was not detectibly impacted (Figure 5I,J),

sharply contrasting with the lethal export defect observed with HSP101 depletion or inactivation

mutants (23-25). Indeed, HSP101-kER parasites could be cultured indefinitely following excision

and displayed sustained ER retention of HSP101 (Figure 5H and S4E,F, mean Rap internal

HSP101 $58.82\pm9.58\%$ at 24 hours vs $54.82\pm9.84\%$ at 7 days). As an orthogonal approach to

monitor protein redistribution by KDEL fusion, we released infected RBCs with saponin to

permeabilize the PVM and quantified fluorescence via flow cytometry with or without proteinase

K treatment. Again, we observed an increase in protected fluorescence following rapamycin

treatment, consistent with redistribution of HSP101 to the ER (Figure S5A). Importantly, PCR and

Western blot indicated that rapamycin-treated parasites had undergone complete excision with

the pre-excised FLAG-tagged version of HSP101 no longer detectible (Figure 5F,G).

Furthermore, both the mNG-FLAG and mNG-HA-KDEL tagged versions of HSP101 migrated as

a single band at the expected size for the full-length fusion proteins with no apparent breakdown

products, ensuring that the mNG signal distribution observed by live microscopy reflected an

intact HSP101 fusion (Figure 5G, S5C and S7B).

While a large fraction of PTEX150 and HSP101 was clearly ER-retained following

knockER induction, some peripheral signal was still apparent after rapamycin treatment (Figure

5C,H). To determine whether the observed phenotypic differences between HSP101 and

PTEX150 were due to a difference in sensitivity to ERD2 surveillance, we quantified the internal

mNG fluorescence signal in HSP101- and PTEX150-kER parasites following KDEL fusion. In

synchronized parasites, a similar increase in internal signal was observed in both lines at 24 hours

post Rap treatment (Figure S4E, mean Rap internal HSP101 58.82±9.58% vs 56.32±18.56% for

PTEX150). Since a comparison could not be made with PTEX150-kER parasites at 7 days due

to parasite death by the second cycle follow excision, we compared HSP101-kER parasites with

PTEX150-kER[attB] control parasites 7 days after excision and again observed a comparable

internal signal redistribution (Figure S4F, mean Rap internal HSP101 54.82±9.84% vs

61.75±16.27% for PTEX150 at 7 days). As differential ER retrieval cannot account for parasite

insensitivity to HSP101-kER, these results indicate vacuolar HSP101 levels are uniquely

maintained in excess from what is required for PTEX function at the PVM.

To determine the amount of HSP101 need to support parasite growth, we combined

knockER with the TetR-DOZI-aptamers system to generate a HSP101-kER line that also allows

for titration of HSP101 expression with aTc (Figure 6A). Like the original HSP101-kER line,

rapamycin-treated HSP101-kER[TetR-DOZI] parasites exhibited ER retention of HSP101 without

impacting parasite growth or export of representative PEXEL and PNEP proteins (Figure 6B-D).

However, HSP101 depletion following aTc washout resulted in a robust block in protein export

and parasite death (Figure 6D and S5B,C), consistent with previous HSP101 knockdown or

inactivation studies (23-25). To determine the level of total HSP101 reduction sufficient to impact

growth, we grew synchronized parasites at a range of aTc concentrations following DMSO or

rapamycin treatment and measured mNG fluorescence via flow cytometry in parallel with parasite

growth (Figure 6E,F). Interestingly, rapamycin-treated parasites displayed a small but

reproducible increase in total HSP101, consistent with a reduction in HSP101 turnover by

endocytosis from the PV to the digestive vacuole following redistribution to the ER (Figure 6E).

More importantly, we determined that a ~70% reduction in total HSP101 levels was necessary to

impact parasite growth, showing that a relatively small amount of HSP101 is needed for survival

in vitro (Figure 6E,F, 5 nM aTc), suggesting that redistribution of HSP101 by knockER does not

meet the threshold required to impact HSP101 PV function and parasite fitness.

If HSP101 functions exclusively in the PV, then knockER-mediated depletion of vacuolar

HSP101 levels should hypersensitize parasites to HSP101 knockdown. To test this, we measured

the dose-response to aTc in DMSO- or rapamycin-treated parasites and strikingly observed no

shift in the EC50, indicating substantial redistribution of HSP101 from the PV to the ER does not

hypersensitize parasites to HSP101 knockdown (Figure 6G). This suggests that the PV-localized

function of HSP101 does not fully account for its role in parasite fitness, implying a distinct

important role in the ER. Taken together, our results provide the first functional genetic evidence

for an extra-vacuolar function of HSP101, consistent with the revised model of protein export (23).

ER retention of HSP101 is lethal *in vivo*

Lack of observable phenotypes *in vitro* can sometimes be masked by the rich resources

available in culturing conditions (64, 65). In addition, exported effectors critical for survival in the

vertebrate host may be dispensable in culture (11). Indeed, export-related defects that escape

detection *in vitro* can be exposed *in vivo* where parasites have to contend with host defenses and

more limited nutrient access (66, 67). To expose any impact on export by ER retention of HSP101

masked during *in vitro* culture, we tagged HSP101 and PTEX150 with kER in a *P. berghei* rodent

malaria DiCre parasite and evaluated the impact of ER retrieval on parasite survival during mouse

infection (Figure S6A). Similar to *P. falciparum* knockER mutants, rapamycin treatment *ex vivo*

produced efficient excision and ER retrieval of PbPTEX150 and PbHSP101 (Figure 6A-D). After

overnight *ex vivo* culture to allow for tag switching, equal numbers of DMSO or rapamycin treated

parasites were IV injected into naïve mice. Strikingly, while DMSO controls became patent by day

two post-injection, neither KDEL-fused PbPTEX150 or PbHSP101 parasites appeared over the

course of 10 days before the experiment was terminated (Figure 7E,F and S6B). That ER-retrieval

of PbHSP101 compromises function sufficiently to prohibit survival in the mammalian host

reinforces the critical role of HSP101 in the PV and indicates that KDEL-mediated depletion of

HSP101 from the *P. falciparum* PV does not sufficiently reduce PTEX activity to impact parasite

fitness under less taxing *in vitro* culture conditions. Collectively, these results provide support for

a distinct HSP101 activity in the ER upstream of its function in PVM translocation at the assembled

PTEX complex.

Discussion

Several genetic tools are now available for protein loss-of-function studies in *Plasmodium*

spp. (68), including knock sideways for conditional sequestration of a target protein away from its

normal site of function (69). While this system is not currently amenable to parasite proteins that

enter the secretory pathway, conditional control of secretory traffic has been achieved in *P.*

falciparum by ligand-mediated alteration of the N-terminal transit peptide that controls apicoplast

targeting, enabling redirection of plastid proteins into the default secretory pathway (70).

Additionally, conditional ER release systems have been developed in model eukaryotes based

on ER retrieval machinery (71). In contrast, knockER provides a simple, DiCre-mediated knock

sideways strategy that exploits the ERD2/KDEL receptor (KDELR) system for conditional ER

retrieval, providing a generalizable approach for interrogating secreted proteins in genetically

tractable eukaryotes.

In this study, we demonstrate knockER can be used for trafficking and loss-of-function

studies. The ability to determine whether a protein traffics from the ER through the *cis* Golgi is an

important consideration for nuclear-encoded apicoplast proteins where Golgi-dependent and -

independent trafficking routes have been reported (33, 35, 72). While these apicoplast trafficking

studies were carried out with reporters and chimeras, knockER allows for monitoring *cis* Golgi

transit of endogenous proteins. The sensitivity of ClpB1 to KDEL-mediated ER retrieval

demonstrates a Golgi-dependent trafficking route to the apicoplast. Moreover, we show for the

first time that this AAA+ chaperone essential for apicoplast maintenance. ClpB1 is one of four

Clp/HSP100 chaperones encoded by *Plasmodium spp.*, three of which localize to the parasite

plastid while the fourth, HSP101 (previously known as ClpB2) localizes to the PV and ER (45).

The other apicoplast-targeted members of this quartet in include the nuclear-encoded ClpC, which

mediates regulated proteolysis together with the serine protease ClpP (73, 74), and ClpM, now

the only uncharacterized member of this group and one of the few proteins encoded on the

apicoplast genome. Similar to the well-studied function of its closest orthologs (75), ClpB1 likely

acts as a disaggregase that, together with ClpC/P and ClpM, forms a critical proteostasis system

required to maintain and regulate the apicoplast proteome (76).

ER perturbations can trigger an integrated stress response leading to cell arrest or death

(77, 78). *Plasmodium* possesses one of the widely conserved ER stress sensors, the PERK-

eIF2α pathway which halts protein translation and flux through the secretory pathway (19, 29, 79-

81), and knockER experiments should be controlled for indirect effects. Importantly, conditional

ER retrieval of proteins produced from their endogenous locus or by heterologous expression was

not generally toxic, enabling loss-of-function studies with knockER. The single notable exception

was EXP2, which rapidly killed parasites when retained in the ER independent to loss of its PVM

transport function. EXP2-KDEL toxicity was dependent on the amphipathic helix that forms the

membrane-spanning region in the oligomeric EXP2 pore (26), suggesting that accumulation of

EXP2 in the ER might trigger an unfolded protein response or lead to premature pore formation

in the early secretory pathway. More work is required to determine if this outcome is specific to

EXP2 or a general problem for unnatural ER-retrieval of membrane proteins. Indirect fitness

defects aside, retrieval of EXP2-KDEL demonstrates that its C-terminus enters the secretory

pathway lumen, consistent with a soluble EXP2 trafficking state that precedes membrane

integration at the PVM. Regardless of the basis for EXP2-kER toxicity, these experiments

highlight that knockER can also be used to investigate membrane topology. While ER-resident

type I membrane proteins utilize a different mechanism for retrieval as their C-termini face the

cytosol of the cell (82, 83), ERD2/KDELR can retrieve membrane proteins whose C-termini face

the lumen (84) and thus knockER should be able to distinguish the topology of single and multi-

pass transmembrane domain-containing proteins.

A unique feature of knockER is the ability to redistribute secreted proteins from their

terminal destination to the ER, providing a method to resolve potential HSP101 functions between

the PV and ER. We tagged both HSP101 and the translocon adaptor PTEX150 with knockER,

reasoning that ER retrieval of either protein would produce a lethal export defect but that

differential impacts on exported cargo trafficking might be exposed. Unexpectedly, ER retrieval of

HSP101 had no impact on *P. falciparum* growth or export of representative PEXEL and PNEP

proteins *in vitro*. Similar PV depletion of PTEX150 produced the anticipated lethal export block,

revealing that HSP101 PV levels are uniquely maintained in excess. As a AAA+ ATPase, HSP101

is the only enzyme in the PTEX core complex and its activity may exceed what is required to

support parasite fitness in the less demanding *in vitro* culture environment, as previously seen for

the aspartic protease Plasmepsin 5 that licenses proteins for export in the ER (30, 85). Indeed,

while critical to avoiding host defenses *in vivo*, most exported proteins are dispensable during *in*

vitro culture (11). Not surprisingly, ER retrieval of *P. berghei* HSP101 abolished parasite survival

in the mouse host, reinforcing the established PV function of HSP101 in effector translocation at

the PVM. In contrast, depletion of HSP101 PV levels in cultured *P. falciparum* likely impacts

translocon kinetics or export of specific proteins in ways that are not problematic for growth *in*

vitro and thus difficult to detect.

Intriguingly, redistribution of HSP101 from the PV to the ER did not hypersensitize

parasites to HSP101 knockdown, suggesting that the PV-localized function of HSP101 does not

fully account for its contribution to parasite fitness. These results provide support for an important,

additional HSP101 function upstream of the PV, presumably in cargo recognition at the ER and/or

guiding export-destine proteins to PTEX, as recently proposed (23). Future work dissecting the

specific extra-vacuolar role(s) of HSP101 may resolve the long-standing question of how early

events in this pathway that mark proteins for export, such as PEXEL processing during ER entry,

are connected to cargo recognition and ultimate PVM translocation into the host cell at the

assembled PTEX complex.

Methods

Parasite culture. *P. falciparum* NF54[attB-DiCre] and derivatives were cultured in RPMI 1640 medium

supplemented with 27 mM sodium bicarbonate, 11 mM glucose, 0.37 mM hypoxanthine, 10 µg/ml

gentamicin and 0.5% Albumax I (Gibco). Parasites were maintained in deidentified, Institutional

Review Board-exempt RBCs obtained from the American National Red Cross. *P. berghei* ANKA

clone 2.34 and derivatives were maintained in Swiss Webster mice (Charles River). All

experiments involving rodents were reviewed and approved by the Iowa State University

Institutional Animal Care and Use Committee.

Plasmids and genetic modification of *P. falciparum* and *P. berghei*. All cloning was carried

out with NEBuilder HIFI DNA (NEB). Primer sequences are given in Table S1. To generate

NF54[attB-DiCre] parasites, a DiCre expression cassette was inserted into the *pfs47* gene in NF54[attB]

parasites (43) using a marker-free CRISPR/Cas9 editing strategy as described (27). Loss of the

endogenous *pfs47* locus was monitored with primers P1/2 and P3/4 and integration was

confirmed using primers P1/5 and P6/4. To confirm DiCre activity in this line, a loxP-flanked mNG

coding sequence with a 3xHA tag was PCR amplified using primers P19/20 and inserted between

AvrII and AflII in pLN-ENR-GFP cite (86), resulting in the plasmid JRB416. This plasmid was co-

transfected with pINT (86) into NF54[attB-DiCre] parasites and selected for with 2.5 µg/ml blasticidin-

S (BSD) 24-48hrs post-transfection.

To generate the kER fusion cassette, homology flanks targeting the 3' end of *exp2* were

amplified from pyPM2GT-EXP2-3xHA-GFP11 (55) using P21/22 and inserted between XhoI and

AvrII in pPM2GT-HSP101-3xFLAG (55). The loxPint sequence (87) was amplified with P23/24

and inserted immediately upstream of the 3xFLAG tag at AvrII in this vector. A partial loxPint

containing the loxP sequence and 3' portion of the intron followed by a 3xHA-KDEL tag was then

amplified with P25/26 and inserted into EagI, resulting in the plasmid JRB492. Finally, the mNG

coding sequencing was amplified from pyPM2GT-EXP2-mNG (88) with P27/28 and inserted into

AvrII in JRB492, resulting in plasmid JRB508.

To generate the SP-mNG-kER reporter, the *exp2* promoter and signal peptide were

amplified from parasite genomic DNA with P29/30 and the mNG-KER cassette was amplified from

JRB508 with P31/32 and these amplicons were assembled together in a second PCR reaction

with P29/32 and inserted into AscI/ApaI in pKD (42), resulting in the plasmid JRB543. For

expression of a second copy of EXP2-kER from the attB locus, the *exp2* promoter and full-length

gene were amplified from genomic DNA with P29/33 and inserted between AscI/BamHI in

JRB543, resulting in the plasmid JRB526. For expression of a second copy of EXP2-kER lacking

the amphipathic helix, a recoded portion of the *exp2* coding sequence beginning immediately after

the amphipathic helix (codon 72) was amplified from pEXP2^{apt} (55) and inserted at BamHI in

JRB543, resulting in the plasmid JRB544. These plasmids were each co-transfected with pINT

and selected with 2.5 μg/ml blasticidin-S as described above. To relieve translational repression

by TetR-DOZI, parasite cultures were supplemented with 500 nM aTc.

For generation of ClpB1-kER and RON6-kER lines, a Cas9 gRNA target was chosen just

downstream　　(*clpB1*,　　ATTTTTAGAAAAATATTATG)　　or　　just　　upstream　　(*ron6*,

TAATACATACAACGTACCAA) of the stop codon and the gRNA seed sequences were

synthesized as primers P34 and P35, respectively, and inserted into the AflII site of pAIO3 (38),

resulting in plasmids JRB143 and MAF22, respectively. To target the 3' end of clpB1 or ron6, a 5'

homology flank (up to but not including the stop codon) was amplified from genomic DNA using

P36/37 or P38/39, incorporating synonymous shield mutations in the gRNA target sites. A 3'

homology flank was amplified using P40/41 or P42/43. The corresponding 5' and 3' flank

amplicons were assembled in a second PCR reaction using P40/37 and P42/39 and inserted

between XhoI/AvrII in JRB508, resulting in plasmids MAF20 and MAF27, respectively. These

plasmids were linearized with AflII and co-transfected with the corresponding Cas9/gRNA plasmid

into NF54[attB-DiCre]. Selection was applied with 5 nM WR99120 24 hrs post-transfection and clonal

lines were isolated by limiting dilution after parasites returned from selection.

For generation of PTEX150-kER and HSP101-kER lines, flank assemblies targeting the

3' end of each gene were amplified using P44/45 from pyPM2GT-PTEX150-3xHA-GFP11 (55) or

P46/47 from pyPM2GT-HSP101-3xHA-GFP11 (55) and inserted between XhoI/AvrII in JRB508,

resulting in plasmids MAF16 and MAF17, respectively. These plasmids were linearized with AflII

and co-transfected with pAIO-PTEX150-CT-gRNA or pAIO-HSP101-CT-gRNA (55), selected with

5 nM WR99120 and cloned after parasites returned. For generation of EXP2-kER line, plasmid

JRB492 was linearized and co-transfected with pUF-Cas9-EXP2-CT-gRNA (55) and selected

with 5 nM WR99120. For generation of HSP101-kER[TetR-DOZI] lines, flank assembles were amplified

using P48/49 and inserted between AscI/BamHI in JRB526, resulting in plasmid MAF41. This

plasmid was linearized with AflII and co-transfected with pAIO-HSP101-CT-gRNA, selected with

2.5 µg/ml blasticidin-S and 500nM aTc, and cloned after parasites returned.

For expression of a second copy of PTEX150-kER from the *attB* locus, the *ptex150*

promoter and full-length gene were amplified from genomic DNA with P50/51 and the mNG-KER

cassette was amplified from JRB508 with P52/53. These amplicons were assembled together in

a second PCR reaction using P50/53 and inserted between ApaI/AflII in pLN-ENR-GFP (86),

resulting in the plasmid MAF39. This plasmid was co-transfected with pINT, selected with 2.5

µg/ml blasticidin-S and parasites were cloned after returning.

To generate the line bearing endogenous EXP2-mRuby and HSP101-mNG tags, the

plasmid pbEXP2-mRuby3 (89) was first linearized with AflII and co-transfected with pUF-Cas9-

EXP2-CT-gRNA (55) into NF54[attB:DiCre]. Selection was applied with 2.5 µg/ml blasticidin-S. Next,

the mNG coding sequence was amplified using P54/55 and inserted between AvrII/EagI in

pPM2GT-HSP101-3xFLAG (55) to replace the FLAG tag with mNG, resulting in the plasmid

JRB325. This plasmid was linearized with AflII and co-transfected with pAIO-HSP101-CT-gRNA

(89) into the NF54$^{attB:DiCre}$::EXP2-mRuby parasites. Selection was applied with 5nM WR99120

resulting in the line NF54$^{attB:DiCre}$::EXP2-mRuby+HSP101-mNG.

For generation of PTEX150-kER and HSP101-kER lines in *P. berghei*, the plasmid pBAT

(90) was digested with MluI/XhoI and the primer P56 was used as an insert to remove the

intervening AvrII, SnaBI and XhoI sites from the MCS. Next, the mRuby3 coding sequence fused

to the PbEXP2 signal peptide was amplified with P57/58 and inserted between SwaI/BamHI,

replacing the GFP. The mNG-kER tagging sequence was amplified from plasmid JRB508 using

P59/60 and inserted between SacII/SphI, replacing the mCherry-3xMYC. Finally, 5' flanks to tag

P. berghei hsp101 or *ptex150* with mNG-kER were amplified from *P. berghei* genomic DNA using

P61/62 or P63/64, respectively, and 3' flanks were amplified using P65/66 or P67/68, respectively.

The corresponding 5' and 3' flank amplicons were assembled with an intervening AflII site in a

second PCR reaction using P65/62 or P67/64, respectively, and inserted at SacII, resulting in

plasmids pPbHSP101-kER and pPbPTEX150-kER. The plasmids were linearized at AflII and

transfected as described (91) into the marker-free HP DiCre line (92). Transfected parasites were

reinjected into naïve mice and selection was applied with 0.07 mg/ml pyrimethamine in drinking

water provided *ad libitum* 24 hours later. After returning from selection, transfected populations

were cloned by limiting dilution IV injection into naïve mice to generate isogenic populations

designated PbHSP101-kER and PbPTEX150-kER.

***P. falciparum* growth assays**. Growth assays were done as described previously (93). Briefly,

parasites were treated with either DMSO or 10nM rapamycin for 3hrs and washed 1X. Then,

parasites were seeded between 0.2-1% parasitemia in triplicate and monitored every 48hrs by

flow cytometry on an Attune NxT (ThermoFisher) by nucleic acid staining with PBS containing 0.8

μg/ml acridine orange. For IPP rescue experiments, ClpB1-kER cultures were supplemented with

200μM IPP. For kER fusions expressed from the *attB* locus, cultures were supplemented with

1µM aTc 24hrs prior to rapamycin treatment to relieve TetR-DOZI-aptamer translational

repression. For HSP101-kER[TetR-DOZI] parasites, aTc was removed by washing parasites 7X with

13mL of BSD only media. 500nM aTc was then supplemented back to control parasites. As

required, parasites were sub-cultured to avoid high parasite density, and relative parasitemia at

each time point was back-calculated based on actual parasitemia multiplied by the relevant

dilution factors. Data were modeled using Prism 9 (GraphPad Software).

For EC_{50} shift assays, parasites were treated with DMSO or rapamycin before removal of

aTc. After aTc removal, parasites were seeded at 0.5% parasitemia with 2-fold dilutions of aTc

starting from 500nM, in triplicate and parasite growth was monitored after 72hrs.

P. berghei kER experiments. Parasite-infected blood was collected by cardiac puncture and

cultured in RPMI supplemented with 20% FBS. Cultures at 3-4% were treated with 10 nM

rapamycin or DMSO for 3-4 hours followed by a media change. After 18 hours of additional

culture, parasites were harvested for diagnostic PCR, live fluorescence microscopy and Western

blot. Equal numbers of DMSO or rapamycin-treated infected RBCs (~10,000 parasites per

injection) were IV injected into naïve mice and parasitemia was monitored by giemsa-stained

blood smear every two days.

Microscopy. For live microscopy, _P. falciparum_ parasites were treated with either DMSO or 10nM

rapamycin and viewed 24-48hrs later. Hoescht (33342, Invitrogen; 1:10,000) was used for to

visualize the nucleus by incubating with parasites 2-5min prior to visualization. ER-Tracker Red

(Invitrogen; 1:1000) was used to visualize the ER by incubating with parasites 30min at 37°C prior

to visualization. For constructs expressed using the _attB_ locus, parasites were grown in 1µM aTc

for 24hrs prior to incubation with DMSO or Rap to allow expression of the constructs. For _P._

berghei, parasites were isolated via cardiac puncture, treated with DMSO or rapamycin for 3hrs,

and allowed to develop overnight in culture before visualization. Quantification of MFI was

calculated using the measured Integrated Density value from Fiji (94). The internal MFI was

calculated using an ROI within the parasite PV and subtracting that from the whole parasite MFI

using an ROI outside the PV boundary. The percent MFI was then calculated by dividing the

internal MFI by the whole parasite MFI.

For protease protection assays of live cells, parasites were treated with 0.03% Saponin in

1X PBS for 15min at 4°C. Purified parasites where then washed once with 5mL of 1X PBS and

once with 5mL of digestion buffer (50mM Tris pH 7.5, 150mM NaCl, 1mM $CaCl_2$) (23). Parasite

samples where then split into two identical tubes, one of which was treated with 20µg/mL of

Proteinase K (740506.75, Macherey-Nagel), and incubated at 37°C for 15min. Proteinase K was

removed and samples resuspended with digestion buffer plus protease inhibitor cocktail (Halt

78429, Thermo Scientific) before visualization.

For IFAs of the EXP2- and ClpB1-kER lines, parasites were treated with either DMSO or

10nM rapamycin and fixed 24hrs or 144hrs later as described previously (93). For HSP101-

kER[TetR-DOZI] lines, parasites were synchronized with 5% sorbitol, treated with either DMSO or

10nM rapamycin, grown with or without 500nM aTc, and fixed 48hrs later. Briefly, parasites were

fixed with Fixative Solution (4% paraformaldehyde, 0.0075% glutaraldehyde in PBS) for 25min at

room temperature in the dark. Parasites were then permeabilized with 0.1% Triton in PBS for 10

minutes at room temperature while rocking. Parasites were washed 1X with PBS and then

blocked with 3% BSA in PBS for 1hr at room temperature. Primary antibody was applied in 3%

BSA solution for 1hr at room temperature, washed 2X with PBS, followed by secondary antibody

incubation for 1hr at room temperature. After 2X washes with PBS, parasites were adhered onto

Poly-L-lysine covered coverslips and then mounted onto slides. For parasites probed with SBP1,

parasites were acetone fixed as in (38). Briefly, thin smears were immersed in 100% acetone for

2min. Smears were washed 1X with PBS and then blocked with 3% BSA in PBS for 1hr at room

temperature. Primary antibody was applied in 3% BSA solution for 1hr at room temperature,

washed 2X with PBS, followed by secondary antibody incubation for 1hr at room temperature.

After 2X washes with PBS, slides were mounted with DAPI, covered with a coverslip and sealed.

The antibodies used for IFA were: mouse anti-HA antibody (clone 16B12; BioLegend, 1:200 IFA

and WB), rabbit polyclonal anti-HA SG77 (ThermoFisher; 1:500 IFA and WB), rabbit anti-

PfGRP78 (BiP) (MRA-1246; BEI Resources, NIAID, NIH 1:100), mouse monoclonal anti-FLAG

clone M2 (Sigma-Aldrich, 1:200 IFA and 1:500 WB), rabbit monoclonal anti-Flag 8H8L17

(ThermoFisher; 1:500 IFA and WB), mouse anti-HRP2 clone 2G12 (1:500 IFA) (95), rabbit

polyclonal anti-SBP1 (BR28, 1:500 IFA) (96), mouse monoclonal anti-EXP2 clone 7.7 (1:500 WB)

(97), rabbit polyclonal anti-*Plasmodium* Aldolase ab207494 (Abcam; 1:500 WB) and rabbit

polyclonal anti-ACP1 (1:500 IFA) (39, 98). Secondary antibodies used were anti-mouse antibody

conjugated to Alexa Fluor 488 or 546 and anti-rabbit antibody conjugated to Alexa Fluor 488, (Life

Technologies, 1:1000). Parasites were mounted with ProLong diamond with 4',6'-diamidino-2-

phenylindole (DAPI) (P36931; Invitrogen) and imaged using a using ZEISS Axio Observer. Image

processing, analysis, and display were performed using ZEISS Blue software and Adobe

Photoshop. Adjustments to brightness and contrast were made for display purposes.

Quantification of protein knockdown. To evaluate the level of HSP101 protein knockdown,

HSP101-kER$^{TetR-DOZI}$ parasites were synchronized via magnet purification and treated with DMSO

or rapamycin before removal of aTc. After aTc removal, parasites were seeded at ~1%

parasitemia with different dilutions of aTc and allowed to grow for 48hrs. Using an Attune NxT

(ThermoFisher) cytometer, infected RBCs were gated using Hoechst (34580, Invitrogen) and

mNG fluorescence was subsequently measured on infected RBCs using the blue laser (488nm)

with emission filter (430/30nm). The same procedure was applied for saponin-released parasites

after protease treatment as described above.

Western blotting. Western blotting for *Plasmodium* parasites was performed as described

previously (38, 55). Briefly, parasites were selectively permeabilized by treatment with ice-cold

0.03% saponin in PBS for 15 min followed by lysis with RIPA to remove the hemozoin. For

constructs expressed using the *attB* locus, parasites were grown in 500nM aTc for 24hrs prior to

incubation with DMSO or Rap to allow expression of the constructs. For the HSP101-kER$^{TetR-DOZI}$

lines, parasites were treated with DMSO or rapamycin before removal of aTc. After aTc removal,

parasites were seeded at ~1% parasitemia with different dilutions of aTc and allowed to grow for

48hrs before purification. The antibodies used in this study were mouse anti-HA antibody (clone

16B12; BioLegend, 1:500), mouse anti-FLAG (Clone M2; Sigma-Aldrich, 1:500), and mouse anti-

EXP2 (Clone 7.7; 1:500). The secondary antibodies used are IRDye 680CW goat anti-rabbit IgG

and IRDye 800CW goat anti-mouse IgG (LICOR Biosciences) (1:20,000). Western blot images

were processed using the Odyssey Clx LICOR infrared imaging system software (LICOR

Biosciences). Full length blots can be seen in Figures S5,S7.

Acknowledgements

This work was supported by NIH grant HL133453 to JRB. The funders had no role in study design,

data collection and interpretation, or the decision to submit the work for publication. We thank J.

McBride, D. Cavanaugh and EMRR for the EXP2 antibody, C. Braun-Breton for the SBP1

antibody and A. Waters for the HP DiCre *P. berghei* parasites.

Figures

Figure 1. Conditional ER retrieval of a secreted reporter does not impact parasite fitness.

A) Schematic of reporter expression cassette installed at the *attB* site of chromosome 6. An

mNeonGreen reporter is fused to the signal peptide of EXP2 and expressed under control of the

exp2 promoter. A 3xFLAG tag and stop codon are flanked by loxP-containing introns (triangles)

followed by a 3xHA-KDEL, stop codon and 10X aptamer array. Activation of DiCre by rapamycin

treatment excises the FLAG tag, bringing into frame the HA-KDEL tag. B) PCR showing excision

in mNG-kER parasites 24hrs after rapamycin treatment using primers P7/8. C) Western blot of

reporter 24hrs post treatment with DMSO or rapamycin. Molecular weights after signal peptide

cleavage are predicted to be 30.8 kDa for mNG-3xFLAG and 30.9 kDa for mNG-3xHA-KDEL. D)

Live microscopy of DMSO or rapamycin-treated parasites 24hrs post-treatment. E) Quantification

of percent internal mNG fluorescence 168hrs post-treatment with DMSO or rapamycin. Data are

pooled from 2 independent experiments and bar indicates mean (****, P<0.0001; unpaired t test).

F) Representative growth of asynchronous parasites (n=2 biological replicates) treated with

DMSO or rapamycin. Data are presented as means ± standard deviation from one biological

replicate (n = 3 technical replicates). All cultures were maintained in media supplemented with

500 nM aTc. Scale bar, 5µm.

Figure 2. Lethal ER retention of the apicoplast chaperone ClpB1 is rescued by IPP. A) Schematic showing strategy for appending kER to the C-terminus of *clpB1*. B) IFA of PFA fixed

parasites stained with mouse anti-Flag and rabbit anti-ACP1 antibodies. C) PCR showing excision

in ClpB1-kER parasites 24hrs after rapamycin treatment using primers P7/8. D) Western blot

24hrs post treatment with DMSO or rapamycin. Molecular weights after transit peptide cleavage

are predicted to be 135 kDa for ClpB1-mNG-3xFLAG and 135.1 kDa for ClpB1-mNG-3xHA-KDEL.

(+) represents the mature form of ClpB1 while the (*) represents ClpB1 with the transit peptide

still attached. E) Live microscopy of DMSO or rapamycin-treated parasites 48hrs post-treatment.

F) Representative growth of asynchronous parasites (n=3 biological replicates) treated with

DMSO or rapamycin, with or without supplementation with 200 μM IPP. Data are presented as

means ± standard deviation from one biological replicate (n = 3 technical replicates). G) IFA of

PFA fixed parasites stained with mouse anti-Flag or anti-HA and rabbit anti-ACP1 antibodies after

3 growth cycles where rapamycin treated parasites were supplemented with 200 μM IPP. Scale

bars, 5μm.

Figure 3. ER retention of rhoptry protein RON6 is tolerated by blood stage parasites. A) Schematic of rhoptry neck protein RON6. SP= signal peptide; black bar= site of piggybac insertion in RON6 (48). B) PCR showing excision in RON6-kER parasites 24hrs after treatment with rapamycin using primers P7/8. C) Live microscopy of DMSO or rapamycin-treated RON6-kER parasites 24hrs post-treatment. D) Representative growth of asynchronous RON6-kER parasites (n=3 biological replicates) treated with DMSO or rapamycin. Data are presented as means ± standard deviation from one biological replicate (n = 3 technical replicates). Scale bar, 5μm.

Figure 4. ER retention of EXP2 causes a lethal fitness defect independent of loss of PVM

transport functions. A) Schematic showing strategy for appending kER to the endogenous C-

terminus of EXP2 without mNG. B) PCR showing excision in EXP2-kER 24hrs after treatment

with rapamycin using primers P9/10 C) Western blot of EXP2-kER parasites 24hrs post treatment

with DMSO or rapamycin. Molecular weights after signal peptide cleavage are predicted to be

34.6 kDa for EXP2-3xFLAG and 34.7 kDa for EXP2-3xHA-KDEL. D) IFA of EXP2-kER 24hrs post-

treatment with DMSO or rapamycin. E) Growth of asynchronous parasites (n=2 biological

replicates) treated with DMSO or rapamycin. Data are presented as means ± standard deviation

from one biological replicate (n = 3 technical replicates). F) Live microscopy of parasites

expressing a full-length second copy of EXP2 with an mNG-kER fusion from the *attB* site under

the control of the endogenous *exp2* promoter (EXP2-FL-kER[attB]). Parasites were viewed 24hrs

after treatment with DMSO or rapamycin. G) Representative growth of asynchronous EXP2-FL-

kER[attB] parasites (n=2 biological replicates) treated with DMSO or rapamycin. Data are presented

as means ± standard deviation from one biological replicate (n = 3 technical replicates). H) Live

microscopy of parasites expressing a second copy of EXP2 with mNG-kER fusion and lacking

the amphipathic helix from the *attB* site (EXP2ΔTM-kER[attB]). Parasites were viewed 24hrs after

treatment with DMSO or rapamycin. Two representative examples of unexcised parasites are

presented, showing localization of the ΔTM version of EXP2 to the PV and host cell in some

infected RBCs (arrow). Digestive vacuole fluorescence was also observed as is typical for PV

proteins. I) Representative growth of asynchronous EXP2ΔTM-kER[attB] parasites (n=2 biological

replicates) treated with DMSO or rapamycin. Data are presented as means ± standard deviation

from one biological replicate (n = 3 technical replicates). Scale bar, 5µm.

Figure 5. ER retention of PTEX150 but not HSP101 causes a lethal block in protein export.

A) PCR showing excision in PTEX150-kER parasites 24hrs after rapamycin treatment using

primers P7/8. B) Western blot of PTEX150-kER 24hrs post-treatment with DMSO or rapamycin.

Molecular weights after signal peptide cleavage are predicted to be 141 kDa for PTEX150-mNG-

3xFLAG and 141.1 kDa for PTEX150-mNG-3xHA-KDEL. Note that PTEX150 is observed to

migrate at higher molecular weight than predicted (61). C) Live microscopy of PTEX150-kER

24hrs post-treatment with DMSO or rapamycin-treated. D) Representative growth of

asynchronous PTEX150-kER parasites (n=3 biological replicates) treated with DMSO or

rapamycin. Data are presented as means ± standard deviation from one biological replicate (n = 3

technical replicates). E) IFA of synchronized, ring-stage PTEX150-kER parasites 48hrs post-

treatment with DMSO or rapamycin and probed with anti-HRP2 and anti-FLAG or anti-HA

antibodies. F) PCR showing excision in HSP101-kER parasites 24hrs after rapamycin treatment

using primers P7/8. G) Western blot of HSP101-kER parasites 14 days post-treatment with DMSO

or rapamycin. Molecular weights after signal peptide cleavage are predicted to be 130.5 kDa for

HSP101-mNG-3xFLAG and 130.6 kDa for HSP101-mNG-3xHA-KDEL. H) Live microscopy of

HSP101-kER 24hrs and 168hrs post-treatment with DMSO or rapamycin. I) Representative

growth of asynchronous HSP101-kER parasites (n=3 biological replicates) treated with DMSO or

rapamycin. Data are presented as means ± standard deviation from one biological replicate (n = 3

technical replicates). J) IFA of synchronized, ring-stage HSP101-kER parasites 48hrs post-

treatment with DMSO or rapamycin and probed with anti-HRP2 and anti-FLAG or anti-HA

antibodies. Scale bar, 5μm.

Figure 6. Depletion of HSP101 PV levels does not hypersensitize parasites to HSP101 knockdown. A) Schematic showing strategy for appending kER and the TetR-DOZI aptamers system to the endogenous C-terminus of HSP101. B) PCR showing excision in HSP101-kER$^{TetR-DOZI}$ parasites 24hrs after rapamycin treatment using primers P7/8. C) Live microscopy of HSP101-kER$^{TetR-DOZI}$ parasites 168hrs post-treatment with DMSO or rapamycin. D) IFA detecting the soluble PEXEL protein HRP2 or the transmembrane PNEP SBP1 in synchronized, ring-stage

HSP101-kER$^{TetR-DOZI}$ parasites grown for 48hrs +/- aTc and treated with DMSO or rapamycin 1

cycle prior. E) Quantification of HSP101 levels by flow cytometry measurement of mNeonGreen

fluorescence in HSP101-kER$^{TetR-DOZI}$ parasites grown for 48 hrs at indicated aTc concentrations

and treated with DMSO or rapamycin 2 cycles prior. MFI data were normalized to DMSO-treated

parasites grown in 500 nM aTc (100%) and presented as mean ± SEM (n = 5 biological replicates).

F) Representative growth curves (n = 4 biological replicates) of HSP101-kER$^{TetR-DOZI}$ parasites

grown at indicated aTc concentrations and treated with DMSO or rapamycin 2 cycles prior. Data

are presented as means ± standard deviation from one biological replicate (n = 3 technical

replicates). G) Representative EC$_{50}$ shift assay (n=3 biological replicates) of parasites treated with

DMSO or rapamycin with two-fold dilutions of aTc starting at 500nM. Data are presented as

means ± standard deviation from one biological replicate (n = 3 technical replicates). Scale bars,

5µm.

Figure 7: ER retention of HSP101 is lethal *in vivo*. A) PCR showing excision in PbPTEX150-kER and PbHSP101-kER *P. berghei* parasites treated with rapamycin and cultured 18hrs *ex vivo* using primers P7/P60. B) Western blot of PbPTEX150-kER and PbHSP101-kER parasites 24hrs post-treatment with DMSO or rapamycin. Molecular weights after signal peptide cleavage are predicted to be 130.9 kDa for PbHSP101-mNG-3xFLAG, 131 kDa for PbHSP101-mNG-3xHA-KDEL, 130.5 kDa for PbPTEX150-mNG-3xFLAG and 130.6 kDa for PbPTEX150-mNG-3xHA-KDEL. Note that PbPTEX150 is observed to migrate at higher molecular weight than predicted (99). C,D) Live microscopy of PbHSP101-kER or PbPTEX150-kER parasites 18hrs post-treatment with DMSO or rapamycin. E,F) Parasitemia of PbHSP101-kER or PbPTEX150-kER infected mice inoculated by tail vein-injection of parasites treated *ex vivo* with DMSO or rapamycin. Data are presented as means ± standard deviation of three independent experiments (one mouse infected per treatment in each experiment). No parasites were observed in mice

injected with rapamycin treated parasites except that one mouse injected with rapamycin-treated

PbHSP101-kER parasites became patent on day 10 (#). However, PCR showed these parasites

had not undergone excision (see Figure S6C). Scale bar, 5µm.

Figure S1. Generation and validation of NF54^attB-DiCre parasites. A) Diagnostic PCR showing

DiCre cassette integration into the *pfs47* locus using primers P1/2 and P3/4 to detect unmodified

locus, and primers P1/5 and P6/4 to detect integration. B) Schematic of reporter integrated at *attB*

in *cg6* to test DiCre functionality in NF54^attB-DiCre. C) Western blot and D) live microscopy of

parasites 48hrs after 3hr treatment with DMSO or 10nM rapamycin. Molecular weights are

predicted to be 30 kDa for mNG-3xHA and 40.1 kDa for aldolase.

 Figure S2. Validation of DiCre-mediated excision in second copy EXP2-kER parasites. A)

Schematic showing cassette for second copy expression of full-length EXP2 or a version lacking

the amphipathic helix with kER fusion from the *attB* site of chromosome 6. B) PCR showing

excision in EXP2-FL-kER[attB] and EXP2ΔTM-kER[attB] parasites 24hrs after rapamycin treatment

using primers P7/8. C) Western blots from second copy EXP2-kER parasite samples taken 24hrs

after DMSO or rapamycin treatment showing conversion from Flag to HA.

Figure S3. Protease protection assay on live cells verifies an internal, perinuclear pool of

HSP101 but not EXP2 or PTEX150. Live microscopy of parasite lines with endogenous

fluorescent protein fusions A) PTEX150-mNG-3xFLAG or B) EXP2-mRuby3 and HSP101-mNG

treated with saponin +/- Proteinase K. Images are representative of n=2 biological replicates.

Arrowhead: protease-protected fluorescence within the digestive vacuole from endocytosis

typically seen for PV proteins; arrow: protease-protected perinuclear fluorescence unique to

HSP101. Scale bar, 5μm.

Figure S4. Supporting data for Figure 5. A) Schematic showing cassette for second copy expression of PTEX150-kER from the *attB* site of chromosome 6. B) PCR showing excision in PTEX150-kER[attB] parasites 24hrs after rapamycin treatment using primers P7/8. C) Representative growth curves of asynchronous, second copy PTEX150-kER parasites (n=3 biological replicates) treated with DMSO or rapamycin. Data are presented as means ± standard deviation from one biological replicate (n = 3 technical replicates). D) Live microscopy of second copy PTEX150-kER parasites 24hrs post-treatment with DMSO or rapamycin. E) Quantification

of percent internal mNG fluorescence in live microcopy images of endogenously tagged HSP101-

and PTEX150-kER parasites 24hrs post-treatment with DMSO or rapamycin. Data are pooled

from 3 (HSP101) or 2 (PTEX150) independent experiments and bars indicate mean ± standard

deviation (****, P<0.0001; unpaired t test). F) Quantification of percent internal mNG fluorescence

in live microcopy images of endogenously tagged HSP101-kER and second copy PTEX150-kER

parasites lines 168hrs post-treatment with DMSO or rapamycin. Data are pooled from 2

independent experiments and bars indicate represented as means ± standard deviation (****,

P<0.0001; unpaired t test).

Figure S5. Supporting data for Figure 6. A) Mean fluorescence intensity quantification from

proteinase K treated parasites. The amount of protected fluorescent signal (percent of total before

protease treatment) was determined by dividing MFI after proteinase K treatment from before

proteinase K treatment. Colors represent different replicates. B) Representative growth curve (n

= 3 biological replicates) of HSP101-kER$^{TetR-DOZI}$ parasites grown with or without aTc. Data are

presented as means ± standard deviation of one biological replicate (n = 3 technical replicates).

C) Western blot of HSP101-kER$^{TetR-DOZI}$ parasites grown 48hrs at indicated aTc concentrations.

Parasites grown in 500 nM aTc were treated with DMSO or rapamycin 24hrs before aTc was

washed out and restored at indicated concentrations. Results are representative of 4 independent

experiments.

Figure S6. Supporting data for Figure 7. A) Schematic showing strategy for appending kER to

the endogenous C-terminus of *P. berghei hsp101* and *ptex150*. B) PCR using P7/P69 to evaluate

excision of the kER cassette in parasites taken from mice injected with DMSO (parasite collected

on day 8) or rapamycin-treated PbHSP101-kER (taken from the single mouse where parasites

appeared on day 10). The results indicate that the kER cassette was unexcised in the parasites

observed in the mouse that became patent on day 10.

Figure S7. Integration PCRs and Uncropped Western blots A) Integration PCR for

endogenous kER tags using primers P11/P12 (HSP101), P13/P12 (ClpB1), P14/P12 (PTEX150),

P15/P16 (RON6), and P17/P8 (EXP2). B) Uncropped Western blots. For RON6-kER, molecular

weights after signal peptide cleavage are predicted to be 141.3 kDa for RON6-mNG-3xFLAG and

141.4 kDa for RON6-mNG-3xHA-KDEL. Arrows indicate RON6 signal; notice the low expression

of RON6, consistent with low expression levels in the blood stage (PlasmoDB). For EXP2-FL-

kER[attB], molecular weights after signal peptide cleavage are predicted to be 61.1 kDa for EXP2-

mNG-3xFLAG and 61.2 kDa for EXP2-mNG-3xHA-KDEL. For EXP2ΔTM-kER[attB], molecular

weights after signal peptide cleavage are predicted to be 54.2 kDa for EXP2ΔTM-mNG-3xFLAG

and 54.3 kDa for EXP2ΔTM-3xHA-KDEL.

References

1. O. A. Akuh, R. Elahi, S. T. Prigge, F. Seeber, The ferredoxin redox system - an essential
electron distributing hub in the apicoplast of Apicomplexa. *Trends Parasitol* **38**, 868-881
(2022).

2. M. M. Cova, M. H. Lamarque, M. Lebrun, How Apicomplexa Parasites Secrete and Build
Their Invasion Machinery. *Annual Review of Microbiology* **76**, 619-640 (2022).

3. J. M. Matz, J. R. Beck, M. J. Blackman, The parasitophorous vacuole of the blood-stage
malaria parasite. *Nat Rev Microbiol* 10.1038/s41579-019-0321-3 (2020).

4. M. Garten, J. R. Beck, Structured to conquer: transport across the Plasmodium
parasitophorous vacuole. *Current Opinion in Microbiology* **63**, 181-188 (2021).

5. T. F. De Koning-Ward, M. W. A. Dixon, L. Tilley, P. R. Gilson, Plasmodium species:
master renovators of their host cells. *Nature Reviews Microbiology* **14**, 494-507 (2016).

6. J. R. Beck, C.-M. Ho, Transport mechanisms at the malaria parasite-host cell interface.
PLOS Pathogens **17**, e1009394 (2021).

7. T. J. Sargeant *et al.*, Lineage-specific expansion of proteins exported to erythrocytes in
malaria parasites. *Genome Biol* **7**, R12 (2006).

8. J. A. Boddey *et al.*, Role of Plasmepsin V in Export of Diverse Protein Families from the
Plasmodium falciparum Exportome. *Traffic* **14**, 532-550 (2013).

9. C. Van Ooij *et al.*, The Malaria Secretome: From Algorithms to Essential Function in
Blood Stage Infection. *PLoS Pathogens* **4**, e1000084 (2008).

10. J. Schulze *et al.*, The Plasmodium falciparum exportome contains non-canonical
PEXEL/HT proteins. *Mol Microbiol* **97**, 301-314 (2015).

11. T. K. Jonsdottir, M. Gabriela, B. S. Crabb, F. d. K.-W. T, P. R. Gilson, Defining the
Essential Exportome of the Malaria Parasite. *Trends Parasitol* **37**, 664-675 (2021).

12. S. Tomavo, C. Slomianny, M. Meissner, V. B. Carruthers, Protein trafficking through the
endosomal system prepares intracellular parasites for a home invasion. *PLoS*
Pathogens **9**, e1003629 (2013).

13. A. Florentin, D. W. Cobb, H. M. Kudyba, V. Muralidharan, Directing Traffic: Chaperone-
mediated protein transport in malaria parasites. *Cell Microbiol* 10.1111/cmi.13215,
e13215 (2020).

14. P. Brauer *et al.*, Structural basis for pH-dependent retrieval of ER proteins from the Golgi
by the KDEL receptor. *Science* **363**, 1103-1107 (2019).

15. S. Munro, H. R. B. Pelham, A C-terminal signal prevents secretion of luminal ER
proteins. *Cell* **48**, 899-907 (1987).

16.   A. Gerondopoulos *et al.*, A signal capture and proofreading mechanism for the KDEL-
819   receptor explains selectivity and dynamic range in ER retrieval. *eLife* **10** (2021).

820   17.   H. G. Elmendorf, K. Haldar, Identification and localization of ERD2 in the malaria
821   parasite Plasmodium falciparum: separation from sites of sphingomyelin synthesis and
822   implications for organization of the Golgi. *The EMBO Journal* **12**, 4763-4773 (1993).

823   18.   N. La Greca, A. R. Hibbs, C. Riffkin, M. Foley, L. Tilley, Identification of an endoplasmic
824   reticulum-resident calcium-binding protein with multiple EF-hand motifs in asexual
825   stages of Plasmodium falciparum. *Mol Biochem Parasitol* **89**, 283-293 (1997).

826   19.   H. M. Kudyba *et al.*, The endoplasmic reticulum chaperone PfGRP170 is essential for
827   asexual development and is linked to stress response in malaria parasites. *Cell Microbiol*
828   **21**, e13042 (2019).

829   20.   D. W. Cobb *et al.*, A redox-active crosslinker reveals an essential and inhibitable
830   oxidative folding network in the endoplasmic reticulum of malaria parasites. *PLOS*
831   *Pathogens* **17**, e1009293 (2021).

832   21.   S. Külzer, N. Gehde, J. M. Przyborski, Return to sender: use of Plasmodium ER retrieval
833   sequences to study protein transport in the infected erythrocyte and predict putative ER
834   protein families. *Parasitology Research* **104**, 1535-1541 (2009).

835   22.   M. A. Fierro *et al.*, An ER CREC family protein regulates the egress proteolytic cascade
836   in malaria parasites. *bioRxiv* 10.1101/457481, 457481 (2019).

837   23.   M. Gabriela *et al.*, A revised mechanism for how Plasmodium falciparum recruits and
838   exports proteins into its erythrocytic host cell. *PLOS Pathogens* **18**, e1009977 (2022).

839   24.   J. R. Beck, V. Muralidharan, A. Oksman, D. E. Goldberg, PTEX component HSP101
840   mediates export of diverse malaria effectors into host erythrocytes. *Nature* **511**, 592-595
841   (2014).

842   25.   B. Elsworth *et al.*, PTEX is an essential nexus for protein export in malaria parasites.
843   *Nature* **511**, 587-591 (2014).

844   26.   C. M. Ho *et al.*, Malaria parasite translocon structure and mechanism of effector export.
845   *Nature* **561**, 70-75 (2018).

846   27.   E. Knuepfer, M. Napiorkowska, C. Van Ooij, A. A. Holder, Generating conditional gene
847   knockouts in Plasmodium – a toolkit to produce stable DiCre recombinase-expressing
848   parasite lines using CRISPR/Cas9. *Scientific Reports* **7** (2017).

849   28.   C. R. Collins *et al.*, Robust inducible Cre recombinase activity in the human malaria
850   parasite P lasmodium falciparum enables efficient gene deletion within a single asexual
851   erythrocytic g. *Molecular Microbiology* **88**, 687-701 (2013).

852   29.   S. Chaubey, M. Grover, U. Tatu, Endoplasmic Reticulum Stress Triggers
853   Gametocytogenesis in the Malaria Parasite. *Journal of Biological Chemistry* **289**, 16662-
854   16674 (2014).

30. A. J. Polino, A. S. Nasamu, J. C. Niles, D. E. Goldberg, Assessment of Biological Role
and Insight into Druggability of the Plasmodium falciparum Protease Plasmepsin V. *ACS*
Infect Dis **6**, 738-746 (2020).

31. J. M. Przyborski *et al.*, Trafficking of STEVOR to the Maurer's clefts in Plasmodium
falciparum-infected erythrocytes. *The EMBO Journal* **24**, 2306-2317 (2005).

32. G. G. Van Dooren *et al.*, Development of the endoplasmic reticulum, mitochondrion and
apicoplast during the asexual life cycle ofPlasmodium falciparum. *Molecular*
Microbiology **57**, 405-419 (2005).

33. C. J. Tonkin, N. S. Struck, K. A. Mullin, L. M. Stimmler, G. I. McFadden, Evidence for
Golgi-independent transport from the early secretory pathway to the plastid in malaria
parasites. *Molecular Microbiology* **61**, 614-630 (2006).

34. M. C. S. Lee, P. A. Moura, E. A. Miller, D. A. Fidock, Plasmodium falciparumSec24
marks transitional ER that exports a model cargo via a diacidic motif. *Molecular*
Microbiology **68**, 1535-1546 (2008).

35. S. R. Heiny, S. Pautz, M. Recker, J. M. Przyborski, Protein Traffic to thePlasmodium
falciparumApicoplast: Evidence for a Sorting Branch Point at the Golgi. *Traffic* **15**, 1290-
1304 (2014).

36. M. J. Boucher *et al.*, Integrative proteomics and bioinformatic prediction enable a high-
confidence apicoplast proteome in malaria parasites. *PLoS Biol* **16**, e2005895 (2018).

37. J. Birnbaum *et al.*, A Kelch13-defined endocytosis pathway mediates artemisinin
resistance in malaria parasites. *Science* **367**, 51-59 (2020).

38. T. Nessel *et al.*, EXP1 is required for organisation of EXP2 in the intraerythrocytic
malaria parasite vacuole. *Cell Microbiol* **22**, e13168 (2020).

39. R. F. Waller, M. B. Reed, A. F. Cowman, G. I. McFadden, Protein trafficking to the
plastid of Plasmodium falciparum is via the secretory pathway. *Embo Journal* **19**, 1794-
1802 (2000).

40. A. Adisa *et al.*, The Signal Sequence of Exported Protein-1 Directs the Green
Fluorescent Protein to the Parasitophorous Vacuole of Transfected Malaria Parasites.
Journal of Biological Chemistry **278**, 6532-6542 (2003).

41. S. M. Ganesan, A. Falla, S. J. Goldfless, A. S. Nasamu, J. C. Niles, Synthetic RNA-
protein modules integrated with native translation mechanisms to control gene
expression in malaria parasites. *Nat Commun* **7**, 10727 (2016).

42. K. Rajaram, H. B. Liu, S. T. Prigge, Redesigned TetR-Aptamer System To Control Gene
Expression in Plasmodium falciparum. *mSphere* **5** (2020).

43. S. H. Adjalley *et al.*, Quantitative assessment of Plasmodium falciparum sexual
development reveals potent transmission-blocking activity by methylene blue.
Proceedings of the National Academy of Sciences **108**, E1214-E1223 (2011).

44.   A. P. AhYoung, A. Koehl, D. Cascio, P. F. Egea, Structural mapping of the ClpB
893   ATPases of Plasmodium falciparum: Targeting protein folding and secretion for
894   antimalarial drug design. *Protein Sci* **24**, 1508-1520 (2015).

895   45.   M. El Bakkouri *et al.*, The Clp Chaperones and Proteases of the Human Malaria Parasite
896   Plasmodium falciparum. *Journal of Molecular Biology* **404**, 456-477 (2010).

897   46.   E. Yeh, J. L. Derisi, Chemical Rescue of Malaria Parasites Lacking an Apicoplast
898   Defines Organelle Function in Blood-Stage Plasmodium falciparum. *PLoS Biology* **9**,
899   e1001138 (2011).

900   47.   G. G. Van Dooren, V. Su, M. C. D'Ombrain, G. I. McFadden, Processing of an
901   Apicoplast Leader Sequence inPlasmodium falciparum and the Identification of a
902   Putative Leader Cleavage Enzyme. *Journal of Biological Chemistry* **277**, 23612-23619
903   (2002).

904   48.   M. Zhang *et al.*, Uncovering the essential genes of the human malaria parasite
905   Plasmodium falciparum by saturation mutagenesis. *Science* **360** (2018).

906   49.   E. Bushell *et al.*, Functional Profiling of a Plasmodium Genome Reveals an Abundance
907   of Essential Genes. *Cell* **170**, 260-272 e268 (2017).

908   50.   M. E. Fichera, D. S. Roos, A plastid organelle as a drug target in apicomplexan
909   parasites. *Nature* **390**, 407-409 (1997).

910   51.   E. L. Dahl, P. J. Rosenthal, Multiple antibiotics exert delayed effects against the
911   Plasmodium falciparum apicoplast. *Antimicrob Agents Chemother* **51**, 3485-3490 (2007).

912   52.   N. I. Proellocks *et al.*, Characterisation of PfRON6, a Plasmodium falciparum rhoptry
913   neck protein with a novel cysteine-rich domain. *Int J Parasitol* **39**, 683-692 (2009).

914   53.   S. A. Desai, D. J. Krogstad, E. W. McCleskey, A nutrient-permeable channel on the
915   intraerythrocytic malaria parasite. *Nature* **362**, 643-646 (1993).

916   54.   S. A. Desai, R. L. Rosenberg, Pore size of the malaria parasite's nutrient channel.
917   *Proceedings of the National Academy of Sciences* **94**, 2045-2049 (1997).

918   55.   M. Garten *et al.*, EXP2 is a nutrient-permeable channel in the vacuolar membrane of
919   Plasmodium and is essential for protein export via PTEX. *Nat Microbiol* **3**, 1090-1098
920   (2018).

921   56.   D. A. Gold *et al.*, The Toxoplasma Dense Granule Proteins GRA17 and GRA23 Mediate
922   the Movement of Small Molecules between the Host and the Parasitophorous Vacuole.
923   *Cell Host Microbe* **17**, 642-652 (2015).

924   57.   P. R. Sanders *et al.*, The N-terminus of EXP2 forms the membrane-associated pore of
925   the protein exporting translocon PTEX in Plasmodium falciparum. *The Journal of
926   Biochemistry* **165**, 239-248 (2019).

58. K. Hakamada, H. Watanabe, R. Kawano, K. Noguchi, M. Yohda, Expression and
characterization of the Plasmodium translocon of the exported proteins component
EXP2. *Biochemical and Biophysical Research Communications* **482**, 700-705 (2017).

59. M. D. Peraro, F. G. Van Der Goot, Pore-forming toxins: ancient, but never really out of
fashion. *Nature Reviews Microbiology* **14**, 77-92 (2016).

60. S. C. Charnaud, R. Kumarasingha, H. E. Bullen, B. S. Crabb, P. R. Gilson, Knockdown
of the translocon protein EXP2 in Plasmodium falciparum reduces growth and protein
export. *PLOS ONE* **13**, e0204785 (2018).

61. T. F. De Koning-Ward *et al.*, A newly discovered protein export machine in malaria
parasites. *Nature* **459**, 945-949 (2009).

62. J. M. Matz *et al.*, The Plasmodium berghei translocon of exported proteins reveals
spatiotemporal dynamics of tubular extensions. *Scientific Reports* **5**, 12532 (2015).

63. S. C. Charnaud *et al.*, Spatial organization of protein export in malaria parasite blood
stages. *Traffic* **19**, 605-623 (2018).

64. S. Nair *et al.*, Nutrient Limitation Magnifies Fitness Costs of Antimalarial Drug
Resistance Mutations. *Antimicrob Agents Chemother* **66**, e0152921 (2022).

65. A. Gupta *et al.*, Complex nutrient channel phenotypes despite Mendelian inheritance in a
Plasmodium falciparum genetic cross. *PLOS Pathogens* **16**, e1008363 (2020).

66. J. M. Matz *et al.*, In Vivo Function of PTEX88 in Malaria Parasite Sequestration and
Virulence. *Eukaryot Cell* **14**, 528-534 (2015).

67. S. A. Chisholm *et al.*, Contrasting Inducible Knockdown of the Auxiliary PTEX
Component PTEX88 in P. falciparum and P. berghei Unmasks a Role in Parasite
Virulence. *PLOS ONE* **11**, e0149296 (2016).

68. H. M. Kudyba, D. W. Cobb, J. Vega-Rodríguez, V. Muralidharan, Some conditions apply:
Systems for studying Plasmodium falciparum protein function. *PLOS Pathogens* **17**,
e1009442 (2021).

69. J. Birnbaum *et al.*, A genetic system to study Plasmodium falciparum protein function.
Nature Methods **14**, 450-456 (2017).

70. A. D. Roberts, S. C. Nair, A. J. Guerra, S. T. Prigge, Development of a conditional
localization approach to control apicoplast protein trafficking in malaria parasites. *Traffic*
20, 571-582 (2019).

71. A. Praznik *et al.*, Regulation of protein secretion through chemical regulation of
endoplasmic reticulum retention signal cleavage. *Nature Communications* **13** (2022).

72. A. Prasad, P. Mastud, S. Patankar, Dually localised proteins found in both the apicoplast
and mitochondrion utilize the Golgi-dependent pathway for apicoplast targeting in
Toxoplasma gondii. *Biology of the Cell* **113**, 58-78 (2021).

73. A. Florentin *et al.*, PfClpC Is an Essential Clp Chaperone Required for Plastid Integrity and Clp Protease Stability in Plasmodium falciparum. *Cell Rep* **21**, 1746-1756 (2017).

74. A. Florentin, D. R. Stephens, C. F. Brooks, R. P. Baptista, V. Muralidharan, Plastid biogenesis in malaria parasites requires the interactions and catalytic activity of the Clp proteolytic system. *Proc Natl Acad Sci U S A* **117**, 13719-13729 (2020).

75. A. Mogk, B. Bukau, H. H. Kampinga, Cellular Handling of Protein Aggregates by Disaggregation Machines. *Molecular Cell* **69**, 214-226 (2018).

76. I. Bouchnak, K. J. Van Wijk, Structure, function, and substrates of Clp AAA+ protease systems in cyanobacteria, plastids, and apicoplasts: A comparative analysis. *Journal of Biological Chemistry* **296**, 100338 (2021).

77. K. Pakos-Zebrucka *et al.*, The integrated stress response. *EMBO reports* **17**, 1374-1395 (2016).

78. M. Costa-Mattioli, P. Walter, The integrated stress response: From mechanism to disease. *Science* **368** (2020).

79. M. B. Harbut *et al.*, Targeting the ERAD pathway via inhibition of signal peptide peptidase for antiparasitic therapeutic design. *Proceedings of the National Academy of Sciences* **109**, 21486-21491 (2012).

80. M. Zhang *et al.*, PK4, a eukaryotic initiation factor 2 (eIF2) kinase, is essential for the development of the erythrocytic cycle of Plasmodium. *Proceedings of the National Academy of Sciences* **109**, 3956-3961 (2012).

81. M. Zhang *et al.*, Inhibiting the Plasmodium eIF2α Kinase PK4 Prevents Artemisinin-Induced Latency. *Cell Host & Microbe* **22**, 766-776.e764 (2017).

82. E. C. Gaynor, S. Te Heesen, T. R. Graham, M. Aebi, S. D. Emr, Signal-mediated retrieval of a membrane protein from the Golgi to the ER in yeast. *Journal of Cell Biology* **127**, 653-665 (1994).

83. M. Stornaiuolo *et al.*, KDEL and KKXX Retrieval Signals Appended to the Same Reporter Protein Determine Different Trafficking between Endoplasmic Reticulum, Intermediate Compartment, and Golgi Complex. *Molecular Biology of the Cell* **14**, 889-902 (2003).

84. B. L. Tang, S. H. Wong, S. H. Low, W. Hong, Retention of a type II surface membrane protein in the endoplasmic reticulum by the Lys-Asp-Glu-Leu sequence. *Journal of Biological Chemistry* **267**, 7072-7076 (1992).

85. B. E. Sleebs *et al.*, Inhibition of Plasmepsin V Activity Demonstrates Its Essential Role in Protein Export, PfEMP1 Display, and Survival of Malaria Parasites. *PLoS Biology* **12**, e1001897 (2014).

86. S. H. Adjalley, M. C. S. Lee, D. A. Fidock, "A Method for Rapid Genetic Integration into Plasmodium falciparum Utilizing Mycobacteriophage Bxb1 Integrase". (Humana Press, 2010), 10.1007/978-1-60761-652-8_6, pp. 87-100.

87. M. L. Jones *et al.*, A versatile strategy for rapid conditional genome engineering using
loxP sites in a small synthetic intron in Plasmodium falciparum. **6**, 21800 (2016).

88. S. Glushakova *et al.*, Exploitation of a newly-identified entry pathway into the malaria
parasite-infected erythrocyte to inhibit parasite egress. *Scientific Reports* **7** (2017).

89. A. I. Ahiya, S. Bhatnagar, J. M. Morrisey, J. R. Beck, A. B. Vaidya, Dramatic
Consequences of Reducing Erythrocyte Membrane Cholesterol on Plasmodium
falciparum. *Microbiol Spectr* **10**, e0015822 (2022).

90. T. W. Kooij, M. M. Rauch, K. Matuschewski, Expansion of experimental genetics
approaches for Plasmodium berghei with versatile transfection vectors. *Mol Biochem*
Parasitol **185**, 19-26 (2012).

91. C. J. Janse, J. Ramesar, A. P. Waters, High-efficiency transfection and drug selection of
genetically transformed blood stages of the rodent malaria parasite Plasmodium berghei.
Nature Protocols **1**, 346-356 (2006).

92. R. S. Kent *et al.*, Inducible developmental reprogramming redefines commitment to
sexual development in the malaria parasite Plasmodium berghei. *Nature Microbiology* **3**,
1206-1213 (2018).

93. M. A. Fierro *et al.*, An Endoplasmic Reticulum CREC Family Protein Regulates the
Egress Proteolytic Cascade in Malaria Parasites. *mBio* **11** (2020).

94. J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nature*
Methods **9**, 676-682 (2012).

95. E. P. Rock *et al.*, Comparative analysis of the Plasmodium falciparum histidine-rich
proteins HRP-I, HRP-II and HRP-III in malaria parasites of diverse origin. *Parasitology*
95, 209-227 (1987).

96. T. Blisnick *et al.*, Pfsbp1, a Maurer's cleft Plasmodium falciparum protein, is associated
with the erythrocyte skeleton. *Mol Biochem Parasitol* **111**, 107-121 (2000).

97. R. Hall *et al.*, Antigens of the erythrocytic stages of the human malaria parasite
Plasmodium falciparum detected by monoclonal antibodies. *Molecular and Biochemical*
Parasitology **7**, 247-265 (1983).

98. M. Ponpuak *et al.*, A role for falcilysin in transit peptide degradation in the<i>Plasmodium
falciparum</i>apicoplast. *Molecular Microbiology* **63**, 314-334 (2007).

99. K. Matthews *et al.*, The<i>Plasmodium</i>translocon of exported proteins (PTEX)
component thioredoxin-2 is important for maintaining normal blood-stage growth.
Molecular Microbiology **89**, 1167-1186 (2013).

1034