1

Knock-sideways by inducible ER retrieval enables a novel approach for studying *Plasmodium* secreted proteins

Manuel A Fierro[1], Tahir Hussain[1], Liam J Campin[2] and Josh R Beck[1,2*]

[1]Department of Biomedical Sciences, Iowa State University, Ames, IA, 50011, USA

[2]Roy J. Carver Department of Biochemistry, Biophysics and Molecular Biology, Iowa State University, Ames, IA, 50011, USA

[*]Corresponding author: jrbeck@iastate.edu

Classification: Biological Sciences, Microbiology

Key words: *Plasmodium,* ER, ClpB1, PTEX, knock-sideways

Abstract

Malaria parasites uniquely depend on protein secretion for their obligate intracellular lifestyle but approaches for dissecting *Plasmodium* secreted protein functions are limited. We report knockER, a novel DiCre-mediated knock-sideways approach to sequester secreted proteins in the ER by inducible fusion with a KDEL ER-retrieval sequence. We show conditional ER sequestration of diverse proteins is not generally toxic, enabling loss-of-function studies. We employed knockER in multiple *Plasmodium* species to interrogate the trafficking, topology and function of an assortment of proteins that traverse the secretory pathway to diverse compartments including the apicoplast (ClpB1), rhoptries (RON6), dense granules and parasitophorous vacuole (EXP2, PTEX150, HSP101). Taking advantage of the unique ability to redistribute secreted proteins from their terminal destination to the ER, we reveal vacuolar levels of the PTEX translocon component HSP101 but not PTEX150 are maintained in excess of what is required to sustain effector protein export into the erythrocyte. Intriguingly, vacuole depletion of HSP101 hypersensitized parasites to a destabilization tag that inhibits HSP101-PTEX complex formation but not to translational knockdown of the entire HSP101 pool, illustrating how redistribution of a target protein by knockER can be used to query function in a compartment-specific manner. Collectively, our results establish knockER as a novel tool for dissecting secreted protein function with sub-compartmental resolution that should be widely amenable to genetically tractable eukaryotes.

Significance

Protein trafficking and secretion through the endomembrane system is a defining feature of eukaryotes. The secretory pathway is central to the unique biology and pathology of the obligate intracellular malaria parasite, however tools for studying secreted protein function are limited. Knock-sideways is a powerful mutagenesis strategy that conditionally sequesters a protein away from its site of function but is generally not applicable to secreted proteins. We developed a simple approach to conditionally sequester *Plasmodium* secreted proteins in the ER by inducible C-terminal fusion with a KDEL ER-retrieval sequence that can be used for trafficking, topology and loss-of-function studies. The knockER strategy is broadly applicable to functional dissection of proteins that traverse the eukaryotic secretory pathway.

Introduction

Protein secretion is a fundamental biological process of all living cells. In eukaryotes this is accomplished by a dedicated secretory pathway that begins with import into the Endoplasmic Reticulum (ER) for initial folding and post-translational modification, followed by sorting in the Golgi and post-Golgi compartments. As an obligate intracellular parasite, secreted proteins play a critical role in the biology of the malaria parasite, *Plasmodium*. Through the secretory pathway, proteins are trafficked into both the plasma membrane and several organelles, including a relict plastid called the apicoplast that is critical for parasite metabolism (1, 2). During schizogony, another subset of proteins is packaged into several distinct secretory organelles that are subsequently discharged to facilitate host cell attachment and penetration (3). Invagination of the host membrane by the parasite during invasion generates an intracellular microenvironment called the parasitophorous vacuole (PV) which forms the interface for host-parasite interactions (4). Resident PV proteins secreted into the vacuole perform critical functions at this interface, including nutrient/waste exchange and protein export (5). The latter process enables parasites to deploy a battery of secreted effector proteins across the PV membrane (PVM) and into the host compartment, dramatically remodeling the erythrocyte to meet nutritional demands and avoid host defenses (6, 7). Remarkably, it is estimated that up to 10% of the parasite genome is devoted to encoding exported effectors, further highlighting the central importance of protein secretion to this parasitic lifestyle (8-12).

As with other eukaryotes, entry into the malaria parasite secretory pathway generally begins with co-translational import of proteins containing a signal peptide or transmembrane domain(s) into the ER via the Sec61 complex, followed by sorting from the Golgi to internal compartments, the PV or beyond into the host (13). ER-resident proteins that escape the ER via bulk flow in COPII vesicles are returned to the ER in COPI vesicles by ERD2 surveillance, which binds a C-terminal retrieval sequence ("KDEL" and variants) in the *cis* Golgi and releases it in the ER in a pH-dependent manner (14-16). Similar to other eukaryotes, ER retention of secreted

reporters or chimeric proteins can be achieved in *Plasmodium spp.* by a C-terminal KDEL or

functional variants (17-22).

In this present work, we developed and characterized a novel DiCre-based knock-

sideways tool called knockER that enables conditional fusion of KDEL to the C-terminus of

secreted proteins, subjecting the target protein to ERD2 surveillance in the *cis* Golgi to cause its

retrieval to the ER. Using this approach, we show successful conditional retrieval of five

endogenous *Plasmodium* proteins (ClpB1, RON6, EXP2, PTEX150 and HSP101) that traffic

through the ER to diverse compartments including the apicoplast, rhoptries, dense granules, and

the PV. Importantly, conditional ER retrieval of both highly expressed reporters and endogenously

tagged proteins does not generally impact fitness, indicating knockER-mediated phenotypes are

not the indirect result of an integrated stress response, thus making this system suitable for protein

loss-of-function studies. Taking advantage of the unique ability to redistribute secreted proteins

from their terminal destination to the ER, we reveal a surprising indifference to ER retrieval of

HSP101 in *P. falciparum*, in contrast to the other PTEX components, indicating HSP101 is

maintained in excess in the PV. Interestingly, redistribution of HSP101 from the PV to the ER

hypersensitized parasites to a post-translational destabilization tag that disrupts HSP101

interaction with PTEX but not to translational knockdown, providing support for an HSP101

function in the early secretory pathway in addition to its role in protein translocation across the

PVM in the assembled PTEX complex, as recently proposed (23-25). These results establish

knockER as a novel tool to study eukaryotic secreted proteins.

Results

Conditional ER retrieval of a secreted reporter does not impact parasite fitness

To facilitate conditional ER retrieval of a protein of interest, we designed a strategy based

on the dimerizable Cre recombinase (DiCre) system (26, 27) where a gene of interest is C-

terminally fused to a 3xFLAG tag and a stop codon flanked by *loxP* sequences imbedded in small

introns (28). Activation of DiCre by rapamycin results in excision of the loxP-intervening sequence,

removing the 3xFLAG tag and stop codon and bringing into frame a 3xHA tag followed by the ER-

retention sequence KDEL and a stop codon (Figure 1A). While the ER plays a crucial role in

quality control and is sensitive to added stress, several previous studies have exogenously

expressed various KDEL or SDEL fusion proteins in *P. falciparum* and *P. berghei* without apparent

impact on parasite fitness (19, 29-36). To directly ascertain whether ER retrieval of a highly

expressed secreted protein impacted parasite fitness, we first appended the knockER system to

an mNeonGreen (mNG) reporter and fused this to the *exp2* 5' UTR and signal peptide, enabling

expression from the strong *exp2* promoter and entry into the ER followed by default secretion into

the PV (37-39). This assembly was placed under control of the anhydrotetracycline (aTc)-

responsive TetR-DOZI-aptamers system to prevent expression until aTc is present in the culture

(Figure 1A) (40, 41). To combine the Bxb1 integrase system with DiCre, we first introduced a

DiCre expression cassette at the benign *pfs47* locus (27) in the *P. falciparum* NF54[attB] strain (42)

(Figure S1) and then integrated the mNG-knockER reporter construct at the *attB* site in the *cg6*

locus on chromosome 6 in this NF54[attB-DiCre] line.

Upon treatment with 10nM rapamycin for 3 hours, parasites efficiently excised the

knockER cassette (kER) with essentially complete conversion of the population within one

parasite developmental cycle (48 hrs) resulting in the expected tag switching from FLAG to HA

(Figure 1B,C). Examination of DMSO vehicle-treated parasites by live microscopy showed a

peripheral mNG signal, consistent with secretion to the PV (Figure 1D). In contrast, following

treatment with rapamycin the mNG signal was dramatically altered to an intracellular, perinuclear

pattern that co-localized with ER Tracker, confirming retention in the ER (Figure 1D).

Quantification of the distribution of mNG fluorescence between the parasite periphery and interior

showed substantial, stable re-localization following KDEL fusion, indicating robust retrieval by

ERD2 (Figure 1E, mean % internal mNG at 24hrs: 48.78 ± 5.8% in DMSO vs 76.87 ± 8.9% in

Rap, and at 168hrs: 42.54 ± 6.0% in DMSO vs 84.66 ± 6.9% in Rap). Importantly, ER retention

of mNG had no impact on parasite growth (Figure 1F) showing that knockER can successfully

retrieve a secreted reporter to the ER without impacting parasite fitness.

knockER reveals Golgi-dependent trafficking and essential apicoplast functionality of

ClpB1

Next, we applied knockER to an endogenous gene for conditional loss-of-function

mutagenesis. We choose ClpB1, a poorly characterized, nuclear-encoded AAA+ class 1

Clp/HSP100 chaperone that is targeted to the apicoplast (43). As apicoplast ablation can be

chemically rescued with isopentenyl pyrophosphate (IPP) (44), we reasoned that IPP

supplementation would provide a simple validation that any loss of parasite fitness is a direct

effect of sequestering ClpB1 away from its site of action and not an indirect result of ER stress.

ClpB1 contains both a signal peptide for ER entry and a transit peptide for apicoplast targeting

and localization of ClpB1 to this organelle was previously confirmed by both microscopy and

proteomic analyses (35, 43). Using CRISPR/Cas9 editing, we fused mNG to the C-terminus of

ClpB1 followed by the kER cassette (Figure 2A). As expected, IFA demonstrated co-localization

with the apicoplast marker ACP1, confirming ClpB1 localization to this organelle (Figure 2B).

Treatment with rapamycin resulted in efficient excision (Figure 2C) and switching to the 3xHA-

KDEL tag produced a shift in migration by Western blot consistent with an increase in molecular

weight of ~14 kDa in the rapamycin treated samples, indicating that the N-terminal transit peptide

(residues 24-152, 13.9 kDa) is no longer cleaved in the ClpB1-KDEL fusion (Figure 2D). Since

this maturation event takes place in the apicoplast (45), this suggested that ClpB1 was no longer

being trafficked to this organelle. Indeed, live microscopy showed a dramatic re-localization of

mNG from puncta to a perinuclear distribution that co-localized with ER Tracker, indicative of

successful ER retrieval (Figure 2E).

While its functional importance has not been directly tested, ClpB1 is expected to be

essential for blood-stage survival based on saturating insertional mutagenesis in *P. falciparum*

and pooled knockout screens in *P. berghei* (46, 47). Indeed, ER retention of ClpB1 was lethal and

resulted in delayed parasite death (Figure 2F), consistent with an essential role in the apicoplast

(48, 49). Moreover, supplementation with IPP fully rescued parasite growth, confirming the

observed phenotype was apicoplast-specific and not due to a disruption of ER homeostasis

following ClpB1 retention (Figure 2F). Rapamycin-treated parasites supplemented with IPP

showed dispersed, vesicular ACP1 signal consistent with plastid disruption, further indicating

ClpB1 is required for apicoplast maintenance (Figure 2G). Taken together, these results show for

the first time that ClpB1 is an indispensable component of the apicoplast machinery and illustrate

that knockER can be used for protein loss-of-function studies in *P. falciparum*.

knockER induces ER retrieval of rhoptry proteins

165   Having demonstrated that knockER is suitable to generate loss-of-function phenotypes by

protein mis-localization to the ER, we next tested this knock-sideways approach on an expanded

set of endogenous soluble and membrane proteins that traffic through the secretory pathway to

diverse destinations, including secretory organelles and the PV/PVM. We began by attempting to

retrieve RON6, a soluble rhoptry-neck protein that is injected into the PV during invasion (Figure

S2A,B). Attempts predating CRISPR technology to disrupt *Pfron6* or truncate its Cysteine-rich C-

terminus were unsuccessful (50). More recently, saturating transposon mutagenesis in *P.*

falciparum recovered a single insertion within an intron (46) expected to generate a truncation at

residue 713 of 950, leaving most of the protein intact (Figure S2A). Curiously, RON6 orthologs in

rodent malaria parasites lack the cysteine-rich region which accounts for the C-terminal ~23% of

P. falciparum RON6, roughly corresponding to the truncated region in the insertional mutant. As

RON6 was not targeted in the PlasmoGEM data set, its suspected essentiality remains uncertain

(47). To directly test the importance of RON6, we generated RON6-kER parasites, which

underwent efficient excision upon rapamycin treatment (Figure S2C,D). Rapamycin treatment

resulted in re-localization of RON6-mNG from the rhoptries to the ER and produced a minor but

reproducible growth defect (Figure S2D-F). These results suggest that RON6 may be dispensable

in the blood stage, although we cannot exclude the possibility that small amounts of RON6-KDEL

may escape ERD2 surveillance at levels sufficient to mediate a critical function. Additionally, the

minor fitness defect that results from RON6-KDEL fusion could be the indirect result of modest

ER stress induced by RON6 retention. Nonetheless, these findings clearly demonstrate that

knockER efficiently retrieves secretory proteins synthesized during the brief period of merozoite

formation at the terminal phase of intraerythrocytic development.

ER retention of EXP2 produces a lethal defect independent to loss of PVM transport

function

Having successfully retained several soluble proteins in the ER using knockER, we next

tested conditional retrieval of a membrane protein. Solute permeation across the PVM is mediated

by a nutrient-permeable channel (51, 52) whose identity was recently tied to EXP2, a PVM protein

broadly conserved among vacuole-dwelling apicomplexans (53, 54). EXP2 contains a signal

peptide followed by an N-terminal amphipathic helix and oligomerizes to form a heptameric pore

in the PVM (25, 55, 56). It is currently unknown how pore formation is specifically constrained to

the PVM to avoid perforating other membranes along the secretory pathway or the parasite

plasma membrane. While not structurally homologous to the EXP2 pore, bacterial α-helical pore

forming toxins are functionally analogous and provide a possible model for EXP2 pore formation.

These proteins are typically maintained in a monomeric state that shields a membrane-spanning

amphipathic helix, enabling soluble trafficking until contact with components of a target membrane

triggers conformational rearrangement, leading to membrane insertion and oligomerization (57).

If EXP2 traffics in a similar configuration, then its C-terminus will enter the secretory

pathway lumen and be sensitive to ER-retrieval by KDEL fusion. Indeed, we endogenously tagged

EXP2 with a version of the kER cassette lacking mNG and found that KDEL fusion disrupted

EXP2 trafficking to the PV and resulted in ER retention, confirming its C-terminus resides in the

secretory pathway lumen (Figure 3A-D). Notably, ER retrieval of EXP2 appeared to cause more

rapid parasite death than previously observed by conditional knockdown or knockout approaches

(53, 58, 59) (Figure 3E). To confirm this observation, we engineered parasites to enable DiCre-

mediated conditional knockout of EXP2 (EXP2-cKO) to allow for matched induction kinetics to

EXP2-kER (Figure S3A-C). Following activation of DiCre in synchronized, ring-stage parasites,

the EXP2 knockout progressed to the next cycle with equivalent new ring parasitemia to the

DMSO control before death in the second cycle; in contrast, EXP2-kER parasites died

immediately in the first cycle and did not form new rings (Figure S3D). To test whether parasite

death was solely due to loss of EXP2 transport functions in the PV and not an indirect result of

retaining EXP2 in the ER, we generated parasites expressing a second copy of EXP2-mNG-kER

from the *attB* locus under the control of its own promoter, leaving the endogenous EXP2 locus

unaltered (Figure S3E-G). Retrieval of this second copy of EXP2 to the ER also resulted in

parasite death, indicating that EXP2 retention in the ER is toxic independent of compromised

transport functions at the PVM (Figure 3F,G). In contrast, ER retrieval of a second copy of EXP2

lacking the amphipathic helix did not cause parasite death (Figure 3H,I and S3E-G). As expected,

prior to KDEL fusion the ΔTM version of EXP2 localized to the PV and the digestive vacuole (due

to endocytosis of PV proteins along with RBC cytosol). Surprisingly, most cells also showed

EXP2ΔTM-mNG export into the RBC cytoplasm (arrow, Figure 3H and S3H). The basis for

EXP2ΔTM-mNG export is unclear but may result from interactions with other PTEX components

that somehow result in recognition as cargo in the absence of the amphipathic helix. The

requirement for the amphipathic helix in EXP2-kER toxicity suggests that increasing EXP2 dwell

time in the early secretory pathway may trigger an unfolded protein response or cause aberrant

EXP2 pore formation in the ER membrane. Thus, this may provide a new tool for studying EXP2

pore formation but also indicates knockER may not be appropriate for loss-of-function studies of

certain membrane proteins.

ER retention of PTEX150 but not HSP101 results in loss of PTEX function in *P. falciparum*

In malaria parasites, the nutrient-permeable channel has been additionally functionalized

by the flange-like adaptor PTEX150 which docks the AAA+ chaperone HSP101 onto EXP2 to

form the <u>P</u>*lasmodium* <u>T</u>ranslocon of <u>Ex</u>ported proteins (PTEX), transforming EXP2 into a protein-

conducting pore to translocate effector proteins across the PVM and into the erythrocyte (23-25,

53, 60). Interestingly, HSP101 shows a dual localization to both the PV and the ER as opposed

to EXP2 and PTEX150, which localize exclusively to the PV and form a sub-complex independent

of HSP101 (34, 39, 61, 62). We verified this unique ER localization of HSP101 by protease

protection assays in parasite lines expressing fluorescently tagged versions of each PTEX core

component, confirming an internal, perinuclear pool of HSP101 but not EXP2 or PTEX150 (Figure

S4). This arrangement suggests that in addition to powering translocation in the assembled PTEX

complex, HSP101 may also function upstream of the vacuole, possibly initiating cargo selection

early in the secretory pathway (61). While several HSP101 conditional knockdown (24, 61) or

inactivation (23) strategies have demonstrated an essential role in protein export, it is difficult to

resolve distinct compartmental functions with these mutants given that they impact the total

protein pool and have no sub-compartmental resolution. In contrast, knockER can potentially

resolve functions across distinct compartments by redistributing a target protein pool from its

terminal destination to the ER.

In an attempt to separate HSP101 ER function from its role in the PV, we generated a kER

fusion to the endogenous copy of HSP101, expecting that redistribution of HSP101 to the ER

would be lethal but might provide insight into any ER localized function which could be preserved

compared to the phenotype of HSP101 knockdown mutants. In parallel, we also generated a

PTEX150-kER line, reasoning that since HSP101 and PTEX150 have similar expression timing

and are exclusively involved in protein export but not nutrient uptake (53), PTEX150 would provide

a control for knockER inactivation of PTEX activity independent of ER function. As expected, the

PTEX150-KDEL fusion was efficiently retrieved to the ER, producing a lethal growth defect and

accompanying block in protein export (Figure 4A-E), similar to *glmS* ribozyme-mediated

knockdown of PTEX150 (24). Importantly, ER retention of a second copy of PTEX150-kER in

parasites where the endogenous PTEX150 locus was unmodified did not impact growth,

demonstrating that the phenotype observed upon ER retention of the endogenously tagged

protein is a direct result of functional disruption of PTEX150 (Figure S5A-D).

Unexpectedly, although rapamycin-induced HSP101-KDEL fusion increased ER

localization similar to PTEX150 (Figure 4F-H), parasite growth was completely unaffected and

protein export was not detectibly impacted (Figure 4I,J), sharply contrasting with the lethal export

defect observed with HSP101 depletion or inactivation mutants (23, 24, 61). Indeed, HSP101-

kER parasites could be cultured indefinitely following excision and displayed sustained ER

retention of HSP101 (Figure 4H and S5E,F, mean Rap internal HSP101: 58.82±9.58% at 24 hours

vs 54.82±9.84% at 7 days). As an orthogonal approach to monitor protein redistribution by KDEL

fusion, we released infected RBCs with saponin to permeabilize the PVM and quantified

fluorescence via flow cytometry with or without proteinase K treatment. Again, we observed an

increase in protected fluorescence following rapamycin treatment, consistent with redistribution

of HSP101 to the ER (Figure S5G). Importantly, PCR and Western blot indicated that rapamycin-

treated parasites had undergone complete excision with the pre-excised FLAG-tagged version of

HSP101 no longer detectible (Figure 4F,G). Furthermore, both the mNG-FLAG and mNG-HA-

KDEL tagged versions of HSP101 migrated as a single band at the expected size for the full-

length fusion proteins with no apparent breakdown products, ensuring that the mNG signal

distribution observed by live microscopy reflected an intact HSP101 fusion (Figure 4G, S7B and

S8B).

While a large fraction of PTEX150 and HSP101 was clearly ER-retained following

knockER induction, some peripheral signal was still apparent after rapamycin treatment (Figure

4C,H). To determine whether the observed phenotypic differences between HSP101 and

PTEX150 were due to a difference in sensitivity to ERD2 surveillance, we quantified the internal

mNG fluorescence signal in HSP101- and PTEX150-kER parasites following KDEL fusion. In

synchronized parasites, a similar increase in internal signal was observed in both lines at 24 hours

post Rap treatment (Figure S5E, mean Rap internal HSP101 58.82±9.58% vs 56.32±18.56% for

PTEX150). Since a comparison could not be made with PTEX150-kER parasites at 7 days due

to parasite death by the second cycle following excision, we compared HSP101-kER parasites

with second copy PTEX150-kER^{attB} control parasites 7 days after excision and again observed a

comparable internal signal redistribution (Figure S5F, mean Rap internal HSP101 54.82±9.84%

vs 61.75±16.27% for PTEX150 at 7 days). As major differences in ER retrieval cannot account

for parasite insensitivity to HSP101-kER, these results indicate vacuolar HSP101 levels are

unexpectedly maintained in excess from what is required for PTEX function at the PVM.

ER retention of *P. berghei* HSP101 results in a lethal export defect

To determine if ER retention of HSP101 was similarly tolerated in other *Plasmodium*

species, we tagged HSP101 and PTEX150 with kER in a *P. berghei* rodent malaria DiCre

parasite. We also included a downstream cassette in this tagging plasmid for expression of a

secreted mRuby under control of the constitutive *Pbhsp70* promoter to label the PV (Figure S6A).

Similar to *P. falciparum* knockER mutants, rapamycin treatment produced efficient excision and

ER retrieval of PbPTEX150 and PbHSP101 (Figure 5A-C,F). Curiously, the localization pattern of

KDEL fusion proteins in *P. berghei* parasites was clearly internal to PV-mRuby but more diffuse

than in *P. falciparum*, suggesting distinctive ER morphology between these two species.

Interestingly, PbHSP101-kER and PbPTEX150-kER parasites displayed greater internal retention

than the corresponding *P. falciparum* knockER mutants (HSP101: mean % internal mNG 77.42 ±

7.72% in *P. berghei* vs 58.82±9.58% in *P. falciparum*; and PTEX150: mean % internal mNG 69.55

± 10.86% in *P. berghei* vs 56.32±18.56% in *P. falciparum*), suggesting that KDEL-mediated ER

retrieval is more stringent in rodent parasites (Figure 5D,G). To evaluate protein export, we

generated versions of the PbPTEX150-kER and PbHSP101-kER mutants where the PV-mRuby

reporter was replaced with a cassette for expression of the exported protein IBIS1 fused to mRuby

(Figure S6A) (63). In contrast to *P. falciparum*, KDEL fusion to both PbPTEX150 and PbHSP101

produced a stark block in IBIS1 export (Figure 5I,J). To determine if KDEL fusion to PbHSP101

impacted parasite fitness, parasites were induced *ex vivo* followed by overnight culture to allow

for tag switching before equal numbers of DMSO or rapamycin treated parasites were IV injected

into naïve mice. Strikingly, while DMSO controls became patent by day two post-injection, neither

KDEL-fused PbPTEX150 or PbHSP101 parasites appeared over the course of 10 days before

the experiment was terminated (Figure 5E,H and S6B).

The block in IBIS1 export suggests that PbHSP101-KDEL is sufficiently retained to

produce a lethal export defect, in contrast with PfHSP101-KDEL; however, phenotypes can be

masked by the rich resources available in culture (64, 65) and many exported effectors critical for

survival in the vertebrate host are dispensable *in vitro* (12) where parasites do not have to contend

with host defenses (66, 67). To distinguish whether the fitness defect incurred by ER retention of

PbHSP101 resulted simply from parasite death or from the unique pressures imposed by the host,

we evaluated parasite development of synchronized ring-stage cultures *ex vivo*. Importantly, both

PbHSP101-kER and PbPTEX150-kER parasites also failed to develop into terminal schizonts *in*

vitro following rapamycin treatment, indicating the *in vivo* fitness defect is not merely the result of

host defenses (Figure S6C-E). Taken together, our data show that ER-retention of HSP101-KDEL

is more efficient in *P. berghei,* resulting in a lethal defect in protein export.

ER retention of HSP101 hypersensitizes parasites to a destabilization tag that inhibits

HSP101-PTEX complex formation

The lack of an observable phenotype after ER-retention of HSP101 in *P. falciparum*

suggests that the level of retention does not reach a critical threshold required to impact HSP101

function at the PV. To determine the amount of HSP101 needed to support *P. falciparum* growth

in vitro, we combined knockER with the TetR-DOZI-aptamers system to generate a HSP101-kER

line that also allows for titration of HSP101 expression with aTc (Figure 6A). Like the original

HSP101-kER line, rapamycin-treated HSP101-kER[TetR-DOZI] parasites exhibited ER retention of

HSP101 without impacting parasite growth or export of representative PEXEL and PNEP proteins

(Figure 6B-D). However, HSP101 depletion following aTc washout resulted in a robust block in

protein export and parasite death (Figure 6D and S7A,B), consistent with previous HSP101

knockdown or inactivation studies (23, 24, 61). To determine the level of total HSP101 reduction

sufficient to impact growth, we grew synchronized parasites at a range of aTc concentrations

following DMSO or rapamycin treatment and measured mNG fluorescence via flow cytometry in

parallel with parasite growth (Figure 6E,F). Interestingly, rapamycin-treated parasites grown at

higher aTc concentrations displayed a small but reproducible increase in total HSP101, consistent

with a reduction in HSP101 turnover by endocytosis from the PV to the digestive vacuole following

redistribution to the ER (Figure 6E). More importantly, we determined that a ~70% reduction in

total HSP101 levels was necessary to impact parasite growth, showing that a relatively small

amount of HSP101 is needed for survival (Figure 6E,F, 5nM aTc), consistent with redistribution

of HSP101 by knockER not meeting the threshold required to impact HSP101 PV function and

parasite fitness in *P. falciparum*.

If HSP101 functions exclusively in the PV, then knockER-mediated depletion of vacuolar

HSP101 levels should hypersensitize parasites to HSP101 knockdown. However, dose-response

to aTc was unaltered by rapamycin-treatment (Figure 6G) and the growth defect of KDEL-fused

HSP101 parasites maintained at 5 nM aTc was indistinguishable from the DMSO control even

over an extended 10-day growth assay (Figure 6F), possibly suggesting that the PV-localized

function of HSP101 may not fully account for its role in parasite fitness. To ensure depletion of

HSP101 from the vacuole by knockER sensitizes parasites to compromised HSP101 PV function,

we generated an HSP101-knockER line where mNG was replaced by a ligand-sensitive DHFR-

based destabilization domain (DDD) (Figure 6H) (23). In the absence of trimethoprim (TMP), the

DDD blocks HSP101 interaction with EXP2/PTEX150 but not cargo binding, specifically targeting

function of the assembled PTEX complex in the PV (23). In contrast to translational knockdown,

HSP101-kER[DDD] parasites were hypersensitized to reduction of TMP levels following rapamycin-

treatment, indicating a synergistic effect when the reduction of HSP101 PV levels is combined

with disruption of PTEX complex formation (Figure 6I and S7C-E). Taken together, these results

illustrate how knockER can be used to study protein function with sub compartmental resolution

and may support a distinct HSP101 activity in the ER upstream of its PVM translocation function

at the assembled PTEX complex.

Discussion

Several genetic tools are now available for protein loss-of-function studies in *Plasmodium*

spp. (68), including knock-sideways for conditional sequestration of a target protein away from its

normal site of function (69, 70). While this system is not currently amenable to parasite proteins

that enter the secretory pathway, conditional control of secretory traffic has been achieved in *P.*

falciparum by ligand-mediated alteration of the N-terminal transit peptide that controls apicoplast

targeting, enabling redirection of plastid proteins into the default secretory pathway (71).

Additionally, conditional ER release systems have been developed in model eukaryotes based

on ER retrieval machinery (72). In contrast, knockER provides a simple, DiCre-mediated knock-

sideways strategy that exploits the ERD2/KDEL receptor (KDELR) system for conditional ER

retrieval, providing a generalizable approach for interrogating secreted proteins in genetically

tractable eukaryotes.

In this study, we demonstrate knockER simultaneously allows for trafficking and loss-of-

function studies. The ability to determine whether a protein traffics from the ER through the *cis*

Golgi is an important consideration for nuclear-encoded apicoplast proteins since Golgi-

dependent and -independent trafficking routes have been reported (31, 33, 73). While these

apicoplast trafficking studies were carried out with reporters and chimeras, knockER allows for

monitoring *cis* Golgi transit of endogenous proteins. The sensitivity of ClpB1 to KDEL-mediated

ER retrieval demonstrates a Golgi-dependent trafficking route to the apicoplast. Moreover, we

show for the first time that this AAA+ chaperone is essential for apicoplast maintenance. ClpB1 is

one of four Clp/HSP100 chaperones encoded by *Plasmodium spp.*, three of which localize to the

parasite plastid while the fourth, HSP101 (previously known as ClpB2) localizes to the PV and

ER (43). The other apicoplast-targeted members of this quartet include the nuclear-encoded

ClpC, which mediates regulated proteolysis together with the serine protease ClpP (74, 75), and

ClpM, now the only uncharacterized member of this group and one of the few proteins encoded

on the apicoplast genome. Similar to the well-studied function of its closest orthologs (76), ClpB1

likely acts as a disaggregase that, together with ClpC/P and ClpM, forms a critical proteostasis

system required to maintain and regulate the apicoplast proteome (77).

ER perturbations can trigger an integrated stress response leading to cell arrest or death

(78, 79). *Plasmodium* possesses one of the widely conserved ER stress sensors, the PERK-

eIF2α pathway which halts protein translation and flux through the secretory pathway (20, 80-83).

Importantly, conditional ER retrieval of proteins produced from their endogenous locus or by

heterologous expression was not generally toxic, enabling loss-of-function studies with knockER.

The single notable exception was EXP2, which rapidly killed parasites when retained in the ER

independent to loss of its PVM transport function, indicating knockER experiments should be

controlled for indirect effects. EXP2-KDEL toxicity was dependent on the amphipathic helix that

forms the membrane-spanning region in the oligomeric EXP2 pore (25), suggesting that

accumulation of EXP2 in the ER might trigger an unfolded protein response or lead to premature

pore formation in the early secretory pathway. More work is required to determine if this outcome

is specific to EXP2 but the observation that parasites tolerate SDEL-mediated retention of several

TM domain-containing exported proteins indicates this is not a general problem for unnatural ER-

retrieval of membrane proteins (84). Indirect fitness defects aside, retrieval of EXP2-KDEL

demonstrates that its C-terminus enters the secretory pathway lumen, consistent with a soluble

EXP2 trafficking state that precedes membrane integration at the PVM. Regardless of the basis

for EXP2-kER toxicity, these experiments highlight that knockER can also be used to investigate

membrane topology. While ER-resident type I membrane proteins utilize a different mechanism

for retrieval as their C-termini face the cytosol of the cell (85, 86), ERD2/KDELR can retrieve

membrane proteins whose C-termini face the lumen (87) and thus knockER should be able to

distinguish the topology of single and multi-pass transmembrane domain-containing proteins.

A unique feature of knockER is the ability to redistribute secreted proteins from their

terminal destination to the ER. We tagged both HSP101 and the translocon adaptor PTEX150

with knockER, reasoning that ER retrieval of either protein would produce a lethal export defect

but that differential impacts on exported cargo trafficking might be exposed. Unexpectedly, ER

retrieval of HSP101 had no impact on *P. falciparum* growth or export of representative PEXEL

and PNEP proteins *in vitro*. Similar PV depletion of PTEX150 produced the anticipated lethal

export block, revealing that HSP101 PV levels are uniquely maintained in excess. As a AAA+

ATPase, HSP101 is the only enzyme in the PTEX core complex and its activity may exceed what

is required to support parasite fitness, at least during *in vitro* culture, as previously seen for the

aspartic protease Plasmepsin V that licenses proteins for export in the ER (88, 89). Interestingly,

parallel experiments in *P. berghei* resulted in greater retention and produced a lethal export defect

for both PTEX150 and HSP101, showing knockER is also suitable for conditional mutagenesis in

this important model species and suggesting retrieval may be more stringent in *P. berghei*, at

least for some proteins. Notably, the strength of ERD2/KDELR binding between different C-

terminal retrieval motifs varies by as much as 10-fold in higher eukaryotes (16). While KDEL-

mediated retrieval was generally robust across the diverse set of reporters and endogenous

proteins surveyed in *P. falciparum* and *P. berghei*, a range of functional variants have been

validated or can be inferred in *Plasmodium spp.* (19) and other retrieval motifs may enable tuning

the strength of retention by knockER.

Intriguingly, redistribution of HSP101 from the PV to the ER hypersensitized parasites to

DDD-mediated HSP101 inactivation but not translational knockdown. While TetR-DOZI

knockdown depletes the entire HSP101 pool, HSP101-DDD acts post-translationally by disrupting

complex formation with EXP2/PTEX150 but not cargo recognition (23), specifically targeting

HSP101 function in PVM translocation. While these results do not enable strong conclusions

about potential extra vacuolar activity of HSP101, they suggest that the PV-localized function may

not fully account for HSP101's contribution to parasite fitness, in line with a hypothesized role in

exported cargo recognition and/or trafficking at the ER (61). Notably, IBIS1 was not retained in

the ER with PbHSP101-KDEL but accumulated in the PV and was not exported into the RBC,

indicating that HSP101 is not required to directly chaperone cargo from the ER to the PV. Future

work dissecting the specific extra-vacuolar role(s) of HSP101 may resolve the long-standing

question of how early events in this pathway that mark proteins for export, such as PEXEL

processing during ER entry, are connected to cargo recognition and ultimate PVM translocation

into the host cell at the assembled PTEX complex.

Materials and Methods

Parasite culture. *P. falciparum* NF54[attB-DiCre] and derivatives were cultured in RPMI 1640 medium

supplemented with 27 mM sodium bicarbonate, 11 mM glucose, 0.37 mM hypoxanthine, 10 µg/ml

gentamicin and 0.5% Albumax I (Gibco). Parasites were maintained in deidentified, Institutional

Review Board-exempt RBCs obtained from the American National Red Cross. *P. berghei* ANKA

clone 2.34 and derivatives were maintained in Swiss Webster mice (Charles River). All

experiments involving rodents were reviewed and approved by the Iowa State University

Institutional Animal Care and Use Committee.

Plasmids and genetic modification of *P. falciparum* and *P. berghei*. All cloning was carried

out with NEBuilder HIFI DNA assembly (NEB). Primer sequences are given in Table S1. To

generate NF54[attB-DiCre] parasites, a DiCre expression cassette was inserted into the *pfs47* gene in

NF54[attB] parasites (42) using a marker-free CRISPR/Cas9 editing strategy as described (27). Loss

of the endogenous *pfs47* locus was monitored with primers P1/2 and P3/4 and integration was

confirmed using primers P1/5 and P6/4. To confirm DiCre activity in this line, a loxP-flanked mNG

coding sequence with a 3xHA tag was PCR amplified using primers P19/20 and inserted between

AvrII and AflIII in pLN-ENR-GFP cite (90), resulting in the plasmid JRB416. This plasmid was co-

transfected with pINT (90) into NF54[attB-DiCre] parasites and selection was applied with 2.5 µg/ml

blasticidin-S 24 hrs post-transfection.

To generate the kER fusion cassette, homology flanks targeting the 3' end of *exp2* were

amplified from pyPM2GT-EXP2-3xHA-GFP11 (53) using P21/22 and inserted between XhoI and

AvrII in pPM2GT-HSP101-3xFLAG (53). The loxPint sequence (28) was amplified with P23/24

and inserted immediately upstream of the 3xFLAG tag at AvrII in this vector. A partial loxPint

containing the loxP sequence and 3' portion of the intron followed by a 3xHA-KDEL tag was then

amplified with P25/26 and inserted into EagI, resulting in the plasmid JRB492. Finally, the mNG

coding sequencing was amplified from pyPM2GT-EXP2-mNG (91) with P27/28 and inserted into

AvrII in JRB492, resulting in plasmid JRB508. For generation of the *exp2* DiCre conditional

knockout line, a recoded version of the *exp2* coding sequence with a 3' 3xHA tag and flanked by

loxP sites was amplified using P76/77 from plasmid pEXP2[apt] (53) and inserted between

AvrII/EagI in plasmid JRB492. A 5' flank immediately upstream of the *exp2* start codon was then

amplified from *P. falciparum* gDNA using P78/79 and inserted between AflIII/AvrII, replacing the

AflIII site with AscI and resulting in the plasmid JRB467. This plasmid was linearized with AscI and

co-transfected with pUF-Cas9-EXP2-CT-gRNA into NF54[attB-DiCre]. Selection was applied with 5

nM WR99120 24 hrs post-transfection and clonal lines were isolated by limiting dilution after

parasites returned from selection.

To generate the SP-mNG-kER reporter, the *exp2* promoter and signal peptide were

amplified from parasite genomic DNA with P29/30 and the mNG-KER cassette was amplified from

JRB508 with P31/32 and these amplicons were assembled together in a second PCR reaction

with P29/32 and inserted into AscI/ApaI in pKD (41), resulting in the plasmid JRB543. For

expression of a second copy of EXP2-kER from the attB locus, the *exp2* promoter and full-length

gene were amplified from genomic DNA with P29/33 and inserted between AscI/BamHI in

JRB543, resulting in the plasmid JRB526. For expression of a second copy of EXP2-kER lacking

the amphipathic helix, a recoded portion of the *exp2* coding sequence beginning at codon 86

(after the amphipathic helix and immediately before the helical body domain) was amplified from

pEXP2^{apt} (53) and inserted at BamHI after the EXP2 signal peptide in JRB543, resulting in the

plasmid JRB544. These plasmids were each co-transfected with pINT and selected with 2.5 µg/ml

blasticidin-S as described above. To relieve translational repression by TetR-DOZI, parasite

cultures were supplemented with 500 nM aTc.

For generation of ClpB1-kER and RON6-kER lines, a Cas9 gRNA target was chosen just

downstream (*clpB1*, ATTTTTAGAAAAATATTATG) or just upstream (*ron6*,

TAATACATACAACGTACCAA) of the stop codon and the gRNA seed sequences were

synthesized as primers P34 and P35, respectively, and inserted into the AflII site of pAIO3 (39),

resulting in plasmids JRB143 and MAF22, respectively. To target the 3' end of *clpB1* or *ron6*, a 5'

homology flank (up to but not including the stop codon) was amplified from genomic DNA using

P36/37 or P38/39, incorporating synonymous shield mutations in the gRNA target sites. A 3'

homology flank was amplified using P40/41 or P42/43. The corresponding 5' and 3' flank

amplicons were assembled in a second PCR reaction using P40/37 and P42/39 and inserted

between XhoI/AvrII in JRB508, resulting in plasmids MAF20 and MAF27, respectively. These

plasmids were linearized with AflII and co-transfected with the corresponding Cas9/gRNA plasmid

into NF54$^{attB-DiCre}$. Selection was applied with 5 nM WR99120 24 hrs post-transfection and clonal

lines were isolated by limiting dilution after parasites returned from selection.

For generation of PTEX150-kER and HSP101-kER lines, flank assemblies targeting the

3' end of each gene were amplified using P44/45 from pyPM2GT-PTEX150-3xHA-GFP11 (53) or

P46/47 from pyPM2GT-HSP101-3xHA-GFP11 (53) and inserted between XhoI/AvrII in JRB508,

resulting in plasmids MAF16 and MAF17, respectively. These plasmids were linearized with AflII

and co-transfected with pAIO-PTEX150-CT-gRNA or pAIO-HSP101-CT-gRNA (53), selected with

5 nM WR99120 and cloned after parasites returned. For generation of the EXP2-kER line, plasmid

JRB492 (described above) was linearized and co-transfected with pUF-Cas9-EXP2-CT-gRNA

(53) and selected with 5 nM WR99120. For generation of HSP101-kER$^{TetR-DOZI}$ lines, flank

assembles were amplified using P48/49 and inserted between AscI/BamHI in JRB526 (described

above), resulting in plasmid MAF41. This plasmid was linearized with AflII and co-transfected with

pAIO-HSP101-CT-gRNA, selected with 2.5 µg/ml blasticidin-S and 500 nM aTc, and cloned after

parasites returned. For generation of HSP101-kER^{DDD} lines, the DDD was amplified from plasmid

pPM2GT-HSP101-3xFlag-DDD (53) using P70/71 and inserted between AvrII/BsrGI in MAF17,

replacing the mNG and resulting in plasmid MAF43. This plasmid was linearized with AflII and co-

transfected with pAIO-HSP101-CT-gRNA. Selection was applied with 10 µM TMP 24 hrs post-

transfection.

For expression of a second copy of PTEX150-kER from the *attB* locus, the *ptex150*

promoter and full-length gene were amplified from genomic DNA with P50/51 and the mNG-KER

cassette was amplified from JRB508 with P52/53. These amplicons were assembled together in

a second PCR reaction using P50/53 and inserted between ApaI/AflII in pLN-ENR-GFP (90),

resulting in the plasmid MAF39. This plasmid was co-transfected with pINT, selected with 2.5

µg/ml blasticidin-S and parasites were cloned after returning.

To generate the line bearing endogenous EXP2-mRuby and HSP101-mNG tags, the

plasmid pbEXP2-mRuby3 (92) was first linearized with AflII and co-transfected with pUF-Cas9-

EXP2-CT-gRNA (53) into NF54$^{attB-DiCre}$. Selection was applied with 2.5 µg/ml blasticidin-S. Next,

the mNG coding sequence was amplified using P54/55 and inserted between AvrII/EagI in

pPM2GT-HSP101-3xFLAG (53) to replace the FLAG tag with mNG, resulting in the plasmid

JRB325. This plasmid was linearized with AflII and co-transfected with pAIO-HSP101-CT-gRNA

(92) into the NF54^{attB-DiCre}::EXP2-mRuby parasites. Selection was applied with 5nM WR99120

resulting in the line NF54^{attB-DiCre}::EXP2-mRuby+HSP101-mNG.

For generation of PTEX150-kER and HSP101-kER lines in *P. berghei*, the plasmid pBAT

(93) was digested with MluI/XhoI and the primer P56 was used as an insert to remove the

intervening AvrII, SnaBI and XhoI sites from the MCS. Next, the mRuby3 coding sequence fused

to the PbEXP2 signal peptide was amplified with P57/58 and inserted between SwaI/BamHI,

replacing the GFP. The mNG-kER tagging sequence was amplified from plasmid JRB508 using

P59/60 and inserted between SacII/SphI, replacing the mCherry-3xMYC. Finally, 5' flanks to tag

P. berghei hsp101 or *ptex150* with mNG-kER were amplified from *P. berghei* genomic DNA using

P61/62 or P63/64, respectively, and 3' flanks were amplified using P65/66 or P67/68, respectively.

The corresponding 5' and 3' flank amplicons were assembled with an intervening AflII site in a

second PCR reaction using P65/62 or P67/64, respectively, and inserted at SacII, resulting in

plasmids pPbHSP101-kER+SP-mRuby and pPbPTEX150-kER+SP-mRuby. The plasmids were

linearized at AflII and transfected as described (94) into the marker-free HP DiCre line (95).

Transfected parasites were reinjected into naïve mice and selection was applied with 0.07 mg/ml

pyrimethamine in drinking water provided *ad libitum* 24 hours later. After returning from selection,

transfected populations were cloned by limiting dilution IV injection into naïve mice to generate

isogenic populations designated PbHSP101-kER and PbPTEX150-kER. To monitor export with

an IBIS1-mRuby fusion, the *ibis1* gene and promoter were amplified from *P. berghei* gDNA using

P72/73 and the *mruby* cds was amplified using P74/75. These amplicons were assembled in a

second PCR reaction using P72/75 and inserted between PvuII/BamHI, resulting in plasmids

pPbHSP101-kER+IBIS1-mRuby and pPbPTEX150-kER+IBIS1-mRuby. Plasmids were

linearized at AflII and transfected as above.

P. falciparum growth assays. Growth assays were performed as described previously (22).

Briefly, parasites were treated with either DMSO or 10 nM rapamycin for 3 hrs and washed 1X.

Then, parasites were seeded between 0.2-1% parasitemia in triplicate and monitored every 48

hrs by flow cytometry on an Attune NxT (ThermoFisher) by nucleic acid staining with PBS

containing 0.8 µg/ml acridine orange. For IPP rescue experiments, ClpB1-kER cultures were

supplemented with 200 µM IPP. For kER fusions expressed from the *attB* locus, cultures were

supplemented with 1 µM aTc 24 hrs prior to rapamycin treatment to relieve TetR-DOZI-aptamer

translational repression. For HSP101-kER[TetR-DOZI] parasites, aTc was removed by washing

parasites 7X with 13 mL of media. 500 nM aTc was then supplemented back to control parasites.

As required, parasites were sub-cultured to avoid high parasite density, and relative parasitemia

at each time point was back-calculated based on actual parasitemia multiplied by the relevant

dilution factors. Data were displayed using Prism 9 (GraphPad Software).

For EC_{50} shift assays of HSP101-kER[TetR-DOZI] and HSP101-kER[DDD], asynchronous

parasites were treated with DMSO or rapamycin before removal of aTc or TMP. After agonist

washout as described above, parasites were seeded in triplicate at 0.5% parasitemia with 2-fold

dilutions starting from 500 nM aTc or 250 nM TMP and parasite growth was monitored after 72

hrs.

***P. berghei* kER experiments.** Parasite-infected blood was collected by cardiac puncture when

parasitemia reached 3-4% and cultured in RPMI supplemented with 20% FBS. Cultures were

treated with 10 nM rapamycin or a DMSO vehicle control for 3 hours followed by a media change.

After 18 hours of additional culture, parasites were harvested for diagnostic PCR, live

fluorescence microscopy and Western blot. Equal numbers of DMSO or rapamycin-treated

infected RBCs (~10,000 parasites per injection) were IV injected into naïve mice and parasitemia

was monitored by giemsa-stained blood smear every two days.

For synchronized, *ex vivo* growth assays, parasites were IP injected into mice and allowed

to reach 2-3% parasitemia. Parasites gathered from cardiac puncture were then treated with

DMSO or 10nM rapamycin 3-4hrs, washed, and allowed to mature into schizonts overnight

without further rapamycin treatment. The mature schizonts were magnet purified and tail vein

injected into naïve mice treated with phenylhydrazine (1mg/mice) (Sigma, 114715) to allow

parasites to egress and invade new RBCs in mice. After 3 hrs to allow parasite egress and re-

invasion, synchronous ring-stage parasites were collected by cardiac puncture and allowed to

mature overnight *in vitro*. At 24 hrs post injection, parasite development was assessed in Giemsa-

stained smears and DNA content was evaluated by Hoechst 34580 staining (1:5,000) on an

Attune NxT flow cytometer. To monitor export of IBIS1-mRuby, parasites were IP injected and

allowed to grow to 2-3% parasitemia and imaged from tail-blood. Mice were then injected IP with

rapamycin (1 mg/kg) and parasites were observed again 24hrs later.

Microscopy. For live microscopy, *P. falciparum* parasites were treated with either DMSO or 10

nM rapamycin and viewed 24-48 hrs later. Hoechst (33342, Invitrogen; 1:10,000) was used to

visualize the nucleus by incubating with parasites 2-5 min prior to visualization. ER-Tracker Red

(Invitrogen; 1:1000) was used to visualize the ER by incubating with parasites 30 min at 37°C

prior to visualization. For constructs expressed from the *attB* locus under TetR-DOZI-aptamers

control, parasites were grown in 500 nM aTc for 24 hrs prior to incubation with DMSO or

rapamycin to allow expression. For RON6 parasites, DMSO and rapamycin-treated schizonts

were incubated with 50nM ML10 (obtained through BEI resources, NIAID, NIH and supplied by

LifeArc, Stevenage, UK: MRT-0207065 (ML10), NR-56525) for 4.5 hrs and visualized by

microscopy. For *P. berghei*, synchronized ring-stage parasites were isolated via cardiac puncture,

treated with DMSO or rapamycin for 3 hrs and allowed to develop overnight in culture before

visualization. For quantification of fluorescence signal, MFI was calculated using the measured

Integrated Density value from Fiji (96). The internal MFI was calculated using an ROI within the

parasite PV and subtracting that from the whole parasite MFI using an ROI outside the PV

boundary. The percent internal fluorescence was then calculated by dividing the internal MFI by

the whole parasite MFI.

For protease protection assays on live cells, parasites were treated with 0.03% saponin

(Sigma) in 1X PBS for 15 min at 4°C. Purified parasites where then washed once with 5 mL of 1X

PBS and once with 5 mL of digestion buffer (50 mM Tris pH 7.5, 150 mM NaCl, 1 mM $CaCl_2$) (61).

Parasite samples where then split into two identical tubes, one of which was treated with 20 µg/mL

of Proteinase K (740506.75, Macherey-Nagel), and incubated at 37°C for 15 min. Proteinase K

was removed and samples resuspended with digestion buffer plus protease inhibitor cocktail (Halt

78429, Thermo Scientific) before visualization.

For IFAs of the EXP2- and ClpB1-kER lines, parasites were treated with either DMSO or

10 nM rapamycin and fixed 24 hrs or 144 hrs later as described previously (22). For HSP101-

$kER^{TetR\text{-}DOZI}$ lines, parasites were synchronized with 5% sorbitol, treated with either DMSO or 10

nM rapamycin, grown with or without 500 nM aTc, and fixed 48 hrs later. Briefly, parasites were

fixed with Fixative Solution (4% paraformaldehyde, 0.0075% glutaraldehyde in PBS) for 25 min

at room temperature in the dark. Parasites were then permeabilized with 0.1% Triton in PBS for

10 minutes at room temperature while rocking. Parasites were washed 1X with PBS and then

blocked with 3% BSA in PBS for 1hr at room temperature. Primary antibody was applied in 3%

BSA solution for 1 hr at room temperature, washed 2X with PBS, followed by secondary antibody

incubation for 1 hr at room temperature. After 2X washes with PBS, parasites were adhered onto

Poly-L-lysine coated coverslips and then mounted onto slides. For parasites probed with SBP1,

parasites were acetone fixed as in (39). Briefly, thin smears were immersed in 100% acetone for

2 min. Smears were washed 1X with PBS and then blocked with 3% BSA in PBS for 1 hr at room

temperature. Primary antibody was applied in 3% BSA solution for 1 hr at room temperature,

washed 2X with PBS, followed by secondary antibody incubation for 1 hr at room temperature.

After 2X washes with PBS, slides were mounted with DAPI, covered with a coverslip and sealed.

The antibodies used for IFA were: mouse anti-HA antibody (clone 16B12, BioLegend; 1:200),

rabbit polyclonal anti-HA SG77 (ThermoFisher; 1:500), rabbit anti-PfGRP78 (BiP) (MRA-1246,

BEI Resources, NIAID, NIH; 1:100), mouse monoclonal anti-FLAG clone M2 (Sigma-Aldrich;

1:200), rabbit monoclonal anti-Flag 8H8L17 (ThermoFisher; 1:500), mouse anti-HRP2 clone

2G12 (1:500) (97), rabbit polyclonal anti-SBP1 BR28 (1:500) (98) and rabbit polyclonal anti-ACP1

(1:500 IFA) (37, 99). Secondary antibodies used were anti-mouse antibody conjugated to Alexa

Fluor 488 or 546 and anti-rabbit antibody conjugated to Alexa Fluor 488, (Life Technologies;

1:1000). Parasites were mounted with ProLong diamond with 4',6'-diamidino-2-phenylindole

(DAPI) (P36931; Invitrogen) and imaged on an Axio Observer 7 equipped with an Axiocam 702

mono camera and Zen 2.6 Pro software (Zeiss) using the same exposure times for all images

across sample groups and experimental replicates. Image processing, analysis, and display were

performed using Zeiss Blue software and Adobe Photoshop. Equivalent adjustments to brightness

and contrast were made within sample groups for display purposes.

Quantification of protein knockdown. To evaluate the level of HSP101 protein knockdown,

HSP101-kER$^{TetR-DOZI}$ parasites were synchronized via magnet purification and treated with DMSO

or rapamycin before removal of aTc. After aTc removal, parasites were seeded at ~1%

parasitemia with different dilutions of aTc and allowed to grow for 48 hrs. Using an Attune NxT

flow cytometer, infected RBCs were gated using Hoechst 34580 (Invitrogen; 1:5,000) and mNG

fluorescence of infected RBCs was concurrently measured. The same procedure was applied for

saponin-released parasites after protease treatment as described above.

Western blotting. Western blotting was performed as described previously (39, 53). Briefly,

parasite-infected RBCs were selectively permeabilized by treatment with ice-cold 0.03% saponin

in PBS for 15 min followed by lysis with RIPA and centrifugation to remove hemozoin. For

constructs expressed from the *attB* locus under control of TetR-DOZI-aptamers, parasites were

grown in 500 nM aTc for 24 hrs prior to incubation with DMSO or rapamycin to allow for

expression. For the HSP101-kER$^{TetR-DOZI}$ lines, parasites were treated with DMSO or rapamycin

before removal of aTc. After aTc removal, parasites were seeded at ~1% parasitemia with

different dilutions of aTc and allowed to grow for 48 hrs before processing. The antibodies used

in this study were mouse anti-HA antibody (clone 16B12, BioLegend; 1:500), rabbit polyclonal

anti-HA SG77 (ThermoFisher; 1:500), mouse anti-FLAG (Clone M2, Sigma-Aldrich; 1:500), rabbit

polyclonal anti-*Plasmodium* Aldolase ab207494 (Abcam; 1:500) and mouse monoclonal anti-

EXP2 clone 7.7 (1:500) (100). The secondary antibodies used were IRDye 680CW goat anti-

rabbit IgG and IRDye 800CW goat anti-mouse IgG (Li-COR Biosciences; 1:20,000). Western blot

images were acquired using an Odyssey Clx infrared imaging system and processed with Image

Studio software (Li-COR Biosciences). Full length versions of cropped blots can be seen in Figure

S8.

Acknowledgements

This work was supported by NIH grant HL133453 to JRB. The funders had no role in study design,

data collection and interpretation, or the decision to submit the work for publication. We thank J.

McBride, D. Cavanaugh and EMRR for the EXP2 antibody and A. Waters for the HP DiCre *P.*

berghei parasites.

Author contributions

689   M.A.F. and J.R.B designed research; M.A.F., T.H., L.J.C. and J.R.B performed research; M.A.F.

690   and J.R.B analyzed data; M.A.F. and J.R.B. wrote the paper.

References

1.   G. G. van Dooren, B. Striepen, The algal past and parasite present of the apicoplast.
Annu Rev Microbiol **67**, 271-289 (2013).

2.   O. A. Akuh, R. Elahi, S. T. Prigge, F. Seeber, The ferredoxin redox system - an essential
electron distributing hub in the apicoplast of Apicomplexa. *Trends Parasitol* **38**, 868-881
(2022).

3.   M. M. Cova, M. H. Lamarque, M. Lebrun, How Apicomplexa Parasites Secrete and Build
Their Invasion Machinery. *Annu Rev Microbiol* **76**, 619-640 (2022).

4.   J. M. Matz, J. R. Beck, M. J. Blackman, The parasitophorous vacuole of the blood-stage
malaria parasite. *Nat Rev Microbiol* **18**, 379-391 (2020).

5.   M. Garten, J. R. Beck, Structured to conquer: transport across the Plasmodium
parasitophorous vacuole. *Curr Opin Microbiol* **63**, 181-188 (2021).

6.   T. F. de Koning-Ward, M. W. Dixon, L. Tilley, P. R. Gilson, Plasmodium species: master
renovators of their host cells. *Nat Rev Microbiol* **14**, 494-507 (2016).

7.   J. R. Beck, C. M. Ho, Transport mechanisms at the malaria parasite-host cell interface.
PLoS Pathog **17**, e1009394 (2021).

8.   T. J. Sargeant *et al.*, Lineage-specific expansion of proteins exported to erythrocytes in
malaria parasites. *Genome Biol* **7**, R12 (2006).

9.   J. A. Boddey *et al.*, Role of plasmepsin V in export of diverse protein families from the
Plasmodium falciparum exportome. *Traffic* **14**, 532-550 (2013).

10.   C. van Ooij *et al.*, The malaria secretome: from algorithms to essential function in blood
stage infection. *PLoS Pathog* **4**, e1000084 (2008).

11.   J. Schulze *et al.*, The Plasmodium falciparum exportome contains non-canonical
PEXEL/HT proteins. *Mol Microbiol* **97**, 301-314 (2015).

12.   T. K. Jonsdottir, M. Gabriela, B. S. Crabb, T. F. de Koning-Ward, P. R. Gilson, Defining
the Essential Exportome of the Malaria Parasite. *Trends Parasitol* **37**, 664-675 (2021).

13.   A. Florentin, D. W. Cobb, H. M. Kudyba, V. Muralidharan, Directing traffic: Chaperone-
mediated protein transport in malaria parasites. *Cell Microbiol* **22**, e13215 (2020).

14.   S. Munro, H. R. Pelham, A C-terminal signal prevents secretion of luminal ER proteins.
Cell **48**, 899-907 (1987).

15.   P. Brauer *et al.*, Structural basis for pH-dependent retrieval of ER proteins from the Golgi
by the KDEL receptor. *Science* **363**, 1103-1107 (2019).

16.   A. Gerondopoulos *et al.*, A signal capture and proofreading mechanism for the KDEL-
receptor explains selectivity and dynamic range in ER retrieval. *Elife* **10** (2021).

17.   H. G. Elmendorf, K. Haldar, Identification and localization of ERD2 in the malaria
726   parasite Plasmodium falciparum: separation from sites of sphingomyelin synthesis and
727   implications for organization of the Golgi. *EMBO J* **12**, 4763-4773 (1993).

728   18.   N. La Greca, A. R. Hibbs, C. Riffkin, M. Foley, L. Tilley, Identification of an endoplasmic
729   reticulum-resident calcium-binding protein with multiple EF-hand motifs in asexual
730   stages of Plasmodium falciparum. *Mol Biochem Parasitol* **89**, 283-293 (1997).

731   19.   S. Kulzer, N. Gehde, J. M. Przyborski, Return to sender: use of Plasmodium ER retrieval
732   sequences to study protein transport in the infected erythrocyte and predict putative ER
733   protein families. *Parasitol Res* **104**, 1535-1541 (2009).

734   20.   H. M. Kudyba *et al.*, The endoplasmic reticulum chaperone PfGRP170 is essential for
735   asexual development and is linked to stress response in malaria parasites. *Cell Microbiol*
736   **21**, e13042 (2019).

737   21.   D. W. Cobb *et al.*, A redox-active crosslinker reveals an essential and inhibitable
738   oxidative folding network in the endoplasmic reticulum of malaria parasites. *PLoS*
739   *Pathog* **17**, e1009293 (2021).

740   22.   M. A. Fierro *et al.*, An Endoplasmic Reticulum CREC Family Protein Regulates the
741   Egress Proteolytic Cascade in Malaria Parasites. *mBio* **11** (2020).

742   23.   J. R. Beck, V. Muralidharan, A. Oksman, D. E. Goldberg, PTEX component HSP101
743   mediates export of diverse malaria effectors into host erythrocytes. *Nature* **511**, 592-595
744   (2014).

745   24.   B. Elsworth *et al.*, PTEX is an essential nexus for protein export in malaria parasites.
746   *Nature* **511**, 587-591 (2014).

747   25.   C. M. Ho *et al.*, Malaria parasite translocon structure and mechanism of effector export.
748   *Nature* **561**, 70-75 (2018).

749   26.   C. R. Collins *et al.*, Robust inducible Cre recombinase activity in the human malaria
750   parasite Plasmodium falciparum enables efficient gene deletion within a single asexual
751   erythrocytic growth cycle. *Mol Microbiol* **88**, 687-701 (2013).

752   27.   E. Knuepfer, M. Napiorkowska, C. van Ooij, A. A. Holder, Generating conditional gene
753   knockouts in Plasmodium - a toolkit to produce stable DiCre recombinase-expressing
754   parasite lines using CRISPR/Cas9. *Sci Rep* **7**, 3881 (2017).

755   28.   M. L. Jones *et al.*, A versatile strategy for rapid conditional genome engineering using
756   loxP sites in a small synthetic intron in Plasmodium falciparum. *Sci Rep* **6**, 21800 (2016).

757   29.   J. M. Przyborski *et al.*, Trafficking of STEVOR to the Maurer's clefts in Plasmodium
758   falciparum-infected erythrocytes. *EMBO J* **24**, 2306-2317 (2005).

759   30.   G. G. van Dooren *et al.*, Development of the endoplasmic reticulum, mitochondrion and
760   apicoplast during the asexual life cycle of Plasmodium falciparum. *Mol Microbiol* **57**, 405-
761   419 (2005).

31. C. J. Tonkin, N. S. Struck, K. A. Mullin, L. M. Stimmler, G. I. McFadden, Evidence for
Golgi-independent transport from the early secretory pathway to the plastid in malaria
parasites. *Mol Microbiol* **61**, 614-630 (2006).

32. M. C. Lee, P. A. Moura, E. A. Miller, D. A. Fidock, Plasmodium falciparum Sec24 marks
transitional ER that exports a model cargo via a diacidic motif. *Mol Microbiol* **68**, 1535-
1546 (2008).

33. S. R. Heiny, S. Pautz, M. Recker, J. M. Przyborski, Protein Traffic to the Plasmodium
falciparum apicoplast: evidence for a sorting branch point at the Golgi. *Traffic* **15**, 1290-
1304 (2014).

34. J. M. Matz *et al.*, The Plasmodium berghei translocon of exported proteins reveals
spatiotemporal dynamics of tubular extensions. *Sci Rep* **5**, 12532 (2015).

35. M. J. Boucher *et al.*, Integrative proteomics and bioinformatic prediction enable a high-
confidence apicoplast proteome in malaria parasites. *PLoS Biol* **16**, e2005895 (2018).

36. J. Birnbaum *et al.*, A Kelch13-defined endocytosis pathway mediates artemisinin
resistance in malaria parasites. *Science* **367**, 51-59 (2020).

37. R. F. Waller, M. B. Reed, A. F. Cowman, G. I. McFadden, Protein trafficking to the
plastid of Plasmodium falciparum is via the secretory pathway. *EMBO J* **19**, 1794-1802
(2000).

38. A. Adisa *et al.*, The signal sequence of exported protein-1 directs the green fluorescent
protein to the parasitophorous vacuole of transfected malaria parasites. *J Biol Chem*
278, 6532-6542 (2003).

39. T. Nessel *et al.*, EXP1 is required for organisation of EXP2 in the intraerythrocytic
malaria parasite vacuole. *Cell Microbiol* **22**, e13168 (2020).

40. S. M. Ganesan, A. Falla, S. J. Goldfless, A. S. Nasamu, J. C. Niles, Synthetic RNA-
protein modules integrated with native translation mechanisms to control gene
expression in malaria parasites. *Nat Commun* **7**, 10727 (2016).

41. K. Rajaram, H. B. Liu, S. T. Prigge, Redesigned TetR-Aptamer System To Control Gene
Expression in Plasmodium falciparum. *mSphere* **5** (2020).

42. S. H. Adjalley *et al.*, Quantitative assessment of Plasmodium falciparum sexual
development reveals potent transmission-blocking activity by methylene blue. *Proc Natl
Acad Sci U S A* **108**, E1214-1223 (2011).

43. M. El Bakkouri *et al.*, The Clp chaperones and proteases of the human malaria parasite
Plasmodium falciparum. *J Mol Biol* **404**, 456-477 (2010).

44. E. Yeh, J. L. DeRisi, Chemical rescue of malaria parasites lacking an apicoplast defines
organelle function in blood-stage Plasmodium falciparum. *PLoS Biol* **9**, e1001138
(2011).

45.   G. G. van Dooren, V. Su, M. C. D'Ombrain, G. I. McFadden, Processing of an apicoplast
leader sequence in Plasmodium falciparum and the identification of a putative leader
cleavage enzyme. *J Biol Chem* **277**, 23612-23619 (2002).

46.   M. Zhang *et al.*, Uncovering the essential genes of the human malaria parasite
Plasmodium falciparum by saturation mutagenesis. *Science* **360** (2018).

47.   E. Bushell *et al.*, Functional Profiling of a Plasmodium Genome Reveals an Abundance
of Essential Genes. *Cell* **170**, 260-272 e268 (2017).

48.   M. E. Fichera, D. S. Roos, A plastid organelle as a drug target in apicomplexan
parasites. *Nature* **390**, 407-409 (1997).

49.   E. L. Dahl, P. J. Rosenthal, Multiple antibiotics exert delayed effects against the
Plasmodium falciparum apicoplast. *Antimicrob Agents Chemother* **51**, 3485-3490 (2007).

50.   N. I. Proellocks *et al.*, Characterisation of PfRON6, a Plasmodium falciparum rhoptry
neck protein with a novel cysteine-rich domain. *Int J Parasitol* **39**, 683-692 (2009).

51.   S. A. Desai, D. J. Krogstad, E. W. McCleskey, A nutrient-permeable channel on the
intraerythrocytic malaria parasite. *Nature* **362**, 643-646 (1993).

52.   S. A. Desai, R. L. Rosenberg, Pore size of the malaria parasite's nutrient channel. *Proc
Natl Acad Sci U S A* **94**, 2045-2049 (1997).

53.   M. Garten *et al.*, EXP2 is a nutrient-permeable channel in the vacuolar membrane of
Plasmodium and is essential for protein export via PTEX. *Nat Microbiol* **3**, 1090-1098
(2018).

54.   D. A. Gold *et al.*, The Toxoplasma Dense Granule Proteins GRA17 and GRA23 Mediate
the Movement of Small Molecules between the Host and the Parasitophorous Vacuole.
Cell Host Microbe **17**, 642-652 (2015).

55.   P. R. Sanders *et al.*, The N-terminus of EXP2 forms the membrane-associated pore of
the protein exporting translocon PTEX in Plasmodium falciparum. *J Biochem* **165**, 239-
248 (2019).

56.   K. Hakamada, H. Watanabe, R. Kawano, K. Noguchi, M. Yohda, Expression and
characterization of the Plasmodium translocon of the exported proteins component
EXP2. *Biochem Biophys Res Commun* **482**, 700-705 (2017).

57.   M. Dal Peraro, F. G. van der Goot, Pore-forming toxins: ancient, but never really out of
fashion. *Nat Rev Microbiol* **14**, 77-92 (2016).

58.   S. C. Charnaud, R. Kumarasingha, H. E. Bullen, B. S. Crabb, P. R. Gilson, Knockdown
of the translocon protein EXP2 in Plasmodium falciparum reduces growth and protein
export. *PLoS One* **13**, e0204785 (2018).

59.   P. Mesen-Ramirez *et al.*, EXP1 is critical for nutrient uptake across the parasitophorous
vacuole membrane of malaria parasites. *PLoS Biol* **17**, e3000473 (2019).

60.   T. F. de Koning-Ward *et al.*, A newly discovered protein export machine in malaria
835   parasites. *Nature* **459**, 945-949 (2009).

836   61.   M. Gabriela *et al.*, A revised mechanism for how Plasmodium falciparum recruits and
837   exports proteins into its erythrocytic host cell. *PLoS Pathog* **18**, e1009977 (2022).

838   62.   S. C. Charnaud *et al.*, Spatial organization of protein export in malaria parasite blood
839   stages. *Traffic* **19**, 605-623 (2018).

840   63.   A. Ingmundson, C. Nahar, V. Brinkmann, M. J. Lehmann, K. Matuschewski, The
841   exported Plasmodium berghei protein IBIS1 delineates membranous structures in
842   infected red blood cells. *Mol Microbiol* **83**, 1229-1243 (2012).

843   64.   S. Nair *et al.*, Nutrient Limitation Magnifies Fitness Costs of Antimalarial Drug
844   Resistance Mutations. *Antimicrob Agents Chemother* **66**, e0152921 (2022).

845   65.   A. Gupta *et al.*, Complex nutrient channel phenotypes despite Mendelian inheritance in a
846   Plasmodium falciparum genetic cross. *PLoS Pathog* **16**, e1008363 (2020).

847   66.   J. M. Matz *et al.*, In Vivo Function of PTEX88 in Malaria Parasite Sequestration and
848   Virulence. *Eukaryot Cell* **14**, 528-534 (2015).

849   67.   S. A. Chisholm *et al.*, Contrasting Inducible Knockdown of the Auxiliary PTEX
850   Component PTEX88 in P. falciparum and P. berghei Unmasks a Role in Parasite
851   Virulence. *PLoS One* **11**, e0149296 (2016).

852   68.   H. M. Kudyba, D. W. Cobb, J. Vega-Rodriguez, V. Muralidharan, Some conditions apply:
853   Systems for studying Plasmodium falciparum protein function. *PLoS Pathog* **17**,
854   e1009442 (2021).

855   69.   J. Birnbaum *et al.*, A genetic system to study Plasmodium falciparum protein function.
856   *Nat Methods* **14**, 450-456 (2017).

857   70.   K. R. Hughes, A. P. Waters, Rapid inducible protein displacement in. *Wellcome Open*
858   *Res* **2**, 18 (2017).

859   71.   A. D. Roberts, S. C. Nair, A. J. Guerra, S. T. Prigge, Development of a conditional
860   localization approach to control apicoplast protein trafficking in malaria parasites. *Traffic*
861   **20**, 571-582 (2019).

862   72.   A. Praznik *et al.*, Regulation of protein secretion through chemical regulation of
863   endoplasmic reticulum retention signal cleavage. *Nat Commun* **13**, 1323 (2022).

864   73.   A. Prasad, P. Mastud, S. Patankar, Dually localised proteins found in both the apicoplast
865   and mitochondrion utilize the Golgi-dependent pathway for apicoplast targeting in
866   Toxoplasma gondii. *Biol Cell* **113**, 58-78 (2021).

867   74.   A. Florentin *et al.*, PfClpC Is an Essential Clp Chaperone Required for Plastid Integrity
868   and Clp Protease Stability in Plasmodium falciparum. *Cell Rep* **21**, 1746-1756 (2017).

869   75.   A. Florentin, D. R. Stephens, C. F. Brooks, R. P. Baptista, V. Muralidharan, Plastid
870   biogenesis in malaria parasites requires the interactions and catalytic activity of the Clp
871   proteolytic system. *Proc Natl Acad Sci U S A* **117**, 13719-13729 (2020).

872   76.   A. Mogk, B. Bukau, H. H. Kampinga, Cellular Handling of Protein Aggregates by
873   Disaggregation Machines. *Mol Cell* **69**, 214-226 (2018).

874   77.   I. Bouchnak, K. J. van Wijk, Structure, function, and substrates of Clp AAA+ protease
875   systems in cyanobacteria, plastids, and apicoplasts: A comparative analysis. *J Biol*
876   *Chem* **296**, 100338 (2021).

877   78.   K. Pakos-Zebrucka *et al.*, The integrated stress response. *EMBO Rep* **17**, 1374-1395
878   (2016).

879   79.   M. Costa-Mattioli, P. Walter, The integrated stress response: From mechanism to
880   disease. *Science* **368** (2020).

881   80.   S. Chaubey, M. Grover, U. Tatu, Endoplasmic reticulum stress triggers
882   gametocytogenesis in the malaria parasite. *J Biol Chem* **289**, 16662-16674 (2014).

883   81.   M. B. Harbut *et al.*, Targeting the ERAD pathway via inhibition of signal peptide
884   peptidase for antiparasitic therapeutic design. *Proc Natl Acad Sci U S A* **109**, 21486-
885   21491 (2012).

886   82.   M. Zhang *et al.*, PK4, a eukaryotic initiation factor 2alpha(eIF2alpha) kinase, is essential
887   for the development of the erythrocytic cycle of Plasmodium. *Proc Natl Acad Sci U S A*
888   **109**, 3956-3961 (2012).

889   83.   M. Zhang *et al.*, Inhibiting the Plasmodium eIF2alpha Kinase PK4 Prevents Artemisinin-
890   Induced Latency. *Cell Host Microbe* **22**, 766-776 e764 (2017).

891   84.   Y. S. Levray *et al.*, Formation of ER-lumenal intermediates during export of Plasmodium
892   proteins containing transmembrane-like hydrophobic sequences. *PLoS Pathog* **19**,
893   e1011281 (2023).

894   85.   E. C. Gaynor, S. te Heesen, T. R. Graham, M. Aebi, S. D. Emr, Signal-mediated retrieval
895   of a membrane protein from the Golgi to the ER in yeast. *J Cell Biol* **127**, 653-665
896   (1994).

897   86.   M. Stornaiuolo *et al.*, KDEL and KKXX retrieval signals appended to the same reporter
898   protein determine different trafficking between endoplasmic reticulum, intermediate
899   compartment, and Golgi complex. *Mol Biol Cell* **14**, 889-902 (2003).

900   87.   B. L. Tang, S. H. Wong, S. H. Low, W. J. Hong, Retention of a Type-Ii Surface-
901   Membrane Protein in the Endoplasmic-Reticulum by the Lys-Asp-Glu-Leu Sequence.
902   *Journal of Biological Chemistry* **267**, 7072-7076 (1992).

903   88.   B. E. Sleebs *et al.*, Inhibition of Plasmepsin V activity demonstrates its essential role in
904   protein export, PfEMP1 display, and survival of malaria parasites. *PLoS Biol* **12**,
905   e1001897 (2014).

89. A. J. Polino, A. S. Nasamu, J. C. Niles, D. E. Goldberg, Assessment of Biological Role
and Insight into Druggability of the Plasmodium falciparum Protease Plasmepsin V. *ACS*
Infect Dis **6**, 738-746 (2020).

90. S. H. Adjalley, M. C. S. Lee, D. A. Fidock, "A Method for Rapid Genetic Integration into
Plasmodium falciparum Utilizing Mycobacteriophage Bxb1 Integrase". (Humana Press,
2010), 10.1007/978-1-60761-652-8_6, pp. 87-100.

91. S. Glushakova *et al.*, Exploitation of a newly-identified entry pathway into the malaria
parasite-infected erythrocyte to inhibit parasite egress. *Sci Rep* **7**, 12250 (2017).

92. A. I. Ahiya, S. Bhatnagar, J. M. Morrisey, J. R. Beck, A. B. Vaidya, Dramatic
Consequences of Reducing Erythrocyte Membrane Cholesterol on Plasmodium
falciparum. *Microbiol Spectr* **10**, e0015822 (2022).

93. T. W. Kooij, M. M. Rauch, K. Matuschewski, Expansion of experimental genetics
approaches for Plasmodium berghei with versatile transfection vectors. *Mol Biochem*
Parasitol **185**, 19-26 (2012).

94. C. J. Janse, J. Ramesar, A. P. Waters, High-efficiency transfection and drug selection of
genetically transformed blood stages of the rodent malaria parasite Plasmodium berghei.
Nat Protoc **1**, 346-356 (2006).

95. R. S. Kent *et al.*, Inducible developmental reprogramming redefines commitment to
sexual development in the malaria parasite Plasmodium berghei. *Nat Microbiol* **3**, 1206-
1213 (2018).

96. J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat*
Methods **9**, 676-682 (2012).

97. E. P. Rock *et al.*, Comparative analysis of the Plasmodium falciparum histidine-rich
proteins HRP-I, HRP-II and HRP-III in malaria parasites of diverse origin. *Parasitology*
95 (Pt 2), 209-227 (1987).

98. T. Blisnick *et al.*, Pfsbp1, a Maurer's cleft Plasmodium falciparum protein, is associated
with the erythrocyte skeleton. *Mol Biochem Parasitol* **111**, 107-121 (2000).

99. M. Ponpuak *et al.*, A role for falcilysin in transit peptide degradation in the Plasmodium
falciparum apicoplast. *Mol Microbiol* **63**, 314-334 (2007).

100. R. Hall *et al.*, Antigens of the Erythrocytic Stages of the Human Malaria Parasite
Plasmodium-Falciparum Detected by Monoclonal-Antibodies. *Molecular and*
Biochemical Parasitology **7**, 247-265 (1983).

101. K. Matthews *et al.*, The Plasmodium translocon of exported proteins (PTEX) component
thioredoxin-2 is important for maintaining normal blood-stage growth. *Mol Microbiol* **89**,
1167-1186 (2013).

 Figures

 Figure 1. Conditional ER retrieval of a secreted reporter does not impact parasite fitness.

A) Schematic of reporter expression cassette installed at the *attB* site of chromosome 6. An mNeonGreen reporter is fused to the signal peptide of EXP2 and expressed under control of the *exp2* promoter. A 3xFLAG tag and stop codon are flanked by loxP-containing introns (triangles) followed by a 3xHA-KDEL, stop codon, and 10X aptamer array (10X-Apt). Aptamer interaction with TetR-DOZI (expressed from a cassette in the plasmid not shown) prevents translation of the reporter, which is relieved in the presence of anhydrotetracycline (aTc) to enable expression. Activation of DiCre by rapamycin treatment excises the FLAG tag, bringing into frame the HA-KDEL tag. B) Time course of excision following rapamycin treatment detected by PCR using primers P7/32. C) Western blot of mNG reporter 24hrs post treatment with DMSO or rapamycin. Molecular weights after signal peptide cleavage are predicted to be 30.8 kDa for mNG-3xFLAG and 30.9 kDa for mNG-3xHA-KDEL. D) Live microscopy of DMSO or rapamycin-treated parasites 24hrs post-treatment. E) Quantification of percent internal mNG fluorescence 24hrs and 168hrs post-treatment with DMSO or rapamycin. Data are pooled from 2 independent experiments and

958   bar indicates mean (****, P<0.0001; unpaired t test). F) Representative growth of asynchronous

959   parasites (n=2 biological replicates) treated with DMSO or rapamycin. Data are presented as

960   means ± standard deviation from one biological replicate (n = 3 technical replicates). All cultures

961   were maintained in media supplemented with 500 nM aTc. Scale bar, 5µm.

Figure 2. Lethal ER retention of the apicoplast chaperone ClpB1 is rescued by IPP. A) Schematic showing strategy for appending kER to the C-terminus of *clpB1*. B) IFA of PFA fixed parasites stained with mouse anti-Flag and rabbit anti-ACP1 antibodies. C) PCR showing excision in ClpB1-kER parasites 24hrs after rapamycin treatment using primers P7/8. D) Western blot 24hrs post treatment with DMSO or rapamycin. Molecular weights after transit peptide cleavage are predicted to be 135 kDa for ClpB1-mNG-3xFLAG and 135.1 kDa for ClpB1-mNG-3xHA-KDEL. (+) represents the mature form of ClpB1 while the (*) represents ClpB1 with the transit peptide still attached. E) Live microscopy of DMSO or rapamycin-treated parasites 48hrs post-treatment. F) Representative growth of asynchronous parasites (n=3 biological replicates) treated with DMSO or rapamycin, with or without supplementation with 200 µM IPP. Data are presented as

972   means ± standard deviation from one biological replicate (n = 3 technical replicates). G) IFA of

973   PFA fixed parasites stained with mouse anti-Flag or anti-HA and rabbit anti-ACP1 antibodies after

974   3 growth cycles where rapamycin treated parasites were supplemented with 200 µM IPP. Scale

975   bars, 5µm.

Figure 3. ER retention of EXP2 causes a lethal fitness defect independent of loss of PVM

transport functions. A) Schematic showing strategy for appending kER to the endogenous C-

terminus of EXP2 without mNG. B) PCR showing excision in EXP2-kER 24hrs after treatment

with rapamycin using primers P9/10. C) Western blot of EXP2-kER parasites 24hrs post treatment

with DMSO or rapamycin. Molecular weights after signal peptide cleavage are predicted to be

34.6 kDa for EXP2-3xFLAG and 34.7 kDa for EXP2-3xHA-KDEL. D) IFA of EXP2-kER 24hrs post-

treatment with DMSO or rapamycin. E) Growth of asynchronous parasites (n=2 biological

replicates) treated with DMSO or rapamycin. Data are presented as means ± standard deviation

from one biological replicate (n = 3 technical replicates). F) Live microscopy of parasites

expressing a full-length second copy of EXP2 with an mNG-kER fusion from the *attB* site under

the control of the endogenous *exp2* promoter (EXP2-FL-kER[attB]). Parasites were viewed 24hrs

after treatment with DMSO or rapamycin. G) Representative growth of asynchronous EXP2-FL-

kER[attB] parasites (n=2 biological replicates) treated with DMSO or rapamycin. Data are presented

as means ± standard deviation from one biological replicate (n = 3 technical replicates). H) Live

microscopy of parasites expressing a second copy of EXP2 with mNG-kER fusion and lacking

the amphipathic helix from the *attB* site (EXP2ΔTM-kER[attB]). Parasites were viewed 24hrs after

treatment with DMSO or rapamycin. Two representative examples of unexcised parasites are

presented, showing localization of the ΔTM version of EXP2 to the PV and host cell in some

infected RBCs (arrow). Digestive vacuole fluorescence was also observed as is typical for PV

proteins. I) Representative growth of asynchronous EXP2ΔTM-kER[attB] parasites (n=2 biological

replicates) treated with DMSO or rapamycin. Data are presented as means ± standard deviation

from one biological replicate (n = 3 technical replicates). Second copy EXP2-kER lines were

maintained in media supplemented with 500 nM aTc. Scale bars, 5µm.

Figure 4. ER retention of PTEX150 but not HSP101 causes a lethal block in protein export in *P. falciparum*. A) PCR showing excision in PTEX150-kER parasites 24hrs after rapamycin treatment using primers P7/8. B) Western blot of PTEX150-kER 24hrs post-treatment with DMSO

or rapamycin. Molecular weights after signal peptide cleavage are predicted to be 141 kDa for

PTEX150-mNG-3xFLAG and 141.1 kDa for PTEX150-mNG-3xHA-KDEL. Note that PTEX150 is

observed to migrate at higher molecular weight than predicted (60). C) Live microscopy of

PTEX150-kER 24hrs post-treatment with DMSO or rapamycin. D) Representative growth of

asynchronous PTEX150-kER parasites (n=3 biological replicates) treated with DMSO or

rapamycin. Data are presented as means ± standard deviation from one biological replicate (n = 3

technical replicates). E) IFA of synchronized, ring-stage PTEX150-kER parasites 48hrs post-

treatment with DMSO or rapamycin and probed with anti-HRP2 and anti-FLAG or anti-HA

antibodies. F) PCR showing excision in HSP101-kER parasites 24hrs after rapamycin treatment

using primers P7/8. G) Western blot of HSP101-kER parasites 14 days post-treatment with DMSO

or rapamycin. Molecular weights after signal peptide cleavage are predicted to be 130.5 kDa for

HSP101-mNG-3xFLAG and 130.6 kDa for HSP101-mNG-3xHA-KDEL. H) Live microscopy of

HSP101-kER 24hrs and 168hrs post-treatment with DMSO or rapamycin. I) Representative

growth of asynchronous HSP101-kER parasites (n=3 biological replicates) treated with DMSO or

rapamycin. Data are presented as means ± standard deviation from one biological replicate (n = 3

technical replicates). J) IFA of synchronized, ring-stage HSP101-kER parasites 48hrs post-

treatment with DMSO or rapamycin and probed with anti-HRP2 and anti-FLAG or anti-HA

antibodies. Scale bar, 5μm.

Figure 5: ER retention of HSP101 causes a lethal block in protein export in *P. berghei*. A) PCR showing excision in PbPTEX150-kER and PbHSP101-kER *P. berghei* parasites treated with rapamycin and cultured 18hrs *ex vivo* using primers P7/P60. B) Western blot of PbPTEX150-kER and PbHSP101-kER parasites 24hrs post-treatment with DMSO or rapamycin. Molecular weights after signal peptide cleavage are predicted to be 130.9 kDa for PbHSP101-mNG-3xFLAG, 131 kDa for PbHSP101-mNG-3xHA-KDEL, 130.5 kDa for PbPTEX150-mNG-3xFLAG and 130.6 kDa for PbPTEX150-mNG-3xHA-KDEL. Note that PbPTEX150 is observed to migrate at higher molecular weight than predicted (101). C,F) Live microscopy of PbHSP101-kER or PbPTEX150-kER parasites 18hrs post-treatment with DMSO or rapamycin. These transgenic lines also contain a downstream cassette for expression of mRuby bearing a signal peptide for secretion into the

PV (see Figure S6A). D,G) Quantification of percent internal mNG fluorescence 24hrs post-

treatment with DMSO or rapamycin. Data are pooled from 2 independent experiments and bar

indicates mean (****, $P<0.0001$; unpaired t test). E,H) Parasitemia of PbHSP101-kER or

PbPTEX150-kER infected mice inoculated by tail vein-injection of parasites treated *ex vivo* with

DMSO or rapamycin. Data are presented as means ± standard deviation of three independent

experiments (one mouse infected per treatment in each experiment). No parasites were observed

in mice injected with rapamycin treated parasites except that one mouse injected with rapamycin-

treated PbHSP101-kER parasites became patent on day 10 (#). However, PCR showed these

parasites had not undergone excision (see Figure S6B). I,J) Live microscopy of PbHSP101-kER

or PbPTEX150-kER parasites containing a downstream cassette for expression of an IBIS1-

mRuby fusion that is exported into the RBC (See Figure S6A). Parasites were imaged before or

24hrs after treatment with rapamycin. Dashed lines indicate RBC boundaries traced from DIC

images. Images are representative of n=2 biological replicates. Scale bars, 5µm.

Figure 6. Depletion of HSP101 PV levels hypersensitizes parasites to post-translational

destabilization of the HSP101-PTEX complex but not HSP101 translational knockdown. A)

Schematic showing strategy for appending kER and the TetR-DOZI aptamers system to the

endogenous C-terminus of HSP101. B) PCR showing excision in HSP101-kER[TetR-DOZI] parasites

24hrs after rapamycin treatment using primers P7/8. C) Live microscopy of HSP101-kER[TetR-DOZI]

parasites 168hrs post-treatment with DMSO or rapamycin. D) IFA detecting the soluble PEXEL

protein HRP2 or the transmembrane PNEP SBP1 in synchronized, ring-stage HSP101-kER[TetR-]

[DOZI] parasites grown for 48hrs +/- aTc and treated with DMSO or rapamycin 1 cycle prior. Dashed

lines indicate RBC boundaries traced from DIC images. E) Quantification of HSP101 levels by

flow cytometry measurement of mNeonGreen fluorescence in HSP101-kER[TetR-DOZI] parasites

grown for 48 hrs at indicated aTc concentrations and treated with DMSO or rapamycin 2 cycles

prior. MFI data were normalized to DMSO-treated parasites grown in 500 nM aTc (100%) and

presented as mean ± SEM (n = 5 biological replicates). F) Representative growth curves (n = 4

biological replicates) of HSP101-kER[TetR-DOZI] parasites grown at indicated aTc concentrations and

treated with DMSO or rapamycin 2 cycles prior. Data are presented as means ± standard

deviation from one biological replicate (n = 3 technical replicates). G) Representative EC_{50} assays

(n=4 biological replicates) with two-fold dilutions of aTc starting at 500nM. Parasites were treated

with DMSO or rapamycin 72 hours prior to adjusting aTc concentrations. Data are presented as

means ± standard deviation from one biological replicate (n = 3 technical replicates). EC_{50} graph

inset presented as mean ± SEM (n = 4 biological replicates): DMSO=10.58±1.16 nM and

rapamycin=9.11±0.29 nM. H) Schematic showing strategy for appending kER and the DDD

system to the endogenous C-terminus of HSP101. I) Representative EC_{50} assays (n=4 biological

replicates) with two-fold dilutions of TMP starting at 250nM. Parasites were treated with DMSO

or rapamycin 72 hours prior to adjusting TMP concentrations. Data are presented as means ±

standard deviation from one biological replicate (n = 3 technical replicates). EC_{50} graph inset

presented as mean ± SEM (n = 4 biological replicates): DMSO=3.39±0.48 nM and rapamycin

5.48±0.73 nM. Scale bars, 5μm.