Plant defensin MtDef4-derived antifungal peptide with multiple modes of action and potential as

a bioinspired fungicide

Meenakshi Tetorya[1*], Hui Li[1*], Arnaud Thierry Djami-Tchatchou[1], Garry W. Buchko[3,4], Kirk J.

Czymmek[1,2], Dilip M. Shah[1]

[1]Donald Danforth Plant Science Center, St Louis, MO. 63132, USA.

[2]Advanced Bioimaging Laboratory, Donald Danforth Plant Science Center, St Louis, MO

63132, USA

[3]Earth and Biological Sciences Directorate, Pacific Northwest National Laboratory, Richland,

WA 99354, USA

[4]School of Molecular Biosciences, Washington State University, Pullman, WA 99164, USA

Corresponding Author*:

Email: dshah@danforthcenter.org)

*These authors contributed equally to this work.

Keywords:

Defensin-derived peptides, antifungal activity, modes of action, *Botrytis cinerea*, gray mold

disease, bioinspired fungicides.

Abstract

Chemical fungicides have been instrumental in protecting crops from fungal diseases. However, mounting fungal resistance to many of the single-site chemical fungicides calls for the development of new antifungal agents with novel modes of action (MoA). The sequence-divergent cysteine-rich antifungal defensins with multi-site MoA are promising starting templates for design of novel peptide-based fungicides. Here, we experimentally tested such a set of 17-amino acid peptides containing the γ-core motif of the antifungal plant defensin MtDef4. These designed peptides exhibited antifungal properties different from those of MtDef4. Focused analysis of a lead peptide, GMA4CG_V6, showed it was a random coil in solution with little or no secondary structure elements. Additionally, it exhibited potent cation-tolerant antifungal activity against the plant fungal pathogen *Botrytis cinerea*, causal agent of gray mold disease in fruits and vegetables. Its multi-site MoA involved localization predominantly to the plasma membrane, permeabilization of the plasma membrane, rapid internalization into the vacuole and cytoplasm, and affinity for bioactive phosphoinositides phosphatidylinositol 3-phosphate (PI3P), PI4P, and PI5P. The sequence motif RRRW was identified as a major determinant of the antifungal activity of this peptide. While topical spray-application of GMA4CG_V6 on *Nicotiana benthamiana* and tomato plants provided preventative and curative suppression of gray mold disease symptoms, the peptide was not internalized into plant cells. Our findings open the possibility that truncated and modified defensin-derived peptides containing the γ-core sequence could serve as promising candidates for further development as bioinspired fungicides.

INTRODUCTION

Crop plants are under constant threat from fungal and oomycete pathogens that cause significant pre- and post-harvest economic losses worldwide and challenge food security globally (Fones *et al.*, 2020). This threat is increasing in severity and scale (Fisher *et al.*, 2012, Savary *et al.*, 2019). It is estimated that fungal and oomycete pathogens cause up to 10-23% yield losses in calorie crops (Savary et al., 2019) and postharvest losses of 10% (Fisher *et al.*, 2018). Modern intensive agriculture depends heavily on chemical fungicides for management of fungal and oomycete diseases in commodity crops as well as high value vegetable and fruit crops (Oliver & Hewitt, 2014). Most chemical fungicides in use today have a single biochemical target and thus are prone to resistance development by diverse and rapidly evolving fungal and oomycete strains (Hawkins and Fraiije, 2018). Facing fungal resistance and unwanted side effects, some fungicides are losing their effectiveness in modern intensive monoculture cropping systems. The structurally diverse plant antifungal peptides with potent antifungal activity could potentially serve as structural templates for the design of a new generation of safe and sustainable fungicides (Lobo & Boto, 2022, van der Weerden *et al.*, 2013).

Plants use a variety of different mechanisms to resist fungal and oomycete infections (van Esse *et al.*, 2020). Anti-microbial peptides (AMPs) are ubiquitous throughout the plant kingdom and play important roles in plant defense against fungal and oomycete pathogens. Their potent anti-fungal and -oomycete activity are vital components of the plant innate immune system (Goyal & Mattoo, 2014, van der Weerden et al., 2013). Defensins, in particular, are one of the largest families of AMPs expressed in plants. Because of their potent antifungal activity, diverse amino acid sequences, and multi-target MoA, plant defensins have emerged as potential antifungal agents for agriculture (Cools *et al.*, 2017, Parisi *et al.*, 2019, Leannec-Rialland *et al.*, 2022). Plant

defensins are 45-54 amino acids long, cysteine-rich, cationic peptides characterized by the

presence of a cysteine-stabilized αβ motif. This motif consists of one α-helix and three antiparallel

β-strands linked by three disulfide bonds. The fourth disulfide bond connects near the N- and C-

termini and essentially cyclizes the peptide (Kovaleva *et al.*, 2020). The cationic plant defensins

exhibit potent broad-spectrum antifungal activity against fungal pathogens *in vitro* and *in planta*

(Kaur *et al.*, 2011, Kovaleva et al., 2020, Lacerda *et al.*, 2014, Leannec-Rialland et al., 2022). We

set out to investigate if truncated plant defensins containing the active carboxy-terminal γ-core

motif ($GXCX_{3-9}C$) exhibited potent fungicidal activity and were effective as spray-on fungicides

for management of fungal diseases. We further investigated if the smaller defensin-derived

peptides exhibited MoA different from those of the full-length parental peptide.

MtDef4 is an evolutionarily conserved 47-amino acid defensin from *Medicago truncatula*

that exhibits broad-spectrum antifungal activity against fungal pathogens at low micromolar

concentrations. MtDef4 MoA against the plant pathogen *Fusarium graminearum* involves plasma

membrane permeabilization, binding to bioactive phosphatidic acid (PA), cell entry, and presumed

interaction with intracellular targets (El-Mounadi *et al.*, 2016, Sagaram *et al.*, 2013, Sagaram *et*

al., 2011). Like many other plant defensins, this peptide contains a well-characterized γ-core motif

required for its antifungal activity (Sagaram et al., 2013, Sagaram et al., 2011). A chemically

synthesized C-terminal 16-amino acid sequence, designated GMA4C, containing this motif and a

net charge of +6 inhibited the growth of the fungal pathogen *Fusarium graminearum in vitro*

although its potency was four-fold lower than that of MtDef4 (Sagaram et al., 2013). This peptide

also exhibited antifungal activity against crown rot fungal pathogens of alfalfa and, in addition,

inhibited the growth of plant and human bacterial pathogens *in vitro* (Sathoff *et al.*, 2019).

Herein, we explored the potential of the synthetic, MtDef4-derived, structurally less complex, 17-amino acid GMA4CG variants for development as fungicides. We characterized the antifungal activity of several GMA4CG variants generated via amino acid substitutions, C-terminal amidation, and disulfide cyclization that exhibited potent antifungal activity *in vitro* against an aggressive necrotrophic fungal pathogen *Botrytis cinerea*, the causal agent of a gray mold disease in vegetables, fruits, and flowers. In sharp contrast to MtDef4, an interesting characteristic of the GMA4CG variants was that they retained potent antifungal activity in media with elevated salt concentrations. Based on its superior *in vitro* and semi-*in planta* antifungal activity against *B. cinerea*, we focused on GMA4CG_V6 as the lead peptide for *in planta* antifungal activity and MoA studies. GMA4CG_V6 exhibited multi-faceted MoA and rapidly killed fungal cells. When sprayed on *Nicotiana benthamiana* and tomato plants, GMA4CG_V6 conferred strong preventative and curative antifungal activity against *B. cinerea* under the tested conditions.

RESULTS

Design of short MtDef4-derived synthetic peptide variants

We based our work on the premise that a small MtDef4-derived peptide spanning the active γ-core motif could serve as a promising template for the design of peptide-based fungicides. We further hypothesized that a small antifungal peptide would be more cell permeant and that with targeted amino acid substitutions of the γ-core motif, we could further enhance its antifungal potency. To test this hypothesis, we generated a truncated set of linear and cyclic variants of the 17-amino acid peptide spanning this motif (Gly30-Cys47s designated GMA4CG_WT and tested them for antifungal activity against *B. cinerea* (Figure 1A).

The substitution of each amino acid with all possible alternative amino acids in GMA4CG_WT was not practical. Thus, our strategy was to substitute cationic or hydrophobic residues in specific positions with the expectation that they would have significant influence on the antifungal activity of this peptide. MtDef4 is an evolutionarily conserved defensin in the plant kingdom and sequences of several homologs from various plants have been reported (Sagaram et al., 2011). GMA4CG_V1 is a natural variant derived from the MtDef4 homolog (KHN09675.1) of *Glycine soja*. This natural variant has three amino acid differences (R5K, T15R and H16I) relative to the sequence of GMA4CG_WT (Figure 1A). GMA4CG_V2 was generated to determine if the substitution of one cationic amino acid with a different cationic amino acid had a notable impact on its antifungal activity. Thus, this variant carried a Lys5 to Arg substitution. We also designed peptide variants (GMA4CG_V3 to GMA4CG_V6) to test the influence of increased hydrophobicity on their antifungal activity (Figure 1A). Thus, each of these variants had 29.4% hydrophobic amino acids as compared with 11.8% in GMA4CG_WT. GMA4CG_WT has four cysteine residues (Cys4, Cys11, Cys13 and Cys17) in its sequence, suggesting a potential to generate a pseudo-cyclic structure via disulfide bond formation between Cys4 and Cys17. Cyclization has been reported previously to increase the antimicrobial activity and selectivity of AMPs (Dathe *et al.*, 2004, Tucker *et al.*, 2018). Thus, we had cyclic GMA4CG_V3 and GMA4CG_V6 synthesized with a Cys4-Cys17 disulfide bond and replaced Cys11 and Cys13 with either an aliphatic Val (GMA4CG_V3) or an aromatic Trp (GMA4CG_V6) residue. Trp is more effective in interacting with the interfacial regions of lipid bilayers and it promotes antimicrobial activity of peptides (Deslouches *et al.*, 2005, Chan *et al.*, 2006, Killian & von Heijne, 2000). All peptides were amidated at the C-terminus, a process known to increase the stability of peptides against proteolytic degradation, and therefore, also its biological activity (Lu *et al.*, 2020).

GMA4CG_V6 variants have potent antifungal activity

We assessed the antifungal activity of GMA4CG_WT and its six variants in Synthetic Fungal

Medium (SFM) against *B. cinerea in vitro.* GMA4CG_WT inhibited the growth of *B.cinerea* with

an MIC value of 3 µM, four-fold greater than that of MtDef4 (Figure 1B). Among all variants,

only GMA4CG_V6 displayed two-fold greater antifungal activity than GMA4CG_WT with an

MIC value of 1.5 µM (Figure 1B).

To determine if GMA4CG_V6 had fungicidal and/or fungistatic activity, *B. cinerea*

conidia were treated with 1.5 µM peptide for 1, 6, and 24 h. Following each treatment, the peptide

was removed, and spore germination examined under the microscope. Approximately 85% of the

conidia failed to germinate after 1 h of peptide challenge, nearly 100% of the conidia failed to

germinate after 6 h, and 100% of the conidia failed to germinate after 24 h of treatment (Figure

S1A and B). To assess further the toxicity of GMA4CG_V6, we stained peptide-treated conidial

cells with propidium iodide, a cell death marker, and observed that nearly 80% of the conidia

stained red within 2 h of peptide treatment indicating fungicidal activity for this peptide (Figure

S1C and D).

The GMA4CG variants, but not MtDef4, retain antifungal activity in the presence of salts

Many naturally occurring AMPs, including plant defensins, are known to lose their antifungal

activity in media containing elevated salt concentrations (Kerenga *et al.*, 2019, Li *et al.*, 2019). We

determined the MIC values of GMA4CG_WT and its variants with the R/KGFRRR motif in SFM

containing 100 mM NaCl, 100 mM KCl, or 2 mM CaCl$_2$ and compared them to MtDef4. At the

greatest concentration tested (6 µM), MtDef4 had no activity against *B. cinerea*. However,

GMA4CG_WT and its variants retained their antifungal activity in salt supplemented SFM to

varying degrees with GMA4CG_V4 and GMA4CG_V6 retaining the most potent antifungal

activity (Figures 1B and S2A). GMA4CG_V6 at 3 μM completely inhibited spore germination in

the presence of NaCl and CaCl$_2$ salts (S2B Fig). GMA4CG_V6 was notably more effective than

GMA4CG_V3 in the presence of 100 mM KCl and 2 mM CaCl$_2$ (Figures 1B and S2A) indicating

that the Val to Trp substitution is important for salt-tolerant antifungal activity in these variants.

Collectively, these data revealed that the MtDef4-derived GMA4CG_WT peptide and its variants

with a net charge of +5 to +6 exhibited salt-tolerant antifungal activity. In contrast, the parental

MtDef4 exhibited salt-sensitive antifungal activity at the same concentrations.

GMA4CG_WT and GMA4CG_V6 are predominantly random coil structures

The three-dimensional structure of full length MtDef4 contains a γ-core motif composed of β-

strands β2 and β3 connected by a positively charged RGFRRR loop (Sagaram et al., 2013). Near-

UV CD spectra for GMA4CG_WT and GMA4CG_V6 are shown in Figure S3. Neither spectrum

shows features characteristic of a peptide with substantial α-helical (double minima at

approximately 222 and 208 nm and an extrapolated maximum around 195 nm) or β-strand (single

minimum at around 220 nm and an extrapolated maximum around 195 nm) secondary structure

(Greenfield, 2006, Holzwarth & Doty, 1965). The notable difference between the CD spectra of

the two peptides was an additional minor minimum at ~230 nm for GMA4CG_WT and a minor

maximum at ~ 230 nm for GMA4CG_V6. In combination with the major sub-220 nm minimum,

the former feature is characteristic of a random coil/unordered structure and the latter feature

characteristic of a polyproline II structure suggesting the multiple amino acid substitutions may

induce a transition of some random coil to polyproline II structure in GMA4CG_V6 (Lopes *et al.*,

2014). Our attempts to predict the structures for GMA4CG and GMA4CG_V6 using the artificial

intelligence-based protein structure prediction software AlphaFold (Jumper *et al.*, 2021) yielded

structures with low confidence. Similarly, *ab initio* structure prediction using the Rosetta protein

structure modeling package (Baek *et al.*, 2021) did not predict any consistent fold or high-

confidence secondary structure elements (data not shown). Hence, the *in silico* modeling data

corroborate that experimental CD data, suggesting the peptides have little long-range order and

are likely dynamic and disordered in solution.

The RRRW motif is a major determinant of the salt-tolerant antifungal activity of

GMA4CG_V6 with no disulfide bond

A series of linear alanine-substituted variants of our lead peptide GMA4CG_V6 (Figure S4A)

were tested to identify amino acids important for antifungal activity. The GMA4CG_V6_Ala3

variant carrying the substitution of RRRW with AAAA lost antifungal activity eight-fold, whereas

all other variants lost antifungal activity four-fold (Figures S4A and S4B). In SFM supplemented

with 100 mM NaCl, no further loss in the antifungal activity of the variants occurred except in

GMA4CG_V6_Ala3, which lost antifungal activity another two-fold (Figure S4C). These results

indicated that the RRRW motif is a major determinant of the antifungal activity of GMA4CG_V6

in media supplemented with salts.

The GMA4CG_V4 and GMA4CG_V6 variants potently reduce gray mold disease symptoms

semi-*in planta*

We compared the semi-*in planta* antifungal activity of GMA4CG_WT and its variants using

detached leaves of *N. benthamiana* as described (Li et al., 2019). GMA4CG_WT and variants

GMA4CG_V1 through GMA4CG_V5 were applied as drops of different concentrations on each

leaf with freshly prepared conidial inoculum immediately applied to each drop. Leaves were

assessed for attenuation of gray mold symptoms by each peptide 48 h post-inoculation (hpi)

relative to no peptide controls by measuring lesion sizes and photosynthetic efficiency (as

measured by Fv (variable fluorescence)/Fm (maximal fluorescence)). Increasing concentration of

each peptide significantly ($p \leq 0.0001$) reduced lesion sizes (Figures 2A and B). For all peptides,

no lesions were formed at 6 µM. At 3 µM, only GMA4CG_V3 and GMA4CG_V4 were effective

in suppressing lesion formation. At 1.5 µM, only GMA4CG_V4 completely suppressed disease

symptoms. The photosynthetic efficiency data further supported GMA4CG_V4 as being more

potent than the other five in reducing gray mold lesions (Figure 2A). We then compared the semi-

in planta antifungal activity of GMA4CG_V4 and GMA4CG_V6. Both peptides completely

suppressed disease symptoms at 1.5 and 3 µM under the applied conditions. However, at a lower

concentration of 0.75 µM, GMA4CG_V6 was more effective in suppressing gray mold disease

symptoms than GMA4CG_V4 (Figures 2C and D).

GMA4CG_V6 enhances permeability of the plasma membrane of *B. cinerea* spores and

germlings

To determine if GMA4CG_V6 permeabilizes the fungal plasma membrane, the spores and

germlings of *B. cinerea* were treated with 1.5 µM GMA4CG_V6 for 30 min followed by the

addition of SYTOX^TM Green (SG) Nucleic Acid Stain. SG is a membrane-impermeant dye, which

fluoresces when it binds to RNA and DNA and is a useful indicator of membrane permeability.

SG uptake was observed in nearly all fungal spores and germlings challenged with this peptide,

but not in controls without peptide (Figure 3A). Thus, we concluded that one MoA of

GMA4CG_V6 involved increased plasma membrane permeabilization.

Since antifungal activity of GMA4CG_V6 was cation-tolerant, we determined if

membrane permeability still occurred in the presence of 100 mM KCl. In the absence of this salt,

95% of the conidia exposed to 1.5 µM GMA4CG_V6 for 1 h took up SG, indicating increased

membrane permeability. No further time- or concentration dependent increase in membrane

disruption was observed in the absence of this salt (Figure 3B). By contrast, uptake of SG occurred

in less than 10% of the conidia treated with 1.5 µM GMA4CG_V6 for 1 h in the presence of 100

mM KCl. Even after 5 h, only ~17% of the conidia took up SG. A high concentration of 6 µM

GMA4CG_V6 was needed for the permeabilization of ~80% of conidia after 5 h (Figure 3B).

Thus, the membrane-permeabilization ability of GMA4CG_V6 was negatively impacted by the

presence of cations; however, the peptide was still effective in permeabilizing the fungal plasma

membrane at increased concentrations with longer exposure.

We also used our SG assay to compare the ability of GMA4CG_V6, GMA4CG_WT, and

MtDef4 to permeabilize the plasma membrane in germlings in the absence and presence of 100

mM KCl. In the absence of KCl, MtDef4 was more effective in permeabilizing the plasma

membrane than GMA4CG_WT or GMA4CG_V6. However, in the presence of this salt, MtDef4's

ability to permeabilize the plasma membrane was completely lost at low concentrations (< 3 µM).

In contrast, GMA4CG_WT and GMA4CG_V6 retained their ability to permeabilize the plasma

membrane at < 3 µM (Figure 3C).

GMA4CG_V6 is internalized into the vacuole and cytoplasm of *B. cinerea* in a time and

concentration dependent manner

Tetramethyl rhodamine (TMR)-labeled GMA4CG_V6 was imaged using time-lapse confocal

microscopy to gain insight into the temporal nature of peptide internalization in *B. cinerea*

germlings. At an MIC of 1.5 µM, the labeled peptide first bound to the cell surface within the first

3 min and started translocating into vacuoles from multiple entry foci by 19 min (Figure 4A). At

this time point, bright field images of germlings showed abnormal morphology with vacuoles

becoming visible. At 27 min, the vacuoles began to expand by coalescence of small vacuoles

(Figure 4A, arrows) and the peptide localized into the cytoplasm. Shortly after the peptide's

appearance in the cytoplasm the vacuoles appeared to lyse with the peptide rapidly dispersing

throughout the germling cytoplasm and inducing cell death (Fig 4A, T= 59).

At a sub-lethal concentration of 0.5 µM, the labeled peptide was localized at the germling

periphery and formed presumptive endosomes in the cytoplasm. No further progression of the

peptide in the germling cells occurred during the 1 h challenge (Figure S5A). At high concentration

(20 µM), TMR-labeled GMA4CG_V6 was internalized by germlings as early as 4 min. The

peptide rapidly accumulated into the germling cytoplasm and within 13 min was completely

dispersed throughout the cell's interior (Figure S5B). This suggests that at high concentrations,

GMA4CG_V6 may largely bypass intermediate endosome or vacuole localization and accumulate

directly in the cytoplasm, or highly accelerate the translocation from vacuole to cytoplasm and

finally accelerate cell death.

The subcellular localization of TMR-GMA4CG_V6 was determined using two dyes which

specifically stain the plasma and endosomal membranes (FM4-64) and vacuole lumen (CMAC).

Within 30 min of challenging germlings with peptide, TMR-GMA4CG_V6 co-localized with

CMAC in vacuoles and with FM4-64 at the plasma membrane and entry foci (Figure 4B,

highlighted white). Based on the region of interest (ROI01) in Figure 4B, the fluorescence intensity

curve clearly showed that the peak signal of TMR-GMA4CG_V6 overlapped with the signal of

FM4-64 in membrane and CMAC in vacuole (Figure 4C). To determine whether GMA4CG_V6

bound to the cell wall of *B. cinerea*, a cell wall specific dye Calcofluor White (CFW) was incubated

with TMR-GMA4CG_V6 treated *B. cinerea* germlings. As illustrated in Figures S6A and B, CFW

overlapped with TMR fluorescent signal in images captured by confocal microscopy. Considering

the thickness and proximity of germling cell wall and plasma membrane and the limitation of the

resolution of the confocal microscopy, super-resolution microscopy (specifically, lattice structured

illumination microscopy (SIM) was used to further evaluate co-localization of TMR-

GMA4CG_V6 with the cell wall. The fluorescent signal of CFW relative to TMR showed the

TMR signal predominantly just inside the cell wall (Figures S6C, D and E), as shown by arrows,

indicating that dominant TMR-GMA4CG_V6 signal was consistent with intense plasma

membrane localization at the cell periphery, although low level cell wall binding could not be ruled

out.

To further study the binding affinity of GMA4CG_V6 to specific subcellular components

in *B. cinerea* germlings, a series of Fluorescence Recovery After Photobleaching (FRAP)

experiments were performed (Figure S7A-C). We specifically studied the mobility of TMR-

GMA4CG_V6 to subcellular components including germling cell periphery, vacuoles, and

cytoplasm. In our previous study, a significant difference in the mobility of the antifungal peptide

NCR044 to germling and conidial cell walls was observed (Velivelli *et al.*, 2020a), thus, we

performed FRAP experiments with conidial spore periphery as well. According to the normalized

fluorescence recovery curve, TMR-GMA4CG_V6 recovery in the germling cytoplasm had the

greatest overall mobility among all four targeted components and the mobility in vacuoles was

relatively low (Figure S7A). Immobile fraction and half time of fluorescence recovery data were

also collected as summarized in Figures S7B and C. GMA4CG_V6 in germling vacuoles showed

a significantly higher ($p < 0.0001$) immobile fraction ($82.6 \pm 3.2\%$) than other subcellular

components and the recovery time was much shorter (Tau= 19.27 sec, R^2= 0.8496). In cell

periphery of germlings and conidia, no significant difference was observed in the immobile

fraction (52.5± 4.2% and 51.0 ± 4.9%, respectively, p> 0.05) or in the half recovery time (37.4

sec, R^2= 0.9133 and 38.4 sec, R^2= 0.9257, respectively). The immobile fraction of the peptide was

lower (p= 0.0031) in germling cytoplasm (35.2± 4.3%) as compared to that in germling cell

periphery and vacuole but showed no significant difference with that of the spore periphery. The

mobility of peptide in germling cytoplasm was the highest among the four selected subcellular

components (Tau= 40.68 and R^2= 0.9636). Collectively, the FRAP analysis indicated that the

mobility of the peptide at the cell periphery and cytoplasm was similar but much different in the

vacuoles. The low mobility in vacuoles is likely because transport to the vacuole lumen is not a

continuum like other compartments and multistep mobility into the vacuole itself would be very

high as the Tau values show.

Because of the difference in the cation tolerant antifungal activity of MtDef4 and

GMA4CG, we examined the localization of MtDef4, GMA4CG_WT, and GMA4CG_V6 on the

spores of *B. cinerea* in the presence of 100 mM KCl. The fluorescently labeled MtDef4 localized

to the cell periphery in the absence of this salt, but this localization was significantly reduced in

its presence (p = 0.001). In contrast, cell periphery localization of GMA4CG_WT and

GMA4CG_V6 was not affected by salt (p> 0.05) (Figures 4D and E).

GMA4CG_WT and GMA4CG_V6 bind to multiple phosphoinositides

MtDef4 binds to phosphatidic acid (PA) in fungal membranes as part of its MoA (18). Therefore,

we tested if MtDef4-derived GMA4CG_V6 also binds to PA. A protein-lipid overlay assay was

performed with various biologically relevant phospholipids immobilized on a lipid strip (Echelon

Bioscience, UT). GMA4CG_V6 bound to several phosphoinositides including phosphatidylinstol

(PI) mono- and bisphosphates, PA, and phosphatidylserine, but with higher affinity for PI4P and

PI5P than for PA (Figure S8). GMA4CG_WT also bound to the same phospholipids as

GMA4CG_V6. His$_6$, the C-terminal tag attached to each peptide used in the assay for antibody

detection, did not bind to the tested phospholipids. These results indicated that GMA4CG_V6 was

more promiscuous in its binding to the membrane resident phospholipids than MtDef4 (Sagaram

et al., 2013).

Spray-applied GMA4CG_V6 provided preventative and curative antifungal activity against

the gray mold disease

We tested the potential of GMA4CG_WT and GMA4CG_V6 to provide preventative protection

from gray mold disease by spraying two-week-old *N. benthamiana* plants with each peptide at

concentrations of 12 and 24 µM followed 2 h later with the spray-inoculum of *B. cinerea* spores.

Each peptide spray-applied at 12 or 24 µM reduced gray mold disease symptoms compared with

control plants inoculated with pathogen only (Figures 5A and S9A). Both peptides significantly

prevented the spread of gray mold on *N. benthamiana* plants (Figures 5A and 5B). Notably, plants

sprayed with GMA4CG_V6 and subsequently inoculated with the pathogen looked as symptom

free as non-inoculated mock (water)-sprayed plants. This data clearly demonstrated higher potency

of GMA4CG_V6 over GMA4CG_WT in reducing gray mold disease in *N. benthamiana* plants in

a preventative manner. In addition, GMA4CG_V6 was more effective in limiting disease severity

than GMA4CG (Figures S9A and B).

We also tested GMA4CG_WT and GMA4CG_V6 for their ability to provide curative

protection from this disease at three different time points following fungal infection. *N.*

benthamiana plants were pre-inoculated with freshly prepared *B. cinerea* spores for 2, 15, and 30

h after which each peptide at 12 µM was sprayed onto the plants. *B. cinerea* spores germinate after

12 h (Bi *et al.*, 2021), and thus, fungal infection of the leaf surface was already initiated at 15 and

30 hpi. Disease symptoms were rated at 72 hpi. Both peptides provided significant curative (p =

0.0019) protection from gray mold (Figures 5C and S9C). However, GMA4CG_V6 conferred

significantly greater (p<0.0001) curative antifungal activity than GMA4CG_WT sprayed on plants

pre-inoculated for 15 h (Figure 5D). The reduced disease severity in GMA4CG_V6 sprayed plants

further confirmed greater efficacy of this peptide (Figure S9C). In plants pre-inoculated for 2 h or

30 h, both peptides provided robust control of disease symptoms with similar efficacy (Figures 5D

and S9D). Furthermore, our data showed the potential of GMA4CG_V6 in reducing the spread of

disease symptoms (Figure S9D).

Next, we tested the curative antifungal activity of GMA4CG_WT and GMA4CG_V6 on

tomato (cv. Mountain Spring) plants pre-inoculated for 12 h and subsequently sprayed with each

peptide at a concentration of 6, 12, and 24 µM. Both peptides conferred curative antifungal activity

at all three concentrations (Figure 6A). At concentrations of 12 and 24 µM, no significant

difference was observed between the two peptides in their ability to reduce disease symptoms.

Both peptides provided robust curative protection with pathogen-challenged plants appearing

similar to water-sprayed non-inoculated mock plants. However, at a concentration of 6 µM,

GMA4CG_V6 conferred significantly greater (p< 0.0001) curative antifungal activity than

GMA4CG_WT (Figure 6B).

Visual rating of disease severity clearly revealed superiority of GMA4CG_V6 in

controlling gray mold disease in tomato plants (Figures S10A and B). The cell wall staining of the

tomato leaves from control plants infected with the pathogen alone and from infected plants

sprayed with GMA4CG_V6 confirmed strong inhibition of fungal growth provided by this peptide

(Figure 6C).

GMA4CG_V6 binds strongly to the tomato leaf surface and is not internalized into plant

cells

To determine the distribution of GMA4CG_WT and GMA4CG_V6 on leaf surfaces, fluorescently

labeled peptide was applied onto detached tomato leaves and the uptake of each peptide by the

plant cells was followed using confocal microscopy. In leaves with no applied peptide, only

autofluorescence signal from the plant cells was observed (Figure S11A). Both GMA4CG_WT

and GMA4CG_V6 bound to the exterior of plant cell walls, but no internalization into plant cells

was detected (Figure S11B). After washing the leaves with water, the fluorescence intensity of

GMA4CG_WT was significantly reduced ($p < 0.0001$) but that of GMA4CG_V6 was only slightly

reduced ($p = 0.1146$) (Figure S11C). This result suggests that GMA4CG_V6 adheres to the tomato

leaf surface more than GMA4CG_WT.

DISCUSSION

Modern agriculture relies heavily upon systemic single-site fungicides with preventative and

curative antifungal properties. However, rapid evolution of fungal resistance to fungicides poses a

major threat to crop protection and food security worldwide. New environmentally friendly

antifungal compounds with multifaceted MoA are needed to mitigate resistance development.

Here, inspired by the broad-spectrum antifungal activity, structure, and multi-faceted MoA of the

plant defensin MtDef4, we explored the potential of a short 17-residue peptide for development as

a peptide-based fungicide.

We designed and synthesized six 17-amino acid GMA4CG variants spanning the active γ-

core motif of a plant defensin MtDef4 (El-Mounadi et al., 2016, Sagaram et al., 2013, Sagaram et

al., 2011). We made several modifications of the GMA4CG peptide that maintained or enhanced

its antifungal activity against an economically important fungal pathogen *B. cinerea*. Although we

could not perform complete analysis of all amino acid substitutions, our selected changes revealed

the importance of cationic and hydrophobic residues for antifungal activity of this peptide. Amino

acids we chose to replace were guided partly by the corresponding amino acid sequences of the

homologs of MtDef4 from other plants. In addition, replacement of the internal Cys11 and Cys13

residues to the aromatic hydrophobic amino acid Trp enhanced antifungal activity. The importance

of Trp at these positions was revealed by our observation that an aliphatic hydrophobic residue

Val at these positions was less effective. It is likely that interactions of the two Trp residues with

the Arg residues present in the GMA4CG_V6 primary amino acid sequence contribute

significantly to peptide stability (Shi *et al.*, 2002). Further, Trp is known to insert into membranes

containing anionic phospholipid head groups with high efficiency, thus contributing to the

increased fungal membrane permeability by this designed peptide (Schibli *et al.*, 2002).

We synthesized GMA4CG_V6 as a cyclic peptide with a disulfide bond connecting Cys4

and Cys17 with the notion that it would be more stable, more fungal cell permeable, more readily

internalized into fungal cells, and confer higher antifungal activity than its linear counterpart.

Cyclic peptides are touted as promising antimicrobial molecules since they are more stable to

proteolytic degradation, have high affinity and selectivity for targets, and better cell membrane

permeability (Falanga *et al.*, 2017).

GMA4CG_WT and GMA4CG_V6 each have random coil/unordered structures as

determined experimentally by steady-state wavelength CD spectra showing the absence of any

significant α-helical or β-strand secondary structure content (Greenfield, 2006, Holzwarth & Doty, 1965). Both peptides are predominately random coil in nature with a suggestion of some polyproline II structure in GMA4CG_V6 (Lopes et al., 2014).

Antifungal activity of several cationic plant defensins is abrogated in well-defined synthetic fungal growth medium containing elevated levels of Na^+ or Ca^{2+} (Spelbrink *et al.*, 2004, Terras *et al.*, 1992, Kerenga et al., 2019, Li et al., 2019). At the highest concentration of 6 μM tested, MtDef4 exhibited no antifungal activity against *B. cinerea* in medium containing elevated levels of Na^+, K^+, or Ca^{2+}. The presence of cations likely weakens the electrostatic interactions between this cationic plant defensin and negatively charged fungal membranes. Surprisingly, all six truncated peptide variants derived from MtDef4 exhibited antifungal activity against *B. cinerea* in fungal growth medium containing elevated levels of Na^+, K^+, or Ca^{2+}. Among the six variants, GMA4CG_V6 was the most potent antifungal agent. While the molecular basis for cation-tolerant antifungal activity of these peptides is unclear, the RRRW motif clearly plays an important role in conferring cation-tolerant antifungal activity.

The structure and MoA of GMA4CG_V6 are unlike those of any fungicide currently commercially available. The preventative and curative plant protection properties of GMA4CG_V6 is likely due to its ability to kill fungal pathogens rapidly. It does not have a discrete intracellular molecular target, but instead interferes with multiple cellular processes (Figure 7). A major characteristic of antifungal peptides is their ability to permeabilize the plasma membrane of fungal pathogens. GMA4CG_V6 quickly binds to the cell surface of fungal spores and germlings and subsequently disrupts the plasma membrane of these cells near the entry points. The co-localization of GMA4CG_V6 with the vacuolar dye CMAC revealed that this peptide, at MIC values, localizes to the plasma membrane and vacuoles in germlings of *B. cinerea*. However, it

remains to be determined if vacuole localization is required for the antifungal activity of this
peptide, or simply a fungal detoxification step. It is of interest to note that a small rationally
designed antifungal hexapeptide PAF26 with the sequence Ac-RKKWFW-NH$_2$ also localizes to
vacuoles in the model fungus *Neurospora crassa* (Munoz *et al.*, 2012). Like PAF26,
GMA4CG_V6 at MIC enters vacuoles and causes their coalescence and lysis. It subsequently
accumulates in the cytoplasm. At elevated concentration, GMA4CG_V6 appeared to accumulate
rapidly in the cytoplasm. MtDef4 binds to the bioactive phospholipid PA as part of its MoA (El-
Mounadi et al., 2016, Sagaram et al., 2013). In contrast, GMA4CG_V6 derived from this defensin
binds to multiple bioactive phospholipids including PA. It preferentially binds to the
phosphoinositides PIPs in the phospholipid-protein overlay assay. Further studies are needed to
understand the role promiscuous interactions with multiple phospholipids play in the multi-faceted
MoA of GMA4CG_V6.

Current crop protection research is guided by a strong demand for sustainable fungicides
that are efficacious, less prone to resistance development, environmentally friendly, and
nonphytotoxic. In semi-*in planta* antifungal assays using detached *N. benthamiana* leaves, all six
GMA4CG analogs significantly reduced gray mold disease lesions caused by *B. cinerea* infection
with GMA4CG_V6 proving most effective. Under the conditions employed, spray application of
GMA4CG_V6 conferred strong preventative and curative antifungal activity against this disease
in young *N. benthamiana* and curative antifungal activity in tomato. These fungicidal effects were
observed readily even though the dose, application frequency, and distribution formulation have
yet to be optimized. In our studies, GMA4CG_V6 was more effective than GMA4CG_WT in
reducing disease symptoms suggesting that the amino acid substitutions introduced in
GMA4CG_V6 are important for its *in vitro* as well as *in planta* antifungal activity. Although

defensin-derived peptides offer the benefits of small size, structural simplicity, and efficiency of

peptide modifications, their commercial use as peptide fungicides will require new innovations in

peptide production at scale to be economical. In addition, efficacy of these peptides for fungal

disease control will need to be assessed in large-scale field trials and their stability to proteolytic

degradation and potential phytotoxicity under field conditions will also need to be evaluated.

Recent studies have shown the promise of antifungal peptides as topical fungicides. A

fusion of the animal AMP dermaseptin and thanatin inhibits the germination of the spores of the

obligate biotrophic fungal pathogen *Phakopsora pachyrhizi* and reduces the symptoms of Asian

rust when spray-applied to soybean leaves (Schwinges *et al.*, 2019). Spray-application of the

antifungal peptide NCR044 from *M. truncatula* on tomato and *N. benthamiana* plants provided

protection from the gray mold disease (Velivelli et al., 2020a). The present study showed that a

designed defensin-derived peptide spanning the functionally active γ-core motif with multi-faceted

MoA exhibited improved preventative and curative antifungal activity against the gray mold

disease of plants. Hundreds of sequences divergent antifungal plant defensins in the AMP database

can now be exploited as templates for the design and development of small peptide-based

fungicides with diverse MoA for future use in agriculture (Shafee & Anderson, 2019, Parisi et al.,

2019).

EXPERIMENTAL PROCEDURES

Fungal cultures and spore suspensions

The fungal strain *Botrytis cinerea* T4 was cultured in V8 (20 %) medium for 7-25 days at 25°C.

Fungal spores were harvested by flooding the fungal growth plates with sterile water. The spore

suspension was filtered through two layers of Miracloth, centrifuged at 13,600 rpm for 1 min,

washed, and re-suspended in low-salt Synthetic Fungal Medium (SFM) (Liang *et al.*, 2001). The

spore suspension of each fungal pathogen was adjusted to the desired spore density of ~10^5

spores/mL using a hemocytometer.

Peptide synthesis and purification

The peptides GMA4CG_WT and GMA4CG_V1-V6 were chemically synthesized with an initial

purity of ~80-85% from multiple commercial vendors and further purified using a C-18 reverse

phase HPLC as described (Velivelli *et al.*, 2020b). The HPLC fractions containing the peptide

were lyophilized and resuspended in the nuclease-free water. Peptide concentrations were

determined using the BCA assay. For peptides containing Trp, peptide concentration was also

determined using NanoDrop 2000c Spectrophotometry (Thermo-Fisher Scientific, MA) to confirm

concentrations determined by the BCA assay.

Antifungal activity assays

The minimal inhibitory concentration (MIC) of each peptide was determined using the resazurin

cell viability assay (Chadha & Kale, 2015, Li et al., 2019). Briefly, this involved incubation of the

pathogen/peptide mixture for 48 h, adding 0.01% (w/v) resazurin solution to each well, and re-

incubating overnight. A change in color of the resazurin dye from blue to pink or colorless

indicated the presence of live fungal cells. MIC was determined as the minimal concentration of

peptide at which no change in color was observed. The MIC value of each peptide was also

determined in SFM supplemented with 100 mM NaCl, 100 mM KCl, or 2 mM $CaCl_2$.

Membrane permeabilization assay using SYTOX^{TM} Green (SG) uptake

The fluorescent dye SG (Thermo-Fisher Scientific, MA) was used to identify GMA4CG_V6-

induced membrane permeabilization of *B. cinerea* conidia and germlings as described previously

(Li et al., 2019). After exposure to 1.5 µM GMA4CG_V6 for 30 min, *B. cinerea* spores and

germlings were imaged using fluorescence microscopy on a Leica SP8-X confocal microscope

with an excitation and emission wavelength of 488 and 508-544 nm, respectively.

The GMA4CG_V6 induced membrane permeabilization of conidia was also assessed

quantitatively in the absence and presence of 100 mM KCl. Fresh conidia at a concentration of $1\times$

10^5/mL were incubated with GMA4CG_V6 at $1\times$, $2\times$, and $4\times$ MIC for 1 h in the absence and

presence of 100 mM KCl for 1 h and 5 h, respectively. Samples were deposited into glass-bottom

petri dishes and imaged by confocal microscopy at an excitation wavelength of 488 nm and an

emission wavelength ranging from 508 to 544 nm in the presence of 1 µM SG added 15 min prior

to imaging. Fresh conidia incubated with SG alone were used as negative controls. A Leica SP8-

X confocal microscope was used for all confocal imaging and Leica LASX (version 3.1.5.)

software was used to process the images. The percentage of conidia showing membrane

permeabilization was determined by counting the number of cells showing fluorescent signal

relative to the total number of cells counted.

The effect of MtDef4, GMA4CG_WT, and GMA4CG_V6 on membrane integrity of *B.*

cinerea germlings was determined using a SG (final concentration: 1 µM) uptake quantification

assay in the presence and absence of 100 mM KCl (Li et al., 2019). In brief, membrane

permeabilization in *B. cinerea* germlings treated with different concentrations (0.187, 0.375, 0.75,

1.5, and 3 µM) of MtDef4, GMA4CG_WT, and GMA4CG_V6 was measured at 5 h in the

presence and absence of 100 mM KCl. SG uptake was quantified by measuring the fluorescence

in a SpectraMax® M3 spectrophotometer (excitation: 488 nm; cutoff: 515 nm; emission: 530 nm).

Internalization and subcellular localization of GMA4CG_V6 in *B. cinerea* germlings

Time-lapse confocal laser scanning microscopy was used to monitor uptake and subcellular

localization of fluorescently labeled peptide as described previously (Velivelli et al., 2020a). Crude

TMR fluorophore labeled GMA4CG_V6 (TMR-GMA4CG_V6) was custom synthesized

(Agentide Co, Piscataway, NJ) and purified by HPLC. *B. cinerea* germlings were generated by

incubating conidia overnight in SFM medium. They were challenged with TMR-GMA4CG_V6 at

concentrations of 0.5, 1.5, or 20 µM and imaged by time-lapse confocal microscopy. The excitation

and emission wavelengths for imaging TMR-GMA4CG_V6 were 562 and 580 to 680 nm,

respectively.

The sub-localization of GMA4CG_V6 was determined by visualizing its localization with

the vacuole-specific dye CMAC and the cell membrane specific dye FM4-64. The excitation and

emission wavelengths for CMAC were 353 nm and 405 to 480 nm, respectively, and for FM4-64

were 562 and 680 to 800 nm, respectively. Co-localized pixels (white overlay) of TMR-

GMA4CG_V6 and CMAC/FM4-64 was displayed using Leica LAS-X software. The threshold for

each channel was 50% and the pixels out of this range were considered uncorrelated. The threshold

for the background was set as 20%. Images were captured 30 min after incubation with peptide.

Quantification of cell wall periphery binding intensity

The spores of *B. cinerea* were treated with 1.5 µM DyLight550-labeled MtDef4, TMR-

GMA4CG_WT, and TMR-GMA4CG_V6 for 20 min. Spores of each treatment were imaged using

a C-Apochromat 63× water immersion objective lens in apotome mode with a ZEISS Elyra7 super-

resolution microscope at excitation and emission wavelengths of 561 and 590 nm, respectively.

Images were processed using the SIM module and 3D leap processing with auto sharpness setting

in ZEN Black 3.0 SR FP2 software. For each captured cell, the cell surface binding intensity was

quantified by free hand drawing a line intensity profile along the cell surface. The fluorescence

intensity of each cell wall periphery was normalized by its perimeter length as described (Velivelli

et al., 2020a). The lower threshold was set to 7000 to subtract the signal in the cytoplasm.

Semi-*in planta* antifungal activity assays

The semi-*in planta* antifungal activity of each peptide against *B. cinerea* was determined using *N.*

benthamiana Nb1 detached leaves as described previously (Li et al., 2019). Each peptide was

tested at concentrations ranging from 0.75, 1.5, 3, and 6 µM. Following incubation of each

peptide/fungal spore mixture at room temperature for 48 h, leaves were photographed in white

light. High-resolution fluorescence images were also taken using CropReporter (PhenoVation,

Wageningen, Netherlands). These images depicted the calculated F_V/F_M (maximum quantum yield

of photosystem II) values of diseased area affected by *B. cinerea* infection. Colors in the images

show five different classes ranging from class I to class V (0.000 to 0.700) depicting varying

degrees of tissue damage. On one end of the scale, green represents class V (0.600 to ≥0.700) high

photosynthesis efficiency. On the other end of the scale, red represents class I (0.000 to 0.160) low

photosynthesis efficiency.

In planta antifungal activity assays

To test preventative and curative antifungal activity of the peptides against *B. cinerea*, four-week-old *N. benthamiana* Nb1 plants grown under controlled growth conditions were used. For preventative activity, GMA4CG_WT and GMA4CG_V6 at 12 or 24 µM concentration were sprayed (2 mL/plant) on experimental plants. Control plants were sprayed with 2 mL of water/plant. The peptide solution or water was allowed to dry out on the surface of the leaves. *B. cinerea* spores (~8×10^4) suspended in 0.5 × SFM were sprayed on the experimental and control tomato plants (1 mL/plant), transferred to Ziploc plastic boxes under high humidity, and disease symptoms recorded after 48 h. Photographs were taken at 72 h post-inoculation using both white light and CropReporter (PhenoVation, Wageningen, Netherlands). Quantification of symptoms was made by determining maximum quantum yield of PSII (Fv/Fm) as described above for the semi-*in planta* assays.

For the curative antifungal activity tests, *N. benthamiana* Nb1 plants of similar age were first sprayed with *B. cinerea* spores (~8×10^4) suspended in 0.5× SFM (1 mL/plant). After 2, 15, and 30 h of pathogen challenge, 2 mL of GMA4CG_WT or GMA4CG_V6 was sprayed on the plants at a concentration of 12 µM. Control plants were each sprayed with 2 mL of water. Disease symptoms were observed after 48 h of pathogen challenge. Plants were photographed at 72 h post infection in white light and quantification was done as described above. The curative antifungal activity of GMA4CG_WT and GMA4CG_V6 at concentration of 6, 12, and 24 µM was tested on 3-week-old tomato (*Solanum lycopersicum* L. cv. Mountain Spring) plants pre-inoculated with *B. cinerea* for 12 h. The severity of disease symptoms was rated using the scoring method for each inoculated plant as follows: 0, no symptoms; 1= 1-25%, 2= 26-50%, 3= 51-75% and 4= 76-100% of leaf surface infected (Segarra *et al.*, 2017).

Visualization of fungal growth on tomato leaf surface

Fresh *B. cinerea* conidia were collected and sprayed on leaves of 3-week-old tomato plants and

allowed to germinate for 12 h. After that, GMA4CG_V6 at 6 µM was applied on the same leaves

and incubated in a container under high humidity. Leaves applied with spores and sterile water

were set as negative controls. After 48 h, leaves were fixed in 4% paraformaldehyde, stained with

CFW for 30 min, and images were captured by an Axio Zoom.V16 microscope (ZEISS) with a

PlanNeoFluar Z 1.0x objective under tile mode. The excitation wavelength was 335-383 nm, and

the emission wavelength was 420-470 nm. Images were visualized and processed by Zen Black

3.0 SR FP2 software (ZEISS).

Cell death assay of GMA4CG_V6 with propidium iodide

To determine fungicidal activity, freshly harvested conidia of *B. cinerea* were incubated with 1.5

µM GMA4CG_V6 for 1, 6, and 24 h. After incubation, conidia were centrifuged and washed twice

in 1x SFM and re-suspended in the same medium. The spores were subsequently allowed to

germinate overnight. Microscope images were taken, and germination rate was calculated.

Propidium iodide (PI) at 0.5 µg/mL was used as a marker to count dead spores. Conidia of *B.*

cinerea were incubated with GMA4CG_V6 for 0.25, 0.5, 1, and 2h. PI was added to the

spores/peptide mixture 10 min in advance of detection. Images were captured on the Leica SP8-X

confocal microscope (excitation: 535 nm, emission: 600-650 nm).

Circular dichroism spectroscopy

A steady-state wavelength circular dichroism spectrum was collected for GMA4CG_WT and GMA4CG_V6 (~0.05 mM peptide, 20 mM sodium acetate, pH 5.6) on a calibrated Aviv Model 410 spectropolarimeter (Lakewood, NJ). Using a quartz cell of 0.1 cm path length and a temperature of 20 °C, the data was recorded in 0.5 nm increments between 200 and 260 nm with a bandwidth of 1.0 nm and a time constant of 1.0 s. The reported steady-state wavelength spectrum was the result of averaging two consecutive scans, subtracting a blank spectrum, and automatic line smoothing using Aviv software.

Antifungal activity of the alanine scanning variants of GMA4CG_V6 with no disulfide bond

The minimal inhibitory concentration (MIC) of linear GMA4CG_V6 alanine scanning variants was determined using the resazurin cell viability assay as described in the methods section. The MIC value of all variants was also determined in SFM supplemented with 100 mM NaCl.

Cell wall/surface localization analysis of GMA4CG_V6

B. cinerea germlings were treated with 1.5 μM of TMR-GMA4CG_V6 and, after 5 min, CFW (5 μM) was added into the same petri dish. Samples were observed using a confocal (Leica SP8-X) and super-resolution microscope (Zeiss Elyra7). To avoid bleed-through of the two fluorescent signals, confocal images were captured sequentially using a fast line switch configuration (TMR-GMA4CG_V6 excitation: 542 nm, emission: 580-680 nm, and CFW excitation: 405 nm, emission: 410-557 nm). To obtain higher resolution images, the C-Apochromat 63×1.4 oil immersion objective lens in apotome mode was used with the ZEISS Elyra7 super-resolution microscope (Zeiss). The excitation wavelength was 405 nm for the CFW signal and 561 nm for the TMR signal with emission wavelengths of 420-480 and 570-620 nm, respectively. Images were processed

using the SIM module plus 3D and leap processing with autosharpness. The channels were aligned

based on 0.1 µm TetraSpeck™ microspheres (T7279, Thermo Fisher Scientific) using ZEN Black

3.0 SR FP2 software.

Fluorescence Redistribution After Photobleaching (FRAP) experimental setup and analysis

The FRAP experiments were performed on a Leica SP8-X microscope (FRAP Wizard Module,

LAS X Version 3.1.5.16.16308) using the HC PlanApochromat 63× water immersion objective

len (Numerical aperture = 1.2) with the pinhole set to 1 Airy unit and zoom set to 1.77 (Velivelli

et al., 2020a). Images were acquired for the selected cellular regions (germling cytoplasm, vacuole,

cell periphery, and spore cell wall/periphery) after 1 h incubation with GMA4CG_V6 under the

following identical imaging and FRAP conditions: 10 pre-bleach and 700 post-bleach frames (512

x 64 pixels, pixel size = 204 nm) captured at 173 ms intervals without averaging using 550

excitation and 560-625 nm emission wavelengths with PMT detectors and a transmitted light

channel. The power of the 550 nm laser was set at 3.5% during imaging to prevent detectable

photobleaching for the duration of each FRAP experiment (~3 min). A 1.52 µm^2 circular bleach

region was applied with 100% laser excitation power (550 nm wavelength) for 3.5 s immediately

following pre-bleach and prior to post-bleach imaging. FRAP analysis was performed on 24 data

sets for each subcellular region showing no evidence of sample drift during acquisition. A single

exponential curve fitting function was applied post-acquisition to extract the half-maximum tau

(τ$^1\!/_2$), mobile fraction, and immobile fraction for each condition.

Phospholipid binding assays

His-tagged GMA4CG_WT and GMA4CG_V6 was obtained from Biomatik Inc. (Ontario,

Canada) and further purified by HPLC. The His-tagged peptides (GMA4CG_WT and

GMA4CG_V6) phospholipid binding assays were performed using the PIP Strips[TM] (Echelon

Biosciences Inc., CA) as previously described (Islam *et al.*, 2017). Hexa-His peptide (Genscript,

Inc., NJ) was used as a control. The peptide binding to the membrane strips was detected using 0.5

µg/mL His-tag monoclonal antibody purchased from Thermo-Fisher Scientific, MA. The binding

signals were detected using the SuperSignal West Pico Chemiluminescent Substrate Kit (Thermo-

Fisher Scientific, MA) following the manufacturer's protocol. Quantitative analysis of the signal

was performed using Image J (https://imagej.nih.gov/ij/download.html).

The binding of GMA4CG_WT and GMA4CG_V6 on tomato leaf surfaces

The fluorophore-labeled peptides, TMR-GMA4CG_WT and TMR-GMA4CG_V6, were applied

on the surface of tomato leaves and allowed to air dry for 1 h. The leaf samples were subjected to

microscopic observation using a Leica SP8-X confocal microscope (TMR wavelength: excitation:

562 nm; emission: 580-680 nm, and autofluorescence of tomato cells wavelength: excitation: 405

nm; emission: 416-483 nm) with an APO CS2 40× water objective lens before and after a gentle

rinse twice in sterile water. The average signal intensity from 10-12 ROIs of each image was

collected for quantification and 26-32 images were captured. The signal intensity of the after-

wash-sample was normalized by the signal of the sample before washing.

Statistical analysis

All experiments were performed in three or more technical replicates where appropriate and in

three independent experiments. The phospholipid binding assays and quantitative measurements

of cell membrane permeability in *B. cinerea* were performed using two independent biological

replications. Data were graphed and analyzed in GraphPad Prism (version 8.0 or 9.4.1) software

using unpaired, two-tailed Student's *t* tests, and one-way ANOVA with Tukey's honestly

significant difference test (HSD) for comparison of multiple groups. Statistical significance was

accepted at $p < 0.05$.

Acknowledgements

This work was supported, in part, by the National Science Foundation-EAGER grant IOS:1955461

awarded to D.S. We acknowledge imaging support from the Advanced Bioimaging Laboratory

(RRID:SCR_018951) at the Danforth Plant Science Center and usage of the Leica SP8-X confocal

microscope acquired through an NSF Major Research Instrumentation grant (DBI-1337680) and

usage of the ZEISS Elyra 7 Super-Resolution Microscope acquired through an NSF Major

Research Instrumentation grant (DBI-2018962). We are thankful to Dr. Saurav Misra of the

Danforth Center for his attempts to generate structures of the peptides using AlphaFold. This work

was funded by the Wells Fargo IN^2 grant awarded to Peptyde Bio. Part of this research was

performed at the Pacific Northwest National Laboratory (PNNL), a facility operated by Battelle

for the U.S. Department of Energy, including access to the W.R. Wiley Environmental Molecular

Sciences Laboratory (EMSL), a national scientific user facility sponsored by the U.S. DOE

Biological and Environmental Research program. Battelle operates PNNL for the U.S. Department

of Energy under contract DE-AC05-76RL01830.

Author Contributions

Conceptualization: Meenakshi Tetorya, Hui Li, Kirk J. Czymmek, Dilip M. Shah

Funding acquisition: Dilip M. Shah

Methodology: Meenakshi Tetorya, Hui Li, Kirk J. Czymmek, Arnaud Thierry Djami-Tchatchou,

Garry W. Buchko, Dilip M. Shah

Project administration: Dilip M. Shah

Supervision: Kirk J. Czymmek, Dilip Shah

Validation: Kirk J. Czymmek, Dilip Shah

Writing-original draft: Meenakshi Tetorya, Hui Li, Garry Buchko

Writing- review & editing: Dilip M. Shah, Kirk J. Czymmek, Garry W. Buchko

REFERENCES

Baek, M., DiMaio, F., Anishchenko, I., Dauparas, J., Ovchinnikov, S., Lee, G. R*., et al.* (2021)
Accurate prediction of protein structures and interactions using a three-track neural
network. *Science,* **373,** 871-876.
Bi, K., Scalschi, L., Jaiswal, N., Mengiste, T., Fried, R., Sanz, A. B*., et al.* (2021) The Botrytis
cinerea Crh1 transglycosylase is a cytoplasmic effector triggering plant cell death and
defense response. *Nat Commun,* **12,** 2166.
Chadha, S. and Kale, S. P. (2015) Simple fluorescence-based high throughput cell viability assay
for filamentous fungi. *Lett Appl Microbiol,* **61,** 238-244.
Chan, D. I., Prenner, E. J. and Vogel, H. J. (2006) Tryptophan- and arginine-rich antimicrobial
peptides: structures and mechanisms of action. *Biochimica et biophysica acta,* **1758,** 1184-
1202.
Cools, T. L., Struyfs, C., Cammue, B. P. and Thevissen, K. (2017) Antifungal plant defensins:
increased insight in their mode of action as a basis for their use to combat fungal infections.
Future Microbiol, **12,** 441-454.
Dathe, M., Nikolenko, H., Klose, J. and Bienert, M. (2004) Cyclization increases the antimicrobial
activity and selectivity of arginine- and tryptophan-containing hexapeptides. *Biochemistry,*
43, 9140-9150.
Deslouches, B., Phadke, S. M., Lazarevic, V., Cascio, M., Islam, K., Montelaro, R. C*., et al.* (2005)
De novo generation of cationic antimicrobial peptides: influence of length and tryptophan
substitution on antimicrobial activity. *Antimicrobial agents and chemotherapy,* **49,** 316-
322.
El-Mounadi, K., Islam, K. T., Hernandez-Ortiz, P., Read, N. D. and Shah, D. M. (2016) Antifungal
mechanisms of a plant defensin MtDef4 are not conserved between the ascomycete fungi
Neurospora crassa and Fusarium graminearum. *Mol Microbiol,* **100,** 542-559.

Falanga, A., Nigro, E., De Biasi, M. G., Daniele, A., Morelli, G., Galdiero, S., *et al.* (2017) Cyclic Peptides as Novel Therapeutic Microbicides: Engineering of Human Defensin Mimetics. *Molecules,* **22**.

Fisher, M. C., Hawkins, N. J., Sanglard, D. and Gurr, S. J. (2018) Worldwide emergence of resistance to antifungal drugs challenges human health and food security. *Science,* **360,** 739-742.

Fisher, M. C., Henk, D. A., Briggs, C. J., Brownstein, J. S., Madoff, L. C., McCraw, S. L., *et al.* (2012) Emerging fungal threats to animal, plant and ecosystem health. *Nature,* **484,** 186-194.

Fones, H. N., Bebber, D. P., Chaloner, T. M., Kay, W. T., Steinberg, G. and Gurr, S. J. (2020) Threats to global food security from emerging fungal and oomycete crop pathogens. *Nature Food,* **1,** 332-342.

Goyal, R. K. and Mattoo, A. K. (2014) Multitasking antimicrobial peptides in plant development and host defense against biotic/abiotic stress. *Plant science (Shannon, Ireland),* **228,** 135-149.

Greenfield, N. J. (2006) Using circular dichroism spectra to estimate protein secondary structure. . *Nature protocols,* **1,** 2876-2890.

Holzwarth, G. M. and Doty, P. (1965) The ultraviolet circular dichroism of polypeptides. . *J Am Chem Soc,* **87,** 218-228.

Islam, K. T., Velivelli, S. L. S., Berg, R. H., Oakley, B. and Shah, D. M. (2017) A novel bi-domain plant defensin MtDef5 with potent broad-spectrum antifungal activity binds to multiple phospholipids and forms oligomers. *Sci Rep,* **7,** 16157.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., *et al.* (2021) Highly accurate protein structure prediction with AlphaFold. *Nature,* **596,** 583-589.

Kaur, J., Sagaram, U. S. and Shah, D. M. (2011) Can plant defensins be sused to engineer durable commercially useful resistance in crop plants? *Fungal Biology Reviews,* **25,** 128-135.

Kerenga, B. K., McKenna, J. A., Harvey, P. J., Quimbar, P., Garcia-Ceron, D., Lay, F. T., *et al.* (2019) Salt-tolerant antifungal and antibacterial activities of the corn defensin ZmD32. *Frontiers in Microbiology,* **10,** 795.

Killian, J. A. and von Heijne, G. (2000) How proteins adapt to a membrane-water interface. *Trends Biochem Sci,* **25,** 429-434.

Kovaleva, V., Bukhteeva, I., Kit, O. Y. and Nesmelova, I. V. (2020) Plant Defensins from a Structural Perspective. *Int J Mol Sci,* **21**.

Lacerda, A. F., Vasconcelos, E. A., Pelegrini, P. B. and Grossi de Sa, M. F. (2014) Antifungal defensins and their role in plant defense. *Front Microbiol,* **5,** 116.

Leannec-Rialland, V., Atanasova, V., Chereau, S., Tonk-Rugen, M., Cabezas-Cruz, A. and Richard-Forget, F. (2022) Use of Defensins to Develop Eco-Friendly Alternatives to Synthetic Fungicides to Control Phytopathogenic Fungi and Their Mycotoxins. *J Fungi (Basel),* **8**.

Li, H., Velivelli, S. L. and Shah, D. M. (2019) Antifungal potency and modes of action of a novel olive tree defensin against closely related ascomycete fungal pathogens. . *Mol Plant Microbe Interact,* **32,** 164901664.

Liang, J., Shah, D. M., Wu, Y., Rosenberger, C. A. and Hakimi, S. M. (2001) Antifungal polypeptide from alfalfa and methods for controlling plant pathogenic fungi. USA: Monsanto Company.

Lobo, F. and Boto, A. (2022) Host-defense peptides as new generation phytosanitaries: low toxicity and low induction of antimicrobial resistance. *Agronomy,* **12,** 1614.

Lopes, J. L., Miles, A. J., Whitmore, L. and Wallace, B. A. (2014) Distinct circular dichroism spectroscopic signatures of polyproline II and unordered secondary structures: applications in secondary structure analyses. *Protein Sci,* **23,** 1765-1772.

Lu, J., Xu, H., Xia, J., Ma, J., Jun, X., Li, Y*., et al.* (2020) D- and Unnatural Amino Acid Substituted Antimicrobial Peptides With Improved Proteolytic Resistance and Their Proteolytic Degradation Characteristics. *Front Microbiol,* **11,** 563030.

Munoz, A., Marcos, J. F. and Read, N. D. (2012) Concentration-dependent mechanisms of cell penetration and killing by the de novo designed antifungal hexapeptide PAF26. *Mol Microbiol,* **85,** 89-106.

Oliver, R. P. and Hewitt, H. G. (2014) Fungicides in Crop protection. In: *Fungicides in Crop Protection.* CAB International, Oxfordshire, Boston.

Parisi, K., Shafee, T. M. A., Quimbar, P., van der Weerden, N. L., Bleackley, M. R. and Anderson, M. A. (2019) The evolution, function and mechanisms of action for plant defensins. *Semin Cell Dev Biol,* **88,** 107-118.

Sagaram, U., El-Mounadi, K., Buchko, G. W., Berg, H., Kaur, J., Pandurangi, R*., et al.* (2013) Structural and functional studies of a phosphatidic-acid binding antifungal plant defensin MtDef4: Identification of an RGFRRR motif governing fungal cell entry. *PLoS One e82485.*

Sagaram, U. S., Pandurangi, R., Kaur, J., Smith, T. J. and Shah, D. M. (2011) Structure-activity determinants in antifungal plant defensins MsDef1 and MtDef4 with different modes of action against *Fusarium graminearum. PloS one,* **6,** e18550.

Sathoff, A. E., Velivelli, S., Shah, D. M. and Samac, D. A. (2019) Plant Defensin Peptides have Antifungal and Antibacterial Activity Against Human and Plant Pathogens. *Phytopathology,* **109,** 402-408.

Savary, S., Willocquet, L., Pethybridge, S. J., Esker, P., McRoberts, N. and Nelson, A. (2019) The global burden of pathogens and pests on major food crops. *Nature Ecology & Evolution,* **3,** 430-439.

Schibli, D. J., Epand, R. F., Vogel, H. J. and Epand, R. M. (2002) Tryptophan-rich antimicrobial peptides: comparative properties and membrane interactions. *Biochem Cell Biol,* **80,** 667-677.

Schwinges, P., Pariyar, S., Jacob, F., Rahimi, M., Apitius, L., Hunsche, M*., et al.* (2019) A bifunctional dermaseptin-thanantin dipeptide functionalizes the crop surface for sustainable pest management. *Green Chemistry*.

Segarra, G., Casanova, E., Borrero, C., Avile´s, M. and Trillas, I. (2017) The suppressive effects of composts used as growth media against *Botrytis cinerea* in cucumber plants. *Eur J Plant Pathol,* **117,** 393-402.

Shafee, T. M. and Anderson, M. A. (2019) A quantitative map of protein sequence space for the cis-defensin superfamily. . *Bioinformatics,* **35,** 743-752.

Shi, Z., Olson, C. A. and Kallenbach, N. R. (2002) Cation-pi interaction in model alpha-helical peptides. *J Am Chem Soc,* **124,** 3284-3291.

Spelbrink, R. G., Dilmac, N., Allen, A., Smith, T. J., Shah, D. M. and Hockerman, G. H. (2004) Differential antifungal and calcium channel-blocking activity among structurally related plant defensins. *Plant Physiol,* **135,** 2055-2067.

Terras, F. R., Schoofs, H. M., De Bolle, M. F., Van Leuven, F., Rees, S. B., Vanderleyden, J.*, et
al.* (1992) Analysis of two novel classes of plant antifungal proteins from radish (Raphanus
sativus L.) seeds. *J Biol Chem,* **267,** 15301-15309.
Tucker, A. T., Leonard, S. P., DuBois, C. D., Knauf, G. A., Cunningham, A. L., Wilke, C. O.*, et
al.* (2018) Discovery of Next-Generation Antimicrobials through Bacterial Self-Screening
of Surface-Displayed Peptide Libraries. *Cell,* **172,** 618-628 e613.
van der Weerden, N. L., Bleackley, M. R. and Anderson, M. A. (2013) Properties and mechanisms
of action of naturally occurring antifungal peptides. *Cell Mol Life Sci,* **70,** 3545-3570.
van Esse, H. P., Reuber, T. L. and van der Does, D. (2020) Genetic modification to improve disease
resistance in crops. *The New phytologist,* **225,** 70-86.
Velivelli, S. L., Czymmek, C. J., Li, H., Shaw, J. B., Buchko, G. W. and Shah, D. M. (2020a)
Antifungal symbiotic peptide NCR044 exhibits unique structure and multi-faceted
mechanisms   of   action   that   confer   plant   protection.   *PNAS,*
www.pnas.org/cgi/doi/10.1073/pnas2003526117.
Velivelli, S. L. S., Czymmek, K. J., Li, H., Shaw, J. B., Buchko, G. W. and Shah, D. M. (2020b)
Antifungal symbiotic peptide NCR044 exhibits unique structure and multifaceted
mechanisms of action that confer plant protection. *Proc Natl Acad Sci U S A,* **117,** 16043-
16054.

Figures and Legends

Figure 1.

A

```
MtDef4:     RTCESQSHKFKGPCASDHNCASVCQTERFSGGRCRGFRRRCFCTTHC
```

	Sequences	**Net charge**	**% hydrophobic amino acids**
GMA4CG_WT:	GGRCRGFRRRCFCTTHC	4.9	11.8
GMA4CG_V1:	GGRCKGFRRRCFCTRIC	5.8	17.7
GMA4CG_V2:	GGRCRGFRRRCFCTRIC	5.8	17.7
GMA4CG_V3:	GGRCKGFRRRVFVTRIC	5.9	29.4
GMA4CG_V4:	FGRCRGFRRRCFCWRWC	5.8	29.4
GMA4CG_V5*:	FGZCZGFZZZCFCWZWC	5.8	29.4
GMA4CG_V6:	GGRCKGFRRRWFWTRIC	5.9	29.4

B

		B. cinerea MIC (µM)		
Peptide	**SFM**	**SFM+ 100 mM NaCl**	**SFM+ 100 mM KCl**	**SFM+ 2 mM CaCl$_2$**
MtDef4	0.75	>6	>6	>6
GMA4CG_WT	3	3	6	6
GMA4CG_V1	3	1.5	3	3-6
GMA4CG_V2	6	3	6	6
GMA4CG_V3	3	1.5-3	>6	>6
GMA4CG_V4	1.5-3	1.5	1.5	3
GMA4CG_V5	3-6	6	>6	>6
GMA4CG_V6	1.5	1.5	1.5-3	3

Fig 1. Amino acid sequences and antifungal activity of MtDef4-derived GMA4CG_WT and its variants. A. Amino acid sequence, net charge, and percentage of hydrophobic residues for GMA4CG_WT and its variants. The amino acid sequence of MtDef4 is shown on top with its γ-core motif highlighted by a red rectangle. The amino acid substitutions in GMA4CG_WT are shown in red. Disulfide bonds verified in GMA4CG_V3 and GMA4CG_V6 are shown with a link. The letter Z indicates the unnatural amino acid 2, 4-diaminobutyric acid. **B.** The antifungal activity of MtDef4, GMA4CG_WT, and its variants against *B. cinerea* in the absence and presence of cations and against *F. graminearum* in the absence of cations. MIC (Minimal Inhibitory Concentration) of each peptide was determined by the resazurin assay. SFM (Synthetic Fungal Medium) is a low salt medium used in this study.

Figure 2.

Fig 2. Topical application of GMA4CG_WT and its variants on detached *N. benthamiana* leaves confers resistance to gray mold. A. Scheme for application of GMA4CG_WT and GMA4CG_V1-V5 on tobacco leaves. Gray mold lesions on *N. benthamiana* leaves were observed and photographed at 48 hpi. The white light and CropRepoter images of the leaves are shown. **B.** Box & whisker plots were used to represent the quantification of lesion sizes. **C**. Scheme for application of GMA4CG_V4 and GMA4CG_V6 on *N. benthamiana* leaves. The white light and CropReporter images of the infected leaves are shown. **D**. Box & whisker plots were used to represent the quantification of lesion sizes. In **B** & **D**, the horizontal lines represent the median and boxes indicate the 25^{th}-75^{th} percentile. One-way analysis of variance (ANOVA) was used to determine statistical differences in lesion sizes between control (*B. cinerea*) and peptides (GraphPad Prism-version 9.4.1). Three technical replications of tobacco leaves (n = 3) were tested in three independent biological replications. Asterisks represent significant differences between

different groups ($^{ns}p > 0.05$, $*p \leq 0.05$ $**p < 0.01$, $****p \leq 0.0001$) using Tukey's honestly significant
difference test for comparison of multiple groups.

Figure 3.

Fig 3. GMA4CG_V6 permeabilizes cell membrane of *B. cinerea*. A. Confocal microscopy
images of the SG uptake in *B. cinerea* spores and germlings treated with 1.5 µM of GMA4CG_V6
for 30 min. Spores and germlings treated with H₂O served as controls. Scale bars = 20 µm. **B.**
Representative images showing SG signals in *B. cinerea* spores treated with 1.5, 3, and 6 µM of
GMA4CG_V6 for 1 h with no salt, and for 1 h and 5 h with 100 mM KCl. The outline of each
spore is highlighted by a white dashed line. Scale bar = 20 µm. The permeabilization rate (%) of
each treatment is labeled on the images. **C.** Quantitative measurements of cell membrane
permeability in terms of fluorescence emitted by *B. cinerea* germlings treated with various
concentrations of MtDef4 (red), GMA4CG_WT (green), and GMA4CG_V6 (purple) with (dashed
line) and without (solid line) 100 mM KCl. Membrane permeabilization increased with increasing

concentration of each peptide. Values are means of two biological replications. Error bars indicate
standard deviations. Fluorescence measurements were taken at 5 h.

Figure 4.

Fig 4. Internalization of GMA4CG_V6 into *B. cinerea*. A. Time-lapse confocal microscopy images of *B. cinerea* showing internalization of 1.5 µM TMR-GMA4CG_V6 over time (T in min). Images were captured every 4 min starting from 3 min. Arrows indicate vacuoles. Scale bars = 10 µm. **B.** Individual fluorescent (CMAC, TMR, and FM4-64) and channel merged images of *B. cinerea* germlings treated with GMA4CG_V6 for 30 min. The overlap signal of CMAC and TMR or TMR and FM4-64 are shown as white pixels. The threshold for each channel and the background was 50% and 20%, respectively. Scale bars = 10 µm. **C.** The fluorescent signal intensity of the ROI in **B**. Black arrows indicates TMR-GMA4CG_V6 and FM4-64 signals overlap at the membrane and the red arrows indicate TMR-GMA4CG_V6 and CMAC signals overlap at the vacuoles. **D.** Cell surface fluorescence shown in 2.5D. The vertical direction indicates the fluorescence intensity of each pixel with taller peaks representing greater fluorescence intensity. The corresponding 2D images are shown. **E.** Box plot for cell surface fluorescence quantification. Total fluorescence intensity of each cell was normalized to the cell perimeter length. The lower threshold was set to 7000, the substrate signal in the cytoplasm. Statistical analyses were performed using a two-tailed *t*-test assuming unequal variance (GraphPad Prism 8.0). *P*-value over 0.05 indicated no significant difference and less than 0.05 indicates a significant difference. n = 24-39.

Figure 5.

Fig 5. **GMA4CG_V6 exhibits preventative and curative antifungal activity against gray mold**
in *N. benthamiana* plants. A. Preventative activity of GMA4CG_V6. Four-week-old *N.*
benthamiana plants were sprayed with either 2 mL of water or peptide (GMA4CG_WT or
GMA4CG_V6) at 12 µM or 24 µM concentrations prior to spray-inoculation with 1 mL of 8×10^4
spores per mL of *B. cinerea* fresh spore suspension. The mock plants were only sprayed with water
(no fungal spore suspension). The CropReporter images of plants were taken 72 hpi. Maximum
quantum yield of PSII photochemistry (Fv/Fm) was determined at 72 hpi. Red color represents
class I, low photosynthesis efficiency attributed to extensive tissue damage. Green color indicates
class V, high photosynthesis efficiency. **B.** The calculated photosynthetic quantum yield (Fv/Fm)
for individual *N. benthamiana* plants. **C.** Curative antifungal activity of GMA4CG_V6. Four-
week-old *N. benthamiana* plants were first spray-inoculated using 1 mL of 8×10^4 *B. cinerea* spore
suspension. At 2, 15, or 30 hpi the plants were sprayed with 2 mL of peptide (GMA4CG_WT or
GMA4CG_V6) at a concentration of 12 µM or water. The mock plants were only sprayed with

water (no fungal spore suspension). Maximum quantum yield was determined as described above. **D.** The calculated photosynthetic quantum yield (Fv/Fm) for individual *N. benthamiana* plants. In **B** and **D** the horizontal lines represent the median and boxes indicate the 25th and 75th percentiles. One-way analysis of variance (ANOVA) was used to determine statistical difference between control (*B. cinerea* alone), mock (water alone), and peptide-treated inoculated plants (GraphPad Prism-version 9.4.1). Each technical replication included four plants (n = 4). Three biological replications were performed. Asterisks represent significant differences between different groups with *p* values ($^{ns}p > 0.05$, $*p \leq 0.05$ $**p < 0.01$, $****p \leq 0.0001$) determined using Tukey's honestly significant difference test for comparison of multiple groups.

1002　**Figure 6.**

1003

Fig 6. **GMA4CG_V6 exhibits curative antifungal activity against gray mold in tomato plants.**
A. Three-week-old tomato plants were spray-inoculated with 1 mL of 8×10^4 *B. cinerea* spore
suspension. At 12 hpi, plants were sprayed with 2 ml water or 2 ml peptide (GMA4CG_WT or
GMA4CG_V6) at concentrations of 6, 12, or 24 µM. The CropReporter images of the plants were
taken 72 hpi and the maximum quantum yield of PSII photochemistry (Fv/Fm) was determined.
Red color represents class I, low efficiency photosynthesis attributed to extensive tissue damage.
Green color represents class V, high efficiency photosynthesis. **B.** The calculated photosynthetic
quantum yield (Fv/Fm) for individual tomato plants. The horizontal lines represent the median and
boxes indicate the 25^{th} and 75^{th} percentiles. One-way analysis of variance (ANOVA) was used to
determine statistical difference between control (*B. cinerea* alone), mock (water alone), and

peptide-treated inoculated plants (GraphPad Prism-version 9.4.1). Each technical replication included four plants (n= 4). Three biological replications were performed. Asterisks represent significant differences between different groups with *p* values ($^{ns}p>$ 0.05, *$p\leq$ 0.05 **$p<$ 0.01, ****$p\leq$ 0.0001) determined using Tukey's honestly significant difference test for comparison of multiple groups. **C.** Germination and growth of *B. cinerea* spores on the tomato leaf surfaces in the presence of GMA4CG_V6 or water. Tomato plants were inoculated with *B. cinerea* spores. Infection was allowed to progress for 12 h and then the plants were sprayed with water or with GMA4CG_V6. Forty-eight hours post peptide or water treatment, leaf samples were fixed in 4% paraformaldehyde and stained with Calcofluor White for 30 min. ZEISS Axio ZoomV16 was used to capture images. Arrowheads point to the germinated hyphae of *B. cinerea*.

Figure 7.

Fig 7. A proposed model for the modes of antifungal action of GMA4CG_V6 against *B. cinerea*. GMA4CG_V6 sprayed on *B. cinerea* infected leaves first binds predominantly to the

fungal plasma membrane with little or no binding to the cell wall. Binding to plasma membrane-resident phosphoinositides presumably plays a role in membrane permeabilization. At low concentration, GMA4CG_V6 is internalized into cell vacuoles. It remains to be determined if this peptide is internalized into vacuoles through endocytic mechanisms. At high concentration, GMA4CG_V6 targets the cytoplasm immediately following cell entry. Cell death occurs only after GMA4CG_V6 diffuses into the cytoplasm and not while in vacuoles. N: nucleus, V: vacuole, E: endosome.

1062 **Supporting information**

1063 **S1_Fig**

1064

1065

1066 **S1_Fig. GMA4CG_V6 exhibits fungicidal activity.** (**A**) Transmitted light images of the
1067 germination and growth of *B. cinerea* spores treated with 1.5 μM GMA4CG_V6 for 1, 6, and 24

h. After each treatment, unbound peptide was removed, and the spores allowed to germinate overnight. Scale bar = 50 µm. (**B**) Quantification of the germination rate of GMA4CG_V6 treated *B. cinerea* spores. Spores treated with sterile water served as controls. Data were collected from four independent replications (n = 4) and plots are shown in box plot. Individual data were overlaid on plots as red points. Statistical analysis was performed by two-sided Student's t test. (**C**) *B. cinerea* spores were treated with 1.5 µM GMA4CG_V6 and the dead cell staining dye PI for various times. The overlaid images are shown for 0.25, 0.5, 1, and 2 h treatments. Scale bar = 20 µm. (**D**) Quantitative analysis of the percentage of spores stained with PI dye. The top of the columns represents the mean, and the horizontal bars represent the standard deviation (n = 3).

S2_Fig. GMA4CG_V6 retains antifungal activity in the presence of salts. (A) Antifungal
activity of the GMA4CG variants and MtDef4 against *B. cinerea* was determined in 1 x SFM in
the absence or presence of 100 mM NaCl, 100 mM NaCl, or 2 mM CaCl$_2$. After 48 h incubation,
resazurin dye was added to each well to determine cell viability/cell death. A change from blue to
pink/colorless indicated viable fungal cells. **(B)** Microscopic images of *B. cinerea* spores
challenged with various concentrations of GMA4CG_V6 in the presence of different salts (Scale
bar = 50 μm).

1104

1105 **S3_Fig**

1106

1107 **S3_Fig.** Steady-state wavelength CD spectra for GMA4CG_WT (red) and GMA4CG_V6 (blue)
1108 in 20 mM sodium acetate, pH 5.6, at 20 °C.

1109

1110

S4_Fig

A

Peptide	Sequence	SFM MIC (µM)	SFM+100 mM NaCl MIC (µM)
GMA4CG_V6	GGRCKGFRRRWFWTRIC	1.5	1.5
GMA4CG_V6-Ala1	AAAAKGFRRRWFWTRIC	6	6
GMA4CG_V6-Ala2	GGRCAAARRRWFWTRIC	6	6
GMA4CG_V6-Ala3	GGRCKGFAAAAFWTRIC	12	24
GMA4CG_V6-Ala4	GGRCKGFRRRWAAARIC	6	6
GMA4CG_V6-Ala5	GGRCKGFRRRWFWTAAA	6	6

S4_Fig. Antifungal activity of the alanine scanning variants of linear GMA4CG_V6 as
determined by the resazurin cell viability/cell death assay. (A) The primary amino acid
sequences for GMA4CG_V6 and the five Ala scanning mutants and their associated MIC values
in the absence and presence of salt. The GMA4CG_V6 substitutions are highlighted red. **(B)**
Resazurin dye was used to determine cell viability/cell death. A change from blue to pink/colorless
indicated viable fungal cells. Antifungal activity of each variant was performed in 1 x SFM. There
was an eight-fold loss in activity with GMA4CG_V6_Ala3 whereas all other variants showed a
four-fold loss. **(C)** Resazurin cell viability/cell death assay showing the MIC values of alanine
scanning variants of GMA4CG_V6 in the presence of 100 mM NaCl. MIC values of all other
variants remained unchanged except for GMA4CG_V6_Ala3, which increased another two-fold.

1124

1125

1126 **S5_Fig**

1127

1128

S5_Fig. Internalization of TMR-GMA4CG_V6 into *B. cinerea* germlings at sublethal or very
high concentrations. (A) Confocal microscopy images show internalization of 0.5 µM TMR-
GMA4CG_V6. Images were captured 60 min after peptide challenge. **(B)** Time-lapse confocal
microscopy images of *B. cinerea* show internalization of 20 µM TMR-GMA4CG_V6 over time
(T in min) with images captured every 3 min shown. Scale bars = 10 µm.

S6_ Fig

1135

1136

1137

S6_Fig. Co-localization of TMR-GMA4CG_V6 and CFW. (A & C) Representative fluorescent
(CFW and TMR) and channel merged images of *B. cinerea* germlings treated with TMR-
GMA4CG_V6 and captured by a confocal microscope (A) and the overlay image by lattice SIM[2]
super resolution microscope (C). Scale bar is 10 µm in A and 5 µm in C. **(B & E)** The fluorescent
signal intensity of the ROIs in **A** and **C**, respectively. The dash lines in **E** indicate the shift between
the peak of the CFW and TMR signals. **(D)** Enlarged image from the framed area in C. Scale bar
= 2 µm.

1145

1146

1147

1148

1149

1150

1151

1152

1153 **S7_Fig**

1154

1155

S7_Fig. FRAP analysis figure

1156

S7_Fig. FRAP analysis of the subcellular localization of GMA4CG_V6 in *B. cinerea*
germlings and conidia. (A) FRAP plots show the normalized fluorescence recovery of TMR-
GMA4CG_V6 in germlings cytoplasm (blue), cell walls/ periphery (orange), and vacuoles (gray)
and in conidia cell walls/periphery (green) over 120 s. A one-phase association parameter was
used to simulate the fitting curve (solid lines) based on the scatter diagram (GraphPad Prism 8.0).
Higher recovery rates indicate greater overall mobility. **(B)** The immobile fraction (%) of
GMA4CG_V6 localized into the key subcellular compartments of *B. cinerea*. Data are represented
using box plots with individual data overlaid as red points. Data analyzed for each subcellular
compartment of *B. cinerea* are from n = 20 collections from at least three independent experiments.

Asterisks represent significant differences between different subcellular compartments. One-way
analysis of variance (ANOVA) was used to determine the statistical difference with Tukey's
honestly significant difference test (HSD) for comparison of multiple groups. **(C)** A table showing
the half-time of recovery (Tau) of GMA4CG_V6 localized into the key subcellular compartments
of *B. cinerea*. The Tau and R^2 values for each peptide were calculated using a one-phase
association parameter.

1172

1173 **S8_Fig**

1174

1175

S8_Fig. GMA4CG_V6 binding to phospholipids. (A) Lipid overlay assays of GMA4CG_WT,
GMA4CG_V6, and Hexa-His. Both GMA4CG_WT and GMA4CG_V6 show strong binding to
phosphatidylinositol monophosphates PI(3)P, PI(4)P, and PI(5)P while GMA4CG_WT shows
strong binding to additional phospholipids including PA and the phosphatidylinositol diphosphates
PI(3,5)P_2 and PI(4,5)P_2. In contrast, GMA4CG_V6 binds to multiple phospholipids, including
phosphatidylinositol di/triphosphates PI(3,4)P2, PI(3,4,5)P3, and PA, but rather weakly. **(B & C)**
Integrated density (grayscale pixel units) analysis of PIP strip probed with GMA4CG_WT and
GMA4CG_V6. Data are means ± SEM of two independent biological replicates (n = 2).

1184

1185

 S9_ Fig

S9_ Fig. GMA4CG_V6 has preventative and curative antifungal activity against gray mold in *N. benthamiana* plants (A) Preventative activity of GMA4CG_V6. Four-week-old *N. benthamiana* plants were sprayed with either 2 ml water or peptide (GM4CG_WT or GMA4CG_V6) at 12 or 24 µM concentration prior to spray-inoculation with a 1 ml suspension of 8×10^4 *B. cinerea* fungal spores. Plants were observed for symptoms at 72 hpi and photographed in white light. **(B)** Disease severity of tobacco plants inoculated with *B. cinerea* at 72 hpi. The disease severity was rated on a scale of 0 to 4 with 0 = no symptoms; 1 = 1-25%; 2 = 26-50%; 3 = 51–75%; and 4 = 75-100% of the leaf surface infected. One-way analysis of variance (ANOVA)

was used to determine statistical differences between control (*B. cinerea*), mock-treated, and
peptide-treated plants (GraphPad Prism-version 9.4.1). Four *N. benthamiana* plants (n = 4) for
each technical replication were tested in three biological replications. Asterisks represent
significant differences between different groups with P values determined ($^{ns}p>$ 0.05, *$p\leq$ 0.05
**$p<$ 0.01, ****$p\leq$ 0.0001) using Tukey's honestly significant difference test for comparison of
multiple groups. **(C)** Curative antifungal activity of GMA4CG_V6. Four-week-old *N.*
benthamiana plants were spray-inoculated with a 1 mL suspension of 8×10^4 of *B. cinerea* fungal
spores at 2, 15, or 30 h and then sprayed with 2 mL of 12 μM GMA4CG_WT, 12 μM
GMA4CG_V6, or water. Plants were photographed in white light after 72 h. **(D)** Disease severity
was rated as described above. Four *N. benthamiana* plants (n = 4) for each technical replication
were tested in three biological replications. Asterisks represent significant differences between
different groups with *p* values determined ($^{ns}p>$ 0.05, *$p\leq$ 0.05 **$p<$ 0.01, ****$p\leq$ 0.0001) using
Tukey's honestly significant difference test for comparison of multiple groups.

1209

1210 **S10_Fig**

1211

1212

1213 **S10_ Fig. GMA4CG_V6 has curative antifungal activity against gray mold in tomato plants.**
1214 **(A)** Three-week-old tomato (*Solanum lycopersicum* L cv. Mountain Spring) plants were spray-
1215 inoculated with a 1 ml suspension of 8×10^4 fungal spores and subsequently sprayed 12 h later
1216 with 2 ml of GMA4CG_WT or GMA4CG_V6 at a concentration of 6, 12, or 24 μM or with 2 ml
1217 of water (controls). Plants were scored for gray mold symptoms at 72 hpi and photographed in

white light. **(B)** Disease severity of tomato plants inoculated with *B. cinerea* 72 hpi. The disease severity was rated on a scale of 0 to 4 with 0 = no symptoms; 1 = 1-25%; 2 = 26-50%; 3 = 51–75%; and 4 = 75-100% of the leaf surface infected. One-way analysis of variance (ANOVA) was used to determine statistical differences between control (*B. cinerea*), mock, and peptide experiments (GraphPad Prism-version 9.4.1). Four tomato plants (n = 4) for each technical replication were tested in three biological replications. Asterisks represent significant differences between different groups with *p* values ($^{ns}p > 0.05$, $*p \leq 0.05$ $**p < 0.01$, $****p \leq 0.0001$) determined using Tukey's honestly significant difference test for comparison of multiple groups.

S11_ Fig

S11_Fig. Binding of TMR-GMA4CG_WT and TMR-GMA4CG_V6 to tomato leaf surfaces.
(A) Representative confocal microscopy images of tomato leaves with no peptide applied showing the absence of TMR fluorescence signal, background cell wall autofluorescence (cyan), and

transmitted light. Scale bar = 20 µm. **(B)** Confocal microscopy images of plant cell walls stained with TMR-GMA4CG_WT and TMR-GMA4CG_V6 showing no signal observation in the cytoplasm. Scale bar = 20 µm. **(C)** Comparison of TMR signal intensity at the tomato cell wall before and after washing with sterile water. The average signal from leaves treated with each peptide before washing served as the control (100%). Each dot indicates individual data points. Asterisks represent significant differences between different groups (unpaired *t* test, n = 26-32) (GraphPad Prism 8.0).