Title: Plant defensin MtDef4-derived antifungal peptide with multiple modes of action and

potential as a bioinspired fungicide

Authors: Meenakshi Tetorya[a*], Hui Li[a*], Arnaud Thierry Djami-Tchatchou[a], Garry W.

Buchko[c,d], Kirk J. Czymmek[a,b], Dilip M. Shah[a,1]

Author Affiliations:

[a]Donald Danforth Plant Science Center, St Louis, MO. 63132, USA.

[b]Advanced Bioimaging Laboratory, Donald Danforth Plant Science Center, St Louis, MO

63132, USA

[c]Earth and Biological Sciences Directorate, Pacific Northwest National Laboratory, Richland,

WA 99354, USA

[d]School of Molecular Biosciences, Washington State University, Pullman, WA 99164, USA

**Corresponding Author*:*

[1]Correspondence and requests for materials should be addressed to D.M.S. (Email:

dshah@danforthcenter.org)

Author contributions

M.T., H.L, K.C. and D.S. conceived the research. M.T., H.L., A.D., K.C., G.B. and D.S. designed

the research. M.T., H.L., A.D., G.B. and K.C. conducted the experiments. M.T., H.L., A.D., K.C.,

G.B., and D.S. analyzed and interpreted the results. M.T., H.L, G.B., K.C. and D.S. wrote the

manuscript.

*These authors made equal contribution to the manuscript.

Competing Interest Statement

M.T. and D.S. are listed as inventors in a related patent application. The remaining authors declare

no competing interests. M.T., H.L., K.C. and D.S. are affiliated with Peptyde Bio Company.

Classification: Biological Sciences/Plant Biology

Keywords:

Defensin-derived peptides, antifungal activity, modes of action, *Botrytis cinerea*, gray mold

disease, bioinspired fungicides.

Abstract

Chemical fungicides have been instrumental in protecting crops from fungal diseases. However,

mounting fungal resistance to many of the single-site chemical fungicides calls for the

development of new antifungal agents with novel modes of action (MoA). The sequence-divergent

cysteine-rich antifungal defensins with multi-site MoA are promising starting templates for design

of novel peptide-based fungicides. Here, we experimentally tested such a set of 17-amino acid

peptides containing the γ-core motif of the antifungal plant defensin MtDef4. These designed

peptides exhibited antifungal properties different from those of MtDef4. Focused analysis of a lead

peptide, GMA4CG_V6, showed it was a random coil in solution with little or no secondary

structure elements. Additionally, it exhibited potent cation-tolerant antifungal activity against the

plant fungal pathogen *Botrytis cinerea*, causal agent of gray mold disease in fruits and vegetables.

Its multi-site MoA involved localization predominantly to the plasma membrane, permeabilization

of the plasma membrane, rapid internalization into the vacuole and cytoplasm, and affinity for

bioactive phosphoinositides phosphatidylinositol 3-phosphate (PI3P), PI4P, and PI5P. The

sequence motif RRRW was identified as a major determinant of the antifungal activity of this

peptide. While topical spray-application of GMA4CG_V6 on *Nicotiana benthamiana* and tomato

plants provided preventative and curative suppression of gray mold disease symptoms, the peptide

was not internalized into plant cells. Our findings open the possibility that truncated and modified

defensin-derived peptides containing the γ-core sequence could serve as promising candidates for

further development as bioinspired fungicides.

Significance

The rapid evolution of fungal pathogen resistance to single-site chemical fungicides in agriculture

has raised an urgent need for development of safe, sustainable, and cost-effective multi-site

fungicides. Short peptides containing the functionally active γ-core motif of antifungal defensins

hold strong potential for development as efficacious, environmentally friendly biofungicides

replacing or complementing chemical fungicides. A 17-amino acid pseudo-cyclic MtDef4 plant

defensin-derived peptide exhibits potent antifungal activity and deploys multi-target modes of

action against the gray mold fungal pathogen *B. cinerea*. Spray-application of this peptide confers

strong preventative and curative antifungal activity against this pathogen in *Nicotiana*

benthamiana and tomato plants. This work provides evidence for the enormous potential of

defensin-derived peptides as bioinspired fungicides in agriculture.

Crop plants are under constant threat from fungal and oomycete pathogens that cause significant

pre- and post-harvest economic losses worldwide and challenge food security globally (1). This

threat is increasing in severity and scale (2, 3). It is estimated that fungal and oomycete pathogens

cause up to 10-23% yield losses in calorie crops (3) and postharvest losses of 10% (4). Modern

intensive agriculture depends heavily on chemical fungicides for management of fungal and

oomycete diseases in commodity crops as well as high value vegetable and fruit crops (5). Most

chemical fungicides in use today have a single biochemical target and thus are prone to resistance

development by diverse and rapidly evolving fungal and oomycete strains (6). Facing fungal

resistance and unwanted side effects, some fungicides are losing their effectiveness in modern

intensive monoculture cropping systems. The structurally diverse plant antifungal peptides with

potent antifungal activity could potentially serve as structural templates for the design of a new

generation of safe and sustainable fungicides (7, 8).

Plants use a variety of different mechanisms to resist fungal and oomycete infections (9).

Anti-microbial peptides (AMPs) are ubiquitous throughout the plant kingdom and play important

roles in plant defense against fungal and oomycete pathogens. Their potent anti-fungal and -

oomycete activity are vital components of the plant innate immune system (8, 10). Defensins, in

particular, are one of the largest families of AMPs expressed in plants. Because of their potent

antifungal activity, diverse amino acid sequences, and multi-target MoA, plant defensins have

emerged as potential antifungal agents for agriculture (11-13). Plant defensins are 45-54 amino

acids long, cysteine-rich, cationic peptides characterized by the presence of a cysteine-stabilized

αβ motif. This motif consists of one α-helix and three antiparallel β-strands linked by three

disulfide bonds. The fourth disulfide bond connects near the N- and C-termini and essentially

cyclizes the peptide (14). The cationic plant defensins exhibit potent broad-spectrum antifungal

activity against fungal pathogens *in vitro* and *in planta* (13-16). We set out to investigate if truncated plant defensins containing the active carboxy-terminal γ-core motif (GXCX$_{3-9}$C) exhibited potent fungicidal activity and were effective as spray-on fungicides for management of fungal diseases. We further investigated if the smaller defensin-derived peptides exhibited MoA different from those of the full-length parental peptide.

MtDef4 is an evolutionarily conserved 47-amino acid defensin from *Medicago truncatula* that exhibits broad-spectrum antifungal activity against fungal pathogens at low micromolar concentrations. MtDef4 MoA against the plant pathogen *Fusarium graminearum* involves plasma membrane permeabilization, binding to bioactive phosphatidic acid (PA), cell entry, and presumed interaction with intracellular targets (17-19). Like many other plant defensins, this peptide contains a well-characterized γ-core motif required for its antifungal activity (18, 19). A chemically synthesized C-terminal 16-amino acid sequence, designated GMA4C, containing this motif and a net charge of +6 inhibited the growth of the fungal pathogen *Fusarium graminearum in vitro* although its potency was four-fold lower than that of MtDef4 (18). This peptide also exhibited antifungal activity against crown rot fungal pathogens of alfalfa and, in addition, inhibited the growth of plant and human bacterial pathogens *in vitro* (20).

Herein, we explored the potential of the synthetic, MtDef4-derived, structurally less complex, 17-amino acid GMA4CG variants for development as fungicides. We characterized the antifungal activity of several GMA4CG variants generated via amino acid substitutions, C-terminal amidation, and disulfide cyclization that exhibited potent antifungal activity *in vitro* against an aggressive necrotrophic fungal pathogen *Botrytis cinerea*, the causal agent of a gray mold disease in vegetables, fruits, and flowers. In sharp contrast to MtDef4, an interesting characteristic of the GMA4CG variants was that they retained potent antifungal activity in media

with elevated salt concentrations. Based on its superior *in vitro* and semi-*in planta* antifungal

activity against *B. cinerea*, we focused on GMA4CG_V6 as the lead peptide for *in planta*

antifungal activity and MoA studies. GMA4CG_V6 exhibited multi-faceted MoA and rapidly

killed fungal cells. When sprayed on *Nicotiana benthamiana* and tomato plants, GMA4CG_V6

conferred strong preventative and curative antifungal activity against *B. cinerea* under the tested

conditions.

Results

Design of short MtDef4-derived synthetic peptide variants

We based our work on the premise that a small MtDef4-derived peptide spanning the active γ-core

motif could serve as a promising template for the design of peptide-based fungicides. We further

hypothesized that a small antifungal peptide would be more cell permeant and that with targeted

amino acid substitutions of the γ-core motif, we could further enhance its antifungal potency. To

test this hypothesis, we generated a truncated set of linear and cyclic variants of the 17-amino acid

peptide spanning this motif (Gly30-Cys47s designated GMA4CG_WT and tested them for

antifungal activity against *B. cinerea.*

The substitution of each amino acid with all possible alternative amino acids in

GMA4CG_WT was not practical. Thus, our strategy was to substitute cationic or hydrophobic

residues in specific positions with the expectation that they would have significant influence on

the antifungal activity of this peptide. MtDef4 is an evolutionarily conserved defensin in the plant

kingdom and sequences of several homologs from various plants have been reported (19).

GMA4CG_V1 is a natural variant derived from the MtDef4 homolog (KHN09675.1) of *Glycine*

soja. This natural variant has three amino acid differences (R5K, T15R and H16I) relative to the

sequence of GMA4CG_WT. GMA4CG_V2 was generated to determine if the substitution of one cationic amino acid with a different cationic amino acid had a notable impact on its antifungal activity. Thus, this variant carried a Lys5 to Arg substitution. We also designed peptide variants (GMA4CG_V3 to GMA4CG_V6) to test the influence of increased hydrophobicity on their antifungal activity. Thus, each of these variants had 29.4% hydrophobic amino acids as compared with 11.8% in GMA4CG_WT. GMA4CG_WT has four cysteine residues (Cys4, Cys11, Cys13 and Cys17) in its sequence, suggesting a potential to generate a pseudo-cyclic structure via disulfide bond formation between Cys4 and Cys17. Cyclization has been reported previously to increase the antimicrobial activity and selectivity of AMPs (21, 22). Thus, we had cyclic GMA4CG_V3 and GMA4CG_V6 synthesized with a Cys4-Cys17 disulfide bond and replaced Cys11 and Cys13 with either an aliphatic Val (GMA4CG_V3) or an aromatic Trp (GMA4CG_V6) residue. Trp is more effective in interacting with the interfacial regions of lipid bilayers and it promotes antimicrobial activity of peptides (23-25). All peptides were amidated at the C-terminus, a process known to increase the stability of peptides against proteolytic degradation, and therefore, also its biological activity (26).

GMA4CG_V4 and GMA4CG_V6 variants have potent antifungal activity

We assessed the antifungal activity of GMA4CG_WT and its six variants in Synthetic Fungal Medium (SFM) against *B. cinerea in vitro.* GMA4CG_WT inhibited the growth of *B.cinerea* with an MIC value of 3 µM, four-fold greater than that of MtDef4 (Fig. 1*B*). Among all variants, only GMA4CG_V6 displayed two-fold greater antifungal activity than GMA4CG_WT with an MIC value of 1.5 µM. GMA4CG_V6 also inhibited the growth of *F. graminearum* with a MIC value of 1.5-3 µM (Fig. 1*B*).

To determine if GMA4CG_V6 had fungicidal and/or fungistatic activity, *B. cinerea*

conidia were treated with 1.5 µM peptide for 1, 6, and 24 h. Following each treatment, the peptide

was removed, and spore germination examined under the microscope. Approximately 85% of the

conidia failed to germinate after 1 h of peptide challenge, nearly 100% of the conidia failed to

germinate after 6 h, and 100% of the conidia failed to germinate after 24 h of treatment (*SI*

Appendix Fig S1 *A* and *B*). To assess further the toxicity of GMA4CG_V6, we stained peptide-

treated conidial cells with propidium iodide, a cell death marker, and observed that nearly 80% of

the conidia stained red within 2 h of peptide treatment indicating fungicidal activity for this peptide

(*SI Appendix* Fig. S1 *C* and *D*).

The GMA4CG variants, but not MtDef4, retain antifungal activity in the presence of salts

Many naturally occurring AMPs, including plant defensins, are known to lose their antifungal

activity in media containing elevated salt concentrations (27, 28). We determined the MIC values

of GMA4CG_WT and its variants with the R/KGFRRR motif in SFM containing 100 mM NaCl,

100 mM KCl, or 2 mM $CaCl_2$ and compared them to MtDef4. At the greatest concentration tested

(6 µM), MtDef4 had no activity against *B. cinerea*. However, GMA4CG_WT and its variants

retained their antifungal activity in salt supplemented SFM to varying degrees with GMA4CG_V4

and GMA4CG_V6 retaining the most potent antifungal activity (Fig. 1*B* and *SI Appendix* Fig. S2

A). GMA4CG_V6 at 3 µM completely inhibited spore germination in the presence of NaCl and

$CaCl_2$ salts (*SI Appendix* Fig. S2 *B*). GMA4CG_V6 was notably more effective than

GMA4CG_V3 in the presence of 100 mM KCl and 2 mM $CaCl_2$ (Fig.1*B* and *SI Appendix* Fig. S2

A) indicating that the Val to Trp substitution is important for salt-tolerant antifungal activity in

these variants. Collectively, these data revealed that the MtDef4-derived GMA4CG_WT peptide

and its variants with a net charge of +5 to +6 exhibited salt-tolerant antifungal activity. In contrast,

the parental MtDef4 exhibited salt-sensitive antifungal activity at the same concentrations.

GMA4CG_WT and GMA4CG_V6 are predominantly random coil structures

The three-dimensional structure of full length MtDef4 contains a γ-core motif composed of β-

strands β2 and β3 connected by a positively charged RGFRRR loop (18). Near-UV CD spectra

for GMA4CG_WT and GMA4CG_V6 are shown in *SI Appendix* Fig. S3. Neither spectrum shows

features characteristic of a peptide with substantial α-helical (double minima at approximately 222

and 208 nm and an extrapolated maximum around 195 nm) or β-strand (single minimum at around

220 nm and an extrapolated maximum around 195 nm) secondary structure (30, 31). The notable

difference between the CD spectra of the two peptides was an additional minor minimum at ~230

nm for GMA4CG_WT and a minor maximum at ~ 230 nm for GMA4CG_V6. In combination

with the major sub-220 nm minimum, the former feature is characteristic of a random

coil/unordered structure and the latter feature characteristic of a polyproline II structure suggesting

the multiple amino acid substitutions may induce a transition of some random coil to polyproline

II structure in GMA4CG_V6 (32). Our attempts to predict the structures for GMA4CG and

GMA4CG_V6 using the artificial intelligence-based protein structure prediction software

AlphaFold (33) yielded structures with low confidence. Similarly, *ab initio* structure prediction

using the Rosetta protein structure modeling package (34) did not predict any consistent fold or

high-confidence secondary structure elements (data not shown). Hence, the *in silico* modeling data

corroborate that experimental CD data, suggesting the peptides have little long-range order and

are likely dynamic and disordered in solution.

The RRRW motif is a major determinant of the salt-tolerant antifungal activity of GMA4CG_V6 with no disulfide bond

A series of linear alanine-substituted variants of our lead peptide GMA4CG_V6 (*SI Appendix* Fig. S4 *A*) were tested to identify amino acids important for antifungal activity. The GMA4CG_V6_Ala3 variant carrying the substitution of RRRW with AAAA lost antifungal activity eight-fold, whereas all other variants lost antifungal activity four-fold (Table 1, *SI Appendix* Fig S4 *B*). In SFM supplemented with 100 mM NaCl, no further loss in the antifungal activity of the variants occurred except in GMA4CG_V6_Ala3, which lost antifungal activity another two-fold (*SI Appendix* Fig S4 *C*). These results indicated that the RRRW motif is a major determinant of the antifungal activity of GMA4CG_V6 in media supplemented with salts.

The GMA4CG_V4 and GMA4CG_V6 variants potently reduce gray mold disease symptoms semi-*in planta*

We compared the semi-*in planta* antifungal activity of GMA4CG_WT and its variants using detached leaves of *N. benthamiana* as described (28). GMA4CG_WT and variants GMA4CG_V1 through GMA4CG_V5 were applied as drops of different concentrations on each leaf with freshly prepared conidial inoculum immediately applied to each drop. Leaves were assessed for attenuation of gray mold symptoms by each peptide 48 h post-inoculation (hpi) relative to no peptide controls by measuring lesion sizes and photosynthetic efficiency (as measured by Fv (variable fluorescence)/Fm (maximal fluorescence)). Increasing concentration of each peptide significantly ($p \leq 0.0001$) reduced lesion sizes (Fig. 2*A* and *B*). For all peptides, no lesions were formed at 6 µM. At 3 µM, only GMA4CG_V3 and GMA4CG_V4 were effective in suppressing lesion formation. At 1.5 µM, only GMA4CG_V4 completely suppressed disease symptoms. The

photosynthetic efficiency data further supported GMA4CG_V4 as being more potent than the

other five in reducing gray mold lesions (Fig 2*A*). We then compared the semi-*in planta* antifungal

activity of GMA4CG_V4 and GMA4CG_V6. Both peptides completely suppressed disease

symptoms at 1.5 and 3 µM under the applied conditions. However, at a lower concentration of

0.75 µM, GMA4CG_V6 was more effective in suppressing gray mold disease symptoms than

GMA4CG_V4 (Fig. 2*C* and *D*).

GMA4CG_V6 enhances permeability of the plasma membrane of *B. cinerea* spores and

germlings

To determine if GMA4CG_V6 permeabilizes the fungal plasma membrane, the spores and

germlings of *B. cinerea* were treated with 1.5 µM GMA4CG_V6 for 30 min followed by the

addition of SYTOX^{TM} Green (SG) Nucleic Acid Stain. SG is a membrane-impermeant dye, which

fluoresces when it binds to RNA and DNA and is a useful indicator of membrane permeability.

SG uptake was observed in nearly all fungal spores and germlings challenged with this peptide,

but not in controls without peptide (Fig. 3*A*). Thus, we concluded that one MoA of GMA4CG_V6

involved increased plasma membrane permeabilization.

Since antifungal activity of GMA4CG_V6 was cation-tolerant, we determined if

membrane permeability still occurred in the presence of 100 mM KCl. In the absence of this salt,

95% of the conidia exposed to 1.5 µM GMA4CG_V6 for 1 h took up SG, indicating increased

membrane permeability. No further time- or concentration dependent increase in membrane

disruption was observed in the absence of this salt (Fig. 3*B*). By contrast, uptake of SG occurred

in less than 10% of the conidia treated with 1.5 µM GMA4CG_V6 for 1 h in the presence of 100

mM KCl. Even after 5 h, only ~17% of the conidia took up SG. A high concentration of 6 µM

GMA4CG_V6 was needed for the permeabilization of ~80% of conidia after 5 h (Fig. 3*B*). Thus,

the membrane-permeabilization ability of GMA4CG_V6 was negatively impacted by the presence

of cations; however, the peptide was still effective in permeabilizing the fungal plasma membrane

at increased concentrations with longer exposure.

We also used our SG assay to compare the ability of GMA4CG_V6, GMA4CG_WT, and

MtDef4 to permeabilize the plasma membrane in germlings in the absence and presence of 100

mM KCl. In the absence of KCl, MtDef4 was more effective in permeabilizing the plasma

membrane than GMA4CG_WT or GMA4CG_V6. However, in the presence of this salt, MtDef4's

ability to permeabilize the plasma membrane was completely lost at low concentrations ($< 3\,\mu$M).

In contrast, GMA4CG_WT and GMA4CG_V6 retained their ability to permeabilize the plasma

membrane at $< 3\,\mu$M (Fig. 3*C*).

GMA4CG_V6 is internalized into the vacuole and cytoplasm of *B. cinerea* in a time and

concentration dependent manner

Tetramethyl rhodamine (TMR)-labeled GMA4CG_V6 was imaged using time-lapse confocal

microscopy to gain insight into the temporal nature of peptide internalization in *B. cinerea*

germlings. At an MIC of 1.5 μM, the labeled peptide first bound to the cell surface within the first

3 min and started translocating into vacuoles from multiple entry foci by 19 min (Fig. 4*A*). At this

time point, bright field images of germlings showed abnormal morphology with vacuoles

becoming visible. At 27 min, the vacuoles began to expand by coalescence of small vacuoles (Fig.

4A, arrows) and the peptide localized into the cytoplasm. Shortly after the peptide's appearance in

the cytoplasm the vacuoles appeared to lyse with the peptide rapidly dispersing throughout the

germling cytoplasm and inducing cell death (Fig. 4*A*, T= 59).

At a sub-lethal concentration of 0.5 µM, the labeled peptide was localized at the germling

periphery and formed presumptive endosomes in the cytoplasm. No further progression of the

peptide in the germling cells occurred during the 1 h challenge (*SI Appendix* Fig. S5 *A*). At high

concentration (20 µM), TMR-labeled GMA4CG_V6 was internalized by germlings as early as 4

min. The peptide rapidly accumulated into the germling cytoplasm and within 13 min was

completely dispersed throughout the cell's interior (*SI Appendix* Fig. S5 *B*). This suggests that at

high concentrations, GMA4CG_V6 may largely bypass intermediate endosome or vacuole

localization and accumulate directly in the cytoplasm, or highly accelerate the translocation from

vacuole to cytoplasm and finally accelerate cell death.

The subcellular localization of TMR-GMA4CG_V6 was determined using two dyes which

specifically stain the plasma and endosomal membranes (FM4-64) and vacuole lumen (CMAC).

Within 30 min of challenging germlings with peptide, TMR-GMA4CG_V6 co-localized with

CMAC in vacuoles and with FM4-64 at the plasma membrane and entry foci (Fig. 4*B*, highlighted

white). Based on the region of interest (ROI01) in Fig. 4*B*, the fluorescence intensity curve clearly

showed that the peak signal of TMR-GMA4CG_V6 overlapped with the signal of FM4-64 in

membrane and CMAC in vacuole (Fig. 4*C*). To determine whether GMA4CG_V6 bound to the

cell wall of *B. cinerea*, a cell wall specific dye Calcofluor White (CFW) was incubated with TMR-

GMA4CG_V6 treated *B. cinerea* germlings. As illustrated in *SI Appendix* Fig. S6 *A* and *B*, CFW

overlapped with TMR fluorescent signal in images captured by confocal microscopy. Considering

the thickness and proximity of germling cell wall and plasma membrane and the limitation of the

resolution of the confocal microscopy, super-resolution microscopy (specifically, lattice structured

illumination microscopy (SIM) was used to further evaluate co-localization of TMR-

GMA4CG_V6 with the cell wall. The fluorescent signal of CFW relative to TMR showed the

TMR signal predominantly just inside the cell wall (*SI Appendix* Fig. S6 *C*, *D* and *E*), as shown by

arrows, indicating that dominant TMR-GMA4CG_V6 signal was consistent with intense plasma

membrane localization at the cell periphery, although low level cell wall binding could not be ruled

out.

To further study the binding affinity of GMA4CG_V6 to specific subcellular components

in *B. cinerea* germlings, a series of Fluorescence Recovery After Photobleaching (FRAP)

experiments were performed (*SI Appendix* Fig. S7 *A-C*). We specifically studied the mobility of

TMR-GMA4CG_V6 to subcellular components including germling cell periphery, vacuoles, and

cytoplasm. In our previous study, a significant difference in the mobility of the antifungal peptide

NCR044 to germling and conidial cell walls was observed (35), thus, we performed FRAP

experiments with conidial spore periphery as well. According to the normalized fluorescence

recovery curve, TMR-GMA4CG_V6 recovery in the germling cytoplasm had the greatest overall

mobility among all four targeted components and the mobility in vacuoles was relatively low (*SI*

Appendix Fig. S7 *A*). Immobile fraction and half time of fluorescence recovery data were also

collected as summarized in *SI Appendix* Fig. S7 *B* and *C*. GMA4CG_V6 in germling vacuoles

showed a significantly higher ($p < 0.0001$) immobile fraction ($82.6 \pm 3.2\%$) than other subcellular

components and the recovery time was much shorter (Tau= 19.27 sec, $R^2 = 0.8496$). In cell

periphery of germlings and conidia, no significant difference was observed in the immobile

fraction ($52.5 \pm 4.2\%$ and $51.0 \pm 4.9\%$, respectively, $p > 0.05$) or in the half recovery time (37.4

sec, $R^2 = 0.9133$ and 38.4 sec, $R^2 = 0.9257$, respectively). The immobile fraction of the peptide was

lower ($p = 0.0031$) in germling cytoplasm ($35.2 \pm 4.3\%$) as compared to that in germling cell

periphery and vacuole but showed no significant difference with that of the spore periphery. The

mobility of peptide in germling cytoplasm was the highest among the four selected subcellular

components (Tau= 40.68 and R^2= 0.9636). Collectively, the FRAP analysis indicated that the

mobility of the peptide at the cell periphery and cytoplasm was similar but much different in the

vacuoles. The low mobility in vacuoles is likely because transport to the vacuole lumen is not a

continuum like other compartments and multistep mobility into the vacuole itself would be very

high as the Tau values show.

Because of the difference in the cation tolerant antifungal activity of MtDef4 and

GMA4CG, we examined the localization of MtDef4, GMA4CG_WT, and GMA4CG_V6 on the

spores of *B. cinerea* in the presence of 100 mM KCl. The fluorescently labeled MtDef4 localized

to the cell periphery in the absence of this salt, but this localization was significantly reduced in

its presence (p = 0.001). In contrast, cell periphery localization of GMA4CG_WT and

GMA4CG_V6 was not affected by salt (p> 0.05) (Fig. 4*D* and *E*).

GMA4CG_WT and GMA4CG_V6 bind to multiple phosphoinositides

MtDef4 binds to phosphatidic acid (PA) in fungal membranes as part of its MoA (18). Therefore,

we tested if MtDef4-derived GMA4CG_V6 also binds to PA. A protein-lipid overlay assay was

performed with various biologically relevant phospholipids immobilized on a lipid strip (Echelon

Bioscience, UT). GMA4CG_V6 bound to several phosphoinositides including phosphatidylinstol

(PI) mono- and bisphosphates, PA, and phosphatidylserine, but with higher affinity for PI4P and

PI5P than for PA (*SI Appendix* Fig. S8). GMA4CG_WT also bound to the same phospholipids as

GMA4CG_V6. His_6, the C-terminal tag attached to each peptide used in the assay for antibody

detection, did not bind to the tested phospholipids. These results indicated that GMA4CG_V6 was

more promiscuous in its binding to the membrane resident phospholipids than MtDef4 (18).

Spray-applied GMA4CG_V6 provided preventative and curative antifungal activity against

the gray mold disease

We tested the potential of GMA4CG_WT and GMA4CG_V6 to provide preventative protection

from gray mold disease by spraying two-week-old *N. benthamiana* plants with each peptide at

concentrations of 12 and 24 µM followed 2 h later with the spray-inoculum of *B. cinerea* spores.

Each peptide spray-applied at 12 or 24 µM reduced gray mold disease symptoms compared with

control plants inoculated with pathogen only (Fig. 5*A* and *SI Appendix* Fig. S9*A*). Both peptides

significantly prevented the spread of gray mold on *N. benthamiana* plants (Fig. 5*A* and 5*B*).

Notably, plants sprayed with GMA4CG_V6 and subsequently inoculated with the pathogen looked

as symptom free as non-inoculated mock (water)-sprayed plants. This data clearly demonstrated

higher potency of GMA4CG_V6 over GMA4CG_WT in reducing gray mold disease in *N.*

benthamiana plants in a preventative manner. In addition, GMA4CG_V6 was more effective in

limiting disease severity than GMA4CG (*SI Appendix* Fig. S9 *A* and *B*).

We also tested GMA4CG_WT and GMA4CG_V6 for their ability to provide curative

protection from this disease at three different time points following fungal infection. *N.*

benthamiana plants were pre-inoculated with freshly prepared *B. cinerea* spores for 2, 15, and 30

h after which each peptide at 12 µM was sprayed onto the plants. *B. cinerea* spores germinate after

12 h (36), and thus, fungal infection of the leaf surface was already initiated at 15 and 30 hpi.

Disease symptoms were rated at 72 hpi. Both peptides provided significant curative ($p = 0.0019$)

protection from gray mold (Fig. 5*C* and *SI Appendix* Fig. S9 *C*). However, GMA4CG_V6

conferred significantly greater ($p<0.0001$) curative antifungal activity than GMA4CG_WT

sprayed on plants pre-inoculated for 15 h (Fig. 5*D*). The reduced disease severity in GMA4CG_V6

sprayed plants further confirmed greater efficacy of this peptide (*SI Appendix* Fig. S9*C*). In plants

pre-inoculated for 2 h or 30 h, both peptides provided robust control of disease symptoms with

similar efficacy (Fig. 5*D* & *SI Appendix* S9 *D*). Furthermore, our data showed the potential of

GMA4CG_V6 in reducing the spread of disease symptoms (*SI Appendix* Fig. S9 *D*).

Next, we tested the curative antifungal activity of GMA4CG_WT and GMA4CG_V6 on

tomato (cv. Mountain Spring) plants pre-inoculated for 12 h and subsequently sprayed with each

peptide at a concentration of 6, 12, and 24 µM. Both peptides conferred curative antifungal activity

at all three concentrations (Fig. 6*A*). At concentrations of 12 and 24 µM, no significant difference

was observed between the two peptides in their ability to reduce disease symptoms. Both peptides

provided robust curative protection with pathogen-challenged plants appearing similar to water-

sprayed non-inoculated mock plants. However, at a concentration of 6 µM, GMA4CG_V6

conferred significantly greater ($p < 0.0001$) curative antifungal activity than GMA4CG_WT (Fig.

6*B*).

Visual rating of disease severity clearly revealed superiority of GMA4CG_V6 in

controlling gray mold disease in tomato plants (*SI Appendix* Fig. S10 *A* and *B*). The cell wall

staining of the tomato leaves from control plants infected with the pathogen alone and from

infected plants sprayed with GMA4CG_V6 confirmed strong inhibition of fungal growth provided

by this peptide (Fig. 6*C*).

GMA4CG_V6 binds strongly to the tomato leaf surface and is not internalized into plant

cells

To determine the distribution of GMA4CG_WT and GMA4CG_V6 on leaf surfaces, fluorescently

labeled peptide was applied onto detached tomato leaves and the uptake of each peptide by the

plant cells was followed using confocal microscopy. In leaves with no applied peptide, only

autofluorescence signal from the plant cells was observed (*SI Appendix* Fig. S11 *A*). Both

GMA4CG_WT and GMA4CG_V6 bound to the exterior of plant cell walls, but no internalization

into plant cells was detected (*SI Appendix* Fig. S11 *B*). After washing the leaves with water, the

fluorescence intensity of GMA4CG_WT was significantly reduced (*p*< 0.0001) but that of

GMA4CG_V6 was only slightly reduced (*p*= 0.1146) (*SI Appendix* Fig. S11 *C*). This result

suggests that GMA4CG_V6 adheres to the tomato leaf surface more than GMA4CG_WT.

Discussion

Modern agriculture relies heavily upon systemic single-site fungicides with preventative and

curative antifungal properties. However, rapid evolution of fungal resistance to fungicides poses a

major threat to crop protection and food security worldwide. New environmentally friendly

antifungal compounds with multifaceted MoAs are needed to mitigate resistance development.

Here, inspired by the broad-spectrum antifungal activity, structure, and multi-faceted MoA of the

plant defensin MtDef4, we explored the potential of a short 17-residue peptide for development as

a peptide-based fungicide.

We designed and synthesized six 17-amino acid GMA4CG variants spanning the active γ-

core motif of a plant defensin MtDef4 (16-18). We made several modifications of the GMA4CG

peptide that maintained or enhanced its antifungal activity against an economically important

fungal pathogen *B. cinerea*. Although we could not perform complete analysis of all amino acid

substitutions, our selected changes revealed the importance of cationic and hydrophobic residues

for antifungal activity of this peptide. Amino acids we chose to replace were guided partly by the

corresponding amino acid sequences of the homologs of MtDef4 from other plants. In addition,

replacement of the internal Cys11 and Cys13 residues to the aromatic hydrophobic amino acid Trp

enhanced antifungal activity. The importance of Trp at these positions was revealed by our

observation that an aliphatic hydrophobic residue Val at these positions was less effective. It is

likely that interactions of the two Trp residues with the Arg residues present in the GMA4CG_V6

primary amino acid sequence contribute significantly to peptide stability (37). Further, Trp is

known to insert into membranes containing anionic phospholipid head groups with high efficiency,

thus contributing to the increased fungal membrane permeability by this designed peptide (38).

We synthesized GMA4CG_V6 as a cyclic peptide with a disulfide bond connecting Cys4

and Cys17 with the notion that it would be more stable, more fungal cell permeable, more readily

internalized into fungal cells, and confer higher antifungal activity than its linear counterpart.

Cyclic peptides are touted as promising antimicrobial molecules since they are more stable to

proteolytic degradation, have high affinity and selectivity for targets, and better cell membrane

permeability (39).

GMA4CG_WT and GMA4CG_V6 each have random coil/unordered structures as

determined experimentally by steady-state wavelength CD spectra showing the absence of any

significant α-helical or β-strand secondary structure content (30, 31). Both peptides are

predominately random coil in nature with a suggestion of some polyproline II structure in

GMA4CG_V6 (32).

Antifungal activity of several cationic plant defensins is abrogated in well-defined

synthetic fungal growth medium containing elevated levels of Na^+ or Ca^{2+} (27, 28, 40, 41). At the

highest concentration of 6 μM tested, MtDef4 exhibited no antifungal activity against *B. cinerea*

in medium containing elevated levels of Na^+, K^+, or Ca^{2+}. The presence of cations likely weakens

the electrostatic interactions between this cationic plant defensin and negatively charged fungal

membranes. Surprisingly, all six truncated peptide variants derived from MtDef4 exhibited

antifungal activity against *B. cinerea* in fungal growth medium containing elevated levels of Na^+,

K^+, or Ca^{2+}. Among the six variants, GMA4CG_V6 was the most potent antifungal agent. While

the molecular basis for cation-tolerant antifungal activity of these peptides is unclear, the RRRW

motif clearly plays an important role in conferring cation-tolerant antifungal activity.

The structure and MoA of GMA4CG_V6 are unlike those of any fungicide currently

commercially available. The preventative and curative plant protection properties of

GMA4CG_V6 is likely due to its ability to kill fungal pathogens rapidly. It does not have a discrete

intracellular molecular target, but instead interferes with multiple cellular processes (Fig. 7). A

major characteristic of antifungal peptides is their ability to permeabilize the plasma membrane of

fungal pathogens. GMA4CG_V6 quickly binds to the cell surface of fungal spores and germlings

and subsequently disrupts the plasma membrane of these cells near the entry points. The co-

localization of GMA4CG_V6 with the vacuolar dye CMAC revealed that this peptide, at MIC

values, localizes to the plasma membrane and vacuoles in germlings of *B. cinerea*. However, it

remains to be determined if vacuole localization is required for the antifungal activity of this

peptide, or simply a fungal detoxification step. It is of interest to note that a small rationally

designed antifungal hexapeptide PAF26 with the sequence Ac-RKKWFW-NH_2 also localizes to

vacuoles in the model fungus *Neurospora crassa* (42). Like PAF26, GMA4CG_V6 at MIC enters

vacuoles and causes their coalescence and lysis. It subsequently accumulates in the cytoplasm. At

elevated concentration, GMA4CG_V6 appeared to accumulate rapidly in the cytoplasm. MtDef4

binds to the bioactive phospholipid PA as part of its MoA (17, 18). In contrast, GMA4CG_V6

derived from this defensin binds to multiple bioactive phospholipids including PA. It preferentially

binds to the phosphoinositides PIPs in the phospholipid-protein overlay assay. Further studies are

needed to understand the role promiscuous interactions with multiple phospholipids play in the

multi-faceted MoA of GMA4CG_V6.

Current crop protection research is guided by a strong demand for sustainable fungicides

that are efficacious, less prone to resistance development, environmentally friendly, and

nonphytotoxic. In semi-*in planta* antifungal assays using detached *N. benthamiana* leaves, all six

GMA4CG analogs significantly reduced gray mold disease lesions caused by *B. cinerea* infection

with GMA4CG_V6 proving most effective. Under the conditions employed, spray application of

GMA4CG_V6 conferred strong preventative and curative antifungal activity against this disease

in young *N. benthamiana* and curative antifungal activity in tomato. These fungicidal effects were

observed readily even though the dose, application frequency, and distribution formulation have

yet to be optimized. In our studies, GMA4CG_V6 was more effective than GMA4CG_WT in

reducing disease symptoms suggesting that the amino acid substitutions introduced in

GMA4CG_V6 are important for its *in vitro* as well as *in planta* antifungal activity. Although

defensin-derived peptides offer the benefits of small size, structural simplicity, and efficiency of

peptide modifications, their commercial use as peptide fungicides will require new innovations in

peptide production at scale to be economical. In addition, efficacy of these peptides for fungal

disease control will need to be assessed in large-scale field trials and their stability to proteolytic

degradation and potential phytotoxicity under field conditions will also need to be evaluated.

Recent studies have shown the promise of antifungal peptides as topical fungicides. A

fusion of the animal AMP dermaseptin and thanatin inhibits the germination of the spores of the

obligate biotrophic fungal pathogen *Phakopsora pachyrhizi* and reduces the symptoms of Asian

rust when spray-applied to soybean leaves (43). Spray-application of the antifungal peptide

NCR044 from *M. truncatula* on tomato and *N. benthamiana* plants provided protection from the

gray mold disease (35). The present study showed that a designed defensin-derived peptide

spanning the functionally active γ-core motif with multi-faceted MoA exhibited improved

preventative and curative antifungal activity against the gray mold disease of plants. Hundreds of

sequences divergent antifungal plant defensins in the AMP database can now be exploited as

templates for the design and development of small peptide-based fungicides with diverse MoA for

future use in agriculture (12, 44).

Materials and Methods

Fungal cultures and spore suspensions

The fungal strains *Botrytis cinerea* T4 and *Fusarium graminearum* PH-1 were each cultured in

their respective normal growth medium shown in *SI Appendix* Table S1. Fungal spores were

harvested by flooding the fungal growth plates with sterile water. The spore suspension was

filtered through two layers of Miracloth, centrifuged at 13,600 rpm for 1 min, washed, and re-

suspended in low-salt Synthetic Fungal Medium (SFM) (45). The spore suspension of each fungal

pathogen was adjusted to the desired spore density of ~10^5 spores/mL using a hemocytometer.

Peptide synthesis and purification

The peptides GMA4CG_WT and GMA4CG_V1-V6 were chemically synthesized with an initial

purity of ~80-85% from multiple commercial vendors and further purified using a C-18 reverse

phase HPLC as described (35). The HPLC fractions containing the peptide were lyophilized and

resuspended in the nuclease-free water. Peptide concentrations were determined using the BCA

assay. For peptides containing Trp, peptide concentration was also determined using NanoDrop

2000c Spectrophotometry (Thermo-Fisher Scientific, MA) to confirm concentrations determined

by the BCA assay.

Antifungal activity assays

The minimal inhibitory concentration (MIC) of each peptide was determined using the resazurin

cell viability assay (28, 46). Briefly, this involved incubation of the pathogen/peptide mixture for

48 h, adding 0.01% (w/v) resazurin solution to each well, and re-incubating overnight. A change

in color of the resazurin dye from blue to pink or colorless indicated the presence of live fungal

cells. MIC was determined as the minimal concentration of peptide at which no change in color

was observed. The MIC value of each peptide was also determined in SFM supplemented with

100 mM NaCl, 100 mM KCl, or 2 mM CaCl$_2$.

Membrane permeabilization assay using SYTOX^{TM} Green (SG) uptake

The fluorescent dye SG (Thermo-Fisher Scientific, MA) was used to identify GMA4CG_V6-

induced membrane permeabilization of *B. cinerea* conidia and germlings as described previously

(28). After exposure to 1.5 µM GMA4CG_V6 for 30 min, *B. cinerea* spores and germlings were

imaged using fluorescence microscopy on a Leica SP8-X confocal microscope with an excitation

and emission wavelength of 488 and 508-544 nm, respectively.

The GMA4CG_V6 induced membrane permeabilization of conidia was also assessed

quantitatively in the absence and presence of 100 mM KCl. Fresh conidia at a concentration of $1\times$

10^5/mL were incubated with GMA4CG_V6 at $1\times$, $2\times$, and $4\times$ MIC for 1 h in the absence and

presence of 100 mM KCl for 1 h and 5 h, respectively. Samples were deposited into glass-bottom

petri dishes and imaged by confocal microscopy at an excitation wavelength of 488 nm and an

emission wavelength ranging from 508 to 544 nm in the presence of 1 µM SG added 15 min prior

to imaging. Fresh conidia incubated with SG alone were used as negative controls. A Leica SP8-

X confocal microscope was used for all confocal imaging and Leica LASX (version 3.1.5.)

software was used to process the images. The percentage of conidia showing membrane

permeabilization was determined by counting the number of cells showing fluorescent signal

relative to the total number of cells counted.

The effect of MtDef4, GMA4CG_WT, and GMA4CG_V6 on membrane integrity of *B.*

cinerea germlings was determined using a SG (final concentration: 1 µM) uptake quantification

assay in the presence and absence of 100 mM KCl (35). In brief, membrane permeabilization in

B. cinerea germlings treated with different concentrations (0.187, 0.375, 0.75, 1.5, and 3 µM) of

MtDef4, GMA4CG_WT, and GMA4CG_V6 was measured at 5 h in the presence and absence of

100 mM KCl. SG uptake was quantified by measuring the fluorescence in a SpectraMax® M3

spectrophotometer (excitation: 488 nm; cutoff: 515 nm; emission: 530 nm).

Internalization and subcellular localization of GMA4CG_V6 in *B. cinerea* germlings

Time-lapse confocal laser scanning microscopy was used to monitor uptake and subcellular

localization of fluorescently labeled peptide as described previously (35). Crude TMR fluorophore

labeled GMA4CG_V6 (TMR-GMA4CG_V6) was custom synthesized (Agentide Co, Piscataway,

NJ) and purified by HPLC. *B. cinerea* germlings were generated by incubating conidia overnight

in SFM medium. They were challenged with TMR-GMA4CG_V6 at concentrations of 0.5, 1.5, or

20 µM and imaged by time-lapse confocal microscopy. The excitation and emission wavelengths

for imaging TMR-GMA4CG_V6 were 562 and 580 to 680 nm, respectively.

The sub-localization of GMA4CG_V6 was determined by visualizing its localization with

the vacuole-specific dye CMAC and the cell membrane specific dye FM4-64. The excitation and

emission wavelengths for CMAC were 353 nm and 405 to 480 nm, respectively, and for FM4-64

were 562 and 680 to 800 nm, respectively. Co-localized pixels (white overlay) of TMR-

GMA4CG_V6 and CMAC/ FM4-64 was displayed using Leica LAS-X software. The threshold

for each channel was 50% and the pixels out of this range were considered uncorrelated. The

threshold for the background was set as 20%. Images were captured 30 min after incubation with

peptide.

Quantification of cell wall periphery binding intensity

The spores of *B. cinerea* were treated with 1.5 µM DyLight550-labeled MtDef4, TMR-

GMA4CG_WT, and TMR-GMA4CG_V6 for 20 min. Spores of each treatment were imaged using

a C-Apochromat 63× water immersion objective lens in apotome mode with a ZEISS Elyra7 super-

resolution microscope at excitation and emission wavelengths of 561 and 590 nm, respectively.

Images were processed using the SIM module and 3D leap processing with autosharpness setting

in ZEN Black 3.0 SR FP2 software. For each captured cell, the cell surface binding intensity was

quantified by free hand drawing a line intensity profile along the cell surface. The fluorescence

intensity of each cell wall periphery was normalized by its perimeter length as described (47). The

lower threshold was set to 7000 to subtract the signal in the cytoplasm.

Semi-*in planta* antifungal activity assays

The semi-*in planta* antifungal activity of each peptide against *B. cinerea* was determined using *N.*

benthamiana Nb1 detached leaves as described previously (28). Each peptide was tested at

concentrations ranging from 0.75, 1.5, 3, and 6 µM. Following incubation of each peptide/fungal

spore mixture at room temperature for 48 h, leaves were photographed in white light. High-

resolution fluorescence images were also taken using CropReporter (PhenoVation, Wageningen,

Netherlands). These images depicted the calculated F_V/F_M (maximum quantum yield of

photosystem II) values of diseased area affected by *B. cinerea* infection. Colors in the images show

five different classes ranging from class I to class V (0.000 to 0.700) depicting varying degrees of

tissue damage. On one end of the scale, green represents class V (0.600 to ≥0.700) high

photosynthesis efficiency. On the other end of the scale, red represents class I (0.000 to 0.160) low

photosynthesis efficiency.

***In planta* antifungal activity assays**

To test preventative and curative antifungal activity of the peptides against *B. cinerea*, four-week-

old *N. benthamiana* Nb1 plants grown under controlled growth conditions were used. For

preventative activity, GMA4CG_WT and GMA4CG_V6 at 12 or 24 µM concentration were

sprayed (2 mL/plant) on experimental plants. Control plants were sprayed with 2 mL of

water/plant. The peptide solution or water was allowed to dry out on the surface of the leaves. *B.*

cinerea spores ($\sim 8 \times 10^4$) suspended in $0.5 \times$ SFM were sprayed on the experimental and control

tomato plants (1 mL/plant), transferred to Ziploc plastic boxes under high humidity, and disease

symptoms recorded after 48 h. Photographs were taken at 72 h post-inoculation using both white

light and CropReporter (PhenoVation, Wageningen, Netherlands). Quantification of symptoms

was made by determining maximum quantum yield of PSII (Fv/Fm) as described above for the

semi-*in planta* assays.

For the curative antifungal activity tests, *N. benthamiana* Nb1 plants of similar age were

first sprayed with *B. cinerea* spores (~8×10^4) suspended in 0.5× SFM (1 mL/plant). After 2, 15,

and 30 h of pathogen challenge, 2 mL of GMA4CG_WT or GMA4CG_V6 was sprayed on the

plants at a concentration of 12 µM. Control plants were each sprayed with 2 mL of water. Disease

symptoms were observed after 48 h of pathogen challenge. Plants were photographed at 72 h post

infection in white light and quantification was done as described above. The curative antifungal

activity of GMA4CG_WT and GMA4CG_V6 at concentration of 6, 12, and 24 µM was tested on

3-week-old tomato (*Solanum lycopersicum* L. cv. Mountain Spring) plants pre-inoculated with *B.*

cinerea for 12 h. The severity of disease symptoms was rated using the scoring method for each

inoculated plant as follows: 0, no symptoms; 1= 1-25%, 2= 26-50%, 3= 51-75% and 4= 76-100%

of leaf surface infected.

Visualization of fungal growth on tomato leaf surface

Fresh *B. cinerea* conidia were collected and sprayed on leaves of 3-week-old tomato plants and

allowed to germinate for 12 h. After that, GMA4CG_V6 at 6 µM was applied on the same leaves

and incubated in a container under high humidity. Leaves applied with spores and sterile water

were set as negative controls. After 48 h, leaves were fixed in 4% paraformaldehyde, stained with

CFW for 30 min, and images were captured by an Axio Zoom.V16 microscope (ZEISS) with a

PlanNeoFluar Z 1.0x objective under tile mode. The excitation wavelength was 335-383 nm, and

the emission wavelength was 420-470 nm. Images were visualized and processed by Zen Black

3.0 SR FP2 software (ZEISS).

Statistical analysis

All experiments were performed in three or more technical replicates where appropriate and in three independent experiments. The phospholipid binding assays and quantitative measurements of cell membrane permeability in *B. cinerea* were performed using two independent biological replications. Data were graphed and analyzed in GraphPad Prism (version 8.0 or 9.4.1) software using unpaired, two-tailed Student's *t* tests, and one-way ANOVA with Tukey's honestly significant difference test (HSD) for comparison of multiple groups. Statistical significance was accepted at $p < 0.05$.

Acknowledgements

This work was supported, in part, by the National Science Foundation-EAGER grant IOS:1955461 awarded to D.S. We acknowledge imaging support from the Advanced Bioimaging Laboratory (RRID:SCR_018951) at the Danforth Plant Science Center and usage of the Leica SP8-X confocal microscope acquired through an NSF Major Research Instrumentation grant (DBI-1337680) and usage of the ZEISS Elyra 7 Super-Resolution Microscope acquired through an NSF Major Research Instrumentation grant (DBI-2018962). We are thankful to Dr. Saurav Misra of the Danforth Center for his attempts to generate structures of the peptides using AlphaFold. This work was funded by the Wells Fargo IN^2 grant awarded to Peptyde Bio. Part of this research was performed at the Pacific Northwest National Laboratory (PNNL), a facility operated by Battelle for the U.S. Department of Energy, including access to the W.R. Wiley Environmental Molecular Sciences Laboratory (EMSL), a national scientific user facility sponsored by the U.S. DOE Biological and Environmental Research program. Battelle operates PNNL for the U.S. Department of Energy under contract DE-AC05-76RL01830.

References

1. H. N. Fones *et al.*, Threats to global food security from emerging fungal and oomycete crop pathogens. *Nature Food* **1**, 332-342 (2020).

2. M. C. Fisher *et al.*, Emerging fungal threats to animal, plant and ecosystem health. *Nature* **484**, 186-194 (2012).

3. S. Savary *et al.*, The global burden of pathogens and pests on major food crops. *Nature Ecology & Evolution* **3**, 430-439 (2019).

4. M. C. Fisher, N. J. Hawkins, D. Sanglard, S. J. Gurr, Worldwide emergence of resistance to antifungal drugs challenges human health and food security. *Science* **360**, 739-742 (2018).

5. R. P. Oliver, H. G. Hewitt, "Fungicides in Crop protection." in Fungicides in Crop Protection. (CAB International, Oxfordshire, Boston., 2014).

6. N. J. Hawkins, B. A. Fraaije, Contrasting levels of genetic predictability in the evolution of resistance to major classes of fungicides. *Mol. Ecol*. **30**, 5318-5327, doi:10.1111/mec.15877 (2021).

7. F. Lobo, A. Boto, Host-defense peptides as new generation phytosanitaries: low toxicity and low induction of antimicrobial resistance. *Agronomy* **12**, 1614 (2022).

8. N. L. van der Weerden, M. R. Bleackley, M. A. Anderson, Properties and mechanisms of action of naturally occurring antifungal peptides. *Cell Mol Life Sci* **70**, 3545-3570 (2013).

9. H. P. van Esse, T. L. Reuber, D. van der Does, D. Genetic modification to improve disease resistance in crops. *New phytologist* **225**, 70-86 (2020).

10. R. K. Goyal, A. K. Mattoo, Multitasking antimicrobial peptides in plant development and

host defense against biotic/abiotic stress. *Plant Science (Shannon, Ireland)* **228**, 135-149

(2014).

11. T. L. Cools, C. Struyfs, B. P. Cammue, K. Thevissen, Antifungal plant defensins: increased

insight in their mode of action as a basis for their use to combat fungal infections. *Future*

Microbiol **12**, 441-454 (2017).

12. K. Parisi *et al.*, The evolution, function and mechanisms of action for plant defensins.

Semin Cell Dev Biol **88**, 107-118 (2019).

13. V. Leannec-Rialland *et al.*, Use of defensins to develop eco-friendly alternatives to

synthetic fungicides to control phytopathogenic fungi and their mycotoxins. *J Fungi*

(Basel) **8,** doi:10.3390/jof8030229 (2022).

14. V. Kovaleva, I. Bukhteeva, O. Y. Kit, I. V. Nesmelova, Plant defensins from a structural

perspective. *Int J Mol Sci* **21,** doi: 10.3390/ijms 21155307 (2020).

15. J. Kaur, U. S. Sagaram, D. M. Shah, Can plant defensins be sused to engineer durable

commercially useful resistance in crop plants? *Fungal Biology Reviews* **25**, 128-135

(2011).

16. A. F. Lacerda, E. A. Vasconcelos, P. B. Pelegrini, M. F. Grossi de Sa, M.F. Antifungal

defensins and their role in plant defense. *Front Microbiol* **5**, 116 (2014).

17. K. El-Mounadi, K. T. Islam, P. Hernandez-Ortiz, N. D. Read, D. M. Shah, Antifungal

mechanisms of a plant defensin MtDef4 are not conserved between the ascomycete fungi

Neurospora crassa and *Fusarium graminearum*. *Mol Microbiol* **100**, 542-559 (2016).

18. U. Sagaram *et al.*, Structural and functional studies of a phosphatidic-acid binding

antifungal plant defensin MtDef4: Identification of an RGFRRR motif governing fungal

cell entry. *PLoS One 8,* e82485 doi:10.1371/journal.pone 0082485 (2013).

19. U. S. Sagaram, R. Pandurangi, J. Kaur, T. J. Smith, D. M. Shah. Structure-activity

determinants in antifungal plant defensins MsDef1 and MtDef4 with different modes of

action against *Fusarium graminearum. PloS One* **6**, e18550,

doi:10.1371/journal.pone.0018550 (2011).

20. A. E. Sathoff, S. Velivelli, D. M. Shah, D. A. Samac. Plant defensin peptides have

antifungal and antibacterial activity against human and plant pathogens. *Phytopathol* **109**,

402-408 (2019).

21. M. Dathe, H. Nikolenko, J. Klose, M. Bienert, Cyclization increases the antimicrobial

activity and selectivity of arginine- and tryptophan-containing hexapeptides. *Biochemistry*

43, 9140-9150 (2004).

22. A. T. Tucker *et al.*, Discovery of next-generation antimicrobials through bacterial self-

screening of surface-displayed peptide libraries. *Cell* **172**, 618-628, e13 (2018).

23. B. Deslouches *et al.*, De novo generation of cationic antimicrobial peptides: influence of

length and tryptophan substitution on antimicrobial activity. *Antimicrob Agents and*

Chemother **49**, 316-322 (2005).

24. D. I. Chan, E. J. Prenner, H. J. Vogel, Tryptophan- and arginine-rich antimicrobial

peptides: structures and mechanisms of action. *Biochimica et Biophysica Acta* **1758**, 1184-

1202 (2006).

25. J. A. Killian, G. von Heijne, How proteins adapt to a membrane-water interface. *Trends*

Biochem Sci **25**, 429-434 (2000).

26. J. Lu *et al.*, D- and Unnatural amino acid substituted antimicrobial peptides with improved

proteolytic resistance and their proteolytic degradation characteristics. *Front Microbiol* **11**,

563030 (2020).

27. B. K. Kerenga *et al.*, Salt-tolerant antifungal and antibacterial activities of the corn defensin

ZmD32. *Frontiers in Microbiology* **10**, 795 (2019).

28. H. Li, S. L. Velivelli, D. M. Shah, Antifungal potency and modes of action of a novel olive

tree defensin against closely related ascomycete fungal pathogens. *Mol Plant Microbe*

Interact **32**, 1649-1664 (2019).

29. J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,

583-589 (2021).

30. N. J. Greenfield, Using circular dichroism spectra to estimate protein secondary structure.

Nature protocols **1**, 2876-2890 (2006).

31. G. M. Holzwarth, P. Doty, The ultraviolet circular dichroism of polypeptides. *J Am Chem*

Soc. **87**, 218-228 (1965).

32. J.L. Lopes, Miles, A.J. Miles, L. Whitmore, B.A. Wallace Distinct circular dichroism

spectroscopic signatures of polyproline II and unordered secondary structures: applications

in secondary structure analyses. *Protein Sci* **23**, 1765-72 (2014).

33. Jumper, J. *et al*. Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,

583–589 (2021).

34. Baek M. *et al*, (2021) Accurate prediction of protein structures and interactions using a 3-

track network. *Science* **373** (6557), 871-876. DOI: 10.1126/science. abj8754

35. S. L. Velivelli *et al.*, Antifungal symbiotic peptide NCR044 exhibits unique structure and

multi-faceted mechanisms of action that confer plant protection. *PNAS* **117,** 16043-16054

(2020).

36. K. Bi *et al.*, The *Botrytis cinerea* Crh1 transglycosylase is a cytoplasmic effector triggering

plant cell death and defense response. *Nat Commun* **12**, 2166 (2021).

37. Z. Shi, C. A. Olson, N. R. Kallenbach, Cation-pi interaction in model alpha-helical

peptides. *J Am Chem Soc* **124**, 3284-3291 (2002).

38. D. J. Schibli, R. F. Epand, H. J. Vogel, R. M. Epand, Tryptophan-rich antimicrobial

peptides: comparative properties and membrane interactions. *Biochem Cell Biol* **80**, 667-

677 (2002).

39. A. Falanga *et al.*, Cyclic Peptides as novel therapeutic microbicides: Engineering of human

defensin mimetics. *Molecules* **22,** (2017) doi: 10.3390/molecules22071217.

40. R. G. Spelbrink *et al.*, Differential antifungal and calcium channel-blocking activity among

structurally related plant defensins. *Plant Physiol* **135**, 2055-2067 (2004).

41. F. R. Terras *et al.*, Analysis of two novel classes of plant antifungal proteins from radish

(Raphanus sativus L.) seeds. *J Biol Chem* **267**, 15301-15309 (1992).

42. A. Munoz, J. F. Marcos, N. D. Read, Concentration-dependent mechanisms of cell

penetration and killing by the de novo designed antifungal hexapeptide PAF26. *Mol*

Microbiol **85**, 89-106 (2012).

43. P. Schwinges *et al.*, A bifunctional dermaseptin-thanantin dipeptide functionalizes the crop

surface for sustainable pest management. *Green Chemistry* DOI: 10.1039/c9gc00457b

(2019).

44. T. M. Shafee, M. A. Anderson, A quantitative map of protein sequence space for the cis-
defensin superfamily. *Bioinformatics* **35**, 743-752 (2019).

45. J. Liang, D. M. Shah, Y. Wu, C. A. Rosenberger, S. M. Hakimi (2001) Antifungal
polypeptide from alfalfa and methods for controlling plant pathogenic fungi. (Monsanto
Company, USA).

46. S. Chadha, S. P. Kale, Simple fluorescence-based high throughput cell viability assay for
filamentous fungi. *Lett Appl Microbiol* **61**, 238-244 (2015).

47. N. Zhang *et al.*, Comparative phenotyping of two commonly used *Chlamydomonas*
reinhardtii background strains: CC-1690 (21gr) and CC-5325 (The CLiP Mutant Library
Background). *Plants (Basel)* **11,** 585 (2022).

Figure 1.

A

```
MtDef4:    RTCESQSHKFKGPCASDHNCASVCQTERFSGGRCRGFRRRCFCTTHC
```

	Sequences	Net charge	% hydrophobic amino acids
GMA4CG_WT:	GGRCRGFRRRCFCTTHC	4.9	11.8
GMA4CG_V1:	GGRCKGFRRRCFCTRIC	5.8	17.7
GMA4CG_V2:	GGRCRGFRRRCFCTRIC	5.8	17.7
GMA4CG_V3:	GGRCKGFRRRVFVTRIC	5.9	29.4
GMA4CG_V4:	FGRCRGFRRRCFCWRWC	5.8	29.4
GMA4CG_V5*:	FGZCZGFZZZCFCWZWC	5.8	29.4
GMA4CG_V6:	GGRCKGFRRRWFWTRIC	5.9	29.4

B

Peptide	B. cinerea MIC (µM)				F. graminearum MIC (µM)
	SFM	SFM+ 100 mM NaCl	SFM+ 100 mM KCl	SFM+ 100 mM CaCl$_2$	SFM
MtDef4	0.75	>6	>6	>6	1.5
GMA4CG_WT	3	3	6	6	3
GMA4CG_V1	3	1.5	3	3-6	3
GMA4CG_V2	6	3	6	6	6
GMA4CG_V3	3	1.5-3	>6	>6	3
GMA4CG_V4	1.5-3	1.5	1.5	3	3-6
GMA4CG_V5	3-6	6	>6	>6	3-6
GMA4CG_V6	1.5	1.5	1.5-3	3	1.5-3

Figure 1. Amino acid sequences and antifungal activity of MtDef4-derived GMA4CG_WT and its variants. A. Amino acid sequence, net charge, and percentage of hydrophobic residues for GMA4CG_WT and its variants. The amino acid sequence of MtDef4 is shown on top with its γ-core motif highlighted by a red rectangle. The amino acid substitutions in GMA4CG_WT are shown in red. Disulfide bonds verified in GMA4CG_V3 and GMA4CG_V6 are shown with a link. The letter Z indicates the unnatural amino acid 2, 4-diaminobutyric acid. **B.** The antifungal activity of MtDef4, GMA4CG_WT, and its variants against *B. cinerea* in the absence and presence of cations and against *F. graminearum* in the absence of cations. MIC (Minimal Inhibitory Concentration) of each peptide was determined by the resazurin assay. SFM (Synthetic Fungal Medium) is a low salt medium used in this study.

Figure 2.

Figure 2. Topical application of GMA4CG_WT and its variants on detached *N. benthamiana* leaves confers resistance to gray mold. A. Scheme for application of GMA4CG_WT and GMA4CG_V1-V5 on tobacco leaves. Gray mold lesions on *N. benthamiana* leaves were observed and photographed at 48 hpi. The white light and CropRepoter images of the leaves are shown. **B.** Box & whisker plots were used to represent the quantification of lesion sizes. **C**. Scheme for application of GMA4CG_V4 and GMA4CG_V6 on *N. benthamiana* leaves. The white light and CropReporter images of the infected leaves are shown. **D**. Box & whisker plots were used to represent the quantification of lesion sizes. In **B** & **D**, the horizontal lines represent the median and boxes indicate the 25th-75th percentile. One-way analysis of variance (ANOVA) was used to determine statistical differences in lesion sizes between control (*B. cinerea*) and peptides (GraphPad Prism-version 9.4.1). Three technical replications of tobacco leaves (n = 3) were tested in three independent biological replications. Asterisks represent significant differences between

 different groups ($^{ns}p > 0.05$, $*p \leq 0.05$ $**p < 0.01$, $****p \leq 0.0001$) using Tukey's honestly significant
difference test for comparison of multiple groups.

Figure 3.

Figure 3. GMA4CG_V6 permeabilizes cell membrane of *B. cinerea*. A. Confocal microscopy
images of the SG uptake in *B. cinerea* spores and germlings treated with 1.5 µM of GMA4CG_V6
for 30 min. Spores and germlings treated with H₂O served as controls. Scale bars = 20 µm. **B.**
Representative images showing SG signals in *B. cinerea* spores treated with 1.5, 3, and 6 µM of
GMA4CG_V6 for 1 h with no salt, and for 1 h and 5 h with 100 mM KCl. The outline of each
spore is highlighted by a white dashed line. Scale bar = 20 µm. The permeabilization rate (%) of
each treatment is labeled on the images. **C.** Quantitative measurements of cell membrane
permeability in terms of fluorescence emitted by *B. cinerea* germlings treated with various
concentrations of MtDef4 (red), GMA4CG_WT (green), and GMA4CG_V6 (purple) with (dashed
line) and without (solid line) 100 mM KCl. Membrane permeabilization increased with increasing
concentration of each peptide. Values are means of two biological replications. Error bars indicate
standard deviations. Fluorescence measurements were taken at 5 h.

　Figure 4.

Figure 4. Internalization of GMA4CG_V6 into *B. cinerea*. A. Time-lapse confocal microscopy
images of *B. cinerea* showing internalization of 1.5 µM TMR-GMA4CG_V6 over time (T in min).
Images were captured every 4 min starting from 3 min. Arrows indicate vacuoles. Scale bars = 10
µm. **B.** Individual fluorescent (CMAC, TMR, and FM4-64) and channel merged images of *B.*
cinerea germlings treated with GMA4CG_V6 for 30 min. The overlap signal of CMAC and TMR

or TMR and FM4-64 are shown as white pixels. The threshold for each channel and the background was 50% and 20%, respectively. Scale bars = 10 µm. **C.** The fluorescent signal intensity of the ROI in **B**. Black arrows indicates TMR-GMA4CG_V6 and FM4-64 signals overlap at the membrane and the red arrows indicate TMR-GMA4CG_V6 and CMAC signals overlap at the vacuoles. **D.** Cell surface fluorescence shown in 2.5D. The vertical direction indicates the fluorescence intensity of each pixel with taller peaks representing greater fluorescence intensity. The corresponding 2D images are shown. **E.** Box plot for cell surface fluorescence quantification. Total fluorescence intensity of each cell was normalized to the cell perimeter length. The lower threshold was set to 7000, the substrate signal in the cytoplasm. Statistical analyses were performed using a two-tailed *t*-test assuming unequal variance (GraphPad Prism 8.0). *P*-value over 0.05 indicated no significant difference and less than 0.05 indicates a significant difference. n = 24-39.

Figure 5.

Figure 5. **GMA4CG_V6 exhibits preventative and curative antifungal activity against gray mold in *N. benthamiana* plants. A.** Preventative activity of GMA4CG_V6. Four-week-old *N. benthamiana* plants were sprayed with either 2 mL of water or peptide (GMA4CG_WT or GMA4CG_V6) at 12 µM or 24 µM concentrations prior to spray-inoculation with 1 mL of 8×10^4 spores per mL of *B. cinerea* fresh spore suspension. The mock plants were only sprayed with water (no fungal spore suspension). The CropReporter images of plants were taken 72 hpi. Maximum quantum yield of PSII photochemistry (Fv/Fm) was determined at 72 hpi. Red color represents class I, low photosynthesis efficiency attributed to extensive tissue damage. Green color indicates class V, high photosynthesis efficiency. **B.** The calculated photosynthetic quantum yield (Fv/Fm) for individual *N. benthamiana* plants. **C.** Curative antifungal activity of GMA4CG_V6. Four-week-old *N. benthamiana* plants were first spray-inoculated using 1 mL of 8×10^4 *B. cinerea* spore suspension. At 2, 15, or 30 hpi the plants were sprayed with 2 mL of peptide (GMA4CG_WT or GMA4CG_V6) at a concentration of 12 µM or water. The mock plants were only sprayed with

water (no fungal spore suspension). Maximum quantum yield was determined as described above.
D. The calculated photosynthetic quantum yield (Fv/Fm) for individual *N. benthamiana* plants. In
B and **D** the horizontal lines represent the median and boxes indicate the 25^{th} and 75^{th} percentiles.
One-way analysis of variance (ANOVA) was used to determine statistical difference between
control (*B. cinerea* alone), mock (water alone), and peptide-treated inoculated plants (GraphPad
Prism-version 9.4.1). Each technical replication included four plants (n = 4). Three biological
replications were performed. Asterisks represent significant differences between different groups
with *p* values ($^{ns}p > 0.05$, $*p \leq 0.05$ $**p < 0.01$, $****p \leq 0.0001$) determined using Tukey's honestly
significant difference test for comparison of multiple groups.

   Figure 6.

Figure 6. **GMA4CG_V6 exhibits curative antifungal activity against gray mold in tomato plants. A.** Three-week-old tomato plants were spray-inoculated with 1 mL of 8×10^4 *B. cinerea* spore suspension. At 12 hpi, plants were sprayed with 2 ml water or 2 ml peptide (GMA4CG_WT or GMA4CG_V6) at concentrations of 6, 12, or 24 µM. The CropReporter images of the plants were taken 72 hpi and the maximum quantum yield of PSII photochemistry (Fv/Fm) was determined. Red color represents class I, low efficiency photosynthesis attributed to extensive tissue damage. Green color represents class V, high efficiency photosynthesis. **B.** The calculated photosynthetic quantum yield (Fv/Fm) for individual tomato plants. The horizontal lines represent the median and boxes indicate the 25^{th} and 75^{th} percentiles. One-way analysis of variance (ANOVA) was used to determine statistical difference between control (*B. cinerea* alone), mock

 (water alone), and peptide-treated inoculated plants (GraphPad Prism-version 9.4.1). Each
technical replication included four plants (n= 4). Three biological replications were performed.
Asterisks represent significant differences between different groups with p values ($^{ns}p > 0.05$, $*p \leq$
0.05 $**p < 0.01$, $****p \leq 0.0001$) determined using Tukey's honestly significant difference test for
comparison of multiple groups. **C.** Germination and growth of *B. cinerea* spores on the tomato leaf
surfaces in the presence of GMA4CG_V6 or water. Tomato plants were inoculated with *B. cinerea*
spores. Infection was allowed to progress for 12 h and then the plants were sprayed with water or
with GMA4CG_V6. Forty-eight hours post peptide or water treatment, leaf samples were fixed in
4% paraformaldehyde and stained with Calcofluor White for 30 min. ZEISS Axio ZoomV16 was
used to capture images. Arrowheads point to the germinated hyphae of *B. cinerea*.

Figure 7.

Figure 7. A proposed model for the modes of antifungal action of GMA4CG_V6 against *B.*
***cinerea*.** GMA4CG_V6 sprayed on *B. cinerea* infected leaves first binds predominantly to the

fungal plasma membrane with little or no binding to the cell wall. Binding to plasma membrane-resident phosphoinositides presumably plays a role in membrane permeabilization. At low concentration, GMA4CG_V6 is internalized into cell vacuoles. It remains to be determined if this peptide is internalized into vacuoles through endocytic mechanisms. At high concentration, GMA4CG_V6 targets the cytoplasm immediately following cell entry. Cell death occurs only after GMA4CG_V6 diffuses into the cytoplasm and not while in vacuoles. N: nucleus, V: vacuole, E: endosome.

Supplementary information

Extended materials and methods

Cell death assay of GMA4CG_V6 with propidium iodide

To determine fungicidal activity, freshly harvested conidia of *B. cinerea* were incubated with 1.5 μM GMA4CG_V6 for 1, 6, and 24 h. After incubation, conidia were centrifuged and washed twice in 1x SFM and re-suspended in the same medium. The spores were subsequently allowed to germinate overnight. Microscope images were taken, and germination rate was calculated. Propidium iodide (PI) at 0.5 μg/mL was used as a marker to count dead spores. Conidia of *B. cinerea* were incubated with GMA4CG_V6 for 0.25, 0.5, 1, and 2h. PI was added to the spores/peptide mixture 10 min in advance of detection. Images were captured on the Leica SP8-X confocal microscope (excitation: 535 nm, emission: 600-650 nm).

Circular dichroism spectroscopy

A steady-state wavelength circular dichroism spectrum was collected for GMA4CG_WT and GMA4CG_V6 (~0.05 mM peptide, 20 mM sodium acetate, pH 5.6) on a calibrated Aviv Model 410 spectropolarimeter (Lakewood, NJ). Using a quartz cell of 0.1 cm path length and a temperature of 20 °C, the data was recorded in 0.5 nm increments between 200 and 260 nm with a bandwidth of 1.0 nm and a time constant of 1.0 s. The reported steady-state wavelength spectrum

was the result of averaging two consecutive scans, subtracting a blank spectrum, and automatic

line smoothing using Aviv software.

Antifungal activity of the alanine scanning variants of GMA4CG_V6 with no disulfide bond

The minimal inhibitory concentration (MIC) of linear GMA4CG_V6 alanine scanning variants

was determined using the resazurin cell viability assay as described in the methods section. The

MIC value of all variants was also determined in SFM supplemented with 100 mM NaCl.

Cell wall/surface localization analysis of GMA4CG_V6

B. cinerea germlings were treated with 1.5 µM of TMR-GMA4CG_V6 and, after 5 min, CFW (5

µM) was added into the same petri dish. Samples were observed using a confocal (Leica SP8-X)

and super-resolution microscope (Zeiss Elyra7). To avoid bleed-through of the two fluorescent

signals, confocal images were captured sequentially using a fast line switch configuration (TMR-

GMA4CG_V6 excitation: 542 nm, emission: 580-680 nm, and CFW excitation: 405 nm, emission:

410-557 nm). To obtain higher resolution images, the C-Apochromat 63×1.4 oil immersion

objective lens in apotome mode was used with the ZEISS Elyra7 super-resolution microscope

(Zeiss). The excitation wavelength was 405 nm for the CFW signal and 561 nm for the TMR signal

with emission wavelengths of 420-480 and 570-620 nm, respectively. Images were processed

using the SIM module plus 3D and leap processing with autosharpness. The channels were aligned

based on 0.1 µm TetraSpeck™ microspheres (T7279, Thermo Fisher Scientific) using ZEN Black

3.0 SR FP2 software.

Fluorescence Redistribution After Photobleaching (FRAP) experimental setup and analysis

The FRAP experiments were performed on a Leica SP8-X microscope (FRAP Wizard Module, LAS X Version 3.1.5.16.16308) using the HC PlanApochromat 63× water immersion objective len (Numerical aperture = 1.2) with the pinhole set to 1 Airy unit and zoom set to 1.77 (1). Images were acquired for the selected cellular regions (germling cytoplasm, vacuole, cell periphery, and spore cell wall/periphery) after 1 h incubation with GMA4CG_V6 under the following identical imaging and FRAP conditions: 10 pre-bleach and 700 post-bleach frames (512 x 64 pixels, pixel size = 204 nm) captured at 173 ms intervals without averaging using 550 excitation and 560-625 nm emission wavelengths with PMT detectors and a transmitted light channel. The power of the 550 nm laser was set at 3.5% during imaging to prevent detectable photobleaching for the duration of each FRAP experiment (~3 min). A 1.52 μm^2 circular bleach region was applied with 100% laser excitation power (550 nm wavelength) for 3.5 s immediately following pre-bleach and prior to post-bleach imaging. FRAP analysis was performed on 24 data sets for each subcellular region showing no evidence of sample drift during acquisition. A single exponential curve fitting function was applied post-acquisition to extract *the half-maximum tau (τ½), mobile fraction, and immobile fraction for each condition.*

Phospholipid binding assays

His-tagged GMA4CG_WT and GMA4CG_V6 was obtained from Biomatik Inc. (Ontario, Canada) and further purified by HPLC. The His-tagged peptides (GMA4CG_WT and GMA4CG_V6) phospholipid binding assays were performed using the PIP Strips™ (Echelon Biosciences Inc., CA) as previously described (2). Hexa-His peptide (Genscript, Inc., NJ) was used as a control. The peptide binding to the membrane strips was detected using 0.5 µg/mL His-tag

1010 monoclonal antibody purchased from Thermo-Fisher Scientific, MA. The binding signals were

1011 detected using the SuperSignal West Pico Chemiluminescent Substrate Kit (Thermo-Fisher

1012 Scientific, MA) following the manufacturer's protocol. Quantitative analysis of the signal was

1013 performed using Image J (https://imagej.nih.gov/ij/download.html).

1014

The binding of GMA4CG_WT and GMA4CG_V6 on tomato leaf surfaces

The fluorophore-labeled peptides, TMR-GMA4CG_WT and TMR-GMA4CG_V6, were applied

on the surface of tomato leaves and allowed to air dry for 1 h. The leaf samples were subjected to

microscopic observation using a Leica SP8-X confocal microscope (TMR wavelength: excitation:

562 nm; emission: 580-680 nm, and autofluorescence of tomato cells wavelength: excitation: 405

nm; emission: 416-483 nm) with an APO CS2 40× water objective lens before and after a gentle

rinse twice in sterile water. The average signal intensity from 10-12 ROIs of each image was

collected for quantification and 26-32 images were captured. The signal intensity of the after-

wash-sample was normalized by the signal of the sample before washing.

1024

1025

1026

1027

1028

1029

1030

1031

1032

 Figure S1

 Figure S1. GMA4CG_V6 exhibits fungicidal activity. (**A**) Transmitted light images of the
 germination and growth of *B. cinerea* spores treated with 1.5 µM GMA4CG_V6 for 1, 6, and 24
 h. After each treatment, unbound peptide was removed, and the spores allowed to germinate
 overnight. Scale bar = 50 µm. (**B**) Quantification of the germination rate of GMA4CG_V6 treated
 B. cinerea spores. Spores treated with sterile water served as controls. Data were collected from

four independent replications (n = 4) and plots are shown in box plot. Individual data were overlaid
on plots as red points. Statistical analysis was performed by two-sided Student's t test. (**C**) *B.*
cinerea spores were treated with 1.5 μM GMA4CG_V6 and the dead cell staining dye PI for
various times. The overlaid images are shown for 0.25, 0.5, 1, and 2 h treatments. Scale bar = 20
μm. (**D**) Quantitative analysis of the percentage of spores stained with PI dye. The top of the
columns represents the mean, and the horizontal bars represent the standard deviation (n = 3).

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067   **Figure S2**

1068

Figure S2. GMA4CG_V6 retains antifungal activity in the presence of salts. (A) Antifungal
activity of the GMA4CG variants and MtDef4 against *B. cinerea* was determined in 1 x SFM in
the absence or presence of 100 mM NaCl, 100 mM NaCl, or 2 mM CaCl$_2$. After 48 h incubation,
resazurin dye was added to each well to determine cell viability/cell death. A change from blue to
pink/colorless indicated viable fungal cells. **(B)** Microscopic images of *B. cinerea* spores
challenged with various concentrations of GMA4CG_V6 in the presence of different salts (Scale
bar = 50 μm).

1076

1077 **Figure S3**

1078

1079 **Figure S3.** Steady-state wavelength CD spectra for GMA4CG_WT (red) and GMA4CG_V6
1080 (blue) in 20 mM sodium acetate, pH 5.6, at 20 °C.

1081

1082

1083   **Figure S4**

1084

Peptide	Sequence	SFM MIC (µM)	SFM+100 mM NaCl MIC (µM)
A GMA4CG_V6	GGRCKGFRRRWFWTRIC	1.5	1.5
GMA4CG_V6-Ala1	AAAAKGFRRRWFWTRIC	6	6
GMA4CG_V6-Ala2	GGRCAAARRRWFWTRIC	6	6
GMA4CG_V6-Ala3	GGRCKGFAAAAFWTRIC	12	24
GMA4CG_V6-Ala4	GGRCKGFRRRWAAARIC	6	6
GMA4CG_V6-Ala5	GGRCKGFRRRWFWTAAA	6	6

1085

Figure S4. Antifungal activity of the alanine scanning variants of linear GMA4CG_V6 as
determined by the resazurin cell viability/cell death assay. (A) The primary amino acid
sequences for GMA4CG_V6 and the five Ala scanning mutants and their associated MIC values
in the absence and presence of salt. The GMA4CG_V6 substitutions are highlighted red. **(B)**
Resazurin dye was used to determine cell viability/cell death. A change from blue to pink/colorless
indicated viable fungal cells. Antifungal activity of each variant was performed in 1 x SFM. There
was an eight-fold loss in activity with GMA4CG_V6_Ala3 whereas all other variants showed a
four-fold loss. **(C)** Resazurin cell viability/cell death assay showing the MIC values of alanine
scanning variants of GMA4CG_V6 in the presence of 100 mM NaCl. MIC values of all other
variants remained unchanged except for GMA4CG_V6_Ala3, which increased another two-fold.

1096

1097

1098 **Figure S5**

1099

1100

Figure S5. Internalization of TMR-GMA4CG_V6 into *B. cinerea* germlings at sublethal or
very high concentrations. (A) Confocal microscopy images show internalization of 0.5 µM
TMR-GMA4CG_V6. Images were captured 60 min after peptide challenge. **(B)** Time-lapse
confocal microscopy images of *B. cinerea* show internalization of 20 µM TMR-GMA4CG_V6
over time (T in min) with images captured every 3 min shown. Scale bars = 10 µm.

Figure S6

1107

1108

1109

Figure S6. Co-localization of TMR-GMA4CG_V6 and CFW. (A & C) Representative
fluorescent (CFW and TMR) and channel merged images of *B. cinerea* germlings treated with
TMR-GMA4CG_V6 and captured by a confocal microscope (A) and the overlay image by lattice
SIM^2 super resolution microscope (C). Scale bar is 10 µm in A and 5 µm in C. **(B & E)** The
fluorescent signal intensity of the ROIs in **A** and **C**, respectively. The dash lines in **E** indicate the
shift between the peak of the CFW and TMR signals. **(D)** Enlarged image from the framed area in
C. Scale bar = 2 µm.

1117

1118

1119

1120

1121

1122

1123

1124

1125 **Figure S7**

1126

1127

A

Normalized fluorescence recovery

- - - Germling cytoplasm
- - - Germling cell wall/ periphery
- - - Germling vacuole
- - - Spore cell wall/ periphery

0.6
0.4
0.2
0.0

0   50   100

Time (sec)

B

Immobile fraction (%)

$p< 0.0001$
$p< 0.0001$
$p< 0.0001$
$p= 0.0031$

Germling cell wall/ periphery
Germling cytoplasm
Germling vacuole
Spore cell wall/ periphery

C

	Germling cytoplasm	Germling cell wall/ periphery	Germling vacuole	Spore cell wall/ periphery
Tau	40.68	37.45	19.27	38.37
R^2	0.9636	0.9133	0.8496	0.9257

1128

Figure S7. FRAP analysis of the subcellular localization of GMA4CG_V6 in *B. cinerea*
germlings and conidia. (A) FRAP plots show the normalized fluorescence recovery of TMR-
GMA4CG_V6 in germlings cytoplasm (blue), cell walls/ periphery (orange), and vacuoles (gray)
and in conidia cell walls/periphery (green) over 120 s. A one-phase association parameter was
used to simulate the fitting curve (solid lines) based on the scatter diagram (GraphPad Prism 8.0).
Higher recovery rates indicate greater overall mobility. **(B)** The immobile fraction (%) of
GMA4CG_V6 localized into the key subcellular compartments of *B. cinerea*. Data are represented
using box plots with individual data overlaid as red points. Data analyzed for each subcellular
compartment of *B. cinerea* are from n = 20 collections from at least three independent experiments.

Asterisks represent significant differences between different subcellular compartments. One-way
analysis of variance (ANOVA) was used to determine the statistical difference with Tukey's
honestly significant difference test (HSD) for comparison of multiple groups. **(C)** A table showing
the half-time of recovery (Tau) of GMA4CG_V6 localized into the key subcellular compartments
of *B. cinerea*. The Tau and R^2 values for each peptide were calculated using a one-phase
association parameter.

1144

1145

1146   **Figure S8**

1147

1148

Figure S8. GMA4CG_V6 binding to phospholipids. (A) Lipid overlay assays of
GMA4CG_WT, GMA4CG_V6, and Hexa-His. Both GMA4CG_WT and GMA4CG_V6 show
strong binding to phosphatidylinositol monophosphates PI(3)P, PI(4)P, and PI(5)P while
GMA4CG_WT shows strong binding to additional phospholipids including PA and the
phosphatidylinositol diphosphates PI(3,5)P$_2$ and PI(4,5)P$_2$. In contrast, GMA4CG_V6 binds to
multiple phospholipids, including phosphatidylinositol di/triphosphates PI(3,4)P2, PI(3,4,5)P3,
and PA, but rather weakly. **(B & C)** Integrated density (grayscale pixel units) analysis of PIP strip
probed with GMA4CG_WT and GMA4CG_V6. Data are means ± SEM of two independent
biological replicates (n = 2).

Figure S9

1159

Figure S9. GMA4CG_V6 has preventative and curative antifungal activity against gray mold in *N. benthamiana* plants (**A**) Preventative activity of GMA4CG_V6. Four-week-old *N. benthamiana* plants were sprayed with either 2 ml water or peptide (GM4CG_WT or GMA4CG_V6) at 12 or 24 µM concentration prior to spray-inoculation with a 1 ml suspension of 8×10^4 *B. cinerea* fungal spores. Plants were observed for symptoms at 72 hpi and photographed in white light. (**B**) Disease severity of tobacco plants inoculated with *B. cinerea* at 72 hpi. The disease severity was rated on a scale of 0 to 4 with 0 = no symptoms; 1 = 1-25%; 2 = 26-50%; 3 = 51–75%; and 4 = 75-100% of the leaf surface infected. One-way analysis of variance (ANOVA)

was used to determine statistical differences between control (*B. cinerea*), mock-treated, and
peptide-treated plants (GraphPad Prism-version 9.4.1). Four *N. benthamiana* plants (n = 4) for
each technical replication were tested in three biological replications. Asterisks represent
significant differences between different groups with P values determined (^{ns}p> 0.05, *p≤ 0.05
**p< 0.01, ****p≤ 0.0001) using Tukey's honestly significant difference test for comparison of
multiple groups. **(C)** Curative antifungal activity of GMA4CG_V6. Four-week-old *N.*
benthamiana plants were spray-inoculated with a 1 mL suspension of 8×10^4 of *B. cinerea* fungal
spores at 2, 15, or 30 h and then sprayed with 2 mL of 12 μM GMA4CG_WT, 12 μM
GMA4CG_V6, or water. Plants were photographed in white light after 72 h. **(D)** Disease severity
was rated as described above. Four *N. benthamiana* plants (n = 4) for each technical replication
were tested in three biological replications. Asterisks represent significant differences between
different groups with *p* values determined (^{ns}p> 0.05, *p≤ 0.05 **p< 0.01, ****p≤ 0.0001) using
Tukey's honestly significant difference test for comparison of multiple groups.

1181

1182 **Figure S10**

1183

1184

1185 **Figure S10. GMA4CG_V6 has curative antifungal activity against gray mold in tomato**
1186 **plants. (A)** Three-week-old tomato (*Solanum lycopersicum* L cv. Mountain Spring) plants were
1187 spray-inoculated with a 1 ml suspension of 8×10^4 fungal spores and subsequently sprayed 12 h
1188 later with 2 ml of GMA4CG_WT or GMA4CG_V6 at a concentration of 6, 12, or 24 μM or with
1189 2 ml of water (controls). Plants were scored for gray mold symptoms at 72 hpi and photographed

in white light. **(B)** Disease severity of tomato plants inoculated with *B. cinerea* 72 hpi. The disease severity was rated on a scale of 0 to 4 with 0 = no symptoms; 1 = 1-25%; 2 = 26-50%; 3 = 51–75%; and 4 = 75-100% of the leaf surface infected. One-way analysis of variance (ANOVA) was used to determine statistical differences between control (*B. cinerea*), mock, and peptide experiments (GraphPad Prism-version 9.4.1). Four tomato plants (n = 4) for each technical replication were tested in three biological replications. Asterisks represent significant differences between different groups with *p* values ($^{ns}p > 0.05$, $*p \leq 0.05$ $**p < 0.01$, $****p \leq 0.0001$) determined using Tukey's honestly significant difference test for comparison of multiple groups.

Figure S11

Figure S11. Binding of TMR-GMA4CG_WT and TMR-GMA4CG_V6 to tomato leaf surfaces. (A) Representative confocal microscopy images of tomato leaves with no peptide applied showing the absence of TMR fluorescence signal, background cell wall autofluorescence

(cyan), and transmitted light. Scale bar = 20 µm. **(B)** Confocal microscopy images of plant cell
walls stained with TMR-GMA4CG_WT and TMR-GMA4CG_V6 showing no signal observation
in the cytoplasm. Scale bar = 20 µm. **(C)** Comparison of TMR signal intensity at the tomato cell
wall before and after washing with sterile water. The average signal from leaves treated with each
peptide before washing served as the control (100%). Each dot indicates individual data points.
Asterisks represent significant differences between different groups (unpaired *t* test, n = 26-32)
(GraphPad Prism 8.0).

1212

1213

1214 **Table S1.** Growth media and conditions for culturing the fungi used in this study

Strains	Stock	Medium for spore production	Culture conditions
B. cinerea **(T4)**	20% V8	20% V8	7-25 days, 25 ˚C
F. graminearum **(PH-1)**	PDA	CMC Broth	3-5 days, 28 ˚C, 180 rpm

1215

References

1. S. L. Velivelli *et al.* Antifungal symbiotic peptide NCR044 exhibits unique structure and
multi-faceted mechanisms of action that confer plant protection. *PNAS* **117,** 16043-16054
(2020).
2. K.T. Islam *et al*. A novel bi-domain plant defensin MtDef5 with potent broad-spectrum
antifungal activity binds to multiple phospholipids and forms oligomers. *Sci Rep* **7**, 16157
(2017).

1223
1224

1225

1226

1227

1228

1229

1230

1231

1232

1233