Selective accumulation of matrix proteins inside of peroxisomal subdomains

Julia Ast[1,2*], Nils Bäcker[1*], Domenica Martorana[1*], Christian Renicke[1,3], Thorsten Stehlik[1],

Thomas Heimerl[4,5], Christof Taxis[1,6], Gert Bange[4,5], Michael Bölker[1,5], Björn Sandrock[1], Kay

Oliver Schink[7,8§] and Johannes Freitag[1§]

1 Department of Biology, Philipps-University Marburg, Marburg, Germany

2 Institute of Metabolism and Systems Research (IMSR), and Centre of Membrane Proteins and Receptors

(COMPARE), University of Birmingham, Birmingham, United Kingdom

3 Department of Genetics, Stanford University School of Medicine, Stanford, CA, USA.

4 Department of Chemistry, Philipps-University Marburg, Marburg, Germany

5 Center for Synthetic Microbiology (SYNMIKRO), Philipps-University Marburg, Marburg, Germany

6 School of Life Sciences, University of Siegen, Siegen, Germany

7 Centre for Cancer Cell Reprogramming, Faculty of Medicine, University of Oslo, Montebello, Oslo, Norway

8 Department of Molecular Cell Biology, Institute for Cancer Research, Oslo University Hospital, Montebello,

Oslo, Norway

* these authors contributed equally

§ to whom correspondence should be addressed:

k.o.schink@medisin.uio.no

johannes.freitag@biologie.uni-marburg.de

Abstract

Formation of specialized reaction spaces prevents interference between distinct cellular

pathways. Peroxisomes are cellular compartments involved in a large diversity of metabolic

processes. How peroxisomes differentiate into subpopulations and by which mechanism

intraorganellar domains are formed remains largely elusive. Here, we report on enzymes from

the fungus *Ustilago maydis*, which accumulate inside of peroxisomal subdomains. We describe

a short peptide motif (Tyr-Ile-Ile-Val) sufficient to trigger focal localization. Mining for

proteins with similar motifs uncovered several peroxisomal matrix proteins that accumulate in

intraorganellar foci. These foci are enriched in the enzyme urate oxidase – a typical constituent

of the paracrystalline core of peroxisomes. Upon peroxisome proliferation uneven distribution

of focal structures results in the formation of peroxisome subpopulations with different protein

content. The underlying principle of subdomain formation is evolutionary conserved in

mammalian peroxisomes and formation of similar foci was also observed inside of

mitochondria. We propose that peroxisomal proteins show an individual propensity to self-

assemble. This formation of protein aggregates appears to be a ubiquitous driving force to

spatially organize the peroxisomal proteome.

Introduction

Peroxisomes are cellular compartments surrounded by a single membrane. They contain many metabolic enzymes, in particular those required for degradation of fatty acids and detoxification of hydrogen peroxide (Poirier et al., 2006; Smith and Aitchison, 2013). In humans, peroxisomes are essential and mutation of peroxisomal biogenesis proteins (Peroxins or Pex proteins) leads to severe disorders including the Zellweger syndrome (Wanders, 2014). Import of proteins into the matrix of peroxisomes requires either of two cytosolic targeting factors, Pex5 and Pex7 (Francisco et al., 2017; Kim and Hettema, 2015). They recognize specific sequence motifs located either at the Carboxy (*C*)-terminal end of the cargo protein (peroxisomal targeting signal type 1; PTS1) or within the Amino (*N*)-terminal part (peroxisomal targeting signal type 2; PTS2) (Lazarow, 2006; Walter and Erdmann, 2019). The prototype PTS1 bound by Pex5 is a tripeptide with the amino acid sequence Ser-Lys-Leu (Gould et al., 1989).

Fungal peroxisomes are diverse and contain specialized metabolic pathways e.g. for secondary metabolite and siderophore biosynthesis, methanol degradation and carbohydrate metabolism (Bartoszewska et al., 2011; Freitag et al., 2012; Gründlinger et al., 2013; Stehlik et al., 2014; van der Klei and Veenhuis, 2006). A peculiar variant of peroxisomes identified in *Neurospora crassa* are Woronin bodies, which seal septal pores in wounded hypha (Jedd and Chua, 2000; Nguyen et al., 2021). Differentiation of peroxisomes into Woronin bodies is achieved through expression of the Hex1 protein, which forms large crystals in Woronin bodies and is their major constituent (Pieuchot and Jedd, 2012; Tey et al., 2005).

The mechanisms by which peroxisomes differentiate into subpopulations in other biological systems or how they form intraorganellar domains are largely uncharacterized. Isolation and biochemical characterization of rat liver peroxisomes revealed peroxisomal populations with different protein content (Völkl et al., 1999). In yeast, distinct peroxisome populations can be distinguished by means of their age (Kumar et al., 2018). In plants, different types of peroxisomes emerge during development (Hu et al., 2012; Lingard et al., 2009). High-resolution imaging studies uncovered the presence of subdomains inside of the peroxisomal membrane of the yeast *Saccharomyces cerevisiae* (Galiani et al., 2016). Early electron microscopy and biochemical studies demonstrated that the enzyme urate oxidase is a major constituent of an electron-dense paracrystalline core structure (Rouiller & Bernard, 1956; De Duve, 1960; Baudhuin et al, 1965; De Duve & Baudhuin, 1966; Völkl et al, 1988).

Here we show, that a short peptide motif drives the accumulation of peroxisomal matrix proteins into subdomains, which we propose to be related to the electron-dense paracrystalline

core. Focal accumulation determines both the protein distribution within single peroxisomes

and the formation of peroxisomes with different protein content upon organelle proliferation.

Results

Identification of peroxisome subpopulations in *U. maydis*

We have previously shown that in *U. maydis* the acyltransferases, Mac1 and Mac3, involved in

biosynthesis of surface-active mannosylerythritol lipids (MELs), are localized to peroxisomes

(Becker et al., 2021; Bölker et al., 2008; Freitag et al., 2014; Kämper et al., 2006). Upon a

closer inspection of fluorescence microscopic images we noticed that both enzymes are not

homogenously distributed among the entire population of peroxisomes (Fig. 1A).

Figure 1 Acyltransferases Mac1 and Mac3 accumulate in peroxisome subpopulations
U. maydis strains expressing N-terminally GFP-tagged versions of Mac1 and Mac3 (cyan) and the peroxisomal
marker protein mCherry-SKL (magenta) were inspected by epifluorescence microscopy. Representative images
of glucose grown cells (A) or oleate grown cells (B). Full images are shown as overlays of the green and red
channel. For insets single channels and merged channels are depicted. (C) Quantifications show Pearson's
correlation coefficients of GFP and mCherry signals for indicated strains. (D) Representative epifluorescence
image of cells expressing mCherry-Mac3 together with GFP-SKL organized as described above. (E) GFP-SKL
and mCherry-SKL or GFP-Mac3 and mCherry-Mac3 were co-expressed and representative images are shown
organized as described above. (F) Quantification show Pearson's correlation coefficients of GFP and mCherry
signals for indicated strains. (A); (B); (D) and (E) Scale bars: 5 μm.

Green fluorescent protein (GFP) tagged Mac1 or Mac3 showed only partial colocalization with
the red fluorescent peroxisomal reporter mCherry-SKL (Fig. 1A). Oleic acid is known to induce
peroxisome proliferation in various species including *U. maydis* (Camões et al., 2015;
Kohlwein et al., 2013). Four hours of incubation in oleic acid-containing medium substantially
enhanced the observed subpopulation phenotype (Fig. 1B). Quantifications revealed low
correlation coefficients for the red and the green signal (Figs. 1C, S1A and S1B). Reciprocal
exchange of fluorescent tags did not alter our results indicating that it was not the specific
fluorescent protein, which caused the remarkable enrichment in peroxisome subpopulations
(Fig. 1D). As a control, we expressed either the combination of GFP-Mac3 and mCherry-Mac3
or the combination of GFP-SKL and mCherry-SKL and observed a high degree of co-
localization (Fig. 1E). In aggregate these data show that both acyltransferases are localized only
in a fraction of peroxisomes suggesting that in *U. maydis* specific populations of peroxisomes
with different protein content exist.
105

Accumulation of Mac1 and Mac3 inside of a subdomain of peroxisomes
GFP-Mac1 or GFP-Mac3 fluorescence seemed not to fill the entire peroxisome but was
enriched in distinct foci that merged with only a section of the organelle stained by mCherry-
SKL (e.g. Fig. 1B). Therefore, we analyzed suborganellar localization of GFP-Mac1 and GFP-
Mac3 by super-resolution imaging using structured illumination microscopy (SIM)
(Gustafsson, 2000). Analysis of SIM images revealed that both fusion proteins indeed
concentrate in subdomains of peroxisomes, while mCherry-SKL and GFP-SKL were
distributed among the whole organelle (Fig. 2A to 2C). This was as well visualized by 3D-
reconstructions (Fig. 2D). Time-lapse imaging showed that foci containing acyltransferases are
stable over time (Fig. S2A and Movie 1).
Previous studies of mammalian peroxisomes uncovered a paracrystalline core region, which
typically contains the enzyme urate oxidase as a major constituent but also harbors additional
proteins (De Duve & Baudhuin, 1966; Völkl *et al*, 1988). We determined the localization of the
putative *U. maydis* urate oxidase ortholog (UMAG_00672) fused to GFP (GFP-Uox1). GFP-
Uox1 accumulated in foci similar to GFP-Mac1 and GFP-Mac3 and also occurred only in a
subpopulation of peroxisomes (Fig. 2E) suggesting that all three proteins are enriched in
structures probably representing the paracrystalline core of peroxisomes. Indeed, co-expression
of GFP-Uox1 and mCherry-Mac1 or GFP-Uox1 and mCherry-Mac3 resulted in a high degree
of colocalization (Fig. 2E and Fig. S2B).

Figure 2 Acyltransferases Mac1 and Mac3 accumulate in peroxisome subdomains also enriched for urate oxidase Uox1.

Cells were analyzed by super-resolution microscopy using SIM. Representative images of cells expressing GFP-Mac1 and mCherry-SKL (A), GFP-Mac3 and mCherry-SKL (B) and GFP-SKL and mCherry-SKL (C). Full images are shown as overlays of the green and red channel. For insets single channels and merged channels are depicted. Associated graphs show the normalized fluorescence intensity of GFP and mCherry along the indicated lines. (D) 3D-reconstruction of GFP-Mac3 containing peroxisomal subdomains (cyan). Scale bar: 0.5 μm (E) Representative epifluorescence images of cells expressing GFP-Uox1 with mCherry-SKL (left) or mCherry-Mac3 (right). Organization of pictures as described above. Scale bars: 5 μm. (F) Crude organelle preparations of indicated strains were imaged by epifluorescence microscopy after incubation in lysis buffer supplemented with Triton X-100 (right). Preparations incubated in lysis buffer without Triton X-100 served as control (left). Organization of pictures as described above. Scale bars: 5 μm. (A) – (F) Signals of GFP-tagged proteins are shown in cyan and of mCherry-tagged proteins in magenta. (G) Transmission electron micrograph of cells expressing GFP-Mac3. Staining was achieved through labeling of GFP-Mac3 with anti-GFP and subsequent immunogold labeling followed by silver enhancement. Scale bar: 0.2 μm.

It was reported previously that the paracrystalline core of peroxisomes can sustain treatment with nonionic detergent (Tsukada et al., 1966). We prepared crude organelle extracts from

strains expressing mCherry-SKL together with GFP-Mac3, mCherry-Mac3 together with GFP-

Uox1 or mCherry-Mac1 together with GFP-Mac3. Triton X-100-treated or non-treated samples

were inspected by epifluorescence microscopy. While detection of mCherry-SKL was highly

sensitive to treatment with detergent, fluorescently labelled Mac1, Mac3 and Uox1 remained

visible inside of detergent resistant punctate structures (Fig. 2F and Fig. S2C). Furthermore,

transmission electron microscopy in combination with immunogold labeling showed that GFP-

Mac3 is concentrated within a subdomain of peroxisomes (Fig. 2G). Hence, foci enriched for

all three enzymes are structures that likely represent the paracrystalline core of *U. maydis*

peroxisomes.

A short peptide motif sufficient for enrichment of proteins inside of the core region

The C-terminal 12 amino acids contribute to targeting efficiency of PTS1 proteins (Brocard and

Hartig, 2006). To examine whether information within the peroxisomal targeting signal is

involved in formation of subdomains and subpopulations, we analyzed the localization of GFP

fused to the C-terminal PTS1 containing dodecamers of Mac1 and Mac3 (GFP-PTS1$_{Mac1}$ and

GFP-PTS1$_{Mac3}$) (Fig. 3A). Interestingly, only GFP-PTS1$_{Mac3}$ accumulated in subpopulations

and was enriched inside of peroxisomal foci, while GFP-PTS1$_{Mac1}$ colocalized with mCherry-

SKL positive peroxisomes and was homogenously distributed within the organelle (Fig. 3B

and 3C). By truncation analysis we identified a short peptide motif inside of the 12mer derived

from Mac3 consisting of the amino acid stretch Tyr-Ile-Ile-Val (TIIV) as required for efficient

focal accumulation of GFP (Fig. S3A and S3B). Insertion of this sequence motif between GFP

and a canonical SKL containing PTS1 motif induced enrichment of GFP inside of

subpopulations of peroxisomes (Fig. 3D). Fusion of the TIIV-tetrapeptide to the C-terminus of

a PTS2-GFP reporter protein (Ast et al., 2022) resulted in a similar pattern of localization (Fig.

3E and 3F). This suggests, that focal accumulation does not involve a specific pathway for

protein import but probably requires events that occur within the organelle.

To address if a TIIV motif does provoke subdomain formation in a different cellular

compartment or if it only functions inside of peroxisomes we attached it to the C-terminus of a

mitochondrial reporter protein containing an N-terminal mitochondrial targeting signal. This

chimeric fusion protein was detected inside of mitochondria in distinct foci suggesting that the

TIIV peptide triggers subdomain formation independent of the targeted organelle (Fig. 3G).

Figure 3 A short peptide motif sufficient for accumulation of proteins inside the core structure
(A) C-terminal dodecamers of Mac1 and Mac3 containing the characteristic PTS1 motifs. Cells were analyzed by super-resolution microscopy using SIM. Representative images of cells expressing GFP-PTS1$_{Mac1}$ and mCherry-SKL (B) or GFP-PTS1$_{Mac3}$ and mCherry-SKL. (C) Full images are shown as overlays of the green and red channel. For insets single channels and merged channels are depicted. Associated graphs show the normalized fluorescence intensity of GFP and mCherry along the indicated lines. (D) GFP fused to a peptide consisting of the small motif Tyr-Ile-Ile-Val (TIIV) and the prototype PTS1 Ser-Lys-Leu (SKL) was co-expressed with mCherry-SKL. A representative epifluorescence image is shown. Organization of the picture as above. (E) Cells expressing a PTS2-GFP reporter proteins either with or without a C-terminal TIIV motif were co-expressed with mCherry-SKL. Representative epifluorescence images are shown. Organization of the pictures as above. (F) Quantification show Pearson's correlation coefficients of GFP and mCherry signals for indicated strains. (G) GFP containing a mitochondrial targeting signal (MTS) either with or without an additional C-terminal TIIV motif was co-expressed with a MTS-mCherry. Representative epifluorescence images are shown. Organization of the pictures as above. ((A) – (E) and (G) Signals of GFP-tagged proteins are shown in cyan and of mCherry-tagged proteins in magenta. Scale bars: 5 µm.

Identification of additional proteins enriched in the core structure

Intrigued by the identification of a short peptide motif that triggers concentration of GFP inside of intraperoxisomal foci we asked whether this could help uncovering additional protein residents of the peroxisomal core structure.

Figure 4 Identification of additional PTS1 proteins enriched in the peroxisomal core structure
(A) Two candidate proteins containing TIIV-like motifs – D-amino oxidase 1 (Dao1) and malate synthase 1 (Mls1) – were co-expressed with mCherry-SKL. Representative epifluorescence images are shown. Full images are provided as overlays of the green and red channel. For insets single channels and merged channels are depicted.
(B) Two candidate proteins without TIIV-like motif – a putative NADPH dependent beta-ketoacyl reductase (Bkr1) and a putative enoyl-CoA hydratase (Enh1) – were co-expressed with mCherry-SKL. Representative epifluorescence images are shown. (C) Quantification shows the mean value of Pearson's correlation coefficients of GFP signals (candidate proteins) and mCherry-Signal signals (mCherry-SKL). Quantifications for three replicates for each of the analyzed candidate proteins are shown in Fig. S4. (D) Co-localization of indicated fusion protein was analyzed by SIM. Organization of pictures as above. Associated graphs show the normalized fluorescence intensity of GFP and mCherry along the indicated lines.
(A, B and D) Signals of GFP-tagged proteins are shown in cyan and of mCherry-tagged proteins in magenta. Scale bars: 5 μm.

We mined the predicted *U. maydis* proteome for candidates, which contain a TIIV motif or a similar sequence in combination with a PTS1 (Tab. S1). We tested seven full-length candidate proteins fused to GFP. Localization of six candidates resembled the so far identified proteins

Mac1, Mac3 and Uox1. The GFP signal accumulated within peroxisomal foci and in organelle

subpopulations (Fig. 4A and 4C; Fig. S4A). This strengthens our notion that the TIIV-like

motifs are involved in concentrating proteins into the core structure. Proteins from a control set

of randomly chosen PTS1-containing proteins that lack a TIIV-like motif on average showed a

higher co-localization with mCherry-SKL (Fig. 4B and 4C, Fig. S4B). The variability of

correlation coefficients, however, suggests that other features in the proteins contribute focal

enrichment in the core structure and formation of subpopulations as well.

Together, these data demonstrate that a significant number of peroxisomal matrix proteins

accumulates in peroxisomal foci. Other proteins have no or minor affinity to these structures or

are unable to form them (Fig. 4). The variable ability of proteins to form or accumulate in

subdomains likely represents a basic regulatory principle, which determines the luminal

distribution of peroxisomal matrix proteins. Among the enriched proteins were enzymes

involved in different metabolic pathways. These include the predicted D-amino oxidase Dao1

(UMAG_05703) and the malate synthase Mls1 (UMAG_15004) involved in glyoxylate

metabolism (Fig. 4A). Both proteins are also typical residents of peroxisomes in other

organisms (Kunze and Hartig, 2013; Pollegioni et al., 2007), but are not necessarily connected

in a metabolic network. Accordingly, enrichment of a particular protein does not coincide with

a specific peroxisomal pathway, but rather seems to be a feature shared by peroxisomal proteins

with very different biochemical functions. Interestingly, although Mac1, Mac3 and Mls1 are

detected in overlapping peroxisomal subdomains and enrich in similar subpopulations they

appear to be concentrated inside of different regions of the structures and may even exclude

each other to some extent (Fig. 4D), suggesting internal heterogeneity and a complex intrinsic

organization of the core structures.

Peroxisome subpopulations form over time

Enrichment of proteins in peroxisomal subdomains could result in their asymmetric distribution

when peroxisomes divide. This may support the formation of peroxisome subpopulations. To

follow GFP-Mac3 localization over time, GFP-Mac3 was expressed under control of an

arabinose inducible and glucose repressible promoter (Brachmann et al., 2001). As a control,

GFP fused to acyl-CoA oxidase 1 (Aox1; UMAG_02208) was used. GFP-Aox1 was not

enriched inside of the core, but uniformly distributed among peroxisomes upon constitutive

expression (Fig. S5A). Induction of both GFP fusion proteins (GFP-Mac3 or GFP-Aox1) in

strains that also express mCherry-SKL, was accomplished through incubation in arabinose

containing medium (Fig. 5A).

Figure 5 Accumulation in core structures and formation of subpopulations over time.
(A) Scheme of the transcriptional pulse-chase experiment to assess the formation of subdomains and
subpopulations over time. Cells were incubated in arabinose medium for 2 hours to induce expression of GFP-
tagged proteins. Expression was stopped via change to glucose containing medium. (B) Quantification of
colocalization data derived from epifluorescence microscopy depicted in Fig. S5. Each circle represents one cell.
(C and D). Cells were analyzed at indicated time points by super-resolution microscopy using SIM. GFP-Mac3
(C; cyan) was compared to GFP-Aox1 (D; cyan) – a protein not accumulating in the core structure upon
constitutive expression (Fig. S5) – in strains containing mCherry-SKL (magenta). Full images are shown as
overlays of the green and red channel. For insets single channels and merged channels are depicted.
260

After repression of gene expression by shifting cells to glucose medium we followed the

localization of the fluorescent proteins at different time points by SIM and epifluorescence

microscopy (Fig. 5A to 5D). Quantifications of epifluorescence images revealed that GFP-

Mac3 containing subpopulations form over time (Fig. 5B and S5B). Both GFP-Mac3 and GFP-

Aox1 were first uniformly distributed among peroxisomes (Fig. 5C and 5D). At later time

points, GFP-Mac3, but not GFP-Aox1, accumulated in distinct foci within peroxisomes

demonstrating formation of subdomains (Fig. 5C and 5D).

Specialization of peroxisome populations

We reasoned that the propensity of matrix proteins to self-assemble may control the protein

content of specialized peroxisomes. This is reminiscent to the biogenesis of peroxisome derived

Woronin bodies in filamentous ascomycetes (Jedd and Chua, 2000).

Figure 6 The Woronin body protein HexA colocalizes with Mac3
(A) *U. maydis* strains expressing an N-terminally GFP-tagged version of *A. nidulans* HexA (cyan) and mCherry-SKL (magenta) or mCherry-Mac3 (magenta) were inspected by epifluorescence microscopy. Full images are shown as overlays of the green and red channel. For insets single channels and merged channels are depicted. (B) Quantification shows Pearson's correlation coefficients of GFP and mCherry signals for indicated strains. (C) Crude organelle preparations of indicated strains were imaged by epifluorescence microscopy after incubation in lysis buffer supplemented with Triton X-100 (right). Preparations incubated in lysis buffer without Triton X-100 served as control (left). Organization of pictures as described above.

These specialized peroxisome derived compartments are enriched in a protein dubbed Hex1 in

N. crassa, which forms crystalline structures resistant to detergent (Jedd and Chua, 2000; Liu

et al., 2011, 2008; Yuan et al., 2003). We hypothesized that subdomain formation in *U. maydis*

and Woronin body formation represent special adaptations of a more generic mechanism that

regulates formation of distinct peroxisome populations. We, therefore, fused GFP to the

Aspergillus nidulans Woronin body protein HexA (Momany et al., 2002; Zekert et al., 2010)

and co-expressed GFP-HexA in *U. maydis* together with either mCherry-Mac3 or mCherry-

SKL. mCherry-Mac3 and GFP-HexA were detected in highly overlapping foci whereas
mCherry-SKL and GFP-HexA showed a lower degree of co-localization (Figs. 6A and 6B).
mCherry-Mac3 and GFP-HexA also appeared in stable and overlapping structures following
treatment of crude organelle extracts with detergent (Fig. 6C). This data supports the concept
that formation of luminal peroxisomal subdomains in *U. maydis* and formation of Woronin
bodies in ascomycetes follows similar mechanistic principles.
298

TIIV dependent accumulation in peroxisomal subdomains of mammalian cells

300   Finally, we addressed transferability of our finding to a different biological system and
301   expressed GFP combined with a minimal PTS1 ending on SKL and containing the TIIV motif
302   in human retinal pigment epithelial 1 (RPE1) cells.
303

304
305

Figure 7 Similar mechanistic principles for enrichment in the core of *U. maydis* and mammalian peroxisomes

(A) Constructs expressing GFP-TIIV-SKL and mCherry-SKL were transfected into RPE1 cells. Cells were analyzed by SIM. The representative picture shows an overlay of the GFP-signal (cyan) and the mCherry-signal (magenta) in the overview. The signal of single channels as well as the overlay are depicted for magnified insets. (B) *Mus musculus* urate oxidase (*Mm*UOX1) was tagged with mCherry (magenta) and co-expressed with GFP-TIIV-SKL (cyan) in RPE1 cells. Cells were analyzed by SIM. Representative pictures are organized as above. (C) *M. musculus* urate oxidase (*Mm*UOX1) was tagged with GFP (cyan) and expressed in *U. maydis* cells either co-expressing mCherry-SKL (magenta) or mCherry-Mac3 (magenta). Representative pictures are organized as above. (D) Quantifications show Pearson's correlation coefficients of GFP and mCherry signals for indicated strains. (E) Crude organelle preparations of indicated strains were imaged by epifluorescence microscopy after incubation in lysis buffer supplemented with Triton X-100 (right). Preparations incubated in lysis buffer without Triton X-100 served as control (left). Organization of pictures as described above. Scale bars: 5 μm.

mCherry-SKL was again used as marker for peroxisomes and subcellular localization of fluorescent proteins was analyzed by SIM. We obtained similar data as in *U. maydis* and observed accumulation of GFP-TIIV-SKL in defined subdomains of peroxisomes in RPE1 cells (Fig. 7A). To test if these structures are also enriched for a protein found in the native paracrystalline core of mammalian peroxisomes we co-expressed GFP-TIIV-SKL together with mCherry fused to murine urate oxidase 1 (*Mm*UOX1). GFP-TIIV-PTS1 showed a profound co-localization with mCherry-*Mm*UOX1 revealing that both proteins are located in vicinity (Fig. 7B). Thus, the rules underlying protein concentration inside of the paracrystalline core are likely similar in fungi and in humans. To further confirm this theory we performed complementary experiments in *U. maydis* cells and expressed GFP fused to murine *Mm*UOX1 together with either mCherry-SKL or mCherry-Mac3. Again, we detected a higher degree of overlap for GFP-UOX1 and mCherry-Mac3 than for GFP-*Mm*UOX1 and mCherry-SKL (Fig. 7C). In addition, we could observe stability of GFP-*Mm*UOX1 containing structures upon detergent treatment of crude organelle extracts (Fig. 7D). Both results are consistent with our model, that in diverse eukaryotic species formation of the core structure of peroxisomes follows analogous principles and is a major driving force to spatially organize the peroxisomal proteome.

Discussion

Here, we have shown selective accumulation of various proteins in subdomains of peroxisomes likely identical to the paracrystalline core structure. This process turned out as a driving force for the generation of organelle subpopulations with different protein content. Remarkably, independent of the original organism or biochemical function all so far tested proteins enriched in these structures show a high degree of co-localization in *U. maydis* and in human cells (Figs. 1 – 4, Fig. 7). Even the very specialized protein HexA from *A. nidulans* exhibited a very similar pattern of localization suggesting that selective enrichment of this peculiar peroxisomal protein

required for Woronin body biogenesis (Jedd & Chua, 2000) follows parallel mechanistic

principles (Fig. 6).

The variable ability to self-assemble inside of the lumen of the peroxisome may be a

fundamental principle to determine the proteome of every single peroxisome. Peroxisomes

proliferate by fission, constantly divide but in contrast to other organelles may not regularly

fuse at least in their mature form (Islinger et al., 2012). Thus, the asymmetric segregation of the

core will lead to peroxisome subpopulations with different protein content after any fission

event. Combined with altered protein expression levels this will almost certainly derive many

unique peroxisome variants within a single cell. Furthermore, peroxisomes can accumulate

huge amounts of cargo protein (DeLoache et al., 2016; Grewal et al., 2021) and the ability of

proteins to accumulate in the core structure may be one prerequisite or even a catalyzer.

Peroxisomes are attractive organelles for implantation of synthetic pathways (DeLoache &

Dueber, 2013; Simon et al., 2020; Stehlik et al., 2014). Forced concentration of enzymes by

addition of TIIV-like motifs might be another way to optimize or to modulate these pathways.

Protein crystallization for structural determination purposes (Schönherr et al., 2018) might

benefit from our finding as well.

Our data provide first insights into a novel code, which will evolve our understanding of the

spatial organization of the peroxisomal proteome. The identification of sequence motifs

sufficient for focal accumulation is a starting point to systematically deduce structural

differences between proteins that tend to exhibit a homogenous luminal peroxisomal

distribution or reside in peroxisomal subdomains. The core structure is, most probably,

metabolically active as it is a major site of a number of abundant enzymes including catalase,

urate oxidase or alcohol oxidase of methylotrophic yeasts (Hayashi et al., 1976; Veenhuis et

al., 1981; Völkl et al., 1988; Heinze et al., 2000; Schönherr et al., 2018). Given the surprising

variability of proteins with the ability to enrich in this structure (Fig. 3), its internal organization

is of considerable interest. Can proteins go in and out regularly? Are their different layers of

enrichment for particular proteins as indicated by our data for acyltransferases and malate

synthase 1 from *U. maydis* (Fig. 4)? Is there biochemical activity everywhere in the core? The

process of formation may even exhibit functional parallels with phase separation of proteins

(Boeynaems et al., 2018; Hyman et al., 2014).

In summary, our work contributes to a more detailed characterization of a conserved

peroxisomal structure, which is known since the discovery of the peroxisome (De Duve &

Baudhuin, 1966; Rouiller & Bernhard, 1956) but whose formation and composition has not

been addressed in greater detail until now.

Methods

Microorganism, growth conditions and transformation

Escherichia coli strain TOP10 (Invitrogen) was used for all cloning purposes and amplification

of plasmid DNA. *U. maydis* strains generated and used in this study are derivatives of Bub8,

MB215 and Bub8 mCherry-SKL (Freitag et al., 2012; Hewald et al., 2005; Weinzierl et al.,

2002) and listed in Supplementary Table 2 (Tab. S2). All strains are available from the authors

upon reasonable request. *U. maydis* cells were grown at 28°C in YEPSlight medium (Tsukuda

et al., 1988) or yeast nitrogen base medium (YNB; Difco) supplemented either with 2% glucose

and 0.2% ammonium sulfate or 0.2% oleic acid (Roth), 0.1% Tween-40 and 0.2% ammonium

sulfate at pH 5.6. For epifluorescence imaging cells were grown in oleic acid media if not

specified differently. SIM imaging was performed on glucose grown cells to circumvent artifact

of oleic acid addition. For chase experiments (Fig. 5) *U. maydis* cells were grown overnight in

YEPSlight medium. Subsequently, cells were washed and diluted in YNB medium containing

2% arabinose and 0.2% ammonium sulfate and incubated for 2 h. Expression of fusion proteins

was stopped by shifting the cells to YNB medium containing 2% glucose and 0.2 % ammonium

sulfate cells and the localization of the fluorescent proteins was microscopically monitored at

indicated time points. Cell transformation was performed as previously described (Hanahan et

al., 1991; Schulz et al., 1990). DNA was integrated into the *ip*-locus of *U. maydis* cells

(Broomfield & Hargreaves, 1992) or randomly integrated into the genome. Hygromycin (200

µg/ml), carboxin (2 µg/ml) and G418 (400 µg/ml) were used for selection. Genomic DNA was

extracted as previously described (Hoffman and Winston, 1987).

Human cell line and transient transfection of cells

hTert-RPE1 cells (ATCC CRL-4000) were cultured in DMEM-F12 medium (Life

Technologies) supplemented with 10% FCS and Penicillin/Streptomycin (100 U/mL penicillin,

100 µg/mL streptomycin) at 37°C, 5% CO_2. Cells were transiently transfected with the help of

Fugene 6 (Promega) using a ratio of 3:1 of reagent to DNA. Cells were transfected in MatTek

3.5 cm glass-bottom dishes. 12 h after transfection, the medium was replaced to remove

transfection reagent, and the cells were imaged between 16 h and 20 h post transfection**.**

Molecular cloning and nucleic acid procedures

Standard procedures were followed for plasmid generation (Sambrook et al., 1989). Plasmids

used for protein expression in *U. maydis* are derivatives of the plasmids pOTEF-GFP-Ala6-

MMXN and pCRG-GFP-Ala6-MXN (Böhmer et al., 2008). Plasmids used for expression of

fluorescent proteins in hTert-RPE1 cells are derivatives of the plasmid pEGFP-C1 and

pmCherry-C1 (Clontech). Plasmids were analyzed by sequencing. All plasmids and

oligonucleotides used in this study are listed in Supplementary table S2 (Tab. S2). Plasmids are

available from the authors upon reasonable request.

Epifluorescence microscopy

Logarithmically grown *U. maydis* cells were treated with 100 µM Carbonyl cyanide m-

chlorophenyl hydrazine (CCCP) resolved in dimethyl sulfoxide to block peroxisomal

movement. For the majority of experiments using epifluorescence microscopy cells were

incubated in oleic acid-containing media for four hours prior to imaging experiments as

formation of subpopulations was easier to follow under these conditions (exceptions: imaging

experiments underlying Figs. 1A and S5). After 5-10 minutes incubation at room temperature,

the cells were either placed on 1% agarose cushions containing 100 µM Carbonyl cyanide m-

chlorophenyl hydrazine CCCP on a microscope glass slide and covered with a coverslip or

placed on MatTek 3.5 cm glas-bottom dishes and covered with an 1% agarose cushion

containing 100 µM CCCP as well. For time-lapse imaging no CCCP was added.

Epifluorescence microscopy was performed on an Axiovert 200 M inverse microscope (Zeiss)

equipped with a 1394 ORCA ERA CCD camera (Hamamatsu Photonics), filter sets for eGFP

and rhodamine, and a Zeiss 63x Plan Apochromat oil lens (NA 1.4). The software Volocity 5.3

(Perkin-Elmer) was used for image acquisition. The ImageJ plugin DeconvolutionLab with 25

iterations of the Richardson-Lucy algorithm was used for image deconvolution (Vonesch and

Unser, 2008), point-spread functions for each channel were computed by the ImageJ plugin

PSF Generator using the Richards & Wolf model (Kirshner et al., 2013). Alternatively,

epifluorescence microscopy as well as time-lapse experiments (500 ms time-lapse for 60 s)

were performed on a Deltavision OMX V4 microscope (GE Healthcare) equipped with three

water-cooled PCO edges CMOS cameras, a solid-state light source and a laser-based autofocus.

The software softWoRx (Applied Precision, GE Healthcare) was used for image deconvolution.

Images were processed in ImageJ/Fiji (Schneider et al., 2012).

SIM

Three-dimensional structured illumination microscopy (SIM) was performed on a Deltavision

OMX V4 microscope with and Olympus x60 NA 1.42 objective and three PCO.edge sCMOS

cameras and 488 nm and 568 nm laser lines. Cells were illuminated with a grid pattern. 15 raw

images (3 orientations, 5 phases) were taken for each image plane. SoftWoRx software

(Applied Precision, GE Healthcare) was used for image reconstruction, alignment and
projection. Images were processed in ImageJ/Fiji (Schneider et al., 2012). For 3D visualization
of peroxisomal subdomains, surfaces were generated using Imaris.
450

Transmission electron microscopy and Immunogold labelling

For visualization of *U. maydis* cells expressing GFP-Mac3 using transmission electron
microscopy (TEM) 50 ml logarithmically growing cells were harvested and frozen under high-
pressure (Wohlwend HPF Compact 02). After subsequent freeze substitution (using acetone,
containing 0.25% osmium tetroxide, 0.2% uranyl acetate, 0.05% ruthenium red, and 5% water)
(Leica AFS2), cells were embedded in Epon812 substitute resin (Fluka). Embedded cells were
sectioned to 50 nm thin sections (Leica EM UC7 RT), which were used for immunolabeling
with α-GFP (Rockland; dilution 1:500). As a secondary antibody rabbit-anti-goat antibodies
coupled to ultra-small gold particles were used (Aurion, dilution 1:100). Subsequently a silver
enhancement procedure was performed and sections were post-stained with 2% uranyl acetate
and 0.5% lead citrate. Analysis of the samples was conducted with a JEOL JEM2100 TEM
equipped with a fast-scan 2k CCD TVIPS (Gauting, Germany) F214 camera. Images were
processed with ImageJ (Schneider et al., 2012).
464

Preparation of crude organelles

Preparation of post-nuclear supernatants was adapted from previous protocols (Cramer et al.,
2015; Kremp et al., 2020; Stehlik et al., 2020). Briefly, 200 ml of logarithmically growing cells
were harvested for 5 min at 1600 x g at 23°C. Cells were washed two times with deionized
water and two times with sorbitol buffer (1.2 M sorbitol; 20 mM KH_2PO_4 adjusted to pH 7.4
with K_2HPO_4). Formation of spheroplasts was achieved through incubation in sorbitol buffer
containing novozyme (4μg/ml) for 30 – 45 min. Degradation of the cell wall was followed
microscopically. Afterwards, spheroplasts were kept on ice and gently washed two times with
sorbitol buffer – centrifugation for 10 min at 2300 rpm at 4°C. Spheroplasts were gently washed
with lysis buffer (5 mM MES, 0.5 mM EDTA, 1 mM KCl, 0.6 M Sorbitol, 1 mM 4-
aminobenzamidine dihydrochloride, 1 μg/ml aprotinin, 1 μg/ml leupeptin, 1 mM
phenylmethylsulfonyl fluoride, 10 μg/ml N-tosyl-L-phenylalanine chloromethyl ketone, and 1
μg/ml pepstatin) – centrifugation for 10 min at 2300 rpm at 4°C, resuspended in 15 ml lysis
buffer and frozen at -80°C overnight. Homogenization was achieved through 2 x 10 strokes
with a Potter-Elvehjem homogenizer interrupted by chilling the samples on ice for 2 min.

Nuclei and cell debris were removed by two subsequent centrifugations at 1600 × g for 10 min.

Subsequently, the PNS was diluted to an OD600 of 1, aliquoted and frozen at -80°C.

Detergent treatment

PNS fractions were thawed on ice, centrifuged at 10k*g for 10 min at 4°C and concentrated

10x in lysis buffer. Samples were split into halves and aliquots were incubated in lysis buffer

containing 1% Triton-X-100 or in lysis buffer for 45 min. Preparations were immediately

analyzed by epifluorescence microscopy immobilized on 1% agarose cushions.

Quantification of peroxisomal subpopulations and statistical analysis

Microscopic data was collected from at least three independent *U. maydis* cultures (biological

replicates) for each experiment. At least five images per culture with at least ten cells were

quantified. Quantification of co-localization was performed as follows. Detection and

measurement of individual peroxisomes was performed automatically via an ImageJ macro

(Supplementary File 1). Briefly, maximum projections of deconvolved RFP and GFP z-stacks

were created, transformed into 16-bit images and corrected for channel-misalignment by the

"StackReg 2.0.0" plugin using "Rigid Body" transformation (Thevenaz et al., 1998;

https://sites.imagej.net/BIG-EPFL/plugins/) to later serve as the reference images for detection

of peroxisomes; same operations were performed on average projections of the respective non-

deconvolved z-stacks for later fluorescence-intensity measurements. To increase sensitivity and

account for peroxisomes showing only fluorescence in one of the two channels, the resulting

images were merged using the Image Calculator function with the maximum option (detection

based on only RFP or GFP references was still included in the macro but performed less reliable

than using the merge of both references). For detection of peroxisomes, the merged average

projections were used to create a mask of the cells. This mask was used to determine the

background noise outside of the cells in the reference merge. The resulting maximum noise

intensity was used for the Find Maxima function of ImageJ; individual peroxisomes were

detected in the reference merge and marked as point selections. These selections were restored

in the background-corrected average projections of the non-deconvolved stacks of both

channels. Subsequently, circular ROIs of a 3-pixel radius (area of 32 pixels) were created

around the point selections and signal intensities were measured for both channels (as Raw

Integrated Density per ROI) and stored side-by-side in the results file for the whole dataset. A

commented version of the macro is attached to the manuscript. For evaluation of the resulting

data, the statistics software R was used (R Core Team, 2015): measured intensities were

normalized to one and a linear regression analysis was performed with the RFP intensities as independent and GFP intensities as dependent variable, in addition a correlation test was performed (Pearson's). Superplots (Lord et al., 2020) and statistical tests were computed using RStudio 1.2.1335 with R 3.6.0 or GraphPadPrism. Plots are structured as follows: center line, mean; error bars, standard error of the mean; circles, mean of experiments. P-values were calculated using an unpaired, two-sided Student's t-test using Graphpad Prism. For data depicted in Fig. 1 and Fig. S3C, which contain multiple comparisons a 1-Way anova combined with a Tukeys post-test was performed to assess significance of the differences. * refers to a p-value lower than or equal to 0.05; ** refer to a p-value lower than or equal to 0.01 and ***lower than or equal to 0.001. ****lower than 0.0001. The graph shown in Fig. 5 was generated using Graphpad Prism. Each dot represents one cell.

Bioinformatics and accession numbers

U. maydis putative peroxisomal PTS1 protein sequences were retrieved from the *Ustilago maydis* genome database (MUMDB) or broad institute and manipulated with notepad ++ using regular expressions (regex). PTS1 motifs were bookmarked with the regular expression '([SA][RKHQNS][LI]|[SA][RK][MFV]|[PCVGE][RK][LI])*$'. Protein sequences containing a TIIV-like motif were bookmarked with the regular expression '(T [I, L, V] [I, L, V] [I, L, V] [I, L, V] [I, L, V])*$'. Peroxisomal candidate proteins with TIIV-like motif and without motif were randomly chosen and the localization of the peroxisomal proteins to subdomains was analyzed by fluorescence microscopy. All genes can now be accessed via the National Center of Biotechnology Information (NCBI) via the UMAG number.

Data availability

Any supporting original data for this study can be requested from the corresponding authors.

Acknowledgments

We thank Marisa Piscator and Ulrikke Dahl Brinch for excellent technical assistance. We are grateful to Reinhard Fischer and Valentin Wernet for plasmids. JA was supported by a fellowship from Marburg Research Academy. TS acknowledges funding from SYNMIKRO. KS was supported by a Career grant from the South-Eastern Norway Regional Health Authority (2020038) and a Research Grant from the Research Council of Norway (315103). JF acknowledges funding from the DFG (grant ID FR-3586/2-1).

Author contributions

Conceptualization: JA, DM, KS and JF; Data acquisition: JA, NB, DM, TH, BS, KS and JF.
Preparation of figures: JA, NB, DM and KS; Generation of code, bioinformatic and statistical
analysis: JA, NB, TS, CR, KS; Supervision: CT, GB, MB, KS, JF; Writing – original draft: JF;
Writing – revision and editing: All authors. Analysis of data: All authors.

Ast J, Bäcker N, Bittner E, Martorana D, Ahmad H, Bölker M, Freitag J. 2022. Two Pex5 Proteins With
Different Cargo Specificity Are Critical for Peroxisome Function in Ustilago maydis. *Front Cell Dev Biol*
10. doi:10.3389/fcell.2022.858084

Bartoszewska M, Opalinski L., Veenhuis M, van der Klei IJ. 2011. The significance of peroxisomes in
secondary metabolite biosynthesis in filamentous fungi. *Biotechnol Lett* **33**:1921–1931.

Baudhuin P, Beaufay H, de Duve C. 1965. Combined biochemical and morphological study of particulate
fractions from rat liver : Analysis of Preparations Enriched in Lysosomes or in Particles Containing Urate
Oxidase, D-Amino Acid Oxidase, and Catalase . *J Cell Biol* **26**:219–243. doi:10.1083/jcb.26.1.219

Becker F, Stehlik T, Linne U, Bölker M, Freitag J, Sandrock B. 2021. Engineering *Ustilago maydis* for
production of tailor-made mannosylerythritol lipids. *Metab Eng Commun* **12**:e00165.
doi:https://doi.org/10.1016/j.mec.2021.e00165

Boeynaems S, Alberti S, Fawzi NL, Mittag T, Polymenidou M, Rousseau F, Schymkowitz J, Shorter J, Wolozin
B, Van Den Bosch L, others. 2018. Protein phase separation: a new phase in cell biology. *Trends Cell Biol*
28:420–435.

Böhmer C, Böhmer M, Bölker M, Sandrock B. 2008. Cdc42 and the Ste20-like kinase Don3 act independently
in triggering cytokinesis in *Ustilago maydis*. *J Cell Sci* **121**:143–148.

Bölker M, Basse CW, Schirawski J. 2008. *Ustilago maydis* secondary metabolism—From genomics to
biochemistry. *Fungal Genet Biol* **45**:S88--S93.

Bortfeld M, Auffarth K, Kahmann R, Basse CW. 2004. The *Ustilago maydis* a2 mating-type locus genes *lga2*
and *rga2* compromise pathogenicity in the absence of the mitochondrial p32 family protein Mrb1. *Plant*
Cell **16**:2233-48.

Brachmann A, Weinzierl G, Kämper J, Kahmann R. 2001. Identification of genes in the bW/bE regulatory
cascade in *Ustilago maydis*. *Mol Microbiol* **42**:1047–1063.

Brocard C, Hartig A. 2006. Peroxisome targeting signal 1: is it really a simple tripeptide? *Biochim Biophys Acta*
1763:1565–73. doi:10.1016/j.bbamcr.2006.08.022

Broomfield PLE, Hargreaves JA. 1992. A single amino-acid change in the iron-sulphur protein subunit of
succinate dehydrogenase confers resistance to carboxin in Ustilago maydis. *Curr Genet* **22**:117–121.

Camões F, Islinger M, Guimarães SC, Kilaru S, Schuster M, Godinho LF, Steinberg G, Schrader M. 2015. New
insights into the peroxisomal protein inventory: Acyl-CoA oxidases and-dehydrogenases are an ancient
feature of peroxisomes. *Biochim Biophys Acta (BBA)-Molecular Cell Res* **1853**:111–125.

Cramer J, Effelsberg D, Girzalsky W, Erdmann R. 2015. Small-scale purification of peroxisomes for analytical
applications. *Cold Spring Harb Protoc*. doi:10.1101/pdb.prot083717

De Duve C. 1960. Intracellular Localization of Enzymes. *Nature* **187**:836. doi:10.1038/187836a0

De Duve C, Baudhuin P. 1966. Peroxisomes (microbodies and related particles). *Physiol Rev* **46**:323–357.

DeLoache WC, Dueber JE. 2013. Compartmentalizing metabolic pathways in organelles. *Nat Biotechnol*
31:320–321.

DeLoache WC, Russ ZN, Dueber JE. 2016. Towards repurposing the yeast peroxisome for compartmentalizing
heterologous metabolic pathways. *Nat Commun* **7**:11152. doi:10.1038/ncomms11152

Francisco T, Rodrigues TA, Dias AF, Barros-Barbosa A, Bicho D, Azevedo JE. 2017. Protein transport into
peroxisomes: knowns and unknowns. *BioEssays* **39**:1700047.

Freitag J, Ast J, Bölker M. 2012. Cryptic peroxisomal targeting via alternative splicing and stop codon read-

through in fungi. *Nature* **485**:522–525.

Freitag J, Ast J, Linne U, Stehlik T, Martorana D, Bölker M, Sandrock B. 2014. Peroxisomes contribute to
biosynthesis of extracellular glycolipids in fungi. *Mol Microbiol* **93**:24–36.

Galiani S, Waithe D, Reglinski K, Cruz-Zaragoza LD, Garcia E, Clausen MP, Schliebs W, Erdmann R, Eggeling
C. 2016. Super resolution microscopy reveals compartmentalization of peroxisomal membrane proteins. *J*
Biol Chem jbc--M116.

Gould SJ, Keller GA, Hosken N, Wilkinson J, Subramani S. 1989. A conserved tripeptide sorts proteins to
peroxisomes. *J Cell Biol* **108**:1657–1664.

Grewal PS, Samson JA, Baker JJ, Choi B, Dueber JE. 2021. Peroxisome compartmentalization of a toxic
enzyme improves alkaloid production. *Nat Chem Biol* **17**:96–103.

Gründlinger M, Yasmin S, Lechner BE, Geley S, Schrettl M, Hynes M, Haas H. 2013. Fungal siderophore
biosynthesis is partially localized in peroxisomes. *Mol Microbiol*. **88(5):**862-75

Gustafsson MGL. 2000. Surpassing the lateral resolution limit by a factor of two using structured illumination
microscopy. *J Microsc* **198**:82–87.

Hanahan D, Jessee J, Bloom FR. 1991. Plasmid transformation of Escherichia coli and other bacteria. *Methods*
Enzymol **204**:63–113.

Hayashi H, Taya K, Suga T, Niinobe S. 1976. Studies on Peroxisomes. VI. Relationship between the
Peroxisomal Core and Urate Oxidase1. *J Biochem* **79**:1029–1034.
doi:10.1093/oxfordjournals.jbchem.a131143

Heinze M, Reichelt R, Kleff S, Eising R. 2000. High Resolution Scanning Electron Microscopy of Protein
Inclusions (Cores) purified from Peroxisomes of Sunflower (Helianthus annuus L.) Cotyledons. *Cryst Res*
Technol **35**:877–886. doi:https://doi.org/10.1002/1521-4079(200007)35:6/7<877::AID-
CRAT877>3.0.CO;2-S

Hewald S, Josephs K, Bölker M. 2005. Genetic analysis of biosurfactant production in Ustilago maydis. *Appl*
Environ Microbiol **71**:3033–3040.

Hoffman CS, Winston F. 1987. A ten-minute DNA preparation from yeast efficiently releases autonomous
plasmids for transformaion of *Escherichia coli*. *Gene* **57**:267–272.

Hu J, Baker A, Bartel B, Linka N, Mullen RT, Reumann S, Zolman BK. 2012. Plant peroxisomes: biogenesis
and function. *Plant Cell* **24**:2279–2303.

Hyman AA, Weber CA, Jülicher F. 2014. Liquid-liquid phase separation in biology. *Annu Rev Cell Dev Biol*.
doi:10.1146/annurev-cellbio-100913-013325

Islinger M, Grille S, Fahimi HD, Schrader M. 2012. The peroxisome: an update on mysteries. *Histochem Cell*
Biol **137**:547–574.

Jedd G, Chua NH. 2000. A new self-assembled peroxisomal vesicle required for efficient resealing of the plasma
membrane. *Nat Cell Biol* **2**:226–231.

Kämper J, Kahmann R, Bölker M, Ma LJ, Brefort T, Saville BJ, Banuett F, Kronstad JW, Gold SE, Müller O,
others. 2006. Insights from the genome of the biotrophic fungal plant pathogen *Ustilago maydis*. *Nature*
444:97–101.

Kim PK, Hettema EH. 2015. Multiple pathways for protein transport to peroxisomes. *J Mol Biol* **427**:1176–
1190.

Kohlwein SD, Veenhuis M, van der Klei IJ. 2013. Lipid Droplets and Peroxisomes: Key Players in Cellular

Lipid Homeostasis or A Matter of Fat—Store'em Up or Burn'em Down. *Genetics* **193**:1–50.

Kremp M, Bittner E, Martorana D, Klingenberger A, Stehlik T, Bölker M, Freitag J. 2020. Non-AUG

Translation Initiation Generates Peroxisomal Isoforms of 6-Phosphogluconate Dehydrogenase in Fungi.

Front Cell Dev Biol **8**:251. doi:10.3389/fcell.2020.00251

Kumar S, de Boer R, van der Klei IJ. 2018. Yeast cells contain a heterogeneous population of peroxisomes that

segregate asymmetrically during cell division. *J Cell Sci* **131**:jcs207522.

Kunze M, Hartig A. 2013. Permeability of the peroxisomal membrane: lessons from the glyoxylate cycle. *Front*

Physiol **4**:204.

Lazarow PB. 2006. The import receptor Pex7p and the PTS2 targeting sequence. *Biochim Biophys Acta (BBA)-*

Molecular Cell Res **1763**:1599–1604.

Lingard MJ, Monroe-Augustus M, Bartel B. 2009. Peroxisome-associated matrix protein degradation in

Arabidopsis. *Proc Natl Acad Sci U S A*. doi:10.1073/pnas.0811329106

Liu F, Lu Y, Pieuchot L, Dhavale T, Jedd G. 2011. Import oligomers induce positive feedback to promote

peroxisome differentiation and control organelle abundance. *Dev Cell*. doi:10.1016/j.devcel.2011.08.004

Liu F, Ng SK, Lu Y, Low W, Lai J, Jedd G. 2008. Making two organelles from one: Woronin body biogenesis

by peroxisomal protein sorting. *J Cell Biol* **180**:325–339.

Lord SJ, Velle KB, Mullins RD, Fritz-Laylin LK. 2020. SuperPlots: Communicating reproducibility and

variability in cell biology. *J Cell Biol* **219**. doi:10.1083/jcb.202001064

Momany M, Richardson EA, Van Sickle C, Jedd G. 2002. Mapping Woronin body position in *Aspergillus*

nidulans. *Mycologia* **94**:260–266.

Nguyen TA, Le S, Lee M, Fan J-S, Yang D, Yan J, Jedd G. 2021. Fungal Wound Healing through Instantaneous

Protoplasmic Gelation. *Curr Biol* **31**:271-282.e5.

Pieuchot L, Jedd G. 2012. Peroxisome assembly and functional diversity in eukaryotic microorganisms. *Annu*

Rev Microbiol **66**:237–263.

Poirier Y, Antonenkov VD, Glumoff T, Hiltunen JK. 2006. Peroxisomal β-oxidation—a metabolic pathway with

multiple functions. *Biochim Biophys Acta* **1763**:1413–1426.

Pollegioni L, Piubelli L, Sacchi S, Pilone MS, Molla G. 2007. Physiological functions of D-amino acid oxidases:

from yeast to humans. *Cell Mol Life Sci* **64**:1373–1394. doi:10.1007/s00018-007-6558-4

Rouiller C and Bernhard W. 1956 "Microbodies" and the problem of mitochondrial regeneration in liver cells. J

Biophys Biochem Cytol Suppl. 2: 355-359.

Sambrook J, Fritsch EF, Maniatis T. 1989. Molecular Cloning: A Laboratory Manual, Cold Spring Harbor

Laboratory, Molecular Cloning a laboratory manual. Cold Spring Harbor Laboratory Press.

doi:10.1128/AEM.71.8.4602

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nat*

Methods. doi:10.1038/nmeth.2089

Schönherr R, Rudolph JM, Redecke L. 2018. Protein crystallization in living cells. *Biol Chem* **399**:751–772.

doi:doi:10.1515/hsz-2018-0158

Schulz B, Banuett F, Dahl M, Schlesinger R, Schäfer W, Martin T, Herskowitz I, Kahmann R. 1990. The b

alleles of *U. maydis*, whose combinations program pathogenic development, code for polypeptides

containing a homeodomain-related motif. *Cell* **60**:295–306.

Simon D, Thomas WW, Yixuan L, Codruta I, C. KS. 2020. Transforming yeast peroxisomes into microfactories

for the efficient production of high-value isoprenoids. *Proc Natl Acad Sci* **117**:31789–31799.

doi:10.1073/pnas.2013968117

Smith JJ, Aitchison JD. 2013. Peroxisomes take shape. *Nat Rev Mol Cell Biol* **14**:803–817.

Stehlik T, Kremp M, Kahnt J, Bölker M, Freitag J. 2020. Peroxisomal targeting of a protein phosphatase type 2C

via mitochondrial transit. *Nat Commun* **11**:2355. doi:10.1038/s41467-020-16146-3

Stehlik T, Sandrock B, Ast J, Freitag J. 2014. Fungal peroxisomes as biosynthetic organelles. *Curr Opin*

Microbiol **22**:8–14. doi:http://dx.doi.org/10.1016/j.mib.2014.09.011

Tey WK, North AJ, Reyes JL, Lu YF, Jedd G. 2005. Polarized Gene Expression Determines Woronin Body

Formation at the Leading Edge of the Fungal Colony. *Mol Biol Cell* **16**:2651–2659.

Thevenaz P, Ruttimann UE, Unser M. 1998. A pyramid approach to subpixel registration based on intensity.

IEEE Trans Image Process **7**:27–41. doi:10.1109/83.650848

Tsukada H, Mochizuki Y, Fujiwara S. 1966. The nucleoids of rat liver cell microbodies fine structure and

enzymes. *J Cell Biol* **28**:449–460.

Tsukuda T, Carleton S, Fotheringham S, Holloman WK. 1988. Isolation and characterization of an

autonomously replicating sequence from Ustilago maydis. *Mol Cell Biol* **8**:3703–3709.

van der Klei IJ, Veenhuis M. 2006. Yeast and filamentous fungi as model organisms in microbody research.

Biochim Biophys Acta **1763**:1364–1373.

Veenhuis M, Harder W, van Dijken JP, Mayer F. 1981. Substructure of crystalline peroxisomes in methanol-

grown Hansenula polymorpha: evidence for an in vivo crystal of alcohol oxidase. *Mol Cell Biol* **1**:949–

957.

Völkl A, Baumgart E, Fahimi HD. 1988. Localization of urate oxidase in the crystalline cores of rat liver

peroxisomes by immunocytochemistry and immunoblotting. *J Histochem \& Cytochem* **36**:329–336.

doi:10.1177/36.4.3346536

Völkl A, Mohr H, Fahimi H. D. 1999. Peroxisome Subpopulations of the Rat Liver: Isolation by Immune Free

Flow Electrophoresis. *J Histochem Cytochem* **47**:1111–1117. doi:10.1177/002215549904700902

Walter T, Erdmann R. 2019. Current Advances in Protein Import into Peroxisomes. *Protein J* **38**:351–362.

doi:10.1007/s10930-019-09835-6

Wanders RJA. 2014. Metabolic functions of peroxisomes in health and disease. *Biochimie* **98**:36–44.

Weinzierl G, Leveleki L, Hassel A, Kost G, Wanner G, Bölker M. 2002. Regulation of cell separation in the

dimorphic fungus *Ustilago maydis*. *Mol Microbiol* **45**:219–231.

Yuan P, Jedd G, Kumaran D, Swaminathan S, Shio H, Hewitt D, Chua N-H, Swaminathan K. 2003. A HEX-1

crystal lattice required for Woronin body function in *Neurospora crassa*. *Nat Struct Mol Biol* **10**:264–270.

doi:10.1038/nsb910

Zekert N, Veith D, Fischer R. 2010. Interaction of the *Aspergillus nidulans* Microtubule-Organizing Center

(MTOC) Component ApsB with Gamma-Tubulin and Evidence for a Role of a Subclass of Peroxisomes in

the Formation of Septal MTOCs. *Eukaryot Cell* **9**:795–805. doi:10.1128/EC.00058-10

Supplementary figures

Figure S1 Examples for the quantification of co-localization
Representative plots showing the fluorescence signals of all analyzed peroxisomes in a representative replicate.
(A) related to Fig. 1C (B) related to Fig. 1F.

A GFP-Mac3 mCherry-SKL

0s   19s

B GFP-Uox1 mCherry-Mac1

C GFP-Uox1 mCherry-Mac3   GFP-Mac1 mCherry-Mac3

Figure S2 Mac proteins localize in detergent resistant peroxisomal subdomains also enriched for urate
oxidase 1
(A) Time lapse imaging related to Movie 1. Epifluorescence imaging of GFP-Mac3 (cyan) and mCherry-SKL
(magenta). One frame per second was recorded. Shown are overlays of the mCherry and GFP signal. (B) A
representative image of cells expressing GFP-Uox1 and mCherry-Mac1. Full images are shown as overlays of the
green and red channel. For insets single channels and merged channels are depicted. Scale bar: 5 μm. (C) Crude
organelle preparations of indicated strains were imaged by epifluorescence microscopy after incubation in lysis
buffer supplemented with Triton X-100 (right). Preparations incubated in lysis buffer without Triton X-100 served
as control (left). Organization of pictures as described above. Scale bars: 5 μm.

A

B

Figure S3 Truncation analysis of the C-terminal dodecamer of Mac3
(A) Depiction of the 3'end of the *mac3* gene encoding the peroxisomal targeting signal PTS1. Please note that the unusual ASL motif had to be substituted by SKL to perform the subsequent truncation analysis as truncated fragment of the dodecamer did not allow full peroxisomal import, which perturbs co-localization analysis (B) Indicated fragments of the C-terminal dodecamer were fused to GFP (cyan) and analyzed in strains also expressing mCherry-SKL (magenta). Full representative images are shown as overlays of the green and red channel. For insets single channels and merged channels are depicted. Scale bars: 5 μm. (C) Quantification of co-localization of experiments shown in (B).

Figure S4 Analysis of candidate proteins with or without TIIV-like motif
U. maydis strains expressing N-terminally GFP-tagged versions of candidate proteins (cyan) with TIIV like motifs
(A) or without TIIV-like motifs (C) and the peroxisomal marker protein mCherry-SKL (magenta) were inspected
by epifluorescence microscopy. Full representative images are shown as overlays of the green and red channel.
For insets single channels and merged channels are depicted. Scale bars: 5 µm. Umag numbers are gene identifiers
accessible via NCBI. (B) Quantifications show Pearson's correlation coefficients of GFP and mCherry signals for
indicated strains.

A

GFP-Aox1 mCherry-SKL

B

GFP-Mac3 mCherry-SKL

GFP-Aox1 mCherry-SKL

748
749
Fig. S5 Formation of peroxisome subpopulations over time.
(A) *U. maydis* strains expressing an N-terminally GFP-tagged version of Aox1 (cyan) and the peroxisomal marker
protein mCherry-SKL (magenta) were inspected by epifluorescence microscopy. Full images are shown as
overlays of the green and red channel. For insets single channels and merged channels are depicted. Scale bar: 5
µm. **(B)** Cells were analyzed at indicated time points after addition of glucose by epifluorescence microscopy.
GFP-Mac3 (cyan) was compared to GFP-Aox1 (cyan) in strains containing mCherry-SKL (magenta). Full images
are shown as overlays of the green and red channel. For insets single channels and merged channels are depicted.
Quantifications of (B) are shown in Fig. 5.