Leaf hydraulic maze; Differential effect of ABA on bundle-sheath, palisade, and spongy mesophyll hydraulic conductance

Adi Yaaran[1], Eyal Erez[1], Carl Procko[2] and Menachem Moshelion[1]*

[1] Institute of Plant Sciences and Genetics in Agriculture, The Robert H. Smith Faculty of Agriculture, Food and Environment, The Hebrew University of Jerusalem, Rehovot 76100, Israel.

[2] Plant Biology Laboratory, Salk Institute for Biological Studies, La Jolla, California 92037, USA.

*Corresponding author:

Menachem Moshelion,

The Robert H. Smith Faculty of Agriculture, Food and Environment,

The Hebrew University of Jerusalem,

Rehovot, Israel

E-mail: menachem.moshelion@mail.huji.ac.il

Keywords

ABA, leaf hydraulics, spongy mesophyll, palisade mesophyll, bundle sheath, P_f

Running title: Differential effect of ABA on leaf hydraulic conductance

ABSTRACT 21

Leaf hydraulic conductance (K_{leaf}) facilitates the movement of water for transpiration, 22

enabling continual CO_2 uptake while the plant maintains its water status. We 23

hypothesized that bundle-sheath and mesophyll cells play key roles in regulating the 24

radial flow of water out of the xylem under optimal and stress conditions. To examine 25

that hypothesis, we generated transgenic *Arabidopsis* plants that were insensitive to 26

abscisic acid (ABA) in their bundle sheath (BSabi) or mesophyll (MCabi) cells. Both 27

BSabi and MCabi plants showed greater K_{leaf} and transpiration under optimal 28

conditions. Yet, the stomatal apertures, stomatal indices and vein densities of the BSabi 29

plants were similar to those of WT plants. MCabi plants had larger stomatal apertures, 30

a higher stomatal index and greater vascular diameter and biomass, relative to the WT 31

and BSabi. In response to xylem-fed ABA, both transgenic and WT plants reduced their 32

K_{leaf} and transpiration. However, leaf water potential was reduced only in the WT. The 33

membrane osmotic water permeability (P_f) of the WTs' spongy mesophyll was higher 34

than that of its palisade mesophyll. Moreover, only the spongy cells reduced their P_f in 35

response to ABA. ABA-insensitive spongy mesophyll cells had a low P_f; whereas 36

ABA-insensitive bundle-sheath cells had a higher P_f. Palisade cells maintained a low 37

P_f at all ABA levels. ABA increased the symplastic water pathway, but its contribution 38

to K_{leaf} was negligible. We suggest that the bundle sheath–spongy mesophyll pathway 39

may control K_{leaf} to maintain steady-state conditions in the palisade cells and optimal 40

whole-leaf water-use efficiency. 41

INTRODUCTION   43

The movement of CO_2 into the leaf and the loss of water from the leaf both   44

occur through the stomata, creating a strong relationship between transpiration and   45

productivity (de Wit, 1958; Fischer et al., 1998; Richards, 2000; Kemanian et al., 2005; 46

Blum, 2009; Sperry et al., 2017; Sinclair, 2018; Xiong and Nadal, 2020). To support   47

productivity (high transpiration), plants must efficiently conduct water from the soil to 48

the evaporation sites in the leaf. Leaf hydraulic conductance (K_{leaf}) is a substantial 49

bottleneck accounting for approx. 30% (on average) and up to 98% of the total water   50

resistance of well-watered plants (Sack and Holbrook, 2006). K_{leaf} is the ratio of water 51

flow through the leaf (xylem to evaporation site) to the water potential gradient across 52

the leaf (Sack and Holbrook, 2006). K_{leaf} is a dynamic trait, which responds to both 53

abiotic (Tyree et al., 2005; Blackman et al., 2009; Shatil-Cohen et al., 2011; Pou et al., 54

2013; Prado and Maurel, 2013; Sade et al., 2015; Kelly et al., 2017) and biotic stress   55

(Attia et al., 2019; Dalal et al., 2020). The structural-functional regulation of leaf   56

hydraulics involves four main pathways: 1) hollow xylem elements; 2) the apoplast,   57

along the cell walls of the leaf tissues; 3) the symplast, which connects cells via   58

plasmodesmata and 4) transmembrane transport, mainly via aquaporins (AQPs).   59

However, little is known about the relative contributions of the different pathways and   60

the mechanisms that regulate K_{leaf}.   61

A simplified approach divides K_{leaf} into two parts: xylem hydraulic   62

conductance, K_x, known as the axial path, and out-of-xylem hydraulic conductance,   63

K_{ox}, known as the radial path. Leaf vein architecture and density greatly affect K_x 64

(Nardini et al., 2005a; Brodribb et al., 2007; Blackman et al., 2010; Brodribb et al.,   65

2010; Scoffoni et al., 2011; Caringella et al., 2015). As the xylem vessels are composed 66

mainly of dead cells, rapid impairment of K_{leaf} was previously attributed to breaks in 67

the water-column continuum (i.e., embolism). However, in recent decades, research has 68

revealed that living cells are involved in the regulation of K_{ox}, which contributes 69

significantly to K_{leaf} and its dynamics (Shatil-Cohen et al., 2011; Sade et al., 2014; 70

Scoffoni and Sack, 2017; Grunwald et al., 2021). 71

The living vascular parenchyma, bundle-sheath cells (BSC) and mesophyll cells 72

(MC) can potentially regulate K_{ox} according to internal and environmental signals. High 73

K_{ox} resistance may also help leaves to avoid catastrophic xylem failure (Scoffoni et al., 74

2017) by dynamically regulating the resistance of the "next in line" pathway (Yaaran 75

and Moshelion, 2016). Previously, stress-induced changes in K_{ox} were attributed to leaf 76

shrinkage (i.e., physical alterations of the apoplastic water pathway; Scoffoni et al., 77

2014) and/or to changes in AQP expression and/or activity (i.e., alterations in the 78

transmembrane water pathway). The latter has been reported in BSC (Kim and Steudle, 79

2009; Shatil-Cohen et al., 2011; Prado et al., 2013; Sade et al., 2015), MC (Morillon 80

and Chrispeels, 2001) and guard cells, (Grondin et al., 2015) as well as at the whole- 81

leaf level (Nardini et al., 2005b; Cochard et al., 2007; Pou et al., 2013; Kelly et al., 82

2017). 83

The bundle sheath is a single cell layer enwrapping the vasculature, which 84

possesses selective barrier characteristics (Pilot et al., 2004; Leegood, 2008; Galvez- 85

Valdivieso et al., 2009; Grunwald et al., 2021) with low levels of apoplastic transport 86

(Shapira et al., 2009; Shatil-cohen and Moshelion, 2012). It is symplastically isolated 87

from the xylem, which limits the transmembrane movement of water into the leaf, but 88

is connected to neighboring MC by plasmodesmata (Ache et al., 2010). The anatomical 89

structure of the MC correspondingly affects K_{leaf} (Brodribb et al., 2007; Scoffoni et al., 90

2014; Buckley, 2015). The current assumption is that the palisade MC functions as a 91

sort of water capacitor (Zwieniecki et al., 2007); whereas the spongy MC are more 92

dynamic in terms of their water status and play a significant role in the determination 93
of K_{ox} (Buckley, 2015; Xiong and Nadal, 2020). In MC, water can move via all three 94
pathways in parallel. A controlled trade-off between water pathways has been 95
suggested as a mechanism for coping with drought stress (Morillon and Chrispeels, 96
2001), yet supportive evidence for this hypothesis is scarce and the evidence that does 97
exist suggests routing toward the apoplastic pathway (Pou et al., 2013) or an increase 98
in transmembrane pathway (Morillon and Chrispeels, 2001; Martre et al., 2002). 99

The drought-induced phytohormone abscisic acid (ABA) can affect 100
transmembrane transport via the levels of expression, phosphorylation and/or 101
degradation of AQPs (reviewed by Shivaraj et al., 2021). ABA has been shown to 102
reduce the membrane osmotic water permeability (P_f) of BSC via AQPs, to have no 103
effect on the P_f of mesophyll protoplasts (Shatil-Cohen et al., 2011) and to link the 104
transmembrane water pathway and regulation of K_{leaf} via non-stomatal mechanisms 105
(Pantin et al., 2013; Coupel-Ledru et al., 2017). Conversely, ABA fed through the roots 106
was reported to increase the P_f of WT mesophyll protoplasts after 24 h and that of *aba1* 107
(mutants with impaired ABA synthesis) protoplasts after 3 h (Morillon and Chrispeels, 108
2001). 109

Focusing on the symplastic pathway, only a few works have studied the effects 110
of ABA and abiotic stress on plasmodesmata. Those works have reported inconsistent 111
results regarding increased (Erwee and Goodwin, 1984; Schulz, 1995) or reduced (Cui 112
and Lee, 2016; Kitagawa et al., 2019) permeability of the plasmodesmata. We 113
hypothesized that ABA might have a direct effect on the symplastic water pathway, in 114
addition to its effect on the transmembrane water pathway. 115

In addition to its widely studied role as a stress hormone, low basal levels of 116
ABA have recently been shown to play a major role in physiological and developmental 117

mechanisms related to plant water status (reviewed by Yoshida et al., 2019), for example, setting the steady-state stomatal aperture (Merilo et al., 2018; Yaaran et al., 2019). However, to the best of our knowledge, the effects of basal ABA on K_{leaf} and, particularly, on K_{ox} have not yet been investigated.

The general goal of the current study was to improve our understanding of the leaf hydraulic maze, particularly along the BSC and MC continuum. We hypothesized that K_{ox} is dynamically regulated by the BSC and MC symplastic, transmembrane and apoplastic pathways in a manner that is controlled by ABA, from the low basal ABA levels observed under optimal growth conditions to the higher ABA levels that are associated with drought. To the best of our knowledge, this is the first study to investigate and compare the osmotic water permeabilities of spongy mesophyll, palisade mesophyll and BSC in the context of the regulation of leaf water balance under optimal and high-ABA conditions. To test this hypothesis, we constructed tissue-specific ABA-insensitive plants by expressing the negative dominant *abi1-1* mutant gene under a tissue-specific promoter, to generate MCabi plants (plants with ABA-insensitive MC) (Negin et al., 2019) and BSabi plants (plants with ABA-insensitive BSC), both of which maintained a WT-like stomatal response to ABA. These lines exhibited locally suppressed ABA signaling (Hoth, 2002), which may diffuse through two to seven layers of cells (Janacek et al., 2009). In this work, we address the effects of tissue-specific insensitivity to ABA from the cellular level through whole-plant transpiration rates.

MATERIALS AND METHODS

Plant Material and Growing Conditions

All of the *Arabidopsis thaliana* plants used in this study were of the Colombia (col) ecotype (wild type; WT). In addition to WT plants, we also used transgenic FBPase::*abi1-1* (MCabi; Negin et al., 2019) and SCR::*abi1-1* (BSabi) plants. *CalS8-1* (037603C) was generously provided by Jung-Youn Lee and *AtBG_ppap* (SAIL_115_G04) was ordered from SALK (Loughborough, UK). PCR was used to confirm the homozygosity of both of those lines. *CORI3::GUS-mCit* and *IQD22::GUS-mCit* lines were constructed by Carl Procko (Procko et al., 2022) and were crossed with MCabi#2 to generate lines with ABA-insensitive, labeled spongy or palisade MC. We labeled the bundle sheath by crossing BSabi#14 with *SCR::GFP* (Torne et al., 2021).

Arabidopsis plants were grown in 250-mL pots, with two to three plants in each pot. Those pots were filled with soil (Klasmann686, Klasmann-Deilmann; Germany) + 4 g/L Osmocote® 6M. The potted plants were kept in a growth chamber at 22°C and 70% relative humidity, with an 8-h light (09:00 – 17:00) / 16-h dark photoperiod (short day). During the day, the illumination, humidity and vapor pressure deficit (VPD) changed in a pattern resembling natural conditions, as in (Negin and Moshelion, 2017). The illumination intensity provided by LED light strips [EnerLED 24 V-5630, 24 W/m, 3000 K (50%)/6000 K (50%)] reached 150–200 μmol m^{-2} s^{-1} at the plant level at midday.

The FBPase::*abi1-1* and SCR::*abi1-1* Constructs and Plant Transformation

For construct assembly, the MultiSite Gateway Three-Fragment Vector Construction Kit (Invitrogen; Ghent, Belgium) was used according to the manufacturer's instructions. The construction of the MCabi plants with ABA-insensitive mesophyll is described in Negin et al. (2019). (In that work, the MCabi plants are referred as *fa* plants.) Briefly, the *abi1-1* gene (Koornneef et al., 1984) and

the FBPase promoter (Lloyd et al., 1991) were cloned into pDONR plasmids and then 167

inserted into the binary pB7M24GW (Invitrogen) plasmid. As MC and BSC are 168

connected by plasmodesmata, we cannot exclude the diffusion of mutated *abi1-1* genes 169

from MC to BSC. Therefore, we considered MCabi plants to be ABA-insensitive in 170

their MC and BSC. The plants with ABA-insensitive BSC (BSabi plants) were 171

constructed in the same way, using the SCR (Shatil-Cohen et al., 2011) promoter, which 172

has been observed in the BSC associated with all veins in mature leaves (Wysocka- 173

Diller et al., 2000). We cannot rule out the diffusion of *abi1-1* into other tissues in the 174

BSabi plants, yet trans-activation of a hairpin by an enhancer trap (J1511) active in the 175

leaf veins showed silencing spread only two to seven cells away from the vein (Janacek 176

et al., 2009). 177

The dominance of the *abi1-1* mutation is due to its "gain of function" nature, 178

negatively regulating ABA signaling (Gosti et al., 1999), with only 9% of the ABA- 179

responsive genes remaining unaltered in *abi1-1* plants (Hoth, 2002). As such, it has a 180

quantity-dependent attribute (high phosphatase activity represses the phosphorylation 181

of SnRKs and, consequently, the initiation of ABA signal transduction). In our plants, 182

abi1-1 was constitutively expressed under a non-native promotor, in addition to the 183

native ABI1 and other PP2Cs expression, suppressing ABA signaling. Therefore, we 184

expected that its strongest effect would be seen at its expression site. Having said that, 185

the dominance could be trait-dependent and, therefore, we always compared the *abi1-* 186

1 phenotype with the WT control. 187

Binary plasmids were floral-dipped (Clough and Bent, 1998). The presence of 188

the transgene containing the G to A substitution was confirmed by sequencing and NocI 189

digestion (described below). The study was performed on three independent lines of 190

each construct, using homozygous t_3 and t_4 lines. 191

Characterization of Transgenic Plants

Stomatal Aperture and Index

Stomatal aperture and index were measured as described by Yaaran et al. (2019). Briefly, epidermal peels were soaked in closure solutions under light (~150 μmol m^{-2} s^{-1}). After 1.5 h, ABA [(+)-cis, trans abscisic acid; Biosynth; Staad, Switzerland] was added to reach a concentration of 10 μM. DMSO at the same concentration was added to the control. All stomata were photographed under a bright-field inverted microscope (1M7100, Zeiss; Jena, Germany) on which a Hitachi HV-D30 CCD camera (Hitachi; Tokyo, Japan) was mounted. Stomatal images were analyzed to determine aperture size using the ImageJ software. Stomatal index was calculated as the number of stomata/number of epidermal cells in an area of 10 μm^2.

Leaf Vein Density

Leaf vein density was measured as described by Grunwald (2021a). Briefly, leaves of the same age and size as those used in the K_{leaf} assay were immersed in 96% (v/v) ethanol solution and incubated at 50°C. The used ethanol was replaced with fresh ethanol until leaves were completely colorless and then was finally replaced with 88% (v/v) lactic acid solution (Fischer Scientific; UK) for overnight incubation at room temperature. Then, the leaves were rinsed in water and submerged in 0.5% safranin O dissolved in water (Sigma Aldrich cat. no S2255) for 1 min, followed by an overnight incubation in water to remove any excess dye. Then, the leaves were scanned in a water tray using an Epson 12000XL scanner (Seiko Epson; Japan) with a 2500-dpi resolution and two to three patches (~1 cm^2 each) for each leaf (9–10 leaves per line) were analyzed using WinRhizo™ software (https://regent.qc.ca/assets/winrhizo_about.html)

for vein detection. Vein density was calculated by dividing the total vein length by the   216

total leaf area scanned.   217

Leaf area is the projected area of a fully expanded leaf (average of five leaves   218

from each plant, six plants of each of the three independent lines of MCabi and BSabi).   219

Leaf area was assessed using the LI-3100C Area Meter (https://www.licor.com/env/).   220

Leaf cross-sections for anatomical analysis of the veins and mesophyll were   221

free-hand cut using a sharp scalpel. Then, the cross-sections were cleared using   222

ClearSee (Kurihara et al., 2015) and stained in a 1-min incubation in Toluidine Blue O   223

dissolved in water (TB; cat. no. T-3260, https://www.sigmaaldrich.com). They were   224

then washed with water two or three times. Cross-sections were photographed in the   225

same set-up as that used for the analyses of stomatal aperture and index. ImageJ   226

software was used to analyze the images to determine mesophyll length.   227

Leaf Gas-Exchange and Hydraulic-Conductance (K_{leaf}) Measurements   229

Gas exchange and hydraulic conductance were measured as described by Negin et al.   230

(2019), using leaves of 6- to 8-week-old plants grown under short-day conditions.   231

Briefly, two leaves from each plant were excised before dawn and put into tubes   232

containing artificial xylem sap (AXS; Shatil-Cohen et al., 2011) and 0.01% DMSO   233

(control) or 10 μM ABA. The leaves were then put into hermetically sealed transparent   234

boxes and left in the light for a minimum of 1 h. Following that period, the lids were   235

opened for 15 min, after which time gas exchange was measured using the LI-6400xt   236

portable gas-exchange system equipped with the Arabidopsis chamber (LI-COR, Inc.;   237

Lincoln, NE, USA). Chamber conditions were set to 400 ppm CO_2, PAR of 200 $\mu E\,m^{-2}$ 238

s^{-1}, VPD of ~1.2 KPa and flow of 200 µmol/s. In the enclosed gas-exchange chamber,   239

there is a trade-off between the stability of humidity vs. stability of the flow. We choose   240

to keep the flow constant at the expense of humidity stability, to maintain an undisturbed boundary layer.

Immediately following the gas-exchange measurement, the leaf was transferred to a pressure chamber (ARIMAD-3000; MRC, Israel) to measure Ψ_{leaf} and determine K_{leaf} (Sade et al., 2014). K_{leaf} was calculated for each individual leaf by dividing the whole-leaf transpiration rate, E, by the leaf water potential (Martre et al., 2002; Sack and Holbrook, 2006). In our calculation, the leaf water potential gradient equals the Ψ_{leaf} measured in the pressure chamber as the leaf petiole was dipped in AXS at a water potential of 0.027 MPa. These measurements were conducted between 10:00 and 13:00 (1–4 h after the lights were turned on).

Whole-Plant Continuous Transpiration

The whole-plant continuous transpiration rate was measured using a high-throughput telemetric, gravimetric-based phenotyping system (Plantarry 3.0 system, Plant-DiTech; Israel; Dalal, 2020) in the greenhouse of the I-CORE Center for Functional Phenotyping (http://departments.agri.huji.ac.il/plantscience/icore.phpon) as described in (Yaaran et al., 2019; Dalal et al., 2020). The output (weight) of the load cells was monitored every 3 min and analyzed using SPAC analytics (Plant-Ditech). Seedlings were transferred to 1.6-L pots (four plants per pot) and gradually exposed to semi-controlled greenhouse conditions: 24–9°C (day/night) and the natural day length and light conditions prevailing in Rehovot, Israel during December and January of 2021.

After establishment, each pot was placed on a load cell in a randomized block arrangement. Daily measurements were conducted continuously and simultaneously for all of the plants in the array, so that all of the plants were exposed to similar ambient

conditions at each measurement point. The soil surface surrounding each Arabidopsis 266

plant was covered, as well as the crack between the pot and load cell, to prevent 267

evaporation. The transpiration rate was normalized to the total plant weight to 268

determine E, as described in (Halperin et al., 2016; Yaaran et al., 2019). 269

Protoplast Isolation and Measurement of the Membrane Osmotic Water 271
Permeability Coefficient (P_f) 272

Protoplast isolation and measurements of the osmotic water permeability 273

coefficient (P_f) were performed as described by Shatil-Cohen et al., (2014). In brief, 274

protoplasts were isolated from 6- to 8-week-old plants according to the gentle and rapid 275

(<30 min) method using an isotonic solution in the extraction process (pH 5.7, 600 276

mOsmol). For the ABA treatment, protoplasts were incubated in 1 μM ABA for 1–4 h. 277

We observed the protoplasts swelling in response to the hypo-osmotic challenge (of 278

0.37 MPa) generated by changing the bath solutions from isotonic (600 mOsmol) to 279

hypotonic (450 mOsmol). Palisade and spongy mesophyll protoplasts were identified 280

by *mCit* fluorescence. Protoplasts were recorded using an inverted epifluorescent 281

microscope (Nikon Eclipse TS100) with a 20x/NA 0.40 objective, a CCD 12-bit camera 282

Manta G-235B (https://www.alliedvision.com) and image-processing software 283

(AcquireControl® v5.0.0; https://www.alliedvision.com). P_f was determined based on 284

the rate at which cell volume increased for 60 s starting from the hypo-osmotic 285

challenge, using a numerical approach in an offline curve-fitting procedure of the PfFit 286

program. 287

Relative Symplastic Permeability 289

Drop-and-See (DANS) Assays 290

Drop-and-see (DANS) assays were performed as described by Cui and Lee
(2016) with minimal modifications. In brief, two 1-µL droplets of 2 mM 5(6)-
carboxyfluorescein diacetate (CFDA; cat. #19582 https://www.caymanchem.com)
were loaded on the center of the adaxial surface of a mature leaf from a 7- to 8-week-
old plant. This was followed by inverted epifluorescent microscope imaging (described
above for the P_f measurements) of the abaxial surface of the leaf at 10 min after loading.
The non-fluorescent ester CFDA can cross cell membranes passively in its electrically
neutral or near-neutral form. Once inside the cells, it is subject to cleavage by esterases
to form a polar fluorescent compound, CF. The extent of cell-to-cell dye movement is
expected to represent the permeability of the plasmodesmata and was evaluated using
an Intensity-Distance chart that was analyzed using ImageJ. A magnification of 10x
was used, despite the fact that the size of some of the MCabi samples exceeded the
imaging border (maximum of 388% compared to WT under control conditions), in
order to capture the extent of all lines' cell-to-cell fluorescence under the same imaging
conditions. The average of the two droplets was considered the leaf's DANS
measurement.

Both attached and detached leaf measurements were performed between 10:00
and 14:00. The detached-leaf experimental set-up was like the set-up used for the
measurements of hydraulic conductivity (K_{leaf}).

Statistical Analysis 311

The data were analyzed using Student's *t*-test for comparisons of two means 312

and Tukey's HSD test for comparisons of more than two means. When two variables 313

were examined (e.g., line and ABA treatment), the interaction between those factors 314

was evaluated using a two-way ANOVA, followed by Tukey's HSD test. The model 315

effects and different *p*-values attained from the two-way ANOVAs are presented in 316

Supplemental Table S1. All analyses were done with JMP software (SAS; Cary, NC, 317

USA). 318

RESULTS 320

Characterization of the BSabi and MCabi Phenotypes 321

To better understand the effects of ABA on leaf hydraulics, we generated tissue- 322

specific ABA-insensitive lines. We expressed the dominant *abi1-1* mutation under the 323

FBPase promoter (for expression in green tissues, but not in guard cells), to create 324

plants that had suppressed ABA signaling in their MC and BSC (i.e., MCabi plants), or 325

under the SCR promoter (for expression in BSC, but not in MC), to create plants with 326

suppressed ABA signaling in their BSC (i.e., BSabi plants; see Materials and Methods). 327

abi1-1 may diffuse through two to seven layers of cells (Janacek et al., 2009), but its 328

strongest effect is expected at the site at which it is expressed. Three independent lines 329

were selected for each mutation (MCabi: 1, 2 and 7; BSabi: 12, 14 and 99). 330

The MCabi plants were broadly characterized and validated (Negin et al., 2019) 331

and the same validation was performed for the BSabi plants (Supplemental Figure 1). 332

MCabi and BSabi lines exhibited different phenotypes, illustrating the vital 333

implications of expression location. MCabi's leaves were larger and thicker than those 334

of the WT, while BSabi had smaller leaves (Figure 1A, F). BSabi maintained the same 335

proportion of palisade to spongy MC as the WT; whereas MCabi had more space 336

occupied by palisade MC (Figure 1F). Despite the changes in leaf size, the vein 337

densities of BSabi and MCabi remained the same as that of the WT (Figure 1C), with 338

BSabi differing from the WT in its narrower veins, but only MCabi differing from WT 339

in its increased xylem to total vein ratio (Figure 1G). Moreover, the steady-state 340

stomatal aperture (Figure 1D) and stomatal index (Figure E) of MCabi were greater 341

than those of the WT and BSabi. Both MCabi and BSabi plants had a WT-like stomatal 342

response to ABA (Negin et al., 2019; Supplemental Figure 1C) and their average foliar 343

ABA levels were similar to those of the WT (Supplemental Figure 2A; one of the three 344

BSabi lines had ABA levels that were higher than those of the WT). 345

Gas Exchange and Leaf Hydraulics 347

To assess the relative contributions of the MC and BSC to the overall K_{leaf}, we 348

performed gas-exchange measurements and then measured Ψ_{leaf} using a pressure 349

chamber. The experiment was first done for each transgene separately (Supplemental 350

Figure 3). All three independent lines were found to be similar to one another and the 351

experiment was then repeated with MCabi and BSabi together, so that they could be 352

compared at the same time under the same conditions (Figure 2). Under optimal 353

conditions, BSabi and MCabi had similar levels of transpiration (E), which were higher 354

than that of the WT. Yet, despite that higher E, both types of transgenic plants 355

maintained Ψ_{leaf} that was similar to that of the WT, resulting in higher K_{leaf} than the WT 356

(Figure 2A–C). BSabi and MCabi exhibited higher stomatal conductance (g_s) 357

accompanied by only slightly higher net carbon assimilation rate (A_N), resulting in 358

lower intrinsic water-use efficiency (iWUE; Figure 2D–F). Following ABA treatment, 359

all lines exhibited reduced E, g_s and, surprisingly, K_{leaf} (Figure 2A, C). MCabi and 360

BSabi maintained an unaltered stomatal ABA response (Supplemental Figure 1C; Negin 2019) and, therefore, reduced their g_s and E in response to ABA (Figure 2A, D). However, while ABA reduced the Ψ_{leaf} of the WT, the Ψ_{leaf} levels of MCabi and BSabi remained unchanged (Figure 2B). Consequently, the K_{leaf} reduction observed in MCabi and BSabi was mainly related to their reduced transpiration (Figure 2A–C). Overall, this resulted in higher iWUE for the WT, as compared to the two sets of transgenic lines, even following ABA treatment (Figure 2F).

Although MCabi plants had a higher stomatal index and larger stomatal apertures than the BSabi plants (Figure 1D, E), they had similar E and g_s (Figure 2A, D). E was measured in an enclosed gas-exchange chamber, in which it was difficult to maintain a constant VPD at high levels of E (Supplemental Figure 3). To overcome this challenge, we measured whole-plant transpiration in an open space, under natural conditions, using a gravimetric system in which all plants were simultaneously exposed to the same VPD. We found that the MCabi plants had the highest transpiration rate (Figure 3E). This trend was maintained throughout the whole growing period (40 days; Supplemental Figure 5C). When transpiration was normalized to plant weight, MCabi plants still had the highest E, with BSabi plants exhibiting intermediate level E and WT plants exhibiting the lowest E (Figure 3F). The finding that MCabi had the highest E suggests that the mesophyll has an additive effect on leaf water balance.

Differences Between the Palisade and Spongy MC: P_f and ABA Response

To study the mesophyll's contribution to E and K_{leaf}, in general, and the specific contributions of palisade and spongy MC, we used reporter lines with palisade (*IQD22pro::GUS-mCit*) and lower spongy (*CORI3::GUS-mCit*) tissue-specific labeling with the fluorescent protein mCit (Procko et al., 2022) and examined the P_f of

each type of cell. The two reporter lines revealed largely restricted labeling; we did not

find any apparent labeling on any other leaf tissue, including the bundle sheath and

guard cells (Figure 4A, B). Spongy mesophyll protoplasts exhibited higher P_f under

control conditions (Figure 4C) and were smaller than the palisade protoplasts (Figure

4D). The P_f values of both types of protoplasts showed a tailed distribution, with most

of the protoplasts having low P_f values. Yet, even when we considered only 90% of the

population, to exclude the "tails" of the populations, the P_f of the spongy mesophyll

protoplasts remained almost double that of the palisade protoplasts. Moreover, more

spongy mesophyll protoplasts had extreme P_f values of over 60 µm sec^{-1}. In response

to ABA, the higher P_f of the spongy mesophyll protoplasts was reduced by roughly

60% (19.2 $\pm$ 0.7 µm s^{-1} to 7.2 $\pm$ 1.8 µm s^{-1}, respectively); whereas the average P_f of the

palisade protoplasts did not change (Figure 4E).

To examine the effect of ABA insensitivity on MC, we fluorescence-labeled

palisade and spongy mesophyll protoplasts of the MCabi#2 line. Surprisingly, we found

that ABA insensitivity in the spongy mesophyll reduced the P_f of spongy mesophyll

protoplasts, but did not affect the P_f of palisade protoplasts (Figure 5A). Moreover,

MCabi spongy protoplasts maintained their low P_f at increasing concentrations of ABA

and remained insensitive to ABA even when they were exposed to 50 µM ABA (Figure

5C).

We also used fluorescence-labeled ABA-insensitive BSCs of the BSabi#14 line

to investigate the effect of ABA insensitivity in the bundle sheath and found that

insensitivity increased the P_f of the bundle-sheath protoplasts (Figure 5B). BSabi#14

bundle-sheath protoplasts also preserved their ABA insensitivity, and their P_f did not

change in response to increasing ABA concentration (Figure 5D), with average P_f

values lower than those measured in the previous experiment (Figure 5B).

We conducted another experiment with the same lines under the same   411

conditions to determine whether this difference represents natural or experimental   412

variation. The results of that work confirmed that WT bundle-sheath P_f remained lower   413

than the BSabi#14 bundle-sheath P_f, as shown in Figure 5B (WT 8.6$\pm$2, BSabi#14 16.9   414

$\pm$ 6.4). Taken together, both basal and higher levels of ABA appear to affect the   415

movement of water across spongy MC and BS membranes, affecting the   416

transmembrane water pathway. Next, we were interested in exploring the effects of   417

ABA on the movement of water through the symplastic pathway.   418

Effects of ABA on the Symplastic Water Pathway   420

To evaluate the relative contribution of the symplastic water pathway to the total   421

K_{leaf}, we measured the K_{leaf} of known mutants with severely altered plasmodesmatal   422

permeability: *CalS 8-1*, whose plasmodesmatal permeability is ~35% higher than that   423

of the WT (Cui and Lee, 2016) and *AtBG_ppap*, whose plasmodesmatal permeability   424

is ~45% lower than that of the WT (Zavaliev et al., 2013). The K_{leaf} values of both   425

mutants were similar to that of the WT (Figure 6A) and their g_s, E and Ψ_{leaf} were similar   426

as well (Supplemental Figure 6). Next, we used the DANS assay (Cui and Lee 2016) to   427

measure the relative plasmodesmatal permeability of MCabi and BSabi under control   428

conditions and in response to ABA treatment. In principle, the cell-to-cell spread of a   429

fluorescent dye can be used to measure the permeability of the plasmodesmata (this   430

spread can be seen in the top of Figure 6B, marked by yellow circles. For more detailed   431

information, see the Material and Methods). Under control conditions, the relative   432

plasmodesmatal permeability of the MCabi plants was highest, that of the BSabi was   433

intermediate and the WT had the lowest relative plasmodesmatal permeability (Figure   434

6B). This ranking corresponds to the ranking of those lines' E under greenhouse 435

conditions (Figure 3F). 436

Following ABA treatment, only WT leaves exhibited significantly increased 437

relative plasmodesmatal permeability (see Figure 5B and Supplemental Figure 8A for 438

detailed information for each line). Changes in plasmodesmatal permeability can be 439

attributed to the accumulation of callose (a β-1,3 glucan) in the plasmodesmata 440

(Amsbury, 2017). Nevertheless, there were no detected changes in the callose levels in 441

either of the lines in any of the treatments in our experiment (Supplemental Figure 8B). 442

Validation of proper callose staining was done on leaves that had been pre-treated with 443

flagellin, whose reduced relative plasmodesmatal permeability was accompanied by 444

increased callose deposition, as expected (Supplemental Figure 7). 445

DISCUSSION 447

BSC and MC Control K_{leaf} and Transpiration via ABA Regulation 448

The working hypothesis in this study was that K_{ox} is controlled by a series of 449

living cells, including BSC and MC. Our results suggest that the P_f levels of BS and 450

spongy MC are a target of basal and higher ABA levels, while the way that they regulate 451

K_{leaf} may not be straightforward. The increased E and K_{leaf} observed among both BSabi 452

and MCabi under optimal conditions implies that, in the WT, E and K_{leaf} are limited by 453

a non-stomatal mechanism, most likely involving the effect of basal ABA on 454

physiological and developmental regulation. 455

Recently, it has been suggested that the basal levels of ABA expressed under 456

non-stress conditions may limit the steady-state stomatal aperture (Merilo et al., 2018; 457

Yaaran et al., 2019). Our finding that BSC insensitivity to ABA results in increased P_f 458

(Figure 5B) together with BSabi's elevated K_{leaf} under control conditions (Figure 2C) 459

suggest that ABA may play a similar role in BSC-specific regulation. We suggest that 460

this dual restricting role (in the guard cells and in the BSC) of basal ABA serves to 461

preserve leaf water balance by balancing leaf water influx (via the bundle sheath) and 462

efflux (via guard cells) while increasing the P_f of the spongy MC (Figures 4E and 5A), 463

to sustain overall water flow and content between the controlled entry and exit 464

positions. We further suggest that this role of basal ABA is augmented by its activity 465

under stress conditions, when high ABA levels reduce both stomatal apertures and K_{leaf}. 466

It is interesting to note that similarities between the regulation mechanisms of guard 467

cells and BSC have also been reported in the context of blue light signal-transduction 468

pathways, which initiate stomatal opening while increasing bundle-sheath P_f (Iino et 469

al., 1985; Zait et al., 2017; Torne et al., 2021; Grunwald, 2022). Such a role for the 470

bundle sheath also reinforces its importance as a leaf hydraulic checkpoint (Ache et al., 471

2010; Griffiths et al., 2013; Caringella et al., 2015). 472

Nevertheless, we wondered about the benefits of restricting hydraulic 473

conductance (and possibly productivity) under optimal growth conditions. The answer 474

to this question may lie in the reduced iWUE of MCabi and BSabi, which both exhibited 475

a major increase in E as compared to WT [168–180% under controlled growth-chamber 476

conditions (Figure 2A) and 68–80% under open-air greenhouse conditions (Figure 3F)], 477

accompanied by a minor increase in the carbon assimilation rate, A_N (~10%, Figure 478

2E). As E increases linearly with stomatal aperture, while A_N has a maximal saturation 479

point, the optimized WUE falls within the non-maximal gas-exchange range (Yoo et 480

al., 2009). Brodribb et al.'s (2005) idea that the photosynthetic system is saturated at 481

high K_{leaf} levels further supports the potential disadvantage of very high K_{leaf}. 482

Therefore, non-maximal hydraulic conductance and E under optimal growth conditions 483

may have a selective advantage at the evolutionary time scale, preventing excessive water loss that would provide only minimal gains in productivity.

Under ABA treatment, both BSabi and MCabi maintained high, unaltered Ψ_{leaf} (Figure 2B), indicating that BSC and MC contribute to Ψ_{leaf} regulation. Still, these plants exhibited strong reductions in E (46–55%), resulting in K_{leaf} reduction. This is unlike the results presented by Shatil-Cohen et al., (2011), in which ABA caused only a small reduction in E (22–24%) and failed to reduce K_{leaf}. This demonstrates the dual effect of ABA (Pantin et al., 2013): a non-direct hydraulic mechanism enhancing stomatal closure via Ψ_{leaf} regulation (Shatil-Cohen et al., 2011) along with a direct effect on stomatal closure. These results, although surprising, cannot rule out the importance of the bundle sheath as an internal barrier in the leaf, but rather suggest that the MC, especially the spongy MC, acts together with the bundle sheath in a series.

Palisade and Spongy MC Differ in Their Hydraulic Properties

To better understand MC's contribution to leaf hydraulics, we examined palisade and spongy MC separately. Palisade and spongy MC differ in both their structure and their hydraulic properties (Figure 4; (Canny, 2012; Buckley et al., 2015; Álvarez-Arenas et al., 2018; Muries et al., 2019). The smaller spongy MC are distant from each other, forming a tissue with a large surface area, air spaces and low cell-to-cell connectivity (Buckley et al., 2015; Borsuk et al., 2022). Based on our findings, the P_f of the spongy MC under non-stressed conditions was almost twice that of the palisade MC (Figure 4C, E). The spongy mesophyll's vast exposure to the leaf's internal air spaces together with the high P_f of its cells may explain those cells' rapid loss of water and shrinkage in response to high VPD (Canny, 2012) or dehydration (Muries et al., 2019). This is an intriguing point, which may suggest that spongy MC act as hydraulic

sensor in the following way: High P_f enables the rapid loss of water content, resulting   509

in decreased cell volume, which lessens cell-wall and plasma-membrane interactions,   510

to induce ABA production (Bacete et al., 2022). In this way, a physical change, such as   511

high VPD causing an increased rate of water loss from the leaf, can be translated into a   512

physiological response of stomatal closure (Mott, 1991). Surprisingly, MCabi exhibited   513

high K_{leaf} (Figure 2C) despite its spongy MC's low P_f, suggesting that its high K_{leaf} is   514

mainly supported by a bundle sheath that is insensitive to ABA and apoplastic flow, as   515

concluded from its ability to maintain high Ψ_{leaf} (Figure 5B). Nevertheless, the low P_f 516

of MCabi's spongy MC may obstruct their hypothesized role as a hydraulic sensor   517

(mentioned above), resulting in their higher E, g_s, Ψ_{leaf} and K_{leaf} (Figure 2A–D).   518

According to these results, spongy MC P_f dynamics may have a regulatory role rather   519

than directly changing K_{ox}. In response to higher ABA levels, the WT spongy MC   520

reduced its P_f to avoid further water loss, while the palisade P_f remained unchanged   521

(Figure 4E), highlighting tissue-specific ABA regulation.   522

But why would ABA-insensitive spongy MC present a low P_f similar to that of   523

WT spongy mesophyll exposed to ABA (Figures 5A and 4E, respectively)? To answer   524

this question, we propose that spongy MC behave according to an ABA optimization   525

curve (for further elaboration, please see Supplemental Figure 9). Palisade MC's lack   526

of response to ABA supports the idea that these cells play a static hydraulic role   527

(Zwieniecki et al., 2007), in which they serve as a water buffer to maintain leaf water   528

balance (Nardini et al., 2010; Rockwell et al., 2014). The combination of palisade   529

mesophyll with static hydraulic properties and the lack of any direct effect of ABA on   530

photosynthesis (Negin et al., 2019) can be an advantage for continued photosynthesis   531

(Tholen et al., 2012; Buckley, 2015) and position ABA as, primarily, a water-balance   532

regulator (from BSC to guard cells) that controls the transmembrane water pathway.   533

The Role of Plasmodesmata in the Symplastic Water Pathway and K_{ox}   535

To evaluate the roles of plasmodesmata and the symplastic water pathway in   536

K_{leaf}, we used impaired plasmodesmata-permeability mutants (*cals8-1*; (Cui and Lee,   537

2016) and *AtBG_ppap*; (Zavaliev et al., 2013)). Both of those mutants exhibited WT-   538

like K_{leaf} (Figure 5A), indicating that the symplastic pathway has a negligible influence   539

on K_{leaf}. Moreover, the fact that ABA increased the WT's symplastic permeability   540

(Figure 5B), alongside its reduction of K_{leaf}, supports this idea. As high ABA levels   541

reduce P_f while increasing plasmodesmatal permeability, we suggest that the symplastic   542

water flow may provide better cell-to-cell conductivity and turgor sharing and help to   543

maintain a uniform water potential across the tissue. Additionally, it may act as a rapid   544

facilitator of the stress signal by distributing the ABA among the cells.   545

The greater symplastic conductivity of BSabi and the even greater symplastic   546

conductivity of MCabi under control conditions are, therefore, surprising (Figure 6B).   547

Here, we present three possible explanations for this. The first is altered development.   548

During sink-to-source transition, the leaves' plasmodesmata are dramatically reduced   549

(Roberts et al., 2001). Therefore, impaired maturation may sustain a higher number of   550

plasmodesmata. Interestingly, *abi1-1* hybrid aspen trees fail to block their   551

plasmodesmata as they enter dormancy (Tylewicz et al., 2018). The second possible   552

explanation is impaired hormonal crosstalk with salicylic acid. Salicylic acid reduces   553

the permeability of plasmodesmata (wang et al., 2013; Cui and Lee, 2016). ABI1 binds   554

directly to salicylic acid and impairs that crosstalk (Manohar et al., 2017). The relatively   555

low levels of salicylic acid observed in MCabi and BSabi (Supplemental Figure 2B)   556

support the existence of such altered crosstalk, which may affect the permeability of   557

the plasmodesmata. Finally, the *abi1-1* protein may have a signaling function. While   558

abi1-1 mutants lack phosphatase activity, the protein itself may have a signaling 559

function independent of its phosphatase activity (Gosti et al., 1999). If plasmodesmatal 560

permeability is involved in such an ABA-signaling pathway, it may be boosted in the 561

transgenic plants due to the high abundance of *abi1-1* in the specific tissue. Although 562

compelling, we did not pursue these leads as they go beyond the scope of this study. 563

Further research is needed to determine ABA's effect on the permeability of 564

plasmodesmata. 565

A Suggested Model for the Role of ABA in the Regulation of K$_{leaf}$ 567

Integrating our results, we suggest the following hypothetical model of radial 568

water movement within the out-of-xylem leaf tissues (Figure 7). Under optimal 569

conditions, water enters the leaf through the BSC, which selectively regulate 570

transmembrane radial water flux (Leegood, 2008; Galvez-Valdivieso et al., 2009; 571

Shapira et al., 2009; Ache et al., 2010; Shatil-Cohen et al., 2011; Prado and Maurel, 572

2013; Grunwald et al., 2021) based on the osmotic water permeability of their 573

membranes (Figure 5B). Thus, the bundle-sheath barrier serves as the first control 574

point, regulated by basal ABA (Figure 7A, Control Point 1). Then, water proceeds 575

toward and within the MC, with most of the water flowing through the hydraulically 576

dynamic spongy MC (Figure 4C; Buckley et al., 2015; Buckley, 2015; Xiong and 577

Nadal, 2020). The transmembrane pathway (Cochard et al., 2007; Kim and Steudle, 578

2007) and/or the apoplastic pathway (Sack and Holbrook, 2006; Voicu et al., 2009; 579

Buckley, 2015) govern K$_{leaf}$ under optimal conditions, while the contribution of the 580

symplastic pathway is negligible (Figures 6 and 7A, Control Point 2). 581

The high *P$_f$* of the spongy MC may have two advantages, allowing the spongy 582

mesophyll to serve as an interchange point between the parallel water pathways and as 583

a hydraulic sensor. As an interchange point between the parallel water pathways, water   584

crossing a cell membrane may continue toward the next cell in the transmembrane   585

pathway or remain in the apoplast, wetting the cell wall to support the apoplastic water   586

pathway (Maurel, 1997; Steudle and Peterson, 1998; Figure 6A, Control Point 3) and   587

transpiration (Figure 3F and Supplemental Figure 5). As a hydraulic sensor, the spongy   588

mesophyll's surface area combined with its high P_f results in rapid water loss when   589

there is an imbalance between water supply and demand, quickly reducing the cell's   590

water content and volume and offering a conceivable mechanism to capture changes in   591

the rate of water loss from the leaf. The simultaneous limitation of K_{leaf} and stomatal   592

aperture by basal ABA levels maintains the balanced leaf water status that is mirrored   593

in the high Ψ_{leaf} and turgor (Figure 7A).   594

In the presence of a high level of ABA, xylem ABA reduces the BSC's P_f 595

(Shatil-Cohen et al., 2011; Figure 6C, Control Point 1), which limits the movement of   596

water into the leaf. In the immediate term, the stomatal efflux remains unchanged and   597

the spongy MCs' high initial P_f (Figure 4C) results in their rapid water loss, shrinkage   598

(Canny et al., 2012; Muries et al., 2019) and ABA production (McAdam and Brodribb,   599

2016; Sussmilch et al., 2017; Sack et al., 2018; Bacete et al., 2022; Figure 7A, B). When   600

they sense ABA, the spongy MCs decrease their P_f (Figures 4E and 6C, Control Point   601

2, Figure 7B), resulting in the retention of water within the whole mesophyll tissue and   602

reducing the flow over the transmembrane water pathway, cell-wall wetting and,   603

subsequently, the apoplastic flow (Figure 7C, D, Control Point 3). As a result of all the   604

above and the direct effect of ABA on stomata, K_{leaf} and g_s decrease, while the leaf   605

water status deteriorates (e.g., decreased turgor and Ψ_{leaf}). Simultaneously, the high   606

ABA levels increase the symplastic permeability (Figures 6B and 7C, Control Point 2).   607

This may facilitate the cooperation of mesophyll cells and solute transport when E   608

levels are low (Morillon and Chrispeels, 2001) and may enhance the symplastic passage   609

of ABA through the leaf and toward the guard cells.   610

This study highlights the differential effects of ABA on BSCs and spongy and   611

palisade MCs, as well as newly reported effects of those tissues on K_{leaf}. Our efforts to   612

untangle the effects of ABA on leaf hydraulics have revealed that a series of living   613

cells, which are differentially affected by ABA, take part in determining leaf water   614

balance under optimal and high-ABA conditions. We also provide evidence that   615

supports the hypothesis that ABA acts as a central water-balance coordinator and   616

suggest that ABA levels dynamically fine-tune the movement of water into and out of   617

leaves.   618

Acknowledgments and Funding   620

This research was supported by the Israel Science Foundation (Grant No. 1043/20) and   621

NIH (Grant No. 5R35GM122604). We thank Jung-Youn Lee for generously providing   622

us with the *CalS8-1* (037603C) line. We thank Dr. Julius Ben-Ari for the LC-MS/MS   623

analyses of ABA samples and Eduard Belausov for the confocal imaging.   624

Author Contributions   626

Adi Yaaran designed and performed the research, analyzed the data, formulated the   627

hypotheses and wrote the manuscript. Erez Eyal contributed to the characterization of   628

transgenic plants and the P_f measurements. Carl Procko constructed and validated the   629

CORI3::GUS-mCit and *IQD22::GUS-mCit* lines. Menachem Moshelion (the   630

corresponding author) formulated the hypotheses and wrote the manuscript together   631

with Adi Yaaran.   632

Figure 1. MCabi and BSabi phenotypes. Morphological characteristics of 7- to 8-week-old MCabi and BSabi plants 634

grown under short-day conditions, including (A) representative images of the rosettes and fully expanded leaves of 635

each line (bar = 2 cm), (B) leaf area, (C) vein density, (D) stomatal aperture, (E) stomatal index and (F) total leaf 636

thickness divided by the widths occupied by palisade (dark gray) and spongy (light gray) mesophyll. Capital letters 637

refers to significant differences between each mesophyll layer and lower-case letters refer to significant differences 638

in total leaf thickness (Tukey's test, $P < 0.05$). (G) Vein area divided into the area occupied by the xylem (dark gray) 639

and the rest of the vein area (light gray). An asterisk indicates a significantly higher ratio of xylem to total vein area, 640

capital letters refer to significant differences between the xylem and the rest of the vein area and lower-case letters 641

refer to significant differences in total vein area (Tukey's test, $P < 0.05$). (B–E) The box height shows 25–75% of 642

the data range, the symbols indicate the individual data values, the solid line indicates the median, o indicates the 643

mean and the whiskers delimit ±3 times the interquartile range. (F–G) Data are means ± SE. (B) $n = 36$ for WT and 644

90 for MCabi and BSabi, 5 leaves from 6 plants for each transgenic line. (C) $n = 10$ leaves for WT and 28–30 for 645

MCabi and BSabi, 9–10 leaves from each transgenic line, an average of 2–3 patches of ~1 cm^2 per leaf. (D) $n = 80$ 646

stomata for WT and >245 for MCabi and BSabi, ~ 80 stomata from each transgenic line, 20 stomata from 4 leaves 647

for each. (E) $n = 80$ stomata for WT and >245 for MCabi and BSabi, ~ 80 stomata from each transgenic line, 20 648

stomata from 4 different leaves for each. (F, G) $n = 5$ leaves for WT and 15 for MCabi and BSabi, 5 leaves from 649

each transgenic line, an average of 3 cross-sections per leaf. 650

Figure 2. Gas exchange of WT, MCabi and BSabi leaves under non-stressed conditions in the presence of xylem-fed ABA. Leaves of 6- to 8 week-old plants were cut before dawn and petiole-fed AXS (dark gray) or AXS + 10 µM ABA (light gray). After 1–4 h, the (A) whole-leaf transpiration rate, E, and (B) leaf water potential, Ψ_{leaf}, were measured, enabling the calculation of (C) leaf hydraulic conductance (K_{leaf}) for each individual leaf. In addition, the (D) stomatal conductance, g_s, and (E) net carbon assimilation rate, A_N, were measured, enabling the calculation of (F) iWUE for those same leaves. The percentage change following the ABA treatment is indicated as well. Data were analyzed using two-way ANOVA. (All model effects and interactions are presented in Supplemental Table 1.) Different letters indicate significant differences according to Tukey's HSD test ($P < 0.05$); p-values of the effect of each treatment on each genotype according to t-tests are also indicated. The box height shows 25–75% of the data range, the symbols indicate individual data values, the solid line indicates the median, o indicates the mean and the whiskers delimit ±3 times the interquartile range of at least three independent experiments. n = 21 leaves for WT and 23–24 leaves for MCabi and BSabi, 7–8 leaves from each transgenic line.

Figure 3. Continuous measurements of whole plants under greenhouse conditions. (A) Top view and (B) side view 666

of pots with 8-week-old WT Arabidopsis plants, 4 plants per pot. All pots were randomized and measured 667

simultaneously on (C) a lysimeter system located in a greenhouse under semi-controlled conditions. The plants were 668

exposed to (D) ambient vapor pressure deficit (VPD) and PAR conditions as shown here for one representative day, 669

the 11th day after the pots were loaded onto the lysimeters. (E) Whole-plant continuous transpiration rates of WT 670

(black), MCabi (dark gray) and BSabi (light gray). (F) Transpiration of the different lines normalized to plant weight. 671

Data are means + SE (n = 5 pots for WT and 18–19 for MCabi and BSabi, 6–7 pots from each transgenic line, 4 672

plants in each pot). Letters indicate significant differences at 09:00, 12:00 and 15:00, according to Tukey's test ($P <$ 673

0.05). 674

Figure 4. Osmotic water permeability coefficients (P_f) of palisade-mesophyll and spongy-mesophyll protoplasts. 676

(A) Fluorescent images of the stomata and cross-sections of Arabidopsis leaves expressing a GUS-mCitrine reporter 677

protein under a palisade-specific promoter (IQD22) or (B) a spongy MC-specific promoter (CORI3); bar sizes are 678

shown in the images. (C) P_f distributions of WT palisade (black) and spongy (gray) MC protoplasts generated from 679

IQD22::GUS-mCit and CORI3::GUS-mCit reporter lines, respectively. Dashed lines define the 'tail' of the top 10% 680

outstanding P_f values of the 'high-P_f' for both types of protoplasts (>28 μm s^{-1} for palisade and >47 μm s^{-1} for 681

spongy). Means $\pm$ SE of 90% of the populations are also shown. M – median. (D) Volumes of palisade (gray) and 682

spongy (black) protoplasts. The box height shows 25–75% of the data range, the symbols indicate individual data 683

values, the solid line indicates the median, o indicates the mean and the whiskers delimit ±3 times the interquartile 684

range. (E) P_f response to pre-treatment with 1 μM ABA (1–4 h). One asterisk indicates a p-value less than 0.05 and 685

two asterisks indicate a p-value less than 0.01, according to Student's t-test. The p-values are presented in the figure. 686

Protoplast numbers are shown in each bar. 687

Figure. 5. Osmotic water permeability coefficients (P_f) of MCabi and BSabi protoplasts in response to increasing   688

ABA levels. (A) P_f levels of palisade and spongy mesophyll protoplasts of WT and MCabi#2 (ABA-insensitive)   689

carrying the palisade (*IQD22::GUS-mCit*) and spongy (*CORI3::GUS-mCit*) markers under non-stressed conditions..   690

(B) P_f levels of bundle-sheath protoplasts of WT and BSabi#14 (ABA-insensitive) carrying the BSC marker   691

SCR::GFP under non-stressed conditions. (C) P_f responses of palisade mesophyll of MCabi#2 and (D) of the bundle   692

sheath of BSabi#14 to pre-treatment with increasing levels of ABA: 0, 1 and 50 µM ABA (1–4 h). Data are means   693

$\pm$ SE from at least 4 independent experiments. For A and B, an asterisk indicates a *p*-value of less than 0.05,   694

according to Student's *t*-test. The *p*-values are presented in the figure. For C and D, there were no significant   695

differences between the ABA concentrations, according to Tukey's HSD test. Protoplast numbers are presented   696

inside each bar.   697

Figure 6. Effects of ABA on the symplastic and apoplastic pathways. (A) K_{leaf} of mutants with impaired plasmodesmatal (PD) permeability. (B) DANS assays revealed the cell-to-cell spread of fluorescence in WT, MCabi and BSabi plants that were petiole-fed AXS (dark gray) or AXS +10 μM ABA (light gray). Top of B: Yellow circles show the cell-to-cell spread of the fluorescent dye (bar = 20 μm). The close-up images show that the dye was trapped within the cells (bar = 5 μm). The box height shows 25–75% of the data range, the symbols indicate the individual data values, the solid line indicates the median, o indicates the mean and the whiskers delimit ±3 times the interquartile range. Different letters indicate significant differences between genotypes within a treatment (capital letters for control and lower-case letters for ABA treatment) according to Tukey's test. An asterisk indicates a significant difference between treatments within a genotype according to Student's *t*-test ($P < 0.01$). A: n = 15–17 leaves; B: n = 19 for WT and 24–25 for MCabi and BSabi, 8–9 leaves from each transgenic line.

Figure 7. A suggested model for ABA's regulation of K_{leaf}. (A) Under optimal conditions, water flows across the bundle sheath via the transmembrane water pathway, probably through aquaporins (blue dots; Control Point 1) and proceeds toward the mesophyll. In the mesophyll, water may flow in parallel pathways and move between them. The apoplastic (light blue line) and transmembrane pathways are dominant, while the contribution of the symplastic path (dark red line) is relatively small (Control Point 2). (B) The transmembrane water permeability (P_f) of the spongy mesophyll cells can facilitate the export of water from cells, contributing to transmembrane water flow or directing water toward the cell wall to support apoplastic water flow (Control Point 3) and transpiration (E, dashed blue line). Note that under optimal conditions, most water flows through the spongy mesophyll cells, preserving leaf water status (e.g., high Ψ_{leaf} and turgor). (C) High levels of ABA reduce the P_f of the bundle-sheath cells (blue dot, Control Point 1. Shatil-Cohen et al., 2011). In the immediate term, less water enters the leaf while the amount of water leaving through the stomata remains high, which causes the spongy mesophyll cells to shrink (Canny, 2012) due to their high initial P_f, amplifying the stress signal and initiating ABA production (Bactete, 2021). Higher ABA then decreases the spongy mesophyll cells' P_f and (D) decreases the transmembrane water flow and apoplast wetting (Control Points 2 and 3). Diminished flow over the transmembrane and apoplastic water pathways (blue dots and light blue line, respectively) keeps water inside the cells, while ABA increases the amount of water transported via the symplastic pathway (red line; Control Point 2). The effects of all the above combined with the direct effect of ABA on stomata result in reduced K_{leaf} and E (dashed blue line) as the leaf's water status deteriorates (e.g., reduced Ψ_{leaf} and turgor).

Ache P, Bauer H, Kollist H, Al-Rasheid K a S, Lautner S, Hartung W, Hedrich R
(2010) Stomatal action directly feeds back on leaf turgor: new insights into the
regulation of the plant water status from non-invasive pressure probe
measurements. Plant J **62**: 1072–82

**Álvarez-Arenas TEG, Sancho-Knapik D, Peguero-Pina JJ, Gómez-Arroyo A,
Gil-Pelegrín E** (2018) Non-contact ultrasonic resonant spectroscopy resolves the
elastic properties of layered plant tissues. Appl Phys Lett **113**: 1–5

Attia Z, Dalal A, Moshelion M (2019) Vascular bundle sheath and mesophyll cells
modulate leaf water balance in response to chitin. Plant J 1–10

**Bacete L, Schulz J, Engelsdorf T, Bartosova Z, Vaahtera L, Yan G, Gerhold JM,
Ticha T, Øvstebø C, Gigli-Bisceglia N, et al** (2022) THESEUS1 modulates cell
wall stiffness and abscisic acid production in Arabidopsis thaliana. Proc Natl
Acad Sci U S A. e2119258119

Blackman CJ, Brodribb TJ, Jordan GJ (2010) Leaf hydraulic vulnerability is
related to conduit dimensions and drought resistance across a diverse range of
woody angiosperms. New Phytol **188**: 1113–1123

Blackman CJ, Brodribb TJ, Jordan GJ (2009) Leaf hydraulics and drought stress:
Response, recovery and survivorship in four woody temperate plant species.
Plant, Cell Environ **32**: 1584–1595

Blum A (2009) Effective use of water (EUW) and not water-use efficiency (WUE) is
the target of crop yield improvement under drought stress. F Crop Res **112**: 119–
123

Borsuk AM, Roddy AB, Théroux-Rancourt G, Brodersen CR (2022) Structural organization of the spongy mesophyll. New Phytol **234**: 946–960

Brodribb TJ, Feild TS, Jordan GJ (2007) Leaf Maximum Photosynthetic Rate and Venation Are Linked by Hydraulics. Plant Physiol **144**: 1890–1898

Brodribb TJ, Feild TS, Sack L (2010) Viewing leaf structure and evolution from a hydraulic perspective. Funct Plant Biol **37**: 488–498

Brodribb TJ, Holbrook NM, Zwieniecki MA, Palma B (2005) Leaf hydraulic capacity in ferns, conifers and angiosperms: Impacts on photosynthetic maxima. New Phytol **165**: 839–846

Buckley TN (2015) The contributions of apoplastic, symplastic and gas phase pathways for water transport outside the bundle sheath in leaves. Plant, Cell Environ 7–22

Buckley TN, John GP, Scoffoni C, Sack L (2015) How does leaf anatomy influence water transport outside the xylem? Plant Physiol **168**: 1616–1635

Canny M (2012) Water loss from leaf mesophyll stripped of the epidermis. Funct Plant Biol **39**: 421–434

Canny M, Wong SC, Huang C, Miller C (2012) Differential shrinkage of mesophyll cells in transpiring cotton leaves: Implications for static and dynamic pools of water, and for water transport pathways. Funct Plant Biol **39**: 91–102

Caringella MA, Bongers FJ, Sack L (2015) Leaf hydraulic conductance varies with vein anatomy across Arabidopsis thaliana wild-type and leaf vein mutants. **38(12)**: 2735-2746

Clough SJ, Bent AF (1998) Floral dip: A simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. Plant J **16**: 735–743

Cochard H, Venisse JS, Barigah TS, Brunel N, Herbette S, Guilliot A, Tyree MT, Sakr S (2007) Putative role of aquaporins in variable hydraulic conductance of leaves in response to light. Plant Physiol **143**: 122–133

Coupel-Ledru A, Tyerman SD, Masclef D, Lebon E, Christophe A, Edwards EJ, Simonneau T (2017) Abscisic acid down-regulates hydraulic conductance of grapevine leaves in isohydric genotypes only. Plant Physiol **175**: 1121–1134

Cui W, Lee JY (2016) Arabidopsis callose synthases CalS1/8 regulate plasmodesmal permeability during stress. Nat Plants **2**: 1–15

Dalal A, Shenhar I, Bourstein R, Mayo A, Grunwald Y, Averbuch N, Attia Z, Wallach R, Moshelion M (2020) A telemetric, gravimetric platform for real-time physiological phenotyping of plant–environment interactions. J Vis Exp **2020**: 1–28

Erwee MG, Goodwin PB (1984) Characterization of the Egeria densa leaf symplast: Response to plasmolysis, deplasmolysis and to aromatic amino acids. Protoplasma **122**: 162–168

Fischer RA, Rees D, Sayre KD, Lu ZM, Condon AG, Larque Saavedra A (1998) Wheat yield progress associated with higher stomatal conductance and photosynthetic rate, and cooler canopies. Crop Sci **38**: 1467–1475

Galvez-Valdivieso G, Fryer MJ, Lawson T, Slattery K, Truman W, Smirnoff N, Asami T, Davies WJ, Jones AM, Baker NR, et al (2009) The high light

response in Arabidopsis involves ABA signaling between vascular and bundle sheath cells. Plant Cell **21**: 2143–62

Gosti F, Beaudoin N, Serizet C, Webb A a R, Vartanian N, Giraudat J (1999) ABI1 protein phosphatase 2C is a negative regulator of abscisic acid signaling. Plant Cell **11**: 1897–1910

Griffiths H, Weller G, Toy LFM, Dennis RJ (2013) You're so vein: Bundle sheath physiology, phylogeny and evolution in C3 and C4 plants. Plant, Cell Environ **36**: 249–261

Grondin A, Rodrigues O, Verdoucq L, Merlot S, Leonhardt N, Maurel C (2015) Aquaporins Contribute to ABA-Triggered Stomatal Closure through OST1-Mediated Phosphorylation. Plant Cell **27**: 1945–1954

Grunwald Y, Wigoda N, Sade N, Yaaran A, Torne T, Gosa SC, Moran N, Moshelion M (2021) Arabidopsis leaf hydraulic conductance is regulated by xylem sap pH, controlled, in turn, by a P-type H + -ATPase of vascular bundle sheath cells . Plant J **6**: 301–313

Hacke U (2015) Functional and ecological Xylem anatomy. Funct Ecol Xylem Anat 1–281

Halperin O, Gebremedhin A, Wallach R, Moshelion M (2016) High-throughput physiological phenotyping and screening system for the characterization of plant-environment interactions. Plant J. **89(4)**: 839-850

Hoth S (2002) Genome-wide gene expression profiling in Arabidopsis thaliana reveals new targets of abscisic acid and largely impaired gene regulation in the

abi1-1 mutant. J Cell Sci **115**: 4891–4900 826

Iino M, Ogawa T, Zeiger E (1985) Kinetic properties of the blue-light response of 827
stomata. Proc Natl Acad Sci U S A **82**: 8019–8023 828

Janacek SH, Trenkamp S, Palmer B, Brown NJ, Parsley K, Stanley S, Astley 829
HM, Rolfe SA, Paul Quick W, Fernie AR, et al (2009) Photosynthesis in cells 830
around veins of the C3 plant Arabidopsis thaliana is important for both the 831
shikimate pathway and leaf senescence as well as contributing to plant fitness. 832
Plant J **59**: 329–343 833

Kelly G, Sade N, Doron-Faigenboim A, Lerner S, Shatil-Cohen A, Yeselson Y, 834
Egbaria A, Kottapalli J, Schaffer AA, Moshelion M, et al (2017) Sugar and 835
hexokinase suppress expression of PIP aquaporins and reduce leaf hydraulics 836
that preserves leaf water potential. Plant J **91**: 325–339 837

Kemanian AR, Stöckle CO, Huggins DR (2005) Transpiration-use efficiency of 838
barley. Agric For Meteorol **130**: 1–11 839

Kim YX, Steudle E (2009) Gating of aqaporins by light and reactive oxygen species 840
in leaf parenchyma cells of the midrib of Zea mays. J Exp Bot **60**: 547–556 841

Kim YX, Steudle E (2007) Light and turgor affect the water permeability 842
(aquaporins) of parenchyma cells in the midrib of leaves of Zea mays. J Exp Bot 843
58: 4119–4129 844

Kitagawa M, Tomoi T, Fukushima T, Sakata Y, Sato M, Toyooka K, Fujita T, 845
Sakakibara H (2019) Abscisic Acid Acts as a Regulator of Molecular 846
Trafficking through Plasmodesmata in the Moss Physcomitrella patens. Plant 847

Cell Physiol **60**: 738–751   848

Koornneef M, Reuling G, Karssen CM (1984) The isolation and characterization of   849

abscisic acid-insensitive mutants of Arabidopsis thaliana. Physiol Plant **61**: 377–   850

383   851

Kurihara D, Mizuta Y, Sato Y, Higashiyama T (2015) ClearSee: A rapid optical   852

clearing reagent for whole-plant fluorescence imaging. Dev **142**: 4168–4179   853

Leegood RC (2008) Roles of the bundle sheath cells in leaves of C3 plants. J Exp Bot   854

59: 1663–1673   855

Lloyd JC, Raines CA, John UP, Dyer TAA (1991) The chloroplast FBPase gene of   856

wheat: structure and expression of the promoter in photosynthetic and   857

meristematic cells of transgenic tobacco plants. MGG Mol Gen Genet **225**: 209–   858

216   859

Manohar M, Wang D, Manosalva PM, Choi HW, Kombrink E, Klessig DF   860

(2017) Members of the abscisic acid co-receptor PP2C protein family mediate   861

salicylic acid-abscisic acid crosstalk. Plant Direct **1**: e00020   862

Martre P, Morillon R, Barrieu F, North GB, Nobel PS, Chrispeels MJ (2002)   863

Plasma membrane aquaporins play a significant role during recovery from water   864

deficit. Plant Physiol **130**: 2101–2110   865

Maurel C (1997) Aquaporins and water permeability of plant membranes. Annu Rev   866

Plant Biol **48**: 399–429   867

McAdam SAM, Brodribb TJ (2016) Linking turgor with ABA biosynthesis: 868

Implications for stomatal responses to vapor pressure deficit across land plants.   869

Plant Physiol **171**: 2008–2016

Merilo E, Yarmolinsky D, Jalakas P, Parik H, Tulva I, Rasulov B, Kilk K, Kollist H (2018) Stomatal VPD response: There is more to the story than ABA. Plant Physiol **176**: 851–864

Morillon R, Chrispeels MJ (2001) The role of ABA and the transpiration stream in the regulation of the osmotic water permeability of leaf cells. Proc Natl Acad Sci U S A **98**: 14138–14143

Mott KA (1991) Stomatal responses to humidity in air and helox. Plant, Cell Environ **14**: 509–515

Muries B, Mom R, Benoit P, Brunel-michac N, Cochard H, Drevet P, Petel G, Badel E, Fumanal B, Gousset-dupont A, et al (2019) Aquaporins and water control in drought-stressed poplar leaves: A glimpse into the extraxylem vascular territories. Environ Exp Bot **162**: 25–37

Nardini A, Gortan E, Salleo S (2005a) Hydraulic efficiency of the leaf venation system in sun- and shade-adapted species. Funct Plant Biol **32**: 953–961

Nardini A, Raimondo F, Lo Gullo MA, Salleo S (2010) Leafminers help us understand leaf hydraulic design. Plant, Cell Environ **33**: 1091–1100

Nardini A, Salleo S, Andri S (2005b) Circadian regulation of leaf hydraulic conductance in sunflower (Helianthus annuus L. cv Margot). Plant, Cell Environ **28**: 750–759

Negin B, Moshelion M (2017) The advantages of functional phenotyping in pre-field screening for drought-tolerant crops. Funct Plant Biol **44**: 107–118

Negin B, Yaaran A, Kelly G, Zait Y, Moshelion M (2019) Mesophyll abscisic acid restrains early growth and flowering but does not directly suppress photosynthesis. Plant Physiol **180**: 910–925

Pantin F, Monnet F, Jannaud D (2013) The dual effect of abscisic acid on stomata. New Phytol **197**: 65–72

Pilot G, Stransky H, Bushey D (2004) Overexpression of GLUTAMINE DUMPER1 leads to hypersecretion of glutamine from hydathodes of Arabidopsis leaves. Plant Cell **16**: 1827–1840

Pou A, Medrano H, Flexas J, Tyerman SD (2013) A putative role for TIP and PIP aquaporins in dynamics of leaf hydraulic and stomatal conductances in grapevine under water stress and re-watering. Plant, Cell Environ **36**: 828–843

Prado K, Boursiac Y, Tournaire-Roux C, Monneuse J-M, Postaire O, Da Ines O, Schäffner AR, Hem S, Santoni V, Maurel C (2013) Regulation of Arabidopsis leaf hydraulics involves light-dependent phosphorylation of aquaporins in veins. Plant Cell **25**: 1029–39

Prado K, Maurel C (2013) Regulation of leaf hydraulics: from molecular to whole plant levels. Front Plant Sci **4**: 255

Procko C, Lee T, Borsuk A, Bargmann BOR, Dabi T, Nery JR, Estelle M, Baird L, O'Connor C, Brodersen C, et al (2022) Leaf cell-specific and single-cell transcriptional profiling reveals a role for the palisade layer in UV light protection. Plant Cell 1–19

Richards R a (2000) Selectable traits to increase crop photosynthesis and yield of

grain crops. J Exp Bot **51 Spec No**: 447–458   914

Roberts IM, Boevink P, Roberts AG, Sauer N, Reichel C, Oparka KJ (2001)   915
Dynamic changes in the frequency and architecture of plasmodesmata during the   916
sink-source transition in tobacco leaves. Protoplasma **218**: 31–44   917

Rockwell FE, Holbrook NM, Stroock AD (2014) The competition between liquid   918
and vapor transport in transpiring leaves. Plant Physiol **164**: 1741–58   919

Sack L, Holbrook NM (2006) Leaf hydraulics. Annu Rev Plant Biol **57**: 361–81   920

Sack L, John GP, Buckley TN (2018) ABA accumulation in dehydrating leaves is   921
associated with decline in cell volume, not turgor pressure. Plant Physiol **176**:   922
489–493   923

Sade N, Shatil-Cohen A, Attia Z, Maurel C, Boursiac Y, Kelly G, Granot D,   924
Yaaran A, Lerner S, Moshelion M (2014) The Role of Plasma Membrane   925
Aquaporins in Regulating the Bundle Sheath-Mesophyll Continuum and Leaf   926
Hydraulics. Plant Physiol. doi: 10.1104/pp.114.248633   927

Sade N, Shatil-Cohen A, Moshelion M (2015) Bundle-sheath aquaporins play a role   928
in controlling Arabidopsis leaf hydraulic conductivity. Plant Signal Behav **10**: 1–   929
4   930

Schulz A (1995) Plasmodesmal widening accompanies the short-term increase in   931
symplasmic phloem unloading in pea root tips under osmotic stress. Protoplasma   932
188: 22–37   933

Scoffoni C, Albuquerque C, Brodersen CR, Townes S V., John GP, Bartlett MK,   934
Buckley TN, McElrone AJ, Sack L (2017) Outside-Xylem Vulnerability, Not   935

Xylem Embolism, Controls Leaf Hydraulic Decline during Dehydration. Plant
Physiol **173**: 1197–1210

Scoffoni C, Rawls M, Mckown A, Cochard H, Sack L (2011) Decline of leaf
hydraulic conductance with dehydration: Relationship to leaf size and venation
architecture. Plant Physiol **156**: 832–843

Scoffoni C, Sack L (2017) The causes and consequences of leaf hydraulic decline
with dehydration. J Exp Bot **68**: 4479–4496

Scoffoni C, Vuong C, Diep S, Cochard H, Sack L (2014) Leaf shrinkage with
dehydration: Coordination with hydraulic vulnerability and drought tolerance.
Plant Physiol **164**: 1772–1788

Shapira O, Khadka S, Israeli Y, Shani U, Schwartz A (2009) Functional anatomy
controls ion distribution in banana leaves: significance of Na+ seclusion at the
leaf margins. Plant Cell Environ **32**: 476–85

Shatil-Cohen A, Attia Z, Moshelion M (2011) Bundle-sheath cell regulation of
xylem-mesophyll water transport via aquaporins under drought stress: a target of
xylem-borne ABA? Plant J **67**: 72–80

Shatil-cohen A, Moshelion M (2012) The bundle sheath role as xylem-mesophyll
barrier. Plant Signal Behav **7**: 1088–1091

Shatil-Cohen A, Sibony H, Draye X, Chaumont F, Moran N, Moshelion M (2014)
Measuring the osmotic water permeability coefficient (pf) of spherical cells:
Isolated plant protoplasts as an example. J Vis Exp 1–13

Shivaraj SM, Sharma Y, Chaudhary J, Rajora N, Sharma S, Thakral V, Ram H,

Sonah H, Singla-Pareek SL, Sharma TR, et al (2021) Dynamic role of aquaporin transport system under drought stress in plants. Environ Exp Bot **184**: 104367

Sinclair TR (2018) Effective water use required for improving crop growth rather than transpiration efficiency. Front Plant Sci **9**: 1–8

Sperry JS, Venturas MD, Anderegg WRL, Mencuccini M, Mackay DS, Wang Y, Love DM (2017) Predicting stomatal responses to the environment from the optimization of photosynthetic gain and hydraulic cost. Plant Cell Environ **40**: 816–830

Steudle E, Peterson C a. (1998) How does water get through roots? J Exp Bot **49**: 775–788

Sussmilch FC, Brodribb TJ, McAdam SAM (2017) Up-regulation of NCED3 and ABA biosynthesis occur within minutes of a decrease in leaf turgor but AHK1 is not required. J Exp Bot **68**: 2913–2918

Tholen D, Boom C, Zhu XG (2012) Opinion: Prospects for improving photosynthesis by altering leaf anatomy. Plant Sci **197**: 92–101

Torne T, Grunwald Y, Dalal A, Yaaran A, Moran N (2021) Stress is basic: 1 ABA alkalinizes both the xylem sap and the cytosol by inhibiting their P-type H +-ATPase 4 and stimulating their V-type H +-ATPase. bioRxiv 2021-03

Tylewicz S, Petterle A, Marttila S, Miskolczi P, Azeez A, Singh RK, Immanen J, Mähler N, Hvidsten TR, Eklund DM, et al (2018) Photoperiodic control of seasonal growth is mediated by ABA acting on cell-cell communication. Science

(80-) **360**: 212–215   980

Tyree MT, Nardini A, Salleo S, Sack L, El Omari B (2005) The dependence of leaf   981
hydraulic conductance on irradiance during HPFM measurements: any role for   982
stomatal response? J Exp Bot **56**: 737–44   983

Voicu MC, Cooke JEK, Zwiazek JJ (2009) Aquaporin gene expression and   984
apoplastic water flow in bur oak (Quercus macrocarpa) leaves in relation to the   985
light response of leaf hydraulic conductance. J Exp Bot **60**: 4063–4075   986

wang X, sager R, Cui W, Zhang C, Lu H, Lee JY (2013) Salicylic acid regulates   987
plasmodesmata closure during innate immune responses in Arabidopsis. Plant   988
Cell **25**: 2315–2329   989

de Wit C (1958) Transpiration and crop yields. Versl Landbouwk Onderz **64.6** 18–20   990

Wysocka-Diller JW, Helariutta Y, Fukaki H, Malamy JE, Benfey PN (2000)   991
Molecular analysis of SCARECROW function reveals a radial patterning   992
mechanism common to root and shoot. Development **127**: 595–603   993

Xiong D, Nadal M (2020) Linking water relations and hydraulics with   994
photosynthesis. Plant J **101**: 800–815   995

Yaaran A, Moshelion M (2016) Role of aquaporins in a composite model of water   996
transport in the leaf. Int J Mol Sci. **17(7)**: 1045   997

Yaaran A, Negin B, Moshelion M (2019) Plant Science Role of guard-cell ABA in   998
determining steady-state stomatal aperture and prompt vapor-pressure-deficit   999
response. Plant Sci **281**: 31–40   1000

Yael Grunwald, Sanbon Chaka Gosa1, Nava Moran MM (2021) Out of the blue;
Phototropins of the leaf vascular bundle sheath mediate the blue light regulation
of the leaf hydraulic conductance. The Plant Cell **34.6**: 2328-2342.

Yoo CY, Pence HE, Hasegawa PM, Mickelbart M V. (2009) Regulation of
Transpiration to Improve Crop Water Use. CRC Crit Rev Plant Sci **28**: 410–431

Yoshida T, Christmann A, Yamaguchi-Shinozaki K, Grill E, Fernie AR (2019)
Revisiting the Basal Role of ABA – Roles Outside of Stress. Trends Plant Sci
24: 625–635

Zait Y, Shapira O, Schwartz A (2017) The effect of blue light on stomatal
oscillations and leaf turgor pressure in banana leaves. Plant Cell Environ **40.7**:
1143–1152

Zavaliev R, Levy A, Gera A, Epel BL (2013) Subcellular Dynamics and Role of
Arabidopsis β -1 , 3-Glucanases in Cell-to-Cell Movement of Tobamoviruses.
Molecular Plant-Microbe Interactions **26**: 1016–1030

Zwieniecki MA, Brodribb TJ, Holbrook NM (2007) Hydraulic design of leaves:
Insights from rehydration kinetics. Plant, Cell Environ **30**: 910–921