Connectomics of the *Octopus vulgaris* vertical lobe provides insight into conserved and novel
principles of a memory acquisition network

Authors
Flavie Bidel[1,*,+], Yaron Meirovitch[2,*,+], Richard Lee Schalek[2], Xiaotang Lu[2], Elisa Catherine Pavarino[2],
Fuming Yang [2], Adi Peleg[2], Yuelong Wu[2], Tal Shomrat[3], Daniel Raimund Berger[2], Adi Shaked [1], Jeff
William Lichtman[2], Binyamin Hochner[1,+]
7
8 [1] Department of Neurobiology, Silberman Institute of Life Sciences, The Hebrew University, Jerusalem,
9 Israel
10 [2] Department of Molecular and Cellular Biology, Harvard University, Cambridge, MA, United States
11 [3] Faculty of Marine Sciences, Ruppin Academic Center, Michmoret, Israel
12
*Equal contribution.
+Corresponding Authors: bidel.flavie@gmail.com; yaron.mr@gmail.com; benny.hochner@mail.huji.ac.il

Summary

We present the first analysis of the connectome of the vertical lobe (VL) of *Octopus vulgaris,* a brain
structure mediating acquisition of long-term memory in this behaviorally advanced mollusk. Serial section
electron microscopy revealed new types of interneurons, cellular components of extensive modulatory
systems and multiple synaptic motifs. The sensory input to the VL is conveyed via ~1,800,000 axons that
sparsely innervate two parallel and interconnected feedforward networks formed by the two types of
amacrine interneurons (AM), simple AMs (SAMs) and complex AMs (CAMs). SAMs make up 89.3% of
the ~25,000,000 VL cells, each receiving a synaptic input from only a single input neuron on its non-
bifurcating primary neurite, suggesting that each input neuron is represented in only ~12 SAMs. This
synaptic site is likely a "memory site" as it is endowed with LTP. The CAMs, a newly described AM type,
comprise 1.6% of the VL cells. Their bifurcating neurites integrate multiple inputs from the input axons
and SAMs. While the SAM network appears to feedforward sparse "memorizable" sensory representations
into the VL output layer, the CAMs appear to monitor global activity and feedforward a balancing inhibition
for "sharpening" the stimulus-specific VL output. While sharing morphological and wiring features with
circuits supporting associative learning in other animals, the VL has evolved a unique circuit that enables
associative learning based strictly on feedforward information flow.

INTRODUCTION

The advanced learning capabilities of cephalopod mollusks like *Octopus vulgaris* are comparable to those in some vertebrates (Sanders 1975; Mather 1995; Zarrella et al. 2015; Schnell et al. 2016; Hanlon and Messenger 2018). As the last common ancestor of cephalopods and vertebrates existed more than 500 million years ago (Packard 1972; Vitti 2013), the octopus provides a unique opportunity to study the independent evolution of nervous systems subserving sophisticated cognitive abilities (Hochner et al. 2006; Shomrat et al. 2015; Hochner and Glanzman 2016; Turchetti-Maia et al. 2017). The vertical lobe (VL), the highest structure in the hierarchy in the central nervous system of the octopus, is regarded as a high order integrative center involved in visually evoked associative learning and memory acquisition (Boycott and Young 1955; Sanders 1975; Young 1979; Fiorito and Chichery 1995; Hochner 2010, 2012, 2013; Shomrat et al. 2015; Shigeno et al. 2018; Turchetti-Maia et al. 2017). It is exceptionally large, containing 25 million neurons, about half the neurons in the central nervous system. This number is not much smaller than the number of neurons in the human hippocampus (~40 million, West and Gundersen, 1990) and about 4 orders of magnitude greater than the number of neurons in the mushroom body (MB) of *Drosophila* (e.g. Aso et al., 2014) and three orders of magnitude greater than that in the locust MB (Jortner et al. 2007).

Extensive neuroanatomical data on the VL were provided by the work of J. Z. Young and E. G. Gray in the 1960-70's using light and electron microscopy (Gray and Young 1964; Gray 1970; Young 1971). Based on estimated neuron numbers they proposed that the VL network is extremely simple, composed of only two types of morphologically distinguishable neurons: 25 million small amacrine interneurons (AM) and 65,000 large efferent neurons (LN). The main sensory input to the VL is conveyed via 1.8 million axons projecting from the median superior frontal lobe (SFL). These axons compose the SFL-VL tract that runs in the outer neuropil of the five VL lobuli. Each of these axons makes *en passant* synapses with a then unknown number of the AMs. These 25 million AMs converge sharply at a ratio of about 400:1 to innervate the LNs, whose axons were assumed to be the only output of the VL. Young pointed out that the pattern of SFL innervation of the AMs resembles the connectivity matrix in other learning and memory networks like the insect MB and the mammalian hippocampus (Young 1995).

This gross connectivity scheme provided insights that guided more recent physiological studies on the VL. Physiological experiments in the brain-slice preparation (Hochner et al. 2003, 2006) revealed that the *en passant* SFL-to-AM innervation is glutamatergic. It shows a robust activity-dependent NMDA-independent long-term potentiation (LTP) of glutamate release from the SFL presynaptic terminals (Hochner et al. 2003). The AM-to-LN connections are excitatory cholinergic synapses (Shomrat et al. 2011). Recently it was discovered that LTP at the SFL-to-AM synapse is maintained by a novel nitric oxide (NO)-dependent

'molecular switch' mechanism in which NO switches ON NO-synthase (NOS) through a positive feedback

loop of NO-dependent NOS activation (Turchetti-Maia et al. 2018).

Although Gray and Young provided a detailed description of the main neuronal elements constituting the

VL network, the inability to reconstruct individual neurites limited the possibility of providing a detailed

circuit connectivity. Indeed, their observations reported unclassified cell types, thus suggesting that the VL

network is likely composed of more than the 3 neuronal elements they characterized (SFL, AMs and LN).

Previous findings in our group also hinted that the VL is more complex. Firstly, intracellular recording from

LNs revealed that some receive inhibitory inputs (Shomrat et al., 2011). Secondly, a recent wide-ranging

histolabeling study suggested that the AMs may be divided into at least two groups, one cholinergic and

the other GABAergic (Stern-Mentch et al. 2022).

To understand the octopus VL circuitry at a higher resolution the current study utilizes recent advances in

our ability to elucidate synaptic wiring diagrams at the nanoscale level using electron microscopy (EM).

Widely applied to various species and tissues this methodology has contributed much to understanding the

organization of neural networks (Kasthuri et al. 2015; Lee et al. 2016; Morgan et al. 2016; Ryan et al. 2016;

Motta et al. 2019; Li et al. 2020; Macrina et al. 2021; Meirovitch et al. 2021; MICrONS Consortium et al.

2021; Verasztó et al. 2020; Witvliet et al. 2021; Shapson-Coe et al. 2021). Here we sought to produce a

synapse-level wiring diagram of the *Octopus vulgaris* VL from serial EM by tracing out all pre- and

postsynaptic partners of dozens of main afferents of the VL (SFL projecting axons) and AMs. Indeed,

although we found that the gross anatomical organization of octopus VL fits the previous anatomical and

physiological model described above, the nanoscale analysis revealed new types of neurons and

connections that contribute profoundly to a deeper understanding of the functional organization of the

octopus VL and how it implements associative learning and memory.

Our data show that, firstly, in contrast to what was previously thought, the AM interneurons are segregated

into two distinct groups. The majority (~89.3 %) of the VL cells bodies are simple AMs (SAM) that send a

single non-bifurcating neurite into the neuropil. A small fraction (~1.6 %) forms a very small group of

newly discovered complex AMs (CAM) with a branching dendritic tree. Secondly, we found that the SAMs

receive only one synaptic input from a single SFL axon. We propose that this synaptic connection may

serve as a "memory site" as it shows a robust activity-dependent LTP following tetanization of the SFL

axonal tract (Shomrat et al 2011). Thirdly, the segregated SFL-to-SAM expansion format suggests that

each of the 1.8 million SFL neurons is represented extremely sparsely in only around 12 of the 22.3 million

SAMs. Fourthly, in sharp contrast to the SAMs, the CAMs integrate dozen to hundreds of inputs,

suggesting that these interneurons integrate broader incoming SFL activity. As the SAMs and CAMs

converge onto the same output layer of LNs processes, our findings suggest the VL appears to be comprised

of two interconnected feedforward networks - one excitatory, originating from the SAMs (each representing
one SFL) and a second, a parallel feedforward network, likely inhibitory, originating from the CAMs (each
representing several SFLs). We demonstrated that distinct groups of SAMs and CAMs neurites fasciculate
together to form a columnar organization in which each column constitutes canonical microcircuitry that
may function as an "association module", associating SFL sensory features. Lastly, we described an
additional afferent to the VL and two extrinsic putative neuromodulation systems. These interact with the
feedforward networks and thus are likely involved in regulating the activity state of the network and in
conveying reward and punishments signals.

RESULTS

The VL dataset and analysis

The region of interest (ROI) is in the lateral lobule of the VL (**Figure 1A**). We cut, collected and imaged

892 serial sections (30nm section thickness) at 4nm/pixel to generate a traceable 3D image stack covering

a volume of 260x390x27 µm (**Figure 1B; Figure 1- video supplement 1**; 'connectome dataset' available

at https://lichtman.rc.fas.harvard.edu/octopus_connectomes).

110
Figure 1. Reconstruction of octopus VL circuits using volume EM. (A) Low resolution image of the
octopus central nervous system. The VL lies dorsally in the supraesophageal part of the central brain. The
region of interest marked within it (ROI, white rectangle) was scanned with a scanning electron microscope
(SEM) at 4 nm/px. (B) 30 nm sections were aligned in a traceable 3D stack (260x390x27µm. See also
Figure 1–video supplement 1) for machine learning (ML)-aided by manual segmentation and synapse
annotation. (C) The 516 reconstructed cell processes colored according to cell type (See **Figure 1–video**
supplement 2 for all cell types). Total neurite length of the reconstructed cells was 12.8 cm. Reconstructed
cells are superimposed on a single EM image. Inset: spatial distribution of the 13434 postsynaptic sites
annotated on the 516 cells presented as a heat-map. Abbreviations: BL, basal lobe; IFL, inferior frontal
lobe; SFL, superior frontal lobe; SPL, subpedunculate lobe; VL, vertical lobe. See also **Figure 1–figure**
supplement 3 and **Figure 1–figure supplement 4**.
122
To classify the cells we used morphological features (*e.g.,* main neurite orientation, presence of varicosities,

branching pattern), ultrastructural features (*e.g.,* vesicle size, mitochondria shape, cytoplasm staining,

organelles) and connectivity assisted by the 'blueprint' established by Gray and Young (Gray and Young

1964; Gray 1970; Young 1971). Chemical synapses were identified based on features common to

invertebrate synapses - the presence of a cluster of synaptic vesicles (sv) or 'cloud' associated with an active

zone near the presynaptic membrane (Gray 1970; Ryan et al. 2016; White et al. 1986; Zheng et al. 2018;

Meinertzhagen 2019; Witvliet et al. 2021). Synapses were annotated if this accumulation of sv persisted

across at least three sections (> 90 nm). We classified synapses as fast chemical if the presynaptic

component contained sv of 30-60 nm diameter and modulatory synapses when the synaptic component

contained large vesicles (>60nm) (Messenger 1996; Leng and Ludwig 2008; van den Pol 2012; Witvliet et

al. 2021). Consistent with previous studies (Gray and Young 1964; Gray 1970), we did not observe any

obvious postsynaptic specialization (*i.e.,* postsynaptic density) at any chemical synapse and we found no

evidence of gap junctions.

We reconstructed 516 cell processes with a total length of 12.8 cm. The reconstructed cell processes were

classified into 10 cell types (**Figure 1–video supplement 2**) with 13,434 postsynaptic (**Figure 1C, heat**

map) and 2,472 annotated presynaptic sites. Among these 10 cell types, only 7 were involved in the VL

connectivity and therefore were analyzed in depth (see **Figure 1–figure supplement 3** for the 3 cell types

not discussed here). For characterization of the 7 cell types, we reconstructed throughout the volume at least

30 processes per cell type and annotated all their pre- and postsynaptic sites. Additionally, a subset of

cellular compartments and organelles (*i.e.,* cell bodies, nuclei and presynaptic varicosities) were separately

segmented to allow morphological quantification. The core of our connectivity reconstruction included

annotation and neurite reconstructions of all the synaptic partners of a subset of randomly selected 'seed'

cells (Morgan et al., 2016): SFL axons (35 of 84 SFL axons reconstructed) and SAM cells (28 of 108 SAMs

reconstructed).

We divided the analysis according to the three VL regions: 1) The outer cell body cortex; 2) the central

neuropil, and 3) the SFL tract, or outer neuropil, where axons projecting from the SFL are arranged in a

tract forming a ring around the outer zone of neuropil (Gray and Young, 1964; Young, 1971, Stern-Mentch

et al 2022). Note that because most of the cells are not fully contained in the reconstructed volume, it is

impossible to assess the full collection of synapses between pairs of neurons. We therefore used "synaptic

site" to refer to a single anatomical release site or a single synapse.

The VL connectome is characterized by two parallel and interconnected feedforward networks

Figure 2 summarizes the 7 identified neuronal cell types and their connectivity (see also **Figure 2–figure**

supplement 1). None of the 84 reconstructed SFL axons bifurcated within the ROI (~260 μm long)

suggesting a very low rate of bifurcation of the SFL axons, if at all, as suggested by Young (1971). Each

SFL axon innervated *en passant* two distinct sub-populations of AMs; a very large population of

morphologically simple cells, which we termed 'simple amacrine' (SAM) and a small population of

morphologically complex cells that we termed 'complex amacrine' cells (CAM) (**Figure 2 and Figure 3**).

160
Figure 2. The 7 neuronal cell types classified in the ROI and their wiring diagram. The middle panel
shows the wiring diagram, and the other panels depict the 7 neuronal cell types together with the spatial
distribution of their postsynaptic sites (red puncta) and presynaptic sites (blue puncta). The approximate
position of the SFL tract area (green) is shown for orientation and demarcating the outer from the inner
neuropil. In the central panel each gray solid edge represents fast chemical synapses and the arrowhead the
direction of these synapses; line width indicates relative number of synapses calculated from the number of
connections annotated (noted next to each outgoing arrow). Orange dashed edges indicate chemical
synapses observed with modulatory processes. Edges were represented only if more than 10 synapses were

annotated. Note that the connectivity is strictly feedforward (p<0.05). See **Figure 2–figure supplement 1**
for neurite level analysis.
171
The SAM and CAM cell bodies are intermingled in the cortex and their primary neurites run together in the
same bundles, also previously referred to as the *AM tract* (Gray and Young 1964; Gray 1970; Young 1979).
These bundles form a *quasi*-columnar organization toward the neuropil (**Figure 6A, 6B and Figure 7A**).
The SAMs and CAMs showed quite differed connectivity, morphology, and ultrastructure (**Figure 6A and**
6B). The SAMs are monopolar interneurons with a single non-bifurcating neurite. Their cell bodies are
small (mean ± std; $207.7 ± 26.1 \mu m^3$; n=69) and contain almost no cytoplasm. In contrast, the CAMs have
a complex bifurcating dendritic tree, a larger cell body (mean ± std; $337.2 ± 41.6 \ \mu m^3$; n=16) and
cytoplasmic volume (**Figure 4**). Unexpectedly, we found that each SAM receives input from only one SFL
axon (See SAM section for details). In contrast, each CAM integrates synaptic inputs from a dozen to
hundreds of SFL axons and SAMs (**Figure 6B**). Both the CAMs and SAMs consistently converged onto
the LN processes (**Figure 7**). Overall, the transmission from the main afferents (SFLs) is entirely mediated
via the SAMs and CAMs into their shared target neurons - the dendritic processes of the LNs.

A second group of afferent axons innervated exclusively the CAMs. We termed these 'CAM-input-neurons'
(CIN; **Figure 8**). Even though CIN function has yet to be established, their ultrastructure is reminiscent of
the nerve endings which Gray and Young speculated may carry "pain" signals to the VL. They described
them as having neurofilaments ending shortly before the synaptic boutons which were filled with small
vesicles, a structure we also identified in our data. We also identified that the likely fast transmission circuit
(**Figure 2** panels framed with solid line) is overlaid by a diffuse mesh of two types of likely
neuromodulatory processes (NM and AF), each with a typical morphology, ultrastructure, and cell-specific
connections (**Figure 2**; panels framed with broken line). First, the 'neuromodulatory fibers' (NMs) manifest
a large variety of processes with large varicosities filled with large vesicles (60-200nm) that often do not
show the typical structure of a synaptic connection (detailed below). Second, the 'ascending fibers' (AFs)
climb from the neuropil up to the cortex through the *AM tracts* (i.e., a dense bundle of primarily SAM and
CAM neurites; **Figure 9**). A stereotyped feature of all AFs was their "menorah"-shaped bifurcating pattern,
close association with SAM neurites and their intense innervation by the SAM. The ultrastructure and
connectivity of these cells revealed two distinct subcategories, one likely neuromodulatory (AF-Nm,
Figure 9C) and the other yet undefined (AF-Mix, **Figure 9D**). Both the AF-Nm and the NM boutons were
filled with large vesicles and usually lacked synaptic outputs, suggesting that these are release sites of
neuromodulators or neuropeptides that are involved in extrasynaptic volume transmission (Bentley et al.,
2016). Since none of the AF and NM neurites were associated with a reconstructed cell body, we

hypothesize that AFs and NMs are external to the VL and may belong to the widespread ascending fibers
entering the VL from below that have been previously described (Young, 1971).

Overall, we show that the 7 cell types form a feedforward neural network in which only the AFs display
more than 10 reciprocal connections within a cell type (AF-to-AF, **Figure 2**). Analysis at the single neurite
level showed that the number of recurrent connections (cycles of connectivity) in the ROI was significantly
smaller than that in wiring diagrams built from the same set of neurites and synapses, but with otherwise
random directionality determining the pre- and postsynaptic neurons in each connection ($p<0.05$) (see
Methods). Moreover, the connections involved in feedback connectivity (*i.e.,* a cycle of connectivity) were
all associated with reciprocal connections which were rare.

Each SFL axon innervates the SAMs and CAMs via two specialized bouton types

A set of 35 seed SFL afferents was used for a detailed description of the connectivity pattern of the SFL
and its postsynaptic targets (**Figure 3A**). Each SFL axon was found to make two distinct types of synaptic
contacts depending on the target interneuron. First, an average of 28 ± 12 % (mean ± std) of the synapses
of each SFL axon (n=45/161, N=35 SFL) were made on SAMs (**Figure 3B** and **Figure 3–figure**
supplement 1). This synaptic connection is unique as each SAM received only a single SFL input. The
characteristic synaptic structure consisted of an especially large SFL bouton (LB, 6.41 ± 2.1 µm³)
innervating *en passant* a specialized vesicle-filled postsynaptic compartment protruding from the SAM
primary neurite that partially wrapped the SFL LB (**Figure 3A, Figure 3–figure supplement 1** and **Figure**
4). We termed these postsynaptic structures 'palms' due to their somewhat flat and concave morphology.
Second, an average of 66 ± 11% of the synapses (109/161 synapses; N= 35 SFL) of each SFL axon were
made on CAM dendritic endings. These CAM endings were a variety of twig-like structures formed by
variably long (approximately $1 - 30$ µm) thin protrusions emerging from thicker CAM dendritic branches
(**Figure 3A**). These postsynaptic profiles were electron-lucent and filled with polyribosomes (**Figure 3B**).

Unlike the single SFL-to-SAM connection, one SFL axon could make several synapses on the same CAM.
These SFL-to-CAM synapses showed two different arrangements. Either the synapses involved a large
presynaptic bouton (LB) of the SFL manifesting a multisynaptic glomerulus-like structure (**Figure 3A and**
Figure 5) or the synapses involved only a single CAM output with a significantly smaller presynaptic SFL
bouton (SB, 0.95 ± 0.88 µm³; **Figure 3A**). The volume of all boutons (N=295) of all reconstructed SFL
axons (N=84) showed a bimodal distribution corresponding to these two different bouton types (**Figure**
3C).

Figure 3. The SFL axons synapse onto the two populations of amacrine interneurons (SAMs and CAMs) via two distinct populations of presynaptic boutons. (A) Reconstruction of a single SFL axon (green) and its 7 *en passant* outputs within the reconstructed volume: 5 CAMs (various blue hues) and 2 SAMs (yellow). (B) Quantification of the synaptic outputs of the seed SFL axons (n=35) to SAMs (via large boutons; LB), CAMs (via small boutons; SB) and onto unidentified cell types (UN). Upper left shows distribution of connections in percent. On the right high resolution EM images of the SFL synaptic outputs onto SAM and CAM. 1nm/px, dwell time: 200 μs. Scale bar, 500 nm. Empty triangle and asterisk are respectively pre- and postsynaptic profiles. The whisker plot shows the relative density of the different presynaptic boutons along the 35 seed SFLs axons (C) Example of the distribution along an SFL axon segment of presynaptic outputs onto two SAMs (yellow triangles) and CAMs (light blue triangles). The histogram displays the bimodal distribution of bouton volumes of 84 SFL axons into two populations of small boutons (SB-blue) and large boutons (LB-green). The plot of presynaptic sites per SB and LB reveals

245 their monosynaptic and multisynaptic nature, respectively. Asterisks indicate statistical significance ***
246 p<0.001, **<0.01. Abbreviations: sv, synaptic vesicles; AZ-like, active zone-like; Mit, mitochondria, Cis,
247 cisternae; tER, transport endoplasmic reticulum. See also **Figure 3–figure supplement 1** and **Figure 3–**
248 **video supplement 2**.
249
250 **The number of CAMs is extremely low relative to that of SAMs**

251 We were next interested in the relative abundance the SAMs and CAMs. Since we found significant

252 differences in volume and shape of the various cell types, we reconstructed 1482 cell bodies without their

253 projections and used volume and shape descriptors to build a Support Vector Machine classifier to estimate

254 the number of different types of neurons in the VL (**Figure 4**).

255
Figure 4. Cell body composition of VL cortex
A. A subset of 1472 densely packed cell bodies were reconstructed and automatically classified based on
their morphology; SAM, yellow; CAM, blue hues; LN, pink; Others, purple. B-C. The classifier was
designed to capture an observed correlation between cell type and cell body geometry roundish, relatively
smaller cell bodies were associated with SAMs (vol $207.7 \pm 26\mu m^3$, mean $\pm$ STD), often larger and slightly
more elongated cell bodies with CAMs (vol $337.1671 \pm 41.6 \mu m^3$), considerably larger cell bodies with
highly variable volumes with LNs (volume range $464 - 2850 \mu m^3$), and elongated cell bodies with other
cell types including glia and uncharacterized cells. A Support Vector Machine (SVM) was trained to
separate 4 cell categories SAM, CAM, LN, and Others on a set of 110 manually classified cells based on 3
features: volume, sphericity, and surface area, and was validated on an independently prepared validation
set with 47 cells (94.55% training accuracy with ~4.01% error on validation set all associated with SAMs).
267
268 We estimated that 89.3% of the cell bodies in the sampled volume were SAM cell bodies and 1.6% were

269 CAMs (**Figure 4A**) with error rate on validation set smaller than 4.1% (see Methods). Assuming that the

VL is roughly composed of 25 million of cells bodies (Young 1963) and a similar proportion over the VL gives an estimate of ~22.3 ± 1.2 million SAMs and an estimate that the cortex contains fewer than 1.43M CAMs. As we did not observe any false negative or false positive associated with CAMs, the total number is likely close to the classifier's best estimate of 400,000 CAMs which can be further analyzed once this sparse set of cells is sampled on larger EM volumes. Assuming the SFL axons do not bifurcate in the MSF tract, then due to the one-to-one SFL-to-SAM innervation pattern the 1.8 million SFL axons fan-out onto the SAMs at an expansion ratio of about 1-to-12. Since each SAM is activated by a single SFL axon, then information from one SFL axon is represented at the SAM input layer in a non-overlapping manner in only about 12 ± 3.46 SAMs (assuming a Poisson distribution with an average of 12 LB). To estimate the distribution of LBs along the SFL axons we calculated the number of LBs per 100 µm in the 84 reconstructed SFL axonal segments. The results indicate an average and SD of 1.03 ± 0.73 LBs per 100 µm long segments suggesting highly sparse and variable (CV=0.77) distribution along the tract. If this distribution in maintained along large portions of the axon, then the LBs stretch over an average length of 1270 µm which grossly fits the length the of the SFL tract in our preparation (**Figure 1A**). This, in turn, suggests that the LB outputs into the 12 ± 3.46 SAMs is likely maintained along the entire length of the SFL axon rather than being confined in specific area.

The SAMs, CAMs and SFL large boutons are interconnected within synaptic glomeruli

The SAMs-to-CAMs connection forms the interconnection between the two parallel feedforward networks (**Figure 2**). These connections were found locally at a single large synaptic glomerulus encapsulating the SFL LB, the SAM palm and the CAM twig endings (**Figure 5**). Within this glomerulus these 3 neurons were repeatedly connected with multisynaptic motifs: monad, SFL-to-SAM (**Figure 5A**); dyad, SFL-to-CAM and SFL-to-SAM (**Figure 5B**), serial synapses, SFL-to-SAM-to-CAM, and triad where the same SFL presynaptic terminals innervate an adjacent SAM palm and a CAM twig ending (**Figure 5C**), as well as higher order polyads.

This arrangement fits the classical organization of synaptic glomeruli where a number of synapses are congregated in spatially compact structures, so that they may influence each other's functioning (Morgan and Lichtman 2020). Among these arrangements, the SAM palm, packed with vesicles, appears to be a unique bifunctional pre- and postsynaptic compartment, as most of the SAM palms formed serial synaptic connections (82%), receiving a synapse from a single SFL bouton and providing a synapse to 1 - 3 postsynaptic compartments, often twig endings of distinct CAMs. Analyzing a large population of 'long' SAM neurites (SAMs whose primary neurite was fully contained in the SFL tract area, N=56) revealed that each SAM exhibited only a single palm along its inwardly directed primary neurite (N=46/56), rarely two

palms (N=2/56) (**Figure 6C**) and each palm received input from only one SFL large presynaptic bouton
(50/50 palms received a single SFL input).

Figure 5. SFL large boutons (LBs) are found in multisynaptic motifs, forming a synaptic glomerulus
involving serial connectivity with both SAMs and CAMs. Subcellular 3D reconstruction illustrating the
main motifs involving the large varicosity of the SFL axon, which contains glutamate and makes *en passant*
synapses. (A) monad (B) dyad and (C) triad. Note that in (C) only the readily releasable pool of synaptic
vesicles is shown for clarity as otherwise the large "reserve" pool in the SFL LB (>5000 synaptic vesicles)
would obscure other components of the image. The SFL LB were always partly wrapped by at least one
SAM palm (yellow) and they often additionally synapsed onto a postsynaptic CAM twig-like ending (blue).
313

To better describe the structural properties of this unique palm structure we prepared a smaller dataset of

16x16x3.5 µm imaged at 1nm/px (high magnification dataset, see Methods). This revealed the complex

presynaptic structure of the SAM palm which could contain up to 1500 synaptic vesicles (36.8 ± 7.7 nm)

often tightly packed and resembling the "dense-walled" vesicles described by Gray and Young (1964). The

structure included numerous elongated mitochondria and a large network of electron-lucent tubular

organelles (**Figure 5**). These organelles could be small, disconnected units (cisternae, Cis) or large and

continuous, likely part of the endoplasmic reticulum (ER). The main body of the ER usually faced the pre-

and postsynaptic sites, possibly playing a role in vesicle recycling (Ramírez and Couve 2011; Weigel et al.

2021). We also occasionally observed what appeared to be exocytosis at the ER membrane, suggesting the

possibility of a Golgi-like extended role in the synaptic vesicle machinery. It is still not clear whether the

Cis (up to 15 observed in one palm) were part of the larger ER network or were isolated entities that could,

for instance, play a role in keeping vesicles close to the active zone like synaptic ribbons (Dowling et al.

1966; Parsons and Sterling 2003) or the T-bar in Drosophila neurons (Scheffer et al. 2020).

The SAMs distribute their single SFL input onto the LNs, CAMs and AF

We next focused on a detailed reconstruction and proofreading of 28 SAM seed cells. As mentioned above,

each seed SAM neuritic trunk received only one *en passant* SFL LB input (**Figure 6A**). The LB input was

always located at the SAM palm (**Figure 5 and Figure 6A1, C**). Analyzing the neuritic trunks of SAM

seed cells showed that 93% of the seed SAMs appeared to have no other inputs beside the SFL LB.

The SAM outputs were spread evenly along the length of the neuritic trunk. These synaptic outputs targeted

3 cell types (**Figure 6F**). As described by Young, (1971) and confirmed here, many SAMs converged onto

the dendrite-like processes of the LN (LNp), mostly within the central neuropil but occasionally at the level

of the SFL tract (**Figure 6A4 and Figure 6E**).

However, the SAM inputs to the LNp represented only 35% of the SAMs' presynaptic sites (N=109/271).

The remaining connections are described here for the first time. 25% of the SAM outputs targeted the

CAMs, while 26% innervated the processes of the climbing AFs, suggesting the SAMs directly innervate

neuromodulatory processes. The SAM-to-CAM synapses were formed by a variety of dilatations of the

SAM neuritic trunk containing clear, small vesicle-filled profiles synapsing onto a clear postsynaptic profile

of the CAM twig (**Figure 6A2**). This was mainly observed within the SFL tract and in the upper neuropil.

In contrast, the SAM-to-AF synapses (**Figure 6A3**) were spread throughout the dataset, including the

cortex, and the postsynaptic element (AF) was a bouton often filled with dense-core vesicles (**Figure 9**). In

the neuropil and the SFL tract these synapses were arranged in a rosette-like structure in which 2 to 5

adjacent SAMs synapsed locally onto the same AF bouton (**Figure 9**). Strikingly, within the cortex, the

SAM soma occasionally synapsed onto AF varicosities, forming a heterodox somato-neuromodulatory

synapse (**Figure 9A**). We also found rare cases of other synaptic outputs to NM (N=2/271), unidentified

processes (N=19/271) and only one case of recurrent connection of a SAM onto itself (N=1/271) (**Figure**

6F). Reciprocal synaptic contact between SFL and SAM as well as AF and SAM were occasionally

observed.

Figure 6. Stereotypic SFL ('mononeural')-input to the SAMs and multiple ('polyneuronal') inputs to the CAMs. (A,B) 2D projection of a reconstructed SAM (A) and CAM (B), superimposed on a reconstruction of the SFL tract (green). Triangles and puncta represent presynaptic output and postsynaptic input sites, respectively, color coded according to pre- or postsynaptic partner. The insets show the corresponding EM cross-section of each type of synapse. The morphological arrangement of the corresponding synaptic connection is depicted in the colored connectivity scheme next to each image and morphological reconstructions in the corresponding cells' color code (scale bar=500nm). (C) Analysis of the number of inputs onto the SAM 'palm' and its pre- and postsynaptic morphological specialization. 82%

of SAMs had only one palm, each receiving just one input (red bar). Each SAM innervated up to 3 targets
(blue bars). Only SAMs with their primary neurite fully contained in the SFL tract area were considered
(N=56). (D) The majority of the SAMs had only one input located at the palm. (E) Spatial localization of
the input/output of the 28 SAM seed cells within the ROI. Note the restriction of SFL-to-SAM inputs to the
area of the SFL tract. (F) The connectivity diagram of the 28 SAM seed cells shows the fraction of the
inputs (red) and outputs (blue) by neuron type. (G) Histogram showing the wide distribution of the number
of synaptic inputs to the CAMs (dozens to hundreds). (H) The connectivity diagram of 53 CAMs shows the
percentage of total inputs to and outputs from the corresponding cell type (color coded. See B1,2,3,4 for
details). Most of the CAMs' presynaptic partners were not reconstructed but identified based on
characteristic ultrastructure profiles. Profiles which could not be unambiguously identified were classified
as unidentified (Un). (I) Spatial distribution of the CAM inputs and outputs within the ROI. See also **Figure**
6–figure supplement 1, **Figure 6–video supplement 2**, **Figure 6–figure supplement 3** and **Figure 6–**
figure supplement 4.
377
Morphology and connectivity of the CAMs

The CAMs were probably not previously identified due to their small number relative to the huge number

of SAMs. Gray (1970) and Young (1971) may therefore have confused their dendritic arborizations and

electron-lucent ultrastructure within the SFL tract (**Figure 6A2,B1,B2 and B3**) with that of the LN. To

understand the potential functionality of these newly discovered interneurons within the VL circuitry we

annotated all their pre- and postsynaptic sites. Because the SFL, CINs, SAMs AFs and NMs all have

distinctive ultrastructure "fingerprints" (*e.g.,* cytoplasm staining, vesicle size, vesicle type, mitochondria

shape; see Methods), we were able to define their presynaptic partners without reconstruction.

The CAM neurites have 3 distinct regions: the *primary neurite* ramifies into strictly postsynaptic *dendritic*

branches in the SFL tract. The dendritic branches turn into neuritic *axons* in the deeper neuropil (**Figure**

6B and Figure 6- figure supplement 2). Synapses were located preferentially on numerous small twigs

which branched off the thicker dendritic branches. We used 'twig' to define CAM second-order dendritic

branches because their base lies on a microtubule-containing main thick branch(Schneider-Mizell et al.

2016) but, unlike spines that have a maximal length of less than 3 µm (Leiss et al. 2009), the twigs can be

much longer and show a complicated branching pattern. The CAMs showed variable dendritic branch

patterns and orientation, possibly suggesting the existence of CAM subtypes (**Figure 6- figure supplement**

2 and 3). Additionally, the morphology of the CAM twig endings was highly variable with at least 5

categories all somewhat resembling dendritic spine enlargements (**Figure 6B,C**): flatfoot, long neck

mushroom head, talon, palm-like and a claw-like endings that strikingly resembled the claw-like structure

wrapping the projection neurons in the insect MB (Leiss et al., 2009; Schürmann, 2016; Yasuyama et al.,

2002).

In contrast to the SAMs (**Figure 6–figure supplement 1**), each CAM interneuron received dozens to

hundreds of synaptic inputs (**Figure 6G and Figure 6–figure supplement 2 and 3**); the most extreme case

displayed 514 postsynaptic sites over a neurite length of 1.4µm. The CAMs integrate a combination of

inputs (**Figure 6H**), mostly from the SFL (48%, **Figure 6B2 and Figure 6–figure supplement 2G and**

3E) and the SAMs (42%, **Figure 6B1 and Figure 6–figure supplement 2F and 3F**). 5% of the inputs to

the CAMs originated from CINs (**Figure 6B3 and Figure 6H**). The CINs appeared to exclusively innervate

the CAMs with their inputs sparsely distributed over the CAM twig endings (**Figure 8**). Despite the CAMs

showing high diversity in primary neurite projection and dendritic arborizations, all received inputs and

innervated the same cell types and at similar locations within the VL (**Figure 6B and Figure 6–figure**

supplement 3 and 4). The CAMs' distal axonal processes innervated the LNps with many *en passant*

boutons (**Figure 6B4**).

Different convergence patterns of SAMs and CAMs into the LNs

Both the SAMs and CAMs converged onto the LNs, the output cells layer of the VL (**Figure 2, Figure 6**

and Figure 7). To investigate the pattern of LN innervation we first annotated all the synapses of 14 LNs

with reconstructed cell bodies and those on putative LN processes (see below, N=24). The LN cell bodies

(dia. > 15 µm) and their radially directed large primary neurite were poorly contained within our analyzed

volume. However, some of them (N=3/14) gave off dendritic collaterals within the SFL tract area where

they were solely postsynaptic to SAMs (**Figure 2**) and to other unidentified processes (with a total of 115

individual synaptic inputs on the LN processes in the SFL tract).

As suggested by the physiological experiments (Shomrat et al. 2011), we did not find any direct connections

between these LN dendritic collaterals and SFL axons. Among the 7 cell types thoroughly studied, we

consistently observed that the SFL, SAMs, CAMs, CINs, AFs and NMs processes all displayed an EM

profile filled with vesicles within the neuropil. Therefore, by elimination, we considered all electron-lucent

postsynaptic profiles within the neuropil as putative LN dendritic processes (LNp). The 24 LNps that we

examined branched profusely within the neuropil (**Figure 7A**) and were heavily innervated. The histogram

in **Figure 7D** shows that the LNps displayed the largest density of incoming synapses in the dataset with a

median of 162 postsynaptic sites/100 µm, far larger than the densely innervated AF-Nms (41 postsynaptic

sites/100 µm) and the CAMs (25 postsynaptic sites/100 µm that are restricted to the SFL tract; **Figure 7D**).

The SAM and CAM innervation of the LNps often formed rosette-like structures in which 2 - 5 or more

AMs made adjacent synapses onto an LN dendritic process (**Figure 7E**).

Figure 7. Patterns of synaptic convergence of SAMs and CAMs onto LNs. (A) 3D reconstruction of the LN cell bodies, and processes (LNp) (magenta hues) superimposed on SAMs (yellow) and CAMs (cyan). (B) Distributions of the 184 postsynaptic sites on one LN dendrite colored according to the legend. Note the very few CAM inputs relative to those of the SAMs. (C) Convergence of two bundles of intermingled SAM and CAM neurites onto an LNp shown in B. (D) The LNps displayed the greatest postsynaptic density among all dendritic elements. (E) High resolution EM illustrating a rosette-like structure formed by the convergence of 3 SAMs into one LNp. Empty triangle and asterisk are pre- and postsynaptic profiles, respectively. Resolution: 1nm/px. Dwell time: 200 μs.

We next estimated the degree of convergence of the SAMs and CAMs onto the LNs. For this we reconstructed and annotated all the neurites synapsing onto an 81 μm length of LNp dendrite (**Figure 7 B, C**). As shown in **Figure 7C** all the CAM and SAM inputs originated from two distinct fasciculate neuritic bundles (blue and green) that included 160 SAM neurites and only six CAMs. Approximately 90% of the

inputs on LNps were from SAMs (N=160/184 postsynaptic sites), only 3% were from CAMs (n=6/184 postsynaptic sites). **Figure 6B** shows that the CAM and SAM inputs were distributed in a "salt and pepper" manner with the CAM inputs distributed very sparsely and evenly over the LN dendritic process. These data indicate that the VL output neurons integrate massive information from the SAMs and only sparse input from the CAMs. The spatial distribution of the CAM inputs to the LNp suggests a role of the CAM output in adjusting the membrane potential of a large LN dendrite area. Such spatial distribution would be most effective if the CAMs are inhibitory interneurons as we suggest below (see Discussion).

The CINs afferents innervate exclusively CAMs

CIN processes spread throughout the neuropil, but many ran parallel to the SFL axons (**Figure 8A**). Their boutons were filled with many small, rounded mitochondria and the smallest vesicles observed in the VL (≈30nm). Their presynaptic boutons displayed the greatest variability in size with volumes ranging from 0.09 to 29.50 µm^3 (median ± iqr; 1.65 ± 2.23).

Fifty-five of these fibers were reconstructed and their synaptic sites annotated. The CINs were almost exclusively presynaptic (534 sites) with only very few postsynaptic sites (17 sites). All these presynaptic sites were located on large boutons (N=239) that usually made polyadic synapses with several CAMs' postsynaptic structures (*e.g.,* 3 in **Figure 8B** and 4 in **Figure 8C**) with a median of 2.3 ± 1.5 (median ± iqr) synaptic sites per bouton (**Figure 8D**), the most extreme case being 8 presynaptic sites for a single bouton. This synaptic structure may allow a single CIN terminal to influence several CAMs simultaneously.

The CIN innervate *en passant* mainly the CAMs (225 synaptic connections) and only occasionally the AFs (**Figure 2 and Figure 2–figure supplement 1**). Note that the CAMs projected long thin side branches with a morphological specialization at the tip (*i.e.,* long neck with mushroom head, **Figure 8C**) where it received a synaptic input from the CIN bouton. The CINs displayed the highest presynaptic site density found within the VL with a median of 11 presynaptic sites/100µm, twice as high as presynaptic density on SFL axons (**Figure 8D**).

Figure 8. CAM-specific afferents (CINs) innervate exclusively CAMs via large multisynaptic boutons. (A) 3D reconstruction of a CIN (pink) superimposed on the CAMs, their main postsynaptic partners (blue hues). (B, C) 3D arrangement of a large CIN presynaptic bouton innervating multiple CAM dendrites (3 in B; 4 in C). CAM twig endings varied in shape resembling flattened dendritic spine enlargements. As indicated in C their structure can be divided into at least 5 classes. Right inset, 3 consecutives near cross-sections through a CIN bouton and its 3 postsynaptic targets. The CIN profile contained the smallest vesicles, about 30nm dia., of the entire dataset. (D) CIN had the highest presynaptic site density of the dataset. Error bar=SEM.

The neuromodulatory ascending fibers (AF) receive massive synaptic inputs from the SAMs

The AF were highly recognizable in our dataset with their menorah shape, climbing through the neuropil into the cortex while intermingling with SAM and CAM neurites (**Figure 9A**). Their processes displayed large boutons along their length (9.4 boutons/100µm). The reconstruction of the synaptic partners of the SAM seed cells revealed that the AFs receive massive input from the SAMs (**Figure 6F**).

Mapping the AF synaptic sites revealed two clear subcategories of AF based on their connectivity (**Figure 9B**) and EM profile (**Figure 9C,D**). We termed one class AF-neuromodulatory (AF-Nm, n=14). In this class, on the 468 boutons reconstructed we annotated 1460 postsynaptic sites and only 3 presynaptic sites, with a median of 3.2 postsynaptic site per bouton. Each bouton formed a rosette structure of axo-axonal synapses where several SAMs made adjacent synapses onto one AF bouton (**Figure 9C**). The postsynaptic sites were filled mostly with large dense core vesicles (60-80 nm), suggesting a neuromodulatory nature. The synaptic inputs of the SAMs into the AF-Nm boutons may suggest SAMs-induced local extrasynaptic release of neuromodulator (see Discussion). The AFs seem morphologically equivalent to the serotonergic processes ascending into the cell cortex found in immunohistological studies (Shomrat et al. 2010; Shigeno and Ragsdale 2015) and also to tyrosine hydroxylase (catecholaminergic) labeled processes (Stern-Mentch et al., 2022).

In contrast, the second class of AF, the AF-mix (N=18), were more diverse with boutons filled with vesicles of varying shape and size (30-90 nm; N=519), including small dark core vesicles, flat clear core vesicles and larger vesicles. Unlike the AF-Nm, the AF-mix were both presynaptic (median of 0.2 presynaptic sites/bouton) and postsynaptic (median of 0.6 postsynaptic sites/bouton) and occasionally displayed reciprocal contact with a SAM (**Figure 9D**). The diverse nature of the synaptic vesicles in the two subtypes suggests that they convey different neuromodulatory signals and the reciprocal connection with the SAMs suggest their involvement in SAM -induced local release of neuromodulators.

Figure 9. Overlay of two modulatory systems, the ascending fibers (AFs) and widespread neuromodulatory fibers (NMs). (A) 3D reconstruction of a large network of AFs (orange) intermingling with a SAM neuritic bundle (yellow). The AFs showed varicosities along their length filled with large

vesicles and were mainly postsynaptic to the SAMs (C,D) A somatic synaptic input from SAM to AF was occasionally observed (lower left inset). (B) The distribution of the number of pre- and postsynaptic sites per varicosity (blue and red, respectively) revealed two types of AF processes shown in C and D. (C) The AF-Nm type (orange) were exclusively unidirectional synapses forming a rosette-like structure where 2-5 SAMs made adjacent synapses onto one AF-Nm postsynaptic varicosity (asterisk). (D) The AF-mix (light orange) were more diverse. Their boutons were filled with vesicles of various shapes and sizes, ranging from small dark core vesicles to flat clear core vesicles. Unlike the AF-Nm, they could be bidirectional, both pre- and postsynaptic (triangles and puncta, respectively) and occasionally displayed reciprocal contact with a SAM as shown in the EM. (E) 3D reconstruction of the widespread NM fibers color coded by subtype and superimposed on populations of SAMs (yellow) and CAMs (blue). At least 3 distinct subtypes were distinguishable based on vesicle sizes (EM insets: clear core low density, clear core high density, dense core). (F) 3D reconstruction of NM-to-CAM synapses. The NM made multisynaptic contacts. The right inset shows a representative EM of a neuromodulatory synapse. Scale bar EM sections, 500 nm. See **Figure 9–figure supplement 1**

The ubiquitous NM processes appear to be involved in global neuromodulatory signaling via extrasynaptic transmission

A large variety of processes displayed the largest dense vesicles (60-200nm), which appear similar to vesicles thought to be neuromodulatory in other invertebrates (Messenger 1996; Witvliet et al. 2021). The processes were widespread within the neuropil and the SFL tract and their lack of specific orientation made it impossible to assume their origin (**Figure 9E**). At least 3 distinct profiles were clearly distinguishable by vesicle size and appearance, showing either a low density of clear core vesicles, a high density of clear core vesicles or very large dense core vesicles. The boutons of each process contained only one type of vesicle, which were also present within the fiber. Reconstructions of NM processes (N=88) showed that most of the boutons had neither inputs nor outputs (464/632), suggesting that they were neuromodulatory afferents which exert their effects through extrasynaptic volume transmission. Yet, these cells occasionally synapsed with a variety of postsynaptic partners including CAMs, SAMs AFs, NMs themselves, and LNs (**Figure 2 and Figure 2 - figure supplement 1**). Multisynaptic contacts were occasionally observed, for instance, with CAMs (**Figure 9F**), but the density of synaptic sites per bouton was rather low, 52 presynaptic sites and 250 postsynaptic sites on 632 boutons. Accordingly, NM processes may be involved primarily in extrasynaptic transmission and thus may convey a global modulatory signal transmitted though action potentials propagating along them.

DISCUSSION

The novel connectivity and anatomical features discovered here profoundly advance our understanding of the functional organization of the learning and memory circuits in the vertical lobe (VL) of *Octopus vulgaris*. The EM volume reconstruction revealed several novel features that, on one hand, show the VL is composed of several more components than previously thought. On the other hand, these novel cell types and their connectivity help us understand the functional organization of the VL as a learning and memory acquisition network.

The amacrine interneurons in the VL input layer are not a homogeneous group

The first striking finding was that 89% of the ~25 million VL cells (**Figure 4**) send a simple non-bifurcating neurite into the neuropil and were therefore termed simple AMs (SAMs; **Figure 6A**). The SAMs are highly input specific as each received *en passant* input from only a single SFL axon via large presynaptic boutons (LB) (**Figure 3A, Figure 5 and Figure 6A1**). If the pattern of one-to-one innervation is maintained throughout the whole SFL tract and if, as generally believed and as found here, the SFL axons bifurcate very rarely, then every SFL neuron is represented by only around 12 ± 3.46 SAMs (22×10^6 SAMs / 1.8×10^6 SFLs). Moreover, the LBs seem to be sparsely distributed along the SFL axon (1.03 ± 0.73 LBs per 100 µm; see Results), suggesting that each SFL neuron is represented in the SAMs in a highly sparse and non-overlapping fashion.

This extremely low and random topographical expansion of SFL inputs to SAMs is unique for input circuitry. In *Drosophila* mushroom bodies, for example, the expansion ratio is multiplied by a factor of about 6 as six projecting neurons converge onto each Kenyon cell (Li et al. 2020) and in locust the expansion ratio is 400 (Jortner et al. 2007). The unique low expansion of information in the VL is achieved by the formation of a huge number of SAMs, 3-4 orders of magnitude more than the number of the homologous Kenyon cells in the insect mushroom bodies. This mode of expansion may be related to the organization of the octopus visual system where each SFL axon carries visual information that has been highly processed by the 120×10^6 neurons of the two optic lobes and then possibly further classified into lower dimensional visual features in the only 1.8×10^6 cells in the SFL (Young 1971). This unusual expansion pattern resembles the 50×10^{12} granular cells in the human and mice cerebellum - each receiving only 4-6 inputs from the mossy fibers at the expansion ratio of 1-to-200 (D'Angelo 2013; Kawato et al. 2021). An additional striking similarity is that these unique synaptic connections in the cerebellum undergo activity-dependent presynaptic LTP; (Mapelli and D'Angelo 2007) like the SFL-to-SAM synapse in the octopus VL.

Our second novel discovery is that the AMs included a small group of a new class of AM neuron, only 1.6% of the AMs in the VL cortex (**Figure 4**). These neurons showed a bifurcating dendritic structure and

were therefore termed complex AMs (CAMs, **Figure 6B**). The primary neurite segments of the SAMs and

CAMs travel together in the AM bundles within the VL cortex, but the non-bifurcating SAM neuritic trunks

cross the SFL tract to pass into the inner neuropil, while the CAM neurites bifurcate substantially within

the SFL tract (**Figure 2 and Figure 6B**). The CAMs receive inputs *en passant* from multiple SFL axons

either directly or indirectly via a serial connection from the SAMs (**Figure 5 and Figure 6B**). Thus, while

each SAM represents a single SFL neuron, the CAMs integrate information from dozens to hundreds of

SFL neurons together with multiple inputs from nearby SAMs. In addition, as discussed below, the CAMs

receive inputs from afferents from a yet unidentified origin, which exclusively innervate CAMs (CIN,

Figure 6B3 and Figure 8).

The SFL axons innervate the SAMs and CAMs via different classes of presynaptic boutons

SAMs and CAMs are innervated by SFL axons via small (SFL SB) or extremely large boutons (SFL LB –

Figure 3). The LBs are always associated with at least one SAM palm and may additionally innervate a

CAM twig within a dyadic or triadic arrangement (**Figure 5**). By contrast, SBs are always associated with

a single postsynaptic CAM compartment (**Figure 3**). This suggests that the two different types of SFL axon

terminal are functionally specialized. Such functional segregation is also seen in the hippocampal mossy

fiber connections where small boutons target GABAergic interneurons, whereas large boutons contact CA3

pyramidal cells and hilar mossy cells (Acsády et al. 1998). The latter large synapse shares several synaptic

features with the octopus VL synapses (Hochner et al. 2003) – NMDA-independent, presynaptically

expressed LTP and an extremely large range of presynaptic short-and long-term synaptic gain modulations

(Nicoll and Schmitz 2005; Henze et al. 2000; Hainmueller and Bartos 2020; Restivo et al. 2015). The large

synaptic boutons here, like the mossy fiber terminals, may provide an increased dynamical range of synaptic

gain control needed for certain forms of plasticity involving presynaptic modulation of transmitter release

(Orlando et al. 2021; Hainmueller and Bartos 2020; Restivo et al. 2015).

The single SFL synaptic input into the SAM may serve as a gain control site of the stimulus-specific

input to the VL

Each SAM receives synaptic input from only a single SFL neuron, raising the question of the role of such

apparently non-integrating interneurons. Physiological results indicate that the glutamatergic SFL-AM

synapse is the site of activity-dependent LTP in the VL (Hochner et al. 2003; Shomrat et al. 2011; Turchetti-

Maia et al. 2018). The large volume of the SFL-LB relative to that of the SFL-SB found here (on average

6-fold larger) suggests that the recorded synaptic field potential (fPSP) mainly reflects the synaptic input

from the SFL LB to the SAMs. The morphological dimensions of the SAMs revealed here (*e.g.,* a 166µm

long 0.5µm diameter neuritic trunk emerging from a 7µm diameter cell body) together with previous

electrophysiological measurements (>1GΩ input resistance measured at the cell body), suggest that the

inexcitable SAMs (Hochner et al. 2003, 2006)are electrotonically compact (estimated length constant λ >
147 μm) (Hochner and Spira 1987), so that the glutaminergic excitatory postsynaptic potential from the
SFL LB input site is likely sufficient to activate voltage-dependent transmitter release down the SAM
neurite without needing a propagating action potential. These inexcitable properties, which differ drastically
from the excitable properties of the insects Kenyon cells (e.g. Demmer and Kloppenburg 2009), explain
nicely the linear input/output relationship of the SAMs (Shomrat et al. 2011). Since the activity-dependent
changes at the SFL-SAM synapse involve pre- and postsynaptic mechanisms (see below), we suggest that,
rather than being simple relay interneurons, the SAMs provide a synaptic "gain-control relay" of the SFL-
LB input into the CAMs and LNs.

How can specificity and associativity be achieved within these one-to-one SFL-to-SAM connections?

Specificity and *associativity* are essential properties for all associative learning networks (Kandel et al.,
2012; Malenka 2003). To understand how they are achieved within the unique wiring of the VL it is
important to recall that LTP expression in the VL involves presynaptic facilitation of transmitter release
from the SFL boutons. This was suggested to be mediated by an activity-dependent persistence elevation
of NO concentration in the vicinity of the SFL-SAM connection. NO alone cannot induce LTP. Only the
association of strong activation of an SFL- SAM synapse switches ON the NO-dependent NOS activation,
leading to long-term LTP *maintenance* (Turchetti-Maia et al. 2018). The good correspondence of the
SAMs' morphological characteristics with AM-specific labeling by NADPH-diaphorase and cChAT
(choline acetyl transferase) suggests that NOS activity is strongly expressed along the entire length of the
cholinergic SAMs (Stern-Mentch et al. 2022). Accordingly, NO, which can readily diffuse short distances
through cell membranes (Agnati et al., 1995; Garthwaite, 2016, 2020; Turchetti-Maia et al., 2017, 2018)
and which is generated in the SAMs following LTP induction, diffuses retrogradely from the SAM palm
into the presynaptic SFL LB mediating the NO-dependent presynaptic *expression* of LTP. The
characteristic morphological arrangement described here with the SAM palm partly wrapping the SFL LB,
ensures synaptic specificity as the steep NO concertation gradient will affect only these closely attached
presynaptic varicosities (Hall and Garthwaite 2009). The SFL- SAM synapses are organized in a compact
synaptic glomerulus that may facilitate the diffusion of NO within a triadic synaptic organization including
the CAMs. This suggests that NO diffusing from a single postsynaptic SAM palm also modulates the SFL
SB input into the CAMs.

What are the possible mechanisms of LTP *associativity*? The persistence elevation of NO concentration in
the VL is maintained, at least in about half of SFL-SAM synapses, through a molecular switch consisting
of a positive feedback loop of NO-dependent reactivation of NOS (Turchetti-Maia et al. 2018). This
"molecular switch" provides a tentative molecular model for associativity, whereby volume-type

transmission of NO (Garthwaite 2016) between adjacent activated SAMs may mutually facilitate
associative switching ON of NOS in these simultaneously activated SAMs. Our results highlighted that
many SAM neurites converging within the neuropil to form compact bundles of thin intermingling
processes travelling together (**Figure 7A**). Such spatially closed arrangements of thin neurites (e.g., **Figure**
7A, C), which express intense NOS activity (see Turchetti-Maia et al., 2018, Stern-Mentch et al 2022), can
favor mutual NO effects among adjacent SAM neurites.

The CAMs as a putative feedforward inhibitory network balancing the parallel SAM feedforward
excitatory network.

The newly discovered CAMs comprise only 1.6% of the VL cortex. An inhibitory nature of CAMs is
supported by GABA labeling of putative CAM dendritic processes at the level of the SFL tract and in
varicosities in the neuropil (Stern-Mentch et al. 2022). Also, inhibitory CAMs would explain the IPSPs or
EPSP followed with short delay by an IPSP recorded in some of the LNs in response to SFL tract stimulation
(**Figure 7 - figure supplement 1**).

Although the CAMs show morphological diversity (**Figure 6- figure supplement 2 and 3**), they share
similar connectivity, integrating multiple inputs from the SFLs, SAMs and CAM-specific inputs (CIN) in
their bifurcating dendrites (**Figure 6B**). The CAMs innervate the LN dendritic processes only sparsely,
unlike the dense LN innervation by the SAMs (**Figure 7B and 7D**). This innervation pattern suggests that
the LNs integrate stimulus-specific excitatory inputs from many SAMs, while the CAMs, which integrate
ubiquitous and nonselective inputs, are optimally located for inhibitory balancing the overall SAM
excitatory inputs. Yet, the LNp are innervated by only a few CAMs, suggesting that the CAMs inhibitory
inputs are activated simultaneously over a large LN dendritic area causing widespread postsynaptic
inhibition of the LNp.

If the CAMs are indeed inhibitory interneurons, then our results strongly suggest the VL network is
organized into two interconnected parallel feedforward networks, in which the SFL axons conveying
sensory inputs 'fan-out' onto SAMs (in ~1/12 ratio) and 'fan-in' (~4/1) onto the CAMs. Both the SAMs
and CAMs feedforward onto the LNs in the VL readout layer (schematically shown in **Figure 10A**). This
type of organization may sharpen the stimulus specific SAM inputs to the LNs by subtracting the global or
background activity monitored by the inhibitory CAMs. This type of organization may highlight a universal
inhibitory balancing organization in networks with different functions. A similar organization is found in
cortical circuits representing sparse, odor-specific, activity where the sparse activity is generated through
summing unbalanced widespread and broadly tuned inhibition with odor-specific excitation (Poo and
Isaacson 2009). In the octopus VL, such sharpening mechanism may ensure recognition of learned visual
scenarios even under variable light intensities. Theoretical studies have demonstrated the importance of

implementing feedforward inhibition parallel to the feedforward excitation to balance the stable
propagation of an activity wave (Litvak et al., 2003). Feedforward inhibition also appears important for
balancing networks' input/output relationships to avoid attraction of network activity into either up or down
states (Ferrante et al., 2009). Finally, long-term balancing inhibition is believed to be important in ensuring
network homeostasis especially in networks involved in learning and memory, like the hippocampus, where
activity balances may be perturbated by long-term activity-dependent plasticity (Turrigiano and Nelson
2004).

Three candidates for conveying modulatory signals.

Learning and memory networks are usually supervised by neuromodulation. Early anatomical studies by
Young and colleagues proposed several candidates, and our study has refined Young's broad classification
of fibers ascending to the VL. We have described 3 distinct classes of putative extrinsic processes:
neuromodulatory fibers (NM), ascending fibers (AF) and CAM-specific input fibers (CIN). These could
play a role in conveying information on whether the current environment is punitive or rewarding,
reinforcing, or suppressing associative learning.

The rich and widely distributed NMs intermingle with the fast transmitter feedforward network
(schematized in **Figure 10B**). The fibers form a mesh in the neuropil and may be associated with regulation
of overall VL activity. Their exceptionally large boutons, frequently lacking the typical release sites, are
filled with a variety of large dense-core vesicles of the type usually storing and releasing neuropeptides and
non-classical neurotransmitters (Zhu et al. 1986; Leng and Ludwig 2008; van den Pol 2012; Witvliet et al.
2021). A variety of neuropeptides and hormones have been identified in the *Octopus vulgaris* VL -
serotonin, dopamine, octopamine, FMRFamide-related peptides, vasopressin, oxytocin, octopressin, Y-
peptide and more (Shomrat et al. 2010; Shigeno and Ragsdale 2015; Winters et al. 2020; Stern-Mentch et
al. 2022). Tyrosine hydroxylase (TH) - labeled processes show a wide distribution in the VL neuropil
similar to that of the NM processes, suggestive of a neuromodulatory catecholamine innervation (Stern-
Mentch et al. 2022).

As schematized in **Figure 10B**, the AF fibers form close associations with the SAMs (**Figure 9A**), while
the CIN fibers appear to exclusively innervate the CAMs (**Figure 8, schematized in Figure 10A**). The AF
boutons form a rosette-like structure in which several SAMs make adjacent synapses onto one AF bouton,
mostly filled with dense-core vesicles. Such a connectivity motif between neurons and neuromodulatory
processes suggests that co-activated SAMs may activate an AF bouton that, in turn, releases a
neuromodulator. This mechanism can function locally, especially if the AF processes are inexcitable. Local
release of the neuromodulator can heterosynaptically induce synaptic plasticity, similar to the well-known
heterosynaptic serotonin-dependent synaptic plasticity, particularly in the gastropods mollusc *Aplysia*

(Bailey et al. 2000; Kandel et al. 2014). The reciprocal SAM AF connection in the VL network is a plausible

synaptic motif that may suggest a "self" LTP-induction mechanism as shown schematically in Figure 10B.

The feasibility of such a mechanism is hinted at by 5-HT causing both short-term synaptic facilitation of

the input to the AMs and reinforcing LTP induction (Shomrat et al. 2010). Dopamine and octopamine also

affect synaptic physiology but, in contrast to 5-HT, they suppress the transition to LTP maintenance

(unpublished results). Immunohistochemical studies have also revealed a rich distribution of serotonin-

(Shomrat et al. 2010; Shigeno and Ragsdale 2015) and catecholaminergic-rich fibers (Stern-Mentch et al.

2022) that resembles the AF distribution described here, suggesting possible AF involvement in modulation

of the activity-dependent LTP. While Shomrat et al. (2010) found serotonergic immunoreactive cell bodies

outside the VL and in the subvertical lobe, Stern-Mentch et al. (2022) described a group of TH-labeled cell

bodies organized as a "deep nucleus" at the border between the SFL and VL. This suggests that, while

incoming serotonergic innervation may provide supervision signals, like reward and punishment (Turchetti-

Maia et al. 2017), a dopaminergic system may be involved in the regulation of the internal state of the SFL-

VL system.

The exclusive innervation of the very small group of CAMs by a distinct group of processes (CIN) must be

functionally significant (schematically shown in **Figure 10A**). The CIN processes resemble fibers that

Young speculated were ascending fibers carrying "pain signals" from the arms and mantle (Gray, 1970;

Gray and Young, 1964). Unlike the NMs and AFs that barely display any presynaptic profiles, the CIN

outputs to the CAMs have the typical morphological features of fast transmitter synapses (**Figure 8**). The

presynaptic features are similar to vertebrate presynaptic terminals, with small electron-dense synaptic

vesicles, which are occasionally associated with inhibitory synapses (McDonald et al. 2002; Kasthuri et al.

2015). We therefore speculate that, in addition to feedforward excitation from the SFLs and SAMs, the

CAMs integrate inhibitory inputs from the CIN. The CINs may provide negative feedback adjusting the

level of LNs output to that of its targets, such as the subvertical lobe to where the LNs axons project and

which Young described as the origin of the "pain fibers." In the framework of the VL model (Shomrat et

al. 2015; Turchetti-Maia et al. 2017), in which the GABAergic LN output (Stern-Mentch et al. 2022)

inhibits the octopus' tendency to attack, the CIN inhibitory input onto the CAMs would increase the LN

output, resulting in suppression of the attack behavior. We believe this is what occurs when training

octopuses on a passive avoidance task (Shomrat et al. 2008).

Figure 10. Schematic representation of the currently known circuit architecture of the VL. (A) Fast transmitter connectivity. The parallel SFL axonal tract (green) conveys classified sensory information via *en passant* glutamatergic (glut) synapses to a large group of SAMs (yellow) and a very small group of CAMs (cyan). Each SAM has one palm on its neurite innervated by a single large SFL bouton (SFL LB). This connection is endowed with short- and long-term plasticity. The CAMs integrate ongoing activity through multiple inputs from the SFL, SAMs and CINs (light pink). The SAMs and CAMs converge onto the LN processes (LNp, magenta) forming two parallel and interconnected feedforward networks (shown in **Figure 11**). SAM output to the LNs is excitatory cholinergic (ACh). The CAMs inputs to the LN may be inhibitory GABAergic (see text). (B) The two widespread networks of modulatory inputs into the fast transmission circuit (shown in A). The ascending AF processes (orange) make reciprocal connections with the SAM neurites, possibly inducing local release of neuromodulator (likely serotonin). The NMs (brown) spread throughout the neuropil. Physiological and immunohistochemical studies indicate that AFs and NMs release neuromodulators such as serotonin and octopamine and dopamine that likely play a role in reinforcing/suppressing the LTP at the SFL-to-SAM synapse.

Many SAMs together with a few CAMs appear to be organized in a canonical "association module"

Our reconstructions highlighted that ensembles of closely intermingled groups of SAM and CAM neurites form a *quasi*-columnar structural unit (**Figures 7A,7C,9A,9E, Figure - figure supplement 1**). As suggested above, this group of closely packed thin neurites likely functions as an "association module". The neurons in each module may respond to a combinatorial ensemble of sensory features from the SFL, as schematically represented in **Figure 11 inset**. Such combinatorial coding is a common feature in association networks such as the insect and mammalian olfactory systems (Malnic et al. 1999; Lledo et al. 2005; Galizia and Szyszka 2008; Turner et al. 2008; Gruntman and Turner 2013; Kurian et al. 2021). However, the one-to-one SFL-to-SAM connections suggest that each SFL neuron is represented in probably less than 12 ± 3.6 SD SAMs, most of them in different modules spread randomly over the VL cortex (**Figure 11**). This suggests that each of VL association module associates many (i.e., several hundreds of SMAs) random combinations of visual features that are represented in the 1.8 million SFL neurons. As the number

of association modules in the VL is very large (tens of thousands) it may enable the VL to "learn" the huge
amount of data associated with storing the visual environment. In principle the "association module" is
equivalent to that of a single interneuron in the input layer of a classical "fan-out fan-in" classification
network as proposed in Shomrat et al., (2011). However, the processing power provided by the association
module is computationally more elaborated than a single neuron can provide. It is tempting to speculate
that what is achieved in other systems by a single complex neuron is achieved in the octopus by the
construction of a microcircuitry composed of many neurons with simple properties (e.g., a linear
input/output relationship).

Figure 11. Schema of combinatorial coding in possible association modules array. *Inset:* The SAMs
and CAMs appear to be organized in columnar structures that integrate and feedforward "sharpened"
stimulus-specific inputs into the LNs. A specific stimulus (e.g., a visual feature) is fed forward via the
SAMs while the CAMs integrate these and the SAM inputs together and feed a general balancing inhibitory
input forward to the LNs. As the input to the SAMs is endowed with LTP, we suggest that this microcircuit
unit is a canonical "association module" of the VL. *The main scheme* shows how an array of association
modules that integrates a very sparse and randomly distribution of the SFL single input to the SAMs (1.0 3
± 0.73 LBs per 100μm). Each association module contains likely several hundred SAMs and relatively only
few CAMs (1.8%) suggesting that each module associates a variable combinatorial association of multiple
SFL features. Note that computationally an association module replaces a single AM in the input layer of
the originally assumed fan-out fan-in classification/association network (Shomrat et al 2011).
777

In summary, while earlier reports (Gray and Young 1964; Gray 1970; Young 1971) suggested that the VL
is organized in a simple fan-out fan-in feedforward architecture much like an associative "perceptron"
network (Shomrat et al. 2011), our study has revealed that its organization is more complex. Indeed, the
VL circuitry is characterized by two parallel and interconnected feedforward networks that are assembled
in what appear to be association modules, in which the SFL axons that convey processed sensory inputs
fan-out into many SAMs and fan-in into a low number of CAMs. Both SAMs and CAMs feed the excitatory
and inhibitory inputs, respectively, forward into the LN output layer.

785
786 **AKNOWLEDGMENTS**

787 This work was supported by Human Frontier Science Program (HFSP) grant no. RGP0042/2019-102, Israel
788 Science Foundation (ISF) grant no. 1928/15, National Institutes of Health (NIH) grant no. U19 NS104653,
789 U24 NS109102-01, NIH 1U01NS108637. We thank Prof. Tamar Flash for her contribution during the
790 inception phase of the project. We thank Prof. Jenny Kien for editing the manuscript.

791
Author contributions Sample preparation, F.B., Y.M., X.L.; sectioning, R.L.S.; imaging, F.B and R.L.S.;
alignment, A.P. and Y. W.; manual segmentation, F.B, F.Y., X.L. and A.S.; machine learning-aided
segmentation tool development, Y.M. and E.C.P.; Figures, F.B; Videos, Y.M and D.R.B.; statistical
analyses., F.B. and Y.M..; 3D renderings, F.B. and D.R.B.; electrophysiological data., T.S. and B.H.;
writing, F.B, Y.M and B.H.; review, D.R.B.; funding acquisition, J.W.L and B.H.; guidance, J.W.L

Declaration of interests

The authors declare no competing interests.

REFERENCES

Acsády L, Kamondi A, Sík A, et al (1998) GABAergic cells are the major postsynaptic targets of
mossy fibers in the rat hippocampus. J Neurosci 18:3386–3403

Agnati LF, Zoli M, Strömberg I, Fuxe K (1995) Intercellular communication in the brain: Wiring
versus volume transmission. Neuroscience 69:711–726. https://doi.org/10.1016/0306-
4522(95)00308-6

Aso Y, Hattori D, Yu Y, et al (2014) The neuronal architecture of the mushroom body provides a
logic for associative learning. eLife 3:e04577. https://doi.org/10.7554/eLife.04577

Bailey CH, Giustetto M, Huang Y-Y, et al (2000) Is Heterosynaptic modulation essential for
stabilizing hebbian plasiticity and memory. Nat Rev Neurosci 1:11–20.
https://doi.org/10.1038/35036191

Barth W, Mutzel P, Jünger M (2004) Simple and Efficient Bilayer Cross Counting. JGAA 8:179–
194. https://doi.org/10.7155/jgaa.00088

Bentley B, Branicky R, Barnes CL, et al (2016) The Multilayer Connectome of Caenorhabditis
elegans. PLOS Computational Biology 12:e1005283.
https://doi.org/10.1371/journal.pcbi.1005283

Berger DR, Seung HS, Lichtman JW (2018) VAST (Volume Annotation and Segmentation Tool):
Efficient Manual and Semi-Automatic Labeling of Large 3D Image Stacks. Front Neural
Circuits 12:. https://doi.org/10.3389/fncir.2018.00088

Boycott BB, Young JZ (1955) A Memory System in Octopus vulgaris Lamarck. Proceedings of
the Royal Society of London B: Biological Sciences 143:449–480.
https://doi.org/10.1098/rspb.1955.0024

D'Angelo E (2013) Cerebellar Granule Cell. In: Manto M, Schmahmann JD, Rossi F, et al. (eds)
Handbook of the Cerebellum and Cerebellar Disorders. Springer Netherlands,
Dordrecht, pp 765–791

Demmer H, Kloppenburg P (2009) Intrinsic membrane properties and inhibitory synaptic input of
kenyon cells as mechanisms for sparse coding? J Neurophysiol 102:1538–1550.
https://doi.org/10.1152/jn.00183.2009

Dowling JE, Boycott BB, Wells GP (1966) Organization of the primate retina: electron
microscopy. Proceedings of the Royal Society of London Series B Biological Sciences
166:80–111. https://doi.org/10.1098/rspb.1966.0086

Feinstein N, Nesher N, Hochner B (2011) Functional morphology of the neuromuscular system
of the Octopus vulgaris arm. Vie Milieu 11

Ferrante M, Migliore M, Ascoli GA (2009) Feed-forward inhibition as a buffer of the neuronal
input-output relation. PNAS 106:18004–18009. https://doi.org/10.1073/pnas.0904784106

Fiorito G, Chichery R (1995) Lesions of the vertical lobe impair visual discrimination learning by
observation in< i> Octopus vulgaris</i>. Neuroscience letters 192:117–120

Fischler MA, Bolles RC (1981) Random sample consensus: a paradigm for model fitting with
applications to image analysis and automated cartography. Commun ACM 24:381–395.
https://doi.org/10.1145/358669.358692

Fruchterman TMJ, Reingold EM (1991) Graph drawing by force-directed placement. Software:
Practice and Experience 21:1129–1164. https://doi.org/10.1002/spe.4380211102

Galizia CG, Szyszka P (2008) Olfactory coding in the insect brain: molecular receptive ranges,
spatial and temporal coding. Entomologia Experimentalis et Applicata 128:81–92.
https://doi.org/10.1111/j.1570-7458.2007.00661.x

Garthwaite J (2016) From synaptically localized to volume transmission by nitric oxide. The
Journal of Physiology 594:9–18. https://doi.org/10.1113/JP270297

Gray EG (1970) The Fine Structure of the Vertical Lobe of Octopus Brain. Philosophical
Transactions of the Royal Society B: Biological Sciences 258:379–394.
https://doi.org/10.1098/rstb.1970.0040

Gray EG, Young JZ (1964) Electron microscopy of synaptic structure of Octopus brain. J Cell
Biol 21:87–103

Gruntman E, Turner GC (2013) Integration of the olfactory code across dendritic claws of single
mushroom body neurons. Nat Neurosci 16:1821–1829. https://doi.org/10.1038/nn.3547

Hanlon RT, Messenger JB (2018) Cephalopod Behaviour. Cambridge University Press

Hayworth KJ, Morgan JL, Schalek R, et al (2014) Imaging ATUM ultrathin section libraries with
WaferMapper: a multi-scale approach to EM reconstruction of neural circuits. Frontiers in
Neural Circuits 8:. https://doi.org/10.3389/fncir.2014.00068

Hochner B (2010) Functional and comparative assessments of the octopus learning and
memory system. Front Biosci (Schol Ed) 2:764–771

Hochner B (2012) An Embodied View of Octopus Neurobiology. Current Biology 22:R887–
R892. https://doi.org/10.1016/j.cub.2012.09.001

Hochner B (2013) How nervous systems evolve in relation to their embodiment: what we can
learn from octopuses and other molluscs. Brain Behav Evol 82:19–30.
https://doi.org/10.1159/000353419

Hochner B, Brown ER, Langella M, et al (2003) A Learning and Memory Area in the Octopus
Brain Manifests a Vertebrate-Like Long-Term Potentiation. Journal of Neurophysiology
90:3547–3554. https://doi.org/10.1152/jn.00645.2003

Hochner B, Glanzman DL (2016) Evolution of highly diverse forms of behavior in molluscs. Curr
Biol 26:R965–R971. https://doi.org/10.1016/j.cub.2016.08.047

Hochner B, Shomrat T, Fiorito G (2006) The octopus: a model for a comparative analysis of the
evolution of learning and memory mechanisms. Biol Bull 210:308–317.
https://doi.org/10.2307/4134567

Hochner B, Spira ME (1987) Preservation of motoneuron electrotonic characteristics during
postembryonic growth. J Neurosci 7:261–270

Holm S (1979) A simple sequentially rejective multiple test procedure. Scandinavian Journal of
Statistics 6:65–70

Jortner RA, Farivar SS, Laurent G (2007) A Simple Connectivity Scheme for Sparse Coding in
an Olfactory System. J Neurosci 27:1659–1669.
https://doi.org/10.1523/JNEUROSCI.4171-06.2007

Kandel ER, Dudai Y, Mayford MR (2014) The Molecular and Systems Biology of Memory. Cell
157:163–186. https://doi.org/10.1016/j.cell.2014.03.001

Kasthuri N, Hayworth KJ, Berger DR, et al (2015) Saturated Reconstruction of a Volume of
Neocortex. Cell 162:648–661. https://doi.org/10.1016/j.cell.2015.06.054

Kawato M, Ohmae S, Hoang H, Sanger T (2021) 50 Years Since the Marr, Ito, and Albus
Models of the Cerebellum. Neuroscience 462:151–174.
https://doi.org/10.1016/j.neuroscience.2020.06.019

Kier WM (1996) Muscle development in squid: Ultrastructural differentiation of a specialized
muscle fiber type. J Morphol 229:271–288. https://doi.org/10.1002/(SICI)1097-
4687(199609)229:3<271::AID-JMOR3>3.0.CO;2-1

Koren Y (2005) Drawing graphs by eigenvectors: theory and practice. Computers &
Mathematics with Applications 49:1867–1888.
https://doi.org/10.1016/j.camwa.2004.08.015

Kurian SM, Naressi RG, Manoel D, et al (2021) Odor coding in the mammalian olfactory
epithelium. Cell Tissue Res 383:445–456. https://doi.org/10.1007/s00441-020-03327-1

Lee K, Zung J, Li P, et al (2017) Superhuman Accuracy on the SNEMI3D Connectomics
Challenge

Lee W-CA, Bonin V, Reed M, et al (2016) Anatomy and function of an excitatory network in the
visual cortex. Nature 532:370–374. https://doi.org/10.1038/nature17192

Leiss F, Groh C, Butcher NJ, et al (2009) Synaptic organization in the adult Drosophila
mushroom body calyx. Journal of Comparative Neurology 517:808–824.
https://doi.org/10.1002/cne.22184

Leng G, Ludwig M (2008) Neurotransmitters and peptides: whispered secrets and public
announcements. J Physiol 586:5625–5632. https://doi.org/10.1113/jphysiol.2008.159103

Li F, Lindsey JW, Marin EC, et al (2020) The connectome of the adult Drosophila mushroom
body provides insights into function. eLife 9:e62576. https://doi.org/10.7554/eLife.62576

Litvak V, Sompolinsky H, Segev I, Abeles M (2003) On the Transmission of Rate Code in Long
Feedforward Networks with Excitatory–Inhibitory Balance. J Neurosci 23:3006–3015.
https://doi.org/10.1523/JNEUROSCI.23-07-03006.2003

Lledo P-M, Gheusi G, Vincent J-D (2005) Information Processing in the Mammalian Olfactory
System. Physiological Reviews 85:281–317. https://doi.org/10.1152/physrev.00008.2004

Lowe DG (1999) Object recognition from local scale-invariant features. In: Proceedings of the
Seventh IEEE International Conference on Computer Vision. IEEE, Kerkyra, Greece, pp
1150–1157 vol.2

Macrina T, Lee K, Lu R, et al (2021) Petascale neural circuit reconstruction: automated
methods. bioRxiv 2021.08.04.455162. https://doi.org/10.1101/2021.08.04.455162

Malnic B, Hirono J, Sato T, Buck LB (1999) Combinatorial receptor codes for odors. Cell
96:713–723. https://doi.org/10.1016/s0092-8674(00)80581-4

Mather JA (1995) Cognition in Cephalopods. In: Advances in the Study of Behavior. Elsevier, pp
317–353

McDonald AJ, Muller JF, Mascagni F (2002) GABAergic innervation of alpha type II
calcium/calmodulin-dependent protein kinase immunoreactive pyramidal neurons in the
rat basolateral amygdala. J Comp Neurol 446:199–218.
https://doi.org/10.1002/cne.10204

Meinertzhagen IA (2019) Morphology of Invertebrate Neurons and Synapses. The Oxford
Handbook of Invertebrate Neurobiology.
https://doi.org/10.1093/oxfordhb/9780190456757.013.9

Meirovitch Y, Kang K, Draft RW, et al (2021) Neuromuscular connectomes across development
reveal synaptic ordering rules. bioRxiv 2021.09.20.460480.
https://doi.org/10.1101/2021.09.20.460480

Meirovitch Y, Mi L, Saribekyan H, et al (2019) Cross-Classification Clustering: An Efficient Multi-
Object Tracking Technique for 3-D Instance Segmentation in Connectomics. pp 8425–
8435

Messenger JB (1996) Neurotransmitters of cephalopods. Neurotransmitters of cephalopods
2:95–114

MICrONS Consortium, Bae JA, Baptiste M, et al (2021) Functional connectomics spanning
multiple areas of mouse visual cortex. bioRxiv 2021.07.28.454025

Morgan JL, Berger DR, Wetzel AW, Lichtman JW (2016) The Fuzzy Logic of Network
Connectivity in Mouse Visual Thalamus. Cell 165:192–206.
https://doi.org/10.1016/j.cell.2016.02.033

Morgan JL, Lichtman JW (2020) An Individual Interneuron Participates in Many Kinds of
Inhibition and Innervates Much of the Mouse Visual Thalamus. Neuron 106:468-481.e2.
https://doi.org/10.1016/j.neuron.2020.02.001

Motta A, Berning M, Boergens KM, et al (2019) Dense connectomic reconstruction in layer 4 of
the somatosensory cortex. Science 366:. https://doi.org/10.1126/science.aay3134

Packard A (1972) Cephalopods and Fish: The Limits of Convergence. Biological Reviews
47:241–307. https://doi.org/10.1111/j.1469-185X.1972.tb00975.x

Parsons TD, Sterling P (2003) Synaptic Ribbon: Conveyor Belt or Safety Belt? Neuron 37:379–
382. https://doi.org/10.1016/S0896-6273(03)00062-X

Pavarino E, Berger D, Kang K, et al (2019) mEMbrain: an interactive deep learning tool for
labeling and instance segmentation of EM datasets. [Poster]. Bernstein Conference,
September 17-18, Berlin, doi: 10.12751/nncn.bc2019.0083

Ramírez OA, Couve A (2011) The endoplasmic reticulum and protein trafficking in dendrites and
axons. Trends in Cell Biology 21:219–227. https://doi.org/10.1016/j.tcb.2010.12.003

Ronneberger O, Fischer P, Brox T (2015) U-Net: Convolutional Networks for Biomedical Image
Segmentation. In: Navab N, Hornegger J, Wells WM, Frangi AF (eds) Medical Image
Computing and Computer-Assisted Intervention – MICCAI 2015. Springer International
Publishing, Cham, pp 234–241

Ryan K, Lu Z, Meinertzhagen IA (2016) The CNS connectome of a tadpole larva of Ciona
intestinalis (L.) highlights sidedness in the brain of a chordate sibling. eLife 5:e16962.
https://doi.org/10.7554/eLife.16962

Saalfeld S, Fetter R, Cardona A, Tomancak P (2012) Elastic volume reconstruction from series
of ultra-thin microscopy sections. Nat Methods 9:717–720.
https://doi.org/10.1038/nmeth.2072

Sanders GD (1975) The Cephalopods. In: Corning W.C., Dyal J.A. & Willows A.O.D. (eds)
Invertebrate learning, Vol. 3. Plenium Press, New-York, pp 1-101.

Scheffer LK, Xu CS, Januszewski M, et al (2020) A connectome and analysis of the adult
Drosophila central brain. eLife 9:e57443. https://doi.org/10.7554/eLife.57443

Schmidt M, Derby CD (2011) Cytoarchitecture and ultrastructure of neural stem cell niches and
neurogenic complexes maintaining adult neurogenesis in the olfactory midbrain of spiny
lobsters, Panulirus argus. J Comp Neurol 519:2283–2319.
https://doi.org/10.1002/cne.22657

Schneider-Mizell CM, Gerhard S, Longair M, et al (2016) Quantitative neuroanatomy for
connectomics in Drosophila. eLife 5:e12059. https://doi.org/10.7554/eLife.12059

Schnell AK, Hanlon RT, Benkada A, Jozet-Alves C (2016) Lateralization of Eye Use in
Cuttlefish: Opposite Direction for Anti-Predatory and Predatory Behaviors. Front Physiol
7:. https://doi.org/10.3389/fphys.2016.00620

Shapson-Coe A, Januszewski M, Berger DR, et al (2021) A connectomic study of a petascale
fragment of human cerebral cortex. bioRxiv 2021.05.29.446289

Shigeno S, Andrews PLR, Ponte G, Fiorito G (2018) Cephalopod Brains: An Overview of
Current Knowledge to Facilitate Comparison With Vertebrates. Front Physiol 9:.
https://doi.org/10.3389/fphys.2018.00952

Shigeno S, Ragsdale CW (2015) The gyri of the octopus vertical lobe have distinct neurochemical identities. J Comp Neurol 523:1297–1317. https://doi.org/10.1002/cne.23755

Shomrat T, Feinstein N, Klein M, Hochner B (2010) Serotonin is a facilitatory neuromodulator of synaptic transmission and "reinforces" long-term potentiation induction in the vertical lobe of Octopus vulgaris. Neuroscience 169:52–64. https://doi.org/10.1016/j.neuroscience.2010.04.050

Shomrat T, Graindorge N, Bellanger C, et al (2011) Alternative Sites of Synaptic Plasticity in Two Homologous "Fan-out Fan-in" Learning and Memory Networks. Current Biology 21:1773–1782. https://doi.org/10.1016/j.cub.2011.09.011

Shomrat T, Turchetti-Maia AL, Stern-Mentch N, et al (2015) The vertical lobe of cephalopods: an attractive brain structure for understanding the evolution of advanced learning and memory systems. J Comp Physiol A Neuroethol Sens Neural Behav Physiol 201:947–956. https://doi.org/10.1007/s00359-015-1023-6

Shomrat T, Zarrella I, Fiorito G, Hochner B (2008) The Octopus Vertical Lobe Modulates Short-Term Learning Rate and Uses LTP to Acquire Long-Term Memory. Current Biology 18:337–342. https://doi.org/10.1016/j.cub.2008.01.056

Stern-Mentch N, Winters-Bostwick G, Belenky M, et al (2022) Neurotransmission and neuromodulation systems in the learning and memory network of Octopus vulgaris. Journal of Morphology n/a: https://doi.org/10.1002/jmor.21459

Turchetti-Maia A, Shomrat T, Hochner B (2017) The Vertical Lobe of Cephalopods—A Brain Structure Ideal for Exploring the Mechanisms of Complex Forms of Learning and Memory. The Oxford Handbook of Invertebrate Neurobiology. https://doi.org/10.1093/oxfordhb/9780190456757.013.29

Turchetti-Maia AL, Stern-Mentch N, Bidel F, et al (2018) A novel NO-dependent 'molecular-memory-switch' mediates presynaptic expression and postsynaptic maintenance of LTP in the octopus brain. bioRxiv 491340. https://doi.org/10.1101/491340

Turner GC, Bazhenov M, Laurent G (2008) Olfactory Representations by Drosophila Mushroom Body Neurons. Journal of Neurophysiology 99:734–746. https://doi.org/10.1152/jn.01283.2007

Turrigiano GG, Nelson SB (2004) Homeostatic plasticity in the developing nervous system. Nature Reviews Neuroscience 5:97–107. https://doi.org/10.1038/nrn1327

van den Pol AN (2012) Neuropeptide transmission in brain circuits. Neuron 76:98–115. https://doi.org/10.1016/j.neuron.2012.09.014

Verasztó C, Jasek S, Gühmann M, et al (2020) Whole-animal connectome and cell-type complement of the three-segmented Platynereis dumerilii larva. bioRxiv 2020.08.21.260984. https://doi.org/10.1101/2020.08.21.260984

Vitti JJ (2013) Cephalopod Cognition in an Evolutionary Context: Implications for Ethology. Biosemiotics 6:393–401. https://doi.org/10.1007/s12304-013-9175-7

Wadell H (1935) Volume, Shape, and Roundness of Quartz Particles. The Journal of Geology
43:250–280

Weigel AV, Chang C-L, Shtengel G, et al (2021) ER-to-Golgi protein delivery through an
interwoven, tubular network extending from ER. Cell 0:
https://doi.org/10.1016/j.cell.2021.03.035

West MJ, Gundersen HJG (1990) Unbiased stereological estimation of the number of neurons
in the human hippocampus. Journal of Comparative Neurology 296:1–22.
https://doi.org/10.1002/cne.902960102

White JG, Southgate E, Thomson JN, Brenner S (1986) The Structure of the Nervous System of
the Nematode Caenorhabditis elegans. Philosophical Transactions of the Royal Society
B: Biological Sciences 314:1–340. https://doi.org/10.1098/rstb.1986.0056

Winters GC, Polese G, Di Cosmo A, Moroz LL (2020) Mapping of neuropeptide Y expression in
Octopus brains. J Morphol 281:790–801. https://doi.org/10.1002/jmor.21141

Witvliet D, Mulcahy B, Mitchell JK, et al (2021) Connectomes across development reveal
principles of brain maturation. Nature 596:257–261. https://doi.org/10.1038/s41586-021-
03778-8

Young JZ (1979) The nervous system of Loligo. V. The vertical lobe complex. Philosophical
Transactions of the Royal Society of London 285:311–354

Young JZ (1995) Multiple matrices in the memory system of Octopus. In: Cephalopod
Neurobiology. Oxford University Press, Oxford

Young JZ (1963) The Number and Sizes of Nerve Cells in Octopus. Proceedings of the
Zoological Society of London 140:229–254. https://doi.org/10.1111/j.1469-
7998.1963.tb01862.x

Young JZ (1971) The anatomy of the nervous system of Octopus vulgaris. Clarendon Press

Zarrella I, Ponte G, Baldascino E, Fiorito G (2015) Learning and memory in Octopus vulgaris: a
case of biological plasticity. Current Opinion in Neurobiology 35:74–79.
https://doi.org/10.1016/j.conb.2015.06.012

Zheng Z, Lauritzen JS, Perlman E, et al (2018) A Complete Electron Microscopy Volume of the
Brain of Adult Drosophila melanogaster. Cell 0: https://doi.org/10.1016/j.cell.2018.06.019

Zhu PC, Thureson-Klein Å, Klein RL (1986) Exocytosis from large dense cored vesicles outside
the active synaptic zones of terminals within the trigeminal subnucleus caudalis: A
possible mechanism for neuropeptide release. Neuroscience 19:43–54.
https://doi.org/10.1016/0306-4522(86)90004-7

1053

Methods

Tissue preparation

A mature wild *Octopus vulgaris* was collected by local fisherman on the Israeli Mediterranean coast and was held in artificial seawater in an individual aquarium for a few days. As previously published (Shomrat et al. 2011), the animal was deeply anesthetized in seawater containing 55 mM $MgCl_2$ and 2% ethanol. The supraoesophageal mass was removed through a dorsal opening of the cranium (Young 1971) and then quickly sliced into 1 mm sagittal sections using a mouse matrix brain slicer and double- edged razor blade. Three slices containing the vertical lobes were obtained and immersed in fixative solution (20:1 ratio).

Fixation

Because the preservation of the ultrastructure is a central requirement of a connectomics study, 3 fixations protocol were tested. The 3 solutions were chosen for their various aldehyde concentrations and tonicity as well as their previous use with octopus or other marine invertebrate tissue preparations for electron microscopy. Each slice was immersed in one of the following:

Fixative #1 (Kier 1996): 3 % glutaraldehyde, 0.065 M phosphate buffer(PB), 0.5% tannic acid and 6% sucrose for 12 hr at 4°C.

Fixative #2 (Feinstein et al. 2011): a mixture of 3 % glutaraldehyde and 4 % paraformaldehyde formaldehyde in artificial seawater for 4 hr at room temperature (RT).

Fixative #3 (Schmidt and Derby 2011): 5% glutaraldehyde in 0.1 M PB containing 15% sucrose for 4 hr at RT.

After fixation, the slices were rinsed in 0.1M PB 4 × 30 minutes, then transferred to chilled 0.065 M phosphate buffer with 1% glutaraldehyde and 6% sucrose and stored at 4°.

We chose fixative #1 for the volumetric reconstruction, constituting the 'large connectomics dataset' (4nm/px, 200 ns/px), due to good preservation of the macroelements (*i.e.,* cell body) and ultrastructure (*e.g.,* mitochondria, postsynaptic density, various types of synaptic vesicles, etc.) as well as the high contrast between elements. Fixative #3 showed poor preservation of the macroelements but greater expansion (more extracellular space) and overall better fine ultrastuctural preservation. Accordingly, this fixative was used to generate a small 'high-magnification dataset' (1nm/px, 2 μs/px) to examine possible synaptic specialization and the synaptic vesicles at the sites of LTP generation in the VL (Hochner et al., 2003, Shomrat et al., 2011).

EM staining

The samples were stained with a modified *en bloc* staining protocol that was developed by Hua et al. (2015)

for volumetric EM imaging. Before staining, the chemically fixed tissues were rinsed 3 times for 20 min

each in 150mM sodium cacodylate buffer (pH 7.4). The tissues were then stained in 2% w/v

OsO4 buffered with 150mM cacodylate for 2 hours. After being rinsed 3 x 20 min in cacodylate buffer, the

tissues were immersed in 1.5% w/v potassium ferrocyanide buffered with cacodylate for 4 hours. The

tissues were rinsed in deionized water for 3 x 20 min and then incubated in filtered 0.5% w/v

thiocarbohydrazide aqueous solution for 1.5 hours. The tissues were then rinsed in deionized water for 3 x

20 min and stained again with 2% w/v OsO4 aqueous solution for 3 hours. After the second osmication, the

tissues were rinsed in water for 3 x 20 min and transferred to 1% w/v uranyl acetate aqueous solution which

was wrapped in aluminum foil to protect from light for overnight staining (approx.12 hours, RT). On the

next day, the samples had a final rinse in water and were then dehydrated through 25%, 50%, 75%, 100%,

100% acetonitrile solution for 20 min each into pure acetonitrile. The dehydrated samples were infiltrated

with 25% EPON:acetonitrile for 6 hours, 50% EPON:acetonitrile for 12 hours, 75% EPON:acetonitrile for

12 hours, 100% EPON for 12 hours, 100% EPON for another 12 hours. The fully infiltrated samples were

placed in a flat embedding mold and incubated in a 60°C oven for two days.

Collection of serial ultrathin sections using an automated tape-collecting ultramicrotome (ATUM)

The block face of the sample cylinder was trimmed into a 1×1 mm square containing the vertical lobe. 30

nm thin sections for the 'connectomics dataset' and 40nm sections for the 'high magnification dataset' were

cut using a Leica ultramicrotome and collected on a home-built automated tape collecting system using a

plasma-treated polyamide 8 mm wide tape (Kapton, Sheldahl) that was carbon-coated (Hayworth et al.

2014). On collection of the sections, the tape was cut into strips, and these were mounted on 100 mm silicon

wafers. The connectomics dataset was composed of 7 wafers, between 115-140 sections per wafer, (total

892 sections), while the high magnification dataset was composed of a single wafer of 83 sections.

EM imaging

For the connectomics dataset the sections were first mapped at low resolution (**Figure 2A)** and the EM

image stack was aligned by Fiji's plugin "Linear Stack Alignment with SIFT" to form a coherent 3D EM

volume. The VL region free from large cracks in the tissue was chosen for high resolution imaging. The

sections on the wafer were cleaned while in the Magellan microscope using the FEI-supplied plasma

system. A FEI Magellan Scanning Electron Microscope (Thermofisher Scientific) equipped with custom

image acquisition software (WaferMapper, Hayworth, Morgan et al. 2014) was used for high resolution

imaging. A backscattered electron detector in long working distance mode with a ~3.2 NA current at 7 kev

incident electron energy and a dwell time of 200 ns/px was used for imaging at 4 nm/px. For each section

we obtained a 4 by 6 montage of image tiles (each tile was a 16000 x 16000-pixel image). At the end of

each imaging run, out of focus images were manually identified and retaken. Artefacts such as dirt, lead

precipitates, wrinkles or even large folds were seen on many sections and severely damaged sections were

removed from the total stack. This resulted in 51 sections missing from the 892 sections. However, the

occurrence of serious damage in adjacent sections was rare, allowing confident tracing of the elements.

Image stitching and alignment

1121

The stitching and alignment of the dataset were performed using a method inspired by Saalfeld et al (2012).

Each step was parallelized over image tiles and sections. The single-beam electron microscope produced

high resolution image tiles sized 16,000x16,000 pixels whose precise location relative to one another was

not known at the nanometer level. Hence, stitching these tiles together required using image information.

We stitched these image tiles for each of the 892 sections. The process consisted of a coarse-grained

stitching followed by fine-grained stitching. The coarse-grained stitching performed a per-tile affine

transformation, setting the tiles roughly in their final position. SIFT features (Lowe 1999) were computed

for pixels up to 2,400 from the tile boundary. The features of the overlapping tiles were compared and

matched to find point correspondences. To remove outliers of incorrect correspondences, the RANSAC

algorithm (Fischler and Bolles 1981) was used, optimizing for rigid transformations. Finally, the tiles'

affine transformation was optimized by minimizing the sum of squared roots of all correspondences.

To achieve better alignment and compensate for minor non-linear deformations in imaging, a fine-grained

stitching was performed after the coarse-grained stitching. A triangular mesh was laid on each tile and a

1x1 µm2 area around each mesh vertex overlapping with an adjacent tile was cropped. The cropped patch

was matched against a transformed cropped area of the neighboring tile of size 2x2 µm2. The 2D elastic

optimization algorithm described in Saalfeld et al. (2012) was used to minimize the distances of the

matches.

We applied a custom-made 3D alignment algorithm to obtain an image stack of all 892 sections. Once all

the sections were stitched as described above, correspondences between adjacent sections were detected.

The first step consisted of coarse features detection and matching, and subsequently patch matching in small

regions. The matching was performed between all section pairs separated by at most one section (hence

considering matches between non-adjacent sections). Coarse features were obtained from the stitched

sections. The features were detected using the OpenCV SimpleBlobDetector ("OpenCV:

cv:SimpleBlobDetector Class Reference" n.d.). The detector searched for circular blobs that typically

corresponded to blood cells, mitochondria and axonal cross-sections. Features were defined using a SIFT

descriptor (Lowe 1999). After matching the features of each pair of sections, RANSAC filtered out wrong

matches, from which an affine transformation between the sections was computed. In the second step, fine-

grained patch matching at half resolution was performed. A triangular mesh was laid on the sections and a

1.6x1.6 μm^2 area around each mesh vertex was cropped, followed by cross-correlation search against a

transformed cropped area of the neighboring section of size 4x4 µm2. The valid cross-correlation matches

were then used as an input to a 3D elastic optimization algorithm based on Saalfeld et al. (2012) which

minimizes the cross-layer distance of the matches while preserving the 2D structure of the sections.

Semi-automatic segmentation: mEMbrain

To segment the aligned image stack (892 sections, volume 2.7 million μm^3; **Figure 1B**), we applied an

automated reconstruction pipeline to the volume using a machine learning reconstruction pipeline (Lee et

al. 2017; Meirovitch et al. 2019). The inputs to this pipeline required fully annotating the image stacks of 4

manually picked volumes (image stacks ranged in volume from 21.6 to 153 μm^3; **Figure 1-supplement 3**),

which served as ground truth for training the artificial neural network. The annotation of this ground truth

labeled two different categories: intracellular space, excluding cellular membranes (category 1), or any

combination of cellular membrane and ECS (category 2). We leveraged the mEMbrain software package

implemented in MATLAB (Pavarino et al. 2019) to obtain a first template of the cellular processes. This

computation included training a deep neural network (Unet; Ronneberger et al., 2015) to classify pixels

according to the two categories (accuracy above 93%) and produce a 2-dimensional instance segmentation

of the individual cross-sections of all cellular compartments using Watersheds (Pavarino et al. 2019). We

used custom code to merge colored neuronal cross-sections across slices of the image stack if adjacent

cross-sections matched in shape or if adjacent processes had sufficient overlapping cytoplasm far from any

cellular membrane. Applying this conservative merging resulted in a 3-dimensional representation of object

instances that each rarely overlapped two distinct neuronal or glial processes, and often the splitting of

complete neuronal processes into several segmented objects. In most cases, however, the 2-D cross sections

were lawfully (fully) segmented into one object and after conservative agglomeration procedure based on

shape matching these objects were correctly agglomerated over a few consecutive sections. We also applied

3D agglomeration procedures including mean affinity   agglomeration suggested by Lee et al. (2017) and

Meirovitch et al. (2019). However, due to membrane breaks these procedures often resulted in merge errors

that were hard to manually proofread and hence this agglomeration was not included in the circuit analysis.

We found the cause of the merge errors to be associated with specific cellular and compartment types,

including the shafts of the large neurons and many synaptic boutons. Because such preparation artifacts

severely affected the quality of our fully automated reconstruction, we combined the automated

segmentation with manual methods described below.

Membrane-constrained ML-aided reconstruction.

To facilitate manual reconstruction we used the mEMbrain package (Pavarino et al. 2019) implemented in

MATLAB (Mathworks) together with VAST (Volume Annotation and Segmentation Tool

(https://software.rc.fas.harvard.edu/lichtman/vast/, Berger et al., 2018). Using U-net architectures of deep

neural networks (Ronneberger et al. 2015) mEMbrain computed the boundaries of all cellular processes,

scoring the likelihood (0 to 1) that pixels belonged to intracellular space (category 1) or not (category 2).

mEMbrain arranged these scores into a volume of membrane probabilities consistent with the EM image

stack layer in VAST, which allowed tracing cellular processes aided by the pre-computed membrane

probabilities. We constrained the tracing to the membrane probabilities layer generated by mEMbrain using

the "constrained painting" function in VAST, which restricted the tracing to connected components of

pixels far from cellular membranes. This ML-aided tracing allowed the annotator to ignore the location of

the cellular membranes and work in a skeletonization mode if the initial painting stroke overlapped

intracellular space. Tests on small volumes suggested a ~5-10x faster annotation in this mode than manual

painting without constraints, while retraining the accuracy of careful pixel annotation. This membrane-

constrained ML-aided reconstruction was used for the 516 cells described and studied morphologically here

(**Figure 1C**).

Node-based agglomeration of 3D compartments

We also converted the automated neuron reconstruction IDs into a VAST image stack layer holding the

identities of $~16 \times 10^6$ objects to allow even faster semi-automated reconstruction of processes. In this

procedure the annotator used the automatic reconstruction by iteratively agglomerating 3D objects that were

found to belong to a single cellular process. This was done using VAST's "Tool Layer" functionality which

we mapped to the automated segmentation layer and to an additional annotation layer that represented the

annotator's manually made skeletons. The annotator placed single nodes within the objects segmented by

the automated reconstruction pipeline; together the collection of nodes of a single skeleton defined the

agglomeration of all objects straddled by the nodes into a single proofread cell. This membrane-constrained

ML-aided reconstruction was used to quickly reconstruct all the synaptic partners of one LN (**Figure 7C**)

and one AF (**Figure 9 - figure supplement 1**).

Criteria for cell classification

Our morphological reconstruction included 516 cells segments and a wiring length of more than 12.8 cm.

Only neurons that were sufficiently contained within the reconstructed volume were classified. Processes

that could not be followed across the volume or were not wholly contained within it were considered orphan

fragments and fell into the 'unmarked category' with other uncharacterized profiles.

1. Afferent axons

Classified as afferent axons were processes of constant radius (as opposed to dendrites that often taper),
with exclusively presynaptic boutons filled with synaptic vesicles smaller than 60 nm and with no
connection to a cell body. Two processes met these features: the SFL axons (see Results; **Figure 3** and the
CIN (see Results; **Figure 8**). The SFL axons were classified according to several established criteria (Gray
and Young 1964; Gray 1970; Young 1971): the axon projected into the VL lobule within the outer zone of
the neuropil, made *en passant* synapses with AM neurons and at the ultrastructural level their varicosities
showed an electron-lucent cytoplasm with average synaptic vesicle size of 59.7 $\pm$ 1.5nm, and numerous
round mitochondria. Unlike the SFL, CIN were found in the SFL and in the neuropil, displayed the smallest
vesicles in our dataset (38.2 $\pm$ 0.9 nm) and synapsed exclusively with clear postsynaptic profiles.
Additionally, CIN fibers showed a high concentration of neurofilaments which ended where the synaptic
vesicles appeared (Gray 1970). The high neurofilament content of these processes led Gray (1970) to
suggest that these were pain fibers. Both afferents could be easily identified with only electron micrographs.

1225   2. VL Interneurons

Neurons were classified as VL interneuron if they received input from the SFL and/or were previously
described as intrinsic to the lobes. All partners of the SFL axons were the AM cells (Gray and Young
1964; Gray 1970; Young 1971). However, we discovered two types of AMs: SAM and CAM (see Results;
Figure 6). The SAMs showed a single inwardly directed trunk that cannot be defined either as axon or
dendrite. Additionally, their ultrastructural signature was easily identifiable through the presence of
synaptic vesicles (49.5 $\pm$ 1.3 nm) along the length of this trunk and the presence of specific ultrastructural
components such as cisternae and the transport endoplasmic reticulum at every presynaptic site. Unlike the
SAM, the CAM displayed a strict segregation between a dendritic compartment located within the SFL
tract and their axonal process within the neuropil. Their dendritic tree was lightly or heavily branched, with
one or multiple axons arising from it. Their dendritic arborizations displayed a classical electron-lucent
postsynaptic profile filled with polyribosome. Unlike the SAMs, the CAM presynaptic sites in the neuropil
displayed a clearer cytoplasm with no specific ultrastructure and smaller vesicles than the SAMs (42.6 $\pm$
1.1 nm).

We also observed putative interneurons displaying a bifunctional neurite, with both pre- and postsynaptic
sites identified (**Figure 1-figure supplement 3A**). Previously undescribed and poorly represented in our
dataset and apparently poorly involved in the major cell network, all such cells were considered as
unmarked cells in the data analysis.

1243   3. Efferent Fibers

Only the LNs fell into this category. Their cell bodies (diameter > 15 µm; n=14) lay mostly in the inner

zones of the cortex and their large trunks ran radially into the neuropil (see Results; **Figure 7**). While most

of the LNs were poorly contained in the reconstructed volume, some gave off dendritic collaterals within

the SFL tract area where they were solely postsynaptic to the SAMs and unmarked process. Their dendrites

showed a variable diameter along their length and displayed a typical clear postsynaptic profile. In the

neuropil, we classified all clear postsynaptic profiles with convoluted arborizations heavily innervated by

the AMs as LN processes.

4.   Neuromodulatory neurons

Based on known invertebrate modulatory synapses, processes were classified as modulatory if they

contained mostly large vesicles (>60 nm), either dark or clear. We distinguished two types of

neuromodulatory processes (see Results; **Figure 9**). First, the ascending fibers (AF) with a distinctive

morphology - a menorah –shaped tree climbing from the neuropil to the cortex while intermingling with

SAM trunks. Along their length the AF displayed boutons filled either with dense core vesicles (AF-Nm)

and an average size of 74.9 ± 1,6 mm or a mix of smaller vesicle types that could be round, flattened or

irregularly shaped (AF-mix). Various subtypes of AF-mix were observed. The AF-Nm neurons were

heavily innervated by the SAMs without any sign of presynaptic activity. In contrast, the AF-mix were

innervated to lesser extend by the SAMs and displayed some presynaptic sites as well as occasionally

reciprocal contacts.

The second type were neuromodulatory fibers (NMs) which were dispersed more globally in the VL

neuropil and within the SFL tract and lacked a general orientation. Several subtypes were characterized by

the homogeneity of the vesicle types found along their length. Most of their boutons lacked any sign of

synaptic connection but the few pre- or postsynaptic sites observed were unorthodox. These NM processes

displayed the largest vesicles in the entire dataset with diameters up to 200 nm (mean 96.3 ± 3.5 nm) within

their bouton and an accumulation of much smaller vesicles was observed at their rather rare presynaptic

sites.

5.   Glia

Cells were classified as glia if they showed clearly distinguishable gliofibrils (Gray, 1964) and/or did not

receive any synapses. At least two types of protoplasmic glia were identified (**Figure 1-figure supplement**

3C). (1) radial glia (Young, 1971) were full of gliofibrils and characterized by long radial processes and

arborizations infiltrating the neuronal cell cortex. These cells appear to isolate neuronal types from each

other. (2) The astrocyte-like glia had many gliofibrils within their processes, which showed an increased

tortuosity densely infiltrating the SFL tract. Here they appeared to wrap distinct synaptic areas. Through

repeating folds this cell type appears to maximize self-contacts characterized by apposition of their
membranes and accumulation of electron dense materials on both sides, possibly sites for protein exchange.
One astrocyte-like cell with its cell body located near a cell island was partially reconstructed; it covered
an extensive area within the SFL tract. (3) Fibrous glia with cell bodies in the inner cortex were also
observed; these did not receive any synapses in the reconstructed volume.

6. Progenitor-like cells

Cells were classified as progenitor-like only if their cell body was reconstructed (**Figure 1-figure**
supplement 3B). These cells were nested in each other in the inner cortex forming a type of horizontal
layer. There were at least two types of cells in this arrangement. First, the base was formed by elongated
cell bodies whose nuclei showed bumpy irregular membranes often with nuclear folds. These cell bodies
appeared to have very small projections, and, like the glia, they lacked synaptic connections. Above these
elongated cells were more classically spherical cell bodies which appeared somewhat "immature." Their
trunks, extending radially into the neuropil, often lacked synaptic connection and terminated early with
filopodia typical of axon growth cones. Their fibers were rich in grayish filaments that could be filamentous
actin, suggesting that this may be a site of plastic reorganization. Because of the similarity of this
arrangement with neurogenic niches observed in the mammalian dentate gyrus of the hippocampus and
the subventricular zone, we termed these cells "progenitor-like cells." We saw no sign of cell division.

Identification and analysis of synapses

Chemical synapses were identified by features common to invertebrate synapses - the presence of a cluster
of synaptic vesicles (sv) or 'cloud' associated with a lightly active zone near the presynaptic membrane
(Gray 1970; White et al. 1986; Ryan et al. 2016; Zheng et al. 2018; Meinertzhagen 2019; Witvliet et al.
2021). Synapses were annotated if the accumulation of sv persisted through at least 3 sections in a row
(90nm). We distinguished between traditional chemical synapses (fast transmission) with 30-60nm sv and
synapses of modulatory neurons with mostly large vesicles (Witvliet et al. 2021). Consistent with previous
studies (Gray and Young 1964; Gray 1970), we did not find evidence of electrical connections (gap
junction) or any clear postsynaptic density (PSD) at any chemical synapse within the vertical lobe.
However, in the connectomics dataset we observed a rather global membrane thickening at synaptic sites.
This may be partly due to a lack of resolution and the observed global overstaining.

The high magnification dataset was used to confirm the synaptic nature of the connections observed in the
connectomics dataset and the absence of PSD. The absence of PSD made it difficult to annotate polyadic
varicosities (one cluster of vesicles could be associated with two postsynaptic targets). In ambiguous cases
only the most likely partners were identified (larger surface contact), leading to a likely underestimation of

the overall number of synapses. Because most of the cells were not fully contained in the reconstructed

volume, it was not possible to assess the full collection of synapses between a given presynaptic and

postsynaptic neuron. For this reason, we refer to "synaptic site," a single anatomical release site or a single

synapse. Multiple synaptic sites from the same cell were classified as a single input. Additionally, because

cells had different lengths of arbor in the reconstructed volume, we expressed most of the results as synaptic

fraction and number of synaptic sites per 100 µm.

Although multiple annotators would have been ideal, due to project constraints and the difficulty of training

people in an entirely new dataset where most synapses are being described for the first time, all synaptic

annotations were made by a single annotator, giving consistency across annotations.

Clustering of cells based on connectivity

As described above, the reconstructed cells and processes of the vertical lobe were classified based

primarily on morphological criteria. We combined this classification with unsupervised clustering to obtain

a visual summary of the connectivity patterns found in the lobe. First, we produced a cell type wiring

diagram connecting the 7 main cell types (**Figure 2**). The nodes of this graph comprise cell types and the

weighted edges between pairs of cell types represent the number of synapses between the processes that

belonged to these cell types. We used a spring model to minimize the distances between the pairs of nodes

(representing pairs of cell types) that corresponded to a relatively larger number of synapses (Fruchterman

and Reingold 1991). This diagram revealed that synapses within a cell type are rare in the vertical lobe

(mainly in ascending fibers; see Results). In addition, visually inspecting this diagram clarified that the

connectivity was highly structured and far from an "all-to-all" connectivity. Each cell type was connected

to a small number of other cell types (see Results). This observation led us to consider the common inputs

and common outputs of specific processes (described below).

We also discovered that besides these uncommon within-type synapses, there were no connectivity cycles

in the cell-type wiring diagram. This observation led us to analyze the possible cycles in the wiring diagram

while considering individual processes (see above Criteria for cell classification). We then used an

unsupervised approach to visualize the connectivity among individual processes (**Figure 2 – figure**

supplement 1). This was done by considering the adjacency matrix $\mathbf{A_{NxN}}$ whose **(i,j)** entries represent the

number of synapses from the presynaptic neuron **i** to the postsynaptic neuron **j,** considering **N** neurons

(**N**=362). As specific groups of cells tended to synapse into other specific subsets of cells, a stereotypy

observed from the cell-type diagram, we considered the number of shared inputs and shared outputs

between individual processes, mathematically represented by the common output and common input

symmetric matrices, **A*A'** (**presynaptic**) and **A'*A** (**postsynaptic**), respectively. The presynaptic matrix

represents how many common targets exist for pairs of neurons and the postsynaptic matrix represents how

many common inputs exist for pairs of neurons. To visualize the tendencies of neurons to share inputs or

outputs, we considered the matrix **A*A'+ A'*A** which sums for every pair of neurons the number of shared

inputs and outputs, using subspace embedding (Koren 2005) and 100 dimensions, heuristically picking a

dimension that yielded sufficient separation between individual nodes (nearby dimension produced visually

similar outcomes). This embedding of the neurons in the plane resulted in a close positioning with other

neurons with a similarity of inputs and outputs (summed). The position of the entire layout and its

orientation are arbitrary and not used to derive biological observations. To combine the cell type- and

neuron-based diagrams, each neuron was located based on its type diagram shifted by its location in the

neuron-based wiring diagram (**Figure 2**).

Analysis of feed-forwardness

The cell type-based wiring diagram analysis revealed that connectivity cycles were rare and information

flowed in one direction among populations of cells belonging to different cell types (**Figure 2**). To

determine if there were cyclic synaptic information flows in the reconstructed circuit at the level of

individual neurites we used graph algorithms that detect directed cycles in directed graphs. Topological

sorting algorithms (Barth et al. 2004) that determine a linear order in cycle-free directed graphs were used

to detect cycles in the graph. A topological sorting is a sequence of nodes each of which is possibly

connected only to nodes proceeding in the list. Hence, we inspected all the edges in the topological sorting

that went backward. We found a few edges (n= 4) that violated the forwardness of the circuit and each of

these cases was related to a reciprocal synapse between a pair of nodes associated with a single synaptic

site between the cells. We also calculated the probability of observing this level of forwardness in random

graphs that agreed with the wiring diagram, but in which the direction of synaptic transmission was

randomized. This showed that the level of forwardness in the data is unlikely to occur by chance ($p<0.05$).

Classification of cell types from cell body morphology

Our reconstruction revealed that the SAM, CAM, and LN cell bodies lay in the VL cortex along with several

other cell types: glia, progenitor-like and mix cells (Others; **Figure 1 – supplement 4**). We noticed,

consistently with early studies, that the LNs had extremely large cell bodies and that the newly discovered

CAMs tended to have a slightly larger cell bodies than the SAMs. We tested whether these observations

could be used to classify cell types directly from the morphology of their cell bodies. We used 110 cells

whose somata were contained in the volume and their processes were distinctly characterized in the neuropil

by an expert annotator: 7 LNs (6.36%), 69 SAMs (62.73%), 16 CAMs (14.55%) and 18 others (16.36%).

The selection of this subset of cells was agnostic to the details of the classification process described here.

For each of these cells we considered the volume of the cell body as the number of annotated voxels

multiplied by the volume unit of a voxel, and the volume, surface area and sphericity (Wadell 1935) of the

ellipsoid best matching the geometry of the cell body. Ellipsoid approximations were computed from the

eigenvectors and eigenvalues of the covariance matrix of the voxel coordinates of the cell bodies.

Our tests showed that the information gained from these ellipsoids were sufficiently discriminative and

hence these 3 parameters (volumes, surface area and sphericity) were used for further analyses. Support

Vector Machines (SVMs) were used to train, cross-validate and predict the 3 main cell classes found in the

cortex: LN, SAM and CAM and a 4th class, Other, to represent the seemingly small number of other cell

types in the cortex (including glia, progenitor cells and possibly other cells of scarce representation). The

min, max, mean and std of these parameters were computed for each class as well as the training accuracy

(see Results).

In addition, a set of 49 cells (each cell was annotated twice) was used to determine the accuracy of the

SVMs and for leave-one-out cross validation (see Results). Once the SVM model was determined from the

training set, it was applied to a test set, comprising 1447 cells, that were densely annotated in the cortex

(using mEMbrain and VAST as described above). The number and percentage of each of the 4 classes were

calculated (see Results; **Figure 4**).

Wire-length calculation

To calculate the length of the processes (see Results; **Figure 2**), we used a thinning algorithm to map each

volumetric object into a skeleton and used graph theory algorithms to measure the total length of skeletons.

This was done by calculating the graph distance between skeleton nodes representing the end of the

volumetric object (a leaf; a node that is connected to at most one other node, e.g., of the last node in a

branch terminating at a synapse) and skeleton nodes representing branching (nodes that are connected to

more than two other nodes, e.g., a branching point of a dendrite-like structure or axonal branching). The

distances between adjacent skeletal nodes were defined using the Euclidean distance between the voxels in

the 3D 26-neigberhood mask (a 3x3 cube around each voxel). We then summed up all these distances to

measure the wire-length of each process. Occasionally the thinning algorithm produces a biologically

meaningless cycle due to topological holes in the annotated objects (either biological or imperfect

annotation). We guaranteed that only the shortest paths between special skeletal nodes (bifurcations and

leaves) were used for the wire-length calculation using minimum distance graph algorithms between leaf

and bifurcation nodes.

SFL bouton volume calculation

Synaptic boutons are morphologically easy to identify, and they were accordingly individually segmented

to allow volumetric measurements (see Results; **Figure 3**). Volumes of boutons were estimated from the

voxel counts of each bouton. A *post hoc* exact permutation test for independent samples compared the

volumes of the SFL SB and SFL LB. The p-values were adjusted according to the sequential Bonferroni

correction as described by Holm (1979).

1408

Subcellular reconstruction

1409

The geometry of the SFL large synaptic boutons and the postsynaptic structures they innervate were

reconstructed at the subcellular level using the high magnification dataset (see Results; **Figure 5**). Neuronal

elements of interest were manually segmented on VAST. The labeled images and metadata containing the

labels were exported and processed externally or accessed through the VAST API. Vesicles showing a clear

synaptic center in each section were counted by outlining intracellular space containing vesicles (see more

information below). Mitochondria were identified by the presence of an inner membrane with visible

cristae. Transport endoplasm, large electron-lucent tubular organelles, spanned the reconstructed volume

and appeared to be a continuous structure present throughout the length of the AM trunk. They showed

large swellings at the presynaptic site, particularly at the 'palm'. Discontinuous thin lucent tubular

organelles that sometimes appeared as dark lines were annotated as 'cisternae' (Cis). It is not clear whether

the Cis (up to 15 observed in one palm) were part of the larger endoplasmic reticulum network or were

isolated entities.

Counting, modelling, and measuring vesicles

1422

An annotator painted all individual vesicles of several compartments using distinct colors per compartment.

Distinct colors were used to annotate the active zones. The center of mass and areas of the vesicle cross-

sections were used to approximate the centers and spherical representations of individual vesicles, from

which spherical 3D mesh representations were computed. Occasionally, several vesicles of the same

compartment were overlapped spatially. This required us to use distance transforms and watersheds to

achieve separation of vesicles. Vesicle counts were made only after they were separated in space. Due to

the section thickness (30 nm), most of the vesicles were clearly appreciated in a single section, whereas

only the largest vesicles (> 60nm diameter) may have had a strong signal in two adjacent sections. The

annotator attempted to avoid overrepresentation of a vesicle, when possible, but since this task is not

unambiguous our counts likely represent an upper bound. Extremely small vesicle-like structures were not

annotated since it was assumed that these structures were likely represented in adjacent sections.

Rendering

1434

For rendering of all the traced images, 3D surface meshes of labeled objects were generated from VAST

using VastTools (written in MATLAB) and imported into 3D Studio Max 2020 (Autodesk Inc.) to generate

3D renderings of all the traced objects.

Data availability

The 'connectome dataset' is available at https://lichtman.rc.fas.harvard.edu/octopus_connectomes with 32

nm resolution. 8 nm resolution will be provided, and the segmentation files added prior publication.