Pathological crosstalk between AQP4-dependent ATP/Adenosine release and dopamine neurotransmission underlying depressive behavior during cocaine withdrawal

Okada, S.[1], Kobayashi, M.[1], Lee, H.[1], Nuralifah, C.[1,2] and Morita, M.[1]

[1]Department of Biology, Graduate School of Science, Kobe University, Kobe, 657-8501, Japan

2Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Indonesia, Depok, 16424, Indonesia

To whom correspondence should be addressed: Mitsuhiro Morita, Ph.D., Department of Biology, Kobe University, 1-1 Rokkodai Nada Kobe Hyogo 657-8501 Japan, Tel.: +81-78-803-6636; Fax: +81-78-803-6636; E-mail: mmorita@boar.kobe-u,ac,jp

Abstract

Our previous study has demonstrated the AQP4-dependent ATP release and following elevation of extracellular adenosine in hippocampal slices during hypo-osmotic treatment. In the present study, we examined the crosstalk between AQP4-dependent adenosine elevation and dopaminergic neurotransmissions, especially focusing on the mouse depressive behavior during cocaine withdrawal. The depressive behavior was mitigated in AQP4 knockout mice, and was sensitive to adenosine A_1 receptor antagonists. Thus, the influence of cocaine withdrawal on the adenosine or dopamine elevation in the slices from the striatum or medial prefrontal cortex, which is the focus of depression was analyzed by the originally-developed biosensor cells, which elevates intracellular Ca^{2+} in response to adenosine or dopamine. We have found that the evoked-dopamine release in the striatum is tonically suppressed via A_1 receptor in a AQP4 dependent manner, and this suppression is augmented by the cocaine withdrawal. In contrast, the evoked dopamine elevation in the medial prefrontal cortex was suppressed by treating slices with adenosine or an A_1 receptor antagonist, and the cocaine withdrawal abolished the suppression by A_1 receptor antagonist alone in an AQP4-dependent manner. Since a $GABA_A$ or group II metabotropic glutamate receptor antagonist occluded the suppression by A_1 receptor antagonist, it has been suggested that the glutamatergic and GABAergic terminals suppressing dopamine releases have A_1 receptors, which the cocaine withdrawal down-regulates in a AQP4-dependent manner. These results indicate that astrocytes modulate dopaminergic neurotransmissions via A_1 receptor in an AQP4-dependent manner, and suggest that cocaine causes depression by changing this modulation.

Introduction

Adenosine is a neuromodulator, which mediates presynaptic inhibition by A_1 receptor and synaptic

plasticity by A_{2A} receptor (Chen et al., 2001). Despite the associations of extracellular adenosine

concentration in the brain with circadian cycle (*) and pathological conditions including stroke (*)

and epilepsy (*), the pathways for adenosine release in the brain are not fully determined. Our

previous study using a biosensor cell line, which increases intracellular Ca^{2+} in response to

adenosine for measuring the adenosine release in hippocampal slices has demonstrated two novel

distinct pathways, namely the calcium channel-dependent neuronal adenosine release following

electrical stimulation and the AQP4-dependent ATP release and subsequent adenosine elevation due

to ATP degradation following hypoosmotic treatment (*). AQP4 is a water channel exclusively

expressed in astrocytes in the brain (*), thus hypoosmotic treatment likely induces the astrocytic ATP

release due to water influx and subsequent regulatory volume decrease (*). This pathway was further

supported by our related study showing the involvement of volume-regulated anion channel in the

ATP release from cultured astrocytes following mechanical stimulation (*).

Accumulating evidence indicates antagonistic interactions between adenosine and

dopamine in the brain. A_{2A} and D_2 receptors form heterodimer and suppress each other (Hillion et al.,

2002), while A_1 receptor inhibit dopamine release in the striatum and spinal cord (Acton et al., 2018;

Ross and Venton, 2015). The upregulation of dopamine neurotransmission by adenosine receptor

antagonist is a clinical strategy for treating Parkinson's disease patients (Kondo and Mizuno, 2015).

In addition, chemicals, which act on adenosine receptors, as well as modulate extracellular

concentration of adenosine have been shown to affect schizophrenia (*) or bipolar disorders (*),

which involve pathological changes of dopaminergic neurotransmission. Furthermore, astrocytes

likely participate in the crosstalk between adenosine and dopamine, because it has been reported that the A_1 receptor activation by the adenosine from astrocytes tonically inhibit dopamine release (https://pubmed.ncbi.nlm.nih.gov/35042768/) , as well as is associated with the anti-depression effect of sleep deprivation (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3566717/). Thus, the astrocyte atrophy in depression patients (Zhao et al., 2018) may cause symptoms by changing extracellular adenosine, as in the case of epilepsy, which is developed as a consequence of reduced extracellular adenosine due to the pathological change of astrocyte (Fedele et al., 2005).

The present study has hypothesized that the changes of crosstalk between the AQP4-dependent adenosine elevation and dopaminergic neurotransmissions underlie the mouse depression-like behavior during cocaine withdrawal (Filip et al., 2006). Our approach includes the behavioral analysis of wild type and $AQP4^{-/-}$ mice and the measurement of adenosine or dopamine elevation in brain slices from the foci of depression, the striatum and medial frontal cortex (mPFC) by biosensors as in our previous study (*). Once the association between the AQP4-dependnet adenosine elevation and the pathological changes of dopamine neurotransmission in depression is established, astrocytes and water dynamics in the brain will be recognized as a new diagnostic and therapeutic target of psychiatric disorders.

Materials & Methods

Animal experiments

All animal experiments were approved by the institutional animal care and use committee of Kobe University (Permission number: 25-09-04) and performed in compliance with the Kobe University Animal Experimentation Regulations. C57BLC background mice were maintained at 24℃ and

under 12h light/dark cycle. Water and food were provided ad libitum. Mice at least 8 weeks old were used for experiment. AQP4 knock out mice ($AQP4^{-/-}$) (Kitaura et al., 2009) were provided by Dr. Mika Terumitsu (National Center of Neurology and Psychiatry, Tokyo, Japan). Depression-like behavior during cocaine withdrawal was accessed by the forced swim test (FST) 14 to 17 days after five consecutive days of intraperitoneal cocaine injection at 15 mg/kg. For the FST, mice were acclimatized in an experiment room with white noise for one hour, then placed in a transparent cylinder 22 cm in diameter filled with water at 24°C at a height of 12.5 cm (Can et al., 2012). The mobility of mice was recorded as 8 min RGB videos at 15 fps by a web camera, C270n (Logicool, Tokyo, Japan) and quantified as described by Gao et al (Gao et al. 2014). Each frame in red color channel was converted to 8 bit gray-scale and subtracted by previous one. Then, the number of pixels with a value above five were counted and averaged for 15 consecutive frames as the mobility of mice in one second, The immobility time was measured as the duration, in which the mobility was below an arbitrary threshold in 60 secconds.

Biosensor cells

As a biosensor for adenosine or dopamine, a stable CHO cell lines expressing RCaMP, $G_{qi}5$ and mouse A_1 receptor (A_1-CHO) or human D2 receptor (D_2-CHO) were established. CHO cells were maintained in Ham's F12 medium containing 10% fetal bovine serum and antibiotics (50 U/ml penicillin and 50 µg/ml streptomycin). For $[Ca^{2+}]_i$ imaging, cells were seeded on coverslips. Clonal cells were obtained by transfecting with pCAG-cyto-RCaMP (Addgene), pME- $G_{qi}5$ (Yamashiro et al***) and pCMV-mA1 (OriGene) or DRD2-Tango (Addgene) by Lipofectoamine3000 and screening with 200 ug/mL G418. Then, the A_1-CHO or D_2-CHO was selected as a cell line

responding to 100 µM adenosine or 100 µM dopamine, respectively by Ca^{2+} imaging.

Brain slice preparation

Mice were anesthetized mice by intramuscular injection of ketamine (100mg/kg, Daiiichi-sankyo, Tokyo, Japan) and xylazine (10 mg/kg, Bayer Yakuhin, Osaka, Japan) and decapitated. Brains were extracted, put into an ice-cold cutting solution containing (in mM): 2 KCl, 1.5 NaH_2PO_4, 26 $NaHCO_3$, 6 $MgSO_4$, 0.2 $CaCl_2$, 10 D-glucose and 220 sucrose, and saturated with 95% O_2 and 5% CO_2. Coronal slices of 400µm thickness were cut using a microslicer (ZERO-1, Dosaka EM, Kyoto, Japan) and maintained in artificial cerebrospinal fluid (aCSF) containing (in mM): 124 NaCl, 2 KCl, 1.5 NaH_2PO_4, 26 $NaHCO_3$, 1 $MgSO_4$, 2 $CaCl_2$, 10 D-グルコース, for more than one hour before experiment.

Calcium imaging

Slices were placed on the biosensor cells and perfused with aCSF at 5 mL/min. The RCaMP images of the biosensor cells were obtained every one second by using an inverted microscope (IX-70, Olympus, Tokyo, Japan) equipped with an objective (UApo/340 20x / 0.70, Olympus) and a cooled-CCD camera (Orca-R2, Hamamatsu Photonics, Hamamtsu, Japan), and F/F_0 was calculated by ImageJ. Electrical stimulation was given by house-made concentric electrode, isolator (ISO-Flex, Funakoshi, Tokyo, Japan) and pulse generator (Electronic Stimulator, Nihon Koden, Tokyo, Japan). For hypoosmotic treatment, the hypoosmotic aCSF of the osmolarity indicated were prepared by reducing NaCl concentration, and slices were pretreated with the isosmotic low NaCl aCSF, which was supplemented with sucrose in the place of NaCl to achieve iso-osmolarity.

Reagents

If not specified, reagents purchased from Nacalai tesque (Kyoto, Japan).

Statistical analysis

For statistical analysis by Excel 2016, 50 biosensor cells were randomly selected from each slice in Fig2BC and 5AC while eight responding biosensor cells were randomly selected in Fig3, 4, 5BD, 6, 7 and 8.

Results

The involvement of AQP4-dependent adenosine elevation in the mouse depression-like behavior during cocaine withdrawal.

The association of the AQP4-dependent adenosine elevation with the mouse depression-like behavior during cocaine withdrawal was examined by the FST. The immobility times of wild type mice treated with saline (WT / (-)) or cocaine (WT / Cocaine), were respectively 21.33 ± 13.00 sec or 35.82 ± 12.08 sec between 2 and 3 min after starting test, and 19.53 ± 14.09 sec or 36.74 ± 8.59 sec between 3 and 4 min, indicating that the cocaine withdrawal significantly increased the immobility time, thus induced depression-like behavior (Fig. 1A). This depression-like behavior was abolished in AQP4$^{-/-}$ (Fig. 1B). The immobility times of saline (AQP4$^{-/-}$ / (-)) or cocaine (AQP4$^{-/-}$ / Cocaine) treated AQP4$^{-/-}$ mice were respectively 34.77 ± 10.67 sec or 14.83 ± 16.25 sec between 4 and 5 min after stating test, 36.07 ± 10.53 sec or 20.92 ± 15.94 sec between 5 and 6 min, 38.28 ± 12.06 sec or 20.57 ± 14.10 sec between 6 and 7 min, indicating that the cocaine withdrawal significantly reduced the immobility time, thus failed to induce the depression-like behavior in AQP4$^{-/-}$ mice. Then, the involvement of adenosine in the depression-like behavior was examined by

a non-selective adenosine receptor antagonist, caffeine. Saline or 50 mg/kg caffeine was administered to wild type mice during the cocaine withdrawal 40 min prior to the FST. The immobility time of wild type mice during cocaine withdrawal treated with saline (WT / Cocaine / (-)) or caffeine (WT / Cocaine / Caffeine) were respectively 15.64 ± 7.78 sec or 5.74 ± 7.19 sec between 1 and 2 min after starting test, 27.61 ± 6.42 sec or 9.70 ± 7.31 sec between 2 and 3 min, 23.42 ± 5.39 sec or 6.33 ± 8.71 sec between 3 and 4 min, 31.56 ± 9.35 sec or 9.85 ± 11.43 sec between 4 and 5 min, 39.56 ± 8.47 sec or 13.50 ± 10.11 sec between 5 and 6 min, 39.82 ± 10.29 sec or 21.58 ± 13.59 sec between 6 and 7 min, and 44.94 ± 3.37 sec or 19.66 ± 13.22 sec between 7 and 8 min, indicating that caffeine significantly reduced the immobility time as in AQP4$^{-/-}$ mice, thus inhibited the depression-like behavior (Fig. 1C). The effect of caffeine was likely mediated by adenosine A_1 receptor, because an A_1-selective antagonist, 8-Cyclopentyl-1,3-dipropylxanthine (DPCPX), which was administered at 4mg/kg, 30 min prior to the FST also abolished the depression-like behavior (Fig. 1D). The immobility time of wild type mice during cocaine withdrawal treated with saline (WT / Cocaine / (-)) or DPCPX (WT / Cocaine / DPCPX) were respectively 6.48 ± 5.94 sec or 0 ± 0 sec between 1 and 2 min, 21.65 ± 11.55 sec or 6.17 ± 7.01 sec between 3 and 4 min, 37.71 ± 5.49 sec or 5.67 ± 6.39 sec between 5 and 6 min, 36.75 ± 14.42 sec or 5.68 ± 7.04 sec between 6 and 7 min, and 40.28 ± 14.14 sec or 20.97 ± 10.60 sec between 7 and 8 min, indicating that DPCPX also significantly reduced the immobility time, thus inhibited the depression-like behavior. These results indicates the involvement of AQP4 and adenosine A_1 receptor in the depression-like behavior during cocaine withdrawal and suggest that cocaine causes depression by changing the AQP4-dependent adenosine elevation.

The upregulation of AQP4-depedndet adenosine elevation in the striatum during cocaine withdrawal

The adenosine elevation in the ventral part of striatum including nucleus accumbens following electrical stimulation or hypoosmotic treatment was examined by the A_1-CHO. An electrical stimulation (30Hz, 150 pulses), which had induced adenosine elevations in hippocampal slices (Yamashiro et al., 2017) did not elevate adenosine in striatum slices (Fig2A). Thus, it was suggested that neuronal adenosine release is limited the striatum. Meanwhile, adenosine elevations were detected during a hypoosmotic treatment at -124 mOsm. In addition, the striatal slices from mice during cocaine withdrawal showed $223.14 \pm 28.60\%$ significantly larger hypotonically-induced adenosine elevation quantified as the area under the curve (AUG) of Ca^{2+} elevations in the A_1-CHO, than saline treated mice (Fig. 2B). Furthermore, the upregulation of hypoosmotically-induced adenosine elevation during cocaine withdrawal was occluded in AQP4$^{-/-}$ mice (Fig. 2C). Thus, both the increase of adenosine elevation and the depression-like behavior during cocaine withdrawal were AQP4-dependent, suggesting that cocaine develops depression by increasing the AQP4-dependent adenosine elecvation.

The suppression of dopamine release by the AQP4-depedet adenosine tone in the striatum

The influence of adenosine on the dopamine releases in the striatum was examined. An electrical stimulation (40 pulses, 30 Hz) to striatal slices induced dopamine releases, which were measured as the Ca^{2+} elevations in the D_2-CHO. 10 μM adenosine significantly reduced the dopamine release (WT / Ado) to $61.92 \pm 7.44\%$ of control (WT / (-)) (Fig. 3A), while a non-selective adenosine

receptor antagonist, 1 mM caffeine 146.67±23.50% significantly increased dopamine release (WT / Caffeine) to control (WT / (-)) (Fig. 3B). Furthermore, an A_1 antagonist, 10 μM DPCPX 156.32±17.39% significantly increased dopamine release (WT / DPCPX) to control (WT / (-)) (Fig. 3C), while an A_{2A} receptor antagonist, 50 nM SCH58261 did not change the dopamine release (WT / SCH) from control (WT / (-)) (Fig. 3D). These results indicate that exogenous adenosine, as well as endogenous adenosine tone suppress dopamine releases via A_1 receptor in the striatum. The involvement of AQP4 in the adenosine tone suppressing the dopamine releases in the striatum was examined using AQP4$^{-/-}$ mice. The dopamine releases in the striatal slices prepared from AQP4$^{-/-}$ mice (AQP4$^{-/-}$) was 181.38±19.54% significantly larger than control (WT) (Fig. 3E). In addition, DPCPX significantly decreased the dopamine release of AQP4$^{-/-}$ mice (AQP4$^{-/-}$ / DPCPX) to 54.70±8.49% of control (AQP4$^{-/-}$ / (-) (Fig. 3F). Thus DPCPX increased dopamine release in wild type mice, but did not in AQP4$^{-/-}$ mice. These results indicates that the striatal dopamine release is suppressed by the AQP4-dependent adenosine tone and its A_1 receptor activation.

DPCPX increased dopamine release in wild type mice, while decreased in AQP4$^{-/-}$ mice

The further suppression of dopamine release of AQP4$^{-/-}$ mice in the presence of DPCPX propose a facilitation of dopamine release by an A_1 receptor subpopulation, which is activated by adenosine release through other pathway rather than the AQP4 dependent pathway as will be discussed later in mPFC results.

Reduced dopamine release during depressive behavior

Dopamine releases in the striatum during depressive behavior was examined. The dopamine release

of cocaine-treated mice (WT / Cocaine) was significantly smaller to $68.35 \pm 8.53\%$ of control (WT / (-)) (Fig. 4A). DPCPX (WT / Cocaine / DPCPX) $137.99 \pm 18.11\%$ significantly increased the dopamine release of cocaine-treated WT mice (WT / Cocaine / (-)) as in the case of saline-treated WT (Fig. 4B). These results suggests that the suppression of dopamine release by endogenous adenosine via A_1 receptor persists during depressive behavior. In contrast, the dopamine release of cocaine-treated $AQP4^{-/-}$ mice ($AQP4^{-/-}$ / Cocaine) was not significantly different from control ($AQP4^{-/-}$ / (-)) suggesting the ablation of reduced dopamine release by cocaine in $AQP4^{-/-}$ mice (Fig. 4C). These results indicate the lack of depressive behavior and associated reduction of dopamine release in $AQP4^{-/-}$ mice. In addition, DPCX did not change the dopamine release of cocaine-treated $AQP4^{-/-}$ mice ($AQP4^{-/-}$ / Cocaine / DPCPX and $AQP4^{-/-}$ / Cocaine / (-)) (Fig. 4D), indicating the loss of the A_1 receptor-mediated suppression of dopamine release in cocaine treated $AQP4^{-/-}$ mice. From these results, cocaine is supposed to induce depressive behavior by the increase of AQP4-dependent adenosine increase and subsequent suppression of dopamine release by A_1 receptors activation. Furthermore, A_1 receptor antagonists likely suppressed depressive behavior by restoring adenosine tone in the striatum.

Adenosine releases in mPFC

Since the altered neural activities in the mPFC is known to cause depressive behavior (Furuyashiki, 2012), the AQP4-dependent adenosine release in the mPFC was analyzed by using A_1-CHO. Both 150 pulses electrical stimulation at 30 Hz and -82.7 mOsM hypo-osmotic treatment induced adenosine releases in mPFC slices. The AUC after electrical stimulation did not show significant difference between $AQP4^{-/-}$ mice ($AQP4^{-/-}$) and control (WT) (Fig. 5A). In contrast,

hypo-osmotically-induced adenosine release of AQP4$^{-/-}$ mice (AQP4$^{-/-}$) was $64.14 \pm 9.24\%$ significantly reduced to control (WT) (Fig. 5B). These results indicate that adenosine release in the mPFC is in line with our previous study using hippocampal slices, where both electrical stimulation and hypo-osmotic treatment induce adenosine release and the hypo-osmotic release depends on AQP4 (Yamashiro et al., 2017). In order to determine the involvement of mPFC adenosine in depressive behavior, adenosine releases in cocaine-treated mice were examined. As results, both electrically and hypo-osmotically-induced adenosine release of cocaine-treated mice (WT / Cocaine) did not significant difference from those of control (WT / (-)) (Fig. 5C,D). Thus, it was suggested that these adenosine releases in the PFC were not involved in depressive behavior.

The involvement of adenosine in dopamine release in the mPFC

Since the decreased dopamine neurotransmission in the mPFC during depressive behavior is proposed (Furuyashiki, 2012), the involvement of adenosine in evoked-dopamine release in the mPFC was examined by using D$_2$-CHO. As in the striatum, the dopamine releases were induced by 40 pulses electrical stimulation at 30 Hz (arrow) in the mPFC. Exogenous adenosine $67.12 \pm 8.13\%$ significantly reduced dopamine release in the mPFC (WT / Ado) to control (WT / (-)) (Fig. 6A). Meanwhile, caffeine $66.96 \pm 5.82\%$ significantly reduced in dopamine release in the mPFC (WT / Caffeine) to control (WT / (-)) (Fig. 6B). In addition, DPCPX $75.25 \pm 5.45\%$ significantly reduced dopamine release (WT / DPCPX) to control (WT / (-)) (Fig. 6C), while SCH58261 did not change dopamine release (WT / SCH) to control (WT / (-)) (Fig. 6D). These results demonstrate the suppression of dopamine release in the PFC both by exogenous adenosine and A$_1$ receptor that the A$_1$ receptor activation by endogenous adenosine enhance dopamine release as in the opposite case in

the striatum.

Downregulation of A_1 receptor in the mPFC during depressive behavior

Dopamine releases in the mPFC during depressive behavior were examined by D_2-CHO. The electrically induced dopamine releases in the mPFC slices from cocaine treated mice (WT / Cocaine) was not significantly different from control (WT / (-)) (Fig. 7A). This suggests the dopamine release in mPFC was not affected by the cocaine. However, DPCPX, which increased the dopamine releases in the mPFC of naïve mice, did not change the dopamine release in cocaine treated mice (WT / Cocaine / DPCPX and WT / Cocaine / (-)), suggesting the loss of the DPCPX induced increases of dopamine releases by cocaine (Fig. 7B). Since cocaine did not change adenosine release in the mPFC, the failure of DPCPX to change dopamine release in cocaine-treated mPFC is supposed to reflect downregulation of A_1 receptors. This point was examined in AQP4 $AQP4^{-/-}$ mice. Cocaine treatment did not change dopamine release in the mPFC of $AQP4^{-/-}$ mice (($AQP4^{-/-}$ / (-) and $AQP4^{-/-}$ / Cocaine) (Fig. 7C). Meanwhile, DPCPX $61.26 \pm 5.27\%$ significantly reduced dopamine release of cocaine-treated $AQP4^{-/-}$ mice ($AQP4^{-/-}$ / Cocaine / DPCPX) to control ($AQP4^{-/-}$ / Cocaine / (-)) (Fig. 7D). Hence, both the depressive behavior and the down-regulation of A_1 receptors were lost in $AQP4^{-/-}$ mice. All together, these results suggest that cocaine induces depressive behavior by down-regulating A_1 receptor in the mPFC through an AQP4-dependent pathway.

AQP4-dependent adenosine tone facilitates dopamine release in the mPFC.

The involvement of AQP4-dependent adenosine tone in evoked-dopamine release in the mPFC was examined by measuring the evoked-dopamine release of $AQP4^{-/-}$ mice using D_2-CHO. The dopamine

release in the mPFC of AQP4$^{-/-}$ mice (AQP4$^{-/-}$) was not significantly different from wild type mice (WT) (Fig. 8A). The dopamine release in the mPFC of AQP4$^{-/-}$ mice in the presence of DPCPX, which decrease the dopamine release in mPFC of WT mice was also not different from control (AQP4$^{-/-}$ / (-)) (Fig. 8B). These results indicate the lack of A$_1$ receptor-mediated reduction of dopamine release in the mPFC of AQP4$^{-/-}$ mice. Since hypo-osmotically-induced adenosine increase was smaller in the mPFC of AQP$^{-/-}$ mice, the loss of A$_1$ receptor-mediated reduction of dopamine release is likely due to the loss of A$_1$ receptor rather than the change of AQP4-dependent adenosine tone.

Two adenosine-dependent pathways regulating dopamine release.

The present discrepancy that both exogenous adenosine and DPCPX reduce dopamine release in mPFC was further examined. A$_1$ receptor is commonly involved in pre-synaptic inhibition, but the inhibition of A$_1$ receptor suppressed dopamine release in the mPFC. Pre-synaptic inhibition of dopamine release is mediated not only by A$_1$ receptor, but also other receptors, such as metabotropic glutamate receptor (*) or GABA$_A$ receptor (Yonezawa et al., 1998). Thus, one possibility is that a subpopulation of A$_1$ receptor inhibits glutamate or GABA release, which are simultaneously activated with dopamine release by electrical stimulation and pre-synaptically inhibit dopamine release.

To test this possibility, the effects of DPCPX in the presence of mGluR or GABA$_A$ receptors antagonist were examined. The effect of DPCPX was examined in the presence of GABA$_A$ receptor antagonist, 100 µM picrotoxin. In the presence of picrotoxin and DPCPX, the dopamine release (WT / PTX / DPCPX) was not significantly different from the control (WT / PTX / (-)) (Fig. 9A).

Therefore, GABA$_A$ receptor antagonist occlude the effect of DPCPX. Since A$_1$ receptors are mostly resided in excitatory synapses rather than inhibitory ones (Chen et al., 2013), DPCPX supposed to affect on glutamatergic synapses. Since Group II metabotropic glutamate receptors (Group II mGluRs) regulate dopamine release in the mPFC and nucleus accumbens (Cartmell et al., 2000; Gupta & Young, 2018), the possibility that DPCPX affects glutamatergic terminal involving Group II mGluR was examined.

The group II mGluR antagonist LY341495 was used to test this hypothesis and the dopamine release in the presence of 500 nM LY341495 and DPCPX and control (WT / LY / DPCPX and WT / LY / (-)) did not show significant difference (Fig. 9B). Hence, group II mGluR antagonist also occluded the effect of DPCPX on the dopamine release. If the DPCPX-sensitive glutamate release acts on ionotropic glutamate receptor and influences on dopaminergic neuron via neural circuit, AMPA receptor antagonist CNQX and NMDA receptor antagonist APV may affect. However, the dopamine release in the presence of DPCPX, together with 10 µM CNQX and 50 µM APV (WT / CNQX+APV / DPCPX) was 59.82 ± 4.12% significantly reduced to control (WT / CNQX+APV / (-)) (Fig. 9C), excluding the involvement of ionotropic glutamate receptors. These results indicate that the DPCPX affects the glutamate and GABA releases, which act on group II mGLuR or GABA$_A$ receptor locating at the presynaptic terminal releasing dopamine, respectively. Finally, the A$_1$ receptor down-regulation by cocaine treatment at dopamine release terminal, which is sensitive to exogenous adenosine in the mPFC. The dopamine release of cocaine treated mice in the presence of exogenous adenosine (WT / Cocaine / Ado) 73.79±6.82% significantly reduced to control (WT / Cocaine / (-)) (Fig. 9D). This result reflects that cocaine occluded the effect of DPCPX while did not

influence on effect of exogenous adenosine. Overall, the dopamine release in the mPFC were regulated by two adenosine-dependent pathways, one pathways involving glutamate and GABA was down-regulated bycocaine treatment, while another direct adenosine inhibition of dopamine releasing terminal was not affected.

Discussion

Pathological changes in adenosine and dopamine release associated with depressive-like behavior during cocaine withdrawal has been investigated. Depressive-like behavior accessed by the FST suggested that AQP4-mediated increase in adenosine causes depressive-like behavior via A_1 receptors, because genetic disruption of AQP4 and pharmacological inhibition of A_1 receptor suppressed the depressive-like behavior. Adenosine release in the striatum was not detected upon electrical stimulation while was detected by a hypoosmotic treatment, but it required greater osmotic change than in other brain regions, and was not affected by disruption of the AQP4 gene. This indicates the difference of extracellular adenosine dynamics between in the striatum, where A_{2A} receptors are highly expressed (*), and in the cortex and hippocampus. In contrast, adenosine release in the mPFC, adenosine was induced by electrical stimulation and hypoosmotic treatment, and adenosine release upon hypoosmotic treatment was AQP4 dependent, similar to our previous study on the hippocampus (Yamashiro et al., 2017). In contrast, dopamine release upon electrical stimulation was detected in both the striatum and mPFC. In the striatum, dopamine release was inhibited by adenosine, whereas it was enhanced by A_1 receptor antagonist, suggesting tonic inhibition of dopamine release by adenosine. Furthermore, cocaine increased adenosine release

while decreased dopamine release, indicating that cocaine promoted the tonic inhibition of dopamine release by adenosine in the striatum. In contrast, both adenosine and an A_1 receptor antagonist suppressed dopamine release in the mPFC, and cocaine treatment abolished only the effect of A_1 receptor antagonist in a AQP4-dependent manner. Thus, cocaine likely down-regulated A_1 receptors by an AQP4-dependent mechanism in the mPFC, resulting in loss of endogenous adenosine modulation of dopamine release. Since the effects of A_1 receptor antagonist were occluded by $GABA_A$ or gruop II mGluR antagonists, A_1 receptor antagonist likely acted on presynaptic terminal releasing GABA or glutamate, which are known to suppresses dopamine release via GABAA receptor or group II mGluR (*). These results demonstrate modulations of dopamine neurotransmission by AQP4 and adenosine in both the striatum and mPFC, and their involvement in depressive-like behavior. However, depression-like behavior during cocaine withdrawal more likely attributed to the change in the striatum, where A_1 antagonist restored dopamine release as in the case of depression-like behavior, in contrast to the suppression of dopamine release by both exogenous adenosine and A_1 antagonist in the mPFC.

The mitigation of depression-like behavior during cocaine withdrawal in AQP4$_{-/-}$ mice and the suppression of dopamine release by AQP4-dependent increase in adenosine in the striatum are supported by a preceding report of increased dopamine concentrations in the striatum as measured by microdialysis and associated cognitive and psychiatric changes in AQP4$_{-/-}$ mice (Szu and Binder, 2016). The antidepressant effects of caffeine and DPCPX against depression-like behavior during cocaine withdrawal is consistent with previously reported antidepressant effects of these compounds in a forced swimming test using normal mice (Szopa et al., 2016) (Szopa et al., 2018). The present study is the first demonstration of crosstalk between AQP4-dependent increase in

adenosine and pathological change of dopamine neurotransmission in depression by measuring

adenosine and dopamine releases and analyzing their changes during cocaine withdrawal.

The failure of detecting adenosine release upon electrical stimulation in striatal slices is

consistent with a report of low detection success rates in the measurement of similarly-evoked

adenosine release in slices by fast scan cyclic voltammetry (Pajski and Venton, 2013). Meanwhile,

evoked-adenosine releases in the striatum *in vivo* were reported by using microdialysis (Cechova and

Venton, 2008; Cechova et al., 2010). One possible explanation for the failure in slice is depletion of

neuronal adenosine source due to high oxygen and glucose condition for slices. Another explanation

is that AQP4-dependent increase in adenosine is induced by electrical stimulation *in vivo*, but not in

slices lacking blood flow, which is likely essential for water exchange via AQP4 locating at the

interface between astrocytes and blood vessels. We suppose that neuronal activities induce water

influx from blood flow to astrocyte via AQP4, because they increase glucose uptake or glycogen

degradation, both of which increase osmotic pressure in astrocytes. Adenosine release in striatal

slices was induced by hypoosmotic treatment at -120 mOsM, but not at -82.7 mOsM, which was

effective in our previous study using hippocampal slices (*) and the present study using mPFC slices.

In addition, AQP4$^{-/-}$ mice showed reduction of hypoosmotically-induced adenosine release in

cocaine treated mice, but not in normal ones. Thus, it is supposed that astrocytes are capable of

releasing ATP even without AQP4 following intense change of osmolarity, and the availability of

AQP4 in the striatum is upregulated by cocaine treatment. We have been investigating changes of

astrocyte structure and AQP4 expression in the striatum during cocaine withdrawal, however no

significant result is obtained.

Dopamine release in the striatum was suppressed by exogenous adenosine, while

increased by DPCPX and in AQP4$^{-/-}$ mice. These results indicates the tonic suppression of dopamine release in the striatum by adenosine released through AQP4-dependent pathway. This point is consistent with previous reports showing presynaptic inhibition of dopamine release by A$_1$ receptor (Yabuuchi et al., 2006), but not by A$_{2A}$ receptor (Ross and Venton, 2015). In the present study, DCPCX significantly reduced evoked-dopamine release in AQP4-/- striatum. This result can be interpreted as the suppressive effect of DPCPX in the mPFC, namely, adenosine release by an AQP4-indeoendent pathway in the striatum presynaptically suppresses other neuromodulator, such as GABA and glutamate, which negatively regulate dopamine release.

References

A.

B.

C.

D.

Fig. 1 Depressive behavior of cocaine-treated mice were suppressed by adenosine receptor antagonists or in AQP4$^{-/-}$ mice. Depressive behavior during cocaine withdrawal was assessed by the time-dependent changes of immobility time in forced swim test and the influences of pharmacological or genetical interventions were examined. A. Cocaine induced depressive behavior. Wild type mice were treated with saline (WT / (-), n = 8) or cocaine (WT/Cocaine, n=8). Saline or cocaine were treated as described in Materials and Method for inducing depressive behavior during cocaine withdrawal. B. Depressive behaviors of AQP4$^{-/-}$ mice. AQP4$^{-/-}$ mice were treated with saline (AQP4$^{-/-}$ / (-), n = 6) or cocaine (AQP4$^{-/-}$ / cocaine, n = 7). C. The effect of a non-selective adenosine receptor antagonist, caffeine. Saline (WT / Cocaine / (-), n = 6) or 50 mg/kg caffeine (WT / Cocaine / Caffeine, n = 7) were treated 40 min prior to forced swim test. D. The effect of an A$_1$ adenosine receptor selective antagonist DPCPX. Saline (WT / Cocaine / (-), n = 5) or 4 mg/kg DPCPX (WT / Cocaine / DPCPX, n = 5) were treated 30 min prior to forced swim test. Mean ± SD. * $p<0.05$, ** $p<0.01$, *** $p<0.001$, NS, not significant.

Fig. 2 Increased adenosine release in the striatum during depressive behavior. Adenosine

from striatum slices were measured by A_1-CHO (Striatum / A_1-CHO). A. Representative time

dependent changes of $[Ca^{2+}]_i$ during electrical stimulation (arrowhead, 30 Hz, 150 pulses) to slices

prepared from wild type mice (WT). B. Adenosine releases from slices of saline- (WT / (-), n = 6

slices) or cocaine-treated wild type mice (WT / cocaine, n = 6 slices) during hypo-osmotic treatment.

Representative time dependent changes of $[Ca^{2+}]_i$ of A_1-CHO (left, center). Hypo-osmotic treatment

(HO; -124 mOsM) were indicated by lines. $[Ca^{2+}]_i$ increase during HO treatment were shown as relative area under the curves (AUC, right) normalized by control. C. Adenosine releases from slices of saline (AQP4$^{-/-}$ / (-), n = 6 slices) or cocaine (AQP4$^{-/-}$ / Cocaine, n = 6 slices) treated AQP4$^{-/-}$ mice during hypo-osmotic treatment. Mean $\pm$ SE, *** $p<0.001$, NS, not significant.

A. Striatum / D$_2$-CHO

WT / (-)   WT / Ado

$\Delta F/F_0=1$

20s

30 Hz x 40

Relative AUC (%)

**

WT / (-)   WT / Ado

B. Striatum / D$_2$-CHO

WT / (-)   WT / Caffeine

Relative AUC (%)

*

WT / (-)   WT / Caffeine

C. Striatum / D$_2$-CHO

WT / (-)   WT / DPCPX

Relative AUC (%)

**

WT / (-)   WT / DPCPX

D. Striatum / D$_2$-CHO

WT / (-)   WT / SCH

Relative AUC (%)

NS

WT / (-)   WT / SCH

E. Striatum / D$_2$-CHO

WT   AQP4$^{-/-}$

Relative AUC (%)

***

WT   AQP4$^{-/-}$

F. Striatum / D$_2$-CHO

AQP4$^{-/-}$ / (-)   AQP4$^{-/-}$ / DPCPX

Relative AUC (%)

**

AQP4$^{-/-}$ / (-)   AQP4$^{-/-}$ / DPCPX

Fig. 3 Suppression of evoked-dopamine release in the striatum by AQP4 dependent increase of adenosine. Evoked-dopamine releases in striatal slices were measured by D_2-CHO (Striatum / D_2-CHO). A. The effect of adenosine. Slices from wild type mice (WT) were electrically stimulated (30 Hz, 40 pulses) in the absence (WT / (-), n = 4 slices) or presence of 10 µM adenosine (WT / Ado, n = 5 slices). Representative time dependent changes of $[Ca^{2+}]_i$ of D_2-CHO (left, center) during electrical stimulation (arrow). $[Ca^{2+}]_i$ increases of D_2-CHO for one minute after stimulation were shown as relative AUC (right) normalized by control. B. The effect of caffein. Evoked-dopamine release in the absence (WT / (-), n = 6 slices) or presence of 1 mM caffein (WT / Caffeine, n = 6 slices). C. The effect of DPCPX. Evoked-dopamine release in the absence (WT / (-), n = 6 slices) or presence of 10 µM DPCPX (WT / DPCPX, n = 4 slices). D. The effect of an A_{2A} selective antagonist SCH58261. Evoked-dopamine release in the absence (WT / (-), n = 6 slices) or presence of 10 µM DPCPX (WT / DPCPX, n = 4 slices). E. The involvement of AQP4. Evoked-dopamine releases in wild type (WT, n = 6 slices) and AQP4 knock out mice ($AQP4^{-/-}$, n = 7 slices). F. The effect of DPCPX on $AQP^{-/-}$ mice. Evoked-dopamine releases in the absence ($AQP4^{-/-}$ / (-), n = 7 slices) or presence of DPCPX ($AQP4^{-/-}$ / DPCPX, n = 5 slices). Mean ± SE. * $p<0.05$, ** $p<0.01$, *** $p<0.001$, NS, not significant, t-test.

Fig. 4 Reduction of evoked-dopamine release in the striatum of mice during depressive

behavior and its reversal in AQP4 knock out mice. Dopamine releases in striatal slices were

measured by D$_2$-CHO (Striatum / D$_2$-CHO). A. The effect of cocaine. Slices from wild type mice

(WT) with (WT / cocaine, n = 7 slices) or without cocaine treatment (WT / (-), n = 10 slices) were

electrically stimulated (30 Hz, 40 pulses). B. The effect of DPCPX in cocaine treated mice.

Evoked-dopamine release in the absence (WT / Cocaine / (-), n = 6 slices) or presence of 1 mM

caffein (WT / Cocaine / DPCPX, n = 5 slices). C. The effect of cocaine on AQP4$^{-/-}$ mice.

Evoked-dopamine releases in AQP4$^{-/-}$ mice with (AQP4$^{-/-}$ / cocaine, n = 9 slices) or without cocaine

treatment (AQP4$^{-/-}$ / (-), n = 9 slices). D. The effect of DPCPX in cocaine-treated AQP4$^{-/-}$ mice.

Evoked-dopamine release in cocaine-treated AQP4$^{-/-}$ mice in the absence (AQP4$^{-/-}$ / Cocaine /(-), n =

7 slices) or presence of cocaine (AQP4$^{-/-}$ / Cocaine / DPCPX, n = 6 slices). Mean ± SE. * $p < 0.05$,

NS, not significant, t-test.

Fig. 5 Adenosine release in mPFC is not associated with depressive behavior. Adenosine

from medial prefrontal cortex (mPFC) slices were measured by A_1-CHO (mPFC / A_1-CHO). A.

Evoked-adenosine releases from wild type (WT, n = 6 slices) and AQP4$^{-/-}$ mice (AQP4$^{-/-}$, n = 5 slices). Representative time dependent changes of $[Ca^{2+}]_I$ of A_1-CHO (left, center) during electrical stimulation (150 pulses at 30 Hz, arrowhead). $[Ca^{2+}]_i$ increases of A_1-CHO for one minute after stimulation were shown as relative AUC (right) normalized by control. B. Slices from wild type mice (WT, n = 6 slices) and AQP4$^{-/-}$ mice (AQP4$^{-/-}$, n = 6 slices) were hypo-osmotically stimulated (HO; -82.6 mOsM) as indicated by lines. C. Evoked-adenosine releases in the absence (WT / (-), n = 7 slices) or presence of cocaine (WT / Cocaine, n = 6 slices. D. Hypo-osmotically-induced adenosine release in mice with (WT / cocaine, n = 6 slices) or without cocaine treatment (WT / (-), n = 6 slices). Mean ± SE, ** p<0.01, NS, not significant.

Fig. 6 Suppression of evoked-dopamine release in the mPFC by adenosine. Dopamine from

mPFC slices were measured by D_2-CHO (mPFC / D_2-CHO). A. The effect of adenosine. Slices from

wild type mice (WT) were electrically stimulated (30 Hz, 40 pulses) in the absence (WT / (-), n = 8

slices) or presence of adenosine (WT / Ado, n = 9 slices). B. The effect of caffein. Evoked-dopamine

releases in the absence (WT / (-), n = 8 slices) or presence of caffein (WT / Caffeine, n = 8 slices). C.

The effect of DPCPX. Evoked-dopamine releases in the absence (WT / (-), n = 8 slices) or presence

of DPCPX (WT / DPCPX, n = 9 slices). D. The effect of SCH58261. Evoked-dopamine releases in

the absence (WT / (-), n = 10 slices) or presence of SCH58261 (WT / SCH, n = 10 slices). Mean ±

SE. * p<0.05, ** p<0.01, NS, not significant, t-test.

Fig. 7 Downregulation of A_1 receptor in the mPFC of cocaine-treated mice. Dopamine from

mPFC slices were measured by D_2-CHO (mPFC / D_2-CHO). A. The effect of cocaine. Slices from

wild type mice (WT) with (WT / cocaine, n = 12 slices) or without cocaine treatment (WT / (-), n = 9 slices) were electrically stimulated (30 Hz, 40 pulses). B. The effect of DPCPX in cocaine treated mice. Evoked-dopamine releases in the absence (WT / Cocaine / (-), n = 8 slices) or presence of DPCPX (WT / Cocaine / DPCPX, n = 8 slices). C. The effect of cocaine in AQP4$^{-/-}$ mice. Evoked-dopamine releases in absence (AQP4$^{-/-}$ / (-), n = 9 slices) or presence of cocaine (AQP4$^{-/-}$ / cocaine, n = 8 slices). D. The effect of DPCPX in cocaine treated AQP4$^{-/-}$ mice. Evoked-dopamine releases in the absence (AQP4$^{-/-}$ / Cocaine /(-), n = 8 slices) or presence of DPCPX (AQP4$^{-/-}$ / Cocaine / DPCPX, n = 7 slices). Mean ± SE. * $p<0.05$, NS, not significant, t-test.

A. mPFC / D$_2$-CHO

Fig. 8 Facilitation of evoked-dopamine release in the mPFC by AQP4-dependent increase of adenosine. Evoked-dopamine release from mPFC slices were measured by D$_2$-CHO (mPFC / D$_2$-CHO). A. The involvement of AQP4. Slices from wild type mice (WT, n = 7 slices) and AQP4 knock out mice (AQP4$^{-/-}$, n = 7 slices) were electrically stimulated (30 Hz, 40 pulses). B. The effect of DPCPX on AQP4$^{-/-}$ mice. Evoked-dopamine releases in the absence (AQP4$^{-/-}$ / (-), n = 7 slices) or presence of DPCPX (AQP4$^{-/-}$ / DPCPX, n = 8 slices). NS, not significant, t-test.

Fig. 9 Dual regulation of dopamine release in the mPFC by adenosine tone.

Evoked-dopamine releases from mPFC slices were measured by D_2-CHO (mPFC / D_2-CHO). A. The

effect of DPCPX in the presence of a GABA receptor antagonist picrotoxin, PTX. Slices from wild type mice (WT) were electrically stimulated (30 Hz, 40 pulses) in the presence of 100 µM PTX and DPCPX (WT / PTX / DPCPX, n = 6) or PTX alone (WT / PTX / (-), n = 6 slices). B. The effect of Group II mGluR antagonist LY341495 (LY). Evoked-dopamine release int the presence of 500 nM LY and DPCPX (WT / LY / DPCPX, n = 8 slices) or LY alone (WT / LY / (-), n = 8 slices). C. The effect of ion channel type glutamate receptor antagonists, CNQX and APV. Evoked-dopamine releases in the presence of 10 µM CNQX, 50 µM APV and DPCPX (WT / CNQX+APV / DPCPX, n = 9 slices) or CNQX+APV alone (WT / CNQX+APV / (-), n = 10 slices). D. The effect of adenosine on cocaine-treated mice. Evoked-dopamine releases in slices from cocaine-treated wild type mice in the absence (WT / Cocaine / (-), n = 9 slices) of presence of adenosine (WT / Cocaine / Ado, n = 9 slices). NS, not significant, t-test