# Thermoregulation of adult alpine butterflies: microhabitat choice is more important than physical differences

**Irena Kleckova[1], Jan Okrouhlik[2], Tomas Svozil[2], Pavel Matos Maravi[3], Jan Klecka[1]**

[1] Laboratory of Integrative Ecology, Institute of Entomology, Biology Centre of the Czech Academy of Sciences, Branišovská 31, 370 05 České Budějovice, Czech Republic

[2] Faculty of Science, University of South Bohemia, Branišovská 31, 370 05 České Budějovice, Czech Republic

[3] Laboratory of Molecular Ecology and Phylogenetics, Institute of Entomology, Biology Centre of the Czech Academy of Sciences, Branišovská 31, 370 05 České Budějovice, Czech Republic

Correspondence author: Irena Kleckova, e-mail: irena.slamova@gmail.com

**Abstract**

Efficient thermoregulation is crucial for animals living under changing climatic and weather conditions. We used a thermal camera to measure thermoregulation of six co-occurring species of *Erebia* (Lepidoptera: Nymphalidae) mountain butterflies in a laboratory experiment with artificial light and heating source. We studied wild individuals to test whether physical characteristics (body size, wing loading) are responsible for the interspecific differences in thermoregulation detected previously under natural conditions in European Alps. We revealed that physical characteristics had a small effect on explaining interspecific differences compared to distinct thermoregulatory differences recorded in the field. Our results show that larger butterflies, with higher weight and wing loading, heated up more slowly but reached the same asymptotic body temperature as smaller butterflies. Altogether, our results suggest that differences in body temperatures among *Erebia* species observed in the field are likely caused mainly by species-specific differences in microhabitat use and confirm the role of active thermoregulation in adult butterflies.

**Keywords**

alpine species, ecophysiology*,* habitat use, *Lepidoptera*, thermal camera, thermoregulation

## 1. Introduction

Thermal ecophysiology of butterflies is driven by interactions between the environment and species-specific physical (morphology, physiology) and behavioural characteristics. Important morphological traits for butterfly thermoregulation include body size (Heinrich, 1986; Kleckova et al., 2014), wing loading (i.e., the ratio of body weight over wing area) and colouration (Kingsolver, 1983). Behaviourally, insects also thermoregulate by specific body postures (Tsai et al., 2020) or by microhabitat choice (Kleckova & Klecka, 2016). Adult butterflies typically increase their body temperature over the ambient air temperature by basking postures (Heinrich, 1995; Kingsolver, 1985a) and by conduction from warm stones or soil (Kemp & Krockenberger, 2002; Kingsolver, 1985b; Kleckova et al., 2014). Conversely, body temperature can be decreased by a shift to a cooler microhabitat (Dreisig, 1995; Shreeve, 1984; Wickman, 1988). Hence, thermoregulation in butterflies is likely driven by a combination of behaviour, morphological and physiological traits.

Our knowledge about the interactions between behavioural and physiological responses to climate change is generally scarce in insects (Abram et al., 2017) because behavioural and physiological responses are difficult to disentangle by field observations (e.g. Lawson,et al., 2014a). However, it has been established that adult butterflies respond to short-term environmental changes by active thermoregulation based on their species-specific physiological optima (Bladon et al., 2020; Kleckova et al., 2014; Kleckova & Klecka, 2016). Such physiological optima are, in turn, also affected by morphological traits, which affect the choice of ambient temperature and daily activity patterns of different species of butterflies (Kingsolver, 1983) and other insects, such as beetles (Gallego et al., 2018). Similar variation has been observed also at the inter-population level in butterflies (Nève & Després, 2020) and intra-population level in grasshoppers (Forsman, 2000). Moreover, it has been shown that thermoregulatory behaviour of butterflies prevents overheating of living cells distributed in the wings veins (Tsai et al., 2020). Such studies of interactions between behavioural responses, morphological and physiological traits have provided novel insights into the importance of insect thermoregulatory strategies. However, we still know little about the main traits involved in thermoregulation of butterflies occurring in extreme environments such as alpine regions.

To respond to local climatic conditions, butterfly species may evolve different physiological optima (MacLean et al., 2016), disperse to optimal climates (Konvicka et al., 2003) or change their thermoregulatory behaviour, e.g., by shifts of microhabitat use (Kirkpatrick & Sheldon, 2022; Wilson et al., 2015). Species-specific physiological optima evolved under the selection pressures exerted by the environment during the whole life cycle (Radchuk et al., 2013; Willmer, 1982), forming the species' climatic niches (Klečková et al., 2022; Pitteloud et al., 2017). Ultimately, strong selection pressures of local conditions may result in local adaptation of adult thermoregulatory traits across climatic gradients, as has been reported in *Melitaea cinxia* butterflies (Advani et al., 2019; Mattila, 2015). Still, the importance of active behaviour, physical constraints and possible adaptations to local conditions are not fully resolved.

To disentangle the roles of behavioural responses and physical differences in thermoregulation,
we studied adult *Erebia* (Nymphalidae, Satyrinae) butterflies occurring in sympatry in alpine
environments. In the mountains, microclimatic conditions experienced by butterflies are
changing rapidly (Stuhldreher & Fartmann, 2018) and alpine species including *Erebia* are
endangered by climate warming (Konvicka et al., 2021; Sistri et al., 2022). *Erebia* butterflies
have low mobility (Polic et al., 2014; Slamova et al., 2013), which probably leads to slower
uphill shifts in response to climate warming compared to mobile generalist species (Rödder et
al., 2021). On the other hand, at least some populations of high-altitude *Erebia* specialists are
able to use suitable habitats at altitudes below their climatic optimum (Cizek et al., 2003). Our
previous work suggests that inter-specific differences in thermoregulation of co-occurring
closely related *Erebia* species might have been driven mainly by microhabitat use and body
size (Kleckova et al., 2014), but it is unclear what the main thermoregulation strategy is. To test
the relative effects of species identity (a proxy for possible physiological differences) and
morphology (individual weight and wing-loading), we carried out laboratory measurements of
body temperature heating rates under an artificial source of radiation. This allows a better
comparison of the importance of morphological traits and microhabitat choice. Our hypothesis
is that different thermal conditions along the elevational gradient and different microhabitats
inhabited by individual species have driven divergent physiological constraints. We expected
that larger individuals and individuals with higher wing loading would heat up more slowly
than smaller individuals, as detected in *Thymelicus lineola* (Lepidoptera: Hesperiidae) (Pivnick
& McNeil, 1986). Finally, we asked whether microhabitat choice is the primary
thermoregulatory mechanism by comparing data from the laboratory heating experiments to
field measurements of body temperatures (Kleckova et al., 2014).
102
103 **2.  Material and methods**
104
105 2.1. Study site and species
106
We collected adult butterflies of six species of *Erebia* in the Austrian Alps, close to the town
of Sölden, Tirol (1900-2020 m a.s.l., 46°57'37"N 11°01'49"E) and in Hochgurgl (2150 m a.s.l.,
46°54'6"N, 11°03'14"E, 7 km away). The butterflies have brown wings with orange spots.
Species differ in their habitat preferences, altitude of occurrence, body size, and phenology.
Rocky habitats above the treeline host *E. pandrose* (Borkhausen, 1788) and *E. cassioides*
(Reiner & Hochenwarth, 1792). Sub-alpine and alpine grassy meadows above the treeline are
inhabited by *E. alberganus* (de Prunner, 1798) and *E. melampus* (Fuessly, 1775). Subalpine
meadows around the treeline and woodland clearings below the treeline are inhabited by *E.*
*euryale* (Esper, 1805) and *E. ligea* (Linnaeus, 1758). Regarding size differences, *E. cassioides*
and *E. melampus* are small (forewing length less than 20 mm), whereas the other four species
are large (forewing length above 20 mm). Three early flying species, *E. alberganus*, *E.*
*melampus* and *E. pandrose,* were caught between 27th and 29th June 2011. Three late flying
species, *E. ligea, E. cassioides* and *E. euryale,* were caught between 2nd and 4th August 2011.
The butterflies were collected by hand netting and stored alive in envelopes inside a cooling
box with suitable humidity provided by wet tissues. We transported the specimens to the

facilities of the Faculty of Science, University of South Bohemia in České Budějovice, Czech
Republic, for laboratory experiments.
124
125 2.2. Laboratory measurements of body heating
126
Before the experiments, the butterflies were acclimatised overnight, for at least 15 hours, in a
net cage in a climate-controlled room (20°C) with 16 h day and 8 h night photoperiod.
Butterflies were fed ad libitum by 20% sugar solution applied on the net. We measured the
heating rate using a thermal camera (FLIR P660, Flir Systems) in a climate-controlled room.
The thermal camera was mounted on a tripod 0.60 m above a flat wooden arena where the
butterfly was placed. We used wood as material for the arena because it kept a relatively stable
surface temperature during the experiments. On the camera, we set-up the emissivity to 0.85,
humidity to 66 %, and object distance to 0.6 m for all individual specimens. The emissivity
value compensates for the reflection, transmission, and absorption of IR energy and
corresponds to a conservative estimate for insect cuticle, e.g. (Verdu et al., 2012). The humidity
value corresponds to the average air humidity in the experimental room (range 56-82%).
138
The acclimatised butterflies were first cooled at 5 °C for 10 minutes. Then, using cold forceps,
they were placed in the middle of the experimental arena under the two halogen light bulbs (18
W and 160 lm) situated 25 cm above the arena. The butterfly surface body temperature was
recorded every 20 seconds for 10 minutes (Figure 1). All individuals were sitting during the
experiment. Each individual was measured only once. The temperature in the middle of the
mesothorax was recorded using FLIR QuickReport©1.2 software. The air-conditioning system
in the experimental room created a strong draft, so we kept it turned off for the duration of the
measurements. Consequently, the temperature in the experimental room fluctuated from 17.5
to 21.1 °C. To account for these fluctuations in the statistical analysis, room temperature ($T_{room}$)
and *humidity* was measured just before each trial.
149
After the thermal camera measurements, the individual butterflies were sacrificed using ethyl-
acetate. We obtained the weight (mg) of fresh individuals using a laboratory balance (precision
0.0001 g). We also photographed the left side of wings of every individual and calculated the
wing area in the software ImageJ v.1.46 (Rasband, 1997) and multiplied by two to obtain an
estimate of the total wing area. We calculated wing loading (mg/cm$^2$) of every individual as the
ratio of body weight (mg) over wing area (cm$^2$). In total, we measured the heating rate in 78
individuals (Table 1).
157
158 2.3 Statistical analyses
159
160 *Heating curves and species effects*
161
162 All analyses and data visualisation were done in R version 4.2.0 (R Core Team, 2022). First,
163 we analysed heating curves, describing the increase of body temperature $T_b$ in time during
164 basking. The curves were analysed using nonlinear mixed effects models (NLME) with

165 Gaussian error distribution in the nlme package for R (Pinheiro et al., 2022). The heating curve
166 was estimated by two alternative models:

    1. following the modified asymptotic exponential equation $T_b = T_s(1-e^{-b\,(time-c)})$, where $T_s$
       is the estimated asymptotic temperature reached by an individual. The slope of the
       asymptotic exponential curve ($b$) describes the heating rate of the body temperature
       increase in time until it reaches asymptotic body temperature $T_s$.

    2. following the modified Michaelis-Menten equation $T_b = T_s((time-c)/(time+K-c))$, where
       the parameter $T_s$ is the same as above and $K$ is analogous to the Michaelis constant in
       classic Michaelis-Menten equation, but here $K+c$ is the time when $T_b$ reaches 50 % of
       the final $T_s$.

The parameter $c$ is included in both models to account for the fact that the initial $T_b$ at $time=0$
was >0 °C (Figure 2). Specifically, $c$ is the estimate of how long it would take for the butterfly
to increase its body temperature from $T_b=0$ °C to the real initial $T_b$ (which varied among
individuals) under the experimental conditions.

We compared the fit of the two models without any random effects and with two alternative
random effect structures: the random effect of species and the random effect of individual
nested within species. The individual identity was used to account for possible differences of
the heating rate ($b$) and asymptotic body temperature ($T_s$) in different individuals. The model
fits were compared by the Akaike information criterion (AIC).

*Body weight and wing loading effects*

We tested how the shape of the heating curves depends on individual body mass and wing
loading. We used generalised linear mixed effect models (GLMM) in the lme4 (Bates et al.,
2015) and lmerTest libraries for R (Kuznetsova et al., 2017). For every individual, we extracted
the estimates of parameters of asymptotic temperature $T_s$ and heating rate $b$ from the best fitting
NLME model (i.e., modified asymptotic exponential, see Table 2). Then, we correlated the
dependence of individual-level $T_s$ or $b$ on individual body weight and wing-loading. As
additional predictors in the model, we included the $T_{room}$ temperature and *humidity* in the
experimental room recorded during individual measurements. We also compared mean body
weight and wing-loading among species using generalised linear models (GLM) with Gamma
distribution and log link function.

*Thermoregulation in the laboratory vs. in the field*

We intended to compare species-specific values of the asymptotic temperature ($T_s$) reached in
laboratory conditions with body temperatures ($T_{b\_field}$) reached in the field (Kleckova et al.,
2014). Because the range of ambient temperatures experienced in the laboratory differed from
the range in the field, we estimated the standardised asymptotic body temperature $T_s$ and the
standardised body temperature in the field $T_{b\_field}$. The standardised $T_s$ and $T_{b\_field}$ temperatures

were estimated for 20 °C of ambient laboratory ($T_{room}$) and for 20 °C of the two field
temperatures, $T_{air}$ (air temperature at breast height) and $T_{microhabitat}$ (microhabitat temperature)
experienced by individual butterflies in the spot where an individual was sitting (see Kleckova
et al., 2014). For the laboratory measurements, the estimate of $T_s$ at ambient temperature 20°C
was calculated from a generalised linear model $T_s \sim T_{room} + species$ fitted to $T_s$ estimates for all
individuals (see above). We used the *predict()* function in R to obtain the estimate of $T_s$ at
ambient temperature 20°C for each species. For the field observations, we again used the
*predict()* function to infer $T_{b\_field}$ at ambient temperature 20 °C for each species. The dependence
of body temperature $T_{b\_field}$ on both the ambient air ($T_{a\_field}$) and the microhabitat ($T_{m\_field}$)
temperature was obtained from a previous study (Kleckova et al., 2014). Finally, we tested the
relationship between the standardised $T_s$ in the laboratory and the standardised $T_{b\_field}$ reached
in the field for individual species by fitting a generalised linear model with Gaussian error
distribution.
222
223    **3.  Results**
224
225    3.1. Heating curves and species effect
226
The increase of body temperature $T_b$ in the studied *Erebia* butterflies was better described by
the modified asymptotic exponential model (AIC = 583.8), which includes the heating rate
parameter $b$, compared to the modified Michaelis-Menten model (AIC = 1051.3) (Table 2). The
modified asymptotic exponential model predicted that the asymptotic temperature $T_s$ is slightly
more rapidly reached compared to the modified Michaelis-Menten model (Figure 2 and Table
2). The variability of the individual heating curves was best explained by a model including the
random effect of individual identity nested within species (Table 2). However, differences in
the shape of the heating curves among species were modest (Figure 3).
235
236    3.2. Body weight and wing loading effects
237
The six species differed in their body weight and wing loading (Table 1, GLM, $P < 10^{-8}$ in both
cases), although there was considerable intraspecific variation. As expected, wing loading and
body weight were strongly positively correlated (Pearson's correlation coefficient = 0.84, $t_{46}$ =
10.373, $P < 10^{-12}$). Thermoregulation at the individual level was affected by both measures of
body size. The heating rate $b$ decreased with increasing body mass (GLMM, F = 7.17, P =
0.0105) and wing loading (GLMM, F = 4.40, P = 0.0419) (Figure 4). On the other hand, the
asymptotic temperature $T_s$ was independent of body mass (GLMM, F = 0.05, P = 0.8209) and
wing loading (GLMM, F = 0.16, P = 0.6877). The results, thus, demonstrated that larger
butterflies heated up more slowly but reached the same asymptotic temperature as smaller
butterflies. The GLMMs also included the effect of air temperature and humidity measured at
the time of each observation. While $T_s$ increased with the increasing room temperature (F =
46.53, P= <0.0001) and humidity (F = 5.12, P = 0.0294), the heating rate increased with the
increasing room temperature (F = 13.82, P = 0.0011) and was not significantly affected by air
humidity (F = 3.68, P = 0.0612).
252

3.3. Thermoregulation in the laboratory vs. in the field

The standardised asymptotic temperatures $T_s$ measured in the laboratory (Figure 5) were not correlated with the standardised body temperature reached in field ($T_{b\_field}$). There was no statistically significant relationship regardless of whether $T_{b\_field}$ was standardised for air temperature $T_{a\_field}$ (GLM, F = 0.41, P = 0.5580) or for microhabitat temperature $T_{m\_field}$ (F = 0.84, P = 0.4107). It is noteworthy that the two species preferring rocky habitats, *E. cassioides* and *E. pandrose*, kept the highest body temperatures in the field despite their intermediate asymptotic temperatures measured in the laboratory (Figure 5). Also, the subalpine species *E. alberganus* and *E. melampus* kept higher body temperatures than the woodland species *E. euryale* and *E. ligea* measured both in the field and in the laboratory.

## 4. Discussion

4.1. Species and individual effects on thermoregulation under laboratory conditions

The species-specific thermoregulation differences in laboratory conditions were modest (Figure 3), but still detectable (Table 2). The better fit of the individual heating curves by models including the random effect of individual nested in species (Table 2) indicates that there is a noticeable effect of individual-level traits (e.g., species-specific physiological responses) in thermoregulation. However, the observed higher heating rate $b$ in individuals with lower body weight and wing-loading regardless of species identity (Figure 4) points toward a major role of morphological traits. In other words, larger butterflies having higher wing loading heated up more slowly than small species. Our results are in line with (Pivnick & McNeil, 1986), who observed that males of *Thymelicus lineola* (Lepidoptera: Hesperiidae) heated up faster than females, the latter having a higher wing-loading.

In the field, we previously detected a pattern with smaller species reaching higher body temperatures at low ambient temperatures compared to larger species (Kleckova et al., 2014). Thus, our original expectation was that larger species and individuals would accumulate higher body temperatures in the laboratory, as it was also recorded in the butterfly species *Polyommatus icarus* (Lepidoptera: Lycaenidae) (Keyser et al., 2015). Our results, nonetheless, did not support such an expectation in *Erebia*, as we found no relationship between asymptotic temperatures $T_s$ and body size under laboratory conditions. That larger *Erebia* species did not reach higher body temperatures under laboratory conditions might be partly explained by the low light intensity used in our experimental setup (45 W/m$^2$, but solar flux is around 1000 W/m$^2$ (Sterhov & Loshkarev, 2019)). Future experiments should consider the importance of sufficient light intensity because larger butterflies might be able to use their wings as "solar collectors" in natural conditions, but this was not possible to test in our experiment.

Other traits not included in our study, such as fur thickness on the body, may affect heating rates and body temperatures of butterflies. In the laboratory conditions, a small-bodied species, *E. melampus*, warmed up faster than the remaining species (Fig. 3), unlike the other small species, *Erebia cassiodes.* This may be caused by the fact that *E. cassioides* has more "furry"

297 (modified scales) wings than *E. melampus*. We argue that the fur may provide better insulation
298 in cold conditions (Kingsolver, 1988), but it may also slow down heating. Low heating
299 effectiveness can be caused also by different wing emissivity, reflectance and absorbance (Lou
300 et al., 2021; Schmitz, 1994).
301
302 4.2. Habitat use
303
A previous study on thermoregulatory strategies in the field demonstrated that woodland *Erebia*
species kept lower body temperatures compared to species of open and rocky habitats
(Kleckova et al., 2014). In congruence, the smallest difference in asymptotic temperature $T_s$
reached in the laboratory and $T_{b\_field}$ under field conditions was found for the woodland species
*E. ligea*. This indicates low buffering ability of this woodland species. Similarly, small
differences in body temperatures measured in the field and in the laboratory were detected in
another woodland butterfly *Hamaeris lucina* (Riodinidae) (Bladon et al., 2020; Hayes et al.,
2019). In contrast, the greatest difference between $T_s$ and $T_{b\_field}$ was observed for species
inhabiting rocky habitats, *E. cassioides* and *E. pandrose,* indicating their strong ability to
thermoregulate by microclimate choice.
314
315 4.3. Heating curves were better described by the modified asymptotic exponential model
316
There is no consensus in the literature about the most appropriate model for fitting the heating
curves describing the increase of body temperature $T_b$ in time in insects. An exponential model
was used to describe ectotherm thermoregulation, for example, to compare thermoregulation of
colour morphs in the chameleon grasshopper (Umbers et al., 2013) and to infer the increase of
internal body temperature in dung-beetles (Amore et al., 2017). Another model used previously
is  the Michaelis-Menten equation, used e.g. to describe heating curves of a sea snail by Muñoz
et al., 2005. Complex models based on detailed physiological measurements of heat transfer
within animal bodies and between them and their environment have been developed (Dreisig,
1984; Dzialowski & O'Connor, 2001), but they are impractical for use in ecological studies
where detailed physiological parameters are difficult to obtain. In *Erebia* butterflies, the
modified asymptotic exponential model explained the heating of body temperature better than
the modified Michaelis-Menten model (Table 2).
329
330 **5. Conclusions**
331
Studying the interactions of insect physical traits and their behavioural responses to
environmental factors has the potential to contribute to understanding and predicting the
impacts of climate change on biodiversity. When behavioural thermoregulation was excluded
under laboratory conditions, six closely related *Erebia* species differed minimally in
thermoregulation, as measured by heating rate curves (Figure 3). We only detected a negative
effect of individual body size on the rate of heating, but no effect on the asymptotic body
temperate. Thus, the thermal conditions along the elevational gradient and different habitats
have not resulted in divergent physiological constraints among species as we hypothesised, at
least at the adult life stage. However, in the field, when adult butterflies have the opportunity

to actively seek out suitable microclimates, the species differed significantly in thermoregulation (Kleckova et al., 2014). This points toward a more prominent role of behavioural thermoregulation in these closely related butterflies. The key role of behaviour in thermoregulation has been shown also in a territorial butterfly (Rutowski et al., 1994) and in butterflies exposed to extreme temperatures (Tsai et al., 2020). The importance of behavioural thermoregulation found in this study suggests that the adult life stage is not under direct thermal selection induced by ongoing climate change, in agreement with (Buckley et al., 2015). Thus, under increased anthropogenic pressure on *Erebia* populations, it is necessary to safeguard a variety of microhabitats which can facilitate efficient behavioural thermoregulation of adults and increased survival of less mobile developmental stages (Abarca et al., 2019). This is in line with previous calls for management of mountain habitats supporting environmental heterogeneity (Habel et al., 2022) to increase species survival under ongoing climatic and other human-induced environmental changes (Lawson et al. 2014b).

**Acknowledgements**

We are grateful to J. Skála and M. Česánek for recommending the study site, A. Suchačková-Bartoňová for help with the thermal camera work, O. Nedvěd for providing practical advice and to Marie Hronková for borrowing of the thermal camera.

**References**

Abarca, M., Larsen, E. A., & Ries, L. (2019). Heatwaves and Novel Host Consumption Increase Overwinter Mortality of an Imperiled Wetland Butterfly. *Frontiers in Ecology and Evolution*, 7. https://www.frontiersin.org/articles/10.3389/fevo.2019.00193

Abram, P. K., Boivin, G., Moiroux, J., & Brodeur, J. (2017). Behavioural effects of temperature on ectothermic animals: Unifying thermal physiology and behavioural plasticity. *Biological Reviews of the Cambridge Philosophical Society*, *92*(4), 1859–1876. https://doi.org/10.1111/brv.12312

Advani, N. K., Parmesan, C., & Singer, M. C. (2019). Takeoff temperatures in Melitaea cinxia butterflies from latitudinal and elevational range limits: A potential adaptation to solar irradiance. *Ecological Entomology*, *44*(3), 389–396. https://doi.org/10.1111/een.12714

Amore, V., Hernández, M. I. M., Carrascal, L. M., & Lobo, J. M. (2017). Exoskeleton may influence the internal body temperatures of Neotropical dung beetles (Col. Scarabaeinae). *PeerJ*, *5*, e3349. https://doi.org/10.7717/peerj.3349

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *67*, 1–48. https://doi.org/10.18637/jss.v067.i01

Bladon, A. J., Lewis, M., Bladon, E. K., Buckton, S. J., Corbett, S., Ewing, S. R., Hayes, M. P., Hitchcock, G. E., Knock, R., Lucas, C., McVeigh, A., Menéndez, R., Walker, J. M., Fayle, T. M., & Turner, E. C. (2020). How butterflies keep their cool: Physical and ecological traits influence thermoregulatory ability and population trends. *Journal of Animal Ecology*, *89*(11), 2440–2450. https://doi.org/10.1111/1365-2656.13319

Buckley, L. B., Ehrenberger, J. C., & Angilletta Jr, M. J. (2015). Thermoregulatory behaviour
limits local adaptation of thermal niches and confers sensitivity to climate change.
*Functional Ecology*, *29*(8), 1038–1047. https://doi.org/10.1111/1365-2435.12406

Cizek, O., Bakesová, A., Kuras, T., Benes, J., & Konvicka, M. (2003). Vacant niche in alpine
habitat: The case of an introduced population of the butterfly *Erebia epiphron* in the
Krkonoše Mountains. *Acta Oecologica*, *24*(1), 15–23. https://doi.org/10.1016/S1146-
609X(02)00004-8

Dreisig, H. (1984). Control of body temperature in shuttling ectotherms. *Journal of Thermal
Biology*, *9*(4), 229–233. https://doi.org/10.1016/0306-4565(84)90001-9

Dreisig, H. (1995). Thermoregulation and flight activity in territorial male graylings,
*Hipparchia semele* (Satyridae), and large skippers, Ochlodes venata (Hesperiidae).
*Oecologia*, *101*(2), 169–176. https://doi.org/10.1007/BF00317280

Dzialowski, E. M., & O'Connor, M. P. (2001). Thermal time constant estimation in warming
and cooling ectotherms. *Journal of Thermal Biology*, *26*(3), 231–245.
https://doi.org/10.1016/s0306-4565(00)00050-4

Forsman, A. (2000). Some like it hot: Intra-Population Variation in behavioral
Thermoregulation in Color-Polymorphic pygmy Grasshoppers. *Evolutionary Ecology*,
*14*(1), 25–38. https://doi.org/10.1023/A:1011024320725

Gallego, B., Verdú, J. R., & Lobo, J. M. (2018). Comparative thermoregulation between
different species of dung beetles (Coleoptera: Geotrupinae). *Journal of Thermal
Biology*, *74*, 84–91. https://doi.org/10.1016/j.jtherbio.2018.03.009

Habel, J. C., Schmitt, T., Gros, P., & Ulrich, W. (2022). Breakpoints in butterfly decline in
Central Europe over the last century. *Science of The Total Environment*, *851*, 158315.
https://doi.org/10.1016/j.scitotenv.2022.158315

Hayes, M. P., Hitchcock, G. E., Knock, R. I., Lucas, C. B. H., & Turner, E. C. (2019).
Temperature and territoriality in the Duke of Burgundy butterfly, *Hamearis lucina*.
*Journal of Insect Conservation*, *23*(4), 739–750. https://doi.org/10.1007/s10841-019-
00166-6

Heinrich, B. (1986). Thermoregulation and flight activity satyrine, *Coenonympha inornata*
(Lepidoptera: Satyridae). *Ecology*, *67*(3), 594–597. https://doi.org/10.2307/1937682

Heinrich, B. (1995). Insect thermoregulation. *Endeavour*, *19*(1), 28–33.
https://doi.org/10.1016/0160-9327(95)98891-I

Kemp, D. J., & Krockenberger, A. K. (2002). A novel method of behavioural
thermoregulation in butterflies. *Journal of Evolutionary Biology*, *15*(6), 922–929.
https://doi.org/10.1046/j.1420-9101.2002.00470.x

Keyser, R. D., Breuker, C. J., Hails, R. S., Dennis, R. L. H., & Shreeve, T. G. (2015). Why
Small Is Beautiful: Wing Colour Is Free from Thermoregulatory Constraint in the
Small Lycaenid Butterfly, *Polyommatus icarus*. *PLOS ONE*, *10*(4), e0122623.
https://doi.org/10.1371/journal.pone.0122623

Kingsolver, J. G. (1983). Thermoregulation and Flight in *Colias* Butterflies: Elevational
Patterns and Mechanistic Limitations. *Ecology*, *64*(3), 534–545.
https://doi.org/10.2307/1939973

Kingsolver, J. G. (1985a). Thermoregulatory significance of wing melanization in *Pieris*
butterflies (Lepidoptera: Pieridae): physics, posture, and pattern. *Oecologia*, *66*(4),
546–553. https://doi.org/10.1007/BF00379348

Kingsolver, J. G. (1985b). Thermoregulatory significance of wing melanization in *Pieris*
butterflies (Lepidoptera: Pieridae): physics, posture, and pattern. *Oecologia*, *66*(4),
546–553. https://doi.org/10.1007/BF00379348

Kingsolver, J. G. (1988). Thermoregulation, Flight, and the Evolution of Wing Pattern in
Pierid Butterflies: The Topography of Adaptive Landscapes. *American Zoologist*,
*28*(3), 899–912. https://doi.org/10.1093/icb/28.3.899

Kirkpatrick, W. H., & Sheldon, K. S. (2022). Experimental increases in temperature mean and
variance alter reproductive behaviours in the dung beetle *Phanaeus vindex*. *Biology*
*Letters*, *18*(7), 20220109. https://doi.org/10.1098/rsbl.2022.0109

Kleckova, I., & Klecka, J. (2016). Facing the Heat: Thermoregulation and Behaviour of
Lowland Species of a Cold-Dwelling Butterfly Genus, *Erebia*. *PLOS ONE*, *11*(3),
e0150393. https://doi.org/10.1371/journal.pone.0150393

Klečková, I., Klečka, J., Fric, Z. F., Česánek, M., Dutoit, L., Pellissier, L., & Matos-Maraví,
P. (2022). Climatic niche conservatism and ecological diversification in the Holarctic
cold-dwelling butterfly genus *Erebia* (p. 2022.04.12.488065). bioRxiv.
https://doi.org/10.1101/2022.04.12.488065

Kleckova, I., Konvicka, M., & Klecka, J. (2014). Thermoregulation and microhabitat use in
mountain butterflies of the genus *Erebia*: Importance of fine-scale habitat
heterogeneity. *Journal of Thermal Biology*, *41*, 50–58.
https://doi.org/10.1016/j.jtherbio.2014.02.002

Konvicka, M., Kuras, T., Liparova, J., Slezak, V., Horázná, D., Klečka, J., & Kleckova, I.
(2021). Low winter precipitation, but not warm autumns and springs, threatens
mountain butterflies in middle-high mountains. *PeerJ*, *9*, e12021.
https://doi.org/10.7717/peerj.12021

Konvicka, M., Maradova, M., Benes, J., Fric, Z., & Kepka, P. (2003). Uphill shifts in
distribution of butterflies in the Czech Republic: Effects of changing climate detected
on a regional scale. *Global Ecology and Biogeography*, *12*(5), 403–410.
https://doi.org/10.1046/j.1466-822X.2003.00053.x

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in
Linear Mixed Effects Models. *Journal of Statistical Software*, *82*, 1–26.
https://doi.org/10.18637/jss.v082.i13

Lawson, C. R., Bennie, J., Hodgson, J. A., Thomas, C. D., & Wilson, R. J. (2014).
Topographic microclimates drive microhabitat associations at the range margin of a
butterfly. *Ecography*, *37*(8), 732–740. https://doi.org/10.1111/ecog.00535

Lawson, C. R., Bennie, J. J., Thomas, C. D., Hodgson, J. A., & Wilson, R. J. (2014). Active
Management of protected areas enhances metapopulation expansion under climate
change. *Conservation Letters*, *7*(2), 111–118. https://doi.org/10.1111/conl.12036

Lou, C., An, S., Yang, R., Zhu, H., Shen, Q., Jiang, M., Fu, B., Tao, P., Song, C., Deng, T., &
Shang, W. (2021). Enhancement of infrared emissivity by the hierarchical
microstructures from the wing scales of butterfly *Rapala dioetas*. *APL Photonics*, *6*(3),
036101. https://doi.org/10.1063/5.0039079

MacLean, H. J., Higgins, J. K., Buckley, L. B., & Kingsolver, J. G. (2016). Morphological and physiological determinants of local adaptation to climate in Rocky Mountain butterflies. *Conservation Physiology*, *4*(1), cow035. https://doi.org/10.1093/conphys/cow035

Mattila, A. L. K. (2015). Thermal biology of flight in a butterfly: Genotype, flight metabolism, and environmental conditions. *Ecology and Evolution*, *5*(23), 5539–5551. https://doi.org/10.1002/ece3.1758

Muñoz, J. L. P., Randall Finke, G., Camus, P. A., & Bozinovic, F. (2005). Thermoregulatory behavior, heat gain and thermal tolerance in the periwinkle *Echinolittorina peruviana* in central Chile. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, *142*(1), 92–98. https://doi.org/10.1016/j.cbpa.2005.08.002

Nève, G., & Després, L. (2020). Cold adaptation across the elevation gradient in an alpine butterfly species complex. *Ecological Entomology*, *45*(5), 997–1003. https://doi.org/10.1111/een.12875

Pinheiro, J., Bates, D., & R Core Team. (2022). Nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1-157, *https://CRAN.R-project.org/package=nlme*.

Pitteloud, C., Arrigo, N., Suchan, T., Mastretta-Yanes, A., Vila, R., Dincă, V., Hernández-Roldán, J., Brockmann, E., Chittaro, Y., Kleckova, I., Fumagalli, L., Buerki, S., Pellissier, L., & Alvarez, N. (2017). Climatic niche evolution is faster in sympatric than allopatric lineages of the butterfly genus *Pyrgus*. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1852), 20170208. https://doi.org/10.1098/rspb.2017.0208

Pivnick, K. A., & McNeil, J. N. (1986). Sexual Differences in the Thermoregulation of *Thymelicus lineola* Adults (Lepidoptera: Hesperiidae). *Ecology*, *67*(4), 1024–1035. https://doi.org/10.2307/1939825

Polic, D., Fiedler, K., Nell, C., & Grill, A. (2014). Mobility of ringlet butterflies in high-elevation alpine grassland: Effects of habitat barriers, resources and age. *Journal of Insect Conservation*, *18*(6), 1153–1161. https://doi.org/10.1007/s10841-014-9726-5

R Core Team. (2022). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. *Https://www.R-project.org/*.

Radchuk, V., Turlure, C., & Schtickzelle, N. (2013). Each life stage matters: The importance of assessing the response to climate change over the complete life cycle in butterflies. *Journal of Animal Ecology*, *82*(1), 275–285. https://doi.org/10.1111/j.1365-2656.2012.02029.x

Rasband, W. S. (1997). ImageJ, U. S. National Institutes of Health, Bethesda, Maryland, USA (v.1.46). https://imagej.nih.gov/ij/

Rödder, D., Schmitt, T., Gros, P., Ulrich, W., & Habel, J. C. (2021). Climate change drives mountain butterflies towards the summits. *Scientific Reports*, *11*(1), 14382. https://doi.org/10.1038/s41598-021-93826-0

Rutowski, R. L., Demlong, M. J., & Leffingwell, T. (1994). Behavioural thermoregulation at mate encounter sites by male butterflies (*Asterocampa leilia*, Nymphalidae). *Animal Behaviour*, *48*(4), 833–841. https://doi.org/10.1006/anbe.1994.1307

Schmitz, H. (1994). Thermal characterization of butterfly wings—1. Absorption in relation to
different color, surface structure and basking type. *Journal of Thermal Biology*, *19*(6),
403–412. https://doi.org/10.1016/0306-4565(94)90039-6
Shreeve, T. G. (1984). Habitat selection, mate location, and microclimatic constraints on the
activity of the speckled wood butterfly *Pararge aegeria*. *Oikos*, *42*(3), 371–377.
https://doi.org/10.2307/3544407
Sistri, G., Menchetti, M., Santini, L., Pasquali, L., Sapienti, S., Cini, A., Platania, L., Balletto,
E., Barbero, F., Bonelli, S., Casacci, L. P., Dincă, V., Vila, R., Mantoni, C., Fattorini,
S., & Dapporto, L. (2022). The isolated *Erebia pandrose* Apennine population is
genetically unique and endangered by climate change. *Insect Conservation and
Diversity*, *15*(1), 136–148. https://doi.org/10.1111/icad.12538
Slamova, I., Klecka, J., & Konvicka, M. (2013). Woodland and grassland mosaic from a
butterfly perspective: Habitat use by *Erebia aethiops* (Lepidoptera: Satyridae). *Insect
Conservation and Diversity*, *6*(3), 243–254. https://doi.org/10.1111/j.1752-
4598.2012.00212.x
Sterhov, A. I., & Loshkarev, I. Y. (2019). Determination of the proportion of natural light in
solar radiation using the method of conversion of lighting units into energy. *Journal of
Physics: Conference Series*, *1353*(1), 012002. https://doi.org/10.1088/1742-
6596/1353/1/012002
Stuhldreher, G., & Fartmann, T. (2018). Threatened grassland butterflies as indicators of
microclimatic niches along an elevational gradient – Implications for conservation in
times of climate change. *Ecological Indicators*, *94*, 83–98.
https://doi.org/10.1016/j.ecolind.2018.06.043
Tsai, C.-C., Childers, R. A., Nan Shi, N., Ren, C., Pelaez, J. N., Bernard, G. D., Pierce, N. E.,
& Yu, N. (2020). Physical and behavioral adaptations to prevent overheating of the
living wings of butterflies. *Nature Communications*, *11*(1), Article 1.
https://doi.org/10.1038/s41467-020-14408-8
Umbers, K. D. L., Herberstein, M. E., & Madin, J. S. (2013). Colour in insect
thermoregulation: Empirical and theoretical tests in the colour-changing grasshopper,
*Kosciuscola tristis*. *Journal of Insect Physiology*, *59*(1), 81–90.
https://doi.org/10.1016/j.jinsphys.2012.10.016
Verdu, J. R., Alba-Tercedor, J., & Jimenez-Manrique, M. (2012). Evidence of different
thermoregulatory mechanisms between two sympatric Scarabaeus species using
infrared thermography and micro-computer tomography. *PLoS ONE*, *7*(3), e3391.
Wickman, P.-O. (1988). Dynamics of mate-searching behaviour in a hilltopping butterfly,
*Lasiommata megera* (L.): The effects of weather and male density. *Zoological Journal
of the Linnean Society*, *93*(4), 357–377. https://doi.org/10.1111/j.1096-
3642.1988.tb01367.x
Willmer, P. G. (1982). Microclimate and the environmental physiology of insects. In M. J.
Berridge, J. E. Treherne, & V. B. Wigglesworth (Eds.), *Advances in Insect Physiology*
16, 1–57. Academic Press. https://doi.org/10.1016/S0065-2806(08)60151-4
Wilson, R. J., Bennie, J., Lawson, C. R., Pearson, D., Ortúzar-Ugarte, G., & Gutiérrez, D.
(2015). Population turnover, habitat use and microclimate at the contracting range

margin of a butterfly. *Journal of Insect Conservation*, *19*(2), 205–216.

https://doi.org/10.1007/s10841-014-9710-0

**Tables**

**Table 1. The list of study species of the butterfly genus *Erebia*.** For each species, we
provide information about its typical habitat, the number of individuals used for the lab
measurements (N), their mean body weight, total wing area, and wing loading (body
weight/wing area). Species are sorted according to their body weight from lightest to heaviest.

| Species | Habitat | N | Body weight (mg) mean (SD) | Wing area (cm2) mean (SD) | Wing loading (mg/cm2) mean (SD) |
|---|---|---|---|---|---|
| *E. melampus* | Grassland | 11 | 24.4 (4.17) | 7.82 (0.544) | 3.12 (0.402) |
| *E. cassioides* | Rocks | 8 | 34.4 (7.01) | 10.95 (1.019) | 3.15 (0.682) |
| *E. euryale* | Woodland | 10 | 38.5 (5.02) | 15.45 (1.349) | 2.50 (0.359) |
| *E. ligea* | Woodland | 9 | 50.1 (21.76) | 17.94 (1.222) | 2.74 (0.949) |
| *E. alberganus* | Grassland | 9 | 54.0 (11.55) | 12.90 (1.398) | 4.16 (0.636) |
| *E. pandrose* | Rocks | 11 | 68 (14.51) | 13.58 (1.385) | 5.08 (1.286) |

567
568

**Table 2. The comparison of the fit of the modified asymptotic exponential and Michaelis-Menten models describing butterfly heating under laboratory conditions.** The models describe the increase of body temperature ($T_b$) in *time* in 78 individuals of *Erebia* butterflies, which heated up under an artificial source of heating and radiation. Their $T_b$ was measured by thermal camera. The non-linear mixed effect models included the random effects of individual nested in species and the random effect of species. The best model is highlighted **in bold**. Parameters describe maximum body temperature reached during heating (asymptotic temperature $T_s$), heating rate ($b$), Michaelis-Menten constant ($K$), and the estimated time it would take for the butterfly to increase its body temperature from $T_b$=0 °C to the actual initial $T_b$ (c).

| | | Parameter estimates (fixed effects) | | | | | |
|---|---|---|---|---|---|---|---|
| **Model** | **Random effects** | $T_s$ **(SE)** | $b$ **(SE)** | $K$ **(SE)** | $c$ **(SE)** | **AIC** | **ΔAIC** |
| **modified asymptotic exponential** | **Individual nested in species** | **21.00 (0.271)** | **0.687 (0.056)** | | **-1.188 (0.152)** | **583.8** | **0** |
| | Species | 20.93 (0.291) | 0.695 (0.061) | | -1.138 (0.141) | 2375.5 | 1791.7 |
| | None | 20.94 (0.037) | 0.660 (0.013) | | -1.140 (0.033) | 3410 | 2826.2 |
| modified Michaelis-Menten | Individual nested in species | 22.73 (0.334) | | 0.637 (0.073) | -0.559 (0.082) | 1051.3 | 467.5 |
| | Species | 22.46 (0.377) | | 0.587 (0.077) | -0.514 (0.078) | 2485.1 | 1901.3 |
| | None | 22.48 (0.068) | | 0.579 (0.018) | -0.507 (0.020) | 3422.2 | 2838.4 |

582 **Figures**

583
584 **Figure 1.** An example image from the thermal camera displaying the heating of an individual
585 *Erebia* butterfly in *time* (s), blue indicates colder, red warmer temperatures.
586

588
**Figure 2.** The comparison of the fit of the modified asymptotic exponential and Michaelis-
Menten models describing the heating of individuals of *Erebia* butterflies under laboratory
conditions. Data from all individuals of all species are pooled in the plot (grey dots represent
measurements of individual body temperature $T_b$ in *time*). The modified asymptotic
exponential model fits the temporal changes of body temperature during heating of the
butterflies better than the modified Michaelis-Menten model ($\Delta AIC = 467.5$, see Table 2 for
details).

 **Figure 3.** Heating of six *Erebia* butterfly species under artificial source of heat and radiation
under laboratory conditions. The lines display the fit of the modified asymptotic exponential
model. The species-specific parameters of the model were estimated by a nonlinear mixed
effects model with individual and species identity included as nested random effects. The
parameter *b* describes the heating rate, i.e. the increase of the body temperature ($T_b$) in *time*
until it reaches the asymptotic body temperature ($T_s$). The parameter *c* is the estimate of how
long it would take to increase the body temperature from $T_b=0$ °C to the real initial $T_b$.
Equations describing the fitted curves with species-specific parameter estimates are added.

606
**Figure 4.** The effect of body weight and wing loading on the asymptotic body temperature
($T_s$) and on the heating rate ($b$) of individuals of six *Erebia* species. The $T_s$ and $b$ parameters
describing heating of individuals under laboratory conditions were inferred from the modified
asymptotic exponential models (see Figure 3). The y-axis shows partial residuals from a
model which also included the room temperature and humidity at the time of each
measurement. Each dot represents one individual, grouped into species according to colour.
613
614

616
617 **Figure 5.** The correlation between *Erebia* butterfly species body temperatures reached under
618 laboratory conditions ($T_{b\_laboratory}$) and mean body temperatures $T_{b\_field}$ measured in the field at
619 20°C of ambient air (panel a.) and microhabitat (panel b.) temperature. See the Methods
620 section for more details. Estimates of $T_b$ values are accompanied by standard errors.