1

Body size, not species identity, drives body heating in

alpine *Erebia* butterflies

Irena Kleckova[1], Jan Okrouhlik[2], Tomas Svozil[2], Pável Matos-Maraví[1], Jan Klecka[1]

[1] Institute of Entomology, Biology Centre of the Czech Academy of Sciences, Branišovská

31, 370 05 České Budějovice, Czech Republic

[2] Faculty of Science, University of South Bohemia, Branišovská 1760, 370 05 České

Budějovice, Czech Republic

Correspondence author: Irena Kleckova, e-mail: irena.slamova@gmail.com

Abstract

Efficient thermoregulation is crucial for animals living under fluctuating climatic and weather

conditions. We studied the body heating of six butterfly species of the genus *Erebia*

(Lepidoptera: Nymphalidae) that co-occur in the European Alps. We tested whether butterfly

physical characteristics (body size, wing loading) are responsible for the inter-specific

differences in body temperatures recorded previously under natural conditions. We used a

thermal camera to measure body heating of wild butterfly individuals in a laboratory experiment

with artificial light and heating sources. We revealed that physical characteristics had a small

effect on explaining inter-specific differences in mean body temperatures recorded in the field.

Our results show that larger butterflies, with higher weight and wing loading, heated up more

slowly but reached the same asymptotic body temperature as smaller butterflies. Altogether,

our results suggest that differences in body temperatures among *Erebia* species observed in the

field might be caused mainly by species-specific microhabitat use and point towards an

important role of active behavioural thermoregulation in adult butterflies. We speculate that

microclimate heterogeneity in mountain habitats facilitates behavioural thermoregulation of

adults. Similarly, microclimate structuring might also increase survival of less mobile butterfly

life stages, i.e., eggs, larvae and pupae. Thus, landscape heterogeneity in management practices

may facilitate long term survival of montane invertebrates under increased anthropogenic

pressures.

Keywords

alpine species, ecophysiology, habitat use, *Lepidoptera*, thermal camera

1. Introduction

Thermal ecophysiology of butterflies is driven by interactions between the environment and

species-specific physical (morphology, physiology) and behavioural thermoregulation.

Important morphological traits affecting butterfly body temperatures include body size

(Heinrich, 1986; Kleckova et al., 2014), wing loading (i.e., the ratio of body weight over wing

area) and colouration (Kingsolver, 1983). Behaviourally, insects thermoregulate by specific

body postures (Tsai et al., 2020) or by microhabitat choice (Kleckova & Klecka, 2016). Adult

butterflies increase their body temperature over the ambient air temperature by basking postures

(Heinrich, 1995; Kingsolver, 1985a) and by conduction from warm stones or soil (Kemp &

Krockenberger, 2002; Kingsolver, 1985b; Kleckova et al., 2014). Conversely, body

temperature can be decreased by a shift to a cooler microhabitat (Dreisig, 1995; Shreeve, 1984;

Wickman, 1988). Hence, body temperatures in wild butterflies are likely driven by the

interaction of morphological and physical traits with active behavioural thermoregulation.

However, our knowledge about the interactions between behavioural thermoregulation and

physical differences is scarce (Abram et al., 2017) because they are difficult to disentangle by

field observations (e.g. Lawson,et al., 2014a). Adult butterflies likely respond to short-term

environmental changes by active thermoregulation based on their species-specific physiological

optima (Bladon et al., 2020; Kleckova et al., 2014; Kleckova & Klecka, 2016). The

physiological optima are, in turn, also affected by the active choice of ambient temperature and

daily activity patterns of different species of butterflies (Kingsolver, 1983) and other insects,

such as beetles (Gallego et al., 2018). The interplay among behaviour, morphology and

physiology on thermoregulatory strategies has been observed also at the inter-population level

in butterflies (Nève & Després, 2020) and intra-population level in grasshoppers (Forsman,

2000). Further, thermoregulatory behaviour of butterflies is tightly connected with morphology

as behaviour prevents, for example, overheating of living cells distributed in the wings veins

(Tsai et al., 2020). We still know little about the main traits involved in thermoregulation of

butterflies occurring in extreme environments such as alpine regions.

To respond to changes of local climatic conditions in mountains (Rödder et al., 2021),

butterflies may exploit different local physiological optima (MacLean et al., 2016), might

disperse to optimal climates (Konvicka et al., 2003) or change their thermoregulatory

behaviour, e.g., by shifts in the timing of daily activities (Cros et al., 1997; Slamova et al., 2011)

and microhabitat use (Kirkpatrick & Sheldon, 2022; Wilson et al., 2015). Further, climate

change drives the timing of seasonal activities, mainly by affecting less mobile stages of

butterfly life cycle (Konvicka et al., 2021; Radchuk et al., 2013). Species, populations and even

individuals differ in their responses to ambient temperatures. Such selection pressures of local

conditions probably resulted in variable body temperatures necessary for flight initiation in

different populations of *Melitaea cinxia* butterflies (Advani et al., 2019). In addition, Matilla et

al. (2015) detected differences in body temperature in flight among genotypes and sexes of *M.*

cinxia.

To test the role of physical differences for body heating, we studied adult *Erebia* (Nymphalidae,

Satyrinae) butterflies occurring in sympatry in alpine environments. Mountains provide high

heterogeneity of thermal environments for thermoregulation of butterflies (Wickman, 2009).

However, microclimatic conditions are changing rapidly (Stuhldreher & Fartmann, 2018) and

many alpine species, including *Erebia*, are threatened by climate warming (Konvicka et al.,

2021; Sistri et al., 2022). *Erebia* butterflies have low mobility (Polic et al., 2014; Slamova et

al., 2013), which probably leads to slower uphill shifts in response to climate warming

compared to mobile generalist species (Rödder et al., 2021). On the other hand, at least some

populations of high-altitude *Erebia* specialists use suitable habitats at altitudes below their

climatic optimum (Cizek et al., 2003). Inter-specific differences in mean body temperatures of

co-occurring closely related *Erebia* species might have been driven mainly by microhabitat use

and body size (Kleckova et al., 2014), but it is unclear what the main thermoregulation strategy

is. To test the relative effects of species identity (a proxy for possible physiological differences)

and morphology (individual weight and wing-loading as drivers of passive heating), we

measured body temperature heating rates under an artificial source of radiation. Our hypothesis

was that different thermal conditions and microhabitats along elevational gradients might have

driven divergent physiological constraints and heating rates may differ among species. Further,

we expected that larger individuals and individuals with lower wing loading would heat up

more slowly than smaller individuals (Wickman, 2009), as detected in other butterflies such as

Pararge aegeria (Berwaerts et al., 2001) and *Hypolimnas bolina* (Kemp & Krockenberger,

2004). Finally, by comparing data from the laboratory heating experiments to field

measurements of body temperatures (Kleckova et al., 2014), we inferred the relative importance

of microhabitat choice in determining mean adult body temperatures.

2.　Material and methods

2.1. Study site and species

We collected adult butterflies of six species of *Erebia* in the Austrian Alps, close to the town

of Sölden, Tirol (1900-2020 m a.s.l., 46°57'37"N 11°01'49"E) and in Hochgurgl (2150 m a.s.l.,

46°54'6"N, 11°03'14"E, 7 km away). We collected 15-20 individuals of each species in the

field, a subset was used to measure heating rates in the lab (Table 1). The additional specimens

were used for trial measurements and kept as a backup for unexpected events such as escape or

mortality. We collected undamaged butterflies with minimal wing-wear and scored their sex.

Erebia species are protandric (Kuras et al., 2003), and because we were able to access the study

location for a limited period of time in the beginning of the species' flight period, we collected

mainly males. A balanced number of males and females for the experiments was available only

for *E. pandrose* (Table 1).

Species differ in their habitat preferences, altitude of occurrence, body size, and phenology.

Rocky habitats above the treeline host *E. pandrose* (Borkhausen, 1788) and *E. cassioides*

(Reiner & Hochenwarth, 1792), which increase their body temperature in warm stony

microhabitats (Kleckova et al., 2014). Sub-alpine and alpine grassy meadows above the treeline

are inhabited by *E. alberganus* (de Prunner, 1798) and *E. melampus* (Fuessly, 1775), which

typically bask on stems of grasses. Subalpine meadows around the treeline and woodland

clearings below the treeline are inhabited by *E. euryale* (Esper, 1805) and *E. ligea* (Linnaeus,

1758). These species bask on grass blades and on tree branches. All species have wings with

dark brown upper side with orange spots. *E. pandrose* and *E. cassioides* have silver underside

and higher proportion of orange colouration. The remaining four species have brownish

underside. Regarding size differences, *E. cassioides* and *E. melampus* are small (forewing

length less than 20 mm), whereas the other four species are large (forewing length above 20

mm). Three early flying species, *E. alberganus*, *E. melampus* and *E. pandrose,* were caught

between 27th and 29th June 2011. Three late flying species, *E. ligea, E. cassioides* and *E.

euryale,* were caught between 2nd and 4th August 2011. The butterflies were collected by hand

netting and stored alive in envelopes inside a cooling box with suitable humidity provided by

wet tissues (temperature range in the box was 10-15°C). We transported the specimens to the

facilities of the Faculty of Science, University of South Bohemia in České Budějovice, Czech

Republic, for the laboratory experiments.

2.2.Laboratory measurements of body heating

Before the experiments, the butterflies were acclimatised overnight, for at least 15 hours, in a

net cage in a climate-controlled room (20°C) with 16 h day and 8 h night photoperiod.

Butterflies were fed ad libitum by 20% sugar solution applied on the net. We measured the

heating rate using a thermal camera (FLIR P660, Flir Systems) in a climate-controlled room.

The thermal camera was mounted on a tripod 0.60 m above a flat wooden arena where the

butterfly was placed. We used wood as material for the arena because it kept a relatively stable

surface temperature during the experiments. On the camera, we set-up the emissivity to 0.85,

humidity to 66 %, and object distance to 0.60 m for all individual specimens. The emissivity

value compensates for the reflection, transmission, and absorption of IR energy and

corresponds to a conservative estimate for insect cuticle, e.g. Verdu et al. (2012). The humidity

value corresponds to the average air humidity in the experimental room (range 56-82%).

The acclimatised butterflies were first cooled at 5 °C for 10 minutes in a fridge with the lights

on to keep the butterflies active (acclimatization in the dark induced resting behaviour and

butterflies did not stand but fell to one side when placed inside the arena). Then, using cold

forceps, they were placed in the middle of the experimental arena under two halogen light bulbs

(18 W and 160 lm, irradiance at butterfly distance was 45 W/m^2) situated 25 cm above the

arena. However, butterflies were allowed to move across thermally homogeneous arena and

they could change their body posture and open their wings. The butterfly surface body

temperature T_b was recorded every 20 seconds for 10 minutes (Figure 1). In total, we measured

the heating rate in 58 individuals. Nine individuals which escaped from the arena by "jumping"

behaviour sooner than in two minutes (corresponding to 5 thermal camera images) were

removed from subsequent analyses. The remaining 49 individuals were sitting during the

experiment until they took off. Each individual was exposed to the experimental warming only

once. The temperature in a middle point on the mesothorax was recorded using FLIR

QuickReport©1.2 software, using a single manually selected point. The air-conditioning system

in the experimental room created a strong draft, so we kept it turned off for the duration of the

measurements. Consequently, the temperature in the experimental room fluctuated from 17.5

to 21.1 °C. To account for these fluctuations in the statistical analysis, room temperature (T_{room})

and *humidity* was measured just before each trial.

After the thermal camera measurements, the individual butterflies were sacrificed using ethyl-

acetate. We obtained the weight (mg) of fresh individuals using a laboratory balance (precision

0.0001 g). We also photographed the left side of wings of every individual with wings separated

from the body. Photographs of the pair of the wings with a scale were taken by the Panasonic

FZ5 Camera from the distance of 100 cm using focal length 432 mm. Then, we calculated the

wing area in the software ImageJ v.1.46 (Rasband, 1997) and multiplied by two to obtain an

estimate of the total wing area. We calculated wing loading (mg/cm^2) of every individual as the

ratio of body weight (mg) over wing area (cm^2) (Table 1).

2.3 Statistical analyses

Heating curves and species effects

All analyses and data visualisation were done in R version 4.2.0 (R Core Team, 2022). First,

we analysed heating curves, describing the increase of body temperature T_b in time during

basking. The curves were analysed using nonlinear mixed effects models (NLME) with

Gaussian error distribution in the nlme package for R (Pinheiro et al., 2022). The heating curve

was estimated by two alternative models:

1. following the modified asymptotic exponential equation $T_b = T_s\,(1-e^{-b\,(time-c)})$, where T_s

is the estimated asymptotic temperature reached by an individual. The slope of the

asymptotic exponential curve (b) describes the heating rate of the body temperature

increase in time until it reaches asymptotic body temperature T_s.

2. following the modified Michaelis-Menten equation $T_b = T_s((time-c)/(time+K-c))$, where

the parameter T_s is the same as above and K is analogous to the Michaelis constant in

classic Michaelis-Menten equation, but here $K+c$ is the time when T_b reaches 50 % of

the final T_s.

We modified the usual form of the exponential equation and the Michaelis-Menten equation to

include the parameter c which accounts for the fact that the initial T_b at $time=0$ was >0 °C

(Figure 2). Specifically, c is the estimate of how long it would take for the butterfly to increase

its body temperature from $T_b=0$ °C to the real initial T_b (which varied among individuals) under

the experimental conditions. An exponential model (Umbers et al., 2013, Amore et al., 2017)

as well as the Michaelis-Menten equation (Muñoz et al., 2005) were previously used to describe

ectotherm heating, but we are not aware of similar studies comparing the fit of multiple models.

We compared the fit of the two models without any random effects and with two alternative

random effect structures: the random effect of species and the random effect of individual

nested within species. The individual identity was used to account for possible differences of

the heating rate (b) and asymptotic body temperature (T_s) in different individuals. The model

fits were compared by the Akaike information criterion (AIC).

Body weight and wing loading effects

We tested how the shape of the heating curves depends on individual body mass and wing

loading. We used generalised linear mixed effect models (GLMM) in the lme4 (Bates et al.,

2015) and lmerTest libraries for R (Kuznetsova et al., 2017). For every individual, we extracted

the estimates of parameters of asymptotic temperature T_s and heating rate b from the best fitting

NLME model (i.e., modified asymptotic exponential, see Table 2). Then, we correlated the

dependence of individual-level T_s or b on individual body weight and wing-loading. As

additional predictors in the model, we included the T_{room} temperature and *humidity* in the

experimental room recorded during individual measurements. We also compared mean body

weight and wing-loading among species using generalised linear models (GLM) with Gamma

distribution and log link function.

Heating in the laboratory vs. body temperature in the field

We intended to compare species-specific values of the asymptotic temperature (T_s) reached in

laboratory conditions with body temperatures (T_{b_field}) reached in the field (Kleckova et al.,

2014). Because the range of ambient temperatures experienced in the laboratory differed from

the range in the field, we estimated standardised asymptotic body temperature T_s and

standardised body temperature in the field T_{b_field}. In the field, the experienced air and

temperature ranged from 12.0 to 25.0 °C and microhabitat temperature ranged from 11.1 to

31.0 °C. In the laboratory, air temperature ranged 17.5 to 21.1 °C. The standardised T_s and

T_{b_field} temperatures were estimated for 20 °C of ambient laboratory (T_{room}) and for 20 °C of the

two field temperatures, T_{air} (air temperature at breast height) and $T_{microhabitat}$ (microhabitat

temperature) experienced by individual butterflies in the spot where the individuals were

located before capture (see Kleckova et al., 2014). For the laboratory measurements, the

estimate of T_s at ambient temperature 20°C was calculated from a generalised linear model

$T_s\sim T_{room}+species$ fitted to T_s estimates for all individuals (see above). We used the *predict()*

function in R to obtain the estimate of T_s at ambient temperature 20 °C for each species. For

the field observations, we again used the *predict()* function to infer T_{b_field} at ambient

temperature 20 °C for each species. The dependence of body temperature T_{b_field} on both the

ambient air (T_{a_field}) and the microhabitat (T_{m_field}) temperature was obtained from a previous

study (Kleckova et al., 2014). Finally, we tested the relationship between the standardised T_s in

the laboratory and the standardised T_{b_field} reached in the field for individual species by fitting

a generalised linear model with Gaussian error distribution.

3. Results

3.1. Heating curves and species effect

Butterflies typically rested with closed wings after we positioned them in the experimental

arena. Then, they usually opened their wings to a variable degree. Some individuals were

closing and opening their wings repeatedly during the experiments. The wings heated up first

(Figure 1), then, the temperature of the thorax gradually increased. Only a minority (12

individuals) of butterflies took off and flew away from the experimental arena during the

measurements.

The increase of body temperature T_b in the studied *Erebia* butterflies was better described by

the modified asymptotic exponential model (AIC = 583.8), which includes the heating rate

parameter *b*, compared to the modified Michaelis-Menten model (AIC = 1051.3) (Table 2). The

modified asymptotic exponential model predicted that the asymptotic temperature T_s is slightly

more rapidly reached compared to the modified Michaelis-Menten model (Figure 2 and Table

2). The variability of the individual heating curves was best explained by a model including the

random effect of individual identity nested within species (Table 2). However, differences in

the shape of the heating curves among species were modest (Figure 3).

3.2. Body weight and wing loading effects

The six species differed in their body weight ($F_{5,42} = 18.13$, $P < 10^{-8}$) and wing loading ($F_{5,42} =$

13.51, $P < 10^{-7}$, GLM with Gamma error distribution and log link function in both cases),

although there was considerable intraspecific variation (Table 1). As expected, wing loading

and body weight were strongly positively correlated (Pearson's correlation coefficient = 0.84,

$t_{46} = 10.37$, $P < 10^{-12}$). The majority of individuals of all species used in the experiments were

males, except *E. pandrose* for which we measured 4 males and 7 females (Table 1).

Heating at the individual level was affected by both measures of body size. The heating rate *b*

decreased with increasing body mass (GLMM, F = 7.17, P = 0.0105) and wing loading

(GLMM, F = 4.40, P = 0.0419) (Figure 4). On the other hand, the asymptotic temperature T_s

was independent of body mass (GLMM, F = 0.05, P = 0.8209) and wing loading (GLMM, F =

0.16, P = 0.6877). The results, thus, demonstrated that larger butterflies heated up more slowly

but reached the same asymptotic temperature as smaller butterflies. The asymptotic body

temperatures were on average 1.58 ˚C higher than the ambient air temperature (paired t-test, t_{48}

= 13.394, P < 10^{-15}). The GLMMs also included the effect of air temperature and humidity

measured at the time of each observation. While T_s increased with the increasing room

temperature (F = 46.53, P= <0.0001) and humidity (F = 5.12, P = 0.0294), the heating rate

increased with the increasing room temperature (F = 13.82, P = 0.0011) and was not

significantly affected by air humidity (F = 3.68, P = 0.0612).

3.3. Heating in the laboratory vs. body temperature in the field

The standardised asymptotic temperatures T_s measured in the laboratory (Figure 5) were not

correlated with the standardised body temperature reached in field (T_{b_field}). There was no

statistically significant relationship regardless of whether T_{b_field} was standardised for air

temperature T_{a_field} (GLM, F = 0.41, P = 0.5580) or for microhabitat temperature T_{m_field} (F =

0.84, P = 0.4107). It is noteworthy that the two species preferring rocky habitats, *E. cassioides*

and *E. pandrose*, kept the highest body temperatures in the field despite their intermediate

asymptotic temperatures measured in the laboratory (Figure 5). Also, the subalpine species *E.*

alberganus and *E. melampus* kept higher body temperatures than the woodland species *E.*

euryale and *E. ligea* measured both in the field and in the laboratory.

4. Discussion

4.1. Species and individual effects on heating under laboratory conditions

The species-specific differences in body temperature heating in laboratory conditions were

modest (Figure 3), but still detectable (Table 2). The better fit of the individual heating curves

by models including the random effect of individual nested in species (Table 2) indicates that

there is a noticeable effect of individual-level traits (e.g., species-specific physiological

responses) in heating. However, the observed higher heating rate b in individuals with higher

body weight and wing-loading regardless of species identity (Figure 4) points toward a major

role of morphological traits. In other words, larger butterflies heated up more slowly than small

species. This is in agreement with Konvicka et al. (2002), who documented that a large species

in alpine grasslands, *Erebia euryale,* started activity at higher air temperatures and remained

active longer under cloudy weather than a smaller sympatric species, *E. epiphron.* Our results

are also in line with Berwaerts et al. (2001) and Kemp & Krockenberger (2004), who observed

that larger individuals of nymphalid butterflies heated up more slowly than smaller ones. On

the other hand, Neve & Hall (2016) found under controlled experimental conditions that smaller

species of Mediterranean butterflies initiated flight at lower body temperatures than larger ones.

In contrast, smaller *Erebia* species were in flight with higher body temperatures at low ambient

temperatures on mountain slopes of European Alps (Kleckova et al., 2014). By studying the

heating rates of *Erebia* species using a thermal camera in the laboratory, we confirmed that

small body size allows faster heating and likely also facilitates active search for warmer

microclimates during cold mornings or periods of adverse weather.

We have not confirmed our original expectation that larger *Erebia* species and individuals

would accumulate higher body temperatures (higher asymptotic temperatures T_s) in the

laboratory, which was previously recorded in the butterfly species *Polyommatus icarus*

(Lepidoptera: Lycaenidae) (Keyser et al., 2015). However, the similar asymptotic temperatures

reached in our study might be partly explained by the low light intensity used in our

experimental setup 45 W/m^2), while solar flux is around 1000 W/m^2 (Sterhov & Loshkarev,

2019). Future experiments should consider the importance of sufficient light intensity.

Other traits not included in our study, such as wing iridescence (Miaoulis & Heilman, 1998),

emissivity, reflectance and absorbance (Lou et al., 2021; Schmitz, 1994) or fur (modified

scales) thickness (Kingsolver & Moffat, 1982) on the body, may affect heating rates and body

temperatures of butterflies. *Erebia* butterflies are iridescent with metallic reflections in freshly

hatched butterflies, which may increase absorbance of solar influx (Bosi et al., 2008) in

individuals with lower wing-wear. The effect of the density of the thoracic fur has been

demonstrated, e.g., in *Colias* sulphur butterflies, where species inhabiting cold environments

have dense fur on the thorax as it insulates against convective cooling (Watt, 1997). In the

laboratory conditions, a small-bodied species, *E. melampus*, warmed up faster than the

remaining species (Fig. 3), unlike the other small species, *E. cassiodes.* In the studied *Erebia*

species, *E. cassioides* has the most prominent fur on the thorax. We assume that although the

fur may provide better insulation in cold conditions (Kingsolver, 1988), it may also slow down

heating.

The uniform heating rates in the several *Erebia* species in the laboratory conditions with limited

possibilities of microclimate choice are rather expected based on the uniform morphology and

behaviour under the heating source. Although we measured the heating rates in only

approximately ten individuals (Table 1), this main result is robust. The technical restriction

limiting the number of individuals was the availability of a single thermal camera and

temperature-controlled room, which determined the number of individuals we were able to

expose to experimental conditions after collecting them at a field site in Austria, which was

remote from the lab (Czech Republic).

4.2. Habitat use

A previous study on adult microhabitat choice in the field demonstrated that woodland *Erebia* species kept lower body temperatures compared to species of open and rocky habitats (Kleckova et al., 2014). In congruence, the smallest difference in asymptotic temperature T_s reached in the laboratory and T_{b_field} under field conditions was found for the woodland species *E. ligea*. This indicates low buffering ability of this woodland species. Similarly, small differences in body temperatures measured in the field and in the laboratory were detected in another butterfly associated with shrubby grassland habitats and open woodlands, *Hamaeris lucina* (Riodinidae) (Bladon et al., 2020; Hayes et al., 2019). In contrast, the greatest difference between T_s and T_{b_field} was observed for species inhabiting rocky habitats and using stones or gravel as basking sites, *E. cassioides* and *E. pandrose,* indicating a likely active alteration of their body temperature in the wild by microclimate choice.

4.3. Heating curves were better described by the modified asymptotic exponential model

There is no consensus in the literature about the most appropriate model for fitting the heating curves describing the increase of body temperature T_b in time in insects. Exponential models were used to describe ectotherm heating, for example, to compare heating of colour morphs in the chameleon grasshopper (Umbers et al., 2013) and to infer the increase of internal body temperature in dung-beetles (Amore et al., 2017). Another model was the Michaelis-Menten equation, used e.g. to describe heating curves of a sea snail species by Muñoz et al. (2005). Complex models based on detailed physiological measurements of heat transfer within animal bodies and between them and their environment have been developed (Dreisig, 1984;

Dzialowski & O'Connor, 2001), but they are impractical in ecological studies where detailed

physiological parameters are difficult to obtain. In *Erebia* butterflies, the modified asymptotic

exponential model explained the heating of body temperature better than the modified

Michaelis-Menten model (Table 2). The main difference between the two models is that the

exponential model allows for a sharper approach to the asymptote, while the Michaelis-Menten

model is restricted to more gradual approach to the asymptote (see Fig. 2 for an illustrative

example). Heating of the butterflies in our experiments was characterised by a near-linear initial

increase of body temperature followed by a quick stabilisation of the body temperature at the

asymptotic value (Fig. 3), which the exponential model fits better.

5. Implications

Studying the interactions of insect physical traits and their behavioural responses to

environmental factors is important to understand and predict the impacts of climate change on

community composition. When active thermoregulation by microhabitat choice was excluded

under laboratory conditions, six closely related *Erebia* species differed minimally in passive

heating, as measured by heating rate curves (Figure 3). We only detected a negative effect of

individual body size on the rate of heating, but no effect on the asymptotic body temperature.

Thus, the different thermal conditions and habitats found along elevational gradients have not

driven divergent physiological constraints among species as we hypothesised, at least at the

adult life stage. But, in the field, when adult butterflies have the opportunity to actively seek

out suitable microclimates, the study species differed significantly in their body temperatures

(Kleckova et al., 2014). This points toward a more prominent role of behavioural

thermoregulation in these closely related butterflies than the physical traits (body size).

However, further work is needed to test behavioural thermoregulation, e.g., by measuring active

microclimate choice on a thermal gradient. The key role of behaviour in thermoregulation has

been shown in a territorial butterfly (Rutowski et al., 1994) and in butterflies exposed to extreme

temperatures (Tsai et al., 2020). The potential importance of behavioural thermoregulation

implicated from this study suggests that the adult life stage of *Erebia* might not be under direct

thermal selection, in agreement with Buckley et al. (2015). Thus, under increased

anthropogenic pressure on *Erebia* populations, it seems necessary to safeguard a variety of

microhabitats which can facilitate efficient behavioural thermoregulation of adult butterflies

and increased survival of less mobile developmental stages (Abarca et al., 2019). This is in line

with previous calls for management of mountain habitats supporting environmental

heterogeneity (Habel et al., 2022) to increase species survival under ongoing climatic and other

human-induced environmental changes (Lawson et al. 2014b).

6. Conclusions

Heating rates of adult *Erebia* butterflies under laboratory conditions differed slightly among

species. However, smaller individuals heated up more quickly than larger ones regardless of

species identity. This finding implies that small body size likely facilitates earlier initiation of

butterfly daily activities (such as searching for warm microclimates) under unfavourable

weather conditions. This is in an agreement with the observed higher body temperatures of

small butterflies in the field on the mountain slopes of the Alps (Kleckova et al. 2014).

However, the mild inter-specific differences in heating rates contrast with significant inter-

specific differences in body temperatures documented in the field (Kleckova et al. 2014). To

explain this finding, we speculate that microhabitat choice is a primary thermoregulatory

mechanism. Thus, less mobile stages of butterfly developmental cycle are probably more

severely affected by ongoing climatic change, which affects population trends (Konvicka et al.,

2021) and species geographical distributions (Wilson et al., 2007).

Acknowledgements

We are grateful to J. Skála and M. Česánek for recommending the study site, A. Suchačková-

Bartoňová for help with the thermal camera work, O. Nedvěd for providing practical advice, O.

Terland and M. Konvička for the discussion of our results, E. Turner and one anonymous

reviewer for thoughtful comments on a previous version of the manuscript, and to Marie

Hronková for lending us the thermal camera. Funding was provided by the Czech Science

Foundation (GAČR grant: GJ20-18566Y).

Data availability

Data are accessible at https://doi.org/10.6084/m9.figshare.21836103. Full citation for data

repository:

Kleckova I., Okrouhlik J., Svozil T., Matos-Maraví P. & Klecka J. (2022)

Body size, not species identity, drives body heating in alpine *Erebia* butterflies. Figshare

Digital Repository Available at https://doi.org/10.6084/m9.figshare.21836103.

References

Abarca, M., Larsen, E. A., & Ries, L. (2019). Heatwaves and Novel Host Consumption

Increase Overwinter Mortality of an Imperiled Wetland Butterfly. *Frontiers in*

Ecology and Evolution, 7.

https://www.frontiersin.org/articles/10.3389/fevo.2019.00193

Abram, P. K., Boivin, G., Moiroux, J., & Brodeur, J. (2017). Behavioural effects of

temperature on ectothermic animals: Unifying thermal physiology and behavioural

plasticity. *Biological Reviews of the Cambridge Philosophical Society*, *92*(4), 1859–

1876. https://doi.org/10.1111/brv.12312

Advani, N. K., Parmesan, C., & Singer, M. C. (2019). Takeoff temperatures in Melitaea

cinxia butterflies from latitudinal and elevational range limits: A potential adaptation

to solar irradiance. *Ecological Entomology*, *44*(3), 389–396.

https://doi.org/10.1111/een.12714

Amore, V., Hernández, M. I. M., Carrascal, L. M., & Lobo, J. M. (2017). Exoskeleton may

influence the internal body temperatures of Neotropical dung beetles (Col.

Scarabaeinae). *PeerJ*, *5*, e3349. https://doi.org/10.7717/peerj.3349

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects

Models Using lme4. *Journal of Statistical Software*, *67*, 1–48.

https://doi.org/10.18637/jss.v067.i01

Berwaerts, K., Van Dyck, H., Vints, E., & Matthysen, E. (2001). Effect of manipulated wing

characteristics and basking posture on thermal properties of the butterfly Pararge

aegeria (L.). *Journal of Zoology*, *255*(2), 261–267.

https://doi.org/10.1017/S0952836901001327

Bladon, A. J., Lewis, M., Bladon, E. K., Buckton, S. J., Corbett, S., Ewing, S. R., Hayes, M.

P., Hitchcock, G. E., Knock, R., Lucas, C., McVeigh, A., Menéndez, R., Walker, J.

M., Fayle, T. M., & Turner, E. C. (2020). How butterflies keep their cool: Physical

and ecological traits influence thermoregulatory ability and population trends. *Journal*

of Animal Ecology, *89*(11), 2440–2450. https://doi.org/10.1111/1365-2656.13319

Bosi, S. G., Hayes, J., Large, M. C. J., & Poladian, L. (2008). Color, iridescence, and

thermoregulation in *Lepidoptera*. *Applied Optics*, *47*(29), 5235–5241.

https://doi.org/10.1364/AO.47.005235

Buckley, L. B., Ehrenberger, J. C., & Angilletta Jr, M. J. (2015). Thermoregulatory behaviour

limits local adaptation of thermal niches and confers sensitivity to climate change.

Functional Ecology, *29*(8), 1038–1047. https://doi.org/10.1111/1365-2435.12406

Cizek, O., Bakesová, A., Kuras, T., Benes, J., & Konvicka, M. (2003). Vacant niche in alpine

habitat: The case of an introduced population of the butterfly Erebia epiphron in the

Krkonoše Mountains. *Acta Oecologica*, *24*(1), 15–23. https://doi.org/10.1016/S1146-

609X(02)00004-8

Cros, S., Cerdá, X., & Retana, J. (1997). Spatial and temporal variations in the activity

patterns of Mediterranean ant communities. *Écoscience*, *4*(3), 269–278.

https://doi.org/10.1080/11956860.1997.11682405

Dreisig, H. (1984). Control of body temperature in shuttling ectotherms. *Journal of Thermal*

Biology, *9*(4), 229–233. https://doi.org/10.1016/0306-4565(84)90001-9

Dreisig, H. (1995). Thermoregulation and flight activity in territorial male graylings,

Hipparchia semele (Satyridae), and large skippers, Ochlodes venata (Hesperiidae).

Oecologia, *101*(2), 169–176. https://doi.org/10.1007/BF00317280

Dzialowski, E. M., & O'Connor, M. P. (2001). Thermal time constant estimation in warming

and cooling ectotherms. *Journal of Thermal Biology*, *26*(3), 231–245.

https://doi.org/10.1016/s0306-4565(00)00050-4

Forsman, A. (2000). Some like it hot: Intra-Population Variation in behavioral

Thermoregulation in Color-Polymorphic pygmy Grasshoppers. *Evolutionary Ecology*,

14(1), 25–38. https://doi.org/10.1023/A:1011024320725

Gallego, B., Verdú, J. R., & Lobo, J. M. (2018). Comparative thermoregulation between

different species of dung beetles (Coleoptera: Geotrupinae). *Journal of Thermal*

Biology, *74*, 84–91. https://doi.org/10.1016/j.jtherbio.2018.03.009

Habel, J. C., Schmitt, T., Gros, P., & Ulrich, W. (2022). Breakpoints in butterfly decline in

Central Europe over the last century. *Science of The Total Environment*, *851*, 158315.

https://doi.org/10.1016/j.scitotenv.2022.158315

Hayes, M. P., Hitchcock, G. E., Knock, R. I., Lucas, C. B. H., & Turner, E. C. (2019).

Temperature and territoriality in the Duke of Burgundy butterfly, Hamearis lucina.

Journal of Insect Conservation, *23*(4), 739–750. https://doi.org/10.1007/s10841-019-

00166-6

Heinrich, B. (1986). Thermoregulation and Flight Activity Satyrine, Coenonympha Inornata

(Lepidoptera: Satyridae). *Ecology*, *67*(3), 594–597. https://doi.org/10.2307/1937682

Heinrich, B. (1995). Insect thermoregulation. *Endeavour*, *19*(1), 28–33.

https://doi.org/10.1016/0160-9327(95)98891-I

Kemp, D. J., & Krockenberger, A. K. (2002). A novel method of behavioural

thermoregulation in butterflies. *Journal of Evolutionary Biology*, *15*(6), 922–929.

https://doi.org/10.1046/j.1420-9101.2002.00470.x

Kemp, D. J., & Krockenberger, A. K. (2004). Behavioural thermoregulation in butterflies:

The interacting effects of body size and basking posture in Hypolimnas bolina (L.)

(Lepidoptera : Nymphalidae). *Australian Journal of Zoology*, *52*(3), 229–239.

https://doi.org/10.1071/zo03043

Keyser, R. D., Breuker, C. J., Hails, R. S., Dennis, R. L. H., & Shreeve, T. G. (2015). Why

Small Is Beautiful: Wing Colour Is Free from Thermoregulatory Constraint in the

Small Lycaenid Butterfly, Polyommatus icarus. *PLOS ONE*, *10*(4), e0122623.

https://doi.org/10.1371/journal.pone.0122623

Kingsolver, J. G. (1983). Thermoregulation and Flight in Colias Butterflies: Elevational

Patterns and Mechanistic Limitations. *Ecology*, *64*(3), 534–545.

https://doi.org/10.2307/1939973

Kingsolver, J. G. (1985a). Thermoregulatory significance of wing melanization in Pieris

butterflies (Lepidoptera: Pieridae): physics, posture, and pattern. *Oecologia*, *66*(4),

546–553. https://doi.org/10.1007/BF00379348

Kingsolver, J. G. (1985b). Thermoregulatory significance of wing melanization in Pieris

butterflies (Lepidoptera: Pieridae): physics, posture, and pattern. *Oecologia*, *66*(4),

546–553. https://doi.org/10.1007/BF00379348

Kingsolver, J. G. (1988). Thermoregulation, Flight, and the Evolution of Wing Pattern in

Pierid Butterflies: The Topography of Adaptive Landscapes. *American Zoologist*,

28(3), 899–912. https://doi.org/10.1093/icb/28.3.899

Kingsolver, J. G., & Moffat, R. J. (1982). Thermoregulation and the determinants of heat

transfer in Colias butterflies. *Oecologia*, *53*(1), 27–33.

https://doi.org/10.1007/BF00377132

Kirkpatrick, W. H., & Sheldon, K. S. (2022). Experimental increases in temperature mean and

variance alter reproductive behaviours in the dung beetle Phanaeus vindex. *Biology*

Letters, *18*(7), 20220109. https://doi.org/10.1098/rsbl.2022.0109

Kleckova, I., & Klecka, J. (2016). Facing the Heat: Thermoregulation and Behaviour of

Lowland Species of a Cold-Dwelling Butterfly Genus, Erebia. *PLOS ONE*, *11*(3),

e0150393. https://doi.org/10.1371/journal.pone.0150393

Kleckova, I., Konvicka, M., & Klecka, J. (2014). Thermoregulation and microhabitat use in

mountain butterflies of the genus Erebia: Importance of fine-scale habitat

heterogeneity. *Journal of Thermal Biology*, *41*, 50–58.

https://doi.org/10.1016/j.jtherbio.2014.02.002

Konvicka, M., Beneš, J., & Kuras, T. (2002). Microdistribution and diurnal behaviour of two

sympatric mountainous butterflies (Erebia epiphron and E. euryale): Relations to

vegetation and weather. *Biologia*, *57*, 225–235.

Konvicka, M., Kuras, T., Liparova, J., Slezak, V., Horázná, D., Klečka, J., & Kleckova, I.

(2021). Low winter precipitation, but not warm autumns and springs, threatens

mountain butterflies in middle-high mountains. *PeerJ*, *9*, e12021.

https://doi.org/10.7717/peerj.12021

Konvicka, M., Maradova, M., Benes, J., Fric, Z., & Kepka, P. (2003). Uphill shifts in

distribution of butterflies in the Czech Republic: Effects of changing climate detected

on a regional scale. *Global Ecology and Biogeography*, *12*(5), 403–410.

https://doi.org/10.1046/j.1466-822X.2003.00053.x

Kuras, T., Benes, J., Fric, Z., & Konvicka, M. (2003). Dispersal patterns of endemic alpine

butterflies with contrasting population structures: Erebia epiphron and E. sudetica.

Population Ecology, *45*(2), 115–123. https://doi.org/10.1007/s10144-003-0144-x

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in

Linear Mixed Effects Models. *Journal of Statistical Software*, *82*, 1–26.

https://doi.org/10.18637/jss.v082.i13

Lawson, C. R., Bennie, J., Hodgson, J. A., Thomas, C. D., & Wilson, R. J. (2014).

Topographic microclimates drive microhabitat associations at the range margin of a

butterfly. *Ecography*, *37*(8), 732–740. https://doi.org/10.1111/ecog.00535

Lawson, C. R., Bennie, J. J., Thomas, C. D., Hodgson, J. A., & Wilson, R. J. (2014). Active

Management of Protected Areas Enhances Metapopulation Expansion Under Climate

Change. *Conservation Letters*, *7*(2), 111–118. https://doi.org/10.1111/conl.12036

Lou, C., An, S., Yang, R., Zhu, H., Shen, Q., Jiang, M., Fu, B., Tao, P., Song, C., Deng, T., &

Shang, W. (2021). Enhancement of infrared emissivity by the hierarchical

microstructures from the wing scales of butterfly Rapala dioetas. *APL Photonics*, *6*(3),

036101. https://doi.org/10.1063/5.0039079

MacLean, H. J., Higgins, J. K., Buckley, L. B., & Kingsolver, J. G. (2016). Morphological

and physiological determinants of local adaptation to climate in Rocky Mountain

butterflies. *Conservation Physiology*, *4*(1), cow035.

https://doi.org/10.1093/conphys/cow035

Mattila, A. L. K. (2015). Thermal biology of flight in a butterfly: Genotype, flight

metabolism, and environmental conditions. *Ecology and Evolution*, *5*(23), 5539–5551.

https://doi.org/10.1002/ece3.1758

Miaoulis, I. N., & Heilman, B. D. (1998). Butterfly Thin Films Serve as Solar Collectors.

Annals of the Entomological Society of America, *91*(1), 122–127.

https://doi.org/10.1093/aesa/91.1.122

Muñoz, J. L. P., Randall Finke, G., Camus, P. A., & Bozinovic, F. (2005). Thermoregulatory

behavior, heat gain and thermal tolerance in the periwinkle Echinolittorina peruviana

in central Chile. *Comparative Biochemistry and Physiology Part A: Molecular &*

Integrative Physiology, *142*(1), 92–98. https://doi.org/10.1016/j.cbpa.2005.08.002

Nève, G., & Després, L. (2020). Cold adaptation across the elevation gradient in an alpine

butterfly species complex. *Ecological Entomology*, *45*(5), 997–1003.

https://doi.org/10.1111/een.12875

Neve, G., & Hall, C. (2016). Variation of thorax flight temperature among twenty Australian

butterflies (Lepidoptera: Papilionidae, Nymphalidae, Pieridae, Hesperiidae,

Lycaenidae). *European Journal of Entomology*, *113*, 571–578.

https://doi.org/10.14411/eje.2016.077

Pinheiro, J., Bates, D., & R Core Team. (2022). *Nlme: Linear and Nonlinear Mixed Effects*

Models. R package version 3.1-157, https://CRAN.R-project.org/package=nlme.

Pivnick, K. A., & McNeil, J. N. (1986). Sexual Differences in the Thermoregulation of

Thymelicus Lineola Adults (Lepidoptera: Hesperiidae). *Ecology*, *67*(4), 1024–1035.

https://doi.org/10.2307/1939825

Polic, D., Fiedler, K., Nell, C., & Grill, A. (2014). Mobility of ringlet butterflies in high-

elevation alpine grassland: Effects of habitat barriers, resources and age. *Journal of*

Insect Conservation, *18*(6), 1153–1161. https://doi.org/10.1007/s10841-014-9726-5

R Core Team. (2022). *R: A language and environment for statistical computing. R*

Foundation for Statistical Computing, Vienna, Austria. Https://www.R-project.org/.

Radchuk, V., Turlure, C., & Schtickzelle, N. (2013). Each life stage matters: The importance

of assessing the response to climate change over the complete life cycle in butterflies.

Journal of Animal Ecology, *82*(1), 275–285. https://doi.org/10.1111/j.1365-

2656.2012.02029.x

Rasband, W. S. (1997). *ImageJ, U. S. National Institutes of Health, Bethesda, Maryland, USA*

(v.1.46). https://imagej.nih.gov/ij/

Rödder, D., Schmitt, T., Gros, P., Ulrich, W., & Habel, J. C. (2021). Climate change drives

mountain butterflies towards the summits. *Scientific Reports*, *11*(1), 14382.

https://doi.org/10.1038/s41598-021-93826-0

Rutowski, R. L., Demlong, M. J., & Leffingwell, T. (1994). Behavioural thermoregulation at

mate encounter sites by male butterflies (Asterocampa leilia, Nymphalidae). *Animal*

Behaviour, *48*(4), 833–841. https://doi.org/10.1006/anbe.1994.1307

Schmitz, H. (1994). Thermal characterization of butterfly wings—1. Absorption in relation to

different color, surface structure and basking type. *Journal of Thermal Biology*, *19*(6),

403–412. https://doi.org/10.1016/0306-4565(94)90039-6

Shreeve, T. G. (1984). Habitat Selection, Mate Location, and Microclimatic Constraints on

the Activity of the Speckled Wood Butterfly Pararge Aegeria. *Oikos*, *42*(3), 371–377.

https://doi.org/10.2307/3544407

Sistri, G., Menchetti, M., Santini, L., Pasquali, L., Sapienti, S., Cini, A., Platania, L., Balletto,

E., Barbero, F., Bonelli, S., Casacci, L. P., Dincă, V., Vila, R., Mantoni, C., Fattorini,

S., & Dapporto, L. (2022). The isolated Erebia pandrose Apennine population is

genetically unique and endangered by climate change. *Insect Conservation and*

Diversity, *15*(1), 136–148. https://doi.org/10.1111/icad.12538

Slamova, I., Klecka, J., & Konvicka, M. (2011). Diurnal Behavior and Habitat Preferences of

Erebia aethiops, an Aberrant Lowland Species of a Mountain Butterfly Clade. *Journal*

of Insect Behavior, *24*(3), 230–246. https://doi.org/10.1007/s10905-010-9250-8

Slamova, I., Klecka, J., & Konvicka, M. (2013). Woodland and grassland mosaic from a

butterfly perspective: Habitat use by Erebia aethiops (Lepidoptera: Satyridae). *Insect*

Conservation and Diversity, *6*(3), 243–254. https://doi.org/10.1111/j.1752-

4598.2012.00212.x

Sterhov, A. I., & Loshkarev, I. Y. (2019). Determination of the proportion of natural light in

solar radiation using the method of conversion of lighting units into energy. *Journal of*

Physics: Conference Series, *1353*(1), 012002. https://doi.org/10.1088/1742-

6596/1353/1/012002

Stuhldreher, G., & Fartmann, T. (2018). Threatened grassland butterflies as indicators of

microclimatic niches along an elevational gradient – Implications for conservation in

times of climate change. *Ecological Indicators*, *94*, 83–98.

https://doi.org/10.1016/j.ecolind.2018.06.043

Tsai, C.-C., Childers, R. A., Nan Shi, N., Ren, C., Pelaez, J. N., Bernard, G. D., Pierce, N. E.,

& Yu, N. (2020). Physical and behavioral adaptations to prevent overheating of the

living wings of butterflies. *Nature Communications*, *11*(1), Article 1.

https://doi.org/10.1038/s41467-020-14408-8

Umbers, K. D. L., Herberstein, M. E., & Madin, J. S. (2013). Colour in insect

thermoregulation: Empirical and theoretical tests in the colour-changing grasshopper,

Kosciuscola tristis. *Journal of Insect Physiology*, *59*(1), 81–90.

https://doi.org/10.1016/j.jinsphys.2012.10.016

Verdu, J. R., Alba-Tercedor, J., & Jimenez-Manrique, M. (2012). Evidence of Different

Thermoregulatory Mechanisms between Two Sympatric Scarabaeus Species Using

Infrared Thermography and Micro-Computer Tomography | PLOS ONE. *PLoS ONE*,

7(3), | e3391.

Watt, W. B. (1997). Accuracy, Anecdotes, and Artifacts in the Study of Insect Thermal

Ecology. *Oikos*, *80*(2), 399. https://doi.org/10.2307/3546607

Wickman, P. O. (2009). Thermoregulation and habitat use in butterflies. In *Settele, J.,*

Shreeve, T., Konvicka, M., Van Dyck, H. (Eds.). Ecology of Butterflies in Europe (pp.

55–61).

Wickman, P.-O. (1988). Dynamics of mate-searching behaviour in a hilltopping butterfly,

Lasiommata megera (L.): The effects of weather and male density. *Zoological Journal*

of the Linnean Society, *93*(4), 357–377. https://doi.org/10.1111/j.1096-

3642.1988.tb01367.x

Willmer, P. G. (1982). Microclimate and the Environmental Physiology of Insects. In M. J.

Berridge, J. E. Treherne, & V. B. Wigglesworth (Eds.), *Advances in Insect Physiology*

(Vol. 16, pp. 1–57). Academic Press. https://doi.org/10.1016/S0065-2806(08)60151-4

Wilson, R. J., Bennie, J., Lawson, C. R., Pearson, D., Ortúzar-Ugarte, G., & Gutiérrez, D.

(2015). Population turnover, habitat use and microclimate at the contracting range

margin of a butterfly. *Journal of Insect Conservation*, *19*(2), 205–216.

https://doi.org/10.1007/s10841-014-9710-0

Wilson, R. J., Gutiérrez, D., Gutiérrez, J., & Monserrat, V. J. (2007). An elevational shift in

butterfly species richness and composition accompanying recent climate change.

Global Change Biology, *13*(9), 1873–1887. https://doi.org/10.1111/j.1365-

2486.2007.01418.x

Tables

Table 1. The list of study species of the butterfly genus *Erebia*. For each species, we provide

information about its typical habitat, phenology (adults flying early or late in the summer), the

number of individuals used for the lab measurements (N, males (M)/females (F)), their mean

body weight, total wing area, and wing loading (body weight/wing area). Species are sorted

according to their body weight from lightest to heaviest.

Species	Habitat	Phenology	N	Body weight (mg)	Wing area (cm²)	Wing loading (mg/cm²)
			(M/F)	mean (SD)	mean (SD)	mean (SD)
E. melampus	grassland	early	7 (7/0)	24.4 (4.17)	3.91 (0.272)	6.25 (0.805)
E. cassioides	rocks	late	7 (7/0)	34.4 (7.01)	5.48 (0.510)	6.30 (1.365)
E. euryale	woodland	late	8 (8/0)	38.5 (5.02)	7.72 (0.675)	5.01 (0.719)
E. ligea	woodland	late	7 (6/1)	50.1 (21.76)	8.97 (0.611)	5.49 (1.899)
E. alberganus	grassland	early	9 (9/0)	54.0 (11.55)	6.45 (0.699)	8.32 (1.271)
E. pandrose	rocks	early	11 (4/7)	68 (14.51)	6.79 (0.693)	10.15 (2.572)

Table 2. The comparison of the fit of the modified asymptotic exponential and Michaelis-Menten models describing butterfly heating under laboratory conditions. The models describe the increase of body temperature (T_b) in *time* in 49 individuals of *Erebia* butterflies, which heated up under an artificial source of heating and radiation. Their T_b was measured by thermal camera. The non-linear mixed effect models included the random effects of individual nested in species and the random effect of species. The best model is highlighted **in bold**. Parameters describe maximum body temperature reached during heating (asymptotic temperature T_s), heating rate (b), Michaelis-Menten constant (K), and the estimated time it would take for the butterfly to increase its body temperature from $T_b=0$ °C to the actual initial T_b (c).

Model	Random effects	T_s (SE)	b (SE)	K (SE)	c (SE)	AIC	ΔAIC
		Parameter estimates (fixed effects)					
modified asymptotic exponential	**Individual nested in species**	**21.00 (0.271)**	**0.687 (0.056)**		**-1.188 (0.152)**	**583.8**	**0**
	Species	20.93 (0.291)	0.695 (0.061)		-1.138 (0.141)	2375.5	1791.7
	None	20.94 (0.037)	0.660 (0.013)		-1.140 (0.033)	3410	2826.2
modified Michaelis-Menten	Individual nested in species	22.73 (0.334)		0.637 (0.073)	-0.559 (0.082)	1051.3	467.5
	Species	22.46 (0.377)		0.587 (0.077)	-0.514 (0.078)	2485.1	1901.3
	None	22.48 (0.068)		0.579 (0.018)	-0.507 (0.020)	3422.2	2838.4

706 **Figures**

708 **Figure 1.** An example image from the thermal camera displaying the heating of an individual

709 *Erebia* butterfly in *time* (seconds, s), blue indicates colder, red warmer temperatures. Typically,

710 the butterflies were sitting initially with closed wings and open them later.

Figure 2. An example heating curve of a single individual of *Erebia melampus* butterfly with

the comparison of the fit of the modified asymptotic exponential (black line) and Michaelis-

Menten models (orange line). The modified asymptotic exponential model fits the data for this

individual significantly better (ΔAIC = 38.0).

Figure 3. Heating of six *Erebia* butterfly species (49 individuals in total) under artificial source of heat and radiation under laboratory conditions. The lines display the fit of the modified asymptotic exponential model. The species-specific parameters of the model were estimated by a nonlinear mixed effects model with individual and species identity included as nested random effects. The parameter b describes the heating rate, i.e. the increase of the body temperature (T_b) in *time* until it reaches the asymptotic body temperature (T_s). The parameter c is the estimate of how long it would take to increase the body temperature from T_b=0 °C to the real initial T_b. Equations describing the fitted curves with species-specific parameter estimates are added.

Figure 4. The effect of body weight and wing loading on the asymptotic body temperature (T_s)

and on the heating rate (b) of 49 individuals of six *Erebia* species. The T_s and b parameters

describing heating of individuals under laboratory conditions were inferred from the modified

asymptotic exponential models (see Figure 3). The y-axis shows partial residuals from a model

which also included the room temperature and humidity at the time of each measurement. Each

dot represents one individual, grouped into species according to colour.

Figure 5. The correlation between *Erebia* butterfly species body temperatures reached under

laboratory conditions ($T_{b\_laboratory}$) and mean body temperatures T_{b_field} measured in the field at

20°C of ambient air (panel a.) and microhabitat (panel b.) temperature. See the Methods section

for more details. Estimates of T_b values are accompanied by standard errors.

739