1   *Brief Report*

4   **Midbody proteins display distinct temporal dynamics during cytokinesis**

7   Ella F.J. Halcrow[#], Riccardo Mazza[#], Anna Diversi, Anton Enright and Pier Paolo D'Avino[*]

9   Department of Pathology, University of Cambridge, Tennis Court Road, Cambridge, CB2 1QP,

10   UK

18   [#]These two authors contributed equally

20   *Correspondence to: Pier Paolo D'Avino, email: ppd21@cam.ac.uk

Abstract

The midbody is an organelle that forms between the two daughter cells during cytokinesis. It co-ordinates the abscission of the nascent daughter cells and is composed of a multitude of proteins that are meticulously arranged into distinct temporal and spatial localization patterns. However, very little is known about the mechanisms that regulate the localization and function of midbody proteins. Here, we analyzed the temporal and spatial profiles of key midbody proteins during mitotic exit under normal conditions and after treatment with drugs that affect phosphorylation and proteasome-mediated degradation to decipher the impacts of post-translational modifications on midbody protein dynamics. Our results highlighted that midbody proteins show distinct spatio-temporal dynamics during mitotic exit and cytokinesis that depend on both ubiquitin-mediated proteasome degradation and phosphorylation/de-phosphorylation. They also identified two discrete classes of midbody proteins: 'transient' midbody proteins -including Anillin and PRC1- which rapidly accumulate at the midbody after anaphase onset and then slowly disappear, and 'stable' midbody proteins -including CIT-K, KIF14 and KI1F20A- which instead persist at the midbody throughout cytokinesis and also post abscission. These two classes of midbody proteins display distinct interaction networks with ubiquitylation factors, which could potentially explain their different dynamics and stability during cytokinesis.

Keywords: cytokinesis, midbody, phosphorylation, protein dynamics, ubiquitylation.

1. Introduction

Proper execution of cell division regulates growth, development, and reproduction by controlling the partition of genomic and cytoplasmic contents between the two nascent daughter cells. Because of the inaccessibility of the chromatin compacted into chromosomes gene transcription is severely limited during mitosis. Consequently, the vast majority of mitotic processes are regulated by reversible PTMs, including phosphorylation and ubiquitylation, which are mediated by opposing PTM enzymes (e.g., kinases vs. phosphatase and ubiquitin ligases vs. hydrolases) [1,2]. These PTMs control the intricate and finely tuned signals and protein-protein interaction networks that are responsible for the assembly of the mitotic spindle and chromosome alignment in prometaphase and metaphase, and chromosome segregation and daughter cell separation during mitotic exit and cytokinesis. Once chromosomes are properly aligned at the metaphase plate, anaphase onset is triggered by the activation of an E3 ubiquitin ligase, the anaphase promoting complex/cyclosome (APC/C), through its interaction with the cofactor Cdc20 [3]. APC/C^{Cdc20} then targets several proteins for destruction by the 26S proteasome, including cyclin B, which in turn leads to the inactivation of its cyclin-dependent kinase 1 (CDK1) partner. CDK1 inactivation is accompanied by increased activity of PP1 and PP2A serine/threonine phosphatases and changes in the distribution of other serine/threonine mitotic kinases, including Polo-like kinase 1 (PLK1) and Aurora B (AURKB), the kinase component of the chromosomal passenger complex (CPC) [4-9]. Together, these events lead to complex changes in the phosphorylation profiles and activity of a multitude of proteins that propel a remarkable re-organization of the cytoskeleton [10]. Initially, cells determine the position of the cleavage furrow through signals generated by the spindle microtubules (MTs), which are re-organized into an array of antiparallel and interdigitating MTs known as the central spindle. Spindle MTs also promote furrow ingression, which is driven by the assembly and constriction of an actomyosin contractile ring. During furrow ingression, the contractile ring compacts the central spindle and the two daughter cells remain connected by an intercellular bridge, which contains at its center an organelle, the midbody, composed of a multitude of proteins that have diverse functions (Fig. 1A) [11,12]. Some midbody proteins are former components of the contractile ring and central spindle, while others are specifically recruited during the slow midbody maturation process that ultimately leads to the abscission of the two daughter cells [13,14]. All these proteins are arranged in a very precise and stereotyped spatial pattern along the midbody [15], which depends on the multifunctional protein Citron kinase (CIT-K) [16-18]. The proper localization,

regulation and interactions of all these proteins are essential for the execution of abscission

and to prevent incorrect genome segregation [14]. In addition, recent studies have revealed

that the midbody also has important functions after cell division. Following abscission, the

midbody remnants can be either reabsorbed by one of the daughter cells or released into the

extracellular environment and then eventually internalized by another cell [19]. These post-

mitotic midbody remnants have been implicated in disparate biological processes, including

cell fate, pluripotency, apical-basal polarity, tissue organization, cell proliferation, cancer, and

cilium and lumen formation [20]. Finally, some midbody proteins have been linked to brain

development and microcephaly [21]. However, despite the evidence of the involvement of the

midbody in these important processes, our understanding of the mechanisms that regulate

its formation and functions are still very limited. In this study, we report that midbody proteins

show distinct spatio-temporal dynamics that identify two general classes: (i) 'transient'

midbody proteins, including Anillin (ANLN), AURKB, and protein regulator of cytokinesis 1

(PRC1), which rapidly accumulate at the midbody after anaphase onset, but then slowly

disappear; and (ii) 'stable' midbody proteins, including CIT-K, and the kinesins KIF14 and

KIF23/MKLP1, which instead persist to the midbody for much longer, even in post-abscission

midbodies. Furthermore, we present evidence that these different dynamics appears to be

regulated by both phosphorylation and ubiquitylation.

2. Materials and Methods

2.1 Cell culture and treatments

HeLa Kyoto were maintained in DMEM (Sigma) containing 10% Fetal Bovine Serum (Sigma)

and 1% pennicillin/streptomycin (Invitrogen) at 37°C and 5% CO_2. HeLa cell lines stably

expressing GFP-tagged transgenes were described previously [11] and cultured in the same

medium with the addition of appropriate selection antibiotics (puromycin and/or G418).

For RNA interference the following siRNAs were used: scrambled sequence control: 5'-

AACGUACGCGGAAUACUUCGA-3', Anillin (ANLN): 5'-GUAUCGAAACCAAUUGUGAAGUCAA-3',

KIF23/MKLP1:   5'-GCAGUCUUCCAGGUCAUCU-3',   using   Lipofectamine   RNAiMAX

(ThermoFisher) following the manufacturer's instructions.

To synchronize HeLa Kyoto cells at different stages of mitosis, we used a thymidine-

nocodazole block and release procedure essentially as described [11]. Cells were first arrested

in S phase by the addition of 2 mM thymidine (Sigma-Aldrich) for 19 h, washed twice with

phosphate-buffered saline (PBS) and released for 5 h in fresh complete medium. Cells were

then cultured for additional 13 h in fresh complete medium containing 50 ng/ml nocodazole

(Sigma-Aldrich) and then harvested by mitotic shake-off. Mitotic cells were washed five times

with PBS, and released in fresh medium containing either one of the following drugs: 10 µM

MG132 (proteasome inhibitor, Sigma), 50 nM okadaic acid (PP1 and PP2A inhibitor,

Calbiochem), 2 µM tautomycin (PP1 inhibitor, TOCRIS), 2 µM ZM (AURKB inhibitor, TOCRIS) or

the DMSO solvent as control. Cells were then harvested by centrifugation and frozen in dry

ice.

2.2 Fluorescence microscopy

HeLa cells were grown on microscope glass coverslips (Menzel-Gläser) and fixed in either

PHEM buffer (60 mM Pipes, 25 mM Hepes pH 7, 10 mM EGTA, 4 mM MgCl$_2$, 3.7% [v/v]

formaldheyde) for 12 min at room temperature or in ice-cold methanol for 10 min at -20°C.

They were then washed three times for 10 min with PBS and incubated in blocking buffer (PBS,

0.5% [v/v] Triton X-100 and 5% [w/v] BSA) for 1 h at room temperature. Coverslips were

incubated overnight at 4°C with the primary antibodies indicated in the figure legends, diluted

in PBT (PBS, 0.1% [v/v] Triton X-100 and 1% [w/v] BSA). The day after, coverslips were washed

twice for 5 min in PBT, incubated with secondary antibodies diluted in PBT for 2h at RT and

then washed twice with PBT and once with PBS. Coverslips were mounted on SuperFrost

Microscope Slides (VWR) using VECTASHIELD Mounting Medium containing DAPI (Vector

Laboratories). Images were acquired using a Zeiss Axiovert epifluorescence microscope

equipped with MetaMorph software. Fiji [22] was used to generate maximum intensity

projections, which were adjusted for contrast and brightness and assembled using Photoshop.

2.3 Western blot

Cells were centrifuged, resuspended in phosphate buffer saline (PBS) and then an equal

volume of 2x Laemmli buffer was added. Samples were then boiled for 10 min and stored at

-20°C. Proteins were separated by SDS PAGE and then transferred onto PVDF membrane

(Immobilon-P) at 15V for 1 hour. Membranes were blocked overnight at 4°C in PBS + 0.1%

(v/v) Tween (PBST) with 5% (v/v) dry milk powder. After blocking, membranes were washed

once with PBST and then incubated with the appropriate primary antibody diluted in PBST +

3% (v/v) BSA (Sigma) for 2 hours at RT. Membranes were washed 3x5 minutes in PBST and

then incubated with HRPA-conjugated secondary antibodies in PBST + 1% BSA for 1 hour at

room temperature. After further 3x5 min washes in PBST, the signals were detected using the

ECL West Pico substrate (ThermoFisher) and chemiluminescent signals were acquired below

saturation levels using a G:BOX Chemi XRQ (Syngene) and quantified using Fiji [22].

2.4 Antibodies

The following antibodies were used in this study: rabbit polyclonal anti-ANLN (Abcam,

ab154337, dilutions for WB, 1:2000, for IF 1:200), mouse monoclonal anti-Aurora B (clone

AIM-1, BD Transduction Laboratories, 611082 dilutions for WB 1:2000, for IF 1:100), mouse

monoclonal anti-CIT-K (BD Transduction Laboratories, 611377, dilutions for WB 1:1500, for IF

1:200), mouse monoclonal anti-cyclin B1 (clone GNS1, Santa Cruz, sc-245 dilution for WB

1:2000), rabbit monoclonal anti-MKLP1 (Abcam ab174304, dilutions for WB 1:5000, for IF

1:800), rabbit polyclonal anti-KIF20A/MKLP2 (a kind gift of T.U. Mayer [23], dilution for both

WB and IF 1:1000) , mouse monoclonal anti-PRC1 (clone C-1, Santa Cruz, sc-376983 dilutions

for WB 1:5000, for IF 1:100), rabbit polyclonal anti-phospho-histone H3 pS10 (Merck, 06-570

dilution for WB 1:10000), mouse monoclonal anti α-tubulin (clone DM1A, Sigma, T9026

dilutions for WB 1:20000, for IF 1:2000), rabbit polyclonal anti-β-tubulin (Abcam, ab6046

dilutions for WB 1:5000, for IF 1:400). Peroxidase and Alexa-fluor conjugated secondary

antibodies were purchased from Jackson Laboratories and ThermoFisher, respectively.

2.5 Time-lapse imaging

For time-lapse experiments, HeLa cells expressing the different GFP-tagged proteins were

plated on an open μ-Slide with 8 wells (Ibidi, 80826) in complete medium containing 0.5 μM

SiR-DNA dye (SpiroChrome). Images were acquired on a Leica TCS SP8 Inverted Microscope

with a 40x/1.30 NA HC Plan APO CS2 - OIL DIC objective and argon laser power set at 80%. The

Application Suite X software (LAS-X; Leica) for multidimensional image acquisition was used.

Specimens were maintained at 37°C and 5% CO_2 via a chamber, and z-series of ten, 1-μm

sections were captured at 2 min intervals. All images were processed using Fiji [22] to generate

maximum intensity projections, to adjust for brightness and contrast, and to create the final

movies. The fluorescence intensity values shown in Figure 3 were measured from whole cells

(I_C) or midbodies (I_M) at the different time points indicated using Fiji [22]. A background

intensity value, measured at the same time point and from an identically sized area, was

subtracted from each value.

2.6 Computational and statistical analyses

To generate the ubiquitylation midbody sub-network, we searched our midbody interactome

dataset [11] for proteins whose Uniprot protein names field contained the term 'ubiquitin' via

grep in the Unix command line. This generated an initial dataset that was subsequently

manually curated to eliminate proteins that were not directly involved in ubiquitylation. The

final list of 86 proteins (Table S1) was entered into a raw tab-delimited text file and then

imported into Cytoscape to generate the network shown in Figure 4.

Prism 9 (GraphPad) and Excel (Microsoft) were used for statistical analyses and to prepare

graphs.

3. Results and Discussion

3.1 Midbody protein distribution changes during midbody maturation.

Immuno-fluorescence and electron microscopy studies have indicated that the midbody can

be divided in three major regions: (i) the midbody ring, containing former contractile ring

components like Anillin (ANLN) and CIT-K; (ii) the midbody central core, marked by central

spindle proteins such as the centralspindlin complex (an heterotetramer composed of two

subunits of the kinesin KIF23/MKLP1 and two molecules of RacGAP1); and (iii) the midbody

arms, which flank the midbody core and where AURKB and the kinesin KIF20A/MKLP2

accumulate (Fig. 1A) [13,15,18]. However, the midbody is not a static structure, it undergoes a

series of morphological changes during the late stages of cytokinesis, in a process known as

midbody maturation. After completion of furrow ingression, two symmetric constrictions

form at both sides of the midbody ring, making the midbody look similar to a 'bow tie' (Fig. 1).

Subsequently, the microtubule bundles become progressively thinner and ultimately a distinct

abscission site appears usually first at one side of the midbody ring [13]. These changes in

midbody architecture are often reflected by changes in the distribution of midbody proteins.

For example, after completion of furrow ingression ANLN and CIT-K both localize to the

midbody ring -albeit ANLN display a broader distribution (Fig. 1B and C)- but, whilst CIT-K

maintains a ring-like distribution and persists in post-abscission midbody remnants (Fig. 3C)

[17,18], ANLN accumulates to the secondary constriction sites and then disappears from the

midbody before abscission (Fig. 1C) [24]. AURKB localizes to the midbody arms throughout

midbody maturation, although its accumulation slowly decreases during midbody maturation,

similarly to ANLN. The kinesin KIF23/MKLP2 localizes to the midbody core after furrow

ingression and then, starting from the 'bow tie' stage, it forms two juxtaposed discs and

persists in post-abscission midbodies like CIT-K (Fig. 1C and Fig 3D).

Previous studies have indicated that CIT-K plays an important role in establishing and
maintaining the orderly distribution of these midbody proteins, possibly through its direct
interaction with some midbody components, including ANLN, AURKB, KIF14 and KIF23/MKLP1
[11,16-18,25,26]. Nevertheless, we still lack sufficient knowledge of the underlying mechanisms
that control the dynamics and stability of midbody proteins during and after cytokinesis.

3.2 Midbody proteins display different expression profiles during mitotic exit and cytokinesis
To get a detailed understanding of the dynamics of midbody proteins, we first analyzed their
expression profiles by Western blot in synchronized cells as they exited mitosis. HeLa cells
were synchronized in prometaphase by thymidine/nocodazole block and then released to
analyze the levels of midbody proteins (Fig. 2A and B). The specificity of the antibodies used
in this analysis has been previously validated [11,18,27], apart from the anti-ANLN and a new
anti-KIF23/MKLP1 antibody, which we validated using siRNA-mediated depletion
(Supplementary Fig. S1). The levels of cyclin B and histone H3 phosphorylated at S10 (pH3),
which both decrease after anaphase onset, were used to monitor mitotic exit. Cyclin B levels
dropped significantly (more than 80%) 60 min after release from nocodazole, indicating that
the vast majority of cells had exited mitosis at this time point. The levels of KIF14 and
KIF23/MKLP1 remained relatively stable after nocodazole release, whereas the levels of all
other midbody proteins decreased during mitotic exit (Fig. 2A and 2B). ANLN showed a profile
very similar to that of Cyclin B, while the levels of all the other midbody proteins analyzed,
AURKB, CIT-K, KIF20A/MKLP2 and PRC1 decreased more slowly, in parallel with pH3.
As mentioned before, mitotic exit is triggered by ubiquitin-mediated protein degradation and
regulated by phosphorylation/dephosphorylation of cytokinesis proteins. Therefore, to
understand which PTM(s) might be involved in regulating the level of midbody proteins, we
incubated HeLa cells for a short period of time (one hour) with 4 different inhibitors – the
proteasome inhibitor MG132, okadaic acid to inhibit both PP1 and PP2A phosphatases, the
PP1-specific inhibitor tautomycetin [28], and the AURKB inhibitor ZM447439 [29] – starting
from 45 minutes after release from nocodazole, when most cells should be already in, or about
to enter anaphase. The analysis from this indicated that midbody proteins can be divided into
three main groups on the basis of their response to drug treatments (Fig, 2C, 2D and
Supplementary Fig. S2). In one group, ANLN and PRC1 profiles mimicked that of cyclin B, as
they were stabilized after treatment with MG132 (Fig. 2C, 2D and Supplementary Fig. S2A).
This is consistent with the evidence that all these proteins are ubiquitinated to be degraded

during mitotic exit [30-32]. The levels of all these three proteins also appeared to decline faster

after incubation with the AURKB inhibitor ZM447439 (Fig. 2C, 2D and Supplementary Fig. S2A),

which could either reflects a more rapid mitotic exit triggered by AURKB inhibition or indicate

that AURKB activity is required for the stability of these midbody proteins in late telophase.

We favor the latter for two main reasons. Firstly, none of the other proteins showed a similar

response and secondly, ZM447439 was added 45 minutes after nocodazole release, when the

majority of cells were already in anaphase. A second group included AURKB and CIT-K, which

were also stabilized by MG132 treatment (albeit less than ANLN, cyclin B and PRC1), but their

levels did not change after treatment with any of the other inhibitors (Fig. 2C, 2D and

Supplementary Fig. S2B). These results are consistent with the role of the APC/C in the

degradation of Aurora kinases during mitotic exit [33], but reveal for the first time that CIT-K

might also be similarly targeted for degradation via ubiquitination. The final group comprises

the kinesins, KIF14, KIF23/MKLP1, and KIF20A/MKLP2, which do not appear to be particularly

affected by any of the drug treatment, with the exception of KIF23/MKLP1, whose levels are

higher after incubation with the two phosphatase inhibitors (Fig. 2C, 2D and Supplementary

Fig. S2C). This result indicates that KIF23/MKLP1 dephosphorylation might affect its stability,

perhaps by affecting its interaction with other midbody proteins. This is in line with our

previous observation that this kinesin is a PP1 substrate and PP1-mediated dephosphorylation

regulates its interaction with other midbody proteins, including PRC1 [11].

Together, these findings indicate that, in whole cells, the levels of most midbody proteins

decrease during mitotic exit, with the exception of some 'stable' midbody proteins such as

KIF14 and KIF20A. In addition, drug treatments indicate that this may be regulated by both

ubiquitin-mediated proteasome degradation and phosphorylation/de-phosphorylation.

3.3 Midbody proteins display distinct dynamics during mitotic exit and cytokinesis

Our western blot analyses only detected the total amount of midbody proteins in the cell.

Therefore, we next employed time-lapse microscopy to gain more detailed information about

the spatial and temporal dynamics of midbody proteins during mitotic exit and cytokinesis.

We selected cell lines stably expressing GFP-tagged versions of four midbody proteins

representative of the different profiles and responses to drug treatments observed in the

western blot experiments (Fig. 2): AURKB, CIT-K, KIF23/MKLP1 and PRC1 [11]. Cells were also

incubated with a DNA dye to identify cells in mitosis and to simultaneously visualize the

dynamics of chromosomes and midbody proteins. For each GFP-tagged protein, we also

measured fluorescence levels in the whole cell and at the central spindle/cleavage
furrow/midbody. AURKB and PRC1 showed very similar dynamics, as they initially
accumulated at the midbody, but then their levels slowly decreased and low fluorescence
signals were observed at the midbody 1 hour after anaphase onset (Fig. 3A, 3B and 3E; Videos
S1 and S2). Notably, the decrease observed at the central spindle/midbody was more rapid
than that observed in whole cells (Fig. 3E). By contrast, CIT-K and KIF23/MKLP1, after the initial
accumulation to the either the cleavage furrow (CIT-K) or central spindle (MKLP1), persisted
at the midbody for much longer, up to 2 hours after anaphase onset and even in post-mitotic
midbody (Fig. 3C-E and Videos S3 and S4).
These results indicate that midbody proteins can be divided into two distinct classes based on
their dynamics profiles: 'transient' midbody proteins, such as AURKB and PRC1, which after
an initial accumulation at the midbody then slowly dissipate; and 'stable' midbody proteins
that instead persists at the midbody both during and after cytokinesis
289
3.4 Transient and stable midbody proteins display distinct interactions with ubiquitylation
factors
Our findings indicated that transient and stable midbody proteins have different expression
and dynamics profiles (Figs, 2 and 3) and that the levels of some proteins are regulated by
proteasome-mediated degradation (Fig. 2). Therefore, as a first step to understand if and how
ubiquitylation might be involved in regulating midbody protein dynamics and/or stability, we
generated a midbody ubiquitylation protein-protein interaction network (interactome) by
extracting from our previously published midbody interactome dataset [11] proteins whose
full names and/or gene ontology (GO) terms in the fields biological process, cellular
component and molecular function, contained the term "ubiquitin" (Fig. 4 A, Table S1). From
this midbody interactome, we then generated two distinct sub-networks using either
'transient' or 'stable' midbody proteins as baits (Fig. 4B-D). We decided to include CIT-K among
the stable proteins, despite the evidence that its levels decline during mitotic exit (Fig. 2A, 2B)
and are slightly stabilized by MG132 (Fig. 2C and 2D), because time-lapse imaging clearly
indicated that CIT-K stably localized to the midbody during and after cytokinesis (Fig. 3C and
3E). Although the two networks showed considerable overlap (29 out of 76 proteins; Fig. 4B),
they also had distinct interactors (Fig. 4B-D). This would suggest that ubiquitylation of the two
different classes of midbody proteins could be differentially regulated, by specific ubiquitin
ligases and/or deubiquitinating enzymes (deubiquitinases, DUBs). Our observation that CIT-K

persists at the midbody, while its cellular levels decline during mitotic exit and are slightly

stabilized by proteasome inhibition, is consistent with the possibility that one or more DUBs

might be involved in preventing CIT-K degradation specifically at the midbody. Analysis of the

transient and stable midbody ubiquitylation interactomes showed that some DUBs specifically

associated with either transient or stable proteins (Table 1). In particular, USP7 and OTUD4

were pulled down only by KIF23/MKLP1 and USP10 associated specifically with the two close

partners CIT-K and KIF14 [16,26,34]. KIF14 was the only bait that pulled down USP36 and USP9X.

The three DUBs specifically found in the transient midbody ubiquitylation interactome,

UCHL1, USP4 and USP54 all associated with KIF20A/MKLP2, and UCHL1 also interacted with

PRC1. It is noteworthy that some of these DUBs have already been implicated in the regulation

of mitosis. For example, USP7 has been reported to regulate the spindle assembly checkpoint

component BUB3 and the three mitotic kinases Aurora A, CDK1, and PLK1 [35-38]. Moreover,

USP9X has been shown to antagonize the APC/C and to regulate the localization of the CPC

component Survivin [39,40].

In conclusion, our analysis suggests that these specific interactions between DUBs and stable

or transient midbody proteins could potentially explain, at least in part, the different dynamics

and stability of these midbody proteins during cytokinesis. Future studies aimed at defining

the molecular mechanisms underpinning the stability and dynamics of midbody proteins will

undoubtedly help us to understand how this organelle regulates abscission and other

important biological processes, including cell fate, cell proliferation, tissue architecture and

brain development.

Supplementary Materials: The following supporting information can be downloaded at:

www.mdpi.com/xxx/s1, Figure S1: title; Figure S2, Table S1: title; Video S1: time-lapse analysis

of AURKB dynamics during cytokinesis; Video S2: time-lapse analysis of PRC1 dynamics during

cytokinesis;. Video S3: time-lapse analysis of CIT-K dynamics during cytokinesis; Video S4:

time-lapse analysis of KIF23/MKLP1 dynamics during cytokinesis.

Author Contributions: Conceptualization, P.P.D., investigation, A.D., E.F.J.H., R.M. and P.P.D.;

software, A.E.; formal analysis, A.E. and P.P.D.; writing—original draft preparation, P.P.D.;

writing—review and editing, E.F.J.H. and P.P.D.; visualization E.F.J.H., R.M. and P.P.D.,

supervision, P.P.D.; project administration, P.P.D.; funding acquisition, P.P.D. All authors have

read and agreed to the published version of the manuscript.

Funding: This research was funded by a BBSRC grant to PPD (ref: BB/R001227/1). EFJH is supported by a BBSRC DTP studentship and RM was supported through an exchange program between the Collegio Ghislieri (University of Pavia, Italy) and St. John's College (University of Cambridge, UK). AD was supported by an Erasmus traineeship program from the University of Bologna (Italy).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: We are very grateful to P.A. Coelho for help with time-lapse experiments.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Cuijpers, S.A.G.; Vertegaal, A.C.O. Guiding Mitotic Progression by Crosstalk between Post-translational Modifications. *Trends in biochemical sciences* **2018**, *43*, 251-268, doi:10.1016/j.tibs.2018.02.004.
2. Wieser, S.; Pines, J. The biochemistry of mitosis. *Cold Spring Harbor perspectives in biology* **2015**, *7*, a015776, doi:10.1101/cshperspect.a015776.
3. Lindon, C. Control of mitotic exit and cytokinesis by the APC/C. *Biochem Soc Trans* **2008**, *36*, 405-410, doi:10.1042/BST0360405.
4. Holder, J.; Poser, E.; Barr, F.A. Getting out of mitosis: spatial and temporal control of mitotic exit and cytokinesis by PP1 and PP2A. *FEBS Lett* **2019**, *593*, 2908-2924, doi:10.1002/1873-3468.13595.
5. Carmena, M.; Wheelock, M.; Funabiki, H.; Earnshaw, W.C. The chromosomal passenger complex (CPC): from easy rider to the godfather of mitosis. *Nat Rev Mol Cell Biol* **2012**, *13*, 789-803, doi:10.1038/nrm3474.
6. D'Avino, P.P.; Capalbo, L. New Auroras on the Roles of the Chromosomal Passenger Complex in Cytokinesis: Implications for Cancer Therapies. *Front Oncol* **2015**, *5*, 221, doi:10.3389/fonc.2015.00221.
7. Nasa, I.; Kettenbach, A.N. Coordination of Protein Kinase and Phosphoprotein Phosphatase Activities in Mitosis. *Front Cell Dev Biol* **2018**, *6*, 30, doi:10.3389/fcell.2018.00030.
8. Archambault, V.; Glover, D.M. Polo-like kinases: conservation and divergence in their functions and regulation. *Nat Rev Mol Cell Biol* **2009**, *10*, 265-275, doi:10.1038/nrm2653.

9.   Fededa, J.P.; Gerlich, D.W. Molecular control of animal cell cytokinesis. *Nat Cell Biol*
2012, *14*, 440-447, doi:10.1038/ncb2482.
10.   D'Avino, P.P.; Giansanti, M.G.; Petronczki, M. Cytokinesis in animal cells. *Cold Spring*
Harbor perspectives in biology **2015**, *7*, a015834, doi:10.1101/cshperspect.a015834.
11.   Capalbo, L.; Bassi, Z.I.; Geymonat, M.; Todesca, S.; Copoiu, L.; Enright, A.J.; Callaini,
G.; Riparbelli, M.G.; Yu, L.; Choudhary, J.S.; et al. The midbody interactome reveals
unexpected roles for PP1 phosphatases in cytokinesis. *Nature communications* **2019**,
10, 4513, doi:10.1038/s41467-019-12507-9.
12.   Skop, A.R.; Liu, H.; Yates, J., 3rd; Meyer, B.J.; Heald, R. Dissection of the mammalian
midbody proteome reveals conserved cytokinesis mechanisms. *Science* **2004**, *305*,
61-66, doi:10.1126/science.1097931.
13.   D'Avino, P.P.; Capalbo, L. Regulation of midbody formation and function by mitotic
kinases. *Semin Cell Dev Biol* **2016**, *53*, 57-63, doi:10.1016/j.semcdb.2016.01.018.
14.   Mierzwa, B.; Gerlich, D.W. Cytokinetic abscission: molecular mechanisms and
temporal control. *Dev Cell* **2014**, *31*, 525-538, doi:10.1016/j.devcel.2014.11.006.
15.   Hu, C.K.; Coughlin, M.; Mitchison, T.J. Midbody assembly and its regulation during
cytokinesis. *Mol Biol Cell* **2012**, *23*, 1024-1034, doi:10.1091/mbc.E11-08-0721.
16.   Bassi, Z.I.; Audusseau, M.; Riparbelli, M.G.; Callaini, G.; D'Avino, P.P. Citron kinase
controls a molecular network required for midbody formation in cytokinesis. *Proc*
Natl Acad Sci U S A **2013**, *110*, 9782-9787, doi:10.1073/pnas.1301328110.
17.   D'Avino, P.P. Citron kinase - renaissance of a neglected mitotic kinase. *J Cell Sci* **2017**,
130, 1701-1708, doi:10.1242/jcs.200253.
18.   McKenzie, C.; Bassi, Z.I.; Debski, J.; Gottardo, M.; Callaini, G.; Dadlez, M.; D'Avino, P.P.
Cross-regulation between Aurora B and Citron kinase controls midbody architecture
in cytokinesis. *Open Biol* **2016**, *6*, 160019, doi:10.1098/rsob.160019.
19.   Crowell, E.F.; Gaffuri, A.L.; Gayraud-Morel, B.; Tajbakhsh, S.; Echard, A. Engulfment of
the midbody remnant after cytokinesis in mammalian cells. *J Cell Sci* **2014**, *127*, 3840-
3851, doi:10.1242/jcs.154732.
20.   Peterman, E.; Prekeris, R. The postmitotic midbody: Regulating polarity, stemness,
and proliferation. *J Cell Biol* **2019**, *218*, 3903-3911, doi:10.1083/jcb.201906148.
21.   Siskos, N.; Stylianopoulou, E.; Skavdis, G.; Grigoriou, M.E. Molecular Genetics of
Microcephaly Primary Hereditary: An Overview. *Brain Sci* **2021**, *11*,
doi:10.3390/brainsci11050581.
22.   Schindelin, J.; Arganda-Carreras, I.; Frise, E.; Kaynig, V.; Longair, M.; Pietzsch, T.;
Preibisch, S.; Rueden, C.; Saalfeld, S.; Schmid, B.; et al. Fiji: an open-source platform
for biological-image analysis. *Nat Methods* **2012**, *9*, 676-682,
doi:10.1038/nmeth.2019.
23.   Hummer, S.; Mayer, T.U. Cdk1 negatively regulates midzone localization of the
mitotic kinesin Mklp2 and the chromosomal passenger complex. *Curr Biol* **2009**, *19*,
607-612, doi:10.1016/j.cub.2009.02.046.
24.   Renshaw, M.J.; Liu, J.; Lavoie, B.D.; Wilde, A. Anillin-dependent organization of septin
filaments promotes intercellular bridge elongation and Chmp4B targeting to the
abscission site. *Open Biol* **2014**, *4*, 130190, doi:10.1098/rsob.130190.
25.   Gai, M.; Camera, P.; Dema, A.; Bianchi, F.; Berto, G.; Scarpa, E.; Germena, G.; Di
Cunto, F. Citron kinase controls abscission through RhoA and anillin. *Mol Biol Cell*
2011, *22*, 3768-3778, doi:mbc.E10-12-0952 [pii], 10.1091/mbc.E10-12-0952.
26.   Gruneberg, U.; Neef, R.; Li, X.; Chan, E.H.; Chalamalasetty, R.B.; Nigg, E.A.; Barr, F.A.
KIF14 and citron kinase act together to promote efficient cytokinesis. *J Cell Biol* **2006**,
172, 363-372.

27.   Scott, S.J.; Suvarna, K.S.; D'Avino, P.P. Synchronization of human retinal pigment
ephitilial-1 (RPE-1) cells in mitosis. *J Cell Sci* **2020**, doi:10.1242/jcs.247940.
28.   Mitsuhashi, S.; Matsuura, N.; Ubukata, M.; Oikawa, H.; Shima, H.; Kikuchi, K.
Tautomycetin is a novel and specific inhibitor of serine/threonine protein
phosphatase type 1, PP1. *Biochem Biophys Res Commun* **2001**, *287*, 328-331,
doi:10.1006/bbrc.2001.5596.
29.   Ditchfield, C.; Johnson, V.L.; Tighe, A.; Ellston, R.; Haworth, C.; Johnson, T.; Mortlock,
A.; Keen, N.; Taylor, S.S. Aurora B couples chromosome alignment with anaphase by
targeting BubR1, Mad2, and Cenp-E to kinetochores. *J Cell Biol* **2003**, *161*, 267-280,
doi:10.1083/jcb.200208091, jcb.200208091 [pii].
30.   Zhao, W.M.; Fang, G. Anillin is a substrate of anaphase-promoting
complex/cyclosome (APC/C) that controls spatial contractility of myosin during late
cytokinesis. *J Biol Chem* **2005**, *280*, 33516-33524.
31.   Paccosi, E.; Costanzo, F.; Costantino, M.; Balzerano, A.; Monteonofrio, L.; Soddu, S.;
Prantera, G.; Brancorsini, S.; Egly, J.M.; Proietti-De-Santis, L. The Cockayne syndrome
group A and B proteins are part of a ubiquitin-proteasome degradation complex
regulating cell division. *Proc Natl Acad Sci U S A* **2020**, *117*, 30498-30508,
doi:10.1073/pnas.2006543117.
32.   Glotzer, M.; Murray, A.W.; Kirschner, M.W. Cyclin is degraded by the ubiquitin
pathway. *Nature* **1991**, *349*, 132-138, doi:10.1038/349132a0.
33.   Floyd, S.; Pines, J.; Lindon, C. APC/C Cdh1 targets aurora kinase to control
reorganization of the mitotic spindle at anaphase. *Curr Biol* **2008**, *18*, 1649-1658.
34.   Watanabe, S.; De Zan, T.; Ishizaki, T.; Narumiya, S. Citron kinase mediates transition
from constriction to abscission through its coiled-coil domain. *J Cell Sci* **2013**, *126*,
1773-1784, doi:10.1242/jcs.116608.
35.   Galarreta, A.; Valledor, P.; Ubieto-Capella, P.; Lafarga, V.; Zarzuela, E.; Munoz, J.;
Malumbres, M.; Lecona, E.; Fernandez-Capetillo, O. USP7 limits CDK1 activity
throughout the cell cycle. *EMBO J* **2021**, *40*, e99692, doi:10.15252/embj.201899692.
36.   Giovinazzi, S.; Morozov, V.M.; Summers, M.K.; Reinhold, W.C.; Ishov, A.M. USP7 and
Daxx regulate mitosis progression and taxane sensitivity by affecting stability of
Aurora-A kinase. *Cell Death Differ* **2013**, *20*, 721-731, doi:10.1038/cdd.2012.169.
37.   Giovinazzi, S.; Sirleto, P.; Aksenova, V.; Morozov, V.M.; Zori, R.; Reinhold, W.C.; Ishov,
A.M. Usp7 protects genomic stability by regulating Bub3. *Oncotarget* **2014**, *5*, 3728-
3742, doi:10.18632/oncotarget.1989.
38.   Peng, Y.; Liu, Y.; Gao, Y.; Yuan, B.; Qi, X.; Fu, Y.; Zhu, Q.; Cao, T.; Zhang, S.; Yin, L.; et
al. USP7 is a novel Deubiquitinase sustaining PLK1 protein stability and regulating
chromosome alignment in mitosis. *J Exp Clin Cancer Res* **2019**, *38*, 468,
doi:10.1186/s13046-019-1457-8.
39.   Skowyra, A.; Allan, L.A.; Saurin, A.T.; Clarke, P.R. USP9X Limits Mitotic Checkpoint
Complex Turnover to Strengthen the Spindle Assembly Checkpoint and Guard against
Chromosomal Instability. *Cell Rep* **2018**, *23*, 852-865,
doi:10.1016/j.celrep.2018.03.100.
40.   Vong, Q.P.; Cao, K.; Li, H.Y.; Iglesias, P.A.; Zheng, Y. Chromosome alignment and
segregation regulated by ubiquitination of survivin. *Science* **2005**, *310*, 1499-1504,
doi:10.1126/science.1120160.
477

478 **Table 1**. List of deubiquitinases (DUB) identified in the midbody ubiquitylation interactome
479

Group	*Gene Name*	*Protein Name*	*Baits (MASCOT score)*	*Midbody proteome*
Common	VCP	Transitional endoplasmic reticulum ATPase (TER ATPase)	ANLN (84); AURKB (174); KIF14 (76); PRC1 (158)	Yes
	USP5	Ubiquitin carboxyl-terminal hydrolase 5	MKLP1 (41); PRC1 (83)	No
	EIF3F	Eukaryotic translation initiation factor 3 subunit F	ANLN (61); CIT-K (79); KIF14 (59); MKLP1 (73); PRC1 (136)	Yes
	USP9Y	Probable ubiquitin carboxyl-terminal hydrolase FAF-Y	KIF14 (42); KIF20A (36); MKLP1 (68)	No
	USP39	U4/U6.U5 tri-snRNP-associated protein 2	AURKB (36); CIT-K (300); PRC1 (156)	Yes
Transient-specific	UCHL1	Ubiquitin carboxyl-terminal hydrolase isozyme L1	KIF20A (30); PRC1 (392)	No
	USP54	Inactive ubiquitin carboxyl-terminal hydrolase 54	KIF20A (104)	No
	USP4	Ubiquitin carboxyl-terminal hydrolase 4	KIF20A (40)	No
Stable-specific	OTUD4	OTU domain-containing protein 4	MKLP1 (38)	No
	USP7	Ubiquitin carboxyl-terminal hydrolase 7	MKLP1 (51)	Yes
	USP10	Ubiquitin carboxyl-terminal hydrolase 10	CIT-K (59); KIF14 (126)	Yes
	USP36	Ubiquitin carboxyl-terminal hydrolase 36	KIF14 (139)	No
	USP9X	Probable ubiquitin carboxyl-terminal hydrolase FAF-X	KIF14 (50);	No

Figure 1. Changes in midbody protein distribution during its maturation. (A) Schematic
diagram of the midbody showing its different regions and the localization of the proteins
analyzed in this study. (B and C) HeLa cells expressing ANLN::GFP were fixed and stained to
detect GFP (green in the merged panels) Aurora B (blue in the merged panels) and either CIT-
K (red in the merged panel in B) or KIF23/MKLP1 (red in the merged panels in C). Insets show
a 3x magnification of the midbody. Bars, 10 μm.
488

Figure 2. Different expression profiles of midbody proteins during mitotic exit. (A) Time course
analysis of midbody protein expression during mitotic exit. HeLa cells were synchronized by
thymidine/nocodazole block and then collected at the indicate time points after nocodazole
release. Proteins were extracted and used in Western blot analysis to identify the antigens
indicated to the right. The numbers on the left indicate the sizes of the molecular mass marker.
(B) Graph showing the quantification of protein levels, normalized to tubulin and relative to
levels at time 0 min, from at least two different western blots like the one shown in (A) using
protein extracts from two separate experiments. (C) Effect of different inhibitors on midbody
protein levels. HeLa cells were synchronized by thymidine/nocodazole block, released for 45
min in fresh medium, and then incubated for further 60 min in MG132, okadaic acid (OKA),
tautomycetin (TAUT), ZM447439 (ZM), or the solvent DMSO as control. Proteins were
extracted and used in Western blot analysis to identify the antigens indicated to the right. The
numbers on the left indicate the sizes of the molecular mass marker. (D) Graph showing the
quantification of protein levels, normalized to tubulin and relative to DMSO levels, from at
least two different western blots like the one shown in (C) using protein extracts from two
separate experiments.

507
508
Figure 3. Midbody proteins display distinct dynamics during cytokinesis. (A-D) Selected images
from time-lapse recordings of HeLa cells expressing GFP-tagged AURKB (A), PRC1 (B), CIT-K (C)
and KIF23/MKLP1 (D). Chromosomes were visualised using SiR-DNA. GFP-tagged proteins are
in cyan and DNA in magenta in the merged panels. Time is in minutes relative to anaphase
onset (0 time point). The arrowheads in (C) marks CIT-K::GFP localization to post-mitotic
midbodies. Bar, 10 μm. (E) Quantification of midbody proteins during cytokinesis.
Fluorescence intensity values were measured in whole cells and at the central
spindle/cleavage furrow/midbody and then normalized relative to the anaphase onset (0)
time point. Bars indicate SEM, n=10.
518

519
Figure 4. (A) Diagram illustrating the midbody ubiquitylation interactome. Baits are indicated
with blue hexagons, while preys are represented as ovals, either in green, if they were also
found in the midbody proteome, or in orange. The edges connecting the network nodes are
colored according to their Mascot scores as indicated in the color scale bar at the bottom left
(see also [11]). Preys shared by multiple baits are clustered in the center. (B) Proportional Venn
diagram showing the number of proteins present in the transient and stable midbody
ubiquitylation interactomes. (C-D) Diagrams illustrating the midbody ubiquitylation
interactomes generated using either transient (C) or stable (D) midbody proteins as baits.
Baits, preys and edges are as in (A).