Engineering *Escherichia coli* to utilize erythritol as sole carbon source

Fang Ba, Xiangyang Ji, Shuhui Huang, Yufei Zhang, Wan-Qiu Liu and Jian Li[*]

Affiliations:

School of Physical Science and Technology, ShanghaiTech University, Shanghai, 201210, China

*Corresponding author. E-mail: lijian@shanghaitech.edu.cn (J.L.)

Abstract

Erythritol, one of the natural sugar alcohols, is widely used as a sugar substitute sweetener in food industries. Humans and their native gut microbiota lack erythritol catabolic microorganisms to utilize erythritol as carbon source, which inspires us to engineer a model chassis that can respond and utilize erythritol for establishing wide applications in synthetic biology fields. Here, we successfully engineered *Escherichia coli* to respond and utilize erythritol as sole carbon source. First, a heterologous erythritol catabolic gene cluster from *Ochrobactrum spp.* enables *E. coli* to catabolize erythritol into erythrose-4-phosphate and flow into pentose-phosphate pathway. Second, we characterized an erythritol binding transcriptional repressor and its DNA binding site. Then, mRNA transcriptional analysis guided us to build up further metabolic engineering strategies for improving erythritol catabolism. Finally, such engineered *E. coli* strains were applied for *in vitro* simulated intestinal fluid experiments and performed as an erythritol detector to distinguish soda drinks. We anticipate erythritol catabolic *E. coli* strains will expand synthetic biology fields and accelerate applications in biomedical engineering and living therapeutics.

Introduction

The worldwide obesity epidemic and correlative diseases like diabetes, hypertension, and hyperlipidemia encourage people to change their lifestyles by limiting the excessive intake of carbohydrates. Food industries meet people's requirements to produce a series of carbohydrate-reduced foods with sugar alcohols (also called polyols) as additional sweeteners[1]. These compounds contain multiple hydroxyl groups and are usually produced by industrial carbohydrates hydrogenation[2]. Erythritol, the C4 sugar alcohol, naturally exists in small species of vegetables, fruits, mushrooms, and fermented foods[3,4]. It also can be found in human tissues such as semen, lens, cerebrospinal fluid, and serum[1,4,5]. Erythritol industrial manufacturing has been well developed by fermentation processes with engineered lactic acid bacteria and yeast[6,7,8,9,10]. Moreover, erythritol is well-characterized with high stability in a wide range of temperature and pH conditions. Also, it does not take part in Maillard reactions and works as a tremendous non-hygroscopic substance[11]. In general, erythritol works as a stable, cheap, and safe sugar alcohol as a natural food sweetener.

The reason why erythritol is used as a food sweetener because human beings cannot catabolize it. It has been reported that most erythritol can be rapidly absorbed in the small intestine and excreted in urine within 24 h[5,11,12,13,14]. Only a small percent of erythritol flow through intestinal fermentation[13]. However, the human gut microbiota lacks erythritol catabolic microorganisms. Thus, nearly all erythritol cannot be utilized by human beings as energy substance[15,16]. From another aspect, erythritol, such a human-compatible compound, can work as a potential energy substrate by engineered microorganisms that utilize erythritol as energy source, or work as a unique micromolecular compound to regulate human physiological

metabolism.

Up to now, only two distinguishing erythritol degradation pathways have been identified

from microorganisms: The first one was discovered in several *Hyphomicrobiales* species,

including environmental bacterium *Ochrobactrum*[17], pathogenic bacterium

Brucella[18,19,20,21,22,23], nitrogen-fixing plant endosymbionts *Rhizobium*[22], and *Sinorhizobium*[24].

These microorganisms commonly uptake erythritol as the initial substrate and yield erythrose-

4-phosphate via 5-step catalysis[17]. Erythrose-4-phosphate is the intermediate product of

pentose-phosphate pathway and can be utilized as a C4 carbon source. The second pathway was

discovered in *Mycolicibacterium smegmatis*[25]. This pathway is simple 3-step catalysis, and the

final product is also erythrose-4-phosphate. In general, most other organisms are not able to

utilize erythritol as energy source. Meanwhile, synthetic biology takes the core mission of

(re)designing and constructing new biological parts, circuits, and systems for useful purposes.

As such goals inspired, we purposefully aimed to reprogram a general model microorganism to

catabolize erythritol as energy source for potential usage.

To meet this demand, we chose *Escherichia coli*, one of the most used model prokaryote

and GRAS (generally recognized as safe) microorganisms as chassis[26,27]. Afterward, we

separated and identified an environmental bacterium *Ochrobactrum spp.* from outdoor aerosol

and gained an erythritol catabolic gene cluster. Such a 5-step reaction provides *E. coli* capability

to utilize erythritol as sole carbon source. Meanwhile, we characterized the erythritol-responded

genetic repressor eryD and its DNA-binding site. Furthermore, we did mRNA transcriptional

analysis to identify the metabolism of such C4 carbon flux. Relying on the transcriptome results,

we designed genetic circuits to facilitate erythritol catabolism and significantly increased the *E.*

coli growth in M9-erythritol medium. For developing applications of such engineered erythritol catabolic microorganism, probiotic *E. coli* Nissle 1917 (EcN) were designed to utilize erythritol, and such catabolism promoted EcN to utilize erythritol in simulated intestinal fluid (SIF) for growth. Finally, our engineered *E. coli* could be used for distinguishing soda drinks whether they contain high-fructose corn syrup (HFCS, which contains sugar) or erythritol (sugar-free sweetener). We anticipate that our erythritol catabolic *E. coli* strains will accelerate synthetic biology with broad potential research and applications such as carbon cycle, agriculture, genetic circuit design, metabolic engineering, biomedical engineering, and living therapeutics.

Results

Engineered *E. coli* can utilize erythritol as sole carbon source

To gain reliable erythritol catabolic microorganisms from the natural environment, we designed an erythritol-based selective medium derived from M9 minimal medium, one of the chemically defined mediums for *E. coli* incubation. Here we proposed that such erythritol-based selective medium would separate *E. coli*-like gram-negative bacteria that are suitable for further exploration. Hence, we replaced 0.4% glucose (C6 carbon source in M9 minimal medium) with 0.4% erythritol and called it as M9-erythritol medium, while the other components were not changed. After this process, we successfully separated, identified, and characterized an environmental bacterium *Ochrobactrum spp.* from campus aerosol (**Supplementary Fig. 1** and **2**)[28]. Then, ten erythritol catabolism-associated genes were isolated and sequenced from *Ochrobactrum spp.* genome (**Supplementary Fig. 3-5**), DNA sequence alignments suggested that all the genes were almost the same as reported of *Brucella abortus 2308*[29,30], including erythritol catabolic genes eryA, eryB, eryC, eryH, eryI, putative erythritol binding transcriptional repressor eryD, eryR, and erythritol ABC-transporter genes eryE, eryF, eryG (**Fig. 1a**). Thibault *et al.* has been characterized the 5-step erythritol catabolism proteins *in vitro*, which started with an erythritol kinase eryA to produce D-erythritol 1-phosphate, then the phosphorylated intermediate was oxidized by a membrane FAD-dependent dehydrogenase eryB and yielding D-erythrulose 1-phosphate, which would next be isomerized to L-erythrulose 1-phosphate by an isomerase eryC. Next, another isomerase eryH could catalyze phosphate transportation to produce D-erythrulose 4-phosphate, finally, isomerase eryI converted D-erythrulose 4-phosphate to D-erythrose 4-phosphate, which was one of the intermediates in

pentose-phosphate pathway[17]. Such C4 phosphate-substrate could be utilized for carbohydrate

utilization via glycolysis, nucleotide synthesis, and aromatic amino acids biosynthesis via

shikimate pathway (**Fig. 1b**).

Thibault *et al.* has been purified all five erythritol catabolism proteins from *E. coli*-based

protein expression system and successfully characterized their catalytic functions *in vitro*. On

this account, we prospected that the original gene cluster might be functional for *E. coli*

erythritol catabolism *in vivo*. Thus, we initially constructed a plasmid containing erythritol

catabolism genes and transformed it into *E. coli* MG1655 for testing whether engineered *E. coli*

strain could utilize erythritol as carbon source. Surprisingly, such *E. coli* could slowly grow in

liquid M9-erythritol medium (**Fig. 1c**), within 72 h incubation, *E. coli* grew as an "S" curve and

the OD_{600} value finally reached approximately 0.35. To assuredly determine the erythritol

consumption, filtered medium supernatant was collected for high performance liquid

chromatography (HPLC) analysis, results from the refractive index detector (RID) indicated

that during incubation, erythritol was certainly consumed by engineered *E. coli* as carbon

source (**Fig. 1d**).

After simply demonstrating erythritol catabolism *in vivo*, we next characterized this gene

cluster via other aspects. Firstly, the *in vivo* expression of ten proteins was analyzed by SDS-

PAGE and Western-Blot (**Extended Data Fig. 1**). The results showed that except eryC, the

other proteins could be well expressed as a high percentage of soluble fractions *in vivo*[17].

Secondly, all ten genes' promoter and ribosome binding site (RBS) strength were measured

with standardization (**Extended Data Fig. 2** and **3**)[31,32,33]. Thirdly, three erythritol ABC-

transporter proteins eryE, eryF, and eryG were predicted and demonstrated their expression,

transmembrane phenotype, and localization in *E. coli* cells (**Supplementary Fig. 6** and

Extended Data Fig. 4). It was worth noting that *E. coli* MG1655 could transport and catabolize

erythritol without such ABC-transporter system, which means that *E. coli* might carry native

isoenzymes for erythritol transport.

EryD is an erythritol binding transcriptional repressor

In 2000, Sangari *et al*. reported that erythritol might act as an inducer and probably bind

to eryD in *Brucella abortus 2308* to inhibit its repressor activity[19]. However, it lacked more

detailed eryD characterization. Interestingly, our separated erythritol cluster of *Ochrobactrum*

spp. includes an eryD gene whose DNA sequence was totally aligned with eryD in *Brucella*

abortus 2308. In addition, sequence analysis indicated eryR, the other putative DNA-binding

transcriptional repressor might act similarly to eryD. Hence, characterization of such two

proteins should help us figure out their regulation mechanism and expand erythritol-associated

synthetic biology applications. To meet this demand, we initially attempted to determine the

putative DNA-binding sites within two loci: one was the 732 bp gap between eryE and eryA,

and the other is approximately 400 bp gap between eryD and eryR (**Supplementary Fig. 3**).

Here we designed an *in vivo* characterization strategy to precisely identified the DNA-binding

sites: In general, we supposed the repressor's DNA-binding site was localized after a promoter

(generally recognized as a transcriptional unit), thus we brought different putative DNA

sequences into a standardized sfGFP generation plasmid, which consisted of a strong RBS, an

sfGFP coding sequence, and a strong terminator in iGEM standard backbone pSB1C3. Firstly,

we successfully identified two promoters called pEryF-732 (F means forward) and pEryR-732

(R means reverse), which are located in the 732 bp gap between eryE and eryA

(**Supplementary Fig. 7** and **8**). Secondly, to identify the putative DNA-binding sites, other

compatible plasmids were designed to constitutively express eryD or eryR, which consisted of

gradient strength constitutive promoters, strong RBS, eryD/eryR coding sequences, and strong

terminator with compatible vector. The sfGFP fluorescence of cell pellets and Western-Blot

results showed that eryD had a DNA-binding site located in pEryF-732, and eryD expression

level could regulate transcription level, but eryR did not (**Supplementary Fig. 9**). Subsequently,

promoter 5'-region truncation experiments were performed to precisely identified the 5'-site of

eryD DNA-binding site (we called it as erythritol operator, eryO), to achieve this goal, pEryF-

732 were truncated along with the gradient and brought into sfGFP generation plasmids

(**Supplementary Fig. 10-12**). After that, we determined the front site of eryO. Similarly, 3'-

site of eryO were also successfully identified (**Supplementary Fig. 13-15**). Finally, eryO was

identified as a 29 bp sequence located between the -35 and -10 region of promoter pEryF-732

(**Fig. 2a**). To ensure the reliability of *in vivo* results, electrophoretic gel mobility shift assay

(EMSA) was performed in which the FAM-labeled DNA fragments harboring the eryE-eryA

732 bp intergenic region were incubated with an increasing amount of eryD. As expected, eryD

effectively bound to pEryF-732 promoter (**Fig. 2b**), and a further investigation showed that

eryD precisely binds to the 29 bp eryO region (**Supplementary Fig. 16**). In addition, we also

confirmed that there was no eryD or eryR DNA-binding sites in the 400 bp gap between eryD

and eryR (**Supplementary Fig. 17**). Moreover, we concentrated on the eryD protein

characterization. Structure prediction of eryD monomer indicated that eryD was assembled as

an N-terminal helix-turn-helix motif for DNA interaction and a C-terminal domain for erythritol

regulation. SWISS-MODEL prediction showed that eryD is assembled as a homo-tetramer

(**Supplementary Fig. 18**), gel-filtration chromatography and Native-PAGE-based Western-

Blot demonstrated this conclusion (**Extended Data Fig. 5**).

Next, we demonstrated that eryD could be dynamically regulated by gradient

concentration of erythritol (**Fig. 2c-f** and **Supplementary Fig. 19**). For the development of

eryD-based genetic circuit design, T7 RNA polymerase-based erythritol induction system (**Fig.**

2g, h and **Supplementary Fig. 20**), and *E. coli* σ70 factor-based native erythritol induction

systems (**Fig. 2i, j** and **Supplementary Fig. 20**) were constructed, performed, and successfully

demonstrated their function. In the end, the orthogonality of erythritol-eryD interaction was

characterized by a series of generally used carbohydrates and polyols, the results showed that

eryD could interact with erythritol with great orthogonality (**Fig. 2k, l**).

In summary, we demonstrated that eryD is an erythritol-binding transcriptional repressor,

identified its DNA-binding site eryO, demonstrated its concentration-dependent dynamic

response, excellent substrate orthogonality, and applied it to genetic circuit design.

mRNA transcriptional analysis of engineered *E. coli* erythritol catabolism

To determine erythritol catabolism in engineered *E. coli* strains, we performed comparative

mRNA transcriptional analysis on triplicate exponential phase cultures of *E. coli* MG1655 that

grew between M9-erythritol medium and M9-glucose medium (**Extended Data Fig. 6**). Totally,

1298 genes were significantly different between these two mediums, including 759 genes

upregulated, and 539 genes downregulated (**Supplementary Fig. 21**). COG annotations

showed that the significant differences were found in genes related to carbohydrate metabolism,

amino acid metabolism, energy metabolism, transcription, inorganic ion metabolism, and

ribosome-associated translation (**Extended Data Fig. 7**). It was worth noting that most of the

genes related to carbohydrate metabolism were overexpressed, which indicates that erythritol

was not efficiently utilized as carbon source for engineered *E. coli* than glucose. In addition,

nearly all ribosome-associated genes were downregulated, which means that such strain

decreased its translation level to cope with such a low-carbon-source environment. Besides,

KEGG and GO annotation showed similar conclusions.

To take a panoramic view of carbohydrate metabolic pathways of erythritol-utilized *E. coli*

MG1655, transcriptomic data were mapped in a schematic diagram including heterologous

erythritol degradation pathway, pentose-phosphate pathway, glycolysis, and TCA cycle (**Fig.**

3). The end product of erythritol catabolism erythrose-4-phosphate can flow into pentose-

phosphate pathway via two pathways: one way is transaldolases talA and talB convert

erythrose-4-phosphate and β-fructose-6-phosphate into glyceraldehyde-3-phosphate and

sedoheptulose-7-phosphate, another way is transketolase tktA and tktB convert erythrose-4-

phosphate and xylulose-5-phosphate into β-fructose-6-phosphate and glyceraldehyde-3-

phosphate. Transcriptome showed that talA, talB, tktA, and tktB were overexpressed at 5.2, 2.0,

1.3, and 5.9 times, which provided metabolic engineering strategies for erythritol catabolism

improvement. Several carbohydrate degradation pathways were upregulated, including D-

xylose, L-arabinose, lactose, L-rhamnose, and glycerol. However, only galactose and D-

mannose degradation pathways were downregulated. In general, the transcriptome comparison

between erythritol degradation and glucose degradation was not significantly different in

glycolysis and TCA cycle, however, the major distinction was erythrose-4-phosphate

downstream conversion in pentose-phosphate pathway.

Another concentrated point was that both KEGG and COG annotations indicated much

downregulated ribosome-related genes (**Supplementary Information**). In detail, KEGG

annotated 0 ups and 52 downs of ribosome, and COG annotated 4 ups and 70 downs of

translation, ribosomal structure, and biogenesis. These results suggested that erythritol

utilization should cause remarkable reduction of ribosome assembly, which led to suppression

of translation and dispensable protein synthesis.

In addition, after the differentiated gene screening, some distinct upregulated genes and

gene clusters were summarized (**Supplementary Fig. 22**). We noticed that the genes of *E. coli*

glycolate utilization operon *glcCDEFGBA* were all significantly overexpressed, even glcD was

upregulated as 109 times. There was no glycolate or analogue in M9-erythritol medium, so we

considered whether there was a glc cluster transcriptional repressor that might respond to

erythritol catabolism. In this cluster, glcC is a reported DNA-binding transcriptional repressor

that responds to glycolate and acetate[34], hence we did experiments and demonstrated that glcC

could respond to erythrose-4-phosphate or its downstream substrates (**Supplementary Fig. 23**).

Furthermore, another cluster attracted our attention that the genes of phosphonate uptake and

utilization *phn* operon were nearly all upregulated (**Supplementary Fig. 22**)[35], we have

characterized that the H-T-H type transcriptional repressor phnP was not responded to erythritol

catabolism (data not shown), thus we hypothesized that erythritol utilization may request much

phosphonate because erythritol kinase eryA could catalyze erythritol phosphorylation to

produce D-erythritol 1-phosphate.

Metabolic engineering to improve *E. coli* erythritol catabolism

Up to now, we have successfully demonstrated that engineered *E. coli* can utilize erythritol as sole carbon source, and transcriptome analysis provided much more detail to inspire us on improving erythritol catabolism. As mentioned above, wild type *E. coli* MG1655 with native erythritol catabolism plasmid (pFB147) could grow to stationary phase and OD_{600} reached approximately 0.35 in 60 h (**Fig. 1d**), so we attempt to increase its growth rate and cell density. At first, we tried to screen out a better promoter to improve erythritol catabolism cluster transcription, here we chose a constitutive promoter family with gradient strength to replace native promoter pEryF-732 (**Fig. 4a**), however, there was no clear distinction between pEryF-732 and the other two medium strength promoters J23106 and J23105, interestingly, the strongest promoter J23100 and weak promoter j23114 suppressed growth, even the weakest promoter J23109 could not support growth (**Fig. 4b, c**). These results suggested that a suitable transcription level was sufficient for erythritol catabolism. Next, according to transcriptome analysis, transaldolases talA, talB, and transketolase tktA, tktB were upregulated, so we supposed that overexpression of these genes should increase consumption of erythrose-4-phosphate to accelerate carbon flux and cell growth. To demonstrate our hypothesis, we constructed three compatible plasmids that are able to constitutive express talA-tktB, talB, or tktA (**Fig. 4d, f, h**). Additional annotation is that talA is genetically located next to tktB in the genome, so here we only constructed a talA-tktB plasmid rather than two separated plasmids. Surprisingly, the OD_{600} of talA-tktB overexpressed erythritol catabolism *E. coli* could reach approximately 1.0 after 72 h culture (**Fig. 4e**) with optimal promoter J23114. Similarly, the OD_{600} of talB or tktA overexpressed erythritol catabolism *E. coli* could reach approximately 0.8

after 72 h culture (**Fig. 4g, i**). Together, after metabolic engineering, *E. coli* endogenous talA-

tktB overexpression could accelerate its erythritol utilization and cell growth (OD_{600} value

increase from 0.35 to 1.0 after 72 h culture).

Erythritol transport system characterization

As demonstrated above, wild type *E. coli* MG1655 containing erythritol catabolism

plasmid (pFB147) was able to utilize erythritol, which means that some native *E. coli* genes

may act as erythritol transporter. In *Ochrobactrum spp.*, erythritol ABC-transporter system

consisted of eryE, eryF, and eryG[22]. We demonstrated their transmembrane phenotype in *E. coli*

Mach1-T1 cells (**Extended Data Fig. 4**), however, when we tried to bring the original cluster

into different *E. coli* strains, we found that erythritol ABC-transporter may disturb cell division

(**Supplementary Fig. 24**). Therefore, we focused on native genes in *E. coli* MG1655 that assist

erythritol transport.

One way was to find out homologies of erythritol ABC-transporter (**Extended Data Fig.**

8), we firstly summarized all carbohydrate or polyol transport genes in *E. coli* MG1655, then

BLAST protein-protein alignments were proceeded to screen out homologies of erythritol

permease eryG[36,37]. Finally, seven eryG homologous permeases were screened out, including

glycerol facilitator glpF, which has been reported to enable *E. coli* to transport erythritol[38].

Subsequently, we transform pFB147 into seven knock-out strains to characterize whether these

genes could facilitate erythritol transport. Statistical analysis showed that six sugar-ABC

transporter systems might facilitate erythritol transport (**Extended Data Fig. 9**). Another way

was to rely on the transcriptome result. We summarized 34 upregulated putative carbohydrate

transport genes, constructed knock-out strains, and performed the same experiments, however,

statistical analysis showed that there were no putative systems to facilitate erythritol transport

(**Supplementary Fig. 25**). Together, these results demonstrated that part of *E. coli* native

carbohydrate transport systems might facilitate erythritol transport.

Erythritol can facilitate growth of engineered probiotic *E. coli* Nissle 1917 in simulated

intestinal fluid

Following the demonstration, we subsequently attempted to develop and expand its

practical applications. Erythritol is one of the noteworthy and widely-used sweeteners in daily

foods and drinks, and it also can briefly remain in the human GI tract without digestion, hence

we supposed that a kind of simply engineered probiotic which could respond and utilize

erythritol might adjust and balance the gut microbiome. Here we focused on the probiotic

chassis *E. coli* Nissle 1917 (EcN) which has been widely characterized, engineered, and applied

for diagnosis and therapy[27,39,40,41,42]. Firstly, erythritol catabolism plasmid pFB147 was

successfully transformed into EcN and endowed its erythritol utilization (**Supplementary Fig.**

24). Next, such engineered EcN was mixed with simulated intestinal fluid (SIF) in centrifuge

tubes to establish *in vitro* anaerobic culture condition at 37°C. After days of incubation, EcN

cell density could be calculated by counting colony number on LB-agar plates, these colonies

were separated from diluted cell culture (**Fig. 5a**). There were two facts for determination: the

initial cell density and erythritol concentration. At the start, we decided to generate a gradient

cell density of 10^2, 10^4, 10^6, and 10^8 (**Fig. 5b-e**), with 0.4% erythritol (as the same concentration

as M9-erythritol medium). The growth curves showed that engineered EcN could reach a higher

cell density than various initial cell densities. 10^2 to 10^6 incubations could reach the highest cell

density on day 2, however, 10^8 was on day 1 and declined after day 2, which suggested that

high cell density may excessively consume SIF resources. We noticed that the growth curves

between 10^4 and 10^6 were exactly similar, so we decided to use the lower density (10^4 EcN) for

further experiments. Next, six different erythritol concentrations (0%, 0.01%, 0.1%, 1%, 4%,

and 16%) were used in SIF for incubation (**Fig. 5f-k**), we found that 16% erythritol might limit

cell growth probably due to high osmosis pressure. The other five erythritol concentrations

more or less facilitate engineered EcN growth, the highest cell density reached approximately

5×10^9 on day 1 with 0.01% erythritol, and 0.1% erythritol could support EcN alive up to

approximately 6×10^8 on day 3. Nonetheless, 0% additional erythritol could also facilitate

engineered EcN growth, this was probably due to the remained erythritol in bile salts[43].

According to the mentioned incubations, the optimal condition for engineered EcN growth is

SIF with additional 0.1% erythritol and 10^4 initial cell density. In summary, we demonstrated

that erythritol could facilitate erythritol catabolic engineered EcN growth in SIF.

Engineered *E. coli* can distinguish soda drinks

To further expand the applications of erythritol-utilized *E. coli*, we are supposed to build

up a strategy for the distinction between soda drinks. In general, there are three types of sweet

soda drinks: the first one is sugary drinks that generally contain high-fructose corn syrup

(including fructose and glucose) and white sugar (sucrose). The second one is sugar-free drinks

with artificial sweeteners, including acesulfame potassium, sucralose, aspartame, saccharin,

and neotame. The third one is sugar-free drinks with natural sweeteners, including polyols

(erythritol, xylitol, isomaltol, mannitol, sorbitol, allulose, and so on), and other naturally

derived components like stevia and monk fruit sweetener. In general, people cannot

differentiate sweet sodas by tasting, however, the analysis of the specific components relies on

biochemistry analysis by HPLC, GC-MS, and other strategies, which is complicated, costly,

and time-consuming[44]. However, our engineered *E. coli* strain may perform as a sweet soda

detector. Theoretically, wild type *E. coli* strains can utilize fructose and glucose as carbon

sources, and they lack catabolic pathways to utilize sucrose, artificial sweeteners, and some

natural sweeteners like erythritol (**Supplementary Fig. 26**). To summarize, we decided to

distinguish different sweet sodas by engineered *E. coli* strains.

Here, we designed two different *E. coli* strains: one was called Negative Control (NC) due

to it consisted of a weak promoter (J23109) that could not support erythritol catabolism, the

other one was called Erythritol Detector (ED), which could utilize erythritol (**Fig. 6a**). Next,

two strategies were built up via liquid incubation and solid incubation. As for liquid incubation,

to prove the concept, four M9 salt-based liquid mediums were composed of additional ddH_2O,

fructose, glucose, and erythritol. NC and ED were inoculated into these mediums respectively

for days' incubation. As we expected, Both NC and ED could not grow without carbon source

(ddH_2O) but could rapidly grow with fructose and glucose, and only ED could slowly grow

after 36 h with erythritol but NC could not (**Fig. 6b**). After successfully proving the concept,

then we replaced the additional carbon sources by three boughten sodas (10× diluted for use).

Soda 1 was a sugar-free drink that contained two artificial sweeteners: acesulfame potassium

and sucralose. Soda 2 was a sugary drink with fructose and glucose. Soda 3 was a sugar-free

drink that consisted of erythritol and sucralose. After similar incubation, NC could not grow in

soda 3, but ED could utilize erythritol and grow in soda 3 (**Fig. 6c**). These results indicated that

our engineered *E. coli* strains could easily distinguish different sweet soda drinks via liquid

incubation.

In another way, a solid incubation strategy was also built up: prepared M9 salt-agar plates

with additional carbon source liquid could perform as solid medium to verify whether the *E.*

coli could utilize such carbon source as carbon source. As expected, the colony of both NC and

ED could not be observed without carbon source (ddH$_2$O) and could be observed with fructose

and glucose, only ED could form colony after 24 h with erythritol but NC could not (**Fig. 6d**

and Supplementary Fig. 27). Then we used the same three sodas to successfully demonstrated

that solid incubation strategy could also distinguish different sweet soda drinks (**Fig. 6e**). In

summary, we developed simple, visualized, and reliable strategies to verify different soda

drinks by engineered erythritol catabolic *E. coli* strains.

Discussion

In this project, we demonstrated that engineered *E. coli* could transport and utilize the

natural sweetener erythritol as sole carbon source for growth. Firstly, we gained erythritol

catabolism gene cluster from a separated environmental bacterium *Ochrobactrum spp.*.

Secondly, we successfully characterized erythritol catabolism-related genes from three aspects,

including an erythritol-responded transcriptional repressor, an erythritol-catabolic gene cluster,

and an erythritol ABC-transporter system. For in-depth exploration, mRNA transcriptional

analysis was used to compare distinctions between erythritol and glucose as C4 and C6 carbon

sources. Such results inspired us to improve erythritol catabolic efficiency via metabolic

engineering strategies, and we finally remarkably improved the *E. coli* growth in M9-erythritol

medium. Moreover, we applied such engineered strains to expand usages in two fields: On the

one hand, erythritol catabolism enabled engineered probiotic EcN to utilize erythritol in SIF for

increased growth. On the other hand, our engineered *E. coli* performed as "erythritol detector"

to distinguish different sweet soda drinks via two simple strategies.

Overall, our work highlights the potential applications of erythritol catabolism, which will

be a promising addition to the current synthetic biology field. What is more, we developed a

logical research procedure that enables the model chassis to achieve specific goals: Firstly, we

raised the question of how to make full use of erythritol to solve problems and expand

applications rather than just as a natural sweetener in foods and drinks. Secondly, we set a goal

that we want *E. coli* to utilize erythritol as sole carbon source. Thirdly, we acquired resources

of separated *Ochrobactrum spp.* and erythritol catabolism-associated genes. Fourthly, proof of

concept and optimizations were achieved via different strategies. Finally, some expanded

applications might be necessary. In conclusion, such a procedure may inspire synthetic biology

researchers to find out scientific topics and solve practical problems.

Looking forward, we expect our research will provide strategies for expanding synthetic

biology applications, such *E. coli*-based synthetic capability would build up new approaches

for bringing novel abilities to heterologous organisms for useful purposes (**Supplementary Fig.**

28, 29), just as our erythritol catabolism *E. coli* developed connections between lab and daily

life. We can imagine that in the near future, sugar-free drinks will not only provide sweet tastes

but also balance our intestinal flora, and even treat diseases via engineered probiotics.

Furthermore, erythritol will play much more important roles in different fields, including

energy and environment science and engineering, agriculture, food science and technology,

materials science and engineering, and biomedical engineering.

Methods

Strains, vectors, plasmids, primers, and reagents

The details of *E. coli* strains, vectors, plasmids, and primers used in this study are listed in **Supplementary Table 1, 4, 5,** and **6**, respectively. All plasmid sequences are listed in **Supplementary Information**, these plasmids were checked for correctness with Sanger sequencing (GENEWIZ) unless otherwise noted. Oligonucleotides were synthesized by GENEWIZ and listed in Supplementary Table 6. Q5 High-Fidelity DNA Polymerase (New England Biolabs), Phanta Super-Fidelity DNA Polymerase (Vazyme), FastPure Gel DNA Extraction Mini Kit (Vazyme), and ClonExpress Ultra One Step Cloning Kit (Vazyme) were used for molecular cloning. DreamTaq Green PCR Master Mix (Thermo Scientific) was used for colony PCR. Lysogeny Broth (LB) liquid medium contains 10 g tryptone (Oxoid), 5 g yeast extract (Oxoid), and 10 g sodium chloride (Titan/Greagent) in 1 L ddH$_2$O. LB-agar plate was prepared by adding 15 g agar (Sangon Biotech) per liter LB liquid medium. M9 minimal medium (also M9-glucose medium) contains 200 mL 5× M9 salts (15 g/L KH$_2$PO$_4$, 2.5 g/L NaCl, 33.9 g/L Na$_2$HPO$_4$, and 5 g/L NH$_4$Cl), 20 mL 20% w/v glucose, 2 mL 1M MgSO$_4$, and 100 μL 1M CaCl$_2$ in 1 L ddH$_2$O. M9-erythritol medium contains 200 mL 5× M9 salts, 20 mL 20% w/v erythritol, 2 mL 1M MgSO$_4$, and 100 μL 1M CaCl$_2$ in 1 L ddH$_2$O. Antibiotic stocks (1000×) are 100 mg/mL ampicillin, 50 mg/mL kanamycin, and 34 mg/mL chloramphenicol.

Genetic parts and gene clusters

All genetic parts (promoter, RBS, promoter-RBS, terminator, CDS, and protein binding site) and gene clusters used in this study are listed in **Supplementary Table 2,** and **3**. Genetic

parts shorter than 100 bp were synthesized within the oligonucleotides (GENEWIZ) and

inserted into PCR fragments during molecular cloning. Gene clusters were amplified from the

separated *Ochrobactrum spp.* genome, the referenced genome for DNA-DNA alignments was

Brucella anthropi strain T16R-87 chromosome 2 (GenBank: CP044971.1). Gene clusters were

checked for correctness with Sanger sequencing (GENEWIZ), and some of the genetic parts

were amplified from gene clusters.

Plasmids construction

All plasmids were constructed by Gibson Assembly. In brief, all linear polymerase chain

reaction (PCR) products (Q5 High-Fidelity DNA Polymerase, or Phanta Super-Fidelity DNA

Polymerase) were extracted by gel extraction and then assembled by using ClonExpress Ultra

One Step Cloning Kit (Vazyme). After assembly, the reaction mixture was added to 50 μL

competent *E. coli* Mach1-T1 cells, which were incubated overnight on LB-agar plate. Then,

DreamTaq Green PCR Master Mix (Thermo Scientific) was used for colony PCR. The possible

PCR products were sent to GENEWIZ for Sanger Sequencing.

Western-Blot assay

The *E. coli* pellets were collected in 1.5 mL centrifuge tubes at 5000 g and 4°C for 10 min.

The supernatant was discarded and the pellets were resuspended with 1 mL 1× phosphate-

buffered saline (pH 7.4) and lysed by sonication (Q125 sonicator, Qsonica, 10 s on/off, 50% of

amplitude, input energy ~600 Joules). The lysate was then centrifuged at 12000 g and 4°C for

10 min. The total (cell lysate, T), soluble (supernatant, S), and pellet (P) fractions were

separated by SDS-PAGE or Native-PAGE (Omni-Easy One-Step PAGE Gel Fast Preparation

Kit, EpiZyme), followed by wet transferring to PVDF membrane (Bio-Rad) with 1× transfer

buffer (25 mM Tris-HCl, 192 mM glycine and 20% v/v methanol in 1 L ddH$_2$O, pH 8.3). Then,

the PVDF membrane was blocked (Protein Free Rapid Blocking Buffer, Epizyme) for 1 h at

room temperature. After washing thrice with TBST for each 5 min, 1:10000 (TBST buffer-

based) diluted His-Tag Mouse Monoclonal Antibody (Proteintech) solution was added to the

membrane and incubated for 1 h at room temperature. After washing thrice with TBST for each

5 min, 1:10000 (TBST buffer-based) diluted HRP-Goat Anti-Mouse IgG (H+L) Antibody

(Proteintech) solution was added to the membrane and incubated for another 1 h at room

temperature. After the last washing with TBST thrice for each 5 min, the membrane was

visualized using Omni ECL reagent (EpiZyme) under UVP ChemStudio (analytikjena).

Ochrobactrum spp. separation, identification, and characterization

25 mL 1× M9-erythritol medium were added into a 50 mL centrifuge tube, the lid was

replaced by gauze to block aerosol dust and enable microorganisms to drop into the medium.

Such device was stationary-placed outdoors at approximately 25°C for about 7 days. On the

seventh day, the medium became turbid, then the mixture was separated onto an M9-erythritol

agar plate for picking single colonies at 25°C. After three days, some white colonies were

formed and picked up for identification. 16s rRNA was sequenced by primer pair 27F/1492R,

cell morphology was characterized by scanning electron microscope (SEM, JSM-7800F Prime)

and transmission electron microscope (TEM, JEM-2100 Plus). *Ochrobactrum spp.* genome

sequence was referred to *Brucella anthropi* strain T16R-87 chromosome 1 and 2 (GenBank:

CP044970.1 and CP044971.1).

Erythritol catabolic *E. coli* incubation in M9-erythritol liquid medium

All engineered *E. coli* strains that could utilize erythritol as sole carbon source were

cultured and followed the same procedure: Cell pellets from overnight LB culture were

collected in 1.5 mL centrifuge tubes at 5000 g and 4°C for 10 min. The supernatant was

discarded and the pellets were washed thrice by M9-erythritol medium, then the pellets were

resuspended with M9-erythritol and diluted to OD$_{600}$=1.0 for standardization. Then, the

resuspended mixture was inoculated into new M9-erythritol medium (100 mL medium in a 250

mL conical flask, or 1 L medium in a 2 L conical flask) as 1:500 v/v for the following incubation,

in addition, all antibiotic concentration in M9-erythritol medium was diluted for five times for

avoiding growth inhibition in the minimal medium as far as possible.

Erythritol HPLC analysis

1 mL M9-erythritol medium (with *E. coli*) cell cultures were collected in 1.5 mL centrifuge

tubes at 5000 g and 4°C for 10 min, the supernatants were passed through a 0.22 μm filter to

clear cell pellets and other precipitation. Then the samples were loaded for the following HPLC

analysis (**Supplementary Fig. 30**).

1260 Infinity II Prime LC System (Aligent) was used as HPLC instrument, Hi-Plex H

column (Aligent) was used for carbohydrate analysis. The experiment parameters were set as

follows: column temperature: 60°C, mobile phase: 5 mM H_2SO_4 solution, flow rate: 0.6

mL/min, detector: Refractive Index Detector (RID) at 40°C, sample loading volume: 2 μL,

testing time: 20 min for each sample.

eryD binding site characterization *in vivo*

eryD binding site (eryO) was characterized and located in pEryF-732, which was a 732 bp

gap between eryE (5' start) and eryA (3' end). In general, the characterization was performed

as *in vivo* incremental truncation process in two steps:

Firstly, to determine the 5' site, a series of 100 bp-truncated variant plasmids (pFB186 to

pFB191, and pFB210) were constructed and characterized that eryO was located between

pEryF-232 to pEryF-132. Secondly, a series of 10 bp-truncated variant plasmids (pFB192 to

pFB198, pFB208, and pFB209) were constructed and characterized that eryO was located

between pEryF-162 to pEryF-152. Thirdly, a series of 1 bp-truncated variant plasmids (pFB202

to pFB207) were constructed and characterized that eryO 5' site was located in pEryF-159.

Then, to determine the 3' site, the truncation process was the same as mentioned above,

the constructed plasmids (pFB214 to pFB238) were characterized step-by-step. Finally, eryO

3' site was located in pEryF-131, and eryO DNA sequence was characterized as a 29 bp DNA

sequence of 5'-gaaaaaaaatgcgccatctagaaaatttt-3'.

Electrophoretic mobility shift assay (EMSA)

DNA fragment containing the 732 bp eryE-eryA intergenic region (pEryF-732) was amplified with the 5'FAM-labeled primers from plasmid pFB147 and then gel purified. The DNA fragment (200 nM) was incubated with different concentrations of purified eryD in the binding buffer (10 mM Tris-HCl pH 7.5, 1 mM EDTA, 100 mM NaCl, 0.1 mM DTT, 10 μg/mL BSA, 5% v/v glycerol). The 20 μL mixtures were incubated for 30 min at room temperature and then mixed with 5 μL of the same buffer supplemented with 50% v/v glycerol and bromophenol blue. Free DNA and eryD–DNA complexes were separated on 7.5% polyacrylamide gels in 1x TAE buffer (4.844 g/L Tris base, 1.21 mL/L acetate acid, 0.372 g/L EDTA disodium salt dihydrate, pH 8.3). The fluorescence of FAM-labeled bands was visualized under UVP ChemStudio (analytikjena).

eryD orthogonality characterization

E. coli Mach1-T1 with plasmids pFB186 and pFB261 was used for characterization. Inducers (erythritol, glycerol, L-arabinose, D-xylose, D-glucose, D-mannose, D-galactose, D-fructose, L-sorbose, L-rhamnose, D-sucrose, Lactose, Maltodextrin, D-sorbitol, D-mannitol) were dissolved in ddH$_2$O as different concentration (5%, 1%, 0.1%, 0.01%, w/v) solutions, then the solutions were passed through a 0.22 μm filter.

E. coli Mach1-T1 was transformed with the relevant plasmids, starter cultures (LB liquid medium containing 50 μg/mL ampicillin and 17 μg/mL chloramphenicol) were inoculated from a single transformant colony and grew overnight at 37°C for 16 h. The next day, 10 μL starter culture was used to inoculate 5 mL LB medium containing 50 μg/mL ampicillin and 17 μg/mL chloramphenicol in test tubes. The cultures were incubated at 37°C with shaking (250 rpm) until OD$_{600}$ reached 0.6-0.8. The cultures were then added with different inducer solutions and incubated for another 16 h at 37°C and 250 rpm.

The next day, cell pellets were collected in 1.5 mL centrifuge tubes at 5000 g and 4°C for 10 min. The supernatant was discarded and the pellets were resuspended with 1 mL 1× phosphate-buffered saline (pH 7.4). The sfGFP fluorescence of resuspended mixture was

measured by microplate reader (Synergy H1, BioTek) and performed with excitation and

emission wavelengths at 485 and 528 nm, respectively. All measurements were performed at

least in triplicate. Another part of cell pellets was lysed by sonication (Q125 sonicator, Qsonica,

10 s on/off, 50% of amplitude, input energy ~600 Joules). The lysate was then centrifuged at

12000 g and 4°C for 10 min. The total fraction was separated and analyzed by Western-Blot.

eryD structure modeling

Ochrobactrum spp. eryD amino acid sequence was the same as eryD (Uniprot ID: Q2YIQ4)

from *Brucella abortus* (strain 2308). AlphaFold, Robetta, and SWISS-MODEL were used for

structure modeling.

Gel-filtration chromatography

To determine the native molecular weight and assembly model of eryD, gel-filtration

chromatography was performed using a Superdex 200 Increase 10/300 GL column (GE

Healthcare) and standard proteins including Vlm2 (284 kDa) and T7 RNA polymerase (99 kDa).

Purified eryD was concentrated in $1\times$ phosphate-buffered saline (pH 7.4).

erythritol ABC-transporter system transmembrane phenotype characterization

TMHMM-2.0 was used for transmembrane prediction: eryE was predicted as a single

transmembrane protein, and eryG was predicted as a ten-times transmembrane protein. To

determine the transmembrane phenotypes, pFB162 and pFB164 were constructed for eryE-

sfGFP and eryG-sfGFP expression. After transformation into *E. coli* Mach1-T1, starter cultures

(LB liquid medium containing 34 μg/mL chloramphenicol) were inoculated from a single

transformant colony and grew overnight at 37°C for 16 h. The next day, 10 μL starter culture

was used to inoculate 5 mL LB medium containing 34 μg/mL chloramphenicol in test tubes.

The cultures were incubated at 37°C with shaking (250 rpm) until OD_{600} reached 0.6-0.8. The

cultures were then added 1% w/v arabinose and incubated continuously for another 6 h at 30°C

and 250 rpm. Afterward, cell pellets from 1 mL cell culture were collected in 1.5 mL centrifuge

tubes at 5000 g and 4°C for 10 min. The supernatant was discarded and the pellets were

resuspended with 1 mL 1× phosphate-buffered saline (pH 7.4). Then the mixture was imaged

under laser scanning confocal microscope (FV3000, Olympus).

Protein purification

Plasmids pFB161 and pFB167 were used to express and purify eryD and eryR, respectively.

These two proteins were expressed and purified by the same protocol. *E. coli* BL21(DE3) was

transformed with the relevant plasmids, starter cultures (LB liquid medium containing 100

μg/mL ampicillin) were inoculated from a single transformant colony and grew overnight at

37°C for 16 h. 5 mL starter culture was used to inoculate 1 L LB medium containing 100 μg/mL

ampicillin in a 2 L conical flask. The cultures were incubated at 37°C with shaking (250 rpm)

until OD_{600} reached 0.6-0.8. The cultures were then quickly cooled down on ice rapidly to 20°C,

and isopropyl-β-D-thiogalactopyranoside (IPTG) was added for induction at a final

concentration of 0.5 mM. The cultures then grew for another 16 h at 20°C and 220 rpm.

Afterward, cells were harvested by centrifugation (Avanti JXN-26 High-Speed Centrifuge,

Beckman Coulter) at 5000 g and 4°C for 10 min, the pellets were resuspended in Buffer A,

containing 20 mM sodium phosphate (pH 7.4), 1 M sodium chloride, 1 mM dithiothreitol

(DTT), and 50 mM imidazole. The suspension was cooled on ice and then lysed thrice at 1500

bar by ultra-high-pressure homogenization (JNBIO). The lysate was then centrifuged at 4°C

and 20000 g for 30 min. The supernatant was collected and passed through a 0.22 μm filter.

The filtered solution was then purified by Ni^{2+} affinity chromatography using a 1 mL HisTrap

FF column (GE Healthcare). The column was equilibrated with 25 mL Buffer A at a constant

flow rate of 1mL/min. After column equilibration, the filtered protein solution was loaded,

followed by washing with 25 mL Buffer A. Then, bounded proteins were eluted with Buffer B

(Buffer A, but with 500 mM imidazole) and collected into 1.5 mL centrifuge tubes with 0.5 mL

elution. Each 0.5 mL elution was analyzed by SDS-PAGE. Eluted protein samples were

combined in one tube and desalinated using an ultrafiltration tube (Amicon Ultra 3 kDa

molecular weight cut-off, Merck/Millipore) with Buffer C (25 mM Tris-HCl (pH 7.5), 1 mM

DTT, and 1 M sodium chloride). The desalinated and concentrated protein solution was then

mixed with 40% glycerol (v/v in water) with a volume ratio of 1:1. The concentration of the

final protein solution was measured at 280 nm after molar attenuation coefficient correction.

Purified proteins were flash frozen by liquid nitrogen and stored at -80°C until further use.

Protein expression and solubility characterization

Plasmids pFB158 to pFB167 were used to express ten erythritol catabolism-associated

genes. *E. coli* strains were transformed with the relevant plasmids, starter cultures (LB liquid

medium containing 100 μg/mL ampicillin or 34 μg/mL chloramphenicol) were inoculated from

a single transformant colony and grew overnight at 37°C for 16 h. The next day, 10 μL starter

culture was used to inoculate 5 mL LB medium containing 100 μg/mL ampicillin or 34 μg/mL

chloramphenicol in test tubes. The cultures were incubated at 37°C with shaking (250 rpm)

until OD_{600} reached 0.6-0.8. The cultures were then added with inducers (1% w/v arabinose or

0.5 mM IPTG) and incubated continuously. Afterward, cell pellets were collected in 1.5 mL

centrifuge tubes at 5000 g and 4°C for 10 min. The supernatant was discarded and the pellets

were resuspended with 1 mL 1× phosphate-buffered saline (pH 7.4) and lysed by sonication

(Q125 sonicator, Qsonica, 10 s on/off, 50% of amplitude, input energy ~600 Joules). The lysate

was then centrifuged at 12000 g and 4°C for 10 min. The total (cell lysate, T), soluble

(supernatant, S), and pellet (P) fractions were separated and analyzed by SDS-PAGE and

Western-Blot.

Fluorescence measurement standardization

sfGFP (C-terminal 6×His) was used as fluorescence reporter. pFB286 to pFB291 were

performed as standardized sfGFP expression plasmids, these plasmids consisted of four parts

within iGEM standardized vector pSB1C3: *E. coli* constitutive promoter library (J23100,

J23106, J23105, J23114, J23109, J23113, from strong to weak), strong ribosome binding site

B0034, sfGFP, and strong terminator B0015. Other plasmids containing sfGFP were compared

to these six standardized plasmids to determine sfGFP expression level. sfGFP fluorescence

was measured by microplate reader (Synergy H1, BioTek) and performed with excitation and

emission wavelengths at 485 and 528 nm, respectively. All measurements were performed at

least in triplicate.

E. coli genes knock-out

Gene knock-out strategy was derived from Lambda Red recombination. Concretely, plasmids pFB278 to pFB284, and pFB306 to pFB339 were constructed to obtain linear PCR products. The descriptions "homoL" and "homoR" of these plasmids represented "homologous left arm" and "homologous right arm" between the target knock-out gene or gene cluster, these two DNA sequences were designed approximately longer than 300 bp for a higher recombination efficiency. Such linear PCR products (homoL - FRT - KanR - FRT - homoR) were obtained and purified to be used.

Wild type *E. coli* MG1655 was firstly transformed with pKD46, on the next day, a single colony was picked up and inoculated in LB medium (with 1% w/v arabinose) and incubated at 30°C and 250 rpm for approximately 4-6 h, when OD_{600} reached 0.6-0.8, the cell pellets were collected in 1.5 mL centrifuge tubes at 5000 g and 4°C for 10 min. The supernatant was discarded and the pellets were washed thrice with 4°C ddH_2O. Then 1 μg purified linear PCR products were added to the cell mixture and proceeded electro-transformation (Gene Pulser Xcell system (Bio-Rad) and Gene Pulser/Micropulser Electroporation Cuvettes, 0.1 cm gap (Bio-Rad) were used). Subsequently, colony PCR (primer pair: "homoL" forward primer / "homoR" reverse primer) were performed to test whether the gene or gene cluster was deleted. Finally, the expected knock-out *E. coli* MG1655 strains were continuously incubated for days at 37°C for pKD46 (temperature-sensing plasmid) elimination.

Sample preparation and mRNA transcriptome analysis

To process transcriptome analysis, biological triplicate *E. coli* MG1655 (with plasmid pFB147) in both M9-glucose medium (control group) and M9-erythritol medium (experimental group) were performed.

Cell pellets from overnight LB culture (37°C, 250 rpm for 16 h) were collected in 1.5 mL centrifuge tubes at 5000 g and 4°C for 10 min. The supernatant was discarded and the pellets were washed thrice by M9-glucose medium or M9-erythritol medium, then the pellets were resuspended and diluted to $OD_{600}=1.0$ for standardization. Then, such mixtures were inoculated

into new M9-glucose medium or M9-erythritol medium (1 L medium in a 2 L conical flask,

with 7 mg/mL chloramphenicol) as 1:500 v/v for the following incubation.

As for M9-glucose medium, when OD_{600} reached 0.6 (mid-log phase, approximately 6 h

after inoculation), cell pellets were collected. As for M9-erythritol medium, when OD_{600}

reached 0.15 (mid-log phase, approximately 36 h after inoculation), cell pellets were collected.

Then the cell pellets were sent to GENEWIZ for further sample analysis.

Engineered EcN growth in SIF

Probiotic *E. coli* Nissle 1917 (EcN, Mutaflor) were transformed with plasmid pFB147,

such engineered EcN was prepared and diluted in ddH_2O with gradient density (2×10^2, 2×10^4,

2×10^6, 2×10^8) for the following experiment. $2\times$ simulated intestinal fluid (SIF) was prepared

that followed standard protocol with additional erythritol (0, 0.02%, 0.2%, 2%, 8%, 32%,

w/v)[45,46], the solutions were all passed through 0.22 μM filter. 7.5 mL EcN culture and 7.5 mL

$2\times$ SIF were mixed as 1:1 volume to gain a 15 mL culture. such anaerobic cultures were

stationary-placed for incubation for days at 37°C. At each predetermined time point,

approximately 100 μL samples were taken and separated (diluted if necessary) on LB-agar plate

for cell density calculation. All experiments were performed at least in triplicate.

Soda drinks differentiation

E. coli MG1655 with plasmids pFB151 and pFB147 were performed as negative control

(NC) strain and erythritol detector (ED) strain, respectively. Three kinds of sweet soda drinks

were bought from the market. Their ingredients were shown as follows: Soda 1: water, carbon

dioxide, citric acid, potassium citrate, acesulfame potassium, sodium benzoate, sucralose,

sodium gluconate, and essence. Soda 2: water, high-fructose corn syrup (HFCS, including

fructose and glucose), white sugar (sucrose), carbon dioxide, citric acid, potassium citrate,

sodium benzoate, sucralose, acesulfame potassium, essence. Soda 3: water, erythritol, carbon

dioxide, citric acid, potassium citrate, sodium bicarbonate, sucralose, essence. All sodas were

passed through 0.22 μm filter to remove impurities and then shaken to eliminate carbon dioxide.

As for liquid incubation strategy, the sodas were 10 times diluted in ddH_2O to decrease the

ingredient concentration, and the diluted sodas were used as different liquid mediums. NC and ED cell pellets from overnight LB culture (37°C, 250 rpm for 16 h) were collected in 1.5 mL centrifuge tubes at 5000 g and 4°C for 10 min. The supernatant was discarded and the pellets were washed thrice by ddH$_2$O, then the pellets were resuspended and diluted to OD$_{600}$=1.0 for standardization. Then, 10 μL mixtures were inoculated into 5 mL diluted sodas and incubated at 37°C, 250 rpm for 12 h, 24 h, and 36 h. The solutions were imaged at each time point.

As for solid incubation strategy, firstly, M9 salt agar plates were prepared (10 mL mixture in 10 mm plastic plates). Next, 1 mL filtered sodas (without dilution) were added and separated on the solid medium and waited for desiccation. Next, 10 μL NC or ED cell mixtures were inoculated onto the solid medium and incubated at 37°C for 24 h, and 48 h. The bacterial colonies were observed and imaged at each time point.

Acknowledgements

This work was supported by grants from the National Natural Science Foundation of China (Nos. 31971348 and 32171427). J. L. also acknowledges the starting grant of ShanghaiTech University.

Author contributions

J. L. and F. B. designed the experiments. F. B., X. Y. J., S. H. H., J. C. Z., and Y. F. Z. performed experiments. X. Y. J. helped perform cell culture. S. H. H. performed HPLC analysis. J. C. Z. performed electron microscope imaging. Y. F. Z. helped molecular cloning. F. B. analyzed the data and drafted the manuscript. J. L. and W. Q. L. revised and edited the manuscript. J. L. conceived and supervised the study. All authors read and approved the final manuscript.

Competing interests

The authors declare no competing interests.

References

1. Grembecka, M. Sugar alcohols-their role in the modern world of sweeteners: a review. *Eur. Food Res. Technol.* **241**, 15-16 (2015).

2. Shankar, P., Ahuja, S. & Sriram, K. Non-nutritive sweeteners: review and update. *Nutrition* **29**, 1293-1299 (2013).

3. Bernt, W. O., Borzelleca, J. F., Flamm, G., & Munro, I. C. Erythritol: a review of biological and toxicological studies. *Regul. Toxicol. Pharmacol*. **24**, S191–S197 (1996).

4. Munro, I. C., Berndt, W. O., Borzelleca, J. F., Flamm, G., Lynch, B. S., Kennepohl, E., Bär, E. A., & Modderman, J. Erythritol: an interpretive summary of biochemical, metabolic,

toxicological and clinical data. *Food Chem. Toxicol*. **36**, 1139-1174 (1998).

5. Oku, T., & Okazaki M. Laxative threshold of sugar alcohol erythritol in human subjects.
Nutr. Res. **16**, 577-589 (1996).

6. Veiga-da-Cunha, M., Santos, H., & Van Schaftingen, E. Pathway and regulation of
erythritol formation in *Leuconostoc oenos*. *J Bacteriol*. **175**, 3941-3948 (1993).

7. Richter, H., Vlad, D., & Unden, G. Significance of pantothenate for glucose fermentation
by *Oenococcus oeni* and for suppression of the erythritol and acetate production. *Arch.*
Microbiol. **175**, 26-31 (2001).

8. Mirończuk, A., Dobrowolski, A., Rakicka, M., Rywińska, A., & Rymowicz, W. Newly
isolated mutant of *Yarrowia lipolytica* MK1 as a proper host for efficient erythritol
biosynthesis from glycerol. *Process Biochem.* **50**, 61-68 (2015).

9. Rymowicz, W., Rywińska, A., & Marcinkiewicz, M. High-yield production of erythritol
from raw glycerol in fed-batch cultures of *Yarrowia lipolytica*. *Biotechnol. Lett*. **31**, 377-
380 (2009).

10. Qiu, X., Xu, P., Zhao, X., Du, G., Zhang, J., & Li, J. Combining genetically-encoded
biosensors with high throughput strain screening to maximize erythritol production in
Yarrowia lipolytica. *Metab. Eng*. **60**, 66-76 (2020).

11. de Cock, P. Erythritol. in *Sweeteners and Sugar Alternatives in Food Technology* (eds.
O'Donnell, K. & Kearsley, M. W.) 213-241 (Wiley Online Library, 2012).

12. Livesey G. Health potential of polyols as sugar replacers, with emphasis on low glycaemic
properties. *Nutr. Res. Rev*. **16**, 163-191 (2003).

13. Arrigoni, E., Brouns, F., & Amadò, R. Human gut microbiota does not ferment erythritol. *Br.*
J. Nutr. **94**, 643-646 (2005).

14. den Hartog, G. J., Boots, A. W., Adam-Perrot, A., Brouns, F., Verkooijen, I. W., Weseler, A.
R., Haenen, G. R., & Bast, A. Erythritol is a sweet antioxidant. *Nutrition* **26**, 449-458
(2010).

15. Ruiz-Ojeda, F. J., Plaza-Díaz, J., Sáez-Lara, M. J., & Gil, A. Effects of Sweeteners on the
Gut Microbiota: A Review of Experimental Studies and Clinical Trials. *Adv Nutr*. **10**, S31-
S48 (2019).

16. Carocho, M., Morales, P., & Ferreira, I. Sweeteners as food additives in the XXI century:

A review of what is known, and what is to come. *Food Chem. Toxicol*. **107**, 302-317 (2017).

17. Barbier, T., Collard, F., Zúñiga-Ripa, A., Moriyón, I., Godard, T., Becker, J., Wittmann, C.,

Van Schaftingen, E., & Letesson, J. J. Erythritol feeds the pentose phosphate pathway via

three new isomerases leading to D-erythrose-4-phosphate in *Brucella*. *Proc. Natl. Acad.*

Sci. U. S. A. **111**, 17815-17820 (2014).

18. Lillo, A. M., Tetzlaff, C. N., Sangari, F. J., & Cane, D. E. Functional expression and

characterization of EryA, the erythritol kinase of *Brucella abortus*, and enzymatic synthesis

of L-erythritol-4-phosphate. *Bioorg. Med. Chem. Lett*. **13**, 737-739 (2003).

19. Sangari, F. J., Agüero, J., & Garcı A-Lobo, J. M. The genes for erythritol catabolism are

organized as an inducible operon in *Brucella abortus*. *Microbiology (Reading)* **146**, 487-

495 (2000).

20. Sperry, J. F., & Robertson, D. C. Erythritol catabolism by *Brucella abortus*. *J. Bacteriol*.

121, 619-630 (1975).

21. Rodríguez, M. C., Viadas, C., Seoane, A., Sangari, F. J., López-Goñi, I., & García-Lobo, J.

M. Evaluation of the effects of erythritol on gene expression in Brucella abortus. *PLoS One*

7, e50876 (2012).

22. Geddes, B. A., Hausner, G., & Oresnik, I. J. Phylogenetic analysis of erythritol catabolic

loci within the *Rhizobiales* and proteobacteria. *BMC Microbiol*. **13**, 46 (2013).

23. Barbier, T. et al. *Brucella* central carbon metabolism: an update. *Crit. Rev. Microbiol*. **44**,

182-211 (2018).

24. Geddes, B. A., & Oresnik, I. J. Genetic characterization of a complex locus necessary for

the transport and catabolism of erythritol, adonitol and L-arabitol in *Sinorhizobium*

meliloti. *Microbiology (Reading)* **158**, 2180-2191 (2012).

25. Huang, H., Carter, M. S., Vetting, M. W., Al-Obaidi, N., Patskovsky, Y., Almo, S. C., &

Gerlt, J. A. A General Strategy for the Discovery of Metabolic Pathways: d-Threitol, l-

Threitol, and Erythritol Utilization in *Mycobacterium smegmatis*. *J. Am. Chem. Soc*. **137**,

14570-14573 (2015).

26. Adams B. L. The Next Generation of Synthetic Biology Chassis: Moving Synthetic

Biology from the Laboratory to the Field. *ACS Synth. Biol*. **5**, 1328-1330 (2016).

27. Reister, M., Hoffmeier, K., Krezdorn, N., Rotter, B., Liang, C., Rund, S., Dandekar, T.,

Sonnenborn, U., & Oelschlaeger, T. A. Complete genome sequence of the gram-negative probiotic *Escherichia coli* strain Nissle 1917. *J. Biotechnol.* **187**, 106-107 (2014).

28. Bathe, S., Achouak, W., Hartmann, A., Heulin, T., Schloter, M., & Lebuhn, M. Genetic and phenotypic microdiversity of *Ochrobactrum spp. FEMS Microbiol. Ecol.* **56**, 272-280 (2006).

29. Chain, P. S., Comerci, D. J., Tolmasky, M. E., Larimer, F. W., Malfatti, S. A., Vergez, L. M., Aguero, F., Land, M. L., Ugalde, R. A., & Garcia, E. Whole-genome analyses of speciation events in pathogenic Brucellae. *Infect. Immun.* **73**, 8353-8361 (2005).

30. Lamontagne, J., Béland, M., Forest, A., Côté-Martin, A., Nassif, N., Tomaki, F., Moriyón, I., Moreno, E., & Paramithiotis, E. Proteomics-based confirmation of protein expression and correction of annotation errors in the *Brucella abortus* genome. *BMC Genomics* **11**, 300 (2010).

31. Kelly, J. R., Rubin, A. J., Davis, J. H., Ajo-Franklin, C. M., Cumbers, J., Czar, M. J., de Mora, K., Glieberman, A. L., Monie, D. D., & Endy, D. Measuring the activity of BioBrick promoters using an in vivo reference standard. *J. Biol. Eng.* **3**, 4 (2009).

32. Beal, J., Haddock-Angelli, T., Baldwin, G., Gershater, M., Dwijayanti, A., Storch, M., de Mora, K., Lizarazo, M., Rettberg, R., & with the iGEM Interlab Study Contributors. Quantification of bacterial fluorescence using independent calibrants. *PLoS One* **13**, e0199432 (2018).

33. Beal, J. et al. Comparative analysis of three studies measuring fluorescence from engineered bacterial genetic constructs. *PLoS One* **16**, e0252263 (2021).

34. Pellicer, M. T., Fernandez, C., Badía, J., Aguilar, J., Lin, E. C., & Baldom, L. Cross-induction of glc and ace operons of *Escherichia coli* attributable to pathway intersection. Characterization of the glc promoter. *J. Biol. Chem.* **274**, 1745-1752 (1999).

35. Seweryn, P., Van, L. B., Kjeldgaard, M., Russo, C. J., Passmore, L. A., Hove-Jensen, B., Jochimsen, B., & Brodersen, D. E. Structural insights into the bacterial carbon-phosphorus lyase machinery. *Nature* **525**, 68-72 (2015).

36. Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W., & Lipman, D. J. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res*. **25**, 3389–3402 (1997).

37. Johnson, M., Zaretskaya, I., Raytselis, Y., Merezhuk, Y., McGinnis, S., & Madden, T. L. NCBI BLAST: a better web interface. *Nucleic Acids Res*. **36**, W5–W9 (2008).

38. Heller, K. B., Lin, E. C., & Wilson, T. H. Substrate specificity and transport properties of the glycerol facilitator of *Escherichia coli*. *J. Bacteriol*. **144**, 274-278 (1980).

39. Sonnenborn U. *Escherichia coli* strain Nissle 1917-from bench to bedside and back: history of a special *Escherichia coli* strain with probiotic properties. *FEMS Microbiol. Lett*. **363**, fnw212 (2016).

40. Pedrolli, D. B., Ribeiro, N. V., Squizato, P. N., de Jesus, V. N., Cozetto, D. A., & Team AQA Unesp at iGEM 2017. Engineering Microbial Living Therapeutics: The Synthetic Biology Toolbox. *Trends Biotechnol*. **37**, 100-115 (2019).

41. Kelly, V. W., Liang, B. K., & Sirk, S. J. Living Therapeutics: The Next Frontier of Precision Medicine. *ACS Synth. Biol*. **9**, 3184-3201 (2020).

42. Cubillos-Ruiz, A., Guo, T., Sokolovska, A., Miller, P. F., Collins, J. J., Lu, T. K., & Lora, J. M. Engineering living therapeutics with synthetic biology. *Nat. Rev. Drug Discov*. **20**, 941-960 (2021).

43. Forker, E. L. Two sites of bile formation as determined by mannitol and erythritol clearance in the guinea pig. *J. Clin. Invest*. **46**, 1189-1195 (1967).

44. Shindou, T., Sasaki, Y., Eguchi, T., Euguchi, T., Hagiwara, K., & Ichikawa, T. Identification of erythritol by HPLC and GC-MS and quantitative measurement in pulps of various fruits. *J. Agric. Food Chem*. **37**, 1474–1476 (1989).

45. Brodkorb, A. et al. INFOGEST static *in vitro* simulation of gastrointestinal food digestion. *Nat. Protoc*. **14**, 991-1014 (2019).

46. Minekus, M. et al. A standardised static *in vitro* digestion method suitable for food - an international consensus. *Food Funct*. **5**, 1113-1124 (2014).

888
889
Fig. 1 | Engineering *E. coli* to utilize erythritol as sole carbon source.
(**a**) Screened out *Ochrobactrum spp.* provided erythritol catabolism gene cluster and enable *E.*
coli to utilize erythritol as sole carbon source. (**b**) The Erythritol degradation pathway consists
of five genes, which catalyze erythritol to D-erythrose 4-phosphate and then flow into pentose
phosphate pathway. (**c**) *E. coli* MG1655 harboring plasmid pFB147 could grow in liquid M9-
erythritol medium. Each OD_{600} value (average ± standard deviation) was measured with three
biological replicates. (**d**) Double Y-axis graph shows *E. coli* growth in M9-erythritol medium
and erythritol utilization. Erythritol concentration was measured by HPLC analysis with three
biological replicates, the error bar represents the standard deviation (s.d.).

899
900
Fig. 2 | erythritol binding transcriptional repressor eryD characterization.
(**a**) DNA sequence of eryD binding site (erythritol operator, eryO, orange highlighted),
promoter pEryF-732 sequence (-35, -10, and +1 regions were annotated as red, blue, and black
respectively), and their genetic location on erythritol catabolism cluster. (**b**) EMSA analysis of
the interaction between eryD and pEryF-732 DNA fragment (5'FAM-labelled). (**c**) Schematic
shows two compatible plasmids pFB261 and pFB186 co-existed in *E. coli* Mach1-T1, sfGFP
fluorescence indicates the transcription strength of pEryF-732, which may be repressed by eryD.
(**d**)-(**f**) Relative sfGFP expression and fluorescence of *E. coli* Mach1-T1 harboring two
plasmids in (**c**). Pellets were collected from 1.5 mL overnight LB culture (37°C, 250 rpm for
16 h) with added erythritol, then the pellets were measured as normalized fluorescence with
three biological replicates, and Western-Blot analysis (sfGFP with C-terminal 6xHis tag: 27.6
kDa). (**g**) (**h**) Schematic diagram of used strains and plasmids for pT7-eryO operon
characterization. pFB276 and pFB261 co-existed in *E. coli* BL21(DE3), pellets were collected
from 1.5 mL overnight LB culture (37°C, 250 rpm for 16 h) with added erythritol, then the

pellets were measured as normalized fluorescence with three biological replicates. (**i**) (**j**)

Schematic diagram of used strains and plasmids for J231xx-eryO operons characterization.

pFB261 and pFB270-pFB275 co-existed in *E. coli* Mach1-T1, pellets were collected from 1.5

mL overnight LB culture (37°C, 250 rpm for 16 h) with added erythritol, then the pellets were

measured as normalized fluorescence with three biological replicates. (**k**) (**l**) eryD-erythritol

orthogonality test was performed by the same strains and procedure in (**c**). sfGFP expression

was analyzed by normalized fluorescence fold change and Western-Blot. All the error bars

represent the standard deviation (s.d.).

Fig. 3 | Reconstruction of the major carbohydrate metabolic pathways mapped with
mRNA transcriptional analysis.
Reconstructed major carbohydrate metabolic pathways of erythritol catabolic *E. coli* MG1655
(harboring pFB147) were based on the relative differential gene expression profiles between
two cultures (M9-erythritol as the experimental group, and M9-glucose as the control group).
The genes with a significant ($p < 0.05$) differential expression of ≥ 1 log2(fold change) are
indicated with arrow (pathway) in green (up-regulated in M9-erythritol medium) or red (down-
regulated in M9-erythritol medium), the other genes without significant differential expression
are indicated with black arrow (pathway). The details of mRNA transcriptional analysis are
shown in the Supplementary Information.

Fig. 4 | Metabolic engineering for improving *E. coli* erythritol catabolism.
(**a**) Schematic plasmids of pFB148 to pFB152. Strength-gradient constitutive promoters
replaced the native promoter pEryF-732 for erythritol catabolism optimization. (**b**) *E. coli*
MG1655 harboring the above-mentioned plasmids grew on M9-erythritol agar plates after 72
h incubation at 37°C. (**c**) Growth curve of *E. coli* MG1655 harboring the above-mentioned
plasmids in M9-erythritol liquid mediums (5 mL, 37°C, 250 rpm). (**d**) (**f**) (**h**) *E. coli* MG1655
harboring erythritol degradation plasmid pFB147 and other compatible plasmids. (**d**) (**e**)
pFB293-pFB297 co-existed with pFB147 in the strains to constitutively overexpress talA and
tktB to explore whether they could accelerate erythritol catabolism. (**f**) (**g**) pFB298-pFB301
co-existed with pFB147 in the strains to constitutively overexpress talB. (**h**) (**i**) pFB302-
pFB305 co-existed with pFB147 in the strains to constitutively overexpress tktA. All the
measurements were performed with three biological replicates, the error bars represent the
standard deviation (s.d.).

970
971
Fig. 5 | Erythritol can facilitate the growth of engineered probiotic *E. coli* Nissle 1917.
(**a**) Schematic workflow of sample preparation, incubation, and measurement, the mixture was
incubated at 37°C without shaking. (**b**)-(**e**) Growth curves with different inoculated cell
densities of 10^2, 10^4, 10^6, and 10^8, respectively, erythritol was added as 0.4% in SIF. (**f**)-(**k**)
Growth curves with different erythritol concentrations of 0%, 0.01%, 0.1%, 1%, 4%, and 16%
in SIF, respectively, all the initial inoculated cell densities were the same as 10^4. "EcN WT"
means wild type *E. coli* Nissle 1917 without plasmids, "EcN eryABCHI" means wild type *E.*
coli Nissle 1917 harboring pFB147. All the measurements were performed with three biological
replicates, the error bars represent the standard deviation (s.d.).

Fig. 6 | Engineered *E. coli* can distinguish soda drinks.

(**a**) Schematic of two *E. coli* MG1655 strains were used as negative control (NC, with pFB151 that cannot utilize erythritol as sole carbon source), and erythritol detector (ED, with pFB147 that can utilize erythritol as sole carbon source). (**b**) (**c**) The liquid incubation strategy was performed to distinguish synthetic M9-based carbohydrate mediums and soda drinks. (**d**) (**e**) The solid incubation strategy was also performed for distinguishing synthetic M9-based carbohydrate mediums (the same as (**b**)) and soda drinks (the same as (**c**)). All samples are repeated by three biological replicates with similar results.

999
1000
Extended Data Fig. 1 | erythritol catabolism-associated genes *in vivo* expression.
(**a**) SDS-PAGE analysis of gene expression of eryA (with N-terminal 6xHis tag, 56.2 kDa),
eryB (with N-terminal 6xHis tag, 57.2 kDa), and eryC (with N-terminal 6xHis tag, 35.8 kDa).
All these three genes were expressed by plasmids pFB158, pFB159, and pFB160 respectively.
E. coli Mach1-T1 with plasmids were initially inoculated with overnight culture (5 mL LB
medium at 37°C, 250 rpm for 16 h) as a volume ratio of 1:500, and then incubated in 5 mL
new LB medium at 37°C, 250 rpm, when OD_{600} reached 0.6, 1% (w/v) arabinose was added
and incubated for another 16 h at 20°C, 250rpm. Then the cell pellets were resuspended
(equal volume) with 1x phosphate-Buffered Saline (pH 7.4) and lysed by sonication, the total
cell lysate "T", supernatant "S" and pellets "P" were analyzed by SDS-PAGE. The result
shows that eryA and eryB could be well expressed with acceptable percentage of soluble
parts, however, eryC protein bands could not be observed on the gel. It is worth noting that
we initially tried to construct the IPTG-inducted pET22b-eryA/eryB/eryC plasmids, however,

we could not get the correct colonies, the possible reason was that eryA/eryB/eryC leakage
might cause cell toxicity[7], and IPTG-LacI inducible systems always lead to unexpected
protein expression leakage. Thus, we constructed the arabinose-araC inducible systems that
weaken leakage and finally, we gained the correct plasmids (pFB158, pFB159, and pFB160).
(**b**) Western-Blot analysis of the same samples in (**a**), the result indicated that eryC could be
expressed with acceptable percentage of soluble parts, however, the total expression amount
is not enough to be directly observed on SDS-PAGE gel.
(**c**) SDS-PAGE analysis of gene expression of eryD (with N-terminal 6xHis tag, 34.5 kDa),
eryR (with N-terminal 6xHis tag, 24.8 kDa), eryH (with N-terminal 6xHis tag, 28.8 kDa),
eryI (with N-terminal 6xHis tag, 16.6 kDa), eryF (with N-terminal 6xHis tag, 56.2 kDa). All
these five genes were expressed by plasmids pFB161, pFB167, pFB165, pFB166, and
pFB163 respectively. *E. coli* BL21(DE3) with plasmids were initially inoculated with
overnight culture (5 mL LB medium at 37°C, 250 rpm for 16 h) as a volume ratio of 1:500,
and then incubated in 5 mL new LB medium at 37°C, 250 rpm, when OD_{600} reached 0.6, 0.5
mM IPTG was added and incubated for another 16 h at 20°C, 250rpm. Then the cell pellets
were resuspended (equal volume) with 1x phosphate-Buffered Saline (pH 7.4) and lysed by
sonication, the total cell lysate "T", supernatant "S" and pellets "P" were analyzed by SDS-
PAGE. The result shows that all of the five proteins could be well expressed with acceptable
percentage of soluble parts and well observed on the gel.
(**d**) Western-Blot analysis of the same samples in (**c**).
(**e**) SDS-PAGE analysis of gene expression of eryA and eryC. All the procedures were the
same as (**a**) except the arabinose-associated incubation: after 1% (w/v) arabinose addition, the
medium was incubated for another 4 h at 37°C, 250rpm rather than 16 h at 20°C, 250rpm.
(**f**) Western-Blot analysis of the same samples in (**e**). All the results of (**e**) and (**f**) indicated
that in this induction process, both eryA and eryC could be expressed with nearly all insoluble
parts. Moreover, eryC could not be expressed a large amount in either high or low
temperatures.

1057

1058

Extended Data Fig. 2 | Native promoter-RBS (pR-) composite part characterization in
original erythritol cluster.
(**a**) Schematic promoter-RBS (pR-) composite part localization. It is worth mentioning that
we did not successfully characterize the function of eryR, thus here we did not test pR-eryR.
(**b**) Schematic procedure of "pR-ery" reporter plasmids construction (pFB177 to pFB185).
Here we illustrate the construction of pR-eryB-sfGFP reporter plasmid as an example. The
DNA sequence between "eryA CDS except initiation codon ATG" and "the last nucleotide
before eryB CDS" was considered as pR-eryB. This part was then assembled with the
following sfGFP (6xHis) and terminator B0015 in vector pSB1C3 (pFB178).
(**c**) Normalized fluorescence result of all nine "pR-ery" parts. All measurements were
performed in *E. coli* Mach1-T1 with three biological replicates. "NC" represents *E. coli*
Mach1-T1 without plasmids. The other three gray values correspond to pFB291, pFB290, and
pFB289 respectively to act as the positive control (iGEM standard biological parts). The
results indicate that all nine composite parts were not strong enough.

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083

1084

Extended Data Fig. 3 | Native RBS characterization in original erythritol cluster.
(**a**) Schematic RBS localization. It is worth mentioning that we did not successfully
characterize the function of eryR, thus here we did not test eryR-RBS.
(**b**) Schematic procedure of "ery-RBS" reporter plasmids construction (pFB168 to pFB176).
Here we illustrate the construction of J23100-eryA-RBS-sfGFP reporter plasmid as an
example. The proposed "eryA-RBS" sequence (approximately 40 bp) was assembled with
promoter J23100, sfGFP (6xHis), and terminator B0015 in vector pSB1C3 (pFB168).
(**c**) Normalized fluorescence result of all nine RBS parts. All measurements were performed
in *E. coli* Mach1-T1 with three biological replicates. "NC" represents *E. coli* Mach1-T1
without plasmids. The other three gray values correspond to pFB286, pFB288, and pFB289
respectively to act as the positive control (iGEM standard biological parts). The results
indicate that "eryB-RBS" was a lit strong compared with strong ribosome binding site B0034,
but the other eight were not strong enough.

1112

1113

Extended Data Fig. 4 | eryE-sfGFP and eryG-sfGFP characterization.

(**a**) Western-Blot analysis of eryE-sfGFP(6xHis) (50.4 kDa). *E. coli* Mach1-T1 with plasmid
pFB162 were initially inoculated with overnight culture (5 mL LB medium at 37°C, 250 rpm
for 16 h) as a volume ratio of 1:500, and then incubated in 5 mL new LB medium at 37°C, 250
rpm, when OD_{600} reached 0.6, 1% (w/v) arabinose was added and incubated for another 6 h at
30°C, 250rpm. Then the cell pellets were resuspended (equal volume) with 1x phosphate-
Buffered Saline (pH 7.4) and lysed by sonication, the total cell lysate "T", supernatant "S" and
pellets "P" were analyzed by Western-Blot. "NC" represents the total cell lysate of *E. coli*
Mach1-T1 with plasmid pFB162 and without arabinose induction.

(**b**) Confocal laser scanning microscope image of *E. coli* Mach1-T1 that express eryE-sfGFP,
green border indicates the transmembrane localization of eryE-sfGFP.

(**c**) Partially enlarged view of the image in (**b**).

(**d**) Western-Blot analysis of eryG-sfGFP(6xHis) (64.6 kDa). *E. coli* Mach1-T1 with plasmid
pFB164 were initially incubated in 5 mL LB medium at 37°C, 250rpm, when OD_{600} reached
0.6, 1% (w/v) arabinose was added and incubated for another 6 h at 30°C, 250rpm. Then the
cell pellets were resuspended (equal volume) with 1x phosphate-buffered saline (pH 7.4) and
lysed by sonication, the total cell lysate "T", supernatant "S" and pellets "P" were analyzed by
Western-Blot. "NC" represents the total cell lysate of *E. coli* Mach1-T1 with plasmid pFB164
and without arabinose induction.

(**e**) Confocal laser scanning microscope image of *E. coli* Mach1-T1 that express eryG-sfGFP,
green border indicates the transmembrane localization of eryG-sfGFP.

(**f**) Partially enlarged view of the image in (**e**).

1136

1137

1138

Extended Data Fig. 5 | eryD assembled as homotetramer.

(**a**) Gel-filtration chromatography of the purified eryD(6xHis) with Superdex 200 increase

10/300 GL column. eryD homotetramer was eluted as 13.37 mL (red peak), the other two

purified enzymes in our group were used as standard: Vlm2 (284 kDa) was eluted at 10.99

mL, and T7 RNA polymerase (99 kDa) was eluted at 14.22 mL.

(**b**) Native-PAGE gel electrophoresis-based Western-Blot analysis of the purified

eryD(6xHis). the expected homotetramer band (approximately 138 kDa) was shown between

the two markers of 150 kDa and 100 kDa.

1147

a

pEryF-732-eryABCHI (*E. coli* MG1655)

| 36 h | 48 h | 60 h | 72 h |

b

c

1148
1149
Extended Data Fig. 6 | Samples preparation for mRNA transcriptional analysis.
Wild type *E. coli* MG1655 strain with plasmid pFB147 was incubated in 5 mL LB medium
(with 34 μg / mL chloramphenicol) for overnight culture (37°C, 250 rpm for 16 h), on the
second day, 5 mL cell culture was washed with 1x phosphate-Buffered Saline (pH 7.4) by
three times and then resuspended by 5 mL M9-erythritol liquid medium, then the resuspended
cell mixture was inoculated into 1 L M9-erythritol liquid medium (with 6 μg / mL
chloramphenicol) as 1:200 v/v in 2 L shaking flask. After that, the 1 L cell mixture was
incubated for 72 h (37°C, 250 rpm), such process was repeated by three biological replicates.
After these pre-experiments, we found that at 36 h, the cell reached the middle logarithmic
phase (see panel **b**). Afterward, we again processed the same experiments (as three biological
replicates) and collect the cell pellets at 36 h as three parallel RNA-seq samples (experimental
group samples).
(**a**) The cell cultures in pre-experiments reached stationary phase at 72 h.
(**b**) Cell growth curve (OD_{600}-time) of pre-experiment samples. In such a situation, the middle
logarithmic phase was reached at 36 h and the stationary phase was reached at 72 h.

(c) The cell cultures (samples in pre-experiment, at 72 h) were picked and separated on LB-

agar plates (with 34 μg / mL chloramphenicol) to test the cell activity (overnight culture at

37°C for 16 h). The colonies were considered normal and did not lose plasmid pFB147.

1168

1169

1170

1171

1172

1173

1174

1175

1176

1177

1178

1179

1180

1181

1182

1183

1184

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204

1205

1206

1207

1208

a

1209

b

1210

c

Control-VS-Experimental

Control-VS-Experimental

1211
1212
Extended Data Fig. 7 | mRNA transcriptional analysis
Transcriptome results were analyzed via (**a**) COG, (**b**) KEGG, and (**c**) GO databases. The
common different genes were associated with carbohydrate metabolism, amino acid
metabolism, and ribosome. In detail, carbohydrate metabolism was commonly up-regulated,
however, ribosome and translation genes were down-regulated, which indicated that
erythritol-utilized *E. coli* tended to uptake more easily utilized carbohydrates, also suppressed
ribosome assembly and protein translation process to resist such a poor carbohydrate
environment.
1221
1222

1223
1224
Extended Data Fig. 8 | Potential endogenous erythritol transporter homologous
screening.
(**a**) Schematic diagram of screening *E. coli* MG1655 native transporter which could facilitate
erythritol transportation.
(**b**) List of all carbohydrate transportation genes and proteins in *E. coli* MG1655.
(**c**) Screened seven carbohydrate transporter-associated genes. glpF has been reported that is
able to transport erythritol. The other six genes are somewhat homologous to erythritol ABC
transporter gene eryG by BLAST protein-protein alignment analysis.
(**d**) Seven genes' mutated *E. coli* MG1655 strains construction by Lambda-Red
recombination. For each of the seven mutation strains, "NC" means the wild type *E. coli*
MG1655 genome PCR product by primer pair "forward left-homologous-arm" and "reverse
right-homologous-arm", and the other five PCR products were the desired mutation strains by
the same primer pair. The KanR genes were not deleted by pCP20.

1248

Extended Data Fig. 9 | Several eryG homologous permeases in *E. coli* may facilitate
erythritol catabolism.

BLAST protein-protein alignment results showed that there were seven *E. coli* native
carbohydrate ABC-transporter permeases that were homologous to erythritol permease eryG,
we constructed 7 different knock-out *E. coli* MG1655 strains by Lambda-Red recombination.
All these 7 strains and WT (wild type *E. coli* MG1655) strain were then transformed with
pFB147 to catalyze erythritol. After that, all strains were incubated (M9-erythritol medium,
37°C, no shaking) in 96-well plates for 72 h (stationary phase). OD_{600} values were measured
each 12 h. This diagram showed the OD_{600} values at 72 h. All values were measured as three
biological replicates, the error bar represents the standard deviation (s.d.). Student's t-tests are
used for statistical analysis, and $p < 0.05$ indicates statistical significance (*$p < 0.05$, **$p <$
0.01, and ***$p < 0.001$). In conclusion, glpF, rbsC, araH, xylH, mglC, and malG knock-out
strains suppressed their growth rate, in another word, these transport systems could facilitate
erythritol transport.

1263