Welcome to the big leaves: best practices for improving genome annotation in non-model plant

genomes

Vidya S Vuruputoor[1], Daniel Monyak[1], Karl C. Fetter[1], Cynthia Webster[1], Akriti Bhattarai[1],

Bikash Shrestha[1], Sumaira Zaman[1], Jeremy Bennett[1], Susan L. McEvoy[1], Madison Caballero[1]

Jill L. Wegrzyn[1]

[1] Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT 06269

16   Author for correspondence: jill.wegrzyn@uconn.edu

21   ABSTRACT

● **Premise of the study**: Robust standards to evaluate quality and completeness are

lacking for eukaryotic structural genome annotation. Genome annotation software is

developed with model organisms and does not typically include benchmarking to

comprehensively evaluate the quality and accuracy of the final predictions. Plant

genomes are particularly challenging with their large genome sizes, abundant

transposable elements (TEs), and variable ploidies. This study investigates the impact

of genome quality, complexity, sequence read input, and approach on protein-coding

gene prediction.

● **Methods:** The impact of repeat masking, long-read, and short-read inputs, *de novo*, and

genome-guided protein evidence was examined in the context of the popular BRAKER

and MAKER workflows for five plant genomes. Annotations were benchmarked for

structural traits and sequence similarity.

● **Results:** Benchmarks that reflect gene structures, reciprocal similarity search

alignments, and mono-exonic/multi-exonic gene counts provide a more complete view of

annotation accuracy. Transcripts derived from RNA-read alignments alone are not

sufficient for genome annotation. Gene prediction workflows that combine evidence-

based and *ab initio* approaches are recommended, and a combination of short and long-

reads can improve genome annotation. Adding protein evidence from *de novo*

assemblies, genome-guided transcriptome assemblies, or full-length proteins from

OrthoDB generates more putative false positives as implemented in the current

workflows. Post-processing with functional and structural filters is highly recommended.

● **Discussion:** While annotation of non-model plant genomes remains complex, this study

provides recommendations for inputs and methodological approaches. We discuss a set

of best practices to generate an optimal plant genome annotation, and present a more

robust set of metrics to evaluate the resulting predictions.

Keywords: genome annotation, plant genomes, gene identification, BRAKER, TSEBRA,

MAKER, StringTie2

1   INTRODUCTION

The first published plant genome, *Arabidopsis thaliana*, was released in 2000 (Arabidopsis

Genome Initiative, 2000). Its small genome size (135Mb) and minimal repeat content stand in

stark contrast to the plant species sequenced and assembled today (Kress et al., 2022). NCBI's

(https://www.ncbi.nlm.nih.gov/) genome repository contains genomes of over 900 land plant

species, and roughly half of these are assembled to chromosome scale. The total number of

complete reference plant genomes has more than doubled in the last five years (Marks et al.,

2021). Initiatives like the Open Green Genomes (OGG) (https://phytozome-

next.jgi.doe.gov/ogg/), 10KP (Cheng et al., 2018), and the Earth BioGenome Project (Lewin et

al., 2022) are improving the phylogenetic representation of plant genomes by sampling

underrepresented clades. The plant genomes published today are more likely to be polyploids

and/or larger genomes with substantial transposable element content (Sun et al., 2022). The

combination of high throughput sequencing advancements, particularly long reads and

chromosome conformation capture approaches have enabled the completion of these more

challenging assemblies (Pucker et al., 2022).

While genome assembly has seen substantial improvements in accuracy and contiguity,

structural annotation remains challenging. This process delineates the physical positions of

genomic features, including protein-coding genes, promoters, and regulatory elements. It can be

followed by functional annotation, which assigns biological descriptors to the identified features.

The accurate classification of these features provides the basis for questions focused on

species evolution, population dynamics, and functional genomics. Errors in genome annotation

are frequent, even among well-studied models, and are propagated through downstream

analyses (Deutekom et al., 2019; Meyer et al., 2020; Salzberg, 2019). In most eukaryotes,

genome annotation is challenged by partial conservation of sequence patterns, variable lengths

of introns, variable distances between genes, alternative splicing, and higher densities of TEs

and pseudogenes (Kersey, 2019; Salzberg, 2019). As a result of these complexities, the

structural annotation process requires more advanced informatic tools and skills that support the

integration and manipulation of large datasets (Mudge & Harrow, 2016).

Structural and functional genome annotation proceeds in three stages: identifying and masking

noncoding regions (repeats); predicting physical positions of gene structures; and assigning

biological information to the predictions (Jung et al., 2020). Repeat regions are soft-masked

(eg., RepeatMasker (Smit, AFA, Hubley, R & Green, P., 2013-2015) and RepeatModeler2

(Flynn et al., 2020)), which means these regions are indicated but not obscured to annotation

software. This is followed by gene prediction, which may be *ab initio* (evidence-free) or

evidenced-based. Evidence-based approaches use RNA-Seq and protein sequence similarity

search alignments. Evidence-based approaches are often used in combination with *ab initio*

(e.g. AUGUSTUS; (Stanke & Waack, 2003)) to generate models that are trained on patterns

associated with true genes. Given the advanced state of high throughput transcriptome

sequencing, it is common to resolve transcripts from RNA reads through genome-guided

approaches, such as StringTie2 (Kovaka et al., 2019). Long-read cDNA sequencing through

PacBio and Oxford Nanopore can provide additional resolution and improve the identification of

splice variants. When extrinsic evidence from RNA-seq and protein alignments are available,

workflow packages like MAKER (Campbell, Holt, et al., 2014; Cantarel et al., 2008; Holt &

Yandell, 2011) and BRAKER (Brůna et al., 2021; Hoff et al., 2016, 2019) can assist in training

ab initio prediction tools. These packages can leverage sequence data from the target species

as well as external evidence from closely related species. While these workflows can simplify

the integration across external evidence, downstream packages are still required to select or

modify the resulting predictions (Banerjee et al., 2021; Gabriel et al., 2021; Haas et al., 2008).

Here, we provide a comprehensive evaluation of plant genome annotation workflows,

intentionally selecting beyond the typical model species to represent some of the more complex

genomes under investigation today. In doing so, we evaluate the impact of repeat-masking

using two different implementations of the RepeatModeler2 framework (Flynn et al., 2020). This

is followed by exploring the role of read length and accuracy, and the impact of short-read and

long-read data. Finally, we examine the contribution of protein evidence, generated from d*e*

novo assembly of the RNA inputs and a genome-guided assembly. These variations are

examined in the MAKER and BRAKER frameworks to emphasize the importance of defining

benchmarks to guide downstream filtering approaches. Finally, the largest and most repetitive

genome assessed in this study, *Liriodendron chinense*, was used to demonstrate best practices

to refine the predictions.

63   METHODS

Gathering plant genome datasets-

Five plant genomes were chosen for this study, including Chinese tuliptree (*Liriodendron*

chinense) (Chen et al. 2019)*,* black cottonwood (*Populus trichocarpa* v3) (Tuskan et al. 2006)*,*

Chinese rose (*Rosa chinensis*) (Raymond et al. 2018), thale cress (*Arabidopsis thaliana* TAIR

10) (Cheng et al. 2017), and a bryophyte, the common cord-moss (*Funaria hygrometrica*) (Kirbis

et al., 2022) (Table S1). The genomes were selected to represent two model systems (*Populus*

and *Arabidopsis*) with well curated structural annotations and three non-model systems that

exclusively used computational techniques to produce the annotations. Two of these non-

models were also more divergent examples, representing the only sequenced member of their

genus (*Funaria* and *Liriodendron*). The public assembly and annotation for each species were

accessed from NCBI and genome completeness was estimated by searching the genome and

annotation for the conserved single-copy orthologs in the Embryophyta odb10 BUSCO v.5.0.0

(Simão et al., 2015). The contiguity of the reference genomes was assessed with Quast v5.0.2

(Gurevich et al., 2013). Published annotation files were summarized with gFACs (Caballero and

Wegrzyn, 2019).

80
Read sets available through NCBI's Sequence Read Archive (SRA) were accessed to provide

transcriptomic evidence for each species and included a variety of tissue types. The Illumina

short-read libraries were sequenced with Illumina HiSeq 2500 (100bp paired-end). The read

sets included at least four libraries, between 20-82M reads before quality control (QC), and a

minimum of 16M reads after QC. Pacific Biosciences Iso-Seq long-reads were accessed for

Populus and *Liriodendron*, and Oxford PromethION reads were available for *Rosa* and

Arabidopsis. The read sets for long-read data ranged between 161K-41M total reads per

species (Table S2).

Repeat masking and read alignment-

RepeatModeler2 (Flynn et al., 2020) was used to construct repeat libraries with default settings,

and repeats were soft masked with the libraries constructed via RepeatMasker v.4.0.6 (Smit et

al. 2013-2015). The genomes of *Arabidopsis, Funaria, Populus, and Liriodendron* were

additionally masked using RepeatModeler2 with additional LTR identification (-LTRStruct flag).

Quality assessment of the Illumina short reads was performed using FastQC v.0.11.7 (Andrews,

2010) before and after trimming low-quality bases. Sickle v.1.33 (Joshi NA, 2011) was used to

trim low-quality bases with 50bp as the minimum read length threshold. Single end reads

generated post trimming were excluded from RNA alignments and assembly. The trimmed short

reads were aligned against their reference genomes using HISAT2 v2.2.0 (Kim et al., 2019).

HISAT2 was selected for its performance in recent benchmarking studies and as the aligner of

choice for input to Stringtie2 (Corchete et al., 2020; Musich et al., 2021). Long-read RNA data

were obtained for four species: *Arabidopsis* and *Rosa* were sequenced with Oxford Nanopore,

and *Populus* and *Liriodendron* were sequenced with PacBio Sequel. The long-read data sets

were aligned against their respective genomes using Minimap2 v2.1.7 (Li, 2018, 2021).

Generation of protein evidence-

To generate protein evidence, Illumina short reads were assembled *de novo* using Trinity

v.2.8.5 with a minimum contig length of 300 bp (Grabherr et al., 2011). The assembled

transcriptomes for the multiple libraries were combined, and putative coding regions were

predicted using TransDecoder v.5.3.0 (http://transdecoder.github.io). TransDecoder is one of

several frame-selection methods available and performs in a comparable manner but not

always superior in all metrics (Bolger et al., 2018). For this study, it was selected as the most

widely used package for this purpose. Redundancy in the predicted coding regions was reduced

after clustering at 98% identity using UCLUST, a clustering algorithm of USEARCH v.9.0.2132

(Edgar, 2010). Frame-selected transcripts shorter than 300 bp were removed. The remaining

transcripts were aligned to the genome using GMAP v.2019-06-10 (Wu & Watanabe, 2005).

The predicted proteins (from the same Transdecoder run) were aligned to the reference

genome using GenomeThreader v 1.7.1 (Gremme, 2014).

To provide protein evidence from genome-guided sources, the previously aligned Illumina short-

reads (via HISAT2) were constructed into transcripts with StringTie2 v2.2.0 (Kovaka et al., 2019;

M. Pertea et al., 2015). Long-reads were treated similarly, along with a combination of short and

long-reads. The predicted transcripts were extracted using gffRead (G. Pertea & Pertea, 2020)

and frame selected with TransDecoder. The transcriptome alignment annotation file (gff3) was

passed to gFACs for evaluation of gene model statistics. Completeness of the aligned

transcripts and protein sequences were estimated using BUSCO. Finally, to provide evidence

from external sources (not derived from any transcriptomic inputs), full-length protein evidence

from OrthoDB (odb10_plants) was provided for BRAKER/TSEBRA runs.

Genome annotations-

Each genome was tested in four primary open-source annotation softwares to predict gene

models (Table 1). Several different runs of BRAKER v.2.1.5 (Hoff et al., 2019) and

BRAKER/TSEBRA (Gabriel et al., 2021) were used with various combinations of RNA-Seq (long

and short-read inputs) and protein evidence. MAKER v.3.1.3 (Cantarel et al., 2008) was run

once with transcript and protein evidence. Finally, StringTie2 (Kovaka et al., 2019), with

TransDecoder, was used to generate genome-guided predictions from RNA evidence alone

(Table S12).

MAKER annotation-

MAKER (MK) was run on the soft-masked reference genomes of *Arabidopsis, Populus,* and

Funaria with repeats estimated using the additional LTR detection method in RepeatModeler2

(LTRStruct flag; RM2+). This was intended to emulate the MAKER-P (Campbell, Law, et al.,

2014) method since the original repeat and pseudogene identification protocols are deprecated.

MK (RM2+) was executed (i.e., trained) twice. The annotations derived from MK (RM2+) used

protein evidence generated from *de novo* assembled RNA-reads from Trinity. These models

were used to train *ab initio* gene prediction software AUGUSTUS v.3.3.3 (Stanke & Waack,

2003) and SNAP v. 2006-07-28 ((Korf, 2004). The Hidden Markov Models (HMMs) trained using

AUGUSTUS and SNAP were used along with initial aligned evidence (est2genome and

protein2genome parameters) for the second MK (RM2+) run to generate the final gene models.

Assessment of gene predictions-

The quality of genome annotation among different gene prediction methods was evaluated with

three primary metrics: (1) the mono-exonic (single-exon) and multi-exonic (multiple exon) ratio;

(2) conserved single-copy orthologs queried from the predicted gene models using BUSCO

(embryophyta database v10), and (3) gene prediction assessment with EnTAP v0.10.8 (Hart et

al., 2020) using a 70% reciprocal functional annotation approach with NCBI's Refseq Plant and

Uniprot databases. The mono:multi ratio was calculated from the gFACs summary report run

with default parameters (Caballero & Wegrzyn, 2019). We regard a mono: multi ratio near 0.2 to

be ideal and have further validated this with a larger set of model plant genomes (Table S3)

(Jain et al. 2008). The gene prediction assessment was recorded as a percentage of sequence

similarity hits to the total number of genes. The value of the reciprocal BLAST search will be

dependent on the phylogenetic relationships of the target species. While higher annotation rates

indicate better gene prediction assessment, species-specific genes will always be missed

(Armisén et al., 2008), therefore a minimum annotation rate of 80% is a reasonable threshold to

expect when dealing with most non-models. Similarly, a higher BUSCO score indicates a better

annotation since BUSCO utilizes OrthoDB to form its conserved sets and the recommended

target score is >95% for land plants (Manni et al., 2021; Simão et al., 2015).

The sensitivity and precision of the runs for *Arabidopsis* and *Populus* were assessed using

Mikado v2.3.2 (Venturini et al., 2018), by comparing the predicted gene models to the current

reference annotations.

Post-processing filtering-

The predicted gene models for *Liriodendron* were taken a step further to refine the genome

annotation. Post-process filtering was performed using gFACs and assessed for improvement

using BUSCO completeness scores and annotation rates. The mono-exonic and multi-exonic

genes predicted for *Liriodendron* were filtered for unique genes. The mono-exonic genes were

further filtered for the presence of protein domains using InterProScanv.5.35-74.0 and Pfam

(Jones et al., 2014; Quevillon et al., 2005). Multi-exonic genes that did not have an EggNOG or

a sequence similarity hit were removed, and the final annotation was assessed using gFACs

and EnTAP.

45   RESULTS

Genome sizes, repeats, and published annotations-

The genome sizes of the five species assessed represented a 10-fold difference between the

smallest genome of *Arabidopsis* (~119 Mb) and the largest of *Liriodendron* (~1.7 Gb) (Fig 1A;

Table 2). *Liriodendron* (73.18%) and *Rosa* (60.58%) have higher levels of repeat content, and

Arabidopsis has the lowest (23.9%). *Arabidopsis* is the most complete chromosome-scale

genome, with seven contigs reflecting its five chromosomes and two organellar chromosomes.

The other genomes are assembled into pseudochromosomes (except for *Liriodendron*). Once

the genomes were downloaded, contigs < 500 bp were removed. The published genome

assemblies and annotations were compared in terms of completeness via BUSCO (Fig 1, Table

2). When BUSCO is run in genome mode, it searches the genome for the set of 1614 single-

copy orthologs in the embryophyte database. Aside from *Funaria*, which had the lowest

completeness score of 82.4%, the remaining plant genomes ranged from 94% to 99%. When

we evaluated the published annotations for the same species, and ran BUSCO in protein mode,

a slight decrease in completeness was observed in every species except *Funaria* and

Arabidopsis (Fig 1B). The largest reduction in BUSCO score was observed in *Liriodendron*

(98.6% to 75.1%). The discrepancy between the estimated completeness at the genome-level

and most of the published annotations speaks to the challenges of achieving an accurate

structural annotation.

RepeatModeler2 (RM2) with and without the LTRStruct package (the additional LTR masking

module) (Flynn et al. 2020) was used to soft mask repeats in four of the genomes. The increase

in repeat content was marginal in all species, ranging from 1% in *Funaria* to 5% in *Populus*.

Comparisons using the LTRStruct flag were denoted as RM2+ (Table S4).

Transcriptome evidence-

For the subsequent genome annotation analysis, the Illumina RNA short reads were first aligned

to the genome. All libraries, ranging from four to 20 per species, aligned at over 97%, apart from

Rosa (92%) (Table 3; Table S5). Long-read RNA libraries were aligned with Minimap2 for four

species: *Arabidopsis* (Nanopore reads at 97.1%), *Populus* (Iso-Seq reads at 92.01%)*,*

Liriodendron (Iso-Seq reads at 95.5%)*,* and *Rosa* (Nanopore reads at 99%). The N50s for the

long reads range from 976 bp in *Rosa* to 4.6 Kb in *Liriodendron* (Table S6)*.*

Transcript-derived annotations-

The reads were assembled using StringTie2 (ST2) and Trinity. Trinity *de novo* assemblies of the

Illumina short reads generated longer transcripts, with N50s ranging from 1.2 Kb (151,265

transcripts in total) in *Funaria*, to 3.06 Kb (2,839,867 transcripts) in *Liriodendron*. Among

genome-guided assemblies with StringTie2 (ST2(SR)), the range was much smaller, with N50s

ranging from 369 bp (59,741 transcripts in total) in *Funaria* to 2.54 Kb (37,747 transcripts) in

Arabidopsis (Table S6). The StringTie2 (ST2(LR) and ST2(SR/LR)) range was longer, with

N50s ranging from 1.07 Kb (20,633 transcripts in total) in *Rosa* ST2(LR) to 2.36 Kb (45,785

transcripts) in *Liriodendron* ST2 (SR/LR) (Table 3; Table S6). The StringTie2 and Trinity

transcripts were aligned back to the genome using GMAP after frame-selection. BUSCO scores

for the aligned transcriptomes derived from short read data, run in transcriptome mode, ranged

from 73% in *Funaria* to 83% in *Rosa* for Trinity, and 73% in *Liriodendron* to 97% in *Rosa* using

StringTie2 (Table 3). The BUSCO scores were the lowest for the ST2 (LR) runs across all

species as compared to the other StringTie2 only runs. For the ST2 (SR/LR), the BUSCO

scores were lower than ST2 (SR), except for *Rosa*, where the ST2 (SR/LR) was 97.2% as

opposed to 97% in ST2 (SR). In all species, ST2 (SR/LR) had higher BUSCO scores than ST2

(LR). Despite Trinity producing more than double the total transcripts than StringTie2, the

BUSCO completeness score of most StringTie2 runs were much higher than that of Trinity.

Liriodendron remained the only exception with a slightly higher BUSCO score from Trinity.

Arabidopsis and *Populus* were further evaluated with Mikado to compare the sensitivity and

specificity of published annotations (Fig 4B, Table S7). Overall, StringTie2 predictions had

higher sensitivity and precision rates compared to the Trinity runs. From this point, Trinity was

excluded, and StringTie2 runs were compared against BRAKER and TSEBRA predictions (Fig

4B).

The mono: multi ratios produced by StringTie2 ranged from 0.15 in *Populus* (ST2 (LR)) to 0.53

in *Liriodendron* (ST2 (LR)), which were an improvement over the mono: multi ratios produced

from the BRAKER annotations that ranged from 0.37 in *Arabidopsis* (BR (LR)) to 1.27 in *Funaria*

(BR (SR/RM2+)). The BUSCO scores of the proteins predicted from BRAKER were generally

higher than the BUSCO scores from StringTie2. For example, *Arabidopsis* StringTie2 runs

range from 85% (ST2 (LR)) to 95.5% in ST2 (SR), and BRAKER runs ranged from 94% (BR

(LR)) to 95.9% (BR (SR)). However, some runs are comparable, the ST2 (SR) run with a

BUSCO score of 95% was like the BR (SR) run at 95% and the BR (SR/RM2+) run at 95% in

Arabidopsis. StringTie2 predicted models had a higher annotation rate, in general, compared to

BRAKER. For example, the EnTAP annotation rate in *Funaria* was just over 40% post BRAKER

but was near 60% from the StringTie2 runs (Fig 2).

Genome annotation with MAKER -

To replicate MAKER-P's repeat pipeline, the RM2+ genome was used for *Arabidopsis, Populus*,

and *Funaria* for the MAKER runs. BUSCO completeness was low, compared with BRAKER

runs, and ranged from 19.6% in *Populus* to 90.4% in *Arabidopsis* (Fig 3A). The mono: multi ratio

of MAKER(RM2+) for *Arabidopsis* was comparable to the BRAKER runs for the same species

(0.22 for BR (SR) and BR (SR/RM2+), 0.24 for BR (LR), and 0.23 for BR (SR/LR)). The MK

(RM2+) predictions for the total number of genes in *Arabidopsis* and *Funaria* were in the

expected range for these species, 22K and 44K genes, respectively, whereas only 7K genes

were predicted for *Populus*. The gene lengths ranged from 1.8 Kb in *Funaria* to 2.3 Kb in

Arabidopsis (Table S8). The best run for MK (RM2+) was for *Arabidopsis*, with a mono: multi

ratio of 0.22 and a BUSCO score of 90.4%. On the other hand, the mono: multi ratio for *Populus*

was 0.07, and the BUSCO score was 19.6%.

The model systems, *Arabidopsis* and *Populus*, further were evaluated with Mikado to compare

the sensitivity and specificity of the published annotations (Fig 3B; Table S7). The sensitivity

and precision scores for gene predictions were the lowest from MAKER, followed by Trinity, and

were the highest from TSEBRA. StringTie2 and BRAKER yielded similar sensitivity and

specificity scores for *Arabidopsis*, whereas for *Populus* the sensitivity score was lower than

those from BRAKER runs. Given its overall low scores, MAKER was excluded from the

subsequent comparisons. It should be noted, however, that protocols are available that

recommend using GeneMark in MAKER. With this addition to the training protocol, MAKER was

shown to have a higher accuracy in prediction (Brůna et al., 2021).

Genome annotation with BRAKER: -

In general, BUSCO scores were improved in the BRAKER and TSEBRA runs, with the

TSB(SR/OrthoDB) predictions scoring the highest. The best BUSCO scores were from

TSB(SR/OrthoDB), ranging from 98.9% (TSB; SR/OrthoDB) in *Arabidopsis* to 82.2% in *Funaria*.

Overall, the TSEBRA runs fared better with the annotation rates ranging from 99% in

Arabidopsis (TSB; SR/OrthoDB) to 53% in *Funaria*. On the other hand, the TSEBRA runs had

the worst mono:multi ratios. For example, TSB (SR/ST2/RM2+) has the highest ratio at 1.27 for

Funaria (Fig 4).

Overall, the gene models generated by BRAKER for *Arabidopsis* performed similarly across

runs when evaluated by BUSCO scores. The mono: multi ratios across BRAKER runs ranged

between 0.23 to 0.39, and the annotation rates were consistently above 95%.

The annotation rates for *Funaria* were lower than expected, 43% for BR (SR) and 53% for

nearly all methods that included protein-evidence (TSB (SR/TRINITY), TSB (SR/OrthoDB)). The

BUSCO completeness scores of about 85% post BRAKER are comparable to those from

StringTie2.

In the case of *Liriodendron*'s BRAKER and TSEBRA runs, the mono:multi ratios were more

variable when compared to the StringTie2 runs, which ranged from 0.34 BR(SR) to 1.04 with BR

(SR/RM2+). The annotation rates for each run were around 75%, with BUSCO scores from 83%

with TSB (SR/LR/ST2) to 90.8% for BR (SR). *Populus* gene models post-BRAKER without

protein had mono:multi scores around 0.24, and with TSEBRA, the ratio ranged from 0.4 to 0.5.

Annotation rates also differed between BR(SR) (75%) and TSB (SR/ST2) (88%). *Rosa* had

overall consistent scores for BUSCO post BRAKER, ranging around 96%. TSEBRA runs had

higher mono:multi ratios (0.75) compared to the BRAKER runs (0.37) (Table S8).

Annotation with long reads-

For BRAKER runs, the predicted gene lengths from the long-reads were comparable to those

based on short-reads, except for some runs with long-reads within *Populus*. The average gene

length for BRAKER with long-reads for *Populus* ranges from 2.7 Kb to 3.4 Kb, although some

transcripts exceed 6 Kb. The longest average gene lengths were observed in *Liriodendron* (9.3

Kb- BR; SR/LR). The inclusion of long-reads (exclusively) did not improve BUSCO

completeness for any species, apart from *Arabidopsis*, where the BR (LR) BUSCO

completeness was 1% higher than the BR (SR) run. The increase in BUSCO completeness in

Arabidopsis could be due to the large number of long-reads included (23M across four libraries).

However, the quality of genome annotation does not seem correlated with depth of long-read

sequencing; for example, *Rosa* had more reads (41M across 6 libraries), and the BR (LR) run

had a similar BUSCO score to BR (SR) (96%). It should be noted that the long reads for

Arabidopsis and *Rosa* were sequenced with ONT. The ONT reads had higher mapping rates,

compared to Iso-Seq, to their respective genomes, 97.1% in *Arabidopsis* and 99% in *Rosa*

(Table S5). The long-read inputs, regardless of depth or type, lowered the BUSCO

completeness (up to 10%) across all ST2 (LR) runs (Table S9). Finally, we note that the

combination of short-reads and long-reads (BR; SR/LR and TSB; SR/LR/ST2) is comparable to

the BR (SR) reads in terms of BUSCO completeness. Exceptions in this case are *Arabidopsis*

and *Rosa*, with marginally higher BUSCO completeness scores. BR (SR/LR) runs produced

more genes (in total) for *Arabidopsis*, *Populus*, and *Rosa* in comparison to BR (SR); however, in

the TSB (SR/LR/ST2) runs, the total number of genes were lower compared to BR (SR), apart

from *Rosa*. The annotation rate in BR (SR/LR) of *Liriodendron* was higher than BR (SR). The

combination of proteins with SR and LR (TSB(SR/LR/ST2)) resulted in higher annotation rates

across all species. The combination of SR and LR increased mono: multi ratios, therefore,

resulted in worse mono: multi ratios across all species.

Refining the genome annotation for Liriodendron-

The BRAKER runs for *Liriodendron* were filtered with gFACs and InterProScan to remove

unlikely gene models (Table 4). The number of mono-exonic genes was drastically reduced

post-filter with InterProScan. Across all runs, the mono-exonic genes numbered 11K to 25K.

After removing mono-exonics without a protein domain annotation from the Pfam database, they

decreased from 13K to 5K. The decrease in false positive mono-exonics resulted in an

improved mono:multi ratio that range from 0.16 for BR (SR) and BR (SR/RM2+), 0.16 and 0.23

for the StringTie2 runs to 0.43 for the TSEBRA runs. The BUSCO scores decreased slightly

post-filtering (1-2%). EnTAP annotation rates ranged from 66% to 86%, with the TSB

(SR/OrthoDB), which is an improvement from 59% to 68% pre-filtering.

In terms of BUSCO completeness and mono:multi ratios, the two best performing runs (BR (SR)

and BR (SR/LR)) were further filtered (Table 4). In this step, multi-exonic genes without an

EggNOG alignment or a sequence similarity alignment through EnTAP were removed. These

filtered models were re-assessed for mono:multi ratio, BUSCO completeness, and EnTAP

annotation. The BUSCO completeness remained the same for BR(SR), but not for BR(SR/LR).

The EnTAP annotation increased from 66% to 81% in BR (SR), and 67% to 87% in BR (SR/LR).

DISCUSSION

BRAKER (Hoff et al., 2020) and MAKER (Cantarel et al., 2008) are currently the most popular

eukaryotic structural annotation tools, cited 475 and 1,010 times (since 2021, as referenced in

Google Scholar). Processes that select from multiple *ab initio* or aligned forms of evidence are

gaining popularity as well though they add both time and complexity to the analyses (FINDER

cited 22 times, (Banerjee et al., 2021); EVidenceModeler cited 381 times (Haas et al., 2008)).

Finally, as high-throughput transcriptomics, in the form of both short and long-read evidence

become more accessible, rapid approaches like StringTie2 (cited 451 times (Kovaka et al.,

2019)) are occasionally used as the exclusive approach, though more often, used in

combination with the options listed above.

Regardless of the methods selected, recently published benchmarks are challenged to achieve

high values for gene sensitivity in larger genomes (Brůna et al., 2021). Within smaller and less

complex model systems such as *C. elegans* and *D. melanogaster*, *ab initio* prediction results in

gene sensitivity 49.8% and 59.5%, respectively (Brůna et al., 2021). In well studied complex

organisms, such as humans, gene level sensitivity and specificity hovers at 48% and 43%,

respectively (Banerjee et al., 2021). While generating benchmarks with model systems (*A.*

thaliana, *C. elegans*, and *D. melanogaster*) provides more reliable metrics for comparison, they

are infamous for not fully representing the diversity of their respective clades (Chang et al.,

2016).

This study focused on four gene prediction workflows: StringTie2, MAKER, BRAKER, and

BRAKER/TSEBRA, and examined the process across a variety of evidence inputs. Both model

and non-model plant genomes were considered to highlight the challenges and reinforce the

need for downstream filtering.

Genome annotation benchmarks for both models and non-models-

Among plant genomes, the total number of genes is relatively conserved and ranges from

20,000 to just over 40,000. As such, total gene number provides an accessible preliminary

benchmark. However, the number of genes in the reference annotation fails to assess the

overall quality of the annotation. To measure this, we should consider additional metrics. Here,

we describe the utility of BUSCO score, mono-exonic: multi-exonic ratio, and sequence

similarity assessment.

BUSCO allows us to identify complete, duplicated, fragmented, and missing single-copy

orthologs shared by most seed plants (Simão et al. 2015; Seppey, Manni, and Zdobnov 2019).

This provides a reliable benchmark in the absence of a high quality reference annotation and

poor BUSCO scores are immediately indicative of a larger issue. However, a high BUSCO

score is not sufficient to estimate the quality of an annotation (Fig 4B). Six of the 18 BRAKER

runs and four of the 17 StringTie2 runs exceeded 95% completeness. However, total gene

number, gene length, and structure varied considerably.

Repeat content, especially in the form of LTRs, and pseudogenes can lead to inflated gene

model estimates, especially in the form of mono-exonic genes (Scott et al., 2020; Trouern-Trend

et al., 2020). We expect that eukaryotes maintain 20% or less of their gene space as mono-

exonics (Jain et al., 2008; Table S3). Although the BUSCO scores were consistent, we note

tremendous variation in mono- to multi-gene model ratios post-BRAKER. In practice, having a

worse mono: multi ratio is preferable to having a lower BUSCO score, since missing genes,

especially those thought to be conserved, cannot be easily rectified, while putative false

positives can potentially be filtered through other means.

Sequence similarity search metrics are more complex to interpret, but when used with high-

quality and curated databases that contain full-length proteins (e.g. NCBI RefSeq), can provide

a benchmark. Specifically, a reciprocal BLAST search requires that both the query and target in

the search retain a minimum level of coverage in the alignment. For new plant genomes, that

are in the darkest corners of the tree of life, this might be a less reliable metric. For species that

may fare poorly in database comparisons, searches for protein domains can provide some level

of confidence and we demonstrate this as a filter to reduce the mono-exonics in *Liriodendron.*

Masking repeats in plant genomes: Repeat masking is important but may not require

additional LTR resolution to improve performance-

Plant genomes typically contain many repeats, mostly in the form of transposable elements

(TEs), averaging around 46% (Luo et al., 2022). Given the abundance of TEs in genomes, it is

important to mask these in advance of gene prediction. Soft masking involves changing

nucleotides identified as repeats to lowercase (Yandell & Ence, 2012), signaling downstream

programs to ignore these sequences. Of the five genomes included in this study, *Liriodendron*

had the largest genome size and repeat content. Running downstream analyses on an

unmasked genome of *Liriodendron* resulted in a 4-fold increase in gene predictions (Fig 5A).

Many repeats were identified as putative gene models, resulting in a large increase of total

number of genes (Fig 5B).

RepeatModeler2 is a widely used tool for TE discovery (Flynn et al., 2020). The recent release

of RepeatModeler2 includes an optional module for more robust LTR structural detection

(LTRStruct module) that includes the LTRharvest (Ellinghaus et al., 2008), LTRDetector

(Valencia & Girgis, 2019), and LTR_retriever packages (Ou & Jiang, 2018). This is particularly

useful in identifying more divergent LTRs in the genome that may exist in fewer copies (Ou &

Jiang, 2018; Valencia & Girgis, 2019). Among the default packages included, RepeatScout

serves as a fast method to detect young and abundant repeat families in the genome. RECON,

on the other hand, is more computationally intensive and is sensitive enough to detect older TE

families. The LTRStruct module is run on the unmasked genome to identify LTR families that

may be redundant with the families identified by the default package. This creates redundancy

that is resolved through clustering with CD-HIT (Flynn et al., 2020).

In the four species compared, additional repeat masking did not significantly improve gene

predictions (Table S9; Fig 4). The mono: multi ratios across species were consistent before and

after additional LTR masking (Fig 5A). The BUSCO completeness scores remained relatively

the same, with BR (SR/RM2+) being 1% higher than BR (SR) in *Arabidopsis, Funaria* and

Populus. The marginal improvement observed in these genomes could be related to the

structure and type of LTRs, for example, better identification of divergent Ty1-copia elements

described in the *Funaria* genome (Kirbis et al., 2022). While we did not include genomes with

excessive repeat estimates (>70%), our results indicated that the optional LTRStruct module

was not beneficial.

Genome-guided transcriptome assembly for annotation: Transcripts derived directly

from alignments are not sufficient to annotate reference genomes-

Transcriptome assemblers are designed to work with primarily short RNA-Seq reads to

construct full-length transcripts. In the presence of a high-quality reference genome, genome-

guided approaches are preferred as the reads are aligned directly to the target genome in

advance. Aligned RNA evidence provides resolution on exon boundaries, and aids in the

identification of splice variants. *De novo* approaches build graph models directly from the short

(or long) reads to generate transcripts. The latter is much more challenging, computationally

intensive, and prone to error.

We compared the accuracy of the annotations produced by StringTie2; *de novo* assembled

transcripts with Trinity to annotations produced by BRAKER. The selected packages are top

performers when compared in their respective categories of genome-guided and *de novo*

transcriptome assembly (Sahraeian et al., 2017; Venturini et al., 2018). As expected, Trinity

produced a higher number of transcripts than StringTie2, and BUSCO completeness was

consistently lower (Table 3), except for *Liriodendron*. The gene models generated by StringTie2

were more numerous than the BRAKER gene models, more than expected for each species. It

should be noted, however, that StringTie2 identifies splice variants by generating a splice graph

and resolving conflict between multiple potential splice sites (Kovaka et al., 2019), whereas

BRAKER trains an internal algorithm GeneMark-ET to find specific genes with complete support

among all introns to be further used in training Augustus (Hoff et al., 2019).

StringTie2 runs resulted in lower BUSCO completeness when compared to BRAKER and/or

TSEBRA runs (Fig 4A; Table S11). This outcome is supported by the lack of *ab initio* prediction

with genome-guided approaches. Inflated mono-exonic predictions (and lower BUSCO scores)

were also observed in the StringTie2 genome annotation of the water strider (*Microvelia*

longipes) (Toubiana et al., 2021). In our study, *Rosa* ST2 (SR/LR) run was closest with a

BUSCO score of 97.2%, BR (SR/LR) of 96.9%, and TSB (SR/ST2) of 98% (Table S9, S10).

Including proteins: Genome annotations are improved for models,and closely related

species, when full-length proteins sourced from OrthoDB are used in combination with

read data.

The performance of StringTie2 and Trinity-derived protein evidence was assessed on the

predicted gene models using BRAKER and TSEBRA. In this context, the genome-guided or *de*

novo assembled transcripts were translated into proteins and provided as evidence to train the

ab initio component of the pipelines. Adding protein evidence to genome annotation can target

protein-coding genes leading to more accurate predictions than RNA-Seq evidence alone

(Bruna, 2022). This study specifically focused on using protein evidence derived in some

fashion from the transcriptomic inputs but also evaluated the recommendation to include clade

specific OrthoDB protein inputs to the BRAKER/TSEBRA approach.

The TSEBRA runs of the model species, *Arabidopsis* and *Populus* were compared to the

reference annotations. These runs were the best for the model species in terms of sensitivity

and specificity as compared to the MAKER, StringTie2, Trinity and BRAKER runs (Fig 3B). The

model genomes also had very similar BUSCO completeness scores but had different mono:

multi ratios with the addition of protein evidence. As expected, the model genomes benefitted

the most from the inclusion of the external OrthoDB proteins in terms of annotation rate and

BUSCO score (both the *Arabidopsis* and *Populus* reference proteins are contained within this

resource). However, mono: multi ratio challenges remained consistent across the TSEBRA runs

with varying inputs.

In the case of non-model plant genomes, TSEBRA contributed to higher mono:multi ratios, and

this was very evident in *Liriodendron*. However, the BUSCO scores of the non-protein runs were

lower across all runs of *Liriodendron*, compared to the protein runs. The *Rosa* TSB

(SR/OrthoDB) reported the highest BUSCO score across all runs; given the phylogenetic

placement of *Rosa* in comparison to *Arabidopsis*, we believe that the addition of OrthoDB

greatly influenced this result. On the other hand, the annotation rate of TSB (SR/OrthoDB)

remained like the other runs within *Rosa*. The higher quality of the *Rosa* genome assembly,

compared to the other two non-models, could also influence the utility of the protein evidence.

However, mono: multi ratios remained to be high, and annotation rates were like those of the

runs without OrthoDB proteins.

TSEBRA runs with proteins sourced from genome-guided predictions perform similarly, but had

lower BUSCO, higher mono: multi ratios, and total gene number when compared to the short-

read only runs (Fig 4A, Table S9). Among TSEBRA runs, Trinity fares better only for

Liriodendron, which could indicate that genome-guided proteins are not a suitable choice for a

more repetitive genome. This is consistent in independent assessments between *de novo*

transcriptome assemblers and genome-guided assemblers with complex genomes with

fragmented genome assemblies (for example in *Ae. albopictus* (Huang et al., 2016). TSEBRA

with full length proteins sourced from OrthoDB had lower BUSCO scores when compared to

BR(SR) for the non-model species, *Liriodendron* and *Funaria.* In both cases, TSB

(SR/OrthoDB).

The total number of genes predicted by TSEBRA and BRAKER runs remained largely the same

across all species (Table S9). However, the number of mono-exonic genes increased, whereas

the multi-exonic genes decreased across all TSEBRA runs in comparison to the BRAKER runs

without proteins across all species. The average gene lengths also decreased, while average

lengths of mono-exonics remained the same, and the lengths of multi-exonics are higher.

Initial examination of the EnTAP reciprocal BLAST assessment revealed high annotation rates

for the non-model species when protein evidence was included, particularly the multi-exonics

(whereas the mono-exonic percentage remained the same) (Table S9). However, this increase

in multi-exonic annotation proved to be an artifact since the total number of multi-exonic models

was reduced. Direct comparison of the predictions revealed that 40%-52% of the multi-exonics

were split into mono-exonic predictions when comparing the BR(SR) to the TSB (SR/ST2) and

TSB (SR/OrthoDB) gene models predicted using *Liriodendron* (Table S14).

Long-read transcriptomes: Long-reads can be paired with short-reads to improve the

quality of the resulting models-

Long-reads generated from platforms such as Oxford Nanopore or PacBio have the potential to

resolve splice variants and assemble transcripts more accurately than traditional Illumina RNA-

Seq (Amarasinghe et al., 2020). While long-reads can independently generate transcriptomes, it

is recommended to have a combination of short and long reads to achieve greater depth,

improved error profiles, and gain more evidence for splice site resolution (Amarasinghe et al.,

2020; Gonzalez-Ibeas et al., 2016; Watson & Warr, 2019).

In this study, we utilized both ONT and Iso-Seq long-reads. In the latter, we relied on raw reads

(not the error-corrected CCS reads) in our comparisons for genome annotations using long-

reads. In all cases, long-reads (alone) did not outperform short-reads for the BRAKER runs.

However, in some cases, the combination of short-read and long-read inputs was beneficial.

The higher error rate Iso-Seq reads from *Populus* and *Liriodendron* produced comparable, but

lower, BUSCO scores compared to the BR (SR) runs. In contrast, the ONT long-reads used for

Arabidopsis and *Rosa* in the combined runs (BR (SR/LR)) had slightly better BUSCO

completeness as compared to the BR (SR) runs, and similar mono:multi ratios. Overall, the

lower error profile of using ONT reads, supplemented with short-read data, as well as using high

quality reference genomes, support the higher BUSCO completeness scores.

Best Practices for Plant Genome Annotation-

Given existing tools, we recommend that investigators utilize RepeatModeler2 to mask their

genome of interest with the default settings (Flynn et al. 2020). Following soft-masking, RNA-

Seq short reads (between 4-10 libraries, paired-end, minimum 15M reads per library) are

generally sufficient for annotation. While we did not comprehensively investigate the impact of

tissue type, it is recommended to sample from multiple tissues when possible (Kress et al.,

2022). In our study, we did not observe a difference in the annotation completeness among

species with a higher number of short-read libraries, although we did not comprehensively

evaluate the difference of using fewer libraries within a single species.

Sequencing of long-read libraries remains more expensive than generating deep Illumina short-

read RNA-Seq. In most cases, the short reads were sufficient as input. The notable exceptions

include the BR (SR/LR), as they were comparable, and in some cases, slightly improved, over

BR (SR) across all species. The lower error-profile Nanopore reads were more beneficial when

combined with short-reads. However, current long-read technologies available from both

platforms may provide different results.

BRAKER and TSEBRA outperformed runs of MAKER, StringTie2 and Trinity with default

settings. It should be noted that the authors did not comprehensively benchmark MAKER with

more than two training runs of AUGUSTUS as recommended, which could have further

improved results. However, previous benchmarking studies also support lower performance of

MAKER (Banerjee et al., 2021; Hoff et al., 2020). Among the BRAKER runs executed in the

model plants, *Arabidopsis* and *Populus*, the TSEBRA runs were the best runs. TSEBRA,

especially TSB (SR/OrthoDB), performs the best for *Rosa* but would require substantial filtering

to remove false positives. Among the less contiguous and more evolutionary distant species

(*Funaria* and *Liriodendron*), BR (SR) runs performed the best in terms of BUSCO completeness

and mono:multi ratios. Overall EnTAP annotation rates were greatly improved in runs where

OrthoDB proteins were included as evidence. However, when considering BUSCO as well as

mono: multi ratio, especially for non-model species, the BR(SR) runs performed the best. For

more divergent species (as defined by current public databases), BRAKER runs with short-

reads, or short-reads and long-reads, are advised.

Regardless of approach, existing pipelines do not provide appropriate summary statistics or

robust methods for filtering unlikely gene models. All methods produce more putative false

positives than desired. We recommend utilizing reciprocal BLAST searches with well curated

databases containing targets with full-length proteins (such as NCBI's RefSeq) to identify

fragmented models. We also recommend filtering and removing mono-exonics that do not have

a protein domain. Finally, we recommend structural filters to remove unlikely gene structures

(splice sites, start sites, incompletes, etc).

In this study, we demonstrated the impact of post-filtering on the most complex genome

assessed in this study, *Liriodendron*. We improved the published annotation across all

benchmarks evaluated in this study following a new BR(SR) run (Table 4; (Chen et al., 2019)).

The filters reduced the overall number of putative false positives and increased the overall rate

of annotation, with minimal reduction to BUSCO completeness.

Data Availability: All scripts and data used is available through

https://www.protocols.io/blind/3A33C8E3B76511EC84CA0A58A9FEAC02 . The public data

(NCBI SRA and genome assembly accessions) for the reference genomes, short-reads, and

long-reads are listed in Table S2.

Acknowledgements

The authors would like to thank the Institute for Systems Genomics (ISG) and Computational

Biology Core at UConn for HPC services. Thanks go to Dr. Bernard Goffinet for his careful

review of this manuscript. This work was supported by the National Science Foundation awards,

DEB-1753811 and DBI-1943371.

LITERATURE CITED

Amarasinghe, S. L., Su, S., Dong, X., Zappia, L., Ritchie, M. E., & Gouil, Q. (2020).

Opportunities and challenges in long-read sequencing data analysis. *Genome Biology*,

21(1), 30.

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data. Available

online. *Retrieved May*, *17*, 2018.

Arabidopsis Genome Initiative. (2000). Analysis of the genome sequence of the flowering plant

Arabidopsis thaliana. *Nature*, *408*(6814), 796–815.

Armisén, D., Lecharny, A., & Aubourg, S. (2008). Unique genes in plants: specificities and

conserved features throughout evolution. *BMC Evolutionary Biology*, *8*, 280.

Banerjee, S., Bhandary, P., Woodhouse, M., Sen, T. Z., Wise, R. P., & Andorf, C. M. (2021).

FINDER: an automated software package to annotate eukaryotic genes from RNA-Seq

data and associated protein sequences. *BMC Bioinformatics*, *22*(1), 205.

Bolger, M. E., Arsova, B., & Usadel, B. (2018). Plant genome and transcriptome annotations:

from misconceptions to simple solutions. *Briefings in Bioinformatics*, *19*(3), 437–449.

Bruna, T. (2022). *Unsupervised algorithms for automated gene prediction in novel eukaryotic*

genomes. https://smartech.gatech.edu/handle/1853/67297

Brůna, T., Hoff, K. J., Lomsadze, A., Stanke, M., & Borodovsky, M. (2021). BRAKER2:

automatic eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported

by a protein database. *NAR Genomics and Bioinformatics*, *3*(1), lqaa108.

Caballero, M., & Wegrzyn, J. (2019). gFACs: Gene Filtering, Analysis, and Conversion to Unify

Genome Annotations Across Alignment and Gene Prediction Frameworks. *Genomics,*

Proteomics & Bioinformatics, *17*(3), 305–310.

Campbell, M. S., Holt, C., Moore, B., & Yandell, M. (2014). Genome Annotation and Curation

Using MAKER and MAKER-P. *Current Protocols in Bioinformatics / Editoral Board,*

Andreas D. Baxevanis ... [et Al.], *48*, 4.11.1–39.

Campbell, M. S., Law, M., Holt, C., Stein, J. C., Moghe, G. D., Hufnagel, D. E., Lei, J.,

Achawanantakun, R., Jiao, D., Lawrence, C. J., Ware, D., Shiu, S.-H., Childs, K. L., Sun,

Y., Jiang, N., & Yandell, M. (2014). MAKER-P: A Tool Kit for the Rapid Creation,

Management, and Quality Control of Plant Genome Annotations. *Plant Physiology*, *164*(2),

513–524.

Cantarel, B. L., Korf, I., Robb, S. M. C., Parra, G., Ross, E., Moore, B., Holt, C., Sánchez

Alvarado, A., & Yandell, M. (2008). MAKER: an easy-to-use annotation pipeline designed

for emerging model organism genomes. *Genome Research*, *18*(1), 188–196.

Chang, C., Bowman, J. L., & Meyerowitz, E. M. (2016). Field Guide to Plant Model Systems.

Cell, *167*(2), 325–339.

Cheng, S., Melkonian, M., Smith, S. A., Brockington, S., Archibald, J. M., Delaux, P.-M., Li, F.-

W., Melkonian, B., Mavrodiev, E. V., Sun, W., Fu, Y., Yang, H., Soltis, D. E., Graham, S.

W., Soltis, P. S., Liu, X., Xu, X., & Wong, G. K.-S. (2018). 10KP: A phylodiverse genome

sequencing plan. *GigaScience*, *7*(3), 1–9.

Chen, J., Hao, Z., Guang, X., Zhao, C., Wang, P., Xue, L., Zhu, Q., Yang, L., Sheng, Y., Zhou,

Y., Xu, H., Xie, H., Long, X., Zhang, J., Wang, Z., Shi, M., Lu, Y., Liu, S., Guan, L., … Shi,

J. (2019). Liriodendron genome sheds light on angiosperm phylogeny and species-pair

differentiation. *Nature Plants*, *5*(1), 18–25.

Corchete, L. A., Rojas, E. A., Alonso-López, D., De Las Rivas, J., Gutiérrez, N. C., & Burguillo,

F. J. (2020). Systematic comparison and assessment of RNA-seq procedures for gene

expression quantitative analysis. *Scientific Reports*, *10*(1), 19737.

Deutekom, E. S., Vosseberg, J., van Dam, T. J. P., & Snel, B. (2019). Measuring the impact of

gene prediction on gene loss estimates in Eukaryotes by quantifying falsely inferred

absences. *PLoS Computational Biology*, *15*(8), e1007301.

Edgar, R. (2010). *Usearch*. https://www.osti.gov/biblio/1137186

Ellinghaus, D., Kurtz, S., & Willhoeft, U. (2008). LTRharvest, an efficient and flexible software

for de novo detection of LTR retrotransposons. *BMC Bioinformatics*, *9*, 18.

Flynn, J. M., Hubley, R., Goubert, C., Rosen, J., Clark, A. G., Feschotte, C., & Smit, A. F.

(2020). RepeatModeler2 for automated genomic discovery of transposable element

families. *Proceedings of the National Academy of Sciences of the United States of America*,

117(17), 9451–9457.

Gabriel, L., Hoff, K. J., Brůna, T., Borodovsky, M., & Stanke, M. (2021). TSEBRA: transcript

selector for BRAKER. *BMC Bioinformatics*, *22*(1), 566.

Gonzalez-Ibeas, D., Martinez-Garcia, P. J., Famula, R. A., Delfino-Mix, A., Stevens, K. A.,

Loopstra, C. A., Langley, C. H., Neale, D. B., & Wegrzyn, J. L. (2016). Assessing the Gene

Content of the Megagenome: Sugar Pine (Pinus lambertiana). *G3* , *6*(12), 3787–3802.

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X.,

Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A.,

Rhind, N., di Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K., … Regev, A. (2011).

Full-length transcriptome assembly from RNA-Seq data without a reference genome.

Nature Biotechnology, *29*(7), 644–652.

Gremme, G. (2014). *GenomeThreader Gene Prediction Software*.

https://genomethreader.org/doc/gthmanual.pdf

Gurevich, A., Saveliev, V., Vyahhi, N., & Tesler, G. (2013). QUAST: quality assessment tool for

genome assemblies. *Bioinformatics* , *29*(8), 1072–1075.

Haas, B. J., Salzberg, S. L., Zhu, W., Pertea, M., Allen, J. E., Orvis, J., White, O., Buell, C. R., &

Wortman, J. R. (2008). Automated eukaryotic gene structure annotation using

EVidenceModeler and the Program to Assemble Spliced Alignments. *Genome Biology*,

9(1), R7.

Hart, A. J., Ginzburg, S., Xu, M. S., Fisher, C. R., Rahmatpour, N., Mitton, J. B., Paul, R., &

Wegrzyn, J. L. (2020). EnTAP: Bringing faster and smarter functional annotation to non-

model eukaryotic transcriptomes. *Molecular Ecology Resources*, *20*(2), 591–604.

Hoff, K. J., Brŭna, T., Lomsadze, A., & Stanke, M. (2020). Fully Automated and Accurate

Annotation of Eukaryotic Genomes with BRAKER2. *Poster Presented at*.

https://www.researchgate.net/profile/Katharina-Hoff-

2/publication/338831355_Fully_Automated_and_Accurate_Annotation_of_Eukaryotic_Gen

omes_with_BRAKER2/links/5e2d9102299bf152167f6424/Fully-Automated-and-Accurate-

Annotation-of-Eukaryotic-Genomes-with-BRAKER2.pdf

Hoff, K. J., Lange, S., Lomsadze, A., Borodovsky, M., & Stanke, M. (2016). BRAKER1:

Unsupervised RNA-Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS.

Bioinformatics , *32*(5), 767–769.

Hoff, K. J., Lomsadze, A., Borodovsky, M., & Stanke, M. (2019). Whole-Genome Annotation

with BRAKER. *Methods in Molecular Biology* , *1962*, 65–95.

Holt, C., & Yandell, M. (2011). MAKER2: an annotation pipeline and genome-database

management tool for second-generation genome projects. *BMC Bioinformatics*, *12*, 491.

Huang, X., Chen, X.-G., & Armbruster, P. A. (2016). Comparative performance of transcriptome

assembly methods for non-model organisms. *BMC Genomics*, *17*, 523.

Jain, M., Khurana, P., Tyagi, A. K., & Khurana, J. P. (2008). Genome-wide analysis of intronless

genes in rice and Arabidopsis. *Functional & Integrative Genomics*, *8*(1), 69–78.

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J.,

Mitchell, A., Nuka, G., Pesseat, S., Quinn, A. F., Sangrador-Vegas, A., Scheremetjew, M.,

Yong, S.-Y., Lopez, R., & Hunter, S. (2014). InterProScan 5: genome-scale protein function

classification. *Bioinformatics* , *30*(9), 1236–1240.

Joshi NA, F. J. N. (2011). *Sickle: A sliding-window, adaptive, quality-based trimming tool for

FastQ files* (Version Version 1.33) [Computer software]. https://github.com/najoshi/sickle

Jung, H., Ventura, T., Chung, J. S., Kim, W.-J., Nam, B.-H., Kong, H. J., Kim, Y.-O., Jeon, M.-

S., & Eyun, S.-I. (2020). Twelve quick steps for genome assembly and annotation in the

classroom. *PLoS Computational Biology*, *16*(11), e1008325.

Kersey, P. J. (2019). Plant genome sequences: past, present, future. *Current Opinion in Plant*

Biology, *48*, 1–8.

Kim, D., Paggi, J. M., Park, C., Bennett, C., & Salzberg, S. L. (2019). Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology*, *37*(8),

907–915.

Kirbis, A., Rahmatpour, N., Dong, S., Yu, J., van Gessel, N., Waller, M., Reski, R., Lang, D.,

Rensing, S. A., Temsch, E. M., Wegrzyn, J. L., Goffinet, B., Liu, Y., & Szövényi, P. (2022).

Genome dynamics in mosses: Extensive synteny coexists with a highly dynamic gene

space. In *bioRxiv* (p. 2022.05.17.492078). https://doi.org/10.1101/2022.05.17.492078

Korf, I. (2004). Gene finding in novel genomes. *BMC Bioinformatics*, *5*(1), 59.

Kovaka, S., Zimin, A. V., Pertea, G. M., Razaghi, R., Salzberg, S. L., & Pertea, M. (2019).

Transcriptome assembly from long-read RNA-seq alignments with StringTie2. *Genome*

Biology, *20*(1), 278.

Kress, W. J., Soltis, D. E., Kersey, P. J., Wegrzyn, J. L., Leebens-Mack, J. H., Gostel, M. R.,

Liu, X., & Soltis, P. S. (2022). Green plant genomes: What we know in an era of rapidly

expanding opportunities. *Proceedings of the National Academy of Sciences of the United*

States of America, *119*(4). https://doi.org/10.1073/pnas.2115640118

Lewin, H. A., Richards, S., Lieberman Aiden, E., Allende, M. L., Archibald, J. M., Bálint, M.,

Barker, K. B., Baumgartner, B., Belov, K., Bertorelle, G., Blaxter, M. L., Cai, J., Caperello,

N. D., Carlson, K., Castilla-Rubio, J. C., Chaw, S.-M., Chen, L., Childers, A. K., Coddington,

J. A., … Zhang, G. (2022). The Earth BioGenome Project 2020: Starting the clock.

Proceedings of the National Academy of Sciences of the United States of America, *119*(4).

https://doi.org/10.1073/pnas.2115635118

Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* , *34*(18),

3094–3100.

Li, H. (2021). New strategies to improve minimap2 alignment accuracy. *Bioinformatics* .

https://doi.org/10.1093/bioinformatics/btab705

Luo, X., Chen, S., & Zhang, Y. (2022). PlantRep: a database of plant repetitive elements. *Plant*

Cell Reports, *41*(4), 1163–1166.

Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., & Zdobnov, E. M. (2021). BUSCO

Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic

Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Molecular Biology*

and Evolution, *38*(10), 4647–4654.

Marks, R. A., Hotaling, S., Frandsen, P. B., & VanBuren, R. (2021). Representation and

participation across 20 years of plant genome sequencing. *Nature Plants*, *7*(12), 1571–

1578.

Meyer, C., Scalzitti, N., Jeannin-Girardon, A., Collet, P., Poch, O., & Thompson, J. D. (2020).

Understanding the causes of errors in eukaryotic protein-coding gene prediction: a case

study of primate proteomes. *BMC Bioinformatics*, *21*(1), 513.

Mudge, J. M., & Harrow, J. (2016). The state of play in higher eukaryote gene annotation.

Nature Reviews. Genetics, *17*(12), 758–772.

Musich, R., Cadle-Davidson, L., & Osier, M. V. (2021). Comparison of Short-Read Sequence

Aligners Indicates Strengths and Weaknesses for Biologists to Consider. *Frontiers in Plant*

Science, *12*, 657240.

Ou, S., & Jiang, N. (2018). LTR_retriever: A Highly Accurate and Sensitive Program for

Identification of Long Terminal Repeat Retrotransposons. *Plant Physiology*, *176*(2), 1410–

1422.

Pertea, G., & Pertea, M. (2020). GFF Utilities: GffRead and GffCompare. *F1000Research*, *9*.

https://doi.org/10.12688/f1000research.23297.2

Pertea, M., Pertea, G. M., Antonescu, C. M., Chang, T.-C., Mendell, J. T., & Salzberg, S. L.

(2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads.

Nature Biotechnology, *33*(3), 290–295.

Pucker, B., Irisarri, I., de Vries, J., & Xu, B. (2022). Plant genome sequence assembly in the era

of long reads: Progress, challenges and future directions. *Quantitative Plant Biology*, *3*, e5.

Quevillon, E., Silventoinen, V., Pillai, S., Harte, N., Mulder, N., Apweiler, R., & Lopez, R. (2005).

InterProScan: protein domains identifier. *Nucleic Acids Research*, *33*(Web Server issue),

W116–W120.

Sahraeian, S. M. E., Mohiyuddin, M., Sebra, R., Tilgner, H., Afshar, P. T., Au, K. F., Bani Asadi,

N., Gerstein, M. B., Wong, W. H., Snyder, M. P., Schadt, E., & Lam, H. Y. K. (2017).

Gaining comprehensive biological insight into the transcriptome by performing a broad-

spectrum RNA-seq analysis. *Nature Communications*, *8*(1), 59.

Salzberg, S. L. (2019). Next-generation genome annotation: we still struggle to get it right.

Genome Biology, *20*(1), 92.

Scott, A. D., Zimin, A. V., Puiu, D., Workman, R., Britton, M., Zaman, S., Caballero, M., Read, A.

C., Bogdanove, A. J., Burns, E., Wegrzyn, J., Timp, W., Salzberg, S. L., & Neale, D. B.

(2020). A Reference Genome Sequence for Giant Sequoia. *G3* , *10*(11), 3907–3919.

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015).

BUSCO: assessing genome assembly and annotation completeness with single-copy

orthologs. *Bioinformatics* , *31*(19), 3210–3212.

Smit, AFA, Hubley, R & Green, P. (2013-2015). *RepeatMasker Open-4.0*. RepearMasker.

http://www.repeatmasker.org

Stanke, M., & Waack, S. (2003). Gene prediction with a hidden Markov model and a new intron

submodel. *Bioinformatics* , *19 Suppl 2*, ii215–ii225.

Sun, Y., Shang, L., Zhu, Q.-H., Fan, L., & Guo, L. (2022). Twenty years of plant genome

sequencing: achievements and challenges. In *Trends in Plant Science* (Vol. 27, Issue 4, pp.

391–401). https://doi.org/10.1016/j.tplants.2021.10.006

Toubiana, W., Armisén, D., Dechaud, C., Arbore, R., & Khila, A. (2021). Impact of male trait

exaggeration on sex-biased gene expression and genome architecture in a water strider.

BMC Biology, *19*(1), 89.

Trouern-Trend, A. J., Falk, T., Zaman, S., Caballero, M., Neale, D. B., Langley, C. H.,

Dandekar, A. M., Stevens, K. A., & Wegrzyn, J. L. (2020). Comparative genomics of six

Juglans species reveals disease-associated gene family contractions. *The Plant Journal:*

For Cell and Molecular Biology, *102*(2), 410–423.

Valencia, J. D., & Girgis, H. Z. (2019). LtrDetector: A tool-suite for detecting long terminal repeat

retrotransposons de-novo. *BMC Genomics*, *20*(1), 450.

Venturini, L., Caim, S., Kaithakottil, G. G., Mapleson, D. L., & Swarbreck, D. (2018). Leveraging

multiple transcriptome assembly methods for improved gene structure annotation.

GigaScience, *7*(8). https://doi.org/10.1093/gigascience/giy093

Watson, M., & Warr, A. (2019). Errors in long-read assemblies can critically affect protein

prediction [Review of *Errors in long-read assemblies can critically affect protein prediction*].

Nature Biotechnology, *37*(2), 124–126.

Wu, T. D., & Watanabe, C. K. (2005). GMAP: a genomic mapping and alignment program for

mRNA and EST sequences. *Bioinformatics* , *21*(9), 1859–1875.

Yandell, M., & Ence, D. (2012). A beginner's guide to eukaryotic genome annotation. *Nature*

Reviews. Genetics, *13*(5), 329–342.

200   FIGURES

Figure 1: Genome size, repeat content, and BUSCO completeness for the five plant genomes:

Arabidopsis, Populus, Funaria, Rosa, and Liriodendron. Each pie represents the BUSCO

completeness. Green denotes the completeness score, orange indicates the fragmented score,

and blue indicates the missing score from BUSCO. (A) *BUSCO scores estimated from the*

published assemblies. *(B) BUSCO scores estimated from protein-coding gene predictions from*

*the **published annotations**.*

210 *Figure 2: Comparing metrics between BRAKER (blue) and StringTie2 (red) predictions. (A)*

211 *mono:multi ratios, (B) BUSCO comparisons, and (C) EnTAP annotation rates of the gene models.*

212 *The yellow region indicates the ideal value for each of the metrics.*

Figure 3: Comparison of BUSCO, sensitivity, and false positive rates between the Arabidopsis

and Populus annotations (Table S10). (A) BUSCO completeness scores for the MK(SR/RM2+)

and BR(SR/RM2+) runs of Arabidopsis and Populus, green denotes the completeness score,

orange indicates the fragmented score, and blue indicates the missing score from BUSCO (B)

False positive rates and sensitivity scores from Mikado against published annotations for

Arabidopsis (red color) and Populus (gold color) for the MAKER, BRAKER, TSEBRA, Trinity, and

StringTie2 runs. The scores were assessed using MIKADO. Multiple points per run reflect

differences in input read type and repeat-masking.

Figure 4: Comparison of BUSCO completeness *scores (A), mono:multi ratios (B), and EnTAP*

annotation rates (C) across all species between the runs of different input types and software,

i.e., MAKER (MK-green) BRAKER (BK-light blue),TSEBRA (TSB-dark blue) and StringTie2 (ST2-

red). The yellow rectangle represents the target scores for each benchmark. RM2+-

RepeatModeler2 with LTRStruct.

Figure 5: (A) The effect of soft-masking on gene prediction in Liriodendron (Table S13).

Performing structural annotation on the unmasked Liriodendron genome results in inflation in the

mono and multi- exonic genes. Blue denotes the BRAKER (BR) runs for both genomes, SR

denotes short-reads, and LR denotes long reads. The lighter shade represents mono-exonics,

and the darker shade represents the multi-exonics. (B) More genes predicted using the unmasked

genome (blue), as compared to only one gene predicted in this region with the masked genome

(red). The green track shows the LTR elements in the genome as identified by RepeatModeler2.

The RNA alignment reads show a read pile-up at the predicted gene (masked track).

TABLES

Table 1: Notations for the different runs performed for benchmarking.

Run		*Arabidopsis*	*Funaria*	*Populus*	*Liriodendron*	*Rosa*
StringTie2						
ST2 (SR)	Short-reads	X	X	X	X	X
ST2 (LR)	Long-reads	X		X	X	X
ST2(SR/LR)	Short and long-reads	X		X	X	X
BRAKER						
BR (SR)	Short-reads	X	X	X	X	X
BR (LR)	Long-reads	X		X	X	X
BR (SR/LR)	Short and long-reads	X		X	X	X
BR (SR/RM2+)	Short-reads with additional masking for LTRs	X	X	X	X	
TSEBRA						
TSB (SR/TRINITY)	Short-reads and *de novo* proteins	X	X	X	X	X
TSB (SR/ST2)	Short-reads and genome-guided proteins	X	X	X	X	X
TSB (LR/ST2)	Long-reads and genome-guided proteins	X		X	X	X
TSB (SR/LR/ST2)	Short and long-reads and genome-guided proteins	X		X	X	X
TSB (SR/ST2/RM2+)	Short-reads and genome-guided proteins with additional masking for LTRs	X	X	X	X	
TSB (SR/OrthoDB)	Short-reads and OrthoDB proteins	X	X	X	X	X
MAKER						
MK (RM2+)	Short-reads with additional masking for LTRs	X	X	X		

245   SR- Short reads, LR- Long reads, and RM2+- RepeatModeler2 with the additional repeat masking.

247   Table 2: Genome assembly and annotation statistics for the five published plant genomes

Species	Genome size	Total scaffolds (chromosomes)	N50	Repeat content	BUSCO Completeness	
					Genome	Annotation
Arabidopsis thaliana	119 Mb	7 (5)	23.46 Mb	23.6%	99.30%	99.60%
Funaria hygrometrica	327 Mb	687 (26)	1.48 Mb	42.35%	85.60%	86.60%
Liriodendron chinense	1,742 Mb	3,711 (21)	3.53 Mb	73.18%	98.60%	75.10%
Populus trichocarpa	434 Mb	1,446 (19)	19.47 Mb	35.90%	98.80%	98.30%
Rosa chinensis	515 Mb	55 (7)	69.64 Mb	60.53%	98.80%	97.30%

249 Table 3: Comparison between genome-guided (StringTie2- ST2) and de novo (Trinity) genome

250 annotations.

Species	RM %	RM2+ %	Total short-reads (total libraries)	Total long-reads (total libraries)	Total Trinity transcripts (N50)	Total ST2 transcripts (SR) (N50)	Total ST2 transcripts (LR) (N50)	Total ST2 transcripts (SR/LR) (N50)	BUSCO transcript alignments (%)		
									Trinity (SR)	ST2 (SR)	ST2 (SR/LR)
Arabidopsis	15.2	16.5	511,277,126 (9)	23,134,068 (4)	319,434 (2726)	37,747 (2538)	36,241 (1599)	42,265 (1363)	82.7	95.5	93.6
Funaria	42.3	43.1	549,205,030 (9)		151,265 (1198)	59,741 (369)			72.9	84.5	
Liriodendron	73.2	72.7	1,408,831,670 (20)	10,437,029 (1)	2,839,867 (3055)	62,341 (1041)	33,895 (1815)	45,785 (2361)	92.5	87.1	77.3
Populus	35.9	45.1	267,403,772 (5)	161,334 (1)	283,572 (1837)	56,468 (402)	20,633 (1074)	37,222 (1869)	71.3	73.3	65.3
Rosa	60.5		134,461,068 (4)	41,929,383 (6)	812,407 (2187)	53,708 (672)	74397 (1866)	105,639 (1605)	88.8	97	97.2

Species	*Arabidopsis*	*Funaria*	*Liriodendron*	*Populus*	*Rosa*
RM%	23.6	42.35	73.18	35.9	60.53
RM2+%	16.5	43.1	72.7	45.1	
Total short-reads (total libraries)	511,277,126 (9)	549,205,030 (9)	1,408,831,670 (20)	267,403,772 (5)	134,461,068 (4)
Total long-reads (total libraries)	23,134,068 (4)		10,437,029 (1)	161,334 (1)	41,929,383 (6)
Total Trinity transcripts before aligning to genome (N50)	319,434 (2726)	151,265 (1198)	2,839,867 (3055)	283,572 (1837)	812,407 (2187)
Total Trinity transcripts after aligning to genome (N50)	47,024 (1410)	55,085 (1383)	152,255 (1335)	78,774 (1320)	95,016 (1272)
Total ST2 transcripts (SR) (N50)	37,747 (2538)	59,741 (369)	62,341 (1041)	56,468 (402)	53,708 (672)

Vuruputoor et al. *Welcome to the big leaves: best practices for improving genome annotation in non-model plant genomes.*

Total ST2 transcripts (LR) (N50)		36,241 (1599)		33,895 (1815)	20,633 (1074)	74397 (1866)
Total ST2 transcripts (SR/LR) (N50)		42,265 (1363)		45,785 (2361)	37,222 (1869)	105,639 (1605)
BUSCO (%)	**Trinity (SR)**	82.7	72.9	92.5	71.3	88.8
	ST2 (SR)	95.5	84.5	87.1	73.3	97
	ST2 (SR/LR)	93.6		77.3	65.3	97.2

254 SR denotes short reads, LR for Long reads, RM for RepeatModeler2, and RM2+ for RepeatModeler2 with the LTRStruct

255 flag, RM% indicates the repeat content masked with RepeatModeler2, and RM2+% is the repeat content masked with

256 RepeatModeler2 with the LTRStruct flag.

257 Table 4: Gene model statistics for Liriodendron after two rounds of structural and functional filters.

Liriodendron Annotation	Total genes	Mono:Multi Ratio	BUSCO %	EnTAP %
Published annotation	35261	0.7	75.1	63
Mono-exonic filters				
BR (LR)	39031	0.21	87.4	69
BR (SR) *	41065	0.16	90.2	66
BR (SR/LR) *	40420	0.16	90.3	67
BR (SR/RM2+)	40740	0.17	88.2	67
ST2 (SR)	51804	0.16	86.5	80
ST2 (LR)	27012	0.23	65	84
ST2 (SR/LR)	36345	0.24	70.6	82
TSB (LR/ST2)	33132	0.43	82.3	84
TSB (SR/LR/ST2)	33964	0.43	82.4	84
TSB (SR/ST)	32898	0.41	83.4	84
TSB (SR/ST2/RM2+)	33637	0.45	82.8	84
TSB (SR/TRINITY)	34646	0.42	84	83
TSB (SR/OrthoDB)	33667	0.43	86.5	86
+Multi-exonic filters				
BR (SR)	30219	0.24	90.3	81
BR (SR/LR)	30035	0.23	86.9	87

258 ** denotes the two best annotation sets*

259 BR- BRAKER, ST2- StringTie2, TSB- TSEBRA, SR- Short reads, LR- Long-reads, RM2+- RepeatModeler2 with

260 LTRStruct. BUSCO % shows the completeness percentage, and EnTAP % is the annotation rates after reciprocal

261 BLAST.