Captive rearing effects on the methylome of Atlantic salmon after oceanic migration: sex-

specificity and intergenerational stability

Running title: Hatchery-induced DNA methylation in *S. salar*

Clare J. Venney[1,2], Raphaël Bouchard[1,2], Julien April[3], Eric Normandeau[1,2], Laurie Lecomte[1,2],

Guillaume Côté[3], Louis Bernatchez[1,2]

1. Institut de Biologie Intégrative et des Systèmes (IBIS), Université Laval, Québec,

Canada, G1V 0A6

2. Département de Biologie, Université Laval, Québec, Canada, G1V 0A6

3. Direction de l'expertise sur la faune aquatique, Ministère des Forêts, de la Faune et des

Parcs du Québec, Québec, Canada, G1S 4X4

Corresponding author: Clare Venney (clarevenney@gmail.com)

Submitted to Special Issue: *Omics Tools for Species Conservation*

Keywords: hatchery rearing, salmonid, epigenetics, conservation

Abstract

Captive rearing in salmon hatcheries can have considerable impacts on both fish

phenotype and fitness within a single generation, even in the absence of genetic change. Evidence

for hatchery-induced changes in DNA methylation is becoming abundant, though questions

remain on the sex-specificity of these effects, their persistence until spawning, and potential for

transmission to future generations. Here we performed whole genome methylation sequencing of

fin tissue for 16 hatchery and 16 wild Atlantic salmon (*Salmo salar*) returning to spawn in the

Rimouski River, Québec. We identified two cohorts of hatchery-reared salmon through

methylation analysis, one of which was epigenetically similar to wild fish, suggesting that

supplementation efforts may be able to minimize the epigenetic effects of hatchery rearing. We

found considerable sex-specific effects of hatchery rearing, with few genomic regions being

affected in both males and females. We also analysed the methylome of 32 F1 offspring from

four groups (pure wild, pure hatchery origin, and reciprocal hybrids). We found that few

epigenetic changes due to parental hatchery rearing persisted in the F1 offspring though the

patterns of inheritance appear to be complex, involving nonadditive effects. Our results suggest

that the epigenetic effects of hatchery rearing can be minimal in F0. There may also be minimal

epigenetic inheritance and rapid loss of epigenetic changes associated with hatchery rearing.

However, due to sex-specificity and nonadditive patterns of inheritance, methylation changes due

to captive rearing are rather complex and the field would benefit from further research on

minimizing the epigenetic effects of captive rearing in conservation efforts.

Introduction

Many natural populations are in decline due to human activities (Jenkins, 2003), leading to conservation efforts to bolster populations. Certain economically or culturally important fish species are particularly threatened by human-induced changes such as habitat loss, overexploitation, and climate change (Arthington, Dulvy, Gladstone, & Winfield, 2016). Captive breeding is often used both to replenish populations in decline and to supplement populations for recreational fishing purposes (Fraser, 2008), with over 300 species of fish reared in hatcheries for supplementation efforts worldwide (Brown & Day, 2002). This often involves the capture of wild organisms, controlled breeding in hatcheries, and release of offspring into natural systems at various life stages. Salmonids have been a strong focus of supplementation efforts due to their economic and recreational value (Fraser, 2008).

Despite the high costs and ecological importance of salmonid supplementation, hatchery rearing has resulted in unintended phenotypic and fitness consequences. Hatchery-reared salmon often have altered phenotypes and reduced fitness relative to wild fish. A multispecies meta-analysis showed that hatchery fish had 50% lower reproductive success than wild fish (Christie, Ford, & Blouin, 2014) while a multidecade monitoring study in Ireland showed that hatchery-reared Atlantic salmon (*Salmo salar*) had 36% the lifetime reproductive success of wild conspecifics (O'Sullivan et al., 2020). Decreases in reproductive success due to hatchery rearing depend on life stage at release (Bouchard, Wellband, Lecomte, Bernatchez, & April, 2022; Milot, Perrier, Papillon, Dodson, & Bernatchez, 2013) and are generally more pronounced in male salmon, potentially due to reduced sexual selection and higher rates of precocious maturity in hatchery environments (Christie et al., 2014). Hatchery rearing is also associated with phenotypic

changes in salmon, including reduced swimming capacity (Pedersen, Koed, & Malte, 2008),

differences in heart morphology and function (Frisk et al., 2020; Leonard & McCormick, 2001),

and altered gut microbiota (Lavoie, Courcelle, Redivo, & Derome, 2018). These phenotypic

changes often arise after a single generation of captive rearing, prompting interest in the

molecular causes of phenotypic and fitness differences.

Initially, hatchery rearing was thought to induce genetic changes, though genetic changes

are often minimal and transcriptional changes are more pronounced. A single generation of

hatchery rearing is generally insufficient to cause genetic differentiation between captive and

wild salmon (Gavery, Nichols, Goetz, Middleton, & Swanson, 2018; Le Luyer et al., 2017),

though SNPs associated with domestication have been detected after several generations of

hatchery rearing in Atlantic salmon (Harder & Christie, 2022). However, the hatchery

environment has been shown to induce transcriptional changes in steelhead (*Oncorhynchus*

mykiss; Christie, Marine, Fox, French, & Blouin, 2016), Atlantic salmon (Frisk et al., 2020), and

Coho salmon (*O. kisutch*; Leitwein et al., 2022). This led to the idea that epigenetic mechanisms

controlling gene expression, such as DNA methylation, may serve as the molecular mechanisms

underlying rapid phenotypic shifts and fitness declines in hatchery environments.

DNA methylation refers to the addition of methyl groups to the DNA, generally at CpG

cytosines in vertebrates (Anastasiadi, Venney, Bernatchez, & Wellenreuther, 2021). Unlike

genetic variation, DNA methylation can rapidly change in response to environmental cues and

can influence gene expression and phenotype (reviewed in Anastasiadi et al., 2021). Methylation

differences have been reported in captive fish strains reared in typical and enriched environments

(Berbel-Filho et al., 2020; Venney, Wellband, & Heath, 2021) and in European sea bass

(*Dicentrarchus labrax*) undergoing domestication (Anastasiadi, Piferrer, & Wittkopp, 2019).

There is also a growing body of literature supporting the effects of hatchery rearing on the

methylome (Gavery et al., 2019, 2018; Koch, Nuetzel, & Narum, 2022; Le Luyer et al., 2017;

Leitwein et al., 2021; Rodriguez Barreto et al., 2019; Wellband, Roth, Linnansaari, Curry, &

Bernatchez, 2021). A recent review on the epigenetic effects of hatchery rearing found that these

methylation differences were often targeted to genes involved in stress and immune response,

growth, and neural development, consistent with the added stress and environmental simplicity of

hatchery environments (Koch et al., 2022). It is also possible that methylation changes associated

with captive rearing could affect multiple generations as DNA methylation patterns are often

heritable (Anastasiadi et al., 2021). Methylation changes in the sperm of captive-reared fish have

been reported (Gavery et al., 2018; Rodriguez Barreto et al., 2019; Wellband et al., 2021) and can

persist through oceanic migrations until salmon return to spawn (Leitwein et al., 2021). One

study showed epigenetic inheritance due to parental captive rearing in Atlantic salmon, though

this study used smolt-to-adult supplementation, an alternative rearing technique where salmon are

captured as smolts and held until adulthood (Wellband et al., 2021). In this study, many somatic

methylation changes present in F0 were not passed to F1, though novel methylation changes were

present in F1 and were associated with phenotypic changes (Wellband et al., 2021). The evidence

for epigenetic effects of hatchery rearing is growing and has potential long-term,

multigenerational implications for supplemented fish stocks.

However, some knowledge gaps on the effects of captive rearing on the methylome

remain. Despite clear sex-specific effects of hatchery rearing on fitness (Christie et al., 2014), it

remains unclear whether hatchery-induced methylation changes also differ between the sexes. If

sex-specific methylation changes occur and persist until spawning, they could be passed on to

offspring. This could lead to complex patterns of inheritance in F1, particularly when reciprocal

hybrids (hatchery mother x wild father or wild mother x hatchery father) occur in supplemented

populations. Thus, there is a need to assess the sex-specificity of hatchery-induced methylation

changes and the potential for transmission to offspring to understand the stability and complexity

of the effects of captive rearing on natural populations. In this study, we assess (i) whether the

epigenetic effects of early-life hatchery rearing persist until salmon return to freshwater to spawn,

(ii) whether there were sex-specific effects of hatchery rearing on DNA methylation, (iii) to what

extent these methylation changes were passed on to offspring, and (iv) if there were biases in the

inheritance of these marks due to maternal, paternal or nonadditive effects. This study expands

our knowledge of the epigenetic effects of captive rearing that may lead to fitness declines

through supplementation efforts. It also underlines the importance of considering epigenetic

modifications, including their stability and heritability, in conservation management decisions

when assessing the impacts of supplementation efforts.

Materials and Methods

Parent and offspring sampling and DNA extraction

We studied the Rimouski River population of Atlantic salmon (*Salmo salar*), located on

the south shore of the St Lawrence River in Québec, Canada. Stocking has been ongoing in this

river from 1992 to 2021. In this system, wild-caught fish are trapped at an impassable dam each

year and transported past the dam to a 21 km length of upstream spawning grounds. Some of the

captured fish are used as broodstock for the supplementation effort each year and transported to

the Québec Government hatchery in Tadoussac. From 2011 to 2016, the offspring of these

crosses were stocked in the river as young of the year parr with their adipose fins removed for

future identification.

The samples used here were obtained from a previous study on the reproductive success

of wild and hatchery-reared fish, with detailed sampling information available in Bouchard et al.

(2022). In 2018, a fin clip was taken from all returning adults captured and transported up the

dam to their spawning grounds. Hatchery-reared adults were identified by the absence of their

adipose fin. In the following summer (July-August 2019), fry were sampled in the upstream

spawning area through electrofishing and preserved in 95% ethanol. DNA was extracted from

parent and offspring fin tissue using a salt-based DNA extraction method (Aljanabi & Martinez,

1997). While fin is less biologically active than other tissues, its use allowed for non-lethal and

minimally invasive sampling of adults returning to spawn. Due to the tissue specificity of DNA

methylation (Gavery et al., 2018; Venney, Johansson, & Heath, 2016), we used the same tissue

for the offspring to allow comparisons of DNA methylation between generations.

Parentage analysis

Parentage analysis was performed in Bouchard et al. (2022). Briefly, parental sex was

verified using a PCR-based genetic sex marker for Atlantic salmon (King & Stevens, 2020) to

inform parentage possibilities. Parent and offspring samples were amplified at 52 microsatellite

loci using multiplex PCR panels 1a and 1b from Bradbury et al. (2018) in a two-step PCR

protocol (Zhan et al., 2017). The first step used a Qiagen Multiplex Master Mix to amplify each

panel independently using adaptor-tagged primers (for specific PCR conditions and reagent

concentrations, see Bouchard et al. (2022)). The panels were then pooled for each sample and

cleaned using Quanta Bio SparQ PureMag beads. The cleaned products were used in the second

PCR step which ligated barcodes to the primer adaptors to allow multiplexing of samples during

sequencing. Sequencing was performed at the Institut de Biologie Intégrative et des Systèmes

(IBIS) at Université Laval, Québec. Libraries were diluted to 10-12 pM and single-end

sequencing was performed using an Illumina MiSeq with the MiSeq Reagent Kit V3 with 150

cycles and dual indexing to increase multiplexing capacity.

Demultiplexing of sequence data was performed with the MiSeq Sequence Analysis

software. MEGASAT was used to determine allele lengths (Zhan et al., 2017) with a minimum

sequence depth of 20 reads per locus per sample required to call allele sizes. Lengths were

confirmed or modified based on histogram outputs. After all processing, loci were excluded if

they had more than 10% missing data. Parentage analysis was performed in Cervus v3.0

(Kalinowski, Taper, & Marshall, 2007) using a 90% confidence likelihood to find the most

probable mother-offspring dyad.

Whole genome methylation sequencing

Based on parentage analysis, we selected 32 parent and 32 offspring DNA samples for

whole genome bisulfite sequencing (Figure S1). For the parents, we selected 16 distinct breeding

pairs that produced offspring: eight mating pairs of wild fish and eight pairs of hatchery-origin

fish. We selected eight F1 offspring from four distinct groups based on parentage: (i) pure wild

origin, (ii) pure hatchery origin, (iii) wild dam and hatchery-origin sire, and (iv) hatchery-origin

dam and wild sire. The 16 pure strain offspring were the product of the 16 parental breeding

pairs, and no offspring shared parentage. The reciprocal hybrids allowed us to determine if the

effects of hatchery rearing were parent-specific; if these groups differ, they indicate that hatchery

rearing affects males and females differently and these effects can be passed on to offspring. For

each sample, fin DNA extractions used for microsatellite analysis were quantified, quality-

checked, and sent to the Centre d'expertise et de services at Génome Québec, Montréal, Canada,

where library preparation and sequencing were performed. Whole genome Methyl-Seq was

performed across four sequencing lanes on an Illumina NovaSeq6000 using S4 flow cells and

paired-end 150 bp chemistry with anticipated ~15X coverage for each sample.

Methylation data processing

Methylation data processing was performed using a pipeline available at https://github.

com/enormandeau/bwa-meth_pipeline. Sequence data were quality trimmed using fastp (Chen,

Zhou, Chen, & Gu, 2018) to remove sequences under 100 bp and with phred scores less than 25,

and to remove the first and last base of each read. Trimmed data were aligned to the North

American Atlantic salmon genome (GCA_923944775.1) using bwa-meth (https://github.com/

brentp/bwa-meth). Duplicates alignments were removed using Picard tools *MarkDuplicates*

(https://github.com/ broadinstitute/picard). MethylDackel (https://github.com/dpryan79/Methyl-

Dackel) *mbias* was used to trim the beginnings and ends of reads which often result in biased

methylation calls, then *extract* was used to tabulate DNA methylation for each CpG site in each

individual. Paired end reads were then merged to produce bedGraph and methylKit files for

further analysis. Pipeline available at https://github.com/enormandeau/bwa-meth_pipeline.

SNP masking and coverage filtration

C/T SNPs (and A/G SNPs for the G position of CpG sites) prevent accurate methylation

calling. It is not possible to determine whether a T read is due to a true unmethylated cytosine call

or a thymine mutation present at that CpG site. Whole genome SNP data were obtained from an

upcoming study (Lecomte et al. in prep) on two proximal populations of Atlantic salmon in the

Romaine and Puyjalon Rivers, Québec, to mask C/T SNPs from our methylation data. While

whole genome SNP data are not available for the Rimouski River population, these two

populations are geographically and genetically close to our studied population (Figure S2) and

should suffice for masking most SNPs in our dataset. Fin clip DNA was extracted for 29

Romaine River and 31 Puyjalon River Atlantic salmon. Samples were sent for whole genome

sequencing at Génome Québec on an Illumina NovaSeq6000, with anticipated 16X coverage.

Raw sequence data were processed using the pipeline available at

https://github.com/enormandeau/wgs_sample_preparation.

Bcftools *mpileup* and *call* (Danecek et al., 2021) were used to generate genotype

likelihoods from bam files with minimum mapping quality of 5. SNP calls were then filtered with

bcftools *filter* and vcftools (Danecek et al., 2011) to require minimum mapping quality of 30,

minimum quality scores greater than 30, minimum genotype quality over 20, minimum

sequencing depth of five reads, minor allele count of two or greater, maximum two alleles,

maximum 70% missing data, and minor allele frequency ≥ 0.05. SNP calling and filtering scripts

are available at https://github.com/LaurieLecomte/SNP_calling_pipeline_202106. A list of C/T

and A/G SNPs was exported as a bed file and bedtools (Quinlan & Hall, 2010) *intersect -v* was

used to remove these SNPs from bedGraph and methylKit files.

After this, we split the methylation dataset by generation for all future filtering and

analyses. BedGraph and methylKit files were quickly filtered to remove CpG sites with less than

5 or more than 100 reads, then we retained only CpG sites with sufficient coverage in at least

80% of individuals (26 of 32) for each generation. Scripts are available at

https://github.com/cvenney/dss_pipeline.

Principal component and redundancy analyses

We used principal component analysis (PCA) and redundancy analysis (RDA) for each

generation to identify overall trends in whole genome methylation. MethylKit files were

imported into methylKit (Akalin et al., 2012) in R (R Core Team, 2022) and united into a single

data frame for each generation with only CpG sites without missing data included. PCAs were

performed using the *prcomp* function to identify overall trends in the F0 dataset due to parental

sex and origin (i.e., wild or hatchery), and in the F1 dataset due to maternal and paternal origin.

We also performed an RDA for each dataset using vegan (Oksanen et al., 2020) to determine the

significance of model terms at the whole genome level. We tested for the effects of origin, sex,

and their interaction in F0 and the effect of maternal origin, paternal origin, and their interaction

in F1. Model significance was tested with PERMANOVA-like tests with 1000 iterations and the

significance of individual terms was also determined for each analysis.

Coverage filtration and differential methylation analysis

DSS (Feng & Wu, 2019) was used to smooth methylation data over 500 bp regions and

implement models to test for differentially methylated loci (DMLs, i.e., CpG sites) and

differentially methylated regions (DMRs, identified by a high frequency of DMLs within a

region). For all analyses, DMLs were considered significant if they had an FDR-adjusted p-value

less than 0.05. DMRs were called based on DML p-values before FDR adjustment, as

recommended by the developers of DSS, which may lead to the identification of fewer DMLs

than DMRs. We further filtered the DMR results, requiring a minimum length of 100 bp, at least

10 CpGs with at least 50% of them identified as DMLs (p<0.05). Finally, we merged DMRs

within 50 bp of one another.

For the F0 samples, we tested for both overall and sex-specific effects of hatchery rearing.

For the overall effects of hatchery rearing, a generalized linear model (GLM) was used to test for

the influence of origin on DNA methylation in the F0 while controlling for sex and the source by

sex interaction. Based on the results and heatmaps for the F0, we split the hatchery parents into

two cohorts: (i) seven "wildtype" fish whose DMR methylation patterns resembled wild fish, and

(ii) nine "normal" hatchery fish whose DMR methylation patterns differed from wild fish (Figure

S3). We consider epigenetic divergence between wild and hatchery-reared fish to be "normal"

due to considerable evidence for the effects of hatchery rearing on the methylome of salmonids

(e.g., reviewed in Koch et al. (2022)). We reran the DSS analysis for three separate F0

comparisons: comparing each hatchery subset independently to the wild fish and comparing the

hatchery subsets to one another. We also split the full F0 dataset by sex and tested for sex-

specific effects of hatchery rearing using a Wald test for the effect of origin for each sex,

requiring a minimum delta (group difference in methylation) of 10% for both DMLs and DMRs.

A GLM was used to test for maternal and paternal origin effects and their interaction in the F1.

The pipeline is available at https://github.com/cvenney/dss_pipeline.

Annotation and gene ontology enrichment analysis

Gene ontology annotation was performed with the DMR lists obtained from DSS. We

annotated the genome using the GCF_905237065.1_Ssal_v3.1 transcripts file for the publicly

available Atlantic salmon genome (https://www.ncbi.nlm.nih.gov/genome/?term=txid8030[orgn])

with GAWN v0.3.5 (https://github.com/enormandeau/gawn). We identified transcripts proximal

to (±5 kb) the DMR positions from each DSS analysis and performed GO enrichment tests using

the go_enrichment v1.1.0 pipeline (https://github.com/enormandeau/go_enrichment). The

pipeline blasts the targets transcript against the Swiss-Prot database, obtains GO annotation

information from the UniProt database, then uses GOATOOLS (Klopfenstein et al., 2018) to find

significantly enriched GO terms.

Results

We obtained an average of 301 153 191 raw methylation reads per sample and an average

of 173 158 755 successful alignments. After all quality filtering, we retained a total of 25 451 589

CpG sites for the parental analysis and 31 440 242 for offspring (see Table S1 for detailed

coverage information).

Principal component and redundancy analyses

We analyzed 60 179 CpG sites with no missing data for F0 and 65 347 for F1 with PCA

and RDA. We assessed methylation variation using PCA analysis and found whole genome

methylation differences between hatchery reared and wild F0 salmon. The "wildtype" hatchery

cohort identified in DMR analysis, as outlined in the methods, was intermediate between the wild

and "normal" hatchery cohort (Figure 1A). Little differentiation between F1 groups was observed

at the whole genome level (Figure 1B).

The F0 RDA model was significant (p=0.002, adjusted R^2=0.023), as was the effect of

source (p=0.001), though sex and the interaction effect were not significant (p=0.37 and p=0.26,

respectively). The overall F1 RDA model was not significant (p=0.65), nor were the terms in the

model (maternal source p=0.20, paternal source p=0.74, interaction p=0.87) indicating no

significant whole genome effects of parental rearing environment.

275   *Differential methylation analysis: overall F0 effects of hatchery rearing*

276   DSS analysis with all F0 samples identified two cohorts of hatchery fish, one "wildtype"

cohort that clustered with wild fish and one "normal" hatchery cohort that differed from the wild

and wildtype fish when controlling for sex and the source by sex interaction (Figure S3). For the

full dataset, we found 165 DMLs and 266 DMRs due to origin. However, since there may be

multiple age classes present in the hatchery fish and different manipulations have been performed

on hatchery-reared fish by hatchery personnel, we split the F0 dataset into three groups: the two

hatchery cohorts identified in the heatmap and PCA analysis, and the wild fish.

283   We identified 84 431 DMLs and 2 007 DMRs between wild and "normal" hatchery

284   salmon (Figure 2A, Tables S2 and S3), with clear division between the two groups based on the

285   dendrogram. GO enrichment analysis showed the DMRs were associated with organ

286   morphogenesis, skeletal morphogenesis (e.g., GO:0035138 pectoral fin morphogenesis), and

287   metabolic processes (Table S4).

288   We found 34 049 DMLs and 1 285 DMRs between "normal" and "wildtype" hatchery

salmon (Figure 2B, Tables S5 and S6) with the two groups separating as predicted in the

dendrogram other than SASA-31 and SASA-380, which were also the intermediate hatchery

"wild type" points overlapping the hatchery "normal" polygon in Figure 1A. Enriched GO terms

were associated with angiogenesis, intestinal morphology, skeletal morphology, and

transcriptional processes (Table S7).

We only detected 84 DMLs and 156 DMRs between wild and "wildtype" hatchery

salmon (Figure 2C, Tables S8 and S9) with no clear separation of groups in the heatmap,

indicating that the "wildtype" hatchery cohort has a methylation profile similar to that of wild

fish. Despite the small number of DMRs, there was functional enrichment for various

transcriptional processes (Table S10).

Differential methylation analysis: sex-specific F0 effects of hatchery rearing

We found considerable sex-specific effects of hatchery rearing in the F0. We detected

260 844 DMLs and 5 717 DMRs between male hatchery-reared and wild salmon (Figure 3A,

Tables S11 and S12). We observed half the number of DMLs and DMRs due to hatchery rearing

in female fish, with 100 451 DMLs and 2 439 DMRs due to source (Figure 3B, Tables S13 and

S14), though both groups showed clear differentiation based on source in the heatmaps. The

dendrograms also showed separate clustering of "normal" and "wildtype" female fish, though the

cohorts were less differentiated in males (the "wildtype" samples SASA-31 and SASA-380

grouped with the "normal" hatchery males). We found only 734 overlaps between the male and

female sex-specific analyses, indicating that most DMRs identified are sex-specific. There were

39 male-specific and 42 female-specific enriched GO terms with many overlaps between the

sexes. Terms were associated with RNA synthesis and transcription for both sexes (e.g.,

GO:0010468 regulation of gene expression, GO:2001141 regulation of RNA biosynthetic

process, GO:1903506 regulation of nucleic acid-templated transcription, GO:0006357 regulation

of transcription by RNA polymerase II; Tables S15 and S16).

Differential methylation analysis: F1 effects of parental hatchery rearing

The effects of hatchery rearing were diluted in the F1 fish compared to the F0. We

observed 46 DMLs and 284 DMRs due to maternal source (Figure 4A, Tables S17 and S18), with

clear division of offspring with wild dams and hatchery-reared sires, and hatchery-reared dams

and hatchery-reared sires (offspring with wild sires were mixed regardless of maternal source).

We found 11 DMLs and 254 DMRs due to paternal source (Figure 4B, Tables S19 and S20),

with division occurring in offspring with hatchery dams based on paternal source, similar to the

maternal DMRs. There were three DMLs and 271 DMRs due to the interaction between maternal

and paternal source, with clear differentiation between 'pure' (two wild or two hatchery-reared

parents) and hybrid offspring (Figure 4C, Tables S21 and S22). No GO terms were significantly

enriched due to maternal source after filtering results. The paternal source DMRs showed

enrichment for integral component of membrane (GO:0016021) and the interaction effect DMRs

showed enrichment for regulation of cellular process (GO:0050794).

Discussion

While the epigenetic effects of hatchery rearing in salmonids are well-documented

(Gavery et al., 2019, 2018; Koch et al., 2022; Le Luyer et al., 2017; Leitwein et al., 2021, 2022;

Rodriguez Barreto et al., 2019; Venney et al., 2021; Wellband et al., 2021), the long-term

stability of these changes after release into natural systems has not been extensively studied, nor

has the sex-specificity of the effects of hatchery rearing on the methylome. Here we showed that

hatchery rearing led to general and sex-specific changes in Atlantic salmon fin DNA methylation

that persisted until the salmon returned to spawn, consistent with similar results in Coho salmon

returning to freshwater after eighteen months at sea (Leitwein et al., 2021). However, fewer

methylation changes persisted to F1, indicating that the epigenetic effects of hatchery rearing

primarily affected first-generation hatchery fish, though the fitness consequences of hatchery

rearing may be persistent and insidious (O'Sullivan et al., 2020; Willoughby & Christie, 2019).

Methylation analysis identified two hatchery cohorts

Two cohorts of hatchery salmon were identified through both PCA for whole genome

methylation and DMR analysis (Figures 1A, 2A-C, S2) with the "wildtype" cohort more

epigenetically similar to wild fish than the "normal" hatchery cohort. The latter was named

"normal" due to the expectation that hatchery rearing influences the methylome based on the

current literature (e.g., reviewed in Koch et al., 2022). The cohorts were not related to sex or time

spent at sea (one winter vs. multiple winters at sea) and there were no particular rearing

manipulations performed on the F0 during their time in the hatchery. However, hatchery origin

fish were released at different locations down the Rimouski River. Therefore, they could have

experienced different conditions during the final stage of transportation and stocking (e.g., due to

temperature). Each year, some parr were released in the same areas as wild conspecifics whereas

others were stocked in a section of river upstream of a waterfall where wild salmon parr are not

found. Regrettably, we could not determine the release site of each fish as they were not tagged

or otherwise identifiable after release. Thus, it is possible that the cohorts exhibit different

methylation patterns due to differences in early life environment after stocking, with "wildtype"

hatchery salmon assuming methylation patterns similar to wild fish due to a shared environment.

It is also possible that hatchery rearing had a negligible effect on methylation in our study and the

epigenetic differences between groups are due to differences in the environment where they were

released. However, even the "wildtype" hatchery fish show methylation differences compared to

wild fish, and we observed epigenetic inheritance of these effects in F1 regardless of parental

cohort, indicating that the hatchery environment does influence the methylome. Indeed, hatchery

rearing has been shown to affect the methylome even after oceanic migration in Coho salmon

(Leitwein et al., 2021). However, it is possible that the residual effects of hatchery rearing were

diminished due to the shared river environment after release. It is also possible that other tissues

would retain the methylation changes associated with hatchery rearing, though we used fin in our

study to allow nonlethal sampling of F0 during their spawning migration.

The functional significance of DMRs between the two hatchery cohorts was consistent

with previously reported phenotypic effects of hatchery rearing, further suggesting that the

methylation differences were due to the hatchery environment rather than the river environment.

GO enrichment analysis showed that the "normal" and "wildtype" hatchery fish exhibited

methylation differences at regions associated with angiogenesis, intestinal morphology, skeletal

morphology, and transcriptional processes. A previous study comparing mangrove killifish

(*Kryptolebias marmoratus*) reared in both barren tanks and environments enriched with logs and

plants reported methylation changes in genes involved in angiogenesis (Berbel-Filho, Rodríguez-

Barreto, Berry, Garcia De Leaniz, & Consuegra, 2019). Another study using Rimouski salmon

showed that hatchery rearing alters the gut microbiota (Lavoie et al., 2018) which may be

associated with morphological changes (Liu, Wang, Ou, Hou, & He, 2020). Thus, the two

hatchery cohorts may have diverged due to differences in early life environment with the

"normal" cohort harbouring methylation changes associated with the typical phenotypic effects of

hatchery rearing.

Comparisons between wild and "normal" hatchery fish DMRs also showed functional

enrichment for organ and skeletal morphogenesis including pectoral fin morphogenesis, and

metabolic processes. Hatchery-reared Atlantic salmon have lower swimming performance

relative to wild conspecifics (Pedersen et al., 2008). Thus, differences in fin DNA methylation

between "normal" hatchery fish and wild fish are associated with biological functions that may

underly previously reported differences in swimming capacity, though further studies in other

tissues are needed for a mechanistic link.

In contrast, "wildtype" hatchery fish and wild fish were epigenetically similar relative to

other comparisons, though methylation differences were associated with transcriptional

regulation. Therefore, the epigenetic effects of hatchery rearing could potentially be minimized

through supplementation efforts. We suspect this is due to the "wildtype" hatchery fish sharing

an environment with wild fish after stocking, though regrettably we cannot definitively determine

the causative factor(s) leading to epigenetic similarities between "wildtype" hatchery and wild

fish. Further experimental investigation would be needed to determine how to minimize the

epigenetic effects of hatchery rearing. Overall, we show that the epigenetic effects of early

rearing environment are not short-term acclimatory responses, but rather lead to long-term effects

on the methylation states of salmon that persist until spawning (i.e., after two to three years living

in rivers followed by one to two years spent at sea).

Sex-specific effects of hatchery rearing

We provide evidence that hatchery rearing affects the epigenome in a sex-specific way,

further complicating our understanding of the effects of hatchery rearing on the methylome. We

identified 5 717 male-specific and 2 439 female-specific F0 DMRs due to rearing environment,

with only 734 DMRs shared between the sexes. Sex-specific methylation differences have been

shown to occur in fish (Fellous et al., 2018; Laing et al., 2018; Podgorniak, Brockmann,

Konstantinidis, & Fernandes, 2019). In Nile tilapia (*Oreochromis niloticus*), early domestication

(i.e., wild-born fish transferred to a hatchery for domestication) led to sex-specific methylation

responses to the hatchery environment, with only 57 of ~1000 DMLs shared between male and

female fish (Podgorniak et al., 2019). Transcriptional differences between the sexes have been

reported in brook charr (*Salvelinus fontinalis*; Sutherland, Prokkola, Audet, & Bernatchez, 2019)

and are associated with DNA methylation differences in zebrafish (*Danio rerio*; Laing et al.,

2018). While hatchery fish generally have lower reproductive success than wild conspecifics

regardless of sex (Bouchard et al., 2022; Christie et al., 2014; O'Sullivan et al., 2020), a previous

meta-analysis showed that hatchery rearing had a greater effect on the reproductive success of

males than females when compared to wild fish of the same sex (Christie et al., 2014). Female

hatchery fish tend to have higher reproductive success in the wild than male hatchery fish

(Berntson, Carmichael, Flesher, Ward, & Moran, 2011; Christie et al., 2014; Milot et al., 2013;

Thériault, Moyer, Jackson, Blouin, & Banks, 2011; Williamson, Murdoch, Pearsons, Ward, &

Ford, 2010) and thus molecular responses to hatchery rearing are likely complicated by sex.

While the genomic regions affected by hatchery rearing differ between sexes, GO enrichment

analysis showed that the regions were associated with many of the same processes in both males

and females, with 35 of 39 enriched male-specific GO terms also present in the female analysis.

Due to the potential for epigenetic inheritance, sex-specific epigenetic effects could also lead to

parental rearing environment affecting offspring epigenetics in complex ways based on which

parent(s) experience the hatchery environment, and whether maternal or paternal effects occur in

specific regions of the genome.

Dilution of epigenetic effects of hatchery rearing in F1

Despite the persistent effects of hatchery rearing in F0, we observed relatively little

inheritance to F1 based on parental environment: there were few F1 DMLs and DMRs based on

parental source (Figure 4A-B) and there was no evidence for genome-wide methylome

differentiation (Figure 1B). While there is evidence for epigenetic inheritance across taxa, it is

unclear to what extent the epigenome is inherited, though there is often a dilution of inherited

effects over generations (Anastasiadi et al., 2021). Studies have shown that hatchery rearing can

influence sperm methylation (Gavery et al., 2018; Leitwein et al., 2021; Wellband et al., 2021),

though somatic methylation changes are often not present in the F0 germline (Anastasiadi et al.,

2021; Gavery et al., 2019, 2018) and the genomic regions showing differential methylation often

differ between F0 sperm and F1 somatic tissues (Wellband et al., 2021). Thus, despite

considerable epigenetic effects of hatchery rearing in F0, many of these changes are not passed

on to F1.

The few methylation differences that were inherited based on parental rearing

environment exhibited complex, nonadditive effects that may be difficult to account for during

conservation efforts. When testing for maternal influences on offspring methylation, differences

were primarily between offspring from wild and hatchery dams with hatchery-reared sires,

whereas offspring from wild sires were similar regardless of maternal origin (Figure 4A). The

same effect was observed when testing for paternal influences, with paternal rearing environment

mainly affecting offspring methylation when the dams were of hatchery origin (Figure 4B). This

suggests nonadditive epigenetic inheritance based on hatchery rearing wherein the origin of both

parents influences the DNA methylation state of offspring, rather than simple maternal or

paternal effects. A study in Chinook salmon (*Oncorhynchus tshawytscha*) showed that hatchery-

origin fish had greater nonadditive effects on DNA methylation when reared in a semi-natural

environment as opposed to a hatchery environment (Venney et al., 2021), though a study on

reciprocal hybrid steelhead (*O. mykiss*; hatchery dam and wild sire, or wild dam and hatchery

sire) identified additive, heritable effects on gene expression (Christie et al., 2016). Thus, is it

possible that the complexity of the natural environment led to complicated epistatic effects

resulting in nonadditive effects on methylation, though the link between DNA methylation and

gene expression may be complex (Christensen et al., 2021).

Unexpectedly, the maternal by paternal interaction effect DMRs showed two clusters: the

pure strain (hatchery-hatchery and wild-wild) and hybrid offspring. While hybridization can often

cause atypical methylation patterns compared to pure strains (Gimenez, Vazquez, Trepat,

Cambiaso, & Rodríguez, 2021; Raza et al., 2017; Shen et al., 2012), it was surprising that pure

wild and pure hatchery-origin offspring had similar methylation patterns at these loci. It is

possible that hybridization between wild and hatchery salmon led to dysregulation of epigenetic

marks, though we find this an unlikely explanation after a single generation of hatchery rearing.

Overall, these complex nonadditive patterns of inheritance based on parental environment

have the potential to influence offspring phenotype and fitness. Indeed, parental hatchery rearing

has been associated with fitness-related phenotypic variation in F1 in Atlantic salmon (Wellband

et al., 2021), though F1 effects of parental hatchery rearing are complex and weaker than F0

effects in our study. It is possible that slight differences in offspring environment could

contribute to the complex patterns of inheritance we observed. The timing of juvenile release can

also influence the fitness consequences of hatchery rearing (Milot et al., 2013), therefore the

timing of F0 release may influence the extent of epigenetic inheritance due to hatchery rearing

and merits further study. Due to the nonadditive nature of these heritable effects, it may be

difficult to predict epigenetic responses to hatchery rearing beyond F0 and ensure effective

conservation actions, though we come to the encouraging conclusion that the epigenetic effects of

hatchery rearing are clearly diluted in F1 offspring.

Conclusions

We showed that fin DNA methylation changes associated with hatchery rearing persisted

in Atlantic salmon after release into natural systems and until spawning. The detection of two

cohorts of hatchery fish suggests that slightly modified supplementation techniques may be able

to reduce the epigenetic effects of hatchery rearing, though further research is needed to

determine what factors may decrease the extent of these differences. In particular, future studies

attempting to minimize the epigenetic effects of hatchery rearing will need to characterize the

methylome of several different tissues in order to determine the overall systemic implications of

manipulated environments on the methylome once the cost of whole genome methylation

sequencing declines. We also observed sex-specific effects of hatchery rearing in F0 which may

contribute to fitness differences between male and female hatchery-reared fish (Christie et al.,

2014). We showed that few of these effects persisted to F1, suggesting that the effects of hatchery

rearing may disappear rapidly over generations. However, the heritable effects of hatchery

rearing exhibit complex, nonadditive patterns dependent on both maternal and paternal origin

which may affect phenotype and will be difficult to account for in conservation and management

decisions. With increasing investment and research into improving hatchery rearing and

supplementation efforts, understanding and minimizing the detrimental effects of

supplementation on fish is critical to ensure effective conservation efforts.

Acknowledgements

We appreciate Dr Mark Christie's helpful input and discussion of our results. We thank

Gabriel Piette-Lauzière, Charles Babin, and Alexandre Carbonneau for assistance in the

laboratory and sample preparation. We also thank Subject Editor Dr. Luciano Beheregaray and

the anonymous reviewer for their help in improving the manuscript. This study was funded by the

Ministère des Forêts, de la Faune et des Parcs du Québec (MFFP) and by NSERC (RDCPJ

537349–18).

References

Akalin, A., Kormaksson, M., Li, S., Garrett-Bakelman, F. E., Figueroa, M. E., Melnick, A., &
Mason, C. E. (2012). MethylKit: a comprehensive R package for the analysis of genome-
wide DNA methylation profiles. *Genome Biology*, *13*(10). doi: 10.1186/gb-2012-13-10-R87

Aljanabi, S. M., & Martinez, I. (1997). Universal and rapid salt-extraction of high quality
genomic DNA for PCR-based techniques. *Nucleic Acids Research*, *25*(22), 4692–4693.

Anastasiadi, D., Piferrer, F., & Wittkopp, P. (2019). Epimutations in developmental genes
underlie the onset of domestication in farmed European sea bass. *Molecular Biology and
Evolution*, *36*(10), 2252–2264. doi: 10.1093/molbev/msz153

Anastasiadi, D., Venney, C. J., Bernatchez, L., & Wellenreuther, M. (2021). Epigenetic
inheritance and reproductive mode in plants and animals. *Trends in Ecology & Evolution*,
36(12), 1124–1140. doi: 10.1016/j.tree.2021.08.006

Arthington, A. H., Dulvy, N. K., Gladstone, W., & Winfield, I. J. (2016). Fish conservation in
freshwater and marine realms: status, threats and management. *Aquatic Conservation:
Marine and Freshwater Ecosystems*, *26*(5), 838–857. doi: 10.1002/aqc.2712

Berbel-Filho, W. M., Berry, N., Rodríguez-Barreto, D., Rodrigues Teixeira, S., Garcia de Leaniz,
C., & Consuegra, S. (2020). Environmental enrichment induces intergenerational
behavioural and epigenetic effects on fish. *Molecular Ecology*, *29*(12), 2288–2299. doi:
10.1111/mec.15481

Berbel-Filho, W. M., Rodríguez-Barreto, D., Berry, N., Garcia De Leaniz, C., & Consuegra, S.

(2019). Contrasting DNA methylation responses of inbred fish lines to different rearing
environments. *Epigenetics*, *14*(10), 939–948. doi: 10.1080/15592294.2019.1625674

Berntson, E. A., Carmichael, R. W., Flesher, M. W., Ward, E. J., & Moran, P. (2011).
Diminished reproductive success of steelhead from a hatchery supplementation program
(Little Sheep Creek, Imnaha Basin, Oregon). *Transactions of the American Fisheries
Society*, *140*(3), 685–698. doi: 10.1080/00028487.2011.584489

Bouchard, R., Wellband, K., Lecomte, L., Bernatchez, L., & April, J. (2022). Effects of stocking
at the parr stage on the reproductive fitness and genetic diversity of a wild population of
Atlantic salmon (*Salmo salar* L.). *Evolutionary Applications*, *15*, 838–852. doi:
10.1111/eva.13374

Bradbury, I. R., Wringe, B. F., Watson, B., Paterson, I., Horne, J., Beiko, R., … Bentzen, P.
(2018). Genotyping-by-sequencing of genome-wide microsatellite loci reveals fine-scale
harvest composition in a coastal Atlantic salmon fishery. *Evolutionary Applications*, *11*(6),
918–930. doi: 10.1111/eva.12606

Brown, C., & Day, R. L. (2002). The future of stock enhancements: lessons for hatchery practice
from conservation biology. *Fish and Fisheries*, *3*(2), 79–94. Retrieved from
https://www.wellbeingintlstudiesrepository.org/acwp_aff

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics*, *34*(17), i884–i890. doi: 10.1093/bioinformatics/bty560

Christensen, K. A., Luyer, J. Le, Chan, M. T. T., Rondeau, E. B., Koop, B. F., Bernatchez, L., &
Devlin, R. H. (2021). Assessing the effects of genotype-by-environment interaction on
epigenetic, transcriptomic, and phenotypic response in a Pacific salmon. *G3
Genes|Genomes|Genetics*, *0*(January). doi: 10.1093/g3journal/jkab021

Christie, M. R., Ford, M. J., & Blouin, M. S. (2014). On the reproductive success of early-
generation hatchery fish in the wild. *Evolutionary Applications*, *7*(8), 883–896. doi:
10.1111/eva.12183

Christie, M. R., Marine, M. L., Fox, S. E., French, R. A., & Blouin, M. S. (2016). A single
generation of domestication heritably alters the expression of hundreds of genes. *Nature
Communications*, *7*, 1–6. doi: 10.1038/ncomms10676

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., … 1000
Genomes Project Analysis Group. (2011). The variant call format and VCFtools.
Bioinformatics, *27*(15), 2156–2158. doi: 10.1093/bioinformatics/btr330

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., … Li, H. (2021).
Twelve years of SAMtools and BCFtools. *GigaScience*, *10*(2), 1–4. doi:
10.1093/gigascience/giab008

Fellous, A., Labed-Veydert, T., Locrel, M., Voisin, A. S., Earley, R. L., & Silvestre, F. (2018).
DNA methylation in adults and during development of the self-fertilizing mangrove rivulus,
Kryptolebias marmoratus. *Ecology and Evolution*, *8*(12), 6016–6033. doi:

10.1002/ece3.4141

Feng, H., & Wu, H. (2019). Differential methylation analysis for bisulfite sequencing using DSS.
Quantitative Biology, *7*(4), 327–334. doi: 10.1007/s40484-019-0183-8

Fraser, D. J. (2008). How well can captive breeding programs conserve biodiversity? A review of
salmonids. *Evolutionary Applications*, *1*(4), 535–586. doi: 10.1111/j.1752-
4571.2008.00036.x

Frisk, M., Høyland, M., Zhang, L., Vindas, M. A., Øverli, Ø., & Johansen, I. B. (2020). Intensive
smolt production is associated with deviating cardiac morphology in Atlantic salmon (Salmo
salar L.). *Aquaculture*, *529*, 735615. doi: 10.1016/j.aquaculture.2020.735615

Gavery, M. R., Nichols, K. M., Berejikian, B. A., Tatara, C. P., Goetz, G. W., Dickey, J. T., …
Swanson, P. (2019). Temporal dynamics of DNA methylation patterns in response to rearing
juvenile steelhead (*Oncorhynchus mykiss*) in a hatchery versus simulated stream
environment. *Genes*, *10*(5). doi: 10.3390/genes10050356

Gavery, M. R., Nichols, K. M., Goetz, G. W., Middleton, M. A., & Swanson, P. (2018).
Characterization of genetic and epigenetic variation in sperm and red blood cells from adult
hatchery and natural-origin steelhead, *Oncorhynchus mykiss*. *G3: Genes, Genomes,
Genetics*, *8*(11), 3723–3736. doi: 10.1534/g3.118.200458

Gimenez, M. D., Vazquez, D. V., Trepat, F., Cambiaso, V., & Rodríguez, G. R. (2021). Fruit
quality and DNA methylation are affected by parental order in reciprocal crosses of tomato.
Plant Cell Reports, *40*(1), 171–186. doi: 10.1007/s00299-020-02624-x

Harder, A. M., & Christie, M. R. (2022). Genomic signatures of adaptation to novel
environments: hatchery and life history-associated loci in landlocked and anadromous
Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Sciences*, *79*(5),
761–770. doi: 10.1139/cjfas-2021-0066

Jenkins, M. (2003). Prospects for Biodiversity. *Science*, *302*(5648), 1175–1177. doi:
10.1126/science.1088666

Kalinowski, S. T., Taper, M. L., & Marshall, T. C. (2007). Revising how the computer program
CERVUS accommodates genotyping error increases success in paternity assignment.
Molecular Ecology, *16*(5), 1099–1106. doi: 10.1111/j.1365-294X.2007.03089.x

King, R. A., & Stevens, J. R. (2020). An improved genetic sex test for Atlantic salmon (*Salmo
salar* L.). *Conservation Genetics Resources*, *12*(2), 191–193. doi: 10.1007/s12686-019-
01094-y

Klopfenstein, D. V., Zhang, L., Pedersen, B. S., Ramírez, F., Vesztrocy, A. W., Naldi, A., …
Tang, H. (2018). GOATOOLS: A Python library for gene ontology analyses. *Scientific
Reports*, *8*, 10872. doi: 10.1038/s41598-018-28948-z

Koch, I. J., Nuetzel, H. M., & Narum, S. R. (2022). Epigenetic effects associated with salmonid
supplementation and domestication. *Environmental Biology of Fishes*, (0123456789). doi:
10.1007/s10641-022-01278-w

Laing, L. V., Viana, J., Dempster, E., Uren Webster, T. M., van Aerle, R., Mill, J., & Santos, E. M. (2018). Sex-specific transcription and DNA methylation profiles of reproductive and epigenetic associated genes in the gonads and livers of breeding zebrafish. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, *222*, 16–25.

Lavoie, C., Courcelle, M., Redivo, B., & Derome, N. (2018). Structural and compositional mismatch between captive and wild Atlantic salmon (*Salmo salar*) parrs' gut microbiota highlights the relevance of integrating molecular ecology for management and conservation methods. *Evolutionary Applications*, *11*(9), 1671–1685. doi: 10.1111/eva.12658

Le Luyer, J., Laporte, M., Beacham, T. D., Kaukinen, K. H., Withler, R. E., Leong, J. S., … Bernatchez, L. (2017). Parallel epigenetic modifications induced by hatchery rearing in a Pacific salmon. *Proceedings of the National Academy of Sciences*, *114*(49), 12964–12969. doi: 10.1101/148577

Leitwein, M., Laporte, M., Le Luyer, J., Mohns, K., Normandeau, E., Withler, R., & Bernatchez, L. (2021). Epigenomic modifications induced by hatchery rearing persist in germ line cells of adult salmon after their oceanic migration. *Evolutionary Applications*, *14*(10), 2402–2413. doi: 10.1111/eva.13235

Leitwein, M., Wellband, K., Cayuela, H., Luyer, J. Le, Mohns, K., Withler, R., & Bernatchez, L. (2022). Strong parallel differential gene expression induced by hatchery rearing weakly associated with methylation signals in adult Coho salmon (*O. kisutch*). *Genome Biology*, *14*(4), 1–14.

Leonard, J. B. K., & McCormick, S. D. (2001). Metabolic enzyme activity during smolting in stream- and hatchery-reared Atlantic salmon (*Salmo salar*). *Canadian Journal of Fisheries and Aquatic Sciences*, *58*(8), 1585–1593. doi: 10.1139/cjfas-58-8-1585

Liu, Z. Y., Wang, X. L., Ou, S. Q., Hou, D. X., & He, J. H. (2020). Sanguinarine modulate gut microbiome and intestinal morphology to enhance growth performance in broilers. *PLoS ONE*, *15*(6), 1–17. doi: 10.1371/journal.pone.0234920

Milot, E., Perrier, C., Papillon, L., Dodson, J. J., & Bernatchez, L. (2013). Reduced fitness of Atlantic salmon released in the wild after one generation of captive breeding. *Evolutionary Applications*, *6*(3), 472–485. doi: 10.1111/eva.12028

O'Sullivan, R. J., Aykanat, T., Johnston, S. E., Rogan, G., Poole, R., Prodöhl, P. A., … Reed, T. E. (2020). Captive-bred Atlantic salmon released into the wild have fewer offspring than wild-bred fish and decrease population productivity. *Proceedings of the Royal Society B: Biological Sciences*, *287*(1937), 20201671. doi: 10.1098/rspb.2020.1671rspb20201671

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., … Wagner, H. (2020). *vegan: Community ecology package. R package version 2.5-7* (p. https://CRAN.R-project.org/package=vegan). p. https://CRAN.R-project.org/package=vegan.

Pedersen, L. F., Koed, A., & Malte, H. (2008). Swimming performance of wild and F1-hatchery-reared Atlantic salmon (*Salmo salar*) and brown trout (*Salmo trutta*) smolts. *Ecology of Freshwater Fish*, *17*(3), 425–431. doi: 10.1111/j.1600-0633.2008.00293.x

Podgorniak, T., Brockmann, S., Konstantinidis, I., & Fernandes, J. M. O. (2019). Differences in
the fast muscle methylome provide insight into sex-specific epigenetic regulation of growth
in Nile tilapia during early stages of domestication. *Epigenetics*, *14*(8), 818–836. doi:
10.1080/15592294.2019.1618164

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics*, *26*(6), 841–842. doi: 10.1093/bioinformatics/btq033

R Core Team. (2022). *R: A language and environment for statistical computing*. Vienna, Austria:
R Foundation for Statistical Computing. Retrieved from https://www.r-project.org

Raza, M. A., Yu, N., Wang, D., Cao, L., Gan, S., & Chen, L. (2017). Differential DNA
methylation and gene expression in reciprocal hybrids between *Solanum lycopersicum* and
S. pimpinellifolium. *DNA Research*, *24*(6), 597–607. doi: 10.1093/dnares/dsx028

Rodriguez Barreto, D., Garcia De Leaniz, C., Verspoor, E., Sobolewska, H., Coulson, M.,
Consuegra, S., & Mulligan, C. (2019). DNA methylation changes in the sperm of captive-
reared fish: a route to epigenetic introgression in wild populations. *Molecular Biology and
Evolution*, *36*(10), 2205–2211. doi: 10.1093/molbev/msz135

Shen, H., He, H., Li, J., Chen, W., Wang, X., Guo, L., … Deng, X. W. (2012). Genome-wide
analysis of DNA methylation and gene expression changes in two *Arabidopsis* ecotypes and
their reciprocal hybrids. *Plant Cell*, *24*(3), 875–892. doi: 10.1105/tpc.111.094870

Sutherland, B. J. G., Prokkola, J. M., Audet, C., & Bernatchez, L. (2019). Sex-specific co-
expression networks and sex-biased gene expression in the salmonid Brook Charr Salvelinus
fontinalis. *G3: Genes, Genomes, Genetics*, *9*(3), 955–968. doi: 10.1534/g3.118.200910

Thériault, V., Moyer, G. R., Jackson, L. S., Blouin, M. S., & Banks, M. A. (2011). Reduced
reproductive success of hatchery coho salmon in the wild: insights into most likely
mechanisms. *Molecular Ecology*, *20*(9), 1860–1869. doi: 10.1111/j.1365-
294X.2011.05058.x

Venney, C. J., Johansson, M. L., & Heath, D. D. (2016). Inbreeding effects on gene-specific
DNA methylation among tissues of Chinook salmon. *Molecular Ecology*, *25*(18), 4521–
4533. doi: 10.1111/mec.13777

Venney, C. J., Wellband, K. W., & Heath, D. D. (2021). Rearing environment affects the genetic
architecture and plasticity of DNA methylation in Chinook salmon. *Heredity*, *126*(1), 38–49.
doi: 10.1038/s41437-020-0346-4

Wellband, K., Roth, D., Linnansaari, T., Curry, R. A., & Bernatchez, L. (2021). Environment-
driven reprogramming of gamete DNA methylation occurs during maturation and is
transmitted intergenerationally in Atlantic salmon. *G3: Genes, Genomes, Genetics*, *11*(12),
jkab353. doi: 10.1093/g3journal/jkab353

Williamson, K. S., Murdoch, A. R., Pearsons, T. N., Ward, E. J., & Ford, M. J. (2010). Factors
influencing the relative fitness of hatchery and wild spring Chinook salmon (*Oncorhynchus
tshawytscha*) in the Wenatchee River, Washington, USA. *Canadian Journal of Fisheries*

and Aquatic Sciences, *67*(11), 1840–1851. doi: 10.1139/F10-099

Willoughby, J. R., & Christie, M. R. (2019). Long-term demographic and genetic effects of
releasing captive-born individuals into the wild. *Conservation Biology*, *33*(2), 377–388. doi:
10.1111/cobi.13217.This

Zhan, L., Paterson, I. G., Fraser, B. A., Watson, B., Bradbury, I. R., Nadukkalam Ravindran, P.,
… Bentzen, P. (2017). Megasat: automated inference of microsatellite genotypes from
sequence data. *Molecular Ecology Resources*, *17*(2), 247–256. doi: 10.1111/1755-
0998.12561

Data Accessibility and Benefit-Sharing

682   Raw whole genome methylation data and metadata are deposited in the SRA (BioProject

683   PRJNA892473) and will be released publicly upon acceptance. Benefit-sharing statements are

684   not applicable.

Author Contributions

CJV, LB, JA, GC, and RB conceived of the project. CJV, RB, and LL performed the

analyses. CJV wrote the initial manuscript and all authors contributed to the final version.

Funding was provided by the Ministère des Forêts, de la Faune et des Parcs du Québec (MFFP)

and an NSERC grant to LB (RDCPJ 537349 – 18).

Figures

Figure 1: PCA plots for methylation data showing (A) differentiation between F0 hatchery
cohorts and wild fish, and (B) lack of whole genome F1 differentiation based on parental rearing
environment.

Figure 2: F0 differential methylation results between hatchery cohorts ("normal" = orange, "wildtype" = brown) and wild salmon (green). We detected (A) 2 007 DMRs between wild and "normal" hatchery salmon, (B) 1 285 DMRs between "normal" and "wildtype" hatchery cohorts, and (C) only 156 DMRs between the "wildtype" hatchery cohort and wild salmon. Hierarchical clustering was performed based on Euclidean distances. Percent methylation is denoted by the yellow to indigo scale for each DMR (0-100%).

702 **Figure 3:** F0 sex specific DMRs due to rearing environment. We identified (A) 5 717 DMRs
703 between male hatchery-reared and wild salmon, and (B) 2 439 DMRS between female salmon.

Figure 4: F1 DMR analysis identified (A) 284 DMRs due to maternal source, (B) 254 DMRs due to paternal source, and (C) 271 DMRs due to their interaction.