Comprehensive monitoring of tissue composition using in vivo imaging of cell nuclei and deep learning

Amrita Das Gupta[a], Livia Asan[a,d], Jennifer John [a], Carlo A. Beretta [a], Thomas Kuner [a], Johannes Knabbe [a,b]

a Department of Functional Neuroanatomy, Institute for Anatomy and Cell Biology, Heidelberg University, Germany.

b Department of General Psychiatry, Centre for Psychosocial Medicine, Heidelberg University, Germany.

d Current address: Department of Neurology and Center for Translational Neuro- and Behavioral Sciences (C-TNBS), University Hospital Essen, Germany.

Correspondence: johannes.knabbe@uni-heidelberg.de, thomas.kuner@uni-heidelberg.de

Abstract

Comprehensive analysis of tissue composition has so far been limited to ex vivo approaches. Here,

we introduce NuClear (Nucleus-instructed tissue composition using deep learning), an approach

combining in vivo two-photon imaging of histone 2B-eGFP-labeled cell nuclei with subsequent

deep learning-based identification of cell types from structural features of the respective cell

nuclei. This allowed us to classify all cells per imaging volume (0.25 mm^3 containing ~25000 cells)

and identify their position in 3D space in a non-invasive manner using only a single label. NuClear

opens a window to study changes in relative abundance and location of all major brain cell types

in individual mice over extended time periods, enabling comprehensive studies of changes in

cellular composition in physiological and pathophysiological conditions.

121 words

Introduction

Understanding the plasticity and interaction of different brain cell types in vivo has been a limiting

factor for the investigation of large-scale structural brain alterations associated with diverse

physiological and pathophysiological states. Until now, to investigate the effect of an experimental

intervention on all brain cell types and their respective spatial interaction, each cell type would

require an individual labelling strategy using a set of specific promoters and fluorophores. This

would require prioritizing cell types from the outset and thereby limit the scope to a small subset

of the population. With multiple cell types to consider, the number of experiments and animals

investigated may become intractable. So far, studies quantitatively assessing whole tissue

composition have analyzed tissue ex vivo using manual or automated analysis of antibody staining

in tissue slices (Ero, Gewaltig et al. 2018) or after isotropic fractionation (Herculano-Houzel and

Lent 2005), in situ hybridization (Nagendran, Riordan et al. 2018) or single cell sequencing (Zeisel,

Hochgerner et al. 2018). Other studies were using manual (Schmitz and Hof 2007) or automated

stereological approaches (Zhang, Yan et al. 2017, Pakkenberg, Olesen et al. 2019). All these

studies were conducted ex vivo and, in part, require the dissolution of the studied organ's cellular

architecture. The ability to identify all brain cell types and their location in 3D space, here referred

to as 'composition', in the integrity of the living brain to quantify and observe changes over time

is enticing. Here, we propose an experimental approach to study neurons and glial cells,

especially, astroglia, microglia and oligodendroglia, as well as endothelial cells in the same mouse

in vivo, using only one genetically modified mouse line expressing a fusion protein of Histon 2B

and enhanced green fluorescent protein (eGFP) in all cell nuclei. Our approach uses a deep

learning method, implementing artificial neural networks to classify each nucleus belonging to a

specific cell type, making it possible to perform a <u>nu</u>cleus-instructed tissue <u>c</u>omposition analysis

using deep <u>lear</u>ning (NuClear). In addition, knowing the 3D location of every cell opens new

opportunities for further analyses. Apart from the application in longitudinal in vivo imaging of

the mouse brain, the concept of this technique will be applicable to many other cellular imaging

techniques and in different experimental systems, making it a powerful approach for future

analyses of large-scale automated tissue composition.

384 words

Results

For the comprehensive identification and tracking of cell type-composition in the mouse brain

over the course of weeks and even months, a mouse line constitutively expressing a human

Histone 2B-eGFP (H2GFP) fusion protein in all cell nuclei was imaged using 2-photon microscopy

after implantation of a chronic cranial window (Figure 1A, Supplementary Figure 1A, B).

Volumetric images taken from this mouse line resemble the well-known DAPI staining of cell

nuclei (Supplementary Figure 1C). For automated cell type classification of the imaged nuclei, a

neuronal network was trained on images of GFP-labeled nuclei from reporter mouse lines, where

known cell types were additionally labelled with a red fluorescent protein (tdTomato or mCherry),

thus creating a ground truth dataset for supervised training for each cell type (Figure 1B, lower

panel). The trained neuronal network was used to classify cell types based on their GFP-labeled

nuclei (Figure 1B, upper panel). In our study we selected five different cell types for classification

(Figure 2A): Neurons (yellow), astroglia (blue), microglia (green), oligodendroglia (turquoise), and

endothelial cells (red). To generate ground-truth datasets without a bleed through of the strong

tdTomato fluorescence towards the GFP-channel (Supplementary Figure 1D), we chose a method

of inducing the expression of tdTomato for unequivocal cell type identification *after* imaging of

the nuclei. We selected a labelling strategy consisting of a tamoxifen-inducible cell type-specific

Cre-recombinase in conjunction with a floxed-tdTomato mouse line for endothelial cells (Tie2-

CreErt2), astroglia (GFAP-CreErt2) and oligodendroglia (PLP-CreErt2) (Figure 2B). We acquired

volumetric images of nuclei from these mouse lines and induced tdTomato expression via

intraperitoneal injection of tamoxifen (Supplementary Figure 1E). After 3 to 4 days the same

location was imaged again, and the previously imaged nuclei were selected by overlay with the

red fluorescence signal (Figure 2C). As microglia changed their positions after induction with

tamoxifen, we chose acute injection of tomato-lectin coupled to DyLight 594 to label these cells,

which did not produce any signal alterations in the eGFP-channel (Supplementary Figure 1F).

Neurons were labelled using an AAV-mediated expression of the mCherry red fluorescent protein

under the neuron-specific synapsin promotor (syn1-mCherry), which did not affect the eGFP-

signal due to its specific fluorescence properties (Supplementary Figure 1F). Hence, this approach

was used to ascertain images of the labeled nuclei to be undisturbed by signals possibly arising

from the red fluorescence used for unequivocal identification of the cell type.

Automated segmentation of nuclei in the raw data was achieved using the Stardist neuronal

network (Schmidt, Weigert et al. 2018), trained on manually labeled ground truth datasets of the

H2GFP mouse line which achieved a high detection accuracy of 94 % as well as a good shape

segmentation (Supplementary Figure 1G, H). From the binary mask of individual nuclei and their

respective pixel intensities in the raw image, in total 121 features were extracted, including 3D-

diameter, flatness, grey level non-uniformity, entropy, first order minimum or coarseness (Figure

2D) (for a full list see Supplementary Table 1). To reduce overfitting of the classification algorithm,

a subset of 12 features representing the whole feature space was chosen by a feature selection

algorithm (Supplementary Table 2). A polar plot with these nuclear features visualizes the

differences between the cell types, for example neuronal nuclei having a larger 3D diameter and

minor axis length in comparison to microglial and astroglial nuclei that exhibit a larger pixel

entropy (Figure 2E). For certain features, significant differences between the cell types can be

shown as well, for example flatness being significantly higher in nuclei of microglia than in nuclei

of neurons (Figure 2F). Furthermore, combinations of two or three distinctive features allow for

the visual separation of the cell types in the 2D and 3D space, respectively (Supplementary Figure

2). Having demonstrated the ability of nuclear features to distinguish between their

corresponding cell types, we created a neuronal network for cell type classification based on

nucleus features (Supplementary Figure 3). Using data from the five reporter mouse lines, we

trained one classifier per cell type for distinguishing one cell type from the respective other cell

types (Figure 3A). To increase the amount of training data and equalize the nucleus counts for

each cell type to reduce training bias, synthetic data was generated from the features of the

original dataset. Synthetic data distribution fit well to the distribution of the original data for each

cell type (Supplementary Figure 4A, B). Training and validation were performed on a dataset

consisting of original and synthetic data (Figure 3A). After classifying the whole dataset, nuclei

were either assigned to a single class, two or more classes (undecided) or to none (unclassified).

Precision and recall rates for each classifier were high for neurons and endothelial cells. Due to

their relative similarity, glial cells exhibited lower precision and recall rates (Figure 3B). After

combining all classifier output, the classification accuracy for the entire dataset was highest for

neurons (98 %) and endothelial cells (99 %), whereas the classifier showed lower accuracies for

all glial cell types, especially for microglia (96 %) (Figure 3C, D, for the respective confusion plots

see Supplementary Figure 5). A model classifying only neurons, glia (as one common class

irrespective of glial subtype) and endothelial cells showed even higher accuracy, precision and

recall rates for glial cells (data not shown). When applying the classification model to data of

cortical layer 2/3 of the supplementary motor area (SMA) of H2GFP mice, two features not chosen

for classification by the automated feature selection algorithm differed significantly between the

classes: neuronal nucleus volume (Figure 3E) and endothelial mean surface to volume ratio, a

measure for the sphericity of the nucleus (Figure 3F). Both findings are expected for each of the

cell types and hence demonstrate that also features that were not used for classification can be

readily detected. The 3D classification results for one example volume containing 20123 cells are

shown in Supplementary Figure 6 and intuitively visualize the relative abundance and positioning

of cell types within the volume and their mutual spatial relationship (see also Supplementary

Movie 1 for an animated version of the segmentation and classification process). The absolute

nuclei count from the region shows a density of 44000 neurons/mm³ and of 57000 glial cells

(including unclassified cells), resulting in a Glia/Neuron-Ratio of 1.3 (Figure 3G). The number of

cells in a subset of Layer2/3 over 12 weeks remained unchanged after correcting for multiple

comparisons (Figure 3H). When normalizing these counts to baseline, only the number of

astrocytes shows a trend to increase (Figure 3I), which could be either due to astrocytosis caused

by aging or caused by continued reaction to the chronic window implantation. Interestingly,

astrocytic nuclear volume significantly decreases over time (Figure 3J), which might be due to

altered transcriptional activity known to occur in astrocytes with aging (Boisvert, Erikson et al.

2018, Clarke, Liddelow et al. 2018). This latter result illustrates that our method can extract even

subtle and unexpected changes from of a vast parameter space of different aspects of tissue

composition.

1157 words

Discussion

Here, we present a novel method for intravital, longitudinal, comprehensive, large scale and 3D

tissue composition based on a deep learning algorithm trained on radiomics features of cell nuclei

(NuClear -- Nucleus-instructed tissue composition using deep learning). We demonstrate that cell

types can be distinguished via the respective properties of their cell nucleus. This has been mostly

anecdotal anatomical textbook knowledge for a good century, yet it has rarely been quantified

and compared in a systematic manner. Moreover, we demonstrate how this rejuvenated

anatomical knowledge can be converted into a powerful new approach with the help of today's

state-of-the-art imaging and deep learning techniques.

For a demonstration of the usability of NuClear, we analyzed volumetric images of the cortex of

H2GFP mice acquired with in vivo two photon microscopy. To be able to image a large dataset in

a comparatively short time, we chose a feasible imaging resolution with a z-step size of 2 μm and

a lateral resolution of 0.3 μm. We were able to image a whole 3D volume of 700 μm³ consisting

of more than 20000 cells in around 20 minutes, making it possible to perform large scale data

acquisition in a repeated manner, enabling longitudinal intravital monitoring of tissue histology

over time and in response to perturbations, treatments, or adaptive processes.

Neurons and endothelial cells showed the best classification results regarding total classification

accuracy and precision as well as recall rates, as their nucleus feature profile turned out to be

markedly different between each other as well as between the individual glial cell types (Figure

2E, 3B). Differentiation between the three different glial cell types was more difficult to achieve,

resulting in lower overall accuracy rates as well as lower precision and recall rates, due their

respective nucleus feature profiles being more similar. Nevertheless, with an accuracy above 90%

our approach can reliably be used to subclassify glia. We expect that a higher imaging resolution

as well as more training data would increase the classifier performance even more. Another

reason for the comparatively lower classification accuracy in microglia might be that we used

acute injection of tomato-lectin to label these cells, which could have introduced a bias towards

activated microglia. Nevertheless, the overall accuracy of the microglia classifier for the whole

training dataset was around 96%, owing to the fact, that non-microglial nuclei could be identified

correctly most of the time. For astrocytes, we used GFAP-labelled cells for classification, a label

that primarily marks reactive astrocytes and is expressed increasingly during aging. Future studies

could optimize this labelling strategy. Further support for the accuracy of the classifier can be

concluded from analyzing features of nuclei that were not used during training such as mean

nuclear volume and surface to volume ratio. These features showed distinct clustering and low

standard deviation in each class as well as significant differences between the cell types.

When applying the classification model to a Layer 2/3 volume of the supplementary motor area

(SMA) and extrapolating the number of cells per mm³, a marked similarity of the number of

neurons with published results is evident (43822 ± 3747 vs 45374±4894) (Ero, Gewaltig et al.

2018) (Figure 3G). Astrocyte numbers are similar (18882 ± 3848 vs 13258± 1416), microglia

numbers differ more (12573 ± 2840 vs 24584.6± 2687). Oligodendrocyte numbers seem to be

vastly different (4358 ± 586 vs 51436.1± 488), which might stem from the fact that only Layer 2/3

was analyzed, excluding deeper cortical layers that have been shown to contain the majority of

oligodendrocytes (Tan, Kalloniatis et al. 2009). When all glial cells as well as undecided and

unclassified cells are summed up, a Glia/Neuron ratio of 1.3 can be calculated, a result that is in

line with known results of the rodent cortex (Herculano-Houzel 2014). Candidates for unclassified

cells could be the relatively large population of oligodendrocyte precursor cells (about 5% of all

cells in the brain (Dawson, Polito et al. 2003)), which might explain that unclassified nuclei have

a similar volume as oligodendrocytes (Figure 3E). In conclusion, cell type counts are comparable

to published results.

A decisive advantage of NuClear over existing ex vivo methods is the ability to study cell type

changes over time in the same imaging volume and thus achieving a higher statistical power in

experiments with fewer animals, which is crucial for complying with the "3R" rules in animal

research (reduce, replace, refine). In our lab, we were able to perform imaging in selected

locations up to 1 year after the implantation of chronic cranial windows. When analyzing the

effect of time on the imaging volumes from the supplementary motor area (SMA), we were able

to detect a trend towards astrocytosis after 12 weeks, which might be due to the aging process

leading to a higher GFAP-reactivity (Palmer and Ousman 2018). The significant decrease in mean

nucleus volume of astroglia over time could be attributed to altered transcriptional activity since

astrocyte reactivity changes in the cortex over time (Clarke, Liddelow et al. 2018). Another factor

explaining the astrocytosis might be the constant reaction to the chronic cranial window (Guo,

Zou et al. 2017).

Our study illustrates only one possible application of NuClear and subsequent analysis of 3D tissue

composition. Here we focused on the identification of cell types, but the 3D datasets generated

allow for additional analyses such as statistical distribution of cell types with regard to each other,

their nearest neighbor distances or inferences on physical tissue volume (Asan, Falfan-Melgoza et

al. 2021). Our classifiers mostly rely on geometrical parameters such as nucleus shape, but also

considers texture information to a certain degree. An increased sampling resolution may further

increase textural information and thereby yield even higher classification accuracies. NuClear will

be easily applicable to different organs, other organisms, and in vitro preparations such as cell

cultures or organoids, provided that the respective ground truths are available. It will be usable

with a variety of microscopy techniques such as confocal-, light-sheet- or three-photon

microscopy, making large-scale tissue analysis much more accessible. We anticipate that any cell

type will be detected by adding more classifiers trained with appropriate ground truth data. In

this way, we propose an easy and readily usable method to implement large-scale tissue analysis

ex vivo or in vivo to study effects of interventions on cell type compositions of different organs.

1050 words

Materials and Methods

<u>Ethical approval</u>

The entire study was conducted in line with the European Communities Council Directive

(86/609/ECC) to reduce animal pain and/or discomfort. All experiments followed the German

animal welfare guidelines specified in the TierSchG. The study has been approved by the local

animal care and use council (Regierungspräsidium Karlsruhe, Baden-Wuerttemberg) under the

reference numbers G294/15 and G27/20. All experiments followed the 3R principle and complied

with the ARRIVE guidelines (Percie du Sert, Hurst et al. 2020).

<u>Animals</u>

Adult transgenic mice expressing the human histone 2B protein (HIST1H2B) fused with enhanced

green fluorescence protein (eGFP) under the control of the chicken beta actin promoter (CAG-

H2B-eGFP) (Hadjantonakis and Papaioannou 2004) (B6.Cg-Tg(HIST1H2BB/EGFP)1Pa/J, Jackson

Laboratory; # 006069) were used in all animal experiments. For the reporter mouse lines, H2B-

eGFP mice were bred to carry a floxed sequence of the red fluorescent protein tdTomato and a

tamoxifen inducible Cre-recombinase under the expression of different cell type specific

promoters: GFAP   for   astroglia   (HIST1H2BB/EGFP-GFAP/ERT2CRE-

CAG/loxP/STOP/loxP/tdTomato), PLP for oligodendroglia (HIST1H2BB/EGFP-PLP/ERT2CRE-

CAG/loxP/STOP/loxP/tdTomato) and Tie2 for endothelial cells (HIST1H2BB/EGFP-Tie2/ERT2CRE-

CAG/loxP/STOP/loxP/tdTomato). Expression of the Cre-recombinase was achieved over a course

of up to 5 days with daily intraperitoneal injection of two doses of 1mg Tamoxifen (Sigma Life

sciences; dissolved in 1 part ethanol (99.8% absolute) and 10 parts sunflower oil) (supp. Figure 1

D). Microglia were visualized using intracortical injection of Lycopersicon Esculentum (Tomato)

lectin coupled with DyLight 594 during the cranial window surgery after dilution by 1:39 in

150mM NaCl, 2.5 mM KCl, 10mM HEPES at pH 7.4 (Schwendele, Brawek et al. 2012)

(supplementary Figure 1E). For neuronal labelling, an in house produced adeno-associated virus

expressing mCherry under the Synapsin promotor was cortically injected during the cranial

window surgery.

Chronic cranial window implantation

A craniectomy procedure was performed on each mouse to enable in vivo two-photon imaging

following a predefined protocol as described before (Holtmaat, Bonhoeffer et al. 2009)

(Supplementary Figure 1A). Briefly, the mice were anesthetized with an intraperitoneal injection

(i.p.) of a combination of 60 µl Medetomidine (1mg/ml), 160 µl Midazolam (5 mg/ml) and 40 µl

Fentanyl (0.05 mg/ml) at a dosage of 3 µl/g body weight. The head was shaved, and the mice

were fixed with ear bars in the stereotactic apparatus and eye ointment was applied. Xylocain®

1 % (100 µl, Lidocaine hydrochloride) was applied under the cranial skin and 250 µl Carprofen (0.5

mg/ml) was injected subcutaneously (s.c.). The skin was removed to expose the skull and the

surface of the skull was made rough to allow the cement to adhere better. A skull island (approx.

6mm ∅) was drilled centered at 1 mm rostral to bregma using a dental drill and removed with a

forceps (#2 FST by Dumont) making sure not to damage the cortical surface. For improved imaging

condition, the dura was carefully removed from both hemispheres using a fine forceps (#5 FST by

Dumont). Normal rat ringer solution (NRR) was applied on the exposed brain surface to keep it

moist, and a curved cover glass was placed on top to cover it (Asan, Falfan-Melgoza et al. 2021).

With dental acrylic cement (powder: Paladur, Kulzer; activator: Cyano FAST, Hager & Werken

GmbH) the cover glass was sealed, and excess cement was used to cover the exposed skull and

edges of the skin. A custom designed 3-D printed holder was placed on top of the window and

any gaps were filled with cement to ensure maximum adhesion with the skull. After the

procedure, the mice were injected (i.p. / s.c.) with a mix of 30 µl Atipamezole (5 mg/ml), 30 µl

Flumazenil (0.1 mg/m) and 180 µl Naloxon (0.4 mg/ml) at a dosage of 6 µl per gram body weight.

To ensure proper recovery of the mice, 3 more doses of Carprofen were given every 8-12 hours

and the mice were placed onto a heating plate and monitored.

In vivo two photon imaging

Two photon imaging was carried out with a two-photon microscope (TriM Scope II, LaVision

BioTec GmbH) with a pulsed Titanium-Sapphire (Ti:Sa) laser (Chameleon Ultra 2; Coherent) at

excitation wavelengths of 860nm for DyLight 594 and 960nm for H2GFP, tdTomato and mCherry

respectively. A water immersion objective (16x; NA 0.8, Nikon) was used to obtain volumetric 3D

stacks. Individual frames consisted of 694 μm x 694 μm in XY with a resolution of 0.29 μm/pixel.

Stacks were obtained at varying depths up to 702 μm from the cortical surface with a step size of

2 μm in Z. Prior to each imaging session, the laser power was adjusted to achieve the best signal

to noise ratio. Adaptations were made to minimize the effect of laser attenuation due to tissue

depth by creating a z-profile and setting the laser power for different imaging depths while

making sure to minimize oversaturation of pixels.

Mice were initially anesthetized using a vaporizer with 6% isoflurane, eye ointment was applied.

Mice were fixed in a custom-built holder on the microscope stage. Isoflurane was reduced to 0.5

– 1.5% depending on the breathing rate of each mouse to achieve a stable breathing rate of 55 –

65 breaths per minute with an oxygen flow rate of 0.9 – 1.1 l/min. A heating pad was placed

underneath the mouse to regulate body temperature. An infrared camera was used for

monitoring of the mice during the imaging.

Reporter mouse lines for neurons, astroglia, oligodendroglia and endothelial cells were imaged

3-4 weeks after the cranial window surgery. Microglia reporter mice were imaged immediately

after the cranial window surgery for a minimal inflammatory reaction (for a full list of reporter

mice see supplementary Figure 4E). For imaging adult H2GFP mice, 3D volumetric stacks were

imaged in the supplementary motor area (SMA) at two different timepoints. This includes

baseline imaging which was performed at 3-4 weeks after the chronic cranial window surgery and

in week 12 after the baseline. To investigate a possible age effect, mice underwent a sham surgery

at the left hindpaw 1 week after baseline surgery as part of a different study.

Automatic nuclei segmentation using StarDist

Automated nuclei segmentation was achieved using the StarDist algorithm, which implements a

machine learning method using a neuronal network to detect star-convex polygons in 2D and 3D

fluorescence imaging (Schmidt, Weigert et al. 2018). The StarDist model was trained to segment

nuclei in 3D using an in-house developed Jupyter Notebook. Manually segmented and labelled

nuclei from 15 different crops from the two-photon volumetric data with approximately 150 –

200 nuclei each were used for training the StarDist 3D segmentation classifier (supplementary

Figure 1 H).

Detection accuracy and precision were calculated by counting the true positive nuclei (TP), false

positive nuclei (FP) and false negative nuclei (FN). 4 arbitrary volumetric images were selected

containing up to 80 nuclei per crop. Accuracy was calculated as TP / (TP + FP + FN). Precision was

calculated as TP / (TP + FP).

Feature extraction using PyRadiomics

Using the PyRadiomics python package (van Griethuysen, Fedorov et al. 2017), in total 107

radiomics features were extracted for each segmented nucleus including, but not limited, to

morphological features, intensity features and texture features (see supplementary Table 1).

Nuclei touching the borders of the imaging volume were excluded.

<u>Extraction of data for training of the classifier</u>

For supervised training of the deep neural network for cell type classification, a ground truth

dataset was created using the two-photon volumetric fluorescence data and automatically

segmented nuclei. Using the red fluorescence channel, nuclei belonging to a specific cell type

were manually identified by overlaying the green channel data. This allowed for manual

identification of the label ID from the StarDist output by synchronizing the two imaging volumes

in ImageJ/Fiji and extraction of features from PyRadiomics for these individual nuclei.

Approximately 70-150 nuclei were identified for each cell type (supp. Figure 5).

To increase the number of nuclei for training and make it equal in size for all cell types to avoid a

bias in training, synthetic datasets were created from nuclei features using the synthetic data

vault (SDV) package (Patki, Wedge et al. 2016), which utilizes correlations between features of

the original dataset as well as mean, minimum or maximum values and standard deviations. The

synthetic data generated by the model fit to the original dataset with approximately 83% accuracy

(supplementary Figure 5).

<u>Training of the classification model</u>

All training and validation of the classifiers were performed with a custom MATLAB (R2021a,

Mathworks) script, manual validation was performed in ImageJ/Fiji. Original and synthesized data

were merged into a single table and random indices were generated to divide the dataset into a

training, test, and validation dataset with 70%, 15% and 15% of the total dataset, respectively.

The same training, test, and validation datasets were used for training five different classifiers,

one for each cell type. In the training dataset, for each classifier, the class of interest was assigned

a unique identifier and all other cell classes were denoted by another identifier, for example,

when training the neuronal classifier, the neurons were labelled as "Neuron" while all the other

cell types in the dataset were labelled "Other" (supplementary Figure 3). Each classification

network had the same design. Initially, a sequential feature selection algorithm "sequentialfs"

was applied to the whole dataset to extract 12 features with the highest variability to reduce data

overfitting. These were fed into the "feature input layer" of the classifier. The features were z-

score normalized for better compatibility between each class. A "fullyconnected" layer of 50

outputs was created and a mini-batch size of 16 samples was set to reduce the time and memory

consumption during training. A batch normalization layer was added to reduce sensitivity to the

initialization of the network. The activation layer ReLU (Rectified Linear Unit) was added for

thresholding operations. Another "fullyconnected" layer was added, which sets the number of

outputs identical to the number of input classes and a second activation layer "softmax" was

chosen for classifying the output into the two separate groups. The "adam" adaptive moment

estimation optimizer was selected as the solver for the training network which updates the

network weights based on the training data in an iterative manner. The training data and the

validation data were shuffled every epoch before training and network validation, respectively.

After each classifier was trained, further automatic and visual validation was performed to check

for accuracy.

Classification of the H2GFP data

After imaging of the volumetric volume of H2GFP nuclei, automated segmentation of all nuclei

was performed with the StarDist neuronal network (see above, supplementary Figure 6).

Afterwards, features for every segmented nucleus were extracted using PyRadiomics (see

above). Every segmented nucleus was then classified with all the five different classifiers. Thus, a

single nucleus would be either labelled as belonging to one of the five classes or to the "Other"

class. Nuclei that were assigned to multiple classes, were labelled as "Undecided" (UD). Any

nuclei that were identified as "Other" by all the classifiers were labelled as "Unclassified" (UC).

<u>Statistical analysis</u>

All classified nuclei and their features were stored in a local MySQL database (MySQL Workbench

8.0). The data were imported into R (R. Core Team 2021) for statistical analysis. Cell count, mean

nuclei volume and first nearest neighbor distance were calculated for each cell type for different

timepoints. The nearest neighbor distance was calculated using the "spatstat" package for R

(Baddeley and Turner 2005). Depending on the distribution of the data either t-tests or non-

parametric statistical test (Wilcoxon signed-rank test) were used for statistical testing, p-values

were corrected for multiple comparisons using the "Bonferroni"-method. Plots were created

using the R package "ggplot2" (Wickham 2016).

Contributions

ADG: Data acquisition, implementation and development of segmentation and classification

pipeline, training of classifiers, conceptual inputs, generation of figures, writing of the MS.

LA: Acquisition of experimental ground truth data with reporter mouse lines using 2PLSM imaging

and establishment of tomato-lectin labelling, proof-of-concept with initial development of

solutions for segmentation and classification.

CAB: Development of segmentation, application of StarDist, contributions to classification.

JJ: Data acquisition, training of StarDist for segmentation.

TK: Initial project idea and concept, writing of the MS, supervision.

JK: Conceptualization, implementation, experimentation, analysis, writing of the MS, project

management and supervision.

Acknowledgements

We want to thank Michaela Kaiser for her excellent technical support during preparation and

execution of experiments. We gratefully acknowledge the support by the German Research

Foundation (DFG) (SFB1158, project B08 awarded to TK), the data storage service SDS@hd,

supported by the Ministry of Science, Research and the Arts Baden-Württemberg (MWK) and the

DFG through grant INST 35/1314-1 FUGG, as well as the high-performance cluster bwForCluster

MLS&WISO, supported by the MWK and the DFG through Grant INST 35/1134-1 FUGG.

References

Arganda-Carreras, I., C. O. S. Sorzano, J. Kybic and C. Ortiz-de-Solórzano (2008). bUnwarpJ :

Consistent and Elastic Registration in ImageJ . Methods and Applications.

Asan, L., C. Falfan-Melgoza, C. A. Beretta, M. Sack, L. Zheng, W. Weber-Fahr, T. Kuner and J.

Knabbe (2021). "Cellular correlates of gray matter volume changes in magnetic resonance

morphometry identified by two-photon microscopy." Sci Rep 11(1): 4234.

Baddeley, A. and R. Turner (2005). "spatstat: An R Package for Analyzing Spatial Point Patterns."

Journal of Statistical Software 12(6): 1 - 42.

Boisvert, M. M., G. A. Erikson, M. N. Shokhirev and N. J. Allen (2018). "The Aging Astrocyte

Transcriptome from Multiple Regions of the Mouse Brain." Cell Rep 22(1): 269-285.

Clarke, L. E., S. A. Liddelow, C. Chakraborty, A. E. Munch, M. Heiman and B. A. Barres (2018).

"Normal aging induces A1-like astrocyte reactivity." Proc Natl Acad Sci U S A 115(8): E1896-

E1905.

Dawson, M. R., A. Polito, J. M. Levine and R. Reynolds (2003). "NG2-expressing glial progenitor

cells: an abundant and widespread population of cycling cells in the adult rat CNS." Mol Cell

Neurosci 24(2): 476-488.

Ero, C., M. O. Gewaltig, D. Keller and H. Markram (2018). "A Cell Atlas for the Mouse Brain."

Front Neuroinform 12: 84.

Guo, D., J. Zou, N. Rensing and M. Wong (2017). "In Vivo Two-Photon Imaging of Astrocytes in

GFAP-GFP Transgenic Mice." PLoS One 12(1): e0170005.

Hadjantonakis, A. K. and V. E. Papaioannou (2004). "Dynamic in vivo imaging and cell tracking

using a histone fluorescent protein fusion in mice." BMC Biotechnol 4: 33.

Herculano-Houzel, S. (2014). "The glia/neuron ratio: how it varies uniformly across brain

structures and species and what that means for brain physiology and evolution." Glia 62(9):

1377-1391.

Herculano-Houzel, S. and R. Lent (2005). "Isotropic fractionator: a simple, rapid method for the

quantification of total cell and neuron numbers in the brain." J Neurosci 25(10): 2518-2521.

Holtmaat, A., T. Bonhoeffer, D. K. Chow, J. Chuckowree, V. De Paola, S. B. Hofer, M. Hubener, T.

Keck, G. Knott, W. C. Lee, R. Mostany, T. D. Mrsic-Flogel, E. Nedivi, C. Portera-Cailliau, K.

Svoboda, J. T. Trachtenberg and L. Wilbrecht (2009). "Long-term, high-resolution imaging in the

mouse neocortex through a chronic cranial window." Nat Protoc 4(8): 1128-1144.

Nagendran, M., D. P. Riordan, P. B. Harbury and T. J. Desai (2018). "Automated cell-type

classification in intact tissues by single-cell molecular profiling." Elife 7.

Pakkenberg, B., M. V. Olesen, S. S. Kaalund and K. A. Dorph-Petersen (2019). "Editorial:

Neurostereology." Front Neuroanat 13: 42.

Palmer, A. L. and S. S. Ousman (2018). "Astrocytes and Aging." Front Aging Neurosci 10: 337.

Patki, N., R. Wedge and K. Veeramachaneni (2016). The Synthetic Data Vault. 2016 IEEE

International Conference on Data Science and Advanced Analytics (DSAA).

Percie du Sert, N., V. Hurst, A. Ahluwalia, S. Alam, M. T. Avey, M. Baker, W. J. Browne, A. Clark,

I. C. Cuthill, U. Dirnagl, M. Emerson, P. Garner, S. T. Holgate, D. W. Howells, N. A. Karp, S. E.

Lazic, K. Lidster, C. J. MacCallum, M. Macleod, E. J. Pearl, O. H. Petersen, F. Rawle, P. Reynolds,

K. Rooney, E. S. Sena, S. D. Silberberg, T. Steckler and H. Wurbel (2020). "The ARRIVE guidelines

2.0: Updated guidelines for reporting animal research." PLoS Biol 18(7): e3000410.

R. Core Team (2021). "R: A Language and Environment for Statistical Computing."

Schmidt, U., M. Weigert, C. Broaddus and G. Myers (2018). Cell Detection with Star-Convex

Polygons, Cham, Springer International Publishing.

Schmitz, C. and P. R. Hof (2007). Design-Based Stereology in Brain Aging Research. Brain Aging:

Models, Methods, and Mechanisms. D. R. Riddle. Boca Raton (FL).

Schwendele, B., B. Brawek, M. Hermes and O. Garaschuk (2012). "High-resolution in vivo

imaging of microglia using a versatile nongenetically encoded marker." Eur J Immunol 42(8):

2193-2196.

Tan, S. S., M. Kalloniatis, H. T. Truong, M. D. Binder, H. S. Cate, T. J. Kilpatrick and V. E.

Hammond (2009). "Oligodendrocyte positioning in cerebral cortex is independent of projection

neuron layering." Glia 57(9): 1024-1030.

van Griethuysen, J. J. M., A. Fedorov, C. Parmar, A. Hosny, N. Aucoin, V. Narayan, R. G. H. Beets-

Tan, J. C. Fillion-Robin, S. Pieper and H. Aerts (2017). "Computational Radiomics System to

Decode the Radiographic Phenotype." Cancer Res 77(21): e104-e107.

Wickham, H. (2016). "ggplot2: Elegant Graphics for Data Analysis."

Zeisel, A., H. Hochgerner, P. Lonnerberg, A. Johnsson, F. Memic, J. van der Zwan, M. Haring, E.

Braun, L. E. Borm, G. La Manno, S. Codeluppi, A. Furlan, K. Lee, N. Skene, K. D. Harris, J. Hjerling-

Leffler, E. Arenas, P. Ernfors, U. Marklund and S. Linnarsson (2018). "Molecular Architecture of

the Mouse Nervous System." Cell 174(4): 999-1014 e1022.

Zhang, C., C. Yan, M. Ren, A. Li, T. Quan, H. Gong and J. Yuan (2017). "A platform for

stereological quantitative analysis of the brain-wide distribution of type-specific neurons." Sci

Rep 7(1): 14334.

Figures

Figure 1. Comprehensive imaging of all cell nuclei and classification into cell types. (a) In-vivo 2-

photon microscopy of a 700 μm x 700 μm x 700 μm volume of neocortex. 3D reconstruction and

single imaging plane. **(b)** Features of nuclei belonging to identified cell types (red, only one

labeled cell type at a time) are used to train a neural network model for the classification of

H2GFP-labelled nuclei (green).

Figure 2. Features of the cell nucleus reveal the corresponding cell type. (a) Neurons (yellow),

astroglia (blue), oligodendroglia (turquoise), microglia (green), endothelial cells (red) were

labelled using different strategies **(b). (c)** Nuclei belonging to a specific cell type can be identified

by a red fluorescent marker (maximum intensity projections; z = 20 μm). 3D-renderings of

individual nuclei are shown in the lower panel. **(d)** Visualization of a subset of radiomics features

selected by an algorithm for classification. **(e)** Polar plots showing nuclear features of different

cell types. **(f)** Comparison of nuclear features between cell types (Wilcoxon test, n of neurons:

135, n of astroglia: 137, n of microglia: 62, n of oligodendroglia: 72, n of endothelial cells: 155, p

< 0.05 *, p < 0.01 **, p < 0.001 ***), N = Neuron, AGlia = Astroglia, MGlia = Microglia, OGlia =

Oligodendroglia, EC = endothelial cells.

Figure 3. Training a neuronal network for cell type classification. (a) Schematic depiction of the

training process. After segmenting raw data with StarDist, features of nuclei were calculated, and

synthetic data was generated. A separate classification network was trained for each cell type. **(b)**

Precision, recall, false discovery rate and false negative rates for the test dataset. **(c)** Mean

accuracy for all cell types in the training dataset (15 % of the whole dataset). **(d)** Mean accuracy

for all cell types in the test dataset (15 % of the whole dataset). **(e)** Comparison of nucleus volume

of each cell type. n of neurons: 135, n of astroglia: 137, n of microglia: 62, n of oligodendroglia:

72, n of endothelial cells: 155 **(f)** Comparison of mean surface to volume ratio of each cell type.

n of neurons: 135, n of astroglia: 137, n of microglia: 62, n of oligodendroglia: 72, n of endothelial

cells: 155 **(g)** Number of nuclei per mm^3., n per class is 8 **(h)** Comparison of the number of nuclei

for each cell type between baseline and 12-week timepoint, n per class is 8 **(i)** Comparison of

normalized number of nuclei after 12 weeks, n per class is 8 **(j)** Comparison of mean normalized

nucleus volume after 12 weeks, n per class is 8 (Wilcoxon test, p-values were corrected for

multiple comparisons using the Bonferroni-method, $p < 0.05$*, $p < 0.01$ **, $p < 0.001$ ***), N =

Neuron, AGlia = Astroglia, MGlia = Microglia, OGlia = Oligodendroglia, EC = endothelial cells.

a i ii iii iv

b X-Z projection

100 µm

c i Layer 1, Layer 2/3, Layer 5 DAPI ii

500 µm 100 µm

DAPI profile

2P Nuclei density

Depth (µm)

d

	Pre induction	Post induction			Magnified	

eGFP | eGFP | Red labelled | Composite | POST | PRE

Astroglia (GFAP - Tamoxifen - tdTomato)

Oligodendroglia (PLP - Tamoxifen - tdTomato)

Endothelial (Tie2 - Tamoxifen - tdTomato)

50 µm 10 µm

e In-vivo 2P

Intracranial injections

Cranial window implantation

Tamoxifen i.p

f Post injection imaging

eGFP | Red labelled | Composite

Neuron (mcherry)

Microglia (TomatoLectin - Dylight 594)

g Raw | Ground truth

i ii

iii iv

StarDist | Composite

50 µm

h StarDist training

Ground truth segmentation

Number of nuclei

3D volumes

Detection accuracy   Detection precision

Mean = 94.3   Mean = 98.4

Validation crops   Validation crops

Supplementary Figure 1 (a) (i,ii,iii) Chronic cranial window implantation using a curved glass

cover slip and a custom 3D printed holder. **(iv)** Fixation of the anesthetized mouse in the custom

two-photon imaging holder. **(b) (i-iv)** Two-dimensional x-z projection of a two-photon image stack

(c) Imaging position on a DAPI stained brain slice. Slice thickness 50 µm. DAPI intensity profile

showing similar distribution of intensity as a two-photon image of nuclei. The decreasing nucleus

density in the two-photon image stack is a result of reduced fluorescence signal at higher depths.

Inlay: Nucleus density distribution of sub-volumes used for data analysis. **(d)** Visualization of

crosstalk between the eGFP and tdTomato signal after induction with tamoxifen. Overlay of pre

(cyan) and post (yellow) GFP after image alignment with the ImageJ plugin bUnwarpJ (Arganda-

Carreras, Sorzano et al. 2008).

(e) Intracranial and intraperitoneal injections were performed on reporter line mice. **(f)** No

crosstalk between GFP and mcherry signal (upper panel) or eGFP and Tomato lectin-Dylight 594

signal (lower panel) **(g)** Raw data of H2GFP signal (i), manually labelled ground truth (ii), StarDist

segmentation (iii), and overlay of ground truth (red) and StarDist segmentation (green) (iv). **(h)**

Upper panel: Count of manually segmented ground truth nuclei for StarDist training, each color

depicts an individual mouse. Lower panel: StarDist nucleus detection accuracy and precision,

each bar represents an individual imaging volume.

Supplementary Figure 2. Combinations of nuclear morphology, intensity, and texture features

show clear distinction between cell types. **(a)** Flatness, gray level non-uniformity. **(b)** Minor axis

length, Flatness. **(c)** First order minimum, Elongation. **(d)** Large area emphasis, First order 10^{th}

percentile. **(e,f)** Combinations of features in three dimensions. Yellow = Neurons, Blue = Astroglia,

Green = Microglia, Turqouise = Oligodendroglia, Red = Endothelial cells. N = Neuron, AGlia =

Astroglia, MGlia = Microglia, OGlia = Oligodendroglia, EC = endothelial cells.

Supplementary Figure 3 (a) For each cell type, green and red channel data from the two-photon volumetric images were overlayed and nuclei labelled in both channels were identified. A sequential forward feature selection algorithm was applied, which selected 12 features from the 107 radiomics feature. Synthetic data was generated from the radiomics features of the original dataset. The combined dataset was used to train a classifier with the training, validation and test data comprising of 70%, 15%, and 15% of the combined dataset, respectively. **(b)** For

555   classification, volumetric data from H2GFP mice were segmented, radiomics features were

556   extracted and only 12 features from the training were selected. The data was analyzed by all 5

557   classifiers (one for each cell type) and the outputs of the classifier were combined for each

558   nucleus.

Supplementary Figure 4. Synthetic training data was generated from the original dataset. **(a)**

Distribution of synthetic data is comparable to original data, yellow = original data, blue =

synthetic data. **(b)** Statistical analysis of similarity between the distribution of original data and

distribution of synthetic data (K-S test; mean = 83.68%). **(c)** Datasets with different amounts of

synthetic data were created and the training accuracy was compared between them, orig =

original dataset, orig * 2.5 = dataset containing 2.5 times the amount of data as the original

dataset, orig * 2.5 (down sampled) = dataset down sampled to original size from the dataset

containing 2.5 times the amount of data as the original dataset, orig * 9 = dataset containing 9

times the amount of data as the original dataset, orig * 9 (down sampled) = dataset down

sampled to original size from the dataset containing 9 times the amount of data as the original

dataset. **(d)** Mean accuracy of classifiers trained on different datasets, light blue = orig * 2.5, dark

579   blue = orig * 2.5 (downsampled), green = orig * 9, yellow = orig * 9 (down sampled), red = orig.

580   **(e)** Table of manually identified nuclei belonging to each cell type.

Supplementary Figure 5. Confusion matrices for each classifier for the test dataset. On the y-axis

the output class of the classifier is shown, on the x-axis the target class. As the classifier

distinguishes between the desired class and the "other" class, the confusion matrix consists only

of 4 fields. Green fields illustrate correct identification of target and "other" class, blue fields

illustrate erroneous identifications. AGlia = Astroglia, MGlia = Microglia, OGlia = Oligodendroglia,

EC = endothelial cells.

a

Raw data   Segmented nuclei   Classified nuclei

b

Neurons   Astroglia   Microglia   Oligodendroglia   Endothelial

c

Maximum Intensity projection

Raw data   Segmented nuclei   Classified nuclei

50 μm

597   **Supplementary Figure 6. 3D visualization of comprehensive cell type composition. (a)** 3-

598   dimensional visualization of raw, segmented and classified data (700 μm x 700 μm x 700 μm

599   volume). **(b)** Nuclei distribution for each cell type after classification. **(c)** Maximum intensity z-

600   projection of a sub volume (150 μm x 150 μm x 100 μm) showing raw, segmented and classified

601   nuclei.

Radiomics features

Morphology		Intensity	Texture				
2-Dimensional	3-Dimensional	First Order Statistics	Gray Level Co-occurence Matrix (GLCM)	Gray Level Run Length Matrix (GLRLM)	Gray Level Size Zone Matrix (GLSZM)	Neighboring Gray Tone Different Matrix (NGTDM)	Gray Level Dependence Matrix (GLDM)
Mesh Surface	Mesh Volume	Energy	Autocorrelation	Short Run Emphasis (SRE)	Small Area Emphasis (SAE)	Coarseness	Small Dependence Emphasis (SDE)
Pixel Surface	Voxel Volume	Total Energy	Joint Average	Long Run Emphasis (LRE)	Large Area Emphasis (LAE)	Contrast	Large Dependence Emphasis (LDE)
Perimeter	Surface Area	Entropy	Cluster Prominence	Gray Level Non-Uniformity (GLN)	Gray Level Non-Uniformity (GLN)	Busyness	Gray Level Non-Uniformity (GLN)
Perimeter to Surface ratio	Surface Area to Volume ratio	Minimum	Cluster Shade	Gray Level Non-Uniformity Normalized (GLNN)	Gray Level Non-Uniformity Normalized (GLNN)	Complexity	Dependence Non-Uniformity (DN)
Sphericity	Sphericity	10th percentile	Cluster Tendency	Run Length Non-Uniformity (RLN)	Size-Zone Non-Uniformity (SZN)	Strength	Dependence Non-Uniformity Normalized (DNN)
Spherical Disproportion	Compactness 1	90th percentile	Contrast	Run Length Non-Uniformity Normalized (RLNN)	Size-Zone Non-Uniformity Normalized (SZNN)		Gray Level Variance (GLV)
Maximum 2D diameter	Compactness 2	Maximum	Correlation	Run Percentage (RP)	Zone Percentage (ZP)		Dependence Variance (DV)
Major Axis Length	Spherical Disproportion	Mean	Difference Average	Gray Level Variance (GLV)	Gray Level Variance (GLV)		Dependence Entropy (DE)
Minor Axis Length	Maximum 3D diameter	Median	Difference Entropy	Run Variance (RV)	Zone Variance (ZV)		Low Gray Level Emphasis (LGLE)
Elongation	Maximum 2D diameter (Slice)	Interquartile Range	Difference Variance	Run Entropy (RE)	Zone Entropy (ZE)		High Gray Level Emphasis (HGLE)
	Maximum 2D diameter (Column)	Mean Absolute Deviation (MAD)	Joint Energy	Low Gray Level Run Emphasis	Low Gray Level Zone Emphasis		Small Dependence Low Gray Level Emphasis
	Maximum 2D diameter (Row)	Robust Mean Absolute Deviation (rMAD)	Joint Entropy	High Gray Level Run Emphasis	High Gray Level Zone Emphasis		Small Dependence High Gray Level Emphasis
	Major Axis Length	Root Mean Squared (RMS)	Informational Measure of Correlation (IMC) 1	Short Run Low Gray Level Emphasis	Small Area Low Gray Level Emphasis		Large Dependence Low Gray Level Emphasis
	Minor Axis Length	Standard Deviation	Informational Measure of Correlation (IMC) 2	Short Run High Gray Level Emphasis	Small Area High Gray Level Emphasis		Large Dependence High Gray Level Emphasis
	Least Axis Length	Skewness	Inverse Difference Moment (IDM)	Long Run Low Gray Level Emphasis	Large Area Low Gray Level Emphasis		
	Elongation	Kurtosis	Maximal Correlation Coefficient (MCC)	Long Run High Gray Level Emphasis	Large Area High Gray Level Emphasis		
	Flatness	Variance	Inverse Difference Moment Normalized (IDMN)				
		Uniformity	Inverse Difference (ID)				
			Inverse Difference Normalized (IDN)				
			Inverse Variance				
			Maximum Probability				
			Sum Average				
			Sum Entropy				
			Sum of Squares				

610 **Supplementary Table 1:** Features marked in green were used for cell type classification

Selected Features			Definition
Morphology	3-Dimensional	Maximum 3D diameter	Largest pairwise Eucledian distance between two surface vertices
		Maximum 2D diameter (Row)	Largest pairwise Eucledian distance between two surface vertices in the sagittal plane
		Minor Axis Length	Second largest axis length of the ROI-enclosing ellipsoid
		Elongation	Relationship between two largest principal components in the ROI
		Flatness	Relationship between the largest and the smallest principal components in the ROI
Intensity	First order statistics	Minimum	Minimum gray level intensity within the ROI
		10th percentile	10th percentile of all the voxels in the ROI
		Interquartile Range	Difference between the 75th and 25th Percentile
Texture	Gray Level Co-occurence Matrix (GLCM)	Joint Entropy	Measure of randomness or variability in the neighborhood intensity values
	Gray Level Run Length Matrix (GLRLM)	Gray Level Non-Uniformity (GLN)	Similarity of gray-level intensity values in the image
	Gray Level Size Zone Matrix (GLSZM)	Large Area Emphasis (LAE)	Measures the distribution of large area size zones. Higher value indicates more coarseness
	Neighboring Gray Tone Difference Matrix (NGTDM)	Coarseness	Measures the average difference between the center voxel and its neighborhood

613   **Supplementary Table 2:** Morphology, intensity and texture features used for classification

615   **Supplementary Movie 1. Illustration of nucleus segmentation and classification using NuClear.**

616   Raw data of Histon-2B-eGFP-labeled nuclei (green), nucleus segmentation (randomly colored)

617   and classification (apricot = neurons, red = microglia, blue = oligodendrocyts, dark green =

618   astrocytes, magenta = endothelial cells)