**Title page**

**Integrative multispecies omics reveals a hierarchy of cold-responsive regulatory**

**network launched by circadian components in rosids**

Liangyu Guo[1,†], Zhiming Xu[1,†], Shuo Wang[1,†], Yuqi Nie[1], Xiaoxue Ye[2], Xuejiao Jin[1,*],

Jianhua Zhu[3,4,*], and Wenwu Wu[1,*]

[1]State Key Laboratory of Subtropical Silviculture, School of Forestry and

Biotechnology, Zhejiang A&F University, Lin'an, Hangzhou 311300, China.

[2]Institute of Tropical Biosciences and Biotechnology, Chinese Academy of Tropical

Agricultural Sciences, Haikou 571101, Hainan, China

[3]School of Life Sciences, Anhui Agricultural University, Hefei, 230036, China

[4]Department of Plant Science and Landscape Architecture, University of Maryland,

College Park, Maryland 20742, USA

[†]Equal contributors;

*Correspondence should be addressed to X.J ([jinxuejiao1991@cau.edu.cn](mailto:jinxuejiao1991@cau.edu.cn)), J.Z.

([jhzhu@umd.edu](mailto:jhzhu@umd.edu)) or W.W. ([wwwu@zafu.edu.cn](mailto:wwwu@zafu.edu.cn), Dr. Wu is fully responsible for the

distribution of the materials associated with this article).

Running title: A hierarchical cold-responsive regulatory network

Color Figures: 6

Tables: 1

Supporting information: 2 Supplementary Figures and 9 Supplementary Tables

## Abstract

Elucidating regulators and molecular mechanisms underlying gene transcriptional and post-transcriptional co-regulatory network is key to understand plant cold-stress responses. Previous studies were mainly conducted on single species and whether the regulators and mechanisms are conserved across different species remains elusive. Here, we selected three species that diverged at early evolution of rosids (93–115 million years ago) and integrated phylotranscriptome and ChIP/DAP-seq datasets to identify cold-responsive regulators and their regulatory networks. First, we found over ten thousand cold-responsive genes including differentially expressed genes (DEGs) and alternative splicing genes (DASGs) in each species. Among the DEGs, genes encoding a set of transcription factors (TFs) (AP2/ERF, MYB, WRKY, NAC, etc.) and RNA binding proteins (RBPs) (Ribosomal, RRM, DEAD, Helicase_C, etc.) are conserved in cold responses in rosids. Compared to TFs, RBPs show a delayed cold-responsive pattern, likely suggesting a hierarchical regulation of DEGs and DASGs. Between DEGs and DASGs, we identified 259 overlapping DE-DASG orthogroups and interestingly, pathway analysis on each dataset of DEGs, DASGs, and DE-DASGs coincidently shows an enrichment of circadian rhythm. Evidentially, many circadian components are cold-regulated at both transcriptional and post-transcriptional levels. Moreover, we reasoned 226 cold-responsive genes regulated by at least two of five circadian components (CCA1, LHY, RV4, RVE8, and RVE7) in rosids. Finally, we unveiled a conserved hierarchical network in dynamic transcriptional and post-transcriptional regulation of cold-responsive genes launched by circadian components in rosids. Together, our results provide insights into core regulators and mechanisms underlying cold-responsive regulatory network across rosids, despite a long evolutionary history.

**Key words**: transcription factor; RNA binding protein; circadian clock; hierarchical network; cold stress; rosids

## Introduction

For adaptation to adverse environmental conditions, angiosperms have evolved sophisticated and advanced mechanisms and become the largest land plant lineage with more than 350,000 known species widely distributed from tropical to polar terrestrial zones[1]. Among the adverse environmental conditions, seasonal and daily variation of temperature is a fundamental component that affects plant growth, productivity, and geographical distribution[2,3]. Under cold stress, plants undergo physiological and molecular changes that increase the levels of specific proteins, metabolites, and phytohormones, which are generally regulated by transcription factors (TFs) and RNA-binding proteins (RBPs) at transcriptional and post-transcriptional levels[3-6].

Thousands of differentially expressed genes (DEGs) were identified involved in cold stress[7-9] and many TFs have been suggested in regulating and managing their expression[3,4,10]. The best characterized TFs are C-repeat binding factor (*CBF*) genes that are rapidly and transiently induced by cold stress and can enhance freezing tolerance of plants through binding to *cis*-elements (CCGAC) in the promoters of cold-responsive (*COR*) genes[11,12]. RNA-seq analysis of *Arabidopsis cbf123* triple mutant showed that hundreds of *CORs* are regulated through a CBF-dependent pathway[13,14]. Activation of CBF target *CORs*, such as *COR15A*, *COR47*, *KIN1*, and *RD29A*, leads to an increase in plant freezing tolerance[12,15,16]. In addition, cold responsiveness of plants is regulated by circadian clock system, which is involved in plant stress responses through controlling daily energy expenditure and enabling plants to grow and reproduce under seasonal and daily temperature variation[17-21]. In the clock system, some circadian clock-related TFs, such as CIRCADIAN CLOCK ASSOCIATED 1 (CCA1), LATE ELONGATED HYPOCOTYL (LHY), and REVEILLES (RVEs), have been proven involved in cold tolerance by directly regulating the expression of downstream *CORs*, such as *CBFs*, *COR15A/15B*, *COR28,* and *COR27*[2,22-24].

On the other hand, post-transcriptional regulation is also a key process in

reprogramming gene expression for plant to tolerate or survive constantly changing in

temperature[3,4]. Similar to TFs in regulating gene transcription, RBPs are essential in

controlling post-transcriptional RNA processing, including capping, polyadenylation,

alternative splicing (AS), RNA export, translation, and turnover[7,25]. Under cold stress,

upregulation and/or downregulation of RBP genes may cause differentially

post-transcriptional RNA processing. During RNA processing, AS is an important

post-transcriptional mechanism by producing different transcripts and possibly protein

variants from a single gene for increasing transcriptome and proteome diversity[26]. The

concentration, localization, and activity of RBPs determine AS events to produce

different transcripts[27,28]. In plants, extensive AS events have been implicated in

various abiotic stresses, including cold, salt, heat and other environmental

stresses[7,8,29,30]. For example, genome-wide AS profiling in *Arabidopsis* showed a

massive differentially alternative splicing genes (DASGs) after cold treatment and

suggested that plants may fine-tune AS events of gene transcripts for adapting to cold

stress[7]. Among DASGs, *CCA1*, a core component of circadian clock, has two splice

variants: CCA1α and CCA1β. CCA1β competitively inhibits the DNA binding

activity of CCA1α and LHY by forming nonfunctional heterodimers of

CCA1β-CCA1α and CCA1β-LHY[31]. Under cold stress, CCA1β production is

repressed to release the activity of CCA1α and LHY, whereby CCA1α and LHY

promote the expression of downstream *CORs* for improving plant cold tolerance[31].

In the past years, numerous cold-induced DEGs and DASGs were identified in

different plant species by high-throughput methods[7,8,29,30,32]. Identifying core

regulators (e.g., TFs and RBPs) and elucidating molecular mechanisms underlying the

outcome of DEGs and DASGs are key to understand the reprogramming of gene

regulatory network in plant response to cold stress. The previous studies were

generally performed on single plant species, especially the reference model

*Arabidopsis thaliana*, and whether the regulators and mechanisms underlying

transcriptional and post-transcriptional changes are conserved in different plant

lineages remains elusive. In this study, we focused on rosids, a large group of eudicots

in angiosperms, and selected three plant species: *Carya illinoinensis* and *Populus*

*trichocarpa* from fabids (rosid I) and *Arabidopsis thaliana* from malvids (rosid II).

Fabids and malvids are two main clades of rosids, comprising of >40,000 and ~15,000

species, respectively[33]. The three plant species diverged during early evolution of

rosids and evolved independently for over 100 million years[34]. The long evolutionary

history caused great differences in their growth habits and geographic distributions,

and accordingly, we asked whether the regulators and mechanisms underlying gene

transcriptional and post-transcriptional regulation are conserved across rosids in

response to cold stress.

To solve the question, we performed omics analysis of genomes, transcriptomes,

and regulomes and identified common core regulators in regulating cold responses

across the three plant species. First, by RNA-seq experiments, we investigated global

expression profiles of cold-induced DEGs and DASGs in each of the three studied

species. Subsequently, we analyzed cold responses of TFs and RBPs and identified

those TF and RBP families that are conserved in cold responses, which likely

contribute to DEGs and DASGs in rosids. Moreover, we explored overlapping genes

(i.e., DE-DASGs) of DEGs and DASGs and performed homologous analysis to

identify conserved DE-DASG orthogroups in rosids. Finally, we selected five

circadian components and integrated transcriptomes and regulome maps to identify

their target *CORs* for constructing a hierarchical cold-responsive regulatory network.

**Results**

**Global gene expression profiles in rosids under cold stress**

Physiological and molecular changes in plants under cold stress are usually caused by

up- or downregulated expression of genes[3]. To capture the landscape of DEGs under

cold stress, we performed RNA-seq experiments in three rosid species before and

after cold treatments (**see Methods and materials**). Based on RNA-seq analysis, we

identified a huge number of cold-induced DEGs: 7,579, 8,878, and 8,480 respectively

in *A. thaliana*, *C. illinoinensis*, and *P. trichocarpa* (**Supplementary Table 1**). Notably,

more upregulated than downregulated DEGs are observed in all three plants (**Fig 1a**).

Expression profiles show that these upregulated DEGs are mainly clustered in three

expression patterns following three timepoints of cold treatments: 2 hours (h), 24 h,

168 h after cold treatments (**Fig 1b**). Inspecting the expression patterns, we found that

the number of upregulated DEGs in *A. thaliana* is significantly more at 2 h but less at

168 h of cold stress treatment than the counterparts in other two species. Likewise,

there are significantly more downregulated DEGs at 2 h and less downregulated

DEGs at 168 h in *A. thaliana* than those in *C. illinoinensis* and *P. trichocarpa* (**Fig 1b**).

These findings reveal that over seven thousand genes are differentially cold-regulated

in rosids with three common expression patterns but likely in different

cold-responsive speeds: an early extensive response in *A. thaliana* but a delayed

extensive response in *C. illinoinensis* and *P. trichocarpa*.

**Common biological processes of cold responses in rosids**

Despite of different cold-responsive speeds between the three plant species, Gene

Ontology (GO) term enrichment analysis of the DEGs showed a very similar

enrichment of biological processes (**Fig 1c, 1d**). The upregulated DEGs are mainly

enriched in abiotic stress processes, e.g., response to hypoxia, water, and cold, and

circadian rhythm (**Fig 1c**). Additionally, regulations of transcription and RNA

processing are also amongst the most enriched GO terms, which suggests that the

regulators (e.g., TFs and RBPs) at transcriptional and post-transcriptional level are

likely to be more activated under cold stress (**Fig 1c**). The downregulated DEGs are

mainly enriched in processes such as catalytic activity, photosynthesis, developmental

growth, small molecular and/or lipid biosynthetic/metabolic process (**Fig 1d**). It is

reasonable that for improving cold tolerance, plants need to reduce expression of the

genes involved in activities of plant development and growth[3].

**Conserved cold-responsive TF families in rosids**

Differential changes of gene expression at transcriptional level are likely caused by

fluctuations of TFs[3,7]. To detect the roles of TFs under cold stress, we first identified a

total of 1,717, 1,899, and 2,466 TFs in *A. thaliana*, *C. illinoinensis*, and *P. trichocarpa*

genomes, respectively (**Supplementary Table 2, see Methods and materials**).

Transcriptome analysis shows that over five hundred TFs are differentially

cold-regulated, covering a high proportion of the genome TFs, that is, 32%, 35%, and

22% respectively in *A. thaliana*, *C. illinoinensis*, and *P. trichocarpa* (**Fig 2a,**

**Supplementary Table 2**). Notably, there are more upregulated than downregulated

TFs and their expression patterns in cold-responsive speeds agree well with the

aforementioned global expression patterns of DEGs (**Fig 1b, Fig 2b**). This finding

suggests more upregulated TFs might be involved in modulating differentially

regulated genes under cold stress.

Moreover, we classified all TFs into gene families based on their protein domain

architectures (**Supplementary Table 2, see Methods and materials**). Interestingly,

the up- and downregulated TFs are respectively clustered in common gene families in

three plant species (**Fig 2c, 2d**). For example, the main families of upregulated TFs

are AP2/ERF, MYB, WRKY, and NAC, which are the best-known TF families

involved in abiotic stresses[35,36]. The counterparts of downregulated TFs are MYB,

bHLH, Homeobox, and AP2/ERF. Notably, AP2/ERF and MYB are two most

prominent TF families with many more upregulated members but also a considerable

number of downregulated members in all three plants under cold stress. It is

explainable that members from the same TF families can often function in a negative

feedback mechanism in abiotic responses[37,38]. According to the maximum parsimony

principle, we suggest that the conserved cold-responsiveness of those TF families in

three plant species is largely descended from their common ancestor.

**Global alternative splicing profiles in rosids under cold stress**

In addition to transcriptional regulation, we identified an enrichment of

post-transcriptional regulation, such as RNA processing, in response to cold stress

(**Fig 1c**). Based on transcriptome analysis, we detected 2,572, 7,428, and 7,817

cold-induced DASGs, which occupy 10%, 24%, and 18% of coding genes in *A.*

*thaliana*, *C. illinoinensis*, and *P. trichocarpa*, respectively (**Fig 3a**, **Supplementary**

**Table 3**). Among DASGs, many have multiple AS sites, producing 3,192, 11,972, and

13,373 differentially AS events (DASEs), respectively in *A. thaliana*, *C. illinoinensis*,

and *P. trichocarpa* (**Fig 3a**, **Supplementary Table 3**). This is mainly caused by AS

occurrences at the same or different exon/intron boundaries across a gene, which leads

to more diversities of RNA transcripts. In consistence with previous studies[7,30], intron

retention (IR) is the most common AS mode in three plant species (**Fig 3b**). Despite a

relatively small role of skipping exon (SE) in *A. thaliana*, SE provides a crucial role

in other two plants (*C. illinoinensis* and *P. trichocarpa*). Further, GO term enrichment

analysis shows that these DASGs from three different plant species are enriched in

very similar biological processes, such as metabolic processes, gene expression, RNA

splicing, and circadian rhythm (**Fig 3c**). Given a key role of RBPs in RNA

processing[7,25], thousands of cold-induced DASEs might at least partially result from

fluctuations of RBP genes.

**A delayed cold responsivity of RBP genes**

In *A. thaliana*, we collected a total of 788 high-confident RBPs that were

experimentally confirmed by RNA-binding assays[25,39]. Using the RBPs for homologs

searching, we identified 1,110 and 1,379 RBPs in *C. illinoinensis* and *P. trichocarpa*,

respectively (**Fig 4a**, **Supplementary Table 4**). Transcriptome analysis shows that

36%, 31%, and 35% RBPs are differentially regulated by cold stress in *A. thaliana*, *C.*

*illinoinensis*, and *P. trichocarpa*, respectively (**Supplementary Table 4**). Like TFs,

we identified more upregulated RBPs than downregulated ones, especially in *A.*

*thaliana* (**Fig 4a**). Interestingly, in contrast to TF genes, RBP genes show a delayed

responsive pattern after cold treatment in all three plant species (**Fig 2b, 4b**). For

example, only a relatively small number of RBP genes are observed to be

differentially regulated at early stage (2 h) of cold stress. This finding shows that cold

responsivity of RBP genes lags after that of TF genes, which might suggest a

hierarchy of temporal dynamics of transcriptional and post-transcriptional regulatory

network in response to cold stress.

**Conserved cold-responsive RBP families in rosids**

Based on RNA-binding domains, we classified RBP genes into different gene families

(**Supplementary Table 4**). Among the top 15 cold-responsive RBP families,

ribosomal and RRM are two most prominent families with more genes being

upregulated than downregulated under cold stress in all three plant species (**Fig 4c-d**).

DEAD and Helicase_C families are mainly enriched in the upregulated RBPs (**Fig 4c**),

while Chloroa_b-bind and MIP families are more clustered in the downregulated

RBPs (**Fig 4d**). Like TF families, according to the maximum parsimony principle, we

suggest that the conserved cold responsivity of the RBP families in three plant species

is largely descended from their common ancestor.

**Cold association of transcriptional and post-transcriptional activities in rosids**

As described above, thousands of genes were identified as cold-induced DEGs and/or

DASGs in three plants (**Fig 1, Fig 3**). Further, we identified that upregulated DEGs

show a significant overlapping association with DASGs, that is, 558, 1340, and 1419

up-DE-DASGs respectively in *A. thaliana*, *C. illinoinensis*, and *P. trichocarpa* (**Fig 5a**,

**Supplementary Table 5**). Similarly, downregulated DEGs also show a significant

association with DASGs in the three plant species. The findings suggest that many

cold-responsive genes are differentially regulated not only by TFs at transcriptional

level but also by RBPs at post-transcriptional level, and these genes might play an

important role in rosids under cold stress. In the following sections, we determined

whether these genes across the three studied plants are homologous and function in a

conserved way.

**Identification of 259 conserved orthogroups in rosids**

Biased retention and loss of gene duplicates in plants after independent

whole-genome and/or small-scale duplications lead to complex and intertwined

evolution of co-orthologs and in-paralogs across different species[35,40]. To determine

common cold-responsive genes between species, we classified those genes that are

differentially regulated at both transcriptional and post-transcriptional levels into

orthogroups: sets of homologous genes descended from the recent common ancestor

of the plants[41] (**see Methods and materials**). For example, 558, 1,340, and 1,419

up-DE-DASGs are classified into 305, 626, and 625 orthogroups, respectively in *A.*

*thaliana*, *C. illinoinensis*, and *P. trichocarpa* (**Supplementary Table 6**). Among these

orthogroups, Venn diagram analysis shows a significant overlapping of 150

orthogroups among three plants (**Fig 5b, Supplementary Table 6**). From the 150

orthogroups, we extracted 186 *A. thaliana* genes and performed GO term enrichment

analysis (**Fig 5c**). As expected, the genes are mainly enriched in abiotic stresses, such

as response to abiotic stresses (salt and cold stress) and circadian rhythm (**Fig 5c**).

The GO term enrichments of the counterparts in other two species have similar results

(**Supplementary Fig 1**). The same analysis was also conducted for down-DE-DASGs,

and we identified 109 overlapping orthogroups with many genes enriched in diverse

enzymatic activities (**Fig 5d, 5e, Supplementary Fig 1**, **Supplementary Table 7**).

The findings together show a total of 259 orthogroups (150 + 109) that are regulated

at both transcriptional and post-transcriptional levels in a conserved manner in rosids

under low temperature, likely suggesting a conserved role of these orthogroups under

cold stress.

**Seven striking TF orthogroups involved in circadian rhythms**

Among the 259 orthogroups, there are 27 upregulated TF orthogroups

(**Supplementary Table 8**) and it is noteworthy that seven of them encode known

circadian clock components including *CCA1*/*LHY*, *RVE4/8*, *RVE7*, *PRR9*, *BBX19*,

*COL9*, and *JMJD5* genes (**Table 1**). Our results are consistent with previous reports

that *CCA1, LHY* and *PRR9* can be regulated at transcriptional and post-transcriptional

levels under cold stress in *Arabidopsis*[31,42]. We showed in our study these circadian

genes are also cold-regulated in *C. illinoinensis*, and *P. trichocarpa* and we found that

additional circadian clock related genes are differentially cold-regulated at both

transcriptional and post-transcriptional levels in rosids (**Table 1**). This explains well

aforementioned repeated observations that the biological process of circadian rhythm

is enriched (**Fig 1c**, **Fig 3c**, **Fig 5c**). The seven orthogroups are upregulated under

cold stress in all three plant species, and in contrast, AS types of genes in each

orthogroup show a great diversity, which suggest a conservation of gene expression

but a rapid evolution of AS events after rosid radiation (**Table 1**). Moreover, MYB is

the most prominent TF family including three orthogroups (*CCA1/LHY*, *RVE7*, and

*RVE4/8*). The finding suggests that circadian clock components might play important

roles in regulating cold responses in a conserved manner in rosids.

**Cold-responsive genes regulated by circadian clock components**

First, we collected experimentally genome-wide regulome maps of *CCA1*, *LHY*,

*RVE4*, *RVE7*, and *RVE8* in *A. thaliana* with data from chromatin immunoprecipitation

(ChIP-seq)[43] and DNA affinity purification followed by deep sequencing (DAP-seq)[44].

Subsequently, we developed a stringent pipeline to identify cold-responsive genes

regulated by the five TFs (see **Materials and methods**). In brief, we integrated the

ChIP-seq and DAP-seq data to identify *A. thaliana* genes bound by either of the five

TFs and investigated their orthogroup genes in *C. illinoinensis* and *P. trichocarpa*

whose promoters (1-Kb) also have at least one *cis*-regulatory binding site of the TFs.

Finally, and of importance, we integrated phylotranscriptomic data to identify those

orthologous genes that have DNA binding sites of the five TFs and are differentially

cold-regulated in all three plant species. Thus, we designated those orthologous genes

as <u>Co</u>nserved <u>COR</u> gene<u>s</u> (CoCORs) targeted by clock components *CCA1*, *LHY*,

*RVE4*, *RVE7*, or *RVE8* in rosids (**Fig 6a, Supplementary Table 9**).

As shown in Fig 6a, the five circadian components regulate different numbers of

CoCORs but with a common pattern of dominant upregulation in three rosids. For

instance, as compared to only 17 downregulated CoCORs, there are 101 CoCORs

identified to be upregulated by CCA1 (**Fig 6a, Supplementary Table 9**). GO term

analysis further shows that these CoCORs are enriched in environmental stresses,

such as response to light, cold, and circadian rhythm (**Fig 6b**). Moreover, Venn

diagram shows a significance of overlapping CoCORs regulated by the five TFs (**Fig

6c**), and among the overlapping CoCORs, 16 are regulated by all five TFs and 210 are

regulated by at least two of the five TFs. The intertwined DNA binding of CoCORs

by the five circadian TFs suggests a complex regulatory network launched by

circadian clock components.

**A conserved hierarchy of cold-responsive regulatory network in rosids**

To construct the network, we employed the CoCORs involved in abiotic stresses that

are directly regulated by the five circadian components. Taking account of

cold-responsive speeds of the genes under cold treatments at 2, 24, and 168 h and also

their regulatory relationships, we constructed a hierarchical regulatory network with

three layers launched by the five circadian clock components and elaborated this

network as follows.

The first layer is composed of 30 TFs including many well-known cold-regulated

genes, such as *CBFs*, *PRR5*, *ZAT6*, *ZAT10*, *COL9*, *BT2* etc. In this layer, the five

clock components play a dominant role to upregulate the expression of 21 TFs and in

turn work together with these upregulated TFs to further regulate the expression of

downstream genes (e.g., RBPs and structural genes in the second and third layers). In

addition to upregulated TFs launched by circadian clock components, there are four

downregulated TFs (*TCP7*, *EBS*, *IBH1*, and *PIF1*). This demonstrates a potential

negative regulatory loop between differentially regulated TFs for controlling the

balance between plant growth and cold tolerance. Evidentially,

phytochrome-interacting factor 1 (PIF1) negatively regulates plant freezing tolerance

likely through repressing the expression of *CBFs*[45].

The second layer is composed of 14 RBPs. Among the 14 RBPs, 12 are

upregulated by the first layer TFs. These 12 RBPs function in a feedback mechanism

by potentially regulating AS of TFs in the first layer and function in regulating AS of

the downstream layer genes. For example, by analyzing individual nucleotide

resolution crosslinking and immunoprecipitation (iCLIP)-seq experimental data[46], we

found that GRP7 can directly bind to transcripts of *BT2* and *HB6* in the first layer and

*COR27*, *KCS6*, *BAP1*, *COR413-PM1*, *ERD15*, *LTL1*, *PSBTN*, and *MEE14* in the third

layer, and many of these genes undergo differentially AS events under cold stress.

The third layer is composed of 140 *COR*s mainly encoding structural proteins.

These genes are differentially regulated in expression and/or alternative splicing by

the first and/or second layer regulators. Among the genes, overexpression or knockout

of some genes (e.g., *COR47*, *TIL*, *SLD1*, *LOS2*, and *ADC1/2*) can affect plant freezing

tolerance[47-51]. Altogether, our findings suggest that circadian clock components are

actively regulated under cold stress to launch a hierarchy of gene regulatory network

to allow adaptation to cold stress in rosids.

## Discussion

When exposed to cold stress, plants undergo a series of molecular changes to protect

themselves from chilling and/or freezing damage[3,4]. By high-throughput methods,

previous studies provided insights into the reprogramming of gene regulation in

response to cold stress[7,8,29,30,32]. However, the studies were generally performed with

single plant species and focused on the best-known CBF-dependent pathway genes.

Here, we used three plant species (*A. thaliana*, *C. illinoinensis*, and *P. trichocarpa*)

form different rosid lineages which diverged during the early evolution of rosids

(~100 million years ago) and performed RNA-seq experiments with plants subjected

to cold stress to investigate conserved regulators and mechanisms underlying rosids in

response to cold stress. Over seven thousand DEGs were identified in each species

and show common responses of biological processes, such as cold stress process and

circadian rhythm (**Fig 1c**). Interestingly, the three plant species show different

cold-responsive speeds: an early in *A. thaliana* but a delayed extensive cold-response

in *C. illinoinensis* and *P. trichocarpa* (**Fig 1b, 2b**). The difference in cold-responsive

speeds maybe due to the different growth characteristics between annual herbaceous

plants (e.g., *A. thaliana*) and perennial woody plants (e.g., *C. illinoinensis* and *P.*

*trichocarpa*). Compared to woody plants, herbaceous plants are more sensitive to

temperature changes[52], which might cause an early extensive cold response.

In addition to transcriptional regulation, post-transcriptional regulation is a key

process in reprogramming gene expression under cold stress[7,30]. Like DEGs,

thousands of DASGs were identified in rosids under cold stress. Interestingly, the

number of DASGs in *A. thaliana* is significantly much less than the counterparts in *C. illinoinensis* and *P. trichocarpa* (**Fig 3a**). This difference may be due to shorter introns in *Arabidopsis* than in *C. illinoinensis* and *P. trichocarpa* (**Supplementary Fig 2**), and alternative splicing tends to occur around the exons interrupted by long introns[53,54]. In consistence with DEGs, DASGs show an enrichment of components involved in circadian rhythm (**Fig 3c**). Unexpectedly, enrichments of biological processes related to cold or abiotic stress are not observed for DASGs. This is probably because a limited knowledge of cold-induced DASGs and their genetic roles as compared to an extensive study of cold-induced DEGs. Regarding this, we provide a complete list of cold-induced DASGs in three rosid species (**Supplementary Table 3**) and hope this list will be a potential reference for future research of splicing regulation under cold stress.

Moreover, association analysis of transcriptional and post-transcriptional activities revealed 259 conserved DE-DASG orthogroups in rosids. From DE-DASG orthogroup genes, circadian rhythm is again significantly enriched (**Fig 5c**). Circadian clock system can allocate stress-responsive processes to specific times of the day, which provides a control of over daily energy expenditure and enables plants to grow and reproduce under temperature changes[19,55-57]. Here, we found that some conserved circadian clock components are reprogrammed at both transcriptional and post-transcriptional levels under cold stress in rosids (**Fig 5c**). Among them, MYB is the most prominent TF family including three orthogroups with five *Arabidopsis* genes (*CCA1/LHY*, *RVE7*, and *RVE4/8*), which likely contribute to the flexibility and controllability of the cold response network. Thus, by integrative omics of phylotranscriptomes and regulome maps (RNA-seq and ChIP/DAP-seq data), we identified their direct targets and constructed a conserved hierarchy of cold-responsive regulatory network launched by the five MYB TFs. The existence of many well-known cold-response genes, such as *CBFs*, *RBG7/8*, *ZAT6/10*, and *COR27*, in the network demonstrate a high confidence of the network but also imply an importance of circadian clock for plants in response to cold stress. Altogether, our

393    research shed light on conserved hierarchical cold-responsive network in rosids at

394    transcriptional and post-transcriptional levels via circadian clock components, which

395    provide insights into conserved core regulators and ancestral mechanisms underlying

396    cold-responsive gene regulatory network across rosids.

## Materials and Methods

**Identification and characterization of TFs and RBPs**

Proteome sequences and gene annotations of three rosid species (*A. thaliana*, *C.*

*illinoinensis*, and *P. trichocarpa*) were directly downloaded from the public

databases[58,59]. Their divergence times were extracted and assessed from the TimeTree

website[34]. TFs and RBPs in the species were obtained through the following strategies.

For TFs, we downloaded *A. thaliana* and *P. trichocarpa* TFs from plantTFDB 4.0 and

identified TFs in *C. illinoinensis* according to the method as described[60]

(**Supplementary Table 2**). For RBPs, we obtained *A. thaliana* RBPs, which were

experimentally confirmed RNA-binding assays[25,39]. By using *A. thaliana* RBPs as

queries, we employed BLASTP 2.5.0[61] to retrieve RBP homologs in *P. trichocarpa*

and *C. illinoinensis* (E-value < 1e–10, identity > 60%) (**Supplementary Table 4**).

To identify protein families of TFs and RBPs, we downloaded all Hidden Markov

Model (HMM) profiles (>19,000) of protein domains from the Pfam database[62] and

searched them against the proteome sequences of the three plant species by HMMER

3.1b2 with an E-value < 1e–5[63]. Accordingly, we obtained gene families for each TF

and RBP (**Supplementary Table 4**).

**Genome-wide analysis of cold-induced DEGs and DASEs**

To explore molecular changes of rosids under cold stress, we performed RNA-seq

experiments in three above-mentioned plant species with cold treatment (4□) at 0, 2,

24, and 168 hours (h). The experiment and analysis were performed as previously

described[64]. Gene expression under normal and cold treatment conditions was

represented by Trimmed Means of M value (TMM) calculated from the software

StringTie v2.0.3[65]. Subsequently, cold-induced DEGs were defined based on three

criteria: at least one sample with TMM greater than 5, more than two-fold change in

expression levels before and after cold treatment, and a statistical difference with $P$

value < 0.01. Additionally, five types of AS events, including intron retention (IR),

alternative 3' splice site (A3SS), alternative 5' splice site (A5SS), skipping exon (SE),

and mutually exclusive exons (MXE), were identified by rMATS v4.0.2[66] coupled

with CASH v2.2.1[26]. A false discovery rate (FDR) ≤ 0.05 was set as the criterion for

identifying cold-induced DASEs between normal and cold treatments.

**GO term enrichments**

The GO term annotations of *A. thaliana* genes were directly acquired from The

*Arabidopsis* Information Resource (TAIR) database[67]. According to method described

by the previous reports[35,68], the genes in *P. trichocarpa*, and *C. illinoinensis* were

annotated with the highest alignment score matches against *A. thaliana* proteins from

BLASTP searches (E-value < 1e–5)[61]. The GO term enrichment analysis was

performed using the OmicShare tools (www.omicshare.com/tools).

**Identification of DE-DASG orthogroups**

To explore whether DE-DASGs across the rosids are homologous, we classified the

up- and down-DE-DASGs of three rosids into different orthogroups using Orthofinder

v2.3.8[41] with the parameters (-M dendroblast; -S diamond; -A mafft; -I 1.5).

Subsequently, we obtained a total of 1,532 DE-DASG orthogroups and among which,

259 were identified to be present in all three species. In 259 DE-DASG orthogroups,

150 and 109 are respectively upregulated and downregulated under cold stress

(**Supplementary Table 6, 7**).

**Identification of CoCORs targeted by circadian clock components**

To determine CoCORs regulated by circadian clock components in rosids, we

developed the following pipeline. First, we obtained *A. thaliana* ChIP-seq (CCA1)

and DAP-seq (LHY, RVE7, RVE4, and RVE8) datasets[43,44] and performed

ChIP/DAP-seq analysis as described[64]. The genes with at least one binding peak of

the five TFs in the 1-Kb upstream of translation initiation site (TIS) were considered

as direct target genes of the corresponding TFs in *A. thaliana*. Second, we collected

*cis*-regulatory binding motifs of CCA1, *LHY*, *RVE7*, *RVE4*, and *RVE8*[43,44] and

searched them against the 1-Kb promoter sequences of *P. trichocarpa* and *C.*

*illinoinensis* genes using FIMO (E-value <1e–3)[69]. The genes with at least one

*cis*-regulatory binding site for the five TFs were identified as potentially targets of the

TFs in *P. trichocarpa* and *C. illinoinensis*. Subsequently, all coding genes in the three

species were classified into orthogroups according to their homology. The orthogroup

genes that have binding sites of the five TFs and are differentially regulated after cold

stress in all three rosids are defined as CoCORs. Finally, a total of 101/17, 167/67,

193/93, 38/10, and 110/34 CoCORs (**Fig 6a**, **Supplementary Table 9**) were identified

as to be up/down-regulated by CCA1, LHY, RVE7, RVE4 and RVE8, respectively.

**Statistical analysis**

The significance of enrichment analysis between gene groups was calculated by

Fisher Exact test through R programming.

# Acknowledgments

We thank the members from Dr. Wenwu Wu's laboratory in Zhejiang A&F University

for suggestions to improve the quality of the study. This work was supported by the

National Natural Science Foundation of China (grant number 31871233).

# Contributions

W.W., Z.J., and J.X. designed the research; L.G. and X.Z. performed the study; S.W.

did the RNA-seq experiments of the plants and Y.X. performed the RNA-seq analysis;

L.G., W.W. and Z.J. wrote and edited the manuscript.

**Data availability statement**

The RNA-seq datasets of the three plant species including *A. thaliana*, *C. illinoinensis*,

and *P. trichocarpa* under cold stress have been deposited to NCBI BioProject

(Accession: PRJNA767196) and CNGBdb (Accession: CNP0002243).

## Conflict of interests

There are no conflicts of interest.

## Supplementary information

Supplementary Fig 1. The GO term enrichments of conserved up- and down-DE-DASGs in *P. trichocarpa* and *C. illinoinensis*.

Supplementary Fig 2. Alternative splicing seems to occur in the genes with longer introns.

Supplementary Table 1. RNA-seq values of DEGs in rosids at four timepoints (0, 2, 24, and 168 h) of cold treatments (4□).

Supplementary Table 2. TF family genes and expression in rosids under cold stress.

Supplementary Table 3 A list of cold-induced DASGs and their AS types in rosids.

Supplementary Table 4. RBP family genes and expression in rosids under cold stress.

Supplementary Table 5. The detailed information of DE-DASGs.

Supplementary Table 6. The orthogroups of up-DE-DASGs in rosids.

Supplementary Table 7. The orthogroups of down-DE-DASGs in rosids.

Supplementary Table 8. Details of 27 up-DE-DASG TF orthogroups in rosids.

Supplementary Table 9. The predicted CoCORs targeted by clock components of CCA1, LHY, RVE4, RVE8, and RVE7 in rosids.

## References:

1. Wu, S., Han, B. & Jiao, Y. Genetic Contribution of Paleopolyploidy to Adaptive Evolution in Angiosperms. *Molecular Plant* **13**, 59-71 (2020).

2. Creux, N. & Harmer, S. Circadian Rhythms in Plants. *Cold Spring Harb Perspect Biol* **11**(2019).

3. Kidokoro, S., Shinozaki, K. & Yamaguchi-Shinozaki, K. Transcriptional regulatory network of plant cold-stress responses. *Trends in Plant Science* (2022).

4. Ding, Y., Shi, Y. & Yang, S. Advances and challenges in uncovering cold tolerance regulatory mechanisms in plants. *New Phytol* **222**, 1690-1704 (2019).

5. Zhu, J.K. Abiotic Stress Signaling and Responses in Plants. *Cell* **167**, 313-324 (2016).

6. Caldana, C. et al. High-density kinetic analysis of the metabolomic and transcriptomic response of *Arabidopsis* to eight environmental conditions. *Plant J* **67**, 869-84 (2011).

7. Calixto, C. et al. Rapid and Dynamic Alternative Splicing Impacts the *Arabidopsis* Cold Response Transcriptome. *Plant Cell* **30**, 1424-1444 (2018).

8. Zheng, Y. et al. Cold Response Transcriptome Analysis of the Alternative Splicing Events Induced by the Cold Stress in *D. catenatum*. *Int J Mol Sci* **23** (2022).

9. Hannah, M.A., Heyer, A.G. & Hincha, D.K. A global survey of gene regulation during cold acclimation in *Arabidopsis thaliana*. *PLoS Genet* **1**, e26 (2005).

10. Chinnusamy, V., Zhu, J. & Zhu, J. Cold stress regulation of gene expression in plants. *Trends in Plant Science* **12**, 444-451 (2007).

11. Stockinger, E.J., Gilmour, S.J. & Thomashow, M.F. Arabidopsis thaliana CBF1 encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a cis-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. *Proc Natl Acad Sci U S A* **94**, 1035-40 (1997).

12. Jaglo-Ottosen, K.R., Gilmour, S.J., Zarka, D.G., Schabenberger, O. & Thomashow, M.F. Arabidopsis CBF1 overexpression induces COR genes and enhances freezing tolerance. *Science* **280**, 104-6 (1998).

13. Zhao, C. et al. Mutational Evidence for the Critical Role of CBF Transcription Factors in Cold Acclimation in *Arabidopsis*. *Plant Physiol* **171**, 2744-59 (2016).

14. Jia, Y. et al. The cbfs triple mutants reveal the essential functions of CBFs in cold acclimation and allow the definition of CBF regulons in *Arabidopsis*. *New Phytol* **212**, 345-53 (2016).

15. Wang, Y. & Hua, J. A moderate decrease in temperature induces COR15a expression through the CBF signaling cascade and enhances freezing tolerance. *Plant J* **60**, 340-9 (2009).

16. Novillo, F., Medina, J. & Salinas, J. *Arabidopsis* CBF1 and CBF3 have a different function than CBF2 in cold acclimation and define different gene classes in the CBF regulon. *Proc Natl Acad Sci U S A* **104**, 21002-7 (2007).

17. Hsu, P.Y. & Harmer, S.L. Wheels within wheels: the plant circadian system. *Trends in Plant Science* **19**, 240-249 (2014).

18. Greenham, K. & McClung, C.R. Integrating circadian dynamics with physiological processes in plants. *Nature Reviews Genetics* **16**, 598-610 (2015).

19. Markham, K.K. & Greenham, K. Abiotic stress through time. *New Phytol* **231**, 40-46 (2021).

20. Oravec, M.W. & Greenham, K. The adaptive nature of the plant circadian clock in natural environments. *Plant Physiol* **190**, 968-980 (2022).

21. Harmer, S.L. The circadian system in higher plants. *Annu Rev Plant Biol* **60**, 357-77 (2009).

22.  Dong, M.A., Farré, E.M. & Thomashow, M.F. CIRCADIAN CLOCK-ASSOCIATED 1 and LATE
ELONGATED HYPOCOTYL regulate expression of the C-REPEAT BINDING FACTOR (CBF)
pathway in *Arabidopsis*. *Proceedings of the National Academy of Sciences* **108**, 7241-7246 (2011).
23.  Kidokoro, S. et al. Posttranslational regulation of multiple clock-related transcription factors
triggers cold-inducible gene expression in *Arabidopsis*. *Proceedings of the National Academy of*
*Sciences* **118** (2021).
24.  Wang, P. et al. COR27 and COR28 encode nighttime repressors integrating Arabidopsis circadian
clock and cold response. *J Integr Plant Biol* **59**, 78-85 (2017).
25.  Bach-Pages, M. et al. Discovering the RNA-Binding Proteome of Plant Leaves with an Improved
RNA Interactome Capture Method. *Biomolecules* **10** (2020).
26.  Wu, W. et al. CASH: a constructing comprehensive splice site method for detecting alternative
splicing events. *Brief Bioinform* **19**, 905-917 (2018).
27.  Fu, X.D. & Ares, M.J. Context-dependent control of alternative splicing by RNA-binding proteins.
*Nat Rev Genet* **15**, 689-701 (2014).
28.  Lee, Y. & Rio, D.C. Mechanisms and Regulation of Alternative Pre-mRNA Splicing. *Annu Rev*
*Biochem* **84**, 291-323 (2015).
29.  Liu, Z. et al. Global profiling of alternative splicing landscape responsive to drought, heat and their
combination in wheat (*Triticum aestivum* L.). *Plant Biotechnol J* **16**, 714-726 (2018).
30.  Zhao, X. et al. The SR Splicing Factors: Providing Perspectives on Their Evolution, Expression,
Alternative Splicing, and Function in *Populus trichocarpa*. *International Journal of Molecular*
*Sciences* **22**, 11369 (2021).
31.  Seo, P.J. et al. A self-regulatory circuit of CIRCADIAN CLOCK-ASSOCIATED1 underlies the
circadian clock regulation of temperature responses in *Arabidopsis*. *Plant Cell* **24**, 2427-42 (2012).
32.  Wang, Y. et al. Global Survey of the Full-Length Cabbage Transcriptome (*Brassica oleracea* Var.
capitata L.) Reveals Key Alternative Splicing Events Involved in Growth and Disease Response.
*International Journal of Molecular Sciences* **22**, 10443 (2021).
33.  Chase, M.W. & Reveal, J.L. A phylogenetic classification of the land plants to accompany APG III.
*Botanical Journal of the Linnean Society* **161** (2009).
34.  Kumar, S., Stecher, G., Suleski, M. & Hedges, S.B. TimeTree: A Resource for Timelines, Timetrees,
and Divergence Times. *Mol Biol Evol* **34**, 1812-1819 (2017).
35.  Guo, L. et al. Convergent evolution of AP2/ERF III and IX subfamilies through recurrent
polyploidization and tandem duplication during eudicot adaptation to paleoenvironmental changes.
*Plant Commun*, 100420 (2022).
36.  Khan, S.A., Li, M.Z., Wang, S.M. & Yin, H.J. Revisiting the Role of Plant Transcription Factors in
the Battle against Abiotic Stress. *Int J Mol Sci* **19** (2018).
37.  Xie, Z., Nolan, T.M., Jiang, H. & Yin, Y. AP2/ERF Transcription Factor Regulatory Networks in
Hormone and Abiotic Stress Responses in *Arabidopsis*. *Front Plant Sci* **10**, 228 (2019).
38.  Dubos, C. et al. MYB transcription factors in *Arabidopsis*. *Trends Plant Sci* **15**, 573-81 (2010).
39.  Marondedze, C., Thomas, L., Serrano, N.L., Lilley, K.S. & Gehring, C. The RNA-binding protein
repertoire of *Arabidopsis thaliana*. *Sci Rep* **6**, 29766 (2016).
40.  Koonin, E.V. Orthologs, paralogs, and evolutionary genomics. *Annu Rev Genet* **39**, 309-38 (2005).
41.  Emms, D.M. & Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative genomics.
*Genome Biol* **20**, 238 (2019).
42.  James, A.B. et al. Alternative splicing mediates responses of the *Arabidopsis* circadian clock to

temperature changes. *Plant Cell* **24**, 961-81 (2012).

43.  Kamioka, M. et al. Direct Repression of Evening Genes by CIRCADIAN CLOCK-ASSOCIATED1
in the *Arabidopsis* Circadian Clock. *Plant Cell* **28**, 696-711 (2016).

44.  O'Malley, R.C. et al. Cistrome and Epicistrome Features Shape the Regulatory DNA Landscape.
*Cell* **165**, 1280-1292 (2016).

45.  Jiang, B. et al. Cold-Induced CBF-PIF3 Interaction Enhances Freezing Tolerance by Stabilizing the
phyB Thermosensor in *Arabidopsis*. *Mol Plant* **13**, 894-906 (2020).

46.  Meyer, K. et al. Adaptation of iCLIP to plants determines the binding landscape of the
clock-regulated RNA-binding protein AtGRP7. *Genome Biology* **18** (2017).

47.  Lee, H. et al. LOS2, a genetic locus required for cold-responsive gene transcription encodes a
bi-functional enolase. *EMBO J* **21**, 2692-702 (2002).

48.  Cao, S., Ye, M. & Jiang, S. Involvement of GIGANTEA gene in the regulation of the cold stress
response in *Arabidopsis*. *Plant Cell Rep* **24**, 683-90 (2005).

49.  Charron, J.B., Ouellet, F., Houde, M. & Sarhan, F. The plant Apolipoprotein D ortholog protects
*Arabidopsis* against oxidative stress. *BMC Plant Biol* **8**, 86 (2008).

50.  Chen, M., Markham, J.E. & Cahoon, E.B. Sphingolipid Delta8 unsaturation is important for
glucosylceramide biosynthesis and low-temperature performance in *Arabidopsis*. *Plant J* **69**,
769-81 (2012).

51.  Cuevas, J.C. et al. Putrescine is involved in *Arabidopsis* freezing tolerance and cold acclimation by
regulating abscisic acid levels in response to low temperature. *Plant Physiol* **148**, 1094-105 (2008).

52.  Cornelius, C., Estrella, N., Franz, H. & Menzel, A. Linking altitudinal gradients and temperature
responses of plant phenology in the Bavarian Alps. *Plant Biol (Stuttg)* **15 Suppl 1**, 57-69 (2013).

53.  Roy, M., Kim, N., Xing, Y. & Lee, C. The effect of intron length on exon creation ratios during the
evolution of mammalian genomes. *RNA* **14**, 2261-73 (2008).

54.  Fox-Walsh, K.L. et al. The architecture of pre-mRNAs affects mechanisms of splice-site pairing.
*Proc Natl Acad Sci U S A* **102**, 16176-81 (2005).

55.  Espinoza, C., Bieniawska, Z., Hincha, D.K. & Hannah, M.A. Interactions between the circadian
clock and cold-response in *Arabidopsis*. *Plant Signal Behav* **3**, 593-4 (2008).

56.  Espinoza, C. et al. Interaction with diurnal and circadian regulation results in dynamic metabolic
and transcriptional changes during cold acclimation in *Arabidopsis*. *PLoS One* **5**, e14101 (2010).

57.  Bieniawska, Z. et al. Disruption of the Arabidopsis circadian clock is responsible for extensive
variation in the cold-responsive transcriptome. *Plant Physiol* **147**, 263-79 (2008).

58.  Aken, B.L. et al. The Ensembl gene annotation system. *Database (Oxford)* **2016**(2016).

59.  Goodstein, D.M. et al. Phytozome: a comparative platform for green plant genomics. *Nucleic Acids*
*Res* **40**, D1178-86 (2012).

60.  Jin, J. et al. PlantTFDB 4.0: toward a central hub for transcription factors and regulatory interactions
in plants. *Nucleic Acids Res* **45**, D1040-D1045 (2017).

61.  Altschul, S.F., Gish, W., Miller, W., Myers, E.W. & Lipman, D.J. Basic local alignment search tool.
*J Mol Biol* **215**, 403-10 (1990).

62.  Finn, R.D. et al. The Pfam protein families database: towards a more sustainable future. *Nucleic*
*Acids Res* **44**, D279-85 (2016).

63.  Eddy, S.R. Profile hidden Markov models. *Bioinformatics* **14**, 755-63 (1998).

64.  Nie, Y. et al. Innovations and Stepwise Evolution of CBFs/DREB1s and Their Regulatory Networks
in Angiosperms. *J Integr Plant Biol* (2022).

65. Pertea, M. et al. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads.
*Nat Biotechnol* **33**, 290-5 (2015).
66. Shen, S. et al. rMATS: robust and flexible detection of differential alternative splicing from replicate
RNA-Seq data. *Proc Natl Acad Sci U S A* **111**, E5593-601 (2014).
67. Rhee, S.Y. et al. The Arabidopsis Information Resource (TAIR): a model organism database
providing a centralized, curated gateway to Arabidopsis biology, research materials and community.
*Nucleic Acids Res* **31**, 224-8 (2003).
68. Salojärvi, J. et al. Genome sequencing and population genomic analyses provide insights into the
adaptive landscape of silver birch. *Nature Genetics* **49**, 904-912 (2017).
69. Grant, C.E., Bailey, T.L. & Noble, W.S. FIMO: scanning for occurrences of a given motif.
*Bioinformatics* **27**, 1017-8 (2011).
70. Song, Y. et al. The direct targets of CBFs: In cold stress response and beyond. *J Integr Plant Biol* **63**,
1874-1887 (2021).

639
640

## Figure legends

**Fig 1. Identification and enrichment analysis of cold-induced differentially expressed genes (DEGs).**

(**a**) Overview of DEGs under cold stress. The number in parentheses below each species indicates the number of all coding genes in the genome. The evolutionary relationship and divergence times of the species were obtained from the website TimeTree[34]. Bar heights indicate gene numbers of upregulated DEGs (in red) and downregulated DEGs (in green). (**b**) Expression profiles of DEGs before and after cold treatments (4 □). The expression values of DEGs at four different timepoints (0, 2, 24, and 168 h) of cold treatments were normalized by row scale. (**c-d**) GO term enrichments of upregulated DEGs and downregulated DEGs. The top 20 significant GO terms were shown in a conserved manner with common cold-responsive biological processes in three plant species.

**Fig 2. Cold-responsive transcription factors (TFs) in conserved TF families.**

(**a**) Upregulated and downregulated TFs under cold stress. Bar percentage indicates the proportion of upregulated, downregulated, and no significant expression changed TFs to all TFs. The number in parentheses after the species indicates the number of TFs in the genome. (**b**) Expression profiles of up- and downregulated TFs under cold stress. The expression values of the TFs at four different timepoints (0, 2, 24, and 168 h) of cold treatments were normalized by row scale. (**c-d**) The numbers of up- and downregulated TFs in TF families. The top 15 up- and downregulated TF families were selected for visualization.

**Fig 3. Identification and enrichment analysis of cold-induced differentially alternative splicing genes (DASGs).**

(**a**) Overview of DASGs and differentially alternative splicing events (DASEs) under cold stress. The number in circle indicates the number of cold-induced DASGs or DASEs in the corresponding species. (**b**) Proportions of five AS types of cold-induced DASEs. IR: intron retention; SE: skipping exon; A3SS: alternative 3' splicing site; A5SS: alternative 5' splicing site; MXE: mutually exclusive exons. (**c**) GO term enrichments of DASGs. The top 20 significant GO terms were shown in a conserved manner with common cold-responsive biological processes in three species.

**Fig 4. Cold-responsive RNA binding proteins (RBPs) in conserved RBP families.**

(**a**) Upregulated and downregulated RBPs under cold stress. Bar percentage indicates the proportion of upregulated, downregulated, and no significant expression changed RBPs to all RBPs. The number in parentheses after the species indicates the number of RBPs in the genome. (**b**) Expression profiles of up- and downregulated RBPs under cold stress. The expression values of the RBPs at four different timepoints (0, 2, 24, and 168 h) of cold treatments were normalized by row scale. (**c-d**) The numbers of up- and downregulated RBPs in RBP families. The top 15 up- and downregulated RBP families were selected for visualization.

**Fig 5. Association of cold-induced DEGs and DASGs and identification of conserved DE-DASG orthogroups in rosids.**

(**a**) Overlapping and unique genes between DEGs and DASGs in three species. The overlapping significance was assessed by Fisher's Exact Test. (**b-e**) DE-DASG orthogroup identification and enrichment analysis of DE-DASG orthogroup genes in three species. Venn diagram shows overlapping and unique orthogroups of up- (**b**) and down-DE-DASGs (**d**) in three species. For each species, the number in purple after the slash indicates the number of up- or down-regulated DE-DASG genes, while the number in blue before the slash indicates the number of the corresponding DE-DASG orthogroups. In the overlapping 150 up- and 109 down-DE-DASG orthogroups, we obtained 217 (**c**) and 162 (**e**) *C. illinoinensis* genes representatively for GO term analysis. The same analysis was also performed in *A. thaliana* and *P. trichocarpa* (Supplementary Fig 1). The top 15 significant enriched terms were selected to be shown.

**Fig 6. A hierarchical cold-responsive regulatory network launched by circadian clock components in rosids.**

(**a**) Numbers of up- and downregulated CoCORs targeted by CCA1, LHY, RVE4, RVE7, and RVE8. These CoCORs were predicted by a stringent pipeline (see Materials and methods). (**b**) GO enrichment analysis of CoCORs. The top 10 significant GO terms were selected to be shown. (**c**) Overlapping and unique CoCORs regulated by the five circadian-related TFs. (**d**) A hierarchical network launched by the five circadian-related TFs. The network is arranged in three layers (TFs, RBPs, and Structural proteins). To improve the robustness of the network, we also integrated regulations of CBFs, ZAT10, HB6, HB-7, NAC053, WRKY7, MYB73, TCP7, and GRP7 in relation to CoCORs. Dotted lines indicate that the two linked genes are

712 co-regulated under cold stress and their potential regulation remains to be determined.
713 Solid arrow lines indicate that additional binding evidence analyzed from
714 ChIP/DAP-seq and iCLIP-seq experiments supports the linked genes.
715
716 **Tables**

717 **Table 1. Cold-responsive circadian-related TFs in rosids.**

| Family(orthogroup) | Cold regulation *Ath*/*Cil*/*Ptr* | *Ath* (AS-type) | *Cil* (AS-type) | *Ptr* (AS-type) |
|---|---|---|---|---|
| MYB (OG0000072) | UP/UP/UP | *CCA1* (A5SS) *LHY* (A3SS,A5SS,RI,SE) | *CIL1205S0016* (A3SS, A5SS, MXE, RI, SE) | *Potri.014G106800* (A3SS, A5SS, MXE, IR, SE) |
| MYB (OG0000110) | UP/UP/UP | *RVE4 (A3SS, SE)* *RVE8 (A3SS, IR)* | *CIL0982S0042* (A5SS) | *CIL0982S0042* (A5SS) |
| MYB (OG0000043) | UP/UP/UP | *RVE7* (SE) | *CIL1030S0151* (IR) CIL1245S0064 (IR) | *Potri.004G074300 (IR)* *Potri.012G038300 (IR)* |
| PRR (OG0000018) | UP/UP/UP | *PRR9* (A5SS) | *CIL0897S0115* (IR) *CIL1083S0010* (IR) *CIL1194S0023* (IR) | *Potri.002G179800 (A3SS, IR, SE)* *Potri.014G106000 (A3SS, IR, SE)* |
| BBX (OG0000101) | UP/UP/UP | *COL9* (IR) | *CIL1155S0003* (IR) | *Potri.002G214500 (IR)* *Potri.014G170600 (IR)* |
| BBX (OG0000088) | UP/UP/UP | *BBX19* (IR, SE) | *CIL0978S0047* (IR) | *Potri.001G061800* (A3SS) |
| Cupin (OG0000219) | UP/UP/UP | JMJD5 (A3SS, IR) | *CIL1465S0014* (IR) | *Potri.001G016200* (IR) |

718 Note: The short names (*Ath*, *Cil*, and *Ptr*) indicate the species (*A. thaliana*, *C.*
719 *illinoinensis*, and *P. trichocarpa*)

**Fig 1. Identification and enrichment analysis of cold-induced differentially expressed genes (DEGs).**
(a) Overview of DEGs under cold stress. The number in parentheses below each species indicates the number of all coding genes in the genome. The evolutionary relationship and divergence times of the species were obtained from the website TimeTree (Kumar et al., 2017). Bar heights indicate gene numbers of upregulated DEGs (in red) and downregulated DEGs (in green). (b) Expression profiles of DEGs before and after cold treatments (4°C). The expression values of DEGs at four different timepoints (0, 2, 24, and 168 h) of cold treatments were normalized by row scale. (c-d) GO term enrichments of upregulated DEGs and downregulated DEGs. The top 20 significant GO terms were shown in a conserved manner with common cold-responsive biological processes in three species.

**Fig 2. Cold-responsive TFs in conserved TF families.**

(a) Upregulated and downregulated TFs under cold stress. Bar percentage indicates the proportion of upregulated, downregulated, and no significant expression changed TFs to all TFs. The number in parentheses after the species indicates the number of TFs in the genome. (b) Expression profiles of up- and downregulated TFs under cold stress. The expression values of the TFs at four different timepoints (0, 2, 24, and 168 h) of cold treatments were normalized by row scale. (c-d) The numbers of up- and downregulated TFs in TF families. The top 15 up- and downregulated TF families were selected for visualization.

**Fig 3. Identification and enrichment analysis of cold-induced alternative splicing genes (DASGs).**
(a) Overview of DASGs and DASEs under cold stress. The number in circle indicates the number of cold-induced DASGs or DASEs in the corresponding species. (b) Proportions of five AS types of cold-induced DASEs. IR: intron retention; SE: skipping exon; A3SS: alternative 3' splicing site; A5SS: alternative 5' splicing site; MXE: mutually exclusive exons. (c) GO term enrichments of DASGs. The top 20 significant GO terms were shown in a conserved manner with common cold-responsive biological processes in three species.

**Fig 4. Cold-responsive RNA binding proteins (RBPs) in conserved RBP families.**
(a) Upregulated and downregulated RBPs under cold stress. Bar percentage indicates the proportion of upregulated, downregulated, and no significant expression changed RBPs to all RBPs. The number in parentheses after the species indicates the number of RBPs in the genome. (b) Expression profiles of up- and downregulated RBPs under cold stress. The expression values of the RBPs at four different timepoints (0, 2, 24, and 168 h) of cold treatments were normalized by row scale. (c-d) The numbers of up- and downregulated RBPs in RBP families. The top 15 up- and downregulated RBP families were selected for visualization.

**Fig 5. Association of cold-induced DEGs and DASGs and identification of conserved DE-DASG orthogroups in rosids.**

(a) Overlapping and unique genes between DEGs and DASGs in three species. The overlapping significance was assessed by Fisher's Exact Test. (b-e) DE-DASG orthogroup identification and enrichment analysis of DE-DASG orthogroup genes in three species. Venn diagram shows overlapping and unique orthogroups of up- (b) and down-DE-DASGs (d) in three species. For each species, the number in purple after the slash indicates the number of up- or down-regulated DE-DASG genes, while the number in blue before the slash indicates the number of the corresponding DE-DASG orthogroups. In the overlapping 150 up- and 109 down-DE-DASG orthogroups, we obtained 217 (c) and 162 (e) *C. illinoinensis* genes representatively for GO term analysis. The same analysis was also performed in *A. thaliana* and *P. trichocarpa* (Supplementary Fig 1). The top 15 significant enriched terms were select-ed to be shown.

**Fig 6. A hierarchical cold-responsive regulatory network launched by circadian clock components in rosids.**
(a) Numbers of up- and downregulated CoCORs targeted by CCA1, LHY, RVE4, RVE7, and RVE8. These CoCORs were predicted by a stringent pipeline (see Materials and methods). (b) GO enrichment analysis of CoCORs. The top 10 significant GO terms were selected to be shown. (c) Overlapping and unique CoCORs regulated by the five circadian-related TFs. (d) A hierarchical network launched by the five circadian-related TFs. The network is arranged in three layers (TFs, RBPs, and Structural proteins). To improve the robustness of the network, we also integrated regulations of CBFs, ZAT10, HB6, HB-7, NAC053, WRKY7, MYB73, TCP7, and GRP7 in relation to CoCORs. Dotted lines indicate that the two linked genes are co-regulated under cold stress and their potential regulation remains to be determined. Solid arrow lines indicate that additional binding evidence analyzed from ChIP/DAP-seq and iCLIP-seq experiments supports the linked genes.