Title: Environmental filtering of life-history trait diversity in urban populations of *Arabidopsis*

thaliana

Authors: Gregor Schmitz[1], Anja Linstädter[2] , Anke S. K. Frank[1,3], Hannes Dittberner[1], Jessica Thome[1],

Andrea Schrader[1,4], Karl-Heinz Linne von Berg[1], Andrea Fulgione[5], George Coupland[5], Juliette de

Meaux[1],*

[1] Institute for Plant Sciences, Biocenter, Zülpicher Str. 47b, 50674 Cologne, Germany,

[2] Institute of Biochemistry and Biology, University of Potsdam, Maulbeeralle 1, 14469 Potsdam,

Germany,

[3] School of Life and Environmental Sciences, The University of Sydney, Sydney, Australia

[4] RWTH Aachen University, Institute for Biology I, Molecular Botany, Worringer Weg 3, 52062

Aachen, Germany

[5] Max Planck Institute of Plant Breeding Research, Cologne, Germany

*Corresponding author.

Abstract (max 350)

1. The challenges to which plants are exposed in urban environments represent, in miniature,

the challenges plants face as a result of global environmental change. Hence, urban habitats

provide a unique opportunity to assess whether processes of local adaptation are taking

place despite the short temporal and geographical scales that characterize the

Anthropocene.

2. We quantified the ecological diversity of urban habitats hosting *A. thaliana* populations.

Using plant community indicators, we show that these patches differ in their levels of soil

nutrient content and disturbance. Accordingly, plants in each patch displayed a range of

flowering time, size and fitness.

3. Using a deep sampling approach coupled with reduced genome-sequencing, we demonstrate

that most individuals can be assigned to a limited set of clonal lineages; the genetic diversity

of these lineages is a sample of the diversity observed in western European populations of

the species, indicating that established urban populations originate from a broad regional

pool of lineages.

4. We assessed the genetic and phenotypic diversity of these lineages in a set of common

garden experiments. We report marked genetic differences in life-history traits, including

time of primary and secondary dormancy as well as of flowering. These genetic differences in

life-history traits are not randomly distributed but sorted out by ecological differences

among sites of origin.

5. **Synthesis:** Our study shows that the genetically diverse phenology of a regional *A. thaliana*
gene pool is not randomly distributed but filtered by heterogeneity in the urban
environment. To out knowledge, this report is the first to show a pattern indicative of
environmental filtering enhancing local genetic adaptation within urban environments. We
conclude that environmental filtering helps maintain functional diversity within species.

Keywords: urban ecology, local adaptation, environmental filtering, life-history trait
adaptation, ruderal species

1. Introduction

Urban environments provide untapped opportunities to link ecological and phenotypic diversity over the short temporal and spatial scales that define the Anthropocene (Johnson & Munshi-South, 2017; Szulkin et al., 2020). Understanding how plant functional variation is optimized within local urban environments can help understand the potential of species to cope with upcoming global changes.

Compared to that in neighboring rural areas, the climate in urban environments is generally characterized by increased temperature and reduced moisture, and thus a higher drought likelihood (Lambrecht et al. 2016). In addition, small-scale differences within urban microclimates are observable, depending on urban geometry and construction materials (Chatzidimitriou & Yannas, 2015). Urban microhabitats differ in edaphic characteristics: soils vary considerably in their developmental stage, depth, compaction, and surface transformation (Foti, 2017; Johnson & Munshi-South, 2017; Tresch et al., 2018). Shallow, sealed soils with a low capacity for water storage and low nutrient content typify the so-called urban grey spaces associated with built urban elements, such as sidewalks. They occur in close vicinity to nutrient-rich urban green spaces, often with eutrophic or even contaminated soils, which results in huge variation in soil-mediated plant resources over small spatial scales (Bonthoux et al., 2019; Gilbert, 2012; Vega & Küffer, 2021). Cities also host numerous ruderal sites, which are exposed to frequent and often severe disturbances, fragmentation, and high turnover rates among populations (Dubois & Cheptou, 2017; Guo et al., 2018). In addition, human-driven disturbances, such as weeding or mowing, may shorten the time windows for growth and reproduction, and thus modulate further the disturbance patterns.

Major life-history decisions, such as the timing of germination and the timing of flowering, shape the timing of plant phenological transitions and are critical for resource acquisition and reproductive success. They have been shown to contribute to the persistence of populations in heterogeneous urban settings (Liu et al., 2021; Nord & Lynch, 2009). Plant phenology is plastic. It can be adjusted locally in response to the environment because the life-history traits underlying it are modulated both by nutrient resources and by environmental cues, such as temperature and photoperiod (Vidal et al., 2014). For example, flowering is accelerated by winter temperature and resource stress and by high temperatures in spring, and delayed by the shortening days of the fall (Andrés & Coupland, 2012; Li et al., 2019). Plant phenology thus shifts in response to environmental cues that are modified by climate change (Chmielewski & Rötzer, 2001; Piao et al., 2019; Primack & Miller-Rushing, 2012).

The genetic composition of the population can also be tuned by local selective pressures acting on the genetic basis of phenological variation. Adaptation to local environments has been observed experimentally in many organisms (Kawecki & Ebert 2004; Leimu & Fischer 2008). Clearly,

adaptation via new genetic mutations is unlikely within urban settings because urban populations are
small, fragmented, and prone to frequent extinction(Orr, 2005). However, if the pool of potential
migrant genotypes is sufficiently diverse, then individual genotypes may establish or not depending
on their genetic makeup, which results in local adaptation by environmental filtering. Indeed,
patterns of local adaptation detected over small distances suggest that this kind of environmental
filtering could happen in urban habitats (Halbritter et al., 2018; Lenssen et al., 2004; Robionek-
Selosse et al., 2021).

In recent years, several studies have described phenotypic differences and genetic differentiation
between individuals from urban and neighboring rural sites, consistently indicating that populations
adapt rapidly to cities. These studies documented selection for novel optima in life-history traits,
such as a late flowering time and decreased dispersal in holy hawksbeard (*Crepis sancta);* an early
flowering time in common ragweed (*Ambrosia artemisiifolia*); and decreased cyanogenesis in white
clover (*Trifolium repens*) (Cheptou et al., 2008; Gorton et al., 2018; Lambrecht et al., 2016;
Santangelo et al., 2022; Thompson et al., 2016). Altogether, however, the amount of genetic
variation in life-history traits,   the distribution of this variation within the heterogeneous
environments of cities, and its contribution to local adaptation in these environments remain
unknown (Johnson & Munshi-South, 2017).

Urban environments in regions with temperate climates often host populations of short-lived
herbaceous species, including the Brassicaceae species *Arabidopsis thaliana* (Bomblies et al., 2010;
Herben et al., 2016; Huang et al., 2018). This species -- the initial model of plant molecular biology --
has become a powerful system for answering fundamental questions of ecology, evolution, and
global environmental change (Exposito-Alonso et al., 2018; Fournier-Level et al., 2011; Takou et al.,
2019; Vasseur et al., 2018). *A. thaliana*, which is native to large areas of Eurasia and Africa, has
recently colonized many other parts of the world (Durvasula et al., 2017; Lee et al., 2017). The deep
sampling of natural populations across various regions of its area of distribution has shown how
environmental heterogeneity shapes the distribution of genetic diversity in *A. thaliana* (Castilla et al.,
2020; DeLeo et al., 2020; Dubin et al., 2015, p. 20; Luo et al., 2015)*.* Life-history trait adaptation
appears to happen in concert across regional environments (reviewed in Takou et al. 2019). This
includes adaptation in the timing of distinct phenological stages such as germination (Kronholm et
al., 2012), or the onset of flowering (Ågren et al., 2017; Akiyama & Ågren, 2018; Le Corre, 2005), but
also the tuning of other life-history traits such as the rate of rosette growth or leaf economic
spectrum (Debieu et al., 2013; Sartori et al., 2019; Wieters et al., 2021).

In the present study, we determined the relative importance of phenotypic plasticity and genetic
variation in shaping the phenology and fitness of eight spontaneously occurring urban populations of

A. thaliana. We asked i) Are *A. thaliana* urban populations occupying ecologically diverse habitats? ii) Are *A. thaliana* populations genetically diverse? iii) Is this genetic diversity randomly distributed across habitats? By combining plant-community and trait-based ecological measurements with analyses of genetic diversity at genomic and phenological levels, we show that genotypes growing in urban habitats exposed to high resource limitations and increased disturbance levels tend to have traits tailored to local conditions.

2. Materials and Methods

Study area

Our study focused on urban *A. thaliana* populations in the city of Cologne, situated in midwestern Germany (about 50.9°N, 7.0°E). Cologne is Germany's fourth most populous city, with more than one million inhabitants (Ansmann et al., 2021). One of Germany's warmest cities, it has a temperate oceanic climate with a mean annual temperature of 11.7 °C during the day and 6.3 °C at night, and a mean annual precipitation of 840 mm.

Field surveys, phenology monitoring, and functional trait measurements

We surveyed eight *A. thaliana* populations within a 4.3 x 3.7 km area in southwestern Cologne. Study populations were selected based on their accessibility and on their isolation from other populations, with all habitat patches located at least 0.6 km from one another. Urban habitat patches where viable *A. thaliana* populations were observable included green spaces such as disturbed lawns; various habitats within gray spaces, such as sidewalks, wall tops, and bases; and vegetated roadsides (Figure 1a; Figure S1; Table S1). Depending on habitat characteristics, patch size ranged from 4 m^2 for populations on wall tops to > 100 m² for populations in vegetated roadsides, and population size ranged from 15 to 1,000 individuals. All populations were studied throughout their growth period in 2017 (which extended between early March and the end of June), and revisited yearly during five subsequent growth periods (2018-2022) to assess population persistence. No field permissions were necessary for the collection of the plant samples in this study.

We recorded phenological stages and fitness-related plant functional traits of *A. thaliana* individuals distributed over the eight habitat patches. At each patch, we established 10 to 14 observation quadrats (10 x 10 cm), each at least 50-60 cm from one another. We marked from 1 to 3 *A. thaliana* individuals per quadrat (15-24 per habitat patch; 174 in total) and visited them twice a week during the study period to score their development. For all marked individuals, we recorded the day of bolting (i.e., when the inflorescence became visible) and, subsequently, the number of open flowers and the number of closed fruits. Rosette diameter and the number of rosette leaves were measured at the time of bolting. We used these observation data to derive five phenological stages and five fitness-related plant functional traits ("fitness proxies"; see Table S2). The selection of plant

145   functional traits follows recommendations for the assessment of plant fitness (Gibson, 2014), while

146   phenological stages were selected to represent life-history stages of the entire growth period, as

147   recommended for herbaceous species (Huang et al., 2018; Nordt et al., 2021). Germination was

148   generally not possible to determine *in situ*, except in populations growing on wall tops (SGY, KAD,

149   HLU).

Assessment of environmental conditions of habitat patches

Abiotic environmental conditions and characteristics of disturbance regime were recorded at the

level of habitat patches, making use of the indicator function of plant communities (Kollmann &

Fischer, 2003). We used species' Ellenberg indicator values, EIVs (Ellenberg & Leuschner, 2010), as a

proxy for abiotic conditions. EIVs, which express the optimal positions of Central European plant

species along gradients of abiotic factors, provide information about species' niche and habitat

requirements (Diekmann 2003), in both rural and urban habitats (Fanelli et al. 2006). EIVs are

expressed on an ordinal scale ranging from 1 to 9 and refer to light regime (L), temperature (T), and

continentality of climate (K). Edaphic conditions are captured as soil moisture (F), pH (R), and

nutrient availability (N). Likewise, species' disturbance indicator values (DIVs) (Herben et al., 2016),

were used to characterize the disturbance regimes of urban habitat patches. To this end, we

extracted species-level indicators for disturbance severity (DS), disturbance frequency (DF), and

disturbance effects on vegetation structure (DV) from the data compiled in Herben et al., 2016. DIVs

are based on vegetation-plot records of the Czech flora, with a species' DS defined as the mean

disturbance severity of all vegetation classes weighted by its occurrence frequencies in these classes;

DF calculated as the mean of logarithmic disturbance frequency of all vegetation classes weighted by

a species' occurrence frequencies in these classes; and DV calculated from vegetation structural

parameters by summing covers and community-weighted means of all species' growth height per

vegetation class. We determined the composition of plant communities in autumn (late August and

early September) 2017 by recording the presence of all vascular plant species in the herbaceous layer

of the habitat patches. Based on species' specific indicator values (see Table S4) and excluding *A.*

thaliana, weighted average EIVs and DIVs were calculated for the plant community of each habitat

patch.

Assessment of genomic variation

At the end of the 2017 growth period, we harvested seeds from 5 to 17 individuals per population

and used single seeds from each individual to produce progenies. For four of the eight habitat

patches (BER, BIS, KAD, and MIL), we also included seeds from individuals sampled in the year prior

to field data collection, referring to these as population samples (Table S3). In order to broaden our

view of the genomic diversity of *A. thaliana* in Cologne, we also produced progenies from seeds

collected at 30 additional habitat patches distributed throughout the sampling area but not included
in the *in situ* monitoring (hereafter called scattered samples). Field collected seeds were grown in a
randomized setting, under conditions simulating winter, i.e., in growth chambers with 10 h light at
18 °C / 14 h dark at 16 °C, 60 % humidity, for 6 weeks; followed by 8 h light/16 h dark at 4°C, 80 %
humidity, for 6 weeks; and ultimately ripened in the greenhouse in spring-like conditions with 16 h
light, at about 20-24 °C / 8 h dark at about 18-20 °C until harvest. Seeds for progeny were harvested
using Aracons (Betatech, Gent, Belgium).

Three leaves per plant were sampled for DNA extraction, deep frozen, and homogenized using a
Precellys Evolution homogenizer (Thermo Fisher Scientific, Waltham, MA, USA). DNA was isolated
using the NucleoSpin® Plant II kit (Macherey-Nagel, Düren, Germany). Genomic DNA was prepared
for RAD-Seq as described by (Dittberner et al., 2019) using 10 pools with 20 individually barcoded
plant samples each, and sequencing was performed at the Max Planck Institute of Plant Breeding
Research. The bioinformatics pipelines used for read mapping, and identification of each of the 12
genotypes present in the studied habitat patches, population structure and admixture are described
in the supplementary information.

Common garden experiments:

For each identified genotype, two individuals were randomly selected and amplified in the
greenhouse. Since they were raised in the same maternal environment, variance among genotypes
can be assigned to genetic effects and variance among replicate lines of each genotype allows testing
whether all maternal effects were effectively removed when growing parents in common garden
conditions. We then took eight individuals from each of these two lines and grew them in growth
chambers (Johnson Controls, Milwaukee, WI, USA) in three trials under different growth conditions.
Growth conditions were chosen to identify genetic variation in the major pathways regulating
flowering time: constant long days (16 h light at 20°C/8h dark at 18°C); constant short days (8 h light
at 18°C day/16 dark at 16°C night); and short days/vernalization/long days (8 h day light at 18°C /16 h
dark at 16°C for 5 weeks; 8h day light/16h dark at 4 °C for 4 weeks, 16 h day light at 20°C/8h dark at
18°C until ripening). The *A. thaliana* genotypes Col-0 was included as a reference in all experiments.
Because we did not detect differences between replicated lines, we considered the number of
replicates to be 2x8= 16 replicates/genotype. Replicates were sown in individual 6-cm-diameter pots
that were watered regularly by flooding the trays. Flowering time was determined as the time span
between sowing and the first flower to open petals.

Ripe seeds from the short-day/vernalization/long-day experiment were harvested and stored at
room temperature for the germination assay. Germination rates were determined after three
months to quantify primary dormancy (Baskin & Baskin, 2004). Seeds were sown on wet filter paper

and incubated in closed microtiter plates. Before incubation under long-day conditions (16 h light at

20°C/8h dark at 18°C), seeds were incubated in the dark either at 4 °C for 7 days to quantify their

capacity to germinate when dormancy is released, or at -21 °C for 4 d and at 35 °C for 4 d to quantify

variation in secondary dormancy. Germination was scored after 10 days.

To quantify genetic variation under outdoor field conditions, we conducted a set of common garden

experiments. For this, seeds from 4 independent replicate lines per genotype were sown directly at a

density of 8 seeds per pot. In total, we planted 32 pots per genotype and randomized 9-cm- diameter

pots at a field site in the botanical garden of Cologne University. Again, because we found no

significant differences between replicate lines, we determined that the experiment was complete

with 32 replicates per genotype. Seeds were sown in late summer (August 23) and in early autumn

(September 20), mid-autumn (November 8), and late winter (February 26). Rather than being

watered artificially, pots were put on a fabric that retained water after rainfall. The site was

protected against birds and rabbits by a net. Plants were inspected for germination twice a week.

Flowering time was scored when the first flower showed open petals. For fertility measurement,

siliques were counted at the end of each plant's life cycle.

Phenotypic data was analyzed using the glm function in R version 4.1.0 (R Core Team, 2022). We

used one-way ANOVA with habitat patch as the fixed factor to analyze differences in environmental

conditions (EIVs and DIVs of plant species co-occurring with *A. thaliana*); and in the phenological

stages and fitness proxies of *A. thaliana* plants observed *in situ*. Measurements of flowering time in

common garden experiments were analyzed with two-way ANOVA, using genotype and cohort or

growth condition as fixed factors.

For germination assays under controlled indoor conditions, the number of germinants per Petri dish

was analyzed with the R function zeroinfl (Package pscl; Zeileis et al., 2008) to correct for zero

inflation. For germination rates in the outdoor common garden experiment, we proceeded as under

controlled conditions, except that we used a quasipoisson distribution of error. Because the two

replicate lines we used for each genotype never showed significant effects, the line effect was not

included in any of the models. We used the function ANOVA with a Chi-squared test to determine

the overall significance of effects. In all cases, the significance of pairwise differences was tested with

a Tukey post-hoc test, using the function glht from the multcomp package in R (Hothorn et al., 2008).

We used the Pearson coefficient of the cor.test function in R to quantify the correlation between

phenotypes.

Results

Urban habitat patches of *A. thaliana* are ecologically diverse

The eight urban habitat patches differed significantly in environmental conditions related to climate,

soil, and severity of disturbance. Indicator values for light (L), and edaphic conditions -- humidity (F),

and soil nutrients (N) -- differed between sites (Figure 1b, supplemental Figure 1a, b; humidity (F):

$F_{7,113}=2.855$, p=0.00883; soil nutrients (N): $F_{7,93}=2.777$, p=0.0114, light (L): $F_{7,130}=2.329$, p=0.0285).

Indicator values of the accompanying flora for herb-layer disturbance frequency (DF) and herb-layer

structure-based disturbance (DV) differed between sites (Figure 1c, supplemental Figure 1c; DF:

$F_{7,84}=2.897$, p=0.0092; DV: $F_{7,84}=3.108$, p=0.00577), whereas differences in herb-layer disturbance

severity (DS) were not significant. The three values for disturbance were strongly and positively

correlated with each other (range of r: 0.802 – 0.958; p<0.00172) and with the temperature indicator

(range of r: 0.736 – 0.803; p<0.00632; Figure 1d; Figure S3, Table S4) .

Plants differ across habitat patches in their phenology and fitness

258   Across the eight urban *A. thaliana* populations, all phenological stages and plant functional traits

259   measured *in situ* showed significant differences (p<2.43e-06; F=6.094, Table S6; Figure S4). These are

260   exemplified in data on flowering time and rosette size (Figure S1a,b). The diverse traits describing

261   phenological time points, the temporal spacing between phenological states, and the size of the

262   plants also covaried positively (Figure S5).

Community-level indicators of environmental variables explain part of the phenotypic differences
between field sites

In situ observations of phenotypic differences between sampling sites were partly correlated with

the differences in environmental characteristics: as expected, plants from sites with high mean

values for N- and F-indicators, which reflect differences in nutrient and soil moisture levels,

respectively, were on average larger than those from sites with lower N- and F-indicator values (p-

values smaller than 0.05 for F: inflorescence length, maximal number of open flowers, rosette

diameter; for N: rosette diameter at bolting, number of rosette leaves, and number of fruits; Figure

S1c). Plants from sites with higher disturbance values bolted later than plants from other sites, but

formed larger inflorescences bearing more fruits.

Urban populations in Cologne consist of clones of diverse western European genotypes

In order to determine the extent of genomic variation and the history of population establishment,

we genotyped 5-17 individuals per population as well as plants sampled from additional sites

scattered across the study area. In total, we genotyped about 1% of the genome, or 13 Mbp, of 149

individuals using RAD-sequencing (Table S5). Overall, we uncovered 11,489 single nucleotide

polymorphisms (SNPs) in the 8 focal populations, and 17,288 SNPs if we included the scattered

sample. Individuals were overall highly homozygous (>99%), indicating that self-fertilization

dominates in Cologne populations. Populations showed a strong genetic structure, as can be seen

from the genetic distance matrix (Figure 2a): Populations BER, GHU, RKO, and SHY consisted exclusively of genetically identical individuals, with the exception of a single plant with 99% heterozygosity from the GHU site. Populations BIS, HLU, KAS, and MIL each consisted of two genotypes. Overall Fst reached 0.72. Genetic similarity allowed clustering genotypes into three groups: Group 1 comprised RKO and BER, group 2 KAS-1, KAS-2, and BIS-2, group 3 consisted of HLU-1, HLU-2, and SGY. The genotype of plants collected in 2016 was the same as that of individuals collected in 2017 in the same location, confirming that local populations are most often formed from one (maximum two) single genotypes that persist over years.

In order to understand the origin of the genotypes that established populations in the city, we compared the Cologne samples to the European genotypes of the 1001-genome collection (Alonso-Blanco et al., 2016) . We took a subset of representative genotypes from both datasets and first ran a principal component analysis (PCA) (see Materials and Methods). If habitats had been colonized via a stepping stone model, we would expect that the genotypes of the Cologne sample form a cluster of genotypes that stands out of the 1001 genomes. Cologne samples, however, co-localized with western and Central European accessions in the PCA, especially with accessions from Germany, France, the Netherlands, and the UK. They did not form a defined subgroup separate from the other genotypes (Figure 2b, c, Figure S6). Hierarchical clustering using admixture analysis confirmed that individual lineages share ancestry with individuals collected in Germany, the Netherlands or France in variable proportions (Figure 3d). In fact, the average number of pairwise differences between Cologne lineages was comparable to the number of differences between samples of the 1001 genomes (Figure S6). In conclusion, at each site, *A. thaliana* individuals formed either one or two clonal lineages, and collectively they do not form a local endemic group of similar genotypes. In addition, populations displayed no signal of isolation by geographic distance (Figure S6, Mantel test $p>0.05$) nor isolation by environmental distance (Mantel test $p>0.05$ for all environmental factors). We conclude that new populations are often established by long range migration and that the stepping stone model, by which new patches would be colonized by propagules of neighbouring patches, is inconsistent with the pattern of genetic diversity we observed (Duforet-Frebourg & Slatkin, 2016).

Cologne urban populations show genetic variation in flowering time and dormancy regulation

Variation in genetic pathways regulating flowering time and germination may contribute to life-history differences recorded on-site. To link the germination and flowering behavior of the urban genotypes to knowledge about these well-studied pathways, we analyzed flowering under standard conditions in a randomized block design and included the well-studied genotype Col-0 as a reference. In total, three controlled environmental conditions were set: long days (LD), short days (SD), and a

combination of short days/vernalization/long days (SDV). This setup thus allowed us to thoroughly examine genetic variation in the regulation of flowering time. The first two conditions allowed us to derive photoperiod-dependent differences in the regulation of flowering; the third mimicked the succession of environmental cues experienced in the fall, winter, and spring (Andrés & Coupland, 2012). Differences between LD and SDV conditions revealed variation in vernalization requirement, because it reveals the extent to which flowering time differences in LD tend to decrease in SDV as a result of exposure to the cold temperatures.

Flowering times differed between genotypes under all conditions ($F_{12,134to158} > 39.53$, $p < 2$ e^{-16}) and the interaction between genotype and experimental condition was signicant ($F_{24,461} = 36.72$, $p < 2$ e^{-16}). Under LD conditions, BER and RKO plants flowered as early as the early flowering reference genotype Col-0, whereas BIS2 plants had not flowered after 105 days, indicating that the former do not require vernalization to flower (Figure 3a, Table S7). Under SD conditions, several genotypes flowered significantly earlier than Col-0, indicating that natural Cologne genotypes might flower in the short days of the fall or the late winter: RKO plants flowered earliest; BER, KAD1, KAD2, and MIL1 flowered only slightly earlier than Col-0. Both genotypes from the BIS site were unable to flower under SD conditions and were thus unlikely to flower in the fall (Figure 3b). When plants were grown under conditions simulating winter (SDV), most genotypes flowered within a narrow time period, with only BIS1 and BIS2 displaying significantly later flowering (Figure 3c). This experiment demonstrated that there is considerable genetic variation among the 12 Cologne genotypes for the regulation of flowering time. The magnitude of these differences depended on photoperiod and vernalization.

The genotypes also differed in how they regulated germination (Table S8). After three months of seed storage at room temperature, germination assays revealed that the genotypes differed in primary and secondary dormancy. When seeds were untreated, their germination rates indicated levels of primary dormancy (Baskin & Baskin, 2004; Finkelstein et al., 2008). Untreated seeds of BER, BIS2, MIL1, and MIL2 plants had high germination rates, whereas seeds of plants from the other genotypes had a less than 50% germination rate (χ^2, df=11, N=48, $p < 2.2e^{-16}$; Figure 4a). Incubating non-dormant seeds at very low (-21°C) or high (35 °C) temperatures induced variable levels of secondary dormancy in Cologne genotypes, as indicated by a significant interaction between genotypes and germination conditions (likelihood ratio test, $\chi^2 = 83.2$, df=24, $p=1.81e^{-8}$, Figure 4c and supplemental Figure S7).

Outdoor common garden experiments confirm differences in germination and flowering regulation under field conditions

The 12 genotypes were further assayed in an outdoor common garden located in Cologne, and the
genetic variation underlying their differences in survival and fertility was quantified. Here again, we
included the widely used genotype Col-0 in the experiment as a reference (Table S9a-b). Since
phenological variation is known to depend on the timing of germination, seeds were sown directly
into pots at a field site in late August, mid-September, early November (6 weeks, 9 weeks, and 3
months after harvest), and the end of February. As pots were not watered artificially, germination
and further development depended on natural rainfall and temperature. Germination rates among
genotypes strongly differed in the August and September sowing cohorts. In the August cohort, 43
days after seeds were sown, only Col-0 plants had germinated to 100%; BER and MIL1 showed 43 and
28 % germination, respectively, 10 % or fewer of the seeds of the other genotypes had germinated
(differences between the genotypes: $F_{12,195}$=16.97, p<2e-16, Figure 4c). In the September cohort, 15
days after seeds were sown, most genotypes had germinated at 80% or more, and only HLU1, GHU,
and SGY had germinated at less than 60% (after 22 d: $F_{12,195}$=15.84, p<2e-16; Figure 4d). Eventually,
most seed germinated, but the most delayed was the August cohort (Figure S8a-d). Later
germination in this cohort can be explained by heat-induced secondary dormancy. In contrast,
differences in germination for the November and February cohorts were minor (Figure S8e,f).

More than 95 % of the germinated seedlings survived until seed set. Most flowered in spring with the
exception of the Col-0 and BER plants from the August cohort, which germinated almost immediately
and flowered already in late autumn/winter. Differences in flowering time, which were observed for
each of the four cohorts ($F_{12,195}$>13.33, p<2 e^{-16}), revealed a significant interaction between genotype
and sowing date ($F_{34,755}$= 20.89, p<2e^{-16}, Table S3, Figure 3d, Figure S9). Flowering times observed for
the 12 genotypes were generally not correlated across the various common garden cohorts grown
outdoor, with the exception of the February and September cohorts (Figure S10). Fertility among the
genotypes, determined by the number of fruits per pot, differed between the genotypes for the
August, September, and November cohorts ($F_{12,195}$=3.072, p=0.00054; $F_{12,194}$=7.721, p=1.15e^{-12}; and
$F_{12,195}$=4.702, p=1.01e^{-06}). Again, a significant interaction between genotype and seed-sowing date
was observed ($F_{24,584}$=3.534, p= 4.75e^{-08}). For the August cohort, only two genotypes (RKO and SGY)
were less fertile than most of the genotypes (Table S3, Figure S11). Fitness differences for the plants
grown from seeds sown in February, all of which germinated within a week in early March and
flowered 23-39 days later, were not significant. With 52% of the plants dying after an unusually warm
dry spell in May, the average fitness of the spring cohort was low .

In summary, plants from the 12 Cologne genotypes displayed remarkable genetic differences in their
ability to control germination and flowering in common garden experiments. Flowering times
measured in controlled growth chamber and in the common garden were significantly correlated,

but correlation depended on the various experimental indoor conditions and/or the seed dispersal
times characterizing the different common garden cohorts (Figure S10).

Environmental variation among sites of origin correlates with genetic variation observed in
common gardens and controlled environments

The phenotypes of urban *A. thaliana* individuals in situ correlate with the ecological characteristics of
the sites in which they are monitored. We further tested whether the distribution of genetic
differences revealed in common gardens and under lab conditions is independent of the ecological
characteristics at the sites of origin. We found that genetic variation in the rate of germination under
late summer conditions in the August cohort or after cold treatment correlates positively with
variation in the severity of disturbance (r=0.763, p=0.0039 and r=0.816, p=0.0012, respectively).
Furthermore, in the absence of vernalization, variation in the time to flowering correlates negatively
with variation in disturbance severity and temperatures (e.g., indicator value T (flowering date in LD
or SD): r=-0.814 and -0.806; p=0.0013 and 0.0016; Figure 5, Figure S12). Genetic variation in
phenology is thus not randomly distributed across habitats. Its distribution matches the disturbance
gradient along which the *A. thaliana* plant populations monitored in this study are distributed.

Both genetic variation for the regulation of flowering time in LD conditions and phenotypic variation
for flowering time *in situ* were correlated with disturbance frequency, but the slope of the
relationship changed signed ($F_{1,20}$=14.6984, p=0.001, Figure 5, Figure S13). We conclude that genetic
variants accelerating flowering in the absence of vernalization are found in the most disturbed sites,
where plants are observed to flower at a later date.

Discussion

Urban habitats form a mosaic of diverse environments that challenge plant growth in different ways.
However, the impact of this heterogeneity on the distribution of adaptive genetic variation is not
known (Rifkin, 2018). While the influence of urban heterogeneity on gene flow and population
differentiation has been reported in animal and plant species, its role in shaping the distribution of
genetic variation for adaptive traits remains unexplored (Beninde et al., 2015; Gorton et al., 2018;
Rivkin et al., 2019). By combining plant community and trait-based ecological measurements with an
in-depth exploration of genetic diversity at the genomic and phenotypic levels, this study bridges this
knowledge gap. The results provide compelling evidence that the genetic variation in *A. thaliana* is
not randomly distributed across environmentally diverse urban habitats. These findings enhance our
understanding of the actual ecology of a species that serves as a model in plant molecular biology
(Takou et al., 2019), and further provide valuable insights into adaptive genetic traits in urban
environments (Rivkin et al., 2019).

A. thaliana populations sampled here confirm previous reports about urban populations (Bomblies et

al., 2010): Each habitat patch is colonized by one, sometimes two, clonal lineages. An analysis of

genomic variation showed that these lineages belong to the broadly distributed gene pool of

northwestern European *A. thaliana* genotypes, supporting the notion that urban city patches were

established from a pool of regionally diverse migrants. Furthermore, our study shows that the urban

habitats hosting spontaneously occurring *A. thaliana* stands are heterogeneous. The advantage of

ecological characteristics based on community indicators, which leverage the particularly extensive

information available for the German flora, is that they provide robust measures of environmental

variation (Diekmann, 2003; Fanelli et al., 2006). Here, these characteristics reveal that *A. thaliana*

habitats form a continuum of varying levels of humidity and nutrients, as well as varying frequencies

of disturbance. The ecological differences we quantified were clearly associated with the phenotypic

differences displayed by plants *in situ,* as plants growing under better edaphic conditions (more

nutrients, more water) were larger, and plants growing in disturbed habitats bolted later and bore

more fruits. *In situ* observations of plant diversity, however, do not allow disentangling the effects of

plant plasticity to the environment from the effect of genetic differences between populations

(Aguilar-Trigueros et al., 2017; Cadotte & Tucker, 2017; Thakur & Wright, 2017).   Unraveling whether

the successful occupation of diverse urban habitat patches is solely facilitated by plant plasticity

represents a critical step towards understanding the ecological dynamics of biodiversity in urban

environments (Brans et al., 2017; Rivkin et al., 2019).

Focusing on the 12 clonal lineages identified by genomic analysis, we thus explored whether genetic

variation contributes to the phenological variation observed *in situ*.   By comparing the lineages

forming each population patch under common growth conditions, we found strong genetic

differences in how they regulate seed germination and flowering. In total, we relied on at least seven

indoor and outdoor common garden conditions to show that the 12 clonal lineages varied in their

requirements for vernalization and their ability to flower during shortening daylength, as well as in

the ability of their seeds to delay germination after exposure to cold and heat. We confirm previous

findings that the timing of germination plays a major role on life-history traits manifested later in the

life cycle (Donohue, 2009; Wilczek et al., 2009). By contrasting variation in plant traits expressed *in*

situ with *ex situ* observations made in common garden settings, we showed that genetic variation in

phenology tends to buffer the effect that environmental heterogeneity has on trait variation. Indeed,

we found that genotypes that flowered early under LD or SD conditions (*ex situ*) were found in sites

that were more disturbed. Whereas, those from the less disturbed sites flowered later under LD or

SD conditions but flowered early *in situ* (Figure 5). Indeed, the timing of flowering in LD conditions

and *in situ* showed opposite patterns of covariation with disturbance frequency (Figure 5).

Therefore, genotypes growing in habitats with relatively harsher environmental conditions

(characterized by resource stress, frequent disturbance, and a delay in development) tend to have
traits genetically tailored to a life cycle that is shorter in the absence of vernalization. We thereby
reveal a remarkable pattern of counter-gradient adaptation (Conover & Schultz, 1995), wherein
genetic variation for flowering time in LD reduces the extent of phenological variation that would
otherwise typify the heterogeneous environments hosting *A. thaliana* in the city. Therefore, variable
genetic traits may not always be directly observable across environmental gradients. Counter-
gradient variation is indeed a well-known confounding experimental error for *in-situ* studies, that can
only be resolved with genetic studies in common gardens (Conover & Schultz, 1995).

We believe that, in this study, the distribution of genetic differences in the regulation of flowering
reflects the outcome of a process driven by local selective pressures. Indeed, we found no correlation
between genetic distance and geographic or ecological distance, so the non-random pattern of co-
variation between disturbance level and genetic variation in life-history cannot be explained by local
seed dispersal. We identify two other potential explanations: i) specialized genotypes have adapted
via the selection of new mutations affecting life history in an otherwise non-adapted lineage (Orr,
2005), or ii) local environments filtered mis-adapted genotypes out, in the same way that they
exclude mis-adapted species (Guo et al., 2018; Kraft et al., 2015; Laliberté et al., 2014; Yang et al.,
2022). Since many alleles altering the genetics of flowering time have already been characterized in
the western European region from which the populations originate (Le Corre, 2005; Lopez-Arboleda
et al., 2021), it is highly likely that genotypes with diverse regulatory alleles were among the pool of
migrants that populated habitat patches in the city of Cologne. Environmental filtering thus appears
to be the most parsimonious explanation for the non-random distribution of genetic variation in
flowering time documented in this study. Considering that the assessment of variation *in situ* and at
the genetic level constrained us to the study of only 8 populations, our findings suggest that urban
environmental heterogeneity has a strong filtering power on the genetic basis of intraspecific trait
variation. Environmental filters imposed by habitat heterogenetiy in cities have been shown to
diversify plant communities (Aronson et al., 2016). Indeed, cities can preserve species diversity,
depending, for example, on the size and connectivity of habitats within cities (Beninde et al., 2015;
Hahs et al., 2009). To the best of our knowledge, this study is the first to show that environmental
filtering also has the potential to maintain functionally diverse and locally adapted genetic lineages.
Such lineages form *bona fide* ecotypes, a term often used in the literature but seldom scrutinized.

Our assessment of *in situ* and *ex situ* variation in phenology highlights the importance of
interdisciplinary approaches in ecology, quantitative genetics, and genomics. Indeed, our study
identifies a pattern of adaptive genetic trait variation that is not directly observable in situ. With our
sample of a small number of highly structured populations composed of one or two clonal lineages,
we do not yet have the power to identify the loci responsible for adaptive life history variation

(Hancock et al., 2011; Herden et al., 2019). The number of variants that contribute to the genetic

variation of plant phenology in Cologne *A. thaliana* populations will have to be determined in future

studies.

Finally, one can speculate that disturbances may be the driving force behind the environmental

filtering of genetic variation in life history within the city. However, in urban plant communities,

environmental filters are shaped by a complex interplay of various factors, including urban habitat

transformation, fragmentation, typical urban alterations such as heat islands or pollution, and human

preferences(Williams et al., 2009). It is highly likely that each of these factors co-varies with the

degree of disturbance. Consequently, future studies focusing on genetic variation within cities should

intend to disentangle the specific factors filtering genetic variation in life history traits within urban

plant populations and communities.

Acknowledgments

We thank Denitsa Ilieva for help in common garden experiments, and Emily Wheeler, Harwich, MA,

for editorial assistance. We acknowledge funding from the German Science Foundation in the

framework of TRR341 "Plant Ecological Genetics" and from the Center of Excellence in Plant Sciences

(CEPLAS), in Cologne and Düsseldorf.

Authors' Contributions

AL, JM, GS, AS conceived the ideas and designed the methodology; JM, GS and AL wrote the

manuscript; AF, GS, HD, JT, GC collected and/or analyzed the data; all authors contributed to the

draft and gave final approval for publication. Sequencing data was deposited as submission numbers

ERS5063293 to ERS5063441, project PRJEB40091 in the ENA repository. We acknowledge funding

from the Deutsche Forschungsgemeinschaft within the framework of TRR341.

References

Ågren, J., Oakley, C. G., Lundemo, S., & Schemske, D. W. (2017). Adaptive divergence in flowering

time among natural populations of Arabidopsis thaliana: Estimates of selection and QTL

mapping. *Evolution*, *71*(3), Article 3. https://doi.org/10.1111/evo.13126

Aguilar-Trigueros, C. A., Rillig, M. C., & Ballhausen, M.-B. (2017). Environmental Filtering Is a Relic. A

Response to Cadotte and Tucker. *Trends in Ecology & Evolution*, *32*(12), 882–884.

https://doi.org/10.1016/j.tree.2017.09.013

Akiyama, R., & Ågren, J. (2018). Conflicting selection on the timing of germination in a natural

population of Arabidopsis thaliana. *Journal of Evolutionary Biology*, *27*(1), Article 1.

https://doi.org/10.1111/jeb.12293

Alonso-Blanco, C., Andrade, J., Becker, C., Bemm, F., Bergelson, J., Borgwardt, K. M., Cao, J., Chae, E.,

Dezwaan, T. M., Ding, W., Ecker, J. R., Exposito-Alonso, M., Farlow, A., Fitz, J., Gan, X., Grimm,

D. G., Hancock, A. M., Henz, S. R., Holm, S., … Zhou, X. (2016). 1,135 Genomes Reveal the

Global Pattern of Polymorphism in Arabidopsis thaliana. *Cell*, *166*(2), 481–491.

https://doi.org/10.1016/j.cell.2016.05.063

Andrés, F., & Coupland, G. (2012). The genetic basis of flowering responses to seasonal cues. *Nature*

Reviews Genetics, *13*(9), Article 9. https://doi.org/10.1038/nrg3291

Ansmann, L., Vennedey, V., Hillen, H. A., Stock, S., Kuntz, L., Pfaff, H., Mannion, R., Hower, K. I., &

Cologne Research And Development Network CoRe-Net Study Group, C. R. A. D. N. C.-N. S. G.

(2021). Resource dependency and strategy in healthcare organizations during a time of

scarce resources: Evidence from the metropolitan area of cologne. *Journal of Health*

Organization and Management, *35*(9), 211–227. https://doi.org/10.1108/JHOM-12-2020-

0478

Aronson, M. F. J., Nilon, C. H., Lepczyk, C. A., Parker, T. S., Warren, P. S., Cilliers, S. S., Goddard, M. A.,

Hahs, A. K., Herzog, C., Katti, M., La Sorte, F. A., Williams, N. S. G., & Zipperer, W. (2016).

Hierarchical filters determine community assembly of urban species pools. *Ecology*, *97*(11),

2952–2963. https://doi.org/10.1002/ecy.1535

Baskin, J. M., & Baskin, C. C. (2004). A classification system for seed dormancy. *Seed Science*

Research, *14*(1), Article 1.

Beninde, J., Veith, M., & Hochkirch, A. (2015). Biodiversity in cities needs space: A meta-analysis of

factors determining intra-urban biodiversity variation. *Ecology Letters*, *18*(6), 581–592.

https://doi.org/10.1111/ele.12427

Bomblies, K., Yant, L., Laitinen, R. A., Kim, S.-T., Hollister, J. D., Warthmann, N., Fitz, J., & Weigel, D.

(2010). Local-Scale Patterns of Genetic Variability, Outcrossing, and Spatial Structure in

Natural Stands of Arabidopsis thaliana. *PLoS Genetics*, *6*(3), Article 3.

https://doi.org/10.1371/journal.pgen.1000890.g005

Bonthoux, S., Voisin, L., Bouché-Pillon, S., & Chollet, S. (2019). More than weeds: Spontaneous

vegetation in streets as a neglected element of urban biodiversity. *Landscape and Urban*

Planning, *185*, 163–172. https://doi.org/10.1016/j.landurbplan.2019.02.009

Brans, K. I., Jansen, M., Vanoverbeke, J., Tüzün, N., Stoks, R., & De Meester, L. (2017). The heat is on:

Genetic adaptation to urbanization mediated by thermal tolerance and body size. *Global*

Change Biology, *23*(12), 5218–5227. https://doi.org/10.1111/gcb.13784

Cadotte, M. W., & Tucker, C. M. (2017). Should Environmental Filtering be Abandoned? *Trends in*

Ecology & Evolution, *32*(6), 429–437. https://doi.org/10.1016/j.tree.2017.03.004

Castilla, A. R., Méndez-Vigo, B., Marcer, A., Martínez-Minaya, J., Conesa, D., Picó, F. X., & Alonso-

Blanco, C. (2020). Ecological, genetic and evolutionary drivers of regional genetic

differentiation in Arabidopsis thaliana. *BMC Evolutionary Biology*, *20*(1), Article 1.

https://doi.org/10.1186/s12862-020-01635-2

Chatzidimitriou, A., & Yannas, S. (2015). Microclimate development in open urban spaces: The

influence of form and materials. *Energy and Buildings*, *108*, 156–174.

https://doi.org/10.1016/j.enbuild.2015.08.048

Cheptou, P.-O., Carrue, O., Rouifed, S., & Cantarel, A. (2008). Rapid evolution of seed dispersal in an

urban environment in the weed Crepis sancta. *Proceedings of the National Academy of*

Sciences, *105*(10), 3796–3799. https://doi.org/10.1073/pnas.0708446105

Chmielewski, F.-M., & Rötzer, T. (2001). Response of tree phenology to climate change across

Europe. *Agricultural and Forest Meteorology*, *108*(2), 101–112.

https://doi.org/10.1016/S0168-1923(01)00233-7

Conover, D. O., & Schultz, E. T. (1995). Phenotypic similarity and the evolutionary significance of

countergradient variation. *Trends in Ecology & Evolution*, *10*(6), 248–252.

https://doi.org/10.1016/S0169-5347(00)89081-3

Debieu, M., Tang, C., Stich, B., Sikosek, T., EFFGEN, S., Josephs, E., Schmitt, J., Nordborg, M.,

Koornneef, M., & de Meaux, J. (2013). Co-Variation between Seed Dormancy, Growth Rate

and Flowering Time Changes with Latitude in Arabidopsis thaliana. *PLoS ONE*, *8*(5), Article 5.

https://doi.org/10.1371/journal.pone.0061075.s002

DeLeo, V. L., Menge, D. N. L., Hanks, E. M., Juenger, T. E., & Lasky, J. R. (2020). Effects of two

centuries of global environmental variation on phenology and physiology of Arabidopsis

thaliana. *Global Change Biology*, *26*(2), 523–538. https://doi.org/10.1111/gcb.14880

Diekmann, M. (2003). Species indicator values as an important tool in applied plant ecology – a

review. *Basic and Applied Ecology*, *4*(6), 493–506. https://doi.org/10.1078/1439-1791-00185

Dittberner, H., Becker, C., Jiao, W.-B., Schneeberger, K., Hölzel, N., Tellier, A., & Meaux, J. de. (2019).

Strengths and potential pitfalls of hay transfer for ecological restoration revealed by RAD-seq

analysis in floodplain Arabis species. *Molecular Ecology*, *28*(17), Article 17.

https://doi.org/10.1111/mec.15194

Donohue, K. (2009). Some Evolutionary Consequences of Niche Construction with Genotype-

Environment Interaction. *Adaptation and Fitness in Animal Populations*, 131–149.

Dubin, M. J., Zhang, P., Meng, D., Remigereau, M.-S., Osborne, E. J., Paolo Casale, F., Drewe, P.,

Kahles, A., Jean, G., Vilhjálmsson, B., Jagoda, J., Irez, S., Voronin, V., Song, Q., Long, Q.,

Rätsch, G., Stegle, O., Clark, R. M., & Nordborg, M. (2015). DNA methylation in Arabidopsis

has a genetic basis and shows evidence of local adaptation. *ELife*, *4*, e05255.

https://doi.org/10.7554/eLife.05255

Dubois, J., & Cheptou, P.-O. (2017). Effects of fragmentation on plant adaptation to urban

environments. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

372(1712), 20160038. https://doi.org/10.1098/rstb.2016.0038

Duforet-Frebourg, N., & Slatkin, M. (2016). Isolation-by-distance-and-time in a stepping-stone model.

Theoretical Population Biology, *108*, 24–35. https://doi.org/10.1016/j.tpb.2015.11.003

Durvasula, A., Fulgione, A., Gutaker, R. M., Alacakaptan, S. I., Flood, P. J., Neto, C., Tsuchimatsu, T.,

Burbano, H. A., Picó, F. X., Alonso-Blanco, C., & Hancock, A. M. (2017). African genomes

illuminate the early history and transition to selfing in Arabidopsis thaliana. *Proceedings of*

the National Academy of Sciences of the United States of America, *209*, 201616736.

Ellenberg, H., & Leuschner, C. (2010). *Vegetation Mitteleuropas mit den Alpen: In ökologischer,*

dynamischer und historischer Sicht. UTB.

Exposito-Alonso, M., Vasseur, F., Ding, W., Wang, G., Burbano, H. A., & Weigel, D. (2018). Genomic

basis and evolutionary potential for extreme drought adaptation in Arabidopsis thaliana.

Nature Ecology & Evolution, *2*(2), Article 2. https://doi.org/10.1038/s41559-017-0423-0

Fanelli, G., Tescarollo, P., & Testi, A. (2006). Ecological indicators applied to urban and suburban

floras. *Ecological Indicators*, *6*(2), 444–457. https://doi.org/10.1016/j.ecolind.2005.06.002

Finkelstein, R., Reeves, W., Ariizumi, T., & Steber, C. (2008). Molecular Aspects of Seed Dormancy.

Annual Review of Plant Biology, *59*(1), 387–415.

https://doi.org/10.1146/annurev.arplant.59.032607.092740

Foti, L. (2017). *Assesment of the urban pressure effects on soil quality in the Paris region under two*

types of vegetation (lawns and woods). https://tel.archives-ouvertes.fr/tel-

02393689/file/2017PA066481.pdf

Fournier-Level, A., Korte, A., Cooper, M. D., Nordborg, M., Schmitt, J., & Wilczek, A. M. (2011). A map

of local adaptation in Arabidopsis thaliana. *Science*, *334*(6052), Article 6052.

https://doi.org/10.1126/science.1209271

Gibson, D. J. (2014). *Methods in Comparative Plant Population Ecology*.

https://doi.org/10.1093/acprof:oso/9780199671465.001.0001

Gilbert, O. (2012). *The Ecology of Urban Habitats*. Springer Science & Business Media.

Gorton, A. J., Moeller, D. A., & Tiffin, P. (2018). Little plant, big city: A test of adaptation to urban

environments in common ragweed (Ambrosia artemisiifolia). *Proceedings of the Royal*

Society B: Biological Sciences, *285*(1881), 20180968. https://doi.org/10.1098/rspb.2018.0968

Guo, P., Yu, F., Ren, Y., Liu, D., Li, J., Ouyang, Z., & Wang, X. (2018). Response of Ruderal Species

Diversity to an Urban Environment: Implications for Conservation and Management.

International Journal of Environmental Research and Public Health, *15*(12), Article 12. https://doi.org/10.3390/ijerph15122832

Hahs, A. K., McDonnell, M. J., McCarthy, M. A., Vesk, P. A., Corlett, R. T., Norton, B. A., Clemants, S. E., Duncan, R. P., Thompson, K., Schwartz, M. W., & Williams, N. S. G. (2009). A global synthesis of plant extinction rates in urban areas. *Ecology Letters*, *12*(11), 1165–1173. https://doi.org/10.1111/j.1461-0248.2009.01372.x

Halbritter, A. H., Fior, S., Keller, I., Billeter, R., Edwards, P. J., Holderegger, R., Karrenberg, S., Pluess, A. R., Widmer, A., & Alexander, J. M. (2018). Trait differentiation and adaptation of plants along elevation gradients. *Journal of Evolutionary Biology*, *31*(6), 784–800. https://doi.org/10.1111/jeb.13262

Hancock, A. M., Brachi, B., Faure, N., Horton, M. W., Jarymowycz, L. B., Sperone, F. G., Toomajian, C., Roux, F., & Bergelson, J. (2011). Adaptation to climate across the Arabidopsis thaliana genome. *Science*, *334*(6052), Article 6052. https://doi.org/10.1126/science.1209244

Herben, T., Chytrý, M., & Klimešová, J. (2016). A quest for species-level indicator values for disturbance. *Journal of Vegetation Science*, *27*(3), 628–636. https://doi.org/10.1111/jvs.12384

Herden, J., Eckert, S., Stift, M., Joshi, J., & van Kleunen, M. (2019). No evidence for local adaptation and an epigenetic underpinning in native and non-native ruderal plant species in Germany. *Ecology and Evolution*, *9*(17), 9412–9426. https://doi.org/10.1002/ece3.5325

Hothorn, T., Bretz, F., & Westfall, P. (2008). Simultaneous inference in general parametric models. *Biometrical Journal. Biometrische Zeitschrift*, *50*(3), 346–363. https://doi.org/10.1002/bimj.200810425

Huang, L., Koubek, T., Weiser, M., & Herben, T. (2018). Environmental drivers and phylogenetic constraints of growth phenologies across a large set of herbaceous species. *Journal of Ecology*, *106*(4), 1621–1633. https://doi.org/10.1111/1365-2745.12927

Johnson, M. T. J., & Munshi-South, J. (2017). Evolution of life in urban environments. *Science*, *358*(6363), eaam8327. https://doi.org/10.1126/science.aam8327

Kawecki, T. J., & Ebert, D. (2004). Conceptual issues in local adaptation. *Ecology Letters*, *7*(12), Article

12. https://doi.org/10.1111/j.1461-0248.2004.00684.x

Kollmann, J., & Fischer, A. (2003). Vegetation as indicator for habitat quality. *Basic and Applied*

Ecology, *4*(6), 489–491. https://doi.org/10.1078/1439-1791-00209

Kraft, N. J. B., Adler, P. B., Godoy, O., James, E. C., Fuller, S., & Levine, J. M. (2015). Community

assembly, coexistence and the environmental filtering metaphor. *Functional Ecology*, *29*(5),

592–599. https://doi.org/10.1111/1365-2435.12345

Kronholm, I., Picó, F. X., Alonso-Blanco, C., Goudet, J., & Meaux, J. de. (2012). Genetic Basis of

Adaptation in Arabidopsis thaliana: Local Adaptation at the Seed Dormancy QTL DOG1.

Evolution, *66*(7), Article 7. https://doi.org/10.1111/j.1558-5646.2012.01590.x

Laliberté, E., Zemunik, G., & Turner, B. L. (2014). Environmental filtering explains variation in plant

diversity along resource gradients. *Science*, *345*(6204), 1602–1605.

https://doi.org/10.1126/science.1256330

Lambrecht, S. C., Mahieu, S., & Cheptou, P.-O. (2016). Natural selection on plant physiological traits

in an urban environment. *Acta Oecologica*, *77*, 67–74.

https://doi.org/10.1016/j.actao.2016.09.002

Le Corre, V. (2005). Variation at two flowering time genes within and among populations of

Arabidopsis thaliana: Comparison with markers and traits. *Molecular Ecology*, *14*(13), Article

13.

Lee, C.-R., Svardal, H., Farlow, A., Exposito-Alonso, M., Ding, W., Novikova, P., Alonso-Blanco, C.,

Weigel, D., & Nordborg, M. (2017). On the post-glacial spread of human commensal

Arabidopsis thaliana. *Nature Communications*, *8*(1), Article 1.

https://doi.org/10.1038/ncomms14458

Leimu, R., & Fischer, M. (2008). A meta-analysis of local adaptation in plants. *PLoS ONE*, *3*(12), Article

12. https://doi.org/10.1371/journal.pone.0004010

Lenssen, J. P. M., van Kleunen, M., Fischer, M., & de Kroon, H. (2004). Local Adaptation of the Clonal

Plant Ranunculus reptans to Flooding along a Small-Scale Gradient. *Journal of Ecology*, *92*(4),

696–706.

Li, D., Stucky, B. J., Deck, J., Baiser, B., & Guralnick, R. P. (2019). The effect of urbanization on plant

phenology depends on regional temperature. *Nature Ecology & Evolution*, *3*(12), Article 12.

https://doi.org/10.1038/s41559-019-1004-1

Liu, Y., Li, G., Wu, X., Niklas, K. J., Yang, Z., & Sun, S. (2021). Linkage between species traits and plant

phenology in an alpine meadow. *Oecologia*, *195*(2), 409–419.

https://doi.org/10.1007/s00442-020-04846-y

Lopez-Arboleda, W. A., Reinert, S., Nordborg, M., & Korte, A. (2021). Global Genetic Heterogeneity in

Adaptive Traits. *Molecular Biology and Evolution*, *38*(11), 4822–4831.

https://doi.org/10.1093/molbev/msab208

Luo, Y., Widmer, A., & Karrenberg, S. (2015). The roles of genetic drift and natural selection in

quantitative trait divergence along an altitudinal gradient in Arabidopsis thaliana. *Heredity*,

114(2), Article 2. https://doi.org/10.1038/hdy.2014.89

Nord, E. A., & Lynch, J. P. (2009). Plant phenology: A critical controller of soil resource acquisition.

Journal of Experimental Botany, *60*(7), 1927–1937. https://doi.org/10.1093/jxb/erp018

Nordt, B., Hensen, I., Bucher, S. F., Freiberg, M., Primack, R. B., Stevens, A.-D., Bonn, A., Wirth, C.,

Jakubka, D., Plos, C., Sporbert, M., & Römermann, C. (2021). The PhenObs initiative: A

standardised protocol for monitoring phenological responses to climate change using

herbaceous plant species in botanical gardens. *Functional Ecology*, *35*(4), 821–834.

https://doi.org/10.1111/1365-2435.13747

Orr, H. A. (2005). The genetic theory of adaptation: A brief history. *Nature Reviews Genetics*, *6*(2),

Article 2. https://doi.org/10.1038/nrg1523

Piao, S., Liu, Q., Chen, A., Janssens, I. A., Fu, Y., Dai, J., Liu, L., Lian, X., Shen, M., & Zhu, X. (2019). Plant

phenology and global climate change: Current progresses and challenges. *Global Change

Biology*, *25*(6), 1922–1940. https://doi.org/10.1111/gcb.14619

Primack, R. B., & Miller-Rushing, A. J. (2012). Uncovering, Collecting, and Analyzing Records to

Investigate the Ecological Impacts of Climate Change: A Template from Thoreau's Concord.

BioScience, *62*(2), 170–181. https://doi.org/10.1525/bio.2012.62.2.10

R Core Team. (2022). *R: A language and environment for statistical computin.* [Computer software]. R

Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Rivkin, L. R., Santangelo, J. S., Alberti, M., Aronson, M. F. J., de Keyzer, C. W., Diamond, S. E., Fortin,

M.-J., Frazee, L. J., Gorton, A. J., Hendry, A. P., Liu, Y., Losos, J. B., MacIvor, J. S., Martin, R. A.,

McDonnell, M. J., Miles, L. S., Munshi-South, J., Ness, R. W., Newman, A. E. M., … Johnson, M.

T. J. (2019). A roadmap for urban evolutionary ecology. *Evolutionary Applications*, *12*(3),

384–398. https://doi.org/10.1111/eva.12734

Robionek-Selosse, A., Banaś, K., Merdalski, M., & Szmeja, J. (2021). Phenotypic trait variation and life

strategy in Ranunculus reptans L. facing water level changes. *Botany Letters*, *168*(2), 283–

296. https://doi.org/10.1080/23818107.2020.1860815

Santangelo, J. S., Ness, R. W., Cohan, B., Fitzpatrick, C. R., Innes, S. G., Koch, S., Miles, L. S., Munim, S.,

Peres-Neto, P. R., Prashad, C., Tong, A. T., Aguirre, W. E., Akinwole, P. O., Alberti, M., Álvarez,

J., Anderson, J. T., Anderson, J. J., Ando, Y., Andrew, N. R., … Johnson, M. T. J. (2022). Global

urban environmental change drives adaptation in white clover. *Science*, *375*(6586), 1275–

1281. https://doi.org/10.1126/science.abk0989

Sartori, K., Vasseur, F., Violle, C., Baron, E., Gerard, M., Rowe, N., Ayala-Garay, O., Christophe, A.,

Jalón, L. G. de, Masclef, D., Harscouet, E., Granado, M. del R., Chassagneux, A., Kazakou, E., &

Vile, D. (2019). Leaf economics and slow-fast adaptation across the geographic range of

Arabidopsis thaliana. *Scientific Reports*, *9*(1), 10758. https://doi.org/10.1038/s41598-019-

46878-2

Szulkin, M., Munshi-South, J., & Charmantier, A. (2020). *Urban Evolutionary Biology*. Oxford

University Press.

Takou, M., Wieters, B., Kopriva, S., Coupland, G., Linstädter, A., & de Meaux, J. (2019). Linking genes

with ecological strategies in Arabidopsis thaliana. *Journal of Experimental Botany*.70(4),

1141-1151. https://doi.org/10.1093/jxb/ery447

Thakur, M. P., & Wright, A. J. (2017). Environmental Filtering, Niche Construction, and Trait

Variability: The Missing Discussion. *Trends in Ecology & Evolution*, *32*(12), 884–886.

https://doi.org/10.1016/j.tree.2017.09.014

Thompson, K. A., Renaudin, M., & Johnson, M. T. J. (2016). Urbanization drives the evolution of

parallel clines in plant populations. *Proceedings of the Royal Society B: Biological Sciences*,

283(1845), 20162180. https://doi.org/10.1098/rspb.2016.2180

Tresch, S., Moretti, M., Le Bayon, R.-C., Mäder, P., Zanetta, A., Frey, D., & Fliessbach, A. (2018). A

Gardener's Influence on Urban Soil Quality. *Frontiers in Environmental Science*, *6*.

https://www.frontiersin.org/article/10.3389/fenvs.2018.00025

Vasseur, F., Sartori, K., Baron, E., Fort, F., Kazakou, E., Segrestin, J., Garnier, E., Vile, D., & Violle, C.

(2018). Climate as a driver of adaptive variations in ecological strategies in Arabidopsis

thaliana. *Annals of Botany*, *122*(6), Article 6. https://doi.org/10.1093/aob/mcy165

Vega, K. A., & Küffer, C. (2021). Promoting wildflower biodiversity in dense and green cities: The

important role of small vegetation patches. *Urban Forestry & Urban Greening*, *62*, 127165.

https://doi.org/10.1016/j.ufug.2021.127165

Vidal, E. A., Moyano, T. C., Canales, J., & Gutiérrez, R. A. (2014). Nitrogen control of developmental

phase transitions in Arabidopsis thaliana. *Journal of Experimental Botany*, *65*(19), 5611–5618.

https://doi.org/10.1093/jxb/eru326

Wieters, B., Steige, K. A., He, F., Koch, E. M., Ramos-Onsins, S. E., Gu, H., Guo, Y.-L., Sunyaev, S., &

Meaux, J. de. (2021). Polygenic adaptation of rosette growth in Arabidopsis thaliana. *PLOS*

Genetics, *17*(1), Article 1. https://doi.org/10.1371/journal.pgen.1008748

Wilczek, A. M., Roe, J. L., Knapp, M. C., Cooper, M. D., Lopez-Gallego, C., Martin, L. J., Muir, C. D., Sim,

S., Walker, A., Anderson, J., Egan, J. F., Moyers, B. T., Petipas, R., Giakountis, A., Charbit, E.,

Coupland, G., Welch, S. M., & Schmitt, J. (2009). Effects of genetic perturbation on seasonal

life history plasticity. *Science*, *323*(5916), Article 5916.

https://doi.org/10.1126/science.1165826

Williams, N. S. G., Schwartz, M. W., Vesk, P. A., McCarthy, M. A., Hahs, A. K., Clemants, S. E., Corlett,

R. T., Duncan, R. P., Norton, B. A., Thompson, K., & McDonnell, M. J. (2009). A conceptual

framework for predicting the effects of urban environments on floras. *Journal of Ecology*,

97(1), 4–9. https://doi.org/10.1111/j.1365-2745.2008.01460.x

Yang, J., Su, P., Zhou, Z., Shi, R., & Ding, X. (2022). Environmental filtering rather than dispersal

limitation dominated plant community assembly in the Zoige Plateau. *Ecology and Evolution*,

12(7), e9117. https://doi.org/10.1002/ece3.9117

Zeileis, A., Kleiber, C., & Jackman, S. (2008). Regression Models for Count Data in R. *Journal of*

Statistical Software, *27*, 1–25. https://doi.org/10.18637/jss.v027.i08

Figure Legends

Fig. 1: Characteristics of eight urban *Arabidopsis thaliana* populations and their habitat conditions in the city of Cologne, with (a) differences in size and phenological stage between *A. thaliana* from two adjacent habitats (both photographs from 2017-04-03); (b) differences between habitat patches based on plant species' Ellenberg indicator values for soil nutrients and (c) the same based on plant species' Herben indicator for disturbance frequency. Letters indicate significant differences at p<0.05. (d) Similarity of habitat patches based on a PCA with plant communities' average indicator values for abiotic environmental conditions and disturbances. For indicator names, see Materials and Methods.

Fig. 2: a) Analysis of genetic distance between progeny of 85 individual plants from 8 sampling sites in Cologne determined by RAD-sequencing. Individual plants are colour coded according to their origin. b) Principal component analysis of genetic differences between Cologne plants from the 8 sampling sites (blue), plants from additional sites of the sampling area (red) and 357 selected genotypes of the Arabidopsis 1001 genome set (grey). Origin of 1001 genome genotypes is indicated for countries represented by higher number of accessions (AUT-Austria, CZE-Czech Republic, ESP-Spain, FRA-France, GER-Germany, ITA-Italy, NED-The Netherlands, SUI-Switzerland, SWE-Sweden, UK-United Kingdom). c) Magnification of the yellow marked part of the plot b). d) Proportions of ancestry shared between Cologne genomes (CLN) and genomes collected in Europe was estimated by ADMIXTURE. On the basis of minimum cross-validation error, K=8 clusters were identified in the sample of 357 previously published European genomes . From top to bottom are shown genomes from Cologne, then from European dataset. The latter are grouped by country of origin and sorted by ancestry within groups.

Fig. 3: Flowering time (days after sowing) of plants grown in controlled conditions: long days (16 h light/20 °C; 8 h dark/18°C) (a), short days (10 h light/18 °C; 14 h dark/14 °C) (b), or short days followed by vernalisation (8 h light/4 °C; 16 h dark/4 °C) and shift to long days at day 64 (c). Flowering time (days of following year) of plants sown on dry soil on September 20[th] and grown without artificial watering in the botanical garden (d). Letters indicate significant differences at a level p<0,05.

Fig. 4: Germination frequency of seeds from the 12 different Cologne genotypes and Col-0 controls. Seeds were germinated on filter paper at 20 °C in petri dishes after 3-4 moths of afterripening to test for primary dormancy (a) or incubated at 35 °C in the dark for 4 days before incubation at 20 °C to observe heat induced secondary dormancy (b) (4 replicates per genotype). Germination frequency of seeds counted on October 5[th], that were sown on dry soil in a common garden experiment without artificial watering on August 23[rd] (c) or September 20[th] (d) demonstrating different reaction to heat induced secondary dormancy under natural field conditions. Letters indicate significant differences at a level p<0,05.

Fig. 5: Variation of phenology in urban *A. thaliana* genotypes, as observed within urban habitat patches in the City of Cologne (in-situ; blue), and in a common-garden experiment in the same city (ex-situ; red) as a function of variation in the index of Disturbance frequency at the site of origin. The timing of flowering in-situ (blue) increases with disturbance frequency, whereas the opposite trend is observable for the timing of flowering in indoor common garden conditions (red).

distance

0.002 0.003

BER
BIS
GHU
HLU
KAD
MIL
RKO
SGY

(c)

P

SWE

ITA

ER

CZE

ained Variance: 1.6%)

CZE
AUT

UK
NED
FRA
S
GER
ITA

(d)

Fraction of ancestry shared between groups

0   0.5   1

CLN

AUT

CZE

ESP

FRA

GEO

GER

ITA

NED

RUS

SUI

SWE

UK

(c)

(d)

(c)

(d)

5: