The effect of morphine on rat microglial phagocytic activity: an in vitro study of brain region-, plating density-, sex-, morphine concentration-, and receptor-dependency

David N. King'uyu[a*], Lily Nti-Kyemereh[a,b], Jesse L. Bonin[c], Paul J. Feustel[a], Michelle Tram[b], Katherine C. MacNamara[c], Ashley M. Kopec[a*]

a) Department of Neuroscience and Experimental Therapeutics
Albany Medical College
Albany, NY 12208

b) Siena College
Loudonville, NY 12211

c) Department of Immunology and Microbial Disease
Albany Medical College
Albany, NY 12208

*) Corresponding Author
43 New Scotland, NY 12208
ndambud@amc.edu

<u>ABSTRACT</u>
Opioids have long been used for clinical pain management, but also have addictive properties that have contributed to the ongoing opioid epidemic. While opioid activation of opioid receptors is well known to contribute to reward and reinforcement, data now also suggest that opioid activation of immune signaling via toll-like receptor 4 (TLR4) may also play a role in addiction-like processes. TLR4 expression is enriched in immune cells, and in the nervous system is primarily expressed in microglia. Microglial phagocytosis is important for developmental, homeostatic, and pathological processes. To examine how morphine impacts microglial phagocytosis, we isolated microglia from adult male and female rat cortex and striatum and plated them *in vitro* at 10,000 (10K) or 50,000 cells/well densities. Microglia were incubated with neutral fluorescent microbeads to stimulate phagocytosis in the presence of one of four morphine concentrations. We found that the brain region from which microglia are isolated and plating density, but not morphine concentration, impact cell survival *in vitro*. We found that 10^{-12}M morphine, but not higher concentrations, increases phagocytosis in striatal microglia *in vitro* independent of sex and plating density, while 10^{-12}M morphine increased phagocytosis in cortical microglia *in vitro* independent of sex, but contingent on plating density. Finally, we demonstrate that the effect of 10^{-12}M morphine in striatal microglia plated at 10K density is mediated via TLR4, and not µORs. Overall, our data suggest that in rats, a morphine-TLR4 signaling pathway increases phagocytic activity in microglia independent of sex. This may be useful information for better understanding the possible neural outcomes associated with morphine exposures.

INTRODUCTION

Opioids have long been used for clinical pain management, but also have addictive properties (Strang et al., 2020; Volkow et al., 2019; Yoo et al., 2022) that have contributed to the ongoing opioid epidemic (Stoicea et al., 2019; Terplan, 2017). The opioid morphine binds to both opioid receptors (such as mu-opioid receptor, μOR) which are present on several different cell types, and toll-like receptor 4 (TLR4), which is enriched on immune cells (X. Wang et al., 2012; Xie et al., 2017). Opioid receptors are coupled to Gi/o proteins that inhibit adenylate cyclase, decrease cyclic adenosine monophosphate levels, and recruit β-arrestin which activates p38, AKT and ERK pathways (Al-Hasani & Bruchas, 2011; Fu et al., 2014; Valentino & Volkow, 2018). TLR4 is coupled to MyD88 and TRIF which stimulate the translocation of NF-κB to the nucleus leading to production of inflammatory cytokines and type I interferons (Lu et al., 2008; Rahimifard et al., 2017). Functionally, in peripheral immune cells opioid receptor activation induces synthesis and release of opioid peptides (Boué et al., 2011; Schäfer et al., 1994), while TLR4 activation generally stimulates pro-inflammatory and phagocytic clearance activity (Ninkovic et al., 2016; Schaub et al., 2004). Phagocytic activity is a hallmark of immune macrophages that serves to clear damage, debris, and infection (Andoh & Koyama, 2021). While opioid activation of opioid receptors is well known to contribute to reward and reinforcement, recent data suggest that opioid activation of TLR4 also plays a role in addiction-like processes (Hutchinson et al., 2012; Wu & Li, 2020; Zare et al., 2023) via expression of pro-inflammatory cytokines TNFα and IL-1β (Pascual et al., 2012; Rizzo et al., 2018; Xiaohui Wang et al., 2012). In the nervous system, TLR4 expression is primarily expressed in microglia, the resident immune cells of the central nervous system (Zhang et al., 2014; Zhang et al., 2016). Microglial phagocytosis is important for developmental, homeostatic, and pathological processes (Fu et al., 2014; Paolicelli et al., 2022). Thus, understanding the diverse actions of opioids via these different receptors, is important to develop strategies to mitigate their negative consequences.

In peripheral immune cells, morphine reproducibly decreases immune activation and phagocytosis *in vivo* and *in vitro* (Eisenstein, 2019; Ninkovic et al., 2016; Ninković & Roy, 2012; Rojavin et al., 1993) dependent on concentration (Borman et al., 2009) and duration of exposure (Pagán et al., 2001). Unlike peripheral immune cells, the effect of morphine on microglia is less well studied. Morphine has been reported to both increase and decrease microglial phagocytosis (Lipovsky et al., 1998; Peterson et al., 1995; Ryu et al., 2018; G. Sowa et al., 1997). There is no consensus on the modulatory effect of morphine on microglia phagocytosis currently. Several different factors contribute to microglial phagocytic activity, including sex, brain region, and regional microglial density. For example, during neonatal development microglia from female mice are more phagocytically active in the hippocampus (Nelson et al., 2017) and mature faster (Hanamsagar et al., 2017) than microglia from male mice. Morphine increases the number of activated microglia in the female rat periaqueductal gray compared to male rats which correlates to morphine analgesia (Doyle et al., 2017). Morphine also leads to sex specific microglia polarization in the periaqueductal gray (Doyle et al., 2017). There also appears to be heterogeneity in the transcriptional profile of cortical and striatal microglia (De Biase et al., 2017; Grabert et al., 2016). Deep single cell RNA sequencing reveals that microglia transcriptomes are heterogenous (Li et al., 2019) indicating features that may underlie microglia variability across brain regions (H. S. Yang et al., 2021). This heterogeneity suggests that morphine results in different responses depending on the transcriptional identity of the microglia which are spatially regulated (Coffey et al., 2022) potentially modulating the

normal tendency to phagocytose. Finally, the *in vivo* density of microglia varies depending on
brain region with the highest density in the substantia nigra (Lawson et al., 1990; Yang et al.,
2013). A higher number of microglia in a region correlates with increased maladaptive
neuroinflammation and/or microglial 'states' (Bollinger et al., 2017; Furube et al., 2018; Tynan
et al., 2010). Morphine decreases the number of microglia in the substantia nigra (Jokinen et al.,
2018).
99
We hypothesized that morphine would modulate phagocytic activity in microglia *in vitro*, but in
a culture density-, concentration-, region-, and sex-dependent manner. Specifically, we predicted
microglia would be more phagocytically active, and thus less responsive to the suppressive
effects of morphine if they are (1) plated at higher densities relative to lower densities, (2) from
striatum relative to cortex, and (3) from females relative to males. To test this, we isolated
microglia from adult male and female rat cortex and striatum and plated them *in vitro* at a
density of 10,000 or 50,000 cells (10K or 50K hereafter). Microglia were incubated with neutral
fluorescent microbeads to stimulate phagocytosis in the presence of one of four morphine
concentrations (**Fig. 1**). Unexpectedly, we found that we could not directly compare cortical and
striatal microglia, nor the two culture densities, as region and density, but not morphine
concentration, impacted cell survival over the course of the experiment. When we separated
region and density conditions, we observed that 10^{-12}M morphine, our lowest tested
concentration, increased phagocytic activity in 10K striatal, 50K striatal, and 50K cortical cells,
but not 10K cortical cells. The activating effect of 10^{-12}M morphine on 10K striatal, but not 10K
cortical cells, was replicated in separate studies, and we further provide evidence that TLR4s,
and not µORs is the primary receptor mediating this effect. Overall, our data suggest that a
morphine-TLR4 signaling pathway increases phagocytic activity in microglia independent of
sex, which is useful information for better understanding the possible neural outcomes associated
with morphine exposures.
119
MATERIALS AND METHODS
Animals
Young adult (postnatal day (P)70-P90) male and female Sprague Dawley rats were purchased
(Envigo) and group-housed with ad libitum access to food and water for at least 2 weeks.
Conditions were kept at 23°C on a 12:12 light:dark cycle (lights on at 07:00) and cages were
changed twice a week. Experiments and animal care were approved by the Institutional Animal
Care and Use Committee at Albany Medical College.
127
Tissue collection and microglial isolations
Animals were euthanized by CO_2 anesthesia and exsanguination. Saline perfusion was performed
before animals were decapitated and brains extracted. Each brain had cortex and striatum regions
grossly dissected and separated for mechanical and enzymatic breakdown. Brain regions were
minced using a razor into a gelatinous consistency while on a petri dish resting on wet ice.
Minced brain tissue underwent enzymatic dissociation using the Neural Tissue Dissociation Kit
(P) protocol (Miltenyi Biotec, 130-092-628) to isolate neural cells. Neural cells were incubated
with CD11b+ magnetic microbeads (Miltenyi Biotec, CD11b/c Microbeads, rat, 130-105-634)
and CD11b+ cells positively selected through a magnetic column to enrich for microglial cells.
We will refer to CD11b+ cells hereafter as microglia. Microglia comprised on average 28.5% of
live cells in pre-isolation samples, and 72.3% of live cells after CD11b+ isolation

(**Supplemental Fig. 1**).
140

Culture plating
Microglia were incubated in media (Dubelco's Modified Eagle's Medium with 1µM forskolin
and 1% N2 media, 1% L-glutamine, 1% sodium pyruvate, 1% Pen-Strep by volume (v/v)) at
37 °C/5% CO_2 in chambered slides (Thermo Fisher Scientific Nunc Lab-Tek II CC^2 Chamber
Slide, Z734853) at either 10K microglia (10,000 microglia/well or 50,000 microglia/mL) and
50K microglia (50,000 microglia/well or 250,000 microglia/mL) density. Microglia were
counted using trypan blue (live-dead assay) on an automated cell counter (TC20 Automated Cell
counter, Biorad, 508BR08836).
149

150 **Cell survival analysis**
151 Isolated microglia were counted for pre-incubation survival. 10K and 50K microglia were plated
152 in a 96 well plate and incubated at 37⁰C for 2hrs in 200µL media. Morphine sulfate reconstituted
153 in 100µL was added at 10^{-12} M, 10^{-8} M and 10^{-4} M to samples and incubated for 90 mins (final
154 volume 300µL). Microglia were counted for post-incubation survival.
155

Flow cytometry analysis of enrichment
Neural cells were divided into two fractions which we refer to as either pre-enriched or enriched.
Enriched cells underwent magnetic microbead separation to isolate CD11b+ cells. CD11b-
enriched and pre-enriched cells were stained with Zombie Red viability dye (BioLegend Cat. No.
423110) and the surface markers CD11b/c (Thermofisher Scientific Cat. No. 12-011082) and
CD45 (Thermofisher Scientific Cat. No. 17-0461-82). Samples were acquired on a Northern
Lights Spectral Flow Cytometer (Cytek Biosciences, California, US). Data were analyzed using
FlowJo (BD, NJ, US) and statistical analysis was performed using Graphpad Prism 9.5.0 (paired
t-test. $p < 0.05$ was defined as statistically significant.
165

Morphine-fluorescent microbead addition
Microglia were permitted to adhere to the chambered slides for 2hrs in 200µL media/well.
Fluorescent microbeads (Polysciences Inc., Fluoresbrite Carboxylate YG 1.0-micron
microspheres, 15702-10) and morphine sulfate (Mallinckrodt, H11662) at 0 Molar (M), 10^{-12} M,
10^{-8} M and 10^{-4} M were reconstituted in 100µL fresh media/well and was added to seeded
microglia (final volume 300µL) simultaneously. This range of morphine concentration is similar
the range of concentrations previously used *in vitro* to study microglia phagocytosis $\sim 10^{-6} - 10^{-20}$
M (Lipovsky et al., 1998; Peterson et al., 1995; Ryu et al., 2018; G. Sowa et al., 1997). The total
amount of microbeads per well is $\sim 1.0 \times 10^8$ microbeads/well. Microglia were incubated at 37⁰C
for an additional 90 mins, which we have published is sufficient for phagocytic activity to occur
(Kopec et al., 2018).
177

Agonist-antagonist pilot assays
Isolated microglia were permitted to adhere to the chambered slides for 1.5hrs in 100 µL
media/well. Naltrexone hydrochloride (Sigma-Aldrich, Cat. No. N3136) at 0, 50, 100 and 200
µM or *Rhodobacter sphaeroides* lipopolysaccharide (RS-LPS, InvivoGen, Cat. No. tlrl-prslps) at
0, 1, 5, 10 µg/mL were reconstituted in 100 µL fresh media/well and were added to seeded
microglia (volume 200 µL) for 30 mins. (D-Ala(2)-mephe(4)-gly-ol(5))enkephalin (DAMGO,
Sigma-Aldrich, Cat. No. E7384) at 50 µM or lipopolysaccharide (LPS, Sigma-Aldrich, Cat. No.

L4391) at 1 µg/mL was reconstituted with fluorescent microbeads in 100 µL fresh media/well
and were added to naltrexone and RS-LPS pre-incubated microglia respectively (final volume
300 µL) for 1.5hrs.
188
Inhibitor-morphine-bead addition
Isolated microglia were permitted to adhere to the chambered slides for 1.5hrs in 100µL
media/well. Naltrexone at 100µM or *Rhodobacter sphaeroides* lipopolysaccharide (RS-LPS) at
1µg/mL were reconstituted in 100µL fresh media/well and were added to seeded microglia
(volume 200µL) for 30mins. Fluorescent microbeads and morphine sulfate at 10^{-12} M were
reconstituted in 100µL fresh media/well and were added to naltrexone and RS-LPS pre-
incubated microglia (final volume 300µL) for 1.5hrs.
196
Immunocytochemistry
Microglia were rinsed with PBS (2x washes, 300µL/well), then fixed with paraformaldehyde
(300 µL/well PFA; 4%) for 20mins. Microglia were permeabilized and blocked using 0.3%
Triton x-100 and 10% donkey serum in PBS (v/v), and incubated with primary antibody anti-
rabbit Iba1 (1:500; Wako; 019-19741) overnight at 4ºC. The following day, microglia were
rinsed with PBS (3x washes, 300µL/well) and incubated with secondary antibody 1:500 Alexa
Fluor donkey anti-rabbit 568 (ThermoFisher, A21202) at room temperature for 2hrs in the dark.
Microglia were rinsed with PBS (3x washes, 300µL/well) and chambered slide well walls were
removed. Glass slides were dried overnight in the dark. Slides were coverslipped using mounting
agent (Invitrogen, Prolong Glass Antifade Mountant, P36984) the following day.
207
Imaging and Data Acquisition
2-6 images per condition were captured on a ZEISS LSM 880 confocal microscope at 20x
magnification to capture at least 20 Iba1+ structures per sample. The imager was blinded to the
animal, region, and sex of each sample. Morphine concentration and density could not be blinded
due to uniform plating maps to reduce experimental error. 20µm slice z-stacks, 0.5µm step size,
were taken of Texas Red (Iba1+ microglia) and GFP (microbeads) channels. Z-stacks were
maximum projected into 2D images with ImageJ (National Institutes of Health; Bethesda, MD).
Channels were separated and the microglia channel was manually outlined in the absence of
microbead signal. Outlines were drawn by experimenters blinded to the experimental conditions.
After all microglia in the image were outlined, cell outlines were overlayed onto the microbead
channel and mean gray value per Iba1+ area (µm) was measured. Some Iba1+ structures
contained many microbeads, some contained none. A single average value of all GFP+ pixels
within the bounds of the Iba1+ areas was used for each condition. A representative image of the
phagocytic diversity can be found in **Fig. 1B**.
222
Statistical analysis
The endpoint that was statistically analyzed was the amount of phagocytosed beads measured as
bead fluorescent intensity within microglia, which represents microbead internalization. Outliers
were identified with log transformed data due to heteroskedasticity in Graphpad 9.4.1 using the
ROUT method (Q=1%); a single outlier was removed from the data presented in Fig. 4, which is
indicated in the figure legend. Statistics were analyzed with SPSS 28.010 or GraphPad 9.4.1 with
log transformed data due to heteroskedasticity. Either two-tailed paired *t*-tests or one-, two-, or
three-way mixed model ANOVAs (random factor: animal) were performed depending on the

experimental design. Statistically significant *F* values (*p*<0.05) were followed by Tukey's post
hoc tests. Figures were created in Graphpad 9.4.1. Within-group individual differences were
evident, thus for ease of visual interpretation we depict data that is normalized to a within-animal
condition described in each figure legend. Data analysis was performed on raw, non-normalized
data.
236
RESULTS
Region and culture density, but not morphine exposure, influence the survival of microglia
in vitro
Our goal was to conduct a four-variable study directly comparing sex, region, density, and
morphine concentration on microglial phagocytic outcomes. First, to ensure the selected
morphine concentrations would not increase cell death, we measured live cells prior to plating
versus at the conclusion of the 3.5 hr experimental protocol (2 hrs acclimation in media, 1.5 hrs
morphine exposure; **Fig. 2**). Data were analyzed with a three-way ANOVA (drug x region x
density). We unexpectedly found that while morphine did not impact cell survival (main effect of
drug: $F_{3,41}$=0.692, *p*=0.562), there were main effects of both region and density (region:
$F_{1,41}$=4.608, *p*=0.038; density: $F_{1,41}$=124.885, *p*<0.001). Specifically, microglia from striatum
survived better than microglia from cortex, and microglia plated at 50K densities survived better
than microglia plated at 10K densities. There was no significant interaction between variables in
cell survival. Although we were not powered for sex in these studies, we did perform the four-
way ANOVA adding sex as a variable to ensure we were not missing a large effect, which was
indeed absent ($F_{1,2.013}$=0.390, *p*=0.596).
253
10^{-12}M morphine increases phagocytic activity in cortical microglia plated at 50K, but not
10K cells per well, and in striatal microglia at either density, independent of sex
We could not directly compare cortex and striatum, or 10K and 50K densities due to their
differential regulation of microglial survival *in vitro* (**Fig. 2**). Thus, we next sought to determine
whether sex or morphine concentration would play a role in phagocytic activity in 10K cortex,
10K striatum, 50K cortex, and 50K striatum conditions. Microglia in each condition, and from
male or female rats, were concurrently incubated with fluorescent microbeads and either 0M, 10^{-12}M, 10^{-8}M, or 10^{-4}M concentrations of morphine for 1.5 hrs after the 2hr media acclimation. Data
were analyzed with a two-way ANOVA (drug x sex). There was a main effect of drug in 10K
striatum ($F_{3,37}$=3.550, *p*=0.024), 50K cortex ($F_{3,38}$=3.481, *p*=0.025), and 50K striatum groups
($F_{3,39}$=3.102, *p*=0.038), but not in 10K cortex groups ($F_{3,33}$=1.325, *p*=0.283; **Fig. 3**). Specifically,
10^{-12}M morphine increased phagocytic activity relative to all other concentrations in 10K
Striatum and 50K striatum groups and increased phagocytic activity relative to 0M and 10^{-4}M
morphine in 50K cortex groups. There was no significant effect of sex, nor a significant
interaction between sex and drug concentration, in any condition.
269
Morphine-induced increase in phagocytic activity in striatal microglia plated at 10K
density is TLR4-, but not µOR-dependent.
We next wanted to determine whether 10^{-12}M morphine was increasing phagocytosis via TLR4-
or µOR-dependent mechanisms. Pilot studies using DAMGO as an opioid receptor agonist and
naltrexone as a µOR antagonist indicated that 100µM naltrexone would be effective in
precluding µOR-dependent changes in microglial phagocytic activity (**Supplemental Fig. 2A**).
Pilot studies using LPS as a TLR4 agonist and RS-LPS as a TLR4 antagonist indicated that

1µg/mL RS-LPS would be effective in precluding TLR4-dependent changes in microglial
phagocytic activity (**Supplemental Fig. 2B**). Thus, we incubated cortical and striatal microglia
plated at 10K densities with either 100µM naltrexone (or vehicle), 1µg/mL RS-LPS (or vehicle),
the combined inhibitors, or combined vehicles for 30 mins. Then, we added a solution containing
Fluorescent microbeads, 10^{-12}M morphine (final concentration), and the respective inhibitors to
maintain their final concentrations, for an additional 90 mins. There were 8 total treatment
groups, and because we found no significant sex difference in the effects of morphine on
phagocytosis in the studies depicted in **Fig. 3**, we combined the data from male and female rats.
One-way ANOVAs indicated there was no significant difference between Vehicle conditions in
either 10K Cortex ($F_{3,42}$=1.475, p=0.235) or 10K Striatum ($F_{3,43}$=2.382, p=0.083) microglia, and
thus we used a single averaged vehicle value for each animal for data analysis. Paired t-tests
from within-animal vehicle and 10^{-12}M morphine + inhibitor vehicle data indicated that the data
in **Fig. 3** did replicate: we found no significant effect of morphine on phagocytosis in 10K
Cortex conditions (t(14)=1.669, p=0.117), but morphine increased phagocytosis in 10K Striatum
conditions (t(15)=2.301, p=0.0362). A one-way ANOVA indicated there was an effect of
inhibitor on morphine-induced phagocytosis ($F_{3,43}$=7.222, p<0.001). Subsequent Tukey's
posthoc tests indicated the morphine-induced increase in phagocytosis in 10K Striatum microglia
was inhibited by RS-LPS (p=0.002) or RS-LPS+naltrexone (p<0.001), but not by naltrexone
alone (p=0.204). To confirm that sex did not play a role in these data, we also performed a two-
way ANOVA (inhibitor x sex), and sex was not found to play a statistically significant role
independently ($F_{1,14.026}$=3.085, p=0.101) or via interaction with inhibitors ($F_{3,40}$=2.066, p=0.120).

DISCUSSION
Herein, we demonstrate that the brain region from which microglia are isolated and plating
density both impact cell survival *in vitro* (**Fig. 2**). We found that 10^{-12}M morphine, but not higher
concentrations, increases phagocytosis in striatal microglia *in vitro* independent of sex and
plating density (**Fig. 3B, D**). 10^{-12}M morphine also increased phagocytosis in cortical microglia
in vitro independent of sex, but this was contingent on a plating density of 50K (versus 10K; **Fig.
3A, C**). This effect was reliable, as an independent replication of the effect of 10^{-12}M morphine
on 10K cortical and striatal microglia also showed no change in phagocytosis, and increased
phagocytosis, respectively (**Fig. 4A**). Finally, we demonstrate that the effect of 10^{-12}M morphine
in 10K striatal microglia is mediated via TLR4, and not µORs (**Fig. 4B**).

We initially set out to directly compare the effects of culture density, region, sex, and morphine
concentration, as well as their interactions, on microglial phagocytic activity. However, several
unexpected findings limited our ability to directly compare the effects of these variables
concurrently. We predicted that a higher culture density (50K) would increase phagocytic
activity relative to lower (10K) culture density. However, we found that microglia plated at 10K
densities did not survive as well as microglia plated at 50K densities, and thus it was not
appropriate to make this direct statistical comparison (**Fig. 2**). This is important to consider when
reconciling the literature, which include highly variable culture densities with density rarely
included as a variable. In two reports in which microglial phagocytosis is increased by morphine,
the plating density was not reported in one (as far as we could find) and was 30,000 cells/well in
chambered slides (Lipovsky et al., 1998; Peterson et al., 1995). In two reports in which
microglial phagocytosis was decreased by morphine, the plating density was 50,000 cells/ml and
40,000 cells/well (Ryu et al., 2018; G. Sowa et al., 1997). Although we observed that morphine

generally increased phagocytosis in 50K and 10K conditions, this did not reach statistical
significance in cortical microglia plated at 10K density. It is thus possible that cell viability,
perhaps in interaction with region examined, may play a role in these disparate conclusions.
Finally, although we used an artificially dramatic difference in microglial density to test our
hypothesis, it is worth noting that microglia density does vary across the brain, particularly in the
substantia nigra which has the highest density of microglia (Lawson et al., 1990; Mittelbronn et
al., 2001; Wang et al., 2015). Our data might suggest that a higher density of microglia might be
protective in terms of cell viability, but there are several reports indicating that substantia nigra
might be a particularly vulnerable region *because* of its high density of microglia (Flores-
Martinez et al., 2018; Milde et al., 2021; Wang et al., 2014). There is likely to be a difference
between microglial density in acute (e.g., in our experimental design) vs. chronic conditions. It
may be worth varying culture density in long-term cultures to test whether lower microglial
density has a negative effect under chronic activation or inflammatory conditions.

We predicted that striatal microglia would be more phagocytically active than cortical microglia.
Activation of microglia under normal and pathological states increases pro-inflammatory profiles
in striatum more than in cortex (Crapser et al., 2020; Klawonn et al., 2021). We observed that
striatal microglia survived better than cortical microglia, and thus a direct comparison was not
possible (**Fig. 2**). Although this was a main effect, not an interaction between density and region,
visual inspection of the data indicate that this effect is driven primarily by worse survival
outcomes in cortical microglia plated at 10K density. We propose that the beneficial effects of
higher density culture conditions are a result of inter-cellular signaling promoting survival. If this
is the case, cortical microglia may be particularly vulnerable (or striatal microglia more resilient)
to the presence or absence of these signals. There are currently no direct comparisons of cell
survival or death signaling profiles between cortical and striatal microglia. However, microglial
apoptotic regulation is linked to reduced phagocytosis and reduced responses to viral load (Jeong
et al., 2022; Liu et al., 2001). Thus, a few theories might explain the fact that morphine did not
increase phagocytosis in 10K cortical microglia (**Fig. 3A**): (1) reduced survival might bias our
current data, (2) activated apoptotic signaling might inhibit phagocytic potential, and (3)
morphine does not affect microglia in this context. In future studies it will be worthwhile to
examine higher density microglia from different regions to remove the possibility of differential
survival.

We predicted that microglia from female rats would be more phagocytically active than
microglia from male rats. To our surprise, we did not observe any significant effects of sex,
neither main effects nor interactions, in any of our experiments (**Fig. 3**). This may be due to the
rats being young adults, past the developmental plasticity that might be driving sex differences in
microglia (Lynch, 2022; Osborne et al., 2019), and prior to when aging might again cause sex
differences in microglia (Daly et al., 2022; Mela et al., 2022). If true, this raises the question:
what do embryonic or immortalized cultures represent? The general consensus is that these
longer-term culturing methods do not reflect mature cell phenotypes (He et al., 2018; Stansley et
al., 2012). That may be a distinct disadvantage depending on the questions being asked, just as
our current design is unlikely to be reflective of microglial responses at a different life stage. The
lack of a finding of a sex difference may also be due to being underpowered to detect subtle
effects of sex in a complex experimental design. For example, there may be sex x drug
concentration interactions in the 50K Cortex condition that we cannot resolve with the effect

sizes we attained (**Fig. 3C**). Finally, other pharmacological activators of phagocytosis such as
lipopolysaccharide result in sex differences (Loram et al., 2012; Thion et al., 2018), and so does
morphine in peripheral immune cells (Doyle & Murphy, 2017; Rosen et al., 2019). There are
reported sex differences in the response to opioids in humans and rodent models (Bobzean et al.,
2019; Boorman & Keay, 2022; Escorial et al., 2022), but microglial phagocytosis need not
mediate those effects.
375
We hypothesized that morphine would decrease phagocytosis in a dose-dependent manner.
Morphine penetrance into the brain is dependent on concentration, route of administration,
metabolism, and inter-individual factors (Chaves et al., 2017; Ollikainen et al., 2019; Rogers et
al., 2013; Smith, 2009). The statistically significant main effects indicate that 10^{-12}M morphine
increased phagocytosis in 10K Striatum, 50K Cortex, and 50K Striatum conditions (**Fig. 3B-D**).
There was no significant effect of morphine in 10K Cortex conditions (**Fig. 3A**). It is possible
that lower survival in this condition is resulting to the null effect, but visual inspection of the data
indicates that story is likely to be more complicated. While there is no statistically significant
effect of morphine on cortical microglia plated at 10K density, there is a rather clear pattern of
10^{-12}M morphine increasing phagocytosis relative to other concentrations – in females only. This
effect is likely masked (from a statistical analysis perspective) by high variance in male data that
does not follow the same trend. Larger group sizes may permit more nuanced conclusions. We
find that at lower morphine concentrations phagocytosis is increased. This suggests that there is a
window of increased microglia phagocytosis *in vivo* which could be immediate or delayed based
on peak brain morphine concentration as the concentration decays with time. Importantly,
whether there is a sex difference in the effect of morphine on microglial phagocytosis or not, it
does appear that morphine is generally increasing phagocytosis. This could be consequential for
the effect of morphine on synaptic density, as synapses are a frequent target of microglial
phagocytosis (Authement et al., 2016; Kopec et al., 2018; Zhang et al., 2021). In fact, there is
reports that synaptic density is decreased in nucleus accumbens and intermediate medial
interstitial after morphine exposures (Kasture et al., 2009; Wang et al., 2023). An open question
is whether microglia would be a mediator of that effect.
398
Morphine is documented to activate both TLR4 and µOR signaling to mediate its effects. Thus,
in our final set of studies, we tested whether increased phagocytosis in 10^{-12}M morphine treated
striatal microglia plated at 10K density required TLR4s, µORs, or both. The data indicate that
TLR4, but not µORs, is important for this effect (**Fig. 4**). While the expression of TLR4 is
enriched in microglia in neural cell datasets, there has been some controversy over whether
microglia express µORs. The morphine binding affinity for µOR (Human) has been measured at
~1.14nM (Volpe et al., 2011) and morphine levels in the brain peak in the 10-500 ng/ml range
depending on the morphine metabolite measured (Barjavel et al., 1994; Barjavel et al., 1995;
Ederoth et al., 2004; Stain-Texier et al., 1999). However, morphine has been shown to modulate
microglia in femtomolar (10^{-15}) ranges (Grzegorz Sowa et al., 1997). While morphine has been
shown to bind to TLR4 in a parallel manner to its well-known agonist endotoxin
(lipopolysaccharide, specifically) (X. Wang et al., 2012), the binding affinity of morphine for
TLR4 is an ongoing endeavor (J. H. L. Thomas et al., 2022). However, in peripheral
macrophages TLR4 potentiates phagocytosis more robustly than opioid receptors (Ninkovic et

al., 2016) and sex-specifically influences antinociception, tolerance and withdrawal (Jacob H. L. Thomas et al., 2022). Both opioid receptors and TLR4s are reported to influence the effects of morphine on microglial immune signaling (Liang et al., 2016; Lipovsky et al., 1998; X. Wang et al., 2012; Xie et al., 2017; Y. Yang et al., 2021). Our pilot data indicate that microglia respond to a non-specific opioid receptor agonist (DAMGO) with increased phagocytosis, and a µOR-specific antagonist (naltrexone) can inhibit this effect. These data would argue that µORs are, in fact, expressed and functional in microglia. However, this was a pilot study and underpowered for making conclusive statements. We thus cannot rule out that the null effect of naltrexone on morphine-induced phagocytosis is due to no or low expression of µORs on microglia. Regardless, our data do indicate that morphine is activating microglia through canonical immune pathways. TLR4 signaling is associated not just with phagocytic regulation, but also pro-inflammatory cytokine induction. Indeed, morphine potentiates synthesis and release of IL-1β and TNF-α in microglia (Liang et al., 2016; Y. Yang et al., 2021). Whether this would be the case in microglia from either region plated at 50K remains to be tested. It is possible that a µOR- and region-dependent effect would emerge in microglia plated at 50K due to their greater survival (and thus wider sampling), because striatum contains microglia exposed to higher endogenous opioidergic signaling (Rocchi et al., 2020).

Limitations to the current studies
The choice of experimental conditions in studies such as these is not a trivial matter (Bohlen et al., 2017). We made several experimental decisions that could impact our conclusions. Some of the major caveats include: (**1**) Our data comes from microglia from a single rat strain (Sprague-Dawley) and further studies on other rat strains and animal species are needed for generalization of these results. (**2**) Media plays an enormous role in *in vitro* outcomes. Our media does not contain serum, which is known to impact microglial health in culture, but does contain low-concentration (1µM) forskolin. We used forskolin to upregulate cAMP levels in microglia, as cAMP–regulated pathways are essential for physiological cell processes (Sassone-Corsi, 2012) which are stunted during cell isolation (Bohlen et al., 2019). Additionally, cAMP pathways are important for pro-phagocytic responses such as deramification and surveillance, hallmarks of phagocytic activity (Bernier et al., 2019; Ghosh et al., 2020). However, forskolin has been linked to both enhancement and suppression of neuroinflammation (Owona et al., 2016; Veremeyko et al., 2018; Woo et al., 2004). Opioid receptor agonists such as morphine inhibit the generation of cAMP (Zhang et al., 2020), and upregulating cAMP is protective against morphine-induced microglia consequences (He & Whistler, 2005). The 1µM concentration of forskolin used herein is lower than typical studies using forskolin (often 10µM), and there are clear phagocytic responses to morphine in striatum and to DAMGO and LPS in whole brain. Finally (**3**), any isolation protocol will impact the cells in a non-physiological way. We use dissociation techniques reported to perform well in terms of microglial health and viability (Bordt et al., 2020; Nikodemova & Watters, 2012), but there is no way to attain a true *in vivo* physiological setting *in vitro*.

<u>CONCLUSIONS</u>
Overall, our data suggest that a morphine-TLR4 signaling pathway increases phagocytic activity in microglia independent of sex, which is useful information for better understanding the

458 possible neural outcomes associated with morphine exposures.
459

460 **Acknowledgements:** We thank Dr. Justin Bourgeois and Erin Edgar for experimental and
461 quantification assistance. We thank Dr. Margarida Barroso and the AMC Imaging Core Facility
462 for imaging assistance.
463

Funding: This work was supported by the National Institutes of Health [R01DA052889 and
R03AG07011 to AMK and R35GM131842 to KCM] and Albany Medical College Start-up
funds to AMK.

Author Declarations: DNK, LN, JLB, KCM, and AMK designed the experiments. DNK, LN,
JLB, and AMK performed the experiments. DNK, LN, JLB, PJF, and AMK analyzed the
experiments. DNK, MT, and AMK wrote the manuscript. All authors edited the manuscript. The
authors declare no conflicts of interest.

498   **FIGURES**

Figure 1: Experimental Design
(A) The experimental design included sex (male vs. female), brain region (cortex vs. striatum),
density (10,000 microglia/well (10K microglia) vs. 50,000 microglia/well (50K microglia), and
morphine concentration (0, 10^{-12}, 10^{-8}, and 10^{-4} M) as independent variables. Created with
Biorender.com. **(B)** Representative image of Iba1+ microglia (magenta), fluorescent microbeads
(green), and colocalization of microglia and fluorescent microbeads (white) which we refer to as
phagocytosis. Image is representative of all conditions.

Figure 2. Region and culture density, but not morphine exposure, influence the survival of microglia *in vitro*

The percent change in Trypan-Blue indicated live cells from pre-plating to post-experiment was calculated for striatal and cortical microglia plated at 10K or 50K densities and exposed to different concentrations of morphine for 90 mins. $n = 3\text{-}4$ rats (sexes combined). Circles depict average, vertical lines depict standard error of the mean. A three-way mixed model ANOVA revealed region and density, but not drug or interactions between variables, impact microglial survival. * indicates the main effect of density (50K > 10K). *a* and *b* indicate the main effect of region (Striatum > Cortex). Main effects were considered significant if $p<0.05$.

Figure 3: 10^{-12}M morphine increases phagocytic activity in cortical microglia plated at 50K, but not 10K cells per well, and in striatal microglia at either density, independent of sex

(**A**) There was no effect of morphine in cortical microglia plated at 10K density, while (**B**) 10^{-12}M morphine increased phagocytic activity relative to all other concentrations in striatal microglia plated at 10K density. 10^{-12}M morphine also increased phagocytic activity in both (**C**) cortical and (**D**) striatal microglia plated at 50K density (cortical: relative to 0M and 10^{-4}M, but not 10^{-8}M concentrations; striatal: relative to all other concentrations). Sex did not regulate phagocytic activity independently or via interactions with morphine concentration in any context. Circles depict mean, vertical lines depict standard error of the mean. Data were normalized to within-animal 0M conditions for ease of visualization. Data above the gray dotted line would indicate an increase in phagocytic activity above 0M control levels. * = $p < 0.05$; n = 6-8 rats/sex.

A) 10⁻¹² M morphine phagocytic activation

B) Receptor contribution in 10K Striatum condition

566
Figure 4: Morphine-induced increase in phagocytic activity in striatal microglia plated at
10K density is TLR4-, but not μOR-dependent.
(A) A separate set of studies replicated the increased phagocytic effect of 10^{-12}M morphine on
striatal, but not cortical microglia plated at 10K density. Data are normalized to within-animal
vehicle conditions for ease of visualization. Data above the gray dotted line would indicate an
increase in phagocytic activity above 0M control levels and are identified with orange circles;
data below the gray dotted line would indicate a decrease in phagocytic activity and are
identified with blue circles. * = $p<0.05$; n = 15-16 rats (sexes combined). **(B)** Inhibiting TLR4s
with RS-LPS, or TLR4s and μORs with a combination of RS-LPS and naltrexone, but not μORs
(naltrexone) alone, inhibited the effect of 10^{-12}M morphine on phagocytosis in striatal microglia
plated at 10K density. One morphine+naltrexone+RS-LPS- sample was identified as an outlier
and removed from analysis. Data were normalized to within-animal vehicle conditions for ease
of visualization. Data above the gray dotted line would indicate an increase in phagocytic

activity above 0M control levels. Circles depict mean, vertical lines depict standard error of the
mean. * = $p<0.05$; n = 15-16 rats (sexes combined).

Supp. Figure 1: CD11b magnetic isolation enriches for microglia.
Microglia were enriched from single-cell suspension of rat brain tissue using Miltenyi Biotec
MicroBeads. (A) Cells were stained with antibodies against CD11b/c and CD45 and analyzed via
flow cytometry. (B) The frequency of microglia was compared before and after magnetic
microbead enrichment. Data were analyzed with a paired *t*-test. *** = p<0.001; *n*=4 rats (sexes
combined).

A) 50μM DAMGO

Fluorobead pixels per Iba1+ μm² (Normalized to Control)

Naltrexone (μM)

B) 1μg/mL LPS

Fluorobead pixels per Iba1+ μm² (Normalized to Control)

RS-LPS (μg/mL)

Supp. Figure 2: 100μM naltrexone and 1μg/mL RS-LPS reduce DAMGO- and LPS-induced phagocytic activity, respectively

(A) Microglia from whole brain were plated at 50K density and exposed to fluorescent microbeads, Vehicle or 50μM DAMGO, an opioid receptor agonist, and varying concentrations of naltrexone, a μOR antagonist. 100μM naltrexone reduced phagocytosis induced by DAMGO to control levels. **(B)** Microglia from whole brain were plated at 50K density and exposed to fluorescent microbeads, Vehicle or 1μg/mL LPS, a TLR4 agonist, and varying concentrations of RS-LPS, a TLR4 antagonist. 1μg/mL RS-LPS reduced phagocytosis induced by LPS to control levels. Data were normalized to within-animal vehicle conditions (an average of 0M agonist + each inhibitor concentration) for ease of visualization. Data above the gray dotted line would indicate an increase in phagocytic activity above control levels. Circles depict mean, vertical lines depict standard error of the mean. n = 3-4 rats (sexes combined). Statistics were not performed as this was a pilot experiment to choose inhibitor concentrations for the experiment displayed in Figure 5.

REFERENCES

Al-Hasani, R., & Bruchas, M. R. (2011, Dec). Molecular mechanisms of opioid receptor-dependent signaling and behavior. *Anesthesiology, 115*(6), 1363-1381. https://doi.org/10.1097/ALN.0b013e318238bba6

Andoh, M., & Koyama, R. (2021, Sep 27). Comparative Review of Microglia and Monocytes in CNS Phagocytosis. *Cells, 10*(10). https://doi.org/10.3390/cells10102555

Authement, M. E., Langlois, L. D., Kassis, H., Gouty, S., Dacher, M., Shepard, R. D., Cox, B. M., & Nugent, F. S. (2016, Sep 1). Morphine-induced synaptic plasticity in the VTA is reversed by HDAC inhibition. *J Neurophysiol, 116*(3), 1093-1103. https://doi.org/10.1152/jn.00238.2016

Barjavel, M., Sandouk, P., Plotkine, M., & Scherrmann, J. M. (1994). Morphine and morphine metabolite kinetics in the rat brain as assessed by transcortical microdialysis. *Life Sci, 55*(16), 1301-1308. https://doi.org/10.1016/0024-3205(94)90069-8

Barjavel, M. J., Scherrmann, J. M., & Bhargava, H. N. (1995, Dec). Relationship between morphine analgesia and cortical extracellular fluid levels of morphine and its metabolites in the rat: a microdialysis study. *Br J Pharmacol, 116*(8), 3205-3210. https://doi.org/10.1111/j.1476-5381.1995.tb15125.x

Bernier, L. P., Bohlen, C. J., York, E. M., Choi, H. B., Kamyabi, A., Dissing-Olesen, L., Hefendehl, J. K., Collins, H. Y., Stevens, B., Barres, B. A., & MacVicar, B. A. (2019, Jun 4). Nanoscale Surveillance of the Brain by Microglia via cAMP-Regulated Filopodia. *Cell Rep, 27*(10), 2895-2908.e2894. https://doi.org/10.1016/j.celrep.2019.05.010

Bobzean, S. A. M., Kokane, S. S., Butler, B. D., & Perrotti, L. I. (2019, Jul 13). Sex differences in the expression of morphine withdrawal symptoms and associated activity in the tail of the ventral tegmental area. *Neurosci Lett, 705*, 124-130. https://doi.org/10.1016/j.neulet.2019.04.057

Bohlen, C. J., Bennett, F. C., & Bennett, M. L. (2019, Jun). Isolation and Culture of Microglia. *Curr Protoc Immunol, 125*(1), e70. https://doi.org/10.1002/cpim.70

Bohlen, C. J., Bennett, F. C., Tucker, A. F., Collins, H. Y., Mulinyawe, S. B., & Barres, B. A. (2017, May 17). Diverse Requirements for Microglial Survival, Specification, and Function Revealed by Defined-Medium Cultures. *Neuron, 94*(4), 759-773.e758. https://doi.org/10.1016/j.neuron.2017.04.043

Bollinger, J. L., Collins, K. E., Patel, R., & Wellman, C. L. (2017). Behavioral stress alters corticolimbic microglia in a sex- and brain region-specific manner. *PLoS One, 12*(12), e0187631. https://doi.org/10.1371/journal.pone.0187631

715   Boorman, D. C., & Keay, K. A. (2022, Oct). Sex differences in morphine sensitivity are associated with
716   differential glial expression in the brainstem of rats with neuropathic pain. *J Neurosci Res,*
717   *100*(10), 1890-1907. https://doi.org/10.1002/jnr.25103

719   Bordt, E. A., Block, C. L., Petrozziello, T., Sadri-Vakili, G., Smith, C. J., Edlow, A. G., & Bilbo, S. D. (2020, Jun
720   19). Isolation of Microglia from Mouse or Human Tissue. *STAR Protoc, 1*(1).
721   https://doi.org/10.1016/j.xpro.2020.100035

723   Borman, A., Ciepielewski, Z., Wrona, D., Stojek, W., Glac, W., Leszkowicz, E., & Tokarski, J. (2009, Mar).
724   Small doses of morphine can enhance NK cell cytotoxicity in pigs. *Int Immunopharmacol, 9*(3),
725   277-283. https://doi.org/10.1016/j.intimp.2008.11.006

727   Boué, J., Blanpied, C., Brousset, P., Vergnolle, N., & Dietrich, G. (2011, May 1). Endogenous opioid-
728   mediated analgesia is dependent on adaptive T cell response in mice. *J Immunol, 186*(9), 5078-
729   5084. https://doi.org/10.4049/jimmunol.1003335

731   Chaves, C., Remiao, F., Cisternino, S., & Decleves, X. (2017, Nov 14). Opioids and the Blood-Brain Barrier:
732   A Dynamic Interaction with Consequences on Drug Disposition in Brain. *Curr Neuropharmacol,*
733   *15*(8), 1156-1173. https://doi.org/10.2174/1570159x15666170504095823

735   Coffey, K. R., Lesiak, A. J., Marx, R. G., Vo, E. K., Garden, G. A., & Neumaier, J. F. (2022, Apr). A cAMP-
736   Related Gene Network in Microglia Is Inversely Regulated by Morphine Tolerance and
737   Withdrawal. *Biol Psychiatry Glob Open Sci, 2*(2), 180-189.
738   https://doi.org/10.1016/j.bpsgos.2021.07.011

740   Crapser, J. D., Ochaba, J., Soni, N., Reidling, J. C., Thompson, L. M., & Green, K. N. (2020, Jan 1).
741   Microglial depletion prevents extracellular matrix changes and striatal volume reduction in a
742   model of Huntington's disease. *Brain, 143*(1), 266-288. https://doi.org/10.1093/brain/awz363

744   Daly, C. M., Saxena, J., Singh, J., Bullard, M. R., Bondy, E. O., Saxena, A., Buffalino, R. E., Melville, M. F., &
745   Freeman, L. R. (2022, Feb). Sex differences in response to a high fat, high sucrose diet in both
746   the gut microbiome and hypothalamic astrocytes and microglia. *Nutr Neurosci, 25*(2), 321-335.
747   https://doi.org/10.1080/1028415x.2020.1752996

749   De Biase, L. M., Schuebel, K. E., Fusfeld, Z. H., Jair, K., Hawes, I. A., Cimbro, R., Zhang, H. Y., Liu, Q. R.,
750   Shen, H., Xi, Z. X., Goldman, D., & Bonci, A. (2017, Jul 19). Local Cues Establish and Maintain
751   Region-Specific Phenotypes of Basal Ganglia Microglia. *Neuron, 95*(2), 341-356.e346.
752   https://doi.org/10.1016/j.neuron.2017.06.020

754   Doyle, H. H., Eidson, L. N., Sinkiewicz, D. M., & Murphy, A. Z. (2017, Mar 22). Sex Differences in Microglia
755   Activity within the Periaqueductal Gray of the Rat: A Potential Mechanism Driving the Dimorphic

Effects of Morphine. *J Neurosci, 37*(12), 3202-3214. https://doi.org/10.1523/JNEUROSCI.2906-16.2017

Doyle, H. H., & Murphy, A. Z. (2017, Jan 2). Sex differences in innate immunity and its impact on opioid pharmacology. *J Neurosci Res, 95*(1-2), 487-499. https://doi.org/10.1002/jnr.23852

Ederoth, P., Tunblad, K., Bouw, R., Lundberg, C. J., Ungerstedt, U., Nordström, C. H., & Hammarlund-Udenaes, M. (2004, Apr). Blood-brain barrier transport of morphine in patients with severe brain trauma. *Br J Clin Pharmacol, 57*(4), 427-435. https://doi.org/10.1046/j.1365-2125.2003.02032.x

Eisenstein, T. K. (2019). The Role of Opioid Receptors in Immune System Function. *Front Immunol, 10*, 2904. https://doi.org/10.3389/fimmu.2019.02904

Escorial, M., Muriel, J., Margarit, C., Agulló, L., Morales, D., & Peiró Peiró, A. M. (2022, Sep 16). Sex-Differences in Pain and Opioid Use Disorder Management: A Cross-Sectional Real-World Study. *Biomedicines, 10*(9). https://doi.org/10.3390/biomedicines10092302

Flores-Martinez, Y. M., Fernandez-Parrilla, M. A., Ayala-Davila, J., Reyes-Corona, D., Blanco-Alvarez, V. M., Soto-Rojas, L. O., Luna-Herrera, C., Gonzalez-Barrios, J. A., Leon-Chavez, B. A., Gutierrez-Castillo, M. E., Martínez-Dávila, I. A., & Martinez-Fong, D. (2018). Acute Neuroinflammatory Response in the Substantia Nigra Pars Compacta of Rats after a Local Injection of Lipopolysaccharide. *J Immunol Res, 2018*, 1838921. https://doi.org/10.1155/2018/1838921

Fu, R., Shen, Q., Xu, P., Luo, J. J., & Tang, Y. (2014, 2014/06/01). Phagocytosis of Microglia in the Central Nervous System Diseases. *Molecular Neurobiology, 49*(3), 1422-1434. https://doi.org/10.1007/s12035-013-8620-6

Furube, E., Kawai, S., Inagaki, H., Takagi, S., & Miyata, S. (2018, Feb 2). Brain Region-dependent Heterogeneity and Dose-dependent Difference in Transient Microglia Population Increase during Lipopolysaccharide-induced Inflammation. *Sci Rep, 8*(1), 2203. https://doi.org/10.1038/s41598-018-20643-3

Ghosh, S., Stepicheva, N., Yazdankhah, M., Shang, P., Watson, A. M., Hose, S., Liu, H., Weiss, J., Zigler, J. S., Jr., Valapala, M., Watkins, S. C., & Sinha, D. (2020, Mar). The role of lipocalin-2 in age-related macular degeneration (AMD). *Cell Mol Life Sci, 77*(5), 835-851. https://doi.org/10.1007/s00018-019-03423-8

Grabert, K., Michoel, T., Karavolos, M. H., Clohisey, S., Baillie, J. K., Stevens, M. P., Freeman, T. C., Summers, K. M., & McColl, B. W. (2016, Mar). Microglial brain region-dependent diversity and selective regional sensitivities to aging. *Nat Neurosci, 19*(3), 504-516. https://doi.org/10.1038/nn.4222

Hanamsagar, R., Alter, M. D., Block, C. S., Sullivan, H., Bolton, J. L., & Bilbo, S. D. (2017, Sep). Generation
of a microglial developmental index in mice and in humans reveals a sex difference in
maturation and immune reactivity. *Glia, 65*(9), 1504-1520. https://doi.org/10.1002/glia.23176

He, L., & Whistler, J. L. (2005, Jun 7). An opiate cocktail that reduces morphine tolerance and
dependence. *Curr Biol, 15*(11), 1028-1033. https://doi.org/10.1016/j.cub.2005.04.052

He, Y., Yao, X., Taylor, N., Bai, Y., Lovenberg, T., & Bhattacharya, A. (2018, May 22). RNA sequencing
analysis reveals quiescent microglia isolation methods from postnatal mouse brains and
limitations of BV2 cells. *J Neuroinflammation, 15*(1), 153. https://doi.org/10.1186/s12974-018-
1195-4

Hutchinson, M. R., Northcutt, A. L., Hiranita, T., Wang, X., Lewis, S. S., Thomas, J., van Steeg, K., Kopajtic,
T. A., Loram, L. C., Sfregola, C., Galer, E., Miles, N. E., Bland, S. T., Amat, J., Rozeske, R. R.,
Maslanik, T., Chapman, T. R., Strand, K. A., Fleshner, M., Bachtell, R. K., Somogyi, A. A., Yin, H.,
Katz, J. L., Rice, K. C., Maier, S. F., & Watkins, L. R. (2012, Aug 15). Opioid activation of toll-like
receptor 4 contributes to drug reinforcement. *J Neurosci, 32*(33), 11187-11200.
https://doi.org/10.1523/JNEUROSCI.0684-12.2012

Jeong, G. U., Lyu, J., Kim, K. D., Chung, Y. C., Yoon, G. Y., Lee, S., Hwang, I., Shin, W. H., Ko, J., Lee, J. Y., &
Kwon, Y. C. (2022, Jun 29). SARS-CoV-2 Infection of Microglia Elicits Proinflammatory Activation
and Apoptotic Cell Death. *Microbiol Spectr, 10*(3), e0109122.
https://doi.org/10.1128/spectrum.01091-22

Jokinen, V., Sidorova, Y., Viisanen, H., Suleymanova, I., Tiilikainen, H., Li, Z., Lilius, T. O., Mätlik, K.,
Anttila, J. E., Airavaara, M., Tian, L., Rauhala, P. V., & Kalso, E. A. (2018, Apr 1). Differential Spinal
and Supraspinal Activation of Glia in a Rat Model of Morphine Tolerance. *Neuroscience, 375*, 10-
24. https://doi.org/10.1016/j.neuroscience.2018.01.048

Kasture, S., Vinci, S., Ibba, F., Puddu, A., Marongiu, M., Murali, B., Pisanu, A., Lecca, D., Zernig, G., &
Acquas, E. (2009, Nov). Withania somnifera prevents morphine withdrawal-induced decrease in
spine density in nucleus accumbens shell of rats: a confocal laser scanning microscopy study.
Neurotox Res, 16(4), 343-355. https://doi.org/10.1007/s12640-009-9069-2

Klawonn, A. M., Fritz, M., Castany, S., Pignatelli, M., Canal, C., Similä, F., Tejeda, H. A., Levinsson, J.,
Jaarola, M., Jakobsson, J., Hidalgo, J., Heilig, M., Bonci, A., & Engblom, D. (2021, Feb 9).
Microglial activation elicits a negative affective state through prostaglandin-mediated
modulation of striatal neurons. *Immunity, 54*(2), 225-234.e226.
https://doi.org/10.1016/j.immuni.2020.12.016

Kopec, A. M., Smith, C. J., Ayre, N. R., Sweat, S. C., & Bilbo, S. D. (2018, Sep 25). Microglial dopamine
receptor elimination defines sex-specific nucleus accumbens development and social behavior in
adolescent rats. *Nat Commun, 9*(1), 3769. https://doi.org/10.1038/s41467-018-06118-z

Lawson, L. J., Perry, V. H., Dri, P., & Gordon, S. (1990). Heterogeneity in the distribution and morphology
of microglia in the normal adult mouse brain. *Neuroscience, 39*(1), 151-170.
https://doi.org/10.1016/0306-4522(90)90229-w

Li, Q., Cheng, Z., Zhou, L., Darmanis, S., Neff, N. F., Okamoto, J., Gulati, G., Bennett, M. L., Sun, L. O.,
Clarke, L. E., Marschallinger, J., Yu, G., Quake, S. R., Wyss-Coray, T., & Barres, B. A. (2019, Jan
16). Developmental Heterogeneity of Microglia and Brain Myeloid Cells Revealed by Deep
Single-Cell RNA Sequencing. *Neuron, 101*(2), 207-223.e210.
https://doi.org/10.1016/j.neuron.2018.12.006

Liang, Y., Chu, H., Jiang, Y., & Yuan, L. (2016, Dec). Morphine enhances IL-1β release through toll-like
receptor 4-mediated endocytic pathway in microglia. *Purinergic Signal, 12*(4), 637-645.
https://doi.org/10.1007/s11302-016-9525-4

Lipovsky, M. M., Gekker, G., Hu, S., Hoepelman, A. I., & Peterson, P. K. (1998, May). Morphine enhances
complement receptor-mediated phagocytosis of Cryptococcus neoformans by human microglia.
Clin Immunol Immunopathol, 87(2), 163-167. https://www.ncbi.nlm.nih.gov/pubmed/9614931

Liu, B., Wang, K., Gao, H. M., Mandavilli, B., Wang, J. Y., & Hong, J. S. (2001, Apr). Molecular
consequences of activated microglia in the brain: overactivation induces apoptosis. *J
Neurochem, 77*(1), 182-189. https://doi.org/10.1046/j.1471-4159.2001.t01-1-00216.x

Loram, L. C., Sholar, P. W., Taylor, F. R., Wiesler, J. L., Babb, J. A., Strand, K. A., Berkelhammer, D., Day, H.
E., Maier, S. F., & Watkins, L. R. (2012, Oct). Sex and estradiol influence glial pro-inflammatory
responses to lipopolysaccharide in rats. *Psychoneuroendocrinology, 37*(10), 1688-1699.
https://doi.org/10.1016/j.psyneuen.2012.02.018

Lu, Y. C., Yeh, W. C., & Ohashi, P. S. (2008, May). LPS/TLR4 signal transduction pathway. *Cytokine, 42*(2),
145-151. https://doi.org/10.1016/j.cyto.2008.01.006

Lynch, M. A. (2022). Exploring Sex-Related Differences in Microglia May Be a Game-Changer in Precision
Medicine. *Front Aging Neurosci, 14*, 868448. https://doi.org/10.3389/fnagi.2022.868448

Mela, V., Sayd Gaban, A., O'Neill, E., Bechet, S., Walsh, A., & Lynch, M. A. (2022, Feb 18). The
Modulatory Effects of DMF on Microglia in Aged Mice Are Sex-Specific. *Cells, 11*(4).
https://doi.org/10.3390/cells11040729

Milde, S., van Tartwijk, F. W., Vilalta, A., Hornik, T. C., Dundee, J. M., Puigdellívol, M., & Brown, G. C.
(2021, Oct 11). Inflammatory neuronal loss in the substantia nigra induced by systemic
lipopolysaccharide is prevented by knockout of the P2Y(6) receptor in mice. *J*
Neuroinflammation, 18(1), 225. https://doi.org/10.1186/s12974-021-02280-2

884
885 Mittelbronn, M., Dietz, K., Schluesener, H. J., & Meyermann, R. (2001, Mar). Local distribution of
886   microglia in the normal adult human central nervous system differs by up to one order of
887   magnitude. *Acta Neuropathol, 101*(3), 249-255. https://doi.org/10.1007/s004010000284

888
889 Nelson, L. H., Warden, S., & Lenz, K. M. (2017, Aug). Sex differences in microglial phagocytosis in the
890   neonatal hippocampus. *Brain Behav Immun, 64*, 11-22.
891   https://doi.org/10.1016/j.bbi.2017.03.010

892
893 Nikodemova, M., & Watters, J. J. (2012, 2012/06/28). Efficient isolation of live microglia with preserved
894   phenotypes from adult mouse brain. *Journal of Neuroinflammation, 9*(1), 147.
895   https://doi.org/10.1186/1742-2094-9-147

896
897 Ninkovic, J., Anand, V., Dutta, R., Zhang, L., Saluja, A., Meng, J., Koodie, L., Banerjee, S., & Roy, S. (2016,
898   Feb 19). Differential effects of gram-positive and gram-negative bacterial products on morphine
899   induced inhibition of phagocytosis. *Sci Rep, 6*, 21094. https://doi.org/10.1038/srep21094

900
901 Ninković, J., & Roy, S. (2012, Mar). Morphine decreases bacterial phagocytosis by inhibiting actin
902   polymerization through cAMP-, Rac-1-, and p38 MAPK-dependent mechanisms. *Am J Pathol,*
903   *180*(3), 1068-1079. https://doi.org/10.1016/j.ajpath.2011.11.034

904
905 Ollikainen, E., Aitta-Aho, T., Koburg, M., Kostiainen, R., & Sikanen, T. (2019, Mar 1). Rapid analysis of
906   intraperitoneally administered morphine in mouse plasma and brain by microchip
907   electrophoresis-electrochemical detection. *Sci Rep, 9*(1), 3311. https://doi.org/10.1038/s41598-
908   019-40116-5

909
910 Osborne, B. F., Turano, A., Caulfield, J. I., & Schwarz, J. M. (2019, Jan 23). Sex- and region-specific
911   differences in microglia phenotype and characterization of the peripheral immune response
912   following early-life infection in neonatal male and female rats. *Neurosci Lett, 692*, 1-9.
913   https://doi.org/10.1016/j.neulet.2018.10.044

914
915 Owona, B. A., Zug, C., Schluesener, H. J., & Zhang, Z. Y. (2016, Jul). Protective Effects of Forskolin on
916   Behavioral Deficits and Neuropathological Changes in a Mouse Model of Cerebral Amyloidosis. *J*
917   *Neuropathol Exp Neurol, 75*(7), 618-627. https://doi.org/10.1093/jnen/nlw043

918
Pagán, W. E., Figueroa, N. Y., & Renaud, F. L. (2001). Self-enhancement of phagocytosis by murine
resident peritoneal macrophages and its relationship to morphine effects on the process. *Adv*
Exp Med Biol, 493, 111-116. https://doi.org/10.1007/0-306-47611-8_13

Paolicelli, R. C., Sierra, A., Stevens, B., Tremblay, M. E., Aguzzi, A., Ajami, B., Amit, I., Audinat, E.,
Bechmann, I., Bennett, M., Bennett, F., Bessis, A., Biber, K., Bilbo, S., Blurton-Jones, M.,
Boddeke, E., Brites, D., Brône, B., Brown, G. C., Butovsky, O., Carson, M. J., Castellano, B.,
Colonna, M., Cowley, S. A., Cunningham, C., Davalos, D., De Jager, P. L., de Strooper, B., Denes,
A., Eggen, B. J. L., Eyo, U., Galea, E., Garel, S., Ginhoux, F., Glass, C. K., Gokce, O., Gomez-Nicola,
D., González, B., Gordon, S., Graeber, M. B., Greenhalgh, A. D., Gressens, P., Greter, M.,
Gutmann, D. H., Haass, C., Heneka, M. T., Heppner, F. L., Hong, S., Hume, D. A., Jung, S.,
Kettenmann, H., Kipnis, J., Koyama, R., Lemke, G., Lynch, M., Majewska, A., Malcangio, M.,
Malm, T., Mancuso, R., Masuda, T., Matteoli, M., McColl, B. W., Miron, V. E., Molofsky, A. V.,
Monje, M., Mracsko, E., Nadjar, A., Neher, J. J., Neniskyte, U., Neumann, H., Noda, M., Peng, B.,
Peri, F., Perry, V. H., Popovich, P. G., Pridans, C., Priller, J., Prinz, M., Ragozzino, D., Ransohoff, R.
M., Salter, M. W., Schaefer, A., Schafer, D. P., Schwartz, M., Simons, M., Smith, C. J., Streit, W. J.,
Tay, T. L., Tsai, L. H., Verkhratsky, A., von Bernhardi, R., Wake, H., Wittamer, V., Wolf, S. A., Wu,
L. J., & Wyss-Coray, T. (2022, Nov 2). Microglia states and nomenclature: A field at its
crossroads. *Neuron, 110*(21), 3458-3483. https://doi.org/10.1016/j.neuron.2022.10.020

Pascual, O., Ben Achour, S., Rostaing, P., Triller, A., & Bessis, A. (2012, Jan 24). Microglia activation
triggers astrocyte-mediated modulation of excitatory neurotransmission. *Proc Natl Acad Sci U S
A, 109*(4), E197-205. https://doi.org/10.1073/pnas.1111098109

Peterson, P. K., Gekker, G., Hu, S., Sheng, W. S., Molitor, T. W., & Chao, C. C. (1995). Morphine
stimulates phagocytosis of Mycobacterium tuberculosis by human microglial cells: involvement
of a G protein-coupled opiate receptor. *Adv Neuroimmunol, 5*(3), 299-309.
https://www.ncbi.nlm.nih.gov/pubmed/8748073

Rahimifard, M., Maqbool, F., Moeini-Nodeh, S., Niaz, K., Abdollahi, M., Braidy, N., Nabavi, S. M., &
Nabavi, S. F. (2017, Jul). Targeting the TLR4 signaling pathway by polyphenols: A novel
therapeutic strategy for neuroinflammation. *Ageing Res Rev, 36*, 11-19.
https://doi.org/10.1016/j.arr.2017.02.004

Rizzo, F. R., Musella, A., De Vito, F., Fresegna, D., Bullitta, S., Vanni, V., Guadalupi, L., Stampanoni Bassi,
M., Buttari, F., Mandolesi, G., Centonze, D., & Gentile, A. (2018). Tumor Necrosis Factor and
Interleukin-1β Modulate Synaptic Plasticity during Neuroinflammation. *Neural Plast, 2018*,
8430123. https://doi.org/10.1155/2018/8430123

Rocchi, G., Sterlini, B., Tardito, S., Inglese, M., Corradi, A., Filaci, G., Amore, M., Magioncalda, P., &
Martino, M. (2020, Aug). Opioidergic System and Functional Architecture of Intrinsic Brain
Activity: Implications for Psychiatric Disorders. *Neuroscientist, 26*(4), 343-358.
https://doi.org/10.1177/1073858420902360

Rogers, E., Mehta, S., Shengelia, R., & Reid, M. C. (2013, Apr). Four Strategies for Managing Opioid-
Induced Side Effects in Older Adults. *Clin Geriatr, 21*(4).

Rojavin, M., Szabo, I., Bussiere, J. L., Rogers, T. J., Adler, M. W., & Eisenstein, T. K. (1993). Morphine
treatment in vitro or in vivo decreases phagocytic functions of murine macrophages. *Life Sci,*
53(12), 997-1006. https://doi.org/10.1016/0024-3205(93)90122-j

969
970 Rosen, S. F., Ham, B., Haichin, M., Walters, I. C., Tohyama, S., Sotocinal, S. G., & Mogil, J. S. (2019, Feb).
971   Increased pain sensitivity and decreased opioid analgesia in T-cell-deficient mice and
972   implications for sex differences. *Pain, 160*(2), 358-366.
973   https://doi.org/10.1097/j.pain.0000000000001420

974
975 Ryu, J. H., Do, S. H., Han, S. H., & Zuo, Z. (2018, Jul). Morphine reduces mouse microglial engulfment
976   induced by lipopolysaccharide and interferon-γ via δ opioid receptor and p38 mitogen-activated
977   protein kinase. *Neurol Res, 40*(7), 600-606. https://doi.org/10.1080/01616412.2018.1455368

978
979 Sassone-Corsi, P. (2012, Dec 1). The cyclic AMP pathway. *Cold Spring Harb Perspect Biol, 4*(12).
980   https://doi.org/10.1101/cshperspect.a011148

981
982 Schäfer, M., Carter, L., & Stein, C. (1994, May 10). Interleukin 1 beta and corticotropin-releasing factor
983   inhibit pain by releasing opioids from immune cells in inflamed tissue. *Proc Natl Acad Sci U S A,*
984   *91*(10), 4219-4223. https://doi.org/10.1073/pnas.91.10.4219

985
986 Schaub, B., Bellou, A., Gibbons, F. K., Velasco, G., Campo, M., He, H., Liang, Y., Gillman, M. W., Gold, D.,
987   Weiss, S. T., Perkins, D. L., & Finn, P. W. (2004, Sep). TLR2 and TLR4 stimulation differentially
988   induce cytokine secretion in human neonatal, adult, and murine mononuclear cells. *J Interferon*
989   *Cytokine Res, 24*(9), 543-552. https://doi.org/10.1089/jir.2004.24.543

990
991 Smith, H. S. (2009, Jul). Opioid metabolism. *Mayo Clin Proc, 84*(7), 613-624.
992   https://doi.org/10.1016/s0025-6196(11)60750-7

993
994 Sowa, G., Gekker, G., Lipovsky, M. M., Hu, S., Chao, C. C., Molitor, T. W., & Peterson, P. K. (1997, Mar
995   21). Inhibition of swine microglial cell phagocytosis of Cryptococcus neoformans by femtomolar
996   concentrations of morphine. *Biochem Pharmacol, 53*(6), 823-828.
997   https://www.ncbi.nlm.nih.gov/pubmed/9113103

998
999 Sowa, G., Gekker, G., Lipovsky, M. M., Hu, S., Chao, C. C., Molitor, T. W., & Peterson, P. K. (1997,
1000   1997/03/21/). Inhibition of swine microglial cell phagocytosis of cryptococcus neoformans by
1001   femtomolar concentrations of morphine. *Biochemical Pharmacology, 53*(6), 823-828.
1002   https://doi.org/https://doi.org/10.1016/S0006-2952(97)00006-3

1003
1004 Stain-Texier, F., Boschi, G., Sandouk, P., & Scherrmann, J. M. (1999, Oct). Elevated concentrations of
1005   morphine 6-beta-D-glucuronide in brain extracellular fluid despite low blood-brain barrier
1006   permeability. *Br J Pharmacol, 128*(4), 917-924. https://doi.org/10.1038/sj.bjp.0702873

1007

1008 Stansley, B., Post, J., & Hensley, K. (2012, May 31). A comparative review of cell culture systems for the
1009   study of microglial biology in Alzheimer's disease. *J Neuroinflammation, 9*, 115.
1010   https://doi.org/10.1186/1742-2094-9-115

1011
1012 Stoicea, N., Costa, A., Periel, L., Uribe, A., Weaver, T., & Bergese, S. D. (2019, May). Current perspectives
1013   on the opioid crisis in the US healthcare system: A comprehensive literature review. *Medicine
1014   (Baltimore), 98*(20), e15425. https://doi.org/10.1097/md.0000000000015425

1015
1016 Strang, J., Volkow, N. D., Degenhardt, L., Hickman, M., Johnson, K., Koob, G. F., Marshall, B. D. L., Tyndall,
1017   M., & Walsh, S. L. (2020, Jan 9). Opioid use disorder. *Nat Rev Dis Primers, 6*(1), 3.
1018   https://doi.org/10.1038/s41572-019-0137-5

1019
1020 Terplan, M. (2017, Aug). Women and the opioid crisis: historical context and public health solutions.
1021   *Fertil Steril, 108*(2), 195-199. https://doi.org/10.1016/j.fertnstert.2017.06.007

1022
Thion, M. S., Low, D., Silvin, A., Chen, J., Grisel, P., Schulte-Schrepping, J., Blecher, R., Ulas, T., Squarzoni,
P., Hoeffel, G., Coulpier, F., Siopi, E., David, F. S., Scholz, C., Shihui, F., Lum, J., Amoyo, A. A.,
Larbi, A., Poidinger, M., Buttgereit, A., Lledo, P. M., Greter, M., Chan, J. K. Y., Amit, I., Beyer, M.,
Schultze, J. L., Schlitzer, A., Pettersson, S., Ginhoux, F., & Garel, S. (2018, Jan 25). Microbiome
Influences Prenatal and Adult Microglia in a Sex-Specific Manner. *Cell, 172*(3), 500-516.e516.
https://doi.org/10.1016/j.cell.2017.11.042

1029
1030 Thomas, J. H. L., Lui, L., Abell, A., Tieu, W., Somogyi, A. A., Bajic, J. E., & Hutchinson, M. R. (2022, May).
1031   Toll-like receptors change morphine-induced antinociception, tolerance and dependence:
1032   Studies using male and female TLR and signalling gene KO mice. *Brain Behav Immun, 102*, 71-85.
1033   https://doi.org/10.1016/j.bbi.2022.02.001

1034
1035 Thomas, J. H. L., Lui, L., Abell, A., Tieu, W., Somogyi, A. A., Bajic, J. E., & Hutchinson, M. R. (2022,
1036   2022/05/01/). Toll-like receptors change morphine-induced antinociception, tolerance and
1037   dependence: Studies using male and female TLR and signalling gene KO mice. *Brain, Behavior,
1038   and Immunity, 102*, 71-85. https://doi.org/https://doi.org/10.1016/j.bbi.2022.02.001

1039
1040 Tynan, R. J., Naicker, S., Hinwood, M., Nalivaiko, E., Buller, K. M., Pow, D. V., Day, T. A., & Walker, F. R.
1041   (2010, Oct). Chronic stress alters the density and morphology of microglia in a subset of stress-
1042   responsive brain regions. *Brain Behav Immun, 24*(7), 1058-1068.
1043   https://doi.org/10.1016/j.bbi.2010.02.001

1044
1045 Valentino, R. J., & Volkow, N. D. (2018, Dec). Untangling the complexity of opioid receptor function.
1046   *Neuropsychopharmacology, 43*(13), 2514-2520. https://doi.org/10.1038/s41386-018-0225-3

1047
1048 Veremeyko, T., Yung, A. W. Y., Dukhinova, M., Kuznetsova, I. S., Pomytkin, I., Lyundup, A., Strekalova, T.,
1049   Barteneva, N. S., & Ponomarev, E. D. (2018). Cyclic AMP Pathway Suppress Autoimmune

1050   Neuroinflammation by Inhibiting Functions of Encephalitogenic CD4 T Cells and Enhancing M2
1051   Macrophage Polarization at the Site of Inflammation. *Front Immunol, 9*, 50.
1052   https://doi.org/10.3389/fimmu.2018.00050

1053
1054   Volkow, N. D., Jones, E. B., Einstein, E. B., & Wargo, E. M. (2019, Feb 1). Prevention and Treatment of
1055   Opioid Misuse and Addiction: A Review. *JAMA Psychiatry, 76*(2), 208-216.
1056   https://doi.org/10.1001/jamapsychiatry.2018.3126

1057
1058   Volpe, D. A., McMahon Tobin, G. A., Mellon, R. D., Katki, A. G., Parker, R. J., Colatsky, T., Kropp, T. J., &
1059   Verbois, S. L. (2011, Apr). Uniform assessment and ranking of opioid μ receptor binding
1060   constants for selected opioid drugs. *Regul Toxicol Pharmacol, 59*(3), 385-390.
1061   https://doi.org/10.1016/j.yrtph.2010.12.007

1062
1063   Wang, D., Jiang, J., Shang, W., Zhang, J., Jiang, X., Shen, F., Liang, J., Li, Y., Li, M., Wang, M., & Sui, N.
1064   (2023, Mar 8). Effect of early embryonic exposure to morphine on defects in the GABAergic
1065   system of day-old chicks. *Prog Neuropsychopharmacol Biol Psychiatry, 121*, 110657.
1066   https://doi.org/10.1016/j.pnpbp.2022.110657

1067
1068   Wang, Q., Chu, C. H., Qian, L., Chen, S. H., Wilson, B., Oyarzabal, E., Jiang, L., Ali, S., Robinson, B., Kim, H.
1069   C., & Hong, J. S. (2014, Sep 10). Substance P exacerbates dopaminergic neurodegeneration
1070   through neurokinin-1 receptor-independent activation of microglial NADPH oxidase. *J Neurosci,
1071   34*(37), 12490-12503. https://doi.org/10.1523/jneurosci.2238-14.2014

1072
1073   Wang, Q., Oyarzabal, E., Wilson, B., Qian, L., & Hong, J. S. (2015, Oct 1). Substance P enhances microglial
1074   density in the substantia nigra through neurokinin-1 receptor/NADPH oxidase-mediated
1075   chemotaxis in mice. *Clin Sci (Lond), 129*(8), 757-767. https://doi.org/10.1042/cs20150008

1076
1077   Wang, X., Loram, L. C., Ramos, K., de Jesus, A. J., Thomas, J., Cheng, K., Reddy, A., Somogyi, A. A.,
1078   Hutchinson, M. R., Watkins, L. R., & Yin, H. (2012). Morphine activates neuroinflammation in a
1079   manner parallel to endotoxin. *Proceedings of the National Academy of Sciences, 109*(16), 6325-
1080   6330. https://doi.org/doi:10.1073/pnas.1200130109

1081
1082   Wang, X., Loram, L. C., Ramos, K., de Jesus, A. J., Thomas, J., Cheng, K., Reddy, A., Somogyi, A. A.,
1083   Hutchinson, M. R., Watkins, L. R., & Yin, H. (2012, Apr 17). Morphine activates
1084   neuroinflammation in a manner parallel to endotoxin. *Proc Natl Acad Sci U S A, 109*(16), 6325-
1085   6330. https://doi.org/10.1073/pnas.1200130109

1086
1087   Woo, M.-S., Jung, S.-H., Hyun, J.-W., & Kim, H.-S. (2004, 2004/02/19/). Differential regulation of
1088   inducible nitric oxide synthase and cytokine gene expression by forskolin and dibutyryl-cAMP in
1089   lipopolysaccharide-stimulated murine BV2 microglial cells. *Neuroscience Letters, 356*(3), 187-
1090   190. https://doi.org/https://doi.org/10.1016/j.neulet.2003.11.056

1091

1092   Wu, R., & Li, J.-X. (2020, 2020-November-24). Toll-Like Receptor 4 Signaling and Drug Addiction
1093   [Review]. *Frontiers in Pharmacology, 11*. https://doi.org/10.3389/fphar.2020.603445

1094
1095   Xie, N., Gomes, F. P., Deora, V., Gregory, K., Vithanage, T., Nassar, Z. D., Cabot, P. J., Sturgess, D., Shaw,
1096   P. N., & Parat, M. O. (2017, Mar). Activation of μ-opioid receptor and Toll-like receptor 4 by
1097   plasma from morphine-treated mice. *Brain Behav Immun, 61*, 244-258.
1098   https://doi.org/10.1016/j.bbi.2016.12.002

1099
1100   Yang, H. S., Onos, K. D., Choi, K., Keezer, K. J., Skelly, D. A., Carter, G. W., & Howell, G. R. (2021, Feb 9).
1101   Natural genetic variation determines microglia heterogeneity in wild-derived mouse models of
1102   Alzheimer's disease. *Cell Rep, 34*(6), 108739. https://doi.org/10.1016/j.celrep.2021.108739

1103
1104   Yang, T. T., Lin, C., Hsu, C. T., Wang, T. F., Ke, F. Y., & Kuo, Y. M. (2013, Jul). Differential distribution and
1105   activation of microglia in the brain of male C57BL/6J mice. *Brain Struct Funct, 218*(4), 1051-1060.
1106   https://doi.org/10.1007/s00429-012-0446-x

1107
1108   Yang, Y., Sun, Y., Hu, R., Yan, J., Wang, Z., Li, W., & Jiang, H. (2021). Morphine promotes microglial
1109   activation by upregulating the EGFR/ERK signaling pathway. *PLoS One, 16*(9), e0256870.
1110   https://doi.org/10.1371/journal.pone.0256870

1111
1112   Yoo, M., Perret, D., Dusto, N., Lin, K., & Whyte, J. (2022, Jul 1). Pain and Addiction: The Distribution of
1113   Supply and Demand During the Opioid Epidemic. *Am J Phys Med Rehabil, 101*(7), 702-707.
1114   https://doi.org/10.1097/phm.0000000000001914

1115
1116   Zare, N., Pourhadi, M., Vaseghi, G., & Haghjooy Javanmard, S. (2023, Apr). The potential interplay
1117   between opioid and the toll-like receptor 4 (TLR-4). *Immunopharmacol Immunotoxicol, 45*(2),
1118   240-252. https://doi.org/10.1080/08923973.2022.2122500

1119
1120   Zhang, L., Zhang, J. T., Hang, L., & Liu, T. (2020). Mu Opioid Receptor Heterodimers Emerge as Novel
1121   Therapeutic Targets: Recent Progress and Future Perspective. *Front Pharmacol, 11*, 1078.
1122   https://doi.org/10.3389/fphar.2020.01078

1123
1124   Zhang, Y., Chen, K., Sloan, S. A., Bennett, M. L., Scholze, A. R., O'Keeffe, S., Phatnani, H. P., Guarnieri, P.,
1125   Caneda, C., Ruderisch, N., Deng, S., Liddelow, S. A., Zhang, C., Daneman, R., Maniatis, T., Barres,
1126   B. A., & Wu, J. Q. (2014, Sep 3). An RNA-sequencing transcriptome and splicing database of glia,
1127   neurons, and vascular cells of the cerebral cortex. *J Neurosci, 34*(36), 11929-11947.
1128   https://doi.org/10.1523/JNEUROSCI.1860-14.2014

1129
1130   Zhang, Y., Sloan, S. A., Clarke, L. E., Caneda, C., Plaza, C. A., Blumenthal, P. D., Vogel, H., Steinberg, G. K.,
1131   Edwards, M. S., Li, G., Duncan, J. A., 3rd, Cheshier, S. H., Shuer, L. M., Chang, E. F., Grant, G. A.,
1132   Gephart, M. G., & Barres, B. A. (2016, Jan 6). Purification and Characterization of Progenitor and

1133   Mature Human Astrocytes Reveals Transcriptional and Functional Differences with Mouse.
1134   *Neuron, 89*(1), 37-53. https://doi.org/10.1016/j.neuron.2015.11.013

1135

1136 Zhang, Y. Q., Lin, W. P., Huang, L. P., Zhao, B., Zhang, C. C., & Yin, D. M. (2021, Nov 8). Dopamine D2
1137   receptor regulates cortical synaptic pruning in rodents. *Nat Commun, 12*(1), 6444.
1138   https://doi.org/10.1038/s41467-021-26769-9

1139

1140