Distinct phenotype of SARS-CoV-2 Omicron BA.1 in human primary cells but no increased host range in cell lines of putative mammalian reservoir species

Manel Essaidi-Laziosi[1], Francisco Javier Perez Rodriguez[2], Catia Alvarez[1], Pascale Sattonnet-Roche[2], Giulia Torriani[1], Meriem Bekliz[1], Kenneth Adea[1], Matthias Lenk[3], Tasnim Suliman[4], Wolfgang Preiser[5,6], Marcel A. Müller[7], Christian Drosten[7], Laurent Kaiser[8,9,10], Isabella Eckerle[1,9,10*]

[1]Department of Microbiology and Molecular Medicine, University of Geneva, Geneva, Switzerland

[2]Division of Infectious Diseases, Geneva University Hospitals, Geneva, Switzerland.

[3] Friedrich-Loeffler-Institute, Federal Research Institute for Animal Health, Greifswald, Germany

[4]Department of Medical Biosciences, University of the Western Cape, Cape Town, South Africa.

[5]Division of Medical Virology, Faculty of Medicine and Health Sciences, Stellenbosch University, Cape Town, South Africa.

[6]Division of Medical Virology, Tygerberg Hospital, National Health Laboratory Service, Cape Town, South Africa.

[7]Institute of Virology Charité, – Universitätsmedizin Berlin, Berlin, Germany

[8]Laboratory of Virology, Division of Infectious Diseases and Division of Laboratory Medicine, University Hospitals of Geneva & Faculty of Medicine, University of Geneva, 1205 Geneva, Switzerland.

[9]Geneva Centre for Emerging Viral Diseases, University Hospitals Geneva, and University of Geneva, Switzerland.

[10]Division of Infectious Diseases, Geneva University Hospitals, 1205 Geneva, Switzerland.

*corresponding author: Isabella Eckerle, University Hospital of Geneva, Rue Gabrielle-

Perret-Gentil 4, 1205 Geneva, Switzerland, Isabella.eckerle@hcuge.ch , Phone: +41 22 37

29820, Fax: +41 22 37 24097

Running title: Phenotype of SARS-CoV-2 Omicron in cell cultures

Key words: SARS-CoV-2, COVID19, Omicron, variants of concern, primary airway

epithelial cells, reservoir host

Abstract

SARS-CoV-2's genetic plasticity has led to several variants of concern (VOCs). Here we

studied replicative capacity for seven SARS-CoV-2 isolates (B.1, Alpha, Beta, Gamma, Delta,

Zeta, and Omicron BA.1) in primary reconstituted airway epithelia (HAE) and lung-derived

cell lines. Furthermore, to investigate the host range of Delta and Omicron compared to

ancestral SARS-CoV-2, we assessed replication in 17 cell lines from 11 non-primate

mammalian species, including bats, rodents, insectivores and carnivores. Only Omicron's

phenotype differed *in vitro*, with rapid but short replication and efficient production of

infectious virus in nasal HAEs, in contrast to other VOCs, but not in lung cell lines. No

increased infection efficiency for other species was observed, but Delta and Omicron

infection efficiency was increased in A549 cells. Notably replication in A549 and Calu3 cells

was lower than in nasal HAE. Our results suggest better adaptation of VOCs towards humans,

without an extended host range.

Introduction

Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) is the etiological agent of coronavirus disease 19 (COVID19) that has caused one of the biggest public health challenges of modern times (1). Few mutational changes were observed in SARS-CoV-2 during the first year of the pandemic, most notably the Spike D614G mutation. This mutation enhanced angiotensin converting enzyme-2 (ACE-2) binding, providing a fitness advantage, and was responsible for the first pandemic wave (2, 3). By late 2020, SARS-CoV-2 variants had emerged, characterised by numerous mutations, mainly in Spike. These were classified as variants of concern (VOC), variants of interest (VOI) or variants under monitoring (VUMs), based on their genetic, clinical and epidemiological characteristics (4). To date, five VOCs have been designated: Alpha, Beta, Gamma, Delta and Omicron. VOCs are characterized by a rapid increase in case numbers, quickly outcompeting earlier strains in their region of emergence (5-9). Omicron has the most observed mutations, with the majority located in Spike, causing the strongest escape from prior immunity so far (10). It also shows signs of higher transmissibility and secondary attack rates (11). In addition, VUMs and VOIs were co-circulating along with VOCs, including the (former) VOI Zeta used in our study here, that arose alongside the Gamma VOC in South America during a surge in local cases, but nearly all have disappeared (12).

Variant evolution does not yet follow a pattern, and most variants have not evolved from each other, but arose independently from basal circulating viruses (13). Indeed, Omicron is most closely related to a virus that was circulating in mid-2020. There are three main hypotheses on VOC origins (14). The first holds that they developed in populations that were not covered by genomic surveillance. The second involves reverse zoonotic events, with transmission from humans into intermediate hosts, and then back to humans. While a large range of animal species are known to be susceptible to SARS-CoV-2, no plausible VOC progenitors have been found in animals (15). The third hypothesis describes intra-host evolution during chronic infections in immunocompromised patients. Such cases have already been reported, with prolonged shedding of mutated infectious virus (16-18).

VOCs have shown distinct epidemiological and partial clinical differences. While genomic surveillance can inform about mutations and the phylogenetic relationships, it cannot directly conclude on biological properties resulting from these mutations, thus the *in vitro* assessment of phenotypic differences is of immediate relevance whenever new variants arise. Understanding the mechanisms of enhanced fitness, the origins of VOCs and their risk for

reverse zoonotic events are crucial for the further control of this pandemic. Here, we have

assessed replicative capacity and potential *in vitro* phenotypes of ancestral SARS-CoV-2

(B.1), VOCs Alpha, Beta, Gamma, Delta, Omicron (BA.1) and former VOI Zeta on human

primary cells and lung-derived cell lines and on mammalian cell lines derived from bats,

rodents, insectivores and carnivores to investigate potential host range.

Results

Comparative replicative capacity of SARS-CoV-2 variants in human cell culture models

of the respiratory tract

To compare viral phenotypes, we infected primary human airway epithelial cells (HAE),

derived from the nasal epithelium and differentiated *in vitro* in 3-dimensional (3D) air-liquid

interface cultures and monolayer (2D) cultures with SARS-CoV-2 clinical isolates at the

multiplicity of infection (MOI) of 0.1 at 37°C and 33°C (**Figure 1**). Upon infection of HAE

cultures and incubation at 37°C, we observed a rapid increase at 24h post infection for

Omicron, from mean $\log_{10}$ RNA copy numbers/mL (RNAc/mL) of 7.8 to a peak of 11.8,

while all other isolates reached similar peak values at around 96 hours. Omicron, in contrast,

showed a reduction of viral RNAc/mL at 96h to 10.8 $\log_{10}$ RNAc/mL. No differences in

replication kinetics were observed between B.1, Alpha, Beta, Gamma, Delta and Zeta, in all

of which log10 RNAc gradually increased over the first 4 days.

When assessing infectious viral particles by plaque-forming assay at 37°C, Omicron reached

peak infectious viral titer of 7.1 $\log_{10}$ plaque-forming units (PFU) after 24h with a rapid

decline to only 4.3 after 96h. On the contrary, B.1, Beta, Gamma and Delta showed an

increase in PFU from 24 to 96 h, but peak PFU titers stayed below that of Omicron. The

Alpha and the Zeta variants did not show strong increases, with PFUs for 24 and 96h staying

constant and below those of the other variants. The strongest increase was observed for Delta,

which showed the lowest infectious titer at 24h of 3.8, which increased to 6.6 mean $\log_{10}$

PFU/mL.

At 33°C incubation temperature, Omicron also showed a fold increase of 4.0 mean $\log_{10}$

RNAc/mL at 24h, but RNA levels increased further with a peak at 96h of 11.1 mean $\log_{10}$

RNAc/mL. Similarly, the other variants including B.1 reached their peak RNA levels at 96h,

but with higher RNA levels than that of Omicron, ranging between 11.8 and 12.6 mean $\log_{10}$

RNAc/mL. Relative to 37°C, at 33°C viral RNAc/mL were 0.3-0.8 $\log_{10}$ higher for B.1,

Alpha, Beta, Gamma and Zeta, but not for Delta and Omicron, which both showed slightly

lower viral loads at 33°C. Similarly, PFUs were higher for B.1, Alpha, Beta, Gamma and Zeta at 33°C, but similar for Delta and lower for Omicron. No higher PFU titer was observed for Omicron at 24h, and similar infectious titers were found for 24 and 96 hpi (5.8-5.9 mean log10 PFU/mL).

The same experiments were performed in Calu3 cells derived from a human lung adenocarcinoma (**Figure 2**). Replication kinetics at 37°C were similar for all viruses, including Omicron, with peak RNA levels reached at 48hpi. Peak RNA levels were comparable between B.1, Alpha, Beta, Gamma and Zeta, ranging from 10.7 to 11.3, but higher for Delta and Omicron (12.1 for both). No early peak for Omicron, such as in HAE cultures, was observed. In contrast to HAE cultures, PFU titers peaked rapidly in this cell line at 37°C, with higher infectious viral loads at 24h post infection than at 96hpi. Only Omicron did not show an early PFU titer peak, but instead showed an increase from 24 to 96hpi, with peak PFU titers remaining lower than those of the other variants (mean 5.1 $\log_{10}$ PFU/mL).

Viral replication at 33°C showed a homogeneous replication pattern between variants, with peak RNA levels for B.1, Alpha, Beta, Gamma and Zeta reached at 48hpi in the range of 11.1-11.9, while peak RNA levels for Delta and Omicron were delayed with a peak 96hpi with titers of 11.0- 10.1, respectively. Among all variants, Omicron showed the lowest replication with ca. 1 $\log_{10}$ RNA copies/mL lower than Delta and up to 2 $\log_{10}$ RNA copies/mL lower compared to B.1, Alpha, Beta, Gamma and Zeta.

At 33°C compared to 37°C, B.1, Alpha, Beta, Gamma and Zeta had higher peak RNA levels by 0.3 to 0.6 $\log_{10}$ RNAc/mL, while Delta and Omicron had 1.0 and 2.0 $\log_{10}$ RNAc/mL lower peaks, respectively. PFU titers were lower in Calu3 at 33°C, but infectious viral titers 24hpi were still higher than at 96hpi.

Replicative capacity of B.1, Delta and Omicron in mammalian cell lines

In order to assess the replicative capacity of B.1, Delta, and Omicron BA.1, we infected 17 non-human mammalian cell lines, from 11 wild or domestic species (**Table 1**). Species include a range of European wildlife. Cells were inoculated at a MOI of 0.1 and 37°C. The supernatant's viral RNA was quantified after 4 days, along with the PFU titer if viral RNA had increased. Human A549 (with low expression of ACE-2) cell lines and Vero-E6 cell lines (primate) were infected in parallel as controls (**Figure 3**). Efficient replication was observed for all three viruses in a rabbit kidney cell line (RK-13), with RNA levels between 10.3-11.5 $\log_{10}$ RNAc/mL at 4 dpi. Upon titration, infectious titers were between 5.8-7.0 PFU/mL.

Vero-E6 cells, which were infected in parallel, showed an increase in viral RNA between 10.7-11.6 $\log_{10}$ RNA copies/mL, resulting in 6.1-7.0 PFU/mL. No other non-human mammalian cell lines from the species of bats, rodents, minks and foxes showed increased RNA indicating efficient replication, further confirmed by a lack of increase in sub-genomic RNA for all three viruses (**Figure S1**). In contrast, human A549 cells did not show an increase of genomic nor by sub-genomic viral RNA upon infection with B.1, but were efficiently infected by Delta and Omicron with respective increases in $\log_{10}$ RNA copies/mL at 4 dpi of 0.7 and 1.3 and infectious virus titers of 3.9 and 3.2 PFU/mL.

Comparison of ACE-2 orthologs from animal species.

In order to understand differences between ACE-2 receptor binding residues of tested cell lines, we decided to compare ACE-2 amino-acid sequences between the studied species (**Table 2**). These sequences were either available in GenBank/NCBI or obtained by sequencing of the cell lines used when no sequence was available yet (see material and method). The alignment and comparison of key ACE-2 orthologue residues involved in binding to the spike protein showed no difference between *Homo sapiens* and *Chlorocebus aethiops* (Vero-E6). Of the other mammalian species, rabbits and Red foxes showed the highest (76.92%) homology to human ACE-2. Bats, shrews and hedgehogs showed the lowest homology rates (50-65.3%). Amino-acids F28, E37, L45, N53, N330, K353, D355, R357 and R393 were highly conserved between species while amino-acids at positions 38, 41, 83, 27, 322, 329 and 354 were rather variable.

Discussion

Since its emergence in 2019, SARS-CoV-2 has been in constant evolution, frequently giving rise to new variants (19). The emergence of several VOCs, including the latest Omicron has stressed the need for continuous surveillance and rapid phenotypic assessment to guide public health measures. A plethora of studies compared SARS-CoV-2 variants either by live virus or pseudo-viral assays using human *in vitro* models such as cell lines or organoids, or animal models, with the aim to provide rapid answers on the altered biological properties of novel variants (11, 20-22). Differentiated 3D tissues of the human respiratory tract and organoids are considered as one of the most relevant models for such studies as they are human-derived, readily available and most closely recapitulate the *in vivo* situation (23). Nevertheless, widely available immortalized cells are commonly used to study virus tropism and pathogenicity and

decipher the mechanisms involved, and can reflect selected aspects. A range of immortalized
cell lines of wildlife species have become available in recent years, allowing study of the
susceptibility of different hosts, especially those that are not experimental animal models (24-
27). In our study, we extensively compare the replication of isolates of seven SARS-CoV-2
lineages in cell culture models of the human respiratory tract and in cell lines obtained from
domestic and wildlife species, with a focus on European species.

In nasal HAE cells, our findings clearly demonstrate a distinct BA.1 phenotype compared to
the other variants, with shorter but faster replication to higher levels, and early efficient
production of infectious virus. The early and efficient replication, in addition to its immune
evasion properties, could contribute to Omicron's contagiousness and high secondary attack
rate (28, 29). However, in a recent clinical study, we did not observe more infectious viral
shedding for Omicron at symptom onset compared to Delta (30). These differences could be
reflected by the fact that our HAE model allowed studying the very early phase of virus
replication that is undetected by diagnostic testing, as clinical samples are usually collected
after symptom onset. Furthermore, adaptive immunity is lacking in our HAE model, which
may mitigate replication, especially given the infection rates in vaccinated or previously
infected individuals. Although Omicron quickly reaches high viral levels, both RNA levels
and PFU titer rapidly declines in HAE cells, and are lower than levels for Delta at 96 hours, in
agreement with clinical observations.

Agreeing with our findings, early and rapid replication of BA.1 was also seen in human *ex*
vivo bronchus but less in lung parenchyma (21). Less efficient cleavage of the Omicron spike
was suggested to reduce fusogenicity (31). In our study, we have also seen less efficient
replication in lung-derived cells versus HAE cultures, although there are limits to their
comparisons. The less efficient replication in the lower respiratory tract might explain the
reduced clinical severity of BA.1 (22, 31), while the early efficient replication may contribute
to efficient community spread. Additionally, its efficient replication could be explained by an
enhanced cell host entry thanks to improved ACE-2 binding and more efficient endocytosis,
rather than lower sensitivity to interferon responses (32).

Better replication was shown for ancestral SARS-CoV-2 at 33°C in nasal and lung *in vitro*
models (33), and we also observed more efficient infectious virus production at lower
temperature for Alpha, Beta and Zeta, but not for Delta and Omicron (Figure1). Similarly, no
replication differences between 33°C and 37°C were observed for Omicron in organoids of
bronchi and lung in another study (21).

We also investigated host tropism in cell culture models. SARS-CoV-2 has already
demonstrated high promiscuity by infecting multiple animal species, with both animal-human
spill-backs and establishment of new animal reservoirs (15, 34-36). In general, inter-species
transmission is a well-known feature of coronaviruses, considerably contributing to their
pandemic potential, and in the case of SARS-CoV-2 facilitated by conservation of the host
receptor ACE-2 across mammalian species (37-39).

As with any new emerging virus, establishment of new host reservoirs poses a risk for further
mutation of the virus, complicates control of transmission and eventually leads to spill-backs
into humans. Both an origin as well as an increased risk for spill over into new reservoirs of
new variants has been discussed (40). We did not see an increased host range for Delta or
BA.1 in our animal cell lines, which included a range of European small mammals and
several species of bats of the family *Rhinolophidae*. *Rhinolophidae* are known to host diverse
SARS-related coronaviruses, and the closest known relative of SARS-CoV-2 was found in an
Asian species (41 , 42). In contrast, we investigated two European species, *R. ferrumequinum*
and *R. euryale*, and a South African species, *R. clivosus*, with no signs of SARS-CoV-2
replication in kidney and tracheal cell lines from these bats. Nor did the cells of a more
ubiquitous non-*Rhinolophidae* bat, *P. pipistrellus*, show any signs of replication by B.1, Delta
or Omicron. Other studies have shown SARS-CoV-2 does not readily replicate in a range of
bat cell lines including *Rhinolophus* cells (43) while one study found signs of replication in a
kidney cell line from an Asian Rhinolophus species (44). Another study found bat cell lines to
be only weakly susceptible, which could be overcome by expression of hACE-2, similar to
what was observed for SARS-CoV (45, 46).

The only susceptible cell line from our panel was derived from a rabbit kidney, which
replicated equally well in B.1, Delta and Omicron, but no other animal cell lines showed signs
of efficient replication. Indeed, rabbits have been used as susceptible experimental animal
models (47). However, three other rabbit cell lines, did not show signs of replication.
Surprisingly, we were not able to reproduce efficient infection of cells originating from minks
and ferrets, although SARS-CoV-2 was reported to infect both species and has caused
outbreaks in minks farmed for fur (48).

Other studies have assessed the host range of SARS-CoV-2 mostly using ACE2-transgenic
cell lines and pseudotyped viruses, but fewer studies have investigated VOCs, and results
between studies were conflicting (49). A study using lentiviral pseudotypes and transgenic
animal ACE2-expressing cell lines suggested Omicron's tropism extended into domestic

avian species, *Rhinolophus* bats and mice (50). A study using a similar approach suggested broad entry of ancestral SARS-CoV-2 into a range of animal species (cat, dog, cow, horse, camel, hamster, rabbit but not mink) as well as enhanced entry of Alpha and Beta variants (51). One study investigated well-differentiated airway cultures from a range of animals (some of which were also studied here) and found ancestral SARS-CoV-2 replication in cells derived from rhesus macaques and cats, but not in those derived from ferrets, dogs, rabbits, pigs, cattle, goats, camels, llamas and two neotropical bat species (52).

Discrepancies between *in vitro* findings and naturally observed infections or animal experiments could be because: the culture models do not accurately reflect the site of replication *in vivo*, receptor expression is reduced in cell culture, differences in the required infection dose, body temperature, or SARS-CoV-2 uses different receptors in some animal species. Results obtained with pseudotyped viruses and transgenic ACE-2 expression could furthermore differ from wild-type virus assessment in unaltered cell lines.

In addition to infection experiments, several modelling and bioinformatic approaches have tried to identify susceptible species by comparing ACE-2 sequences (37, 53). The *in-silico* prediction of ACE-2 affinity with spike's RBD previously identified of more than 500 animal species as potentially susceptible to SARS-CoV-2 (54). Others have found that the predictive power of ACE-2 sequences is rather low and can be misleading due to biased ACE-2 sequence availability (55). Therefore, phenotypic assessment in cell lines, although not perfectly reflecting the *in vivo* susceptibility, can complement such bioinformatic studies. Cell lines from wildlife species such as bats can provide both biological infection data and a source for ACE-2 sequencing, as done here.

Despite its important variability across species, ACE-2 appears more genetically stable compared to TMPRSS2, a type II transmembrane serine protease involved in virus-cell fusion during virus entry, which had shown less identities between human and animals with possible partial-to-total TMPRSS2 gene loss in some vertebrates (56).

Together with data from animal cells, our observations support a human origin of Omicron rather than a reverse zoonosis. Although bats are considered to be the original hosts, SARS-CoV-2 does not readily replicate in bats cells because of the divergent RBD sequences compared to close relatives in bats, like RaTG13.

In conclusion, using cell models, we showed that Omicron has the strongest phenotypic differences compared to the previous variants, and likely evolved in humans rather than

animals. Such cell culture models can help better understand SARS-CoV-2 infections,

including VOCs, in humans. We also showed the relevance of cell models from a variety of

species, even if not perfectly reflecting *in vivo* situation, to assessing the risk of zoonotic

spillback.

Material and methods

Human and animal Cell lines

Commercially available nasal HAE, called "MucilAir^TM", were purchased from Epithelix

SARL [www.epithelix.com]. They are maintained in air-liquid interface culture where the

medium (MucilAir^TM medium, Epithelix) is supplemented by the basal chamber and the

apical surface of the tissue is in contact with air (see (57) for details).

All used animal cell lines are summarized in Table1. The corresponding species were

confirmed by sequencing of the cytochrome C subunit oxidase I gene (58, 59). Five additional

standard cell lines were used in this study as controls or for viral stock production, including

Calu3 (human lung cancer cell), Vero-E6 (African Green Monkey kidney), A549 (human

lung carcinoma), A549 cells overexpressing ACE-2 receptor and Vero-E6 overexpressing

TMPRSS2. All immortalized cells were cultured in monolayer using medium (MEM

Glutamax, Gibco) supplemented with 10% (cell maintenance) or 2% (infections) Fetal Bovin

Serum (FBS, Gibco), 1X penicillin-streptomycin (Gibco) and 1% non-essential amino-acids

(MEM NEAA, Gibco). All cell cultures were performed at 37 °C under 5% CO_2.

Viruses

SARS-CoV-2 isolates were obtained from clinical samples collected in the outpatient testing

Centre of the University Hospitals of Geneva as described previously (29). All SARS-CoV-2

isolates used are summarized in Table S1. B.1 and the variants Alpha, Gamma, Zeta and Delta

were isolated after one passage in Vero-E6 cells. One additional passage allowed viral stock

production in the same line. Vero-E6 was less permissive to Beta and Omicron variants.

Hence, Beta was isolated in A549-ACE-2, after a second passage in mixed Vero-E6:A549-

ACE-2 (1:1) cells, the viral stock was produced in Vero-E6 cells. Omicron was isolated after

2 passages in Vero-TMPRSS2. All viral stocks were titrated in Vero-E6 and sequenced.

Infections of animal and human cells

Infection assays were performed at 37°C or 33°C at 5% CO_2. HAE reconstituted *in vitro* (3D

culture) or immortalised cell lines in 2D culture, including Calu3 and animal cell lines (Table

1), were tested for infections as previously described (20) at 37°C and an MOI of 0.1 using

seven SARS-CoV-2 isolates (Table S1). Cells were washed after 1h (Calu3 and animal cells)

or 3h (HAE). Supernatant (Calu3) or apical washes (HAE) were collected at different times

after infection. Infected animal cells were lysed using NucliSens easyMAG lysis buffer

(BioMérieux) at 4 dpi.

Quantification of viral RNA

In order to determine the viral load from collected samples, RNA was extracted with

NucliSens easyMAG (BioMérieux) and quantified by quantitative real time PCR (RT-qPCR)

using SuperScript™ III Platinum™ One-Step qRT-PCR Kit (Invitrogen) in a CFX96 Thermal

Cycler (BIORAD). Real time RT-qPCR was performed using specific sets of primers and

probes targeting either genomic or sub-genomic viral RNA as previously described (60, 61)

Assessment of infectious titer

Infectious titer of collected samples was assessed by plaque assays performed at 37°C and 5%

CO_2 as previously described (20). Briefly, Vero-E6 seeded in monolayer of $2x10^5$ cells/mL in

24-well plates were inoculated 2h after using serially diluted samples. The inoculum was

removed and replaced by fresh medium (DMEM 10%FBS, 2mM L-glutamine, 1% penicillin-

streptomycin all from Gibco) 1:1 mixed with 2.4% Avicel (RC-581, FMC biopolymer) one

hour later. Cells were fixed using 6% paraformaldehyde (Sigma) 24h later at least 1h at RT

and stained with crystal violet (Sigma). PFU were counted from each dilution in order to

determine the infectious titer (in PFU/mL) for each sample.

Comparison of ACE-2 orthologues' protein sequences

For the species *Homo sapiens*, *Chlorocebus aethiops, Rhinolophus ferrumequinum*, *Mustela*

*viso*n, *Mustela putorius furo*, *Vulpes vulpes*, *Erinaceus europaeus* and *Oryctolagus cuniculus*,

ACE-2 amino-acid sequences were available in NCBI. *Pipistrellus pipistrellus* ACE-2 protein

sequence has been kindly shared by Joanna Damas and Harris A Lewin (37), obtained from

an original sequence from the Zoonomia Consortium. For the species *Crocidura suaveolens*,

Rhinolophus clivosus and *Rhinolophus euryale* the protein sequences obtained in this study

were predicted from ACE-2 nucleic sequences after RNA trizol-extraction from cell lysates

and amplification by RT-PCR using primers (Microsynth) listed in Table S2. The protein

sequence for *Myodes glareolus*, was predicted from a reference transcriptome (62). Contrarily to other species, ACE-2 from *Crocidura suaveolens* could only be partially sequenced but all key residues were be identified (63). Of note, one sequence from one individual (whom the initial cell line originated from) was determined for each species. Polymorphisms, as described for Rhinolophus, were not considered in this study (64). Multiple ACE-2 polymorphisms were indeed observed only in *Rhinolophus euryale* and only the major amino-acid is shown in the table.

Acknowledgements

The authors acknowledge Erik Boehm for help with editing the manuscript and Pascal Cherpillod for facilitating the work in the BSL-3 laboratory at Geneva University Hospitals. We thank Joanna Damas and Harris A Lewin from the University of California for kindly providing the amino-acid sequence of *Pipistrellus pipistrellus* species (obtained from an original sequence from the Zoonomia Consortium). A549-ACE2 and VeroE6/TMPRSS2 cell lines were kindly provided by Pr Mirco Schmolke and Dr Beryl Sanchez at the University of Geneva.

Funding

This work was supported by the HUG Private Foundation, the Pictet Foundation and the Swiss National Science Foundation under the Special call for coronavirus research (Nr. 196644 and 196383).

Author Bio

Dr Manel Essaidi-Laziosi is a Research and Teaching Fellow at the Faculty of Medicine of Geneva, Department of Microbiology and Molecular Medicine. Her main research interests are the study of the interactions of respiratory viruses with host cells, including differentiated human airway epithelia, during *in-vitro* and *ex-vivo* viral infections.

References

1.	Organization WH. WHO Covid Dashboard. 2022 [cited; Available from: https://covid19.who.int

2.	Zhou B, Thao TTN, Hoffmann D, Taddeo A, Ebert N, Labroussaa F, et al. SARS-CoV-2 spike D614G change enhances replication and transmission. Nature. 2021 Apr;592(7852):122-7.

3.	Plante JA, Liu Y, Liu J, Xia H, Johnson BA, Lokugamage KG, et al. Spike mutation D614G alters SARS-CoV-2 fitness. Nature. 2021 Apr;592(7852):116-21.

4.	Organization WH. Tracking SARS-CoV-2 variants. 2022.

5.	Davies NG, Abbott S, Barnard RC, Jarvis CI, Kucharski AJ, Munday JD, et al. Estimated transmissibility and impact of SARS-CoV-2 lineage B.1.1.7 in England. Science. 2021 Apr 9;372(6538).

6.	Tegally H, Wilkinson E, Giovanetti M, Iranzadeh A, Fonseca V, Giandhari J, et al. Detection of a SARS-CoV-2 variant of concern in South Africa. Nature. 2021 Apr;592(7854):438-43.

7.	Faria NR, Mellan TA, Whittaker C, Claro IM, Candido DDS, Mishra S, et al. Genomics and epidemiology of the P.1 SARS-CoV-2 lineage in Manaus, Brazil. Science. 2021 May 21;372(6544):815-21.

8.	Mlcochova P, Kemp SA, Dhar MS, Papa G, Meng B, Ferreira I, et al. SARS-CoV-2 B.1.617.2 Delta variant replication and immune evasion. Nature. 2021 Nov;599(7883):114-9.

9.	Viana R, Moyo S, Amoako DG, Tegally H, Scheepers C, Althaus CL, et al. Rapid epidemic expansion of the SARS-CoV-2 Omicron variant in southern Africa. Nature. 2022 Jan 7.

10.	Cele S, Jackson L, Khoury DS, Khan K, Moyo-Gwete T, Tegally H, et al. Omicron extensively but incompletely escapes Pfizer BNT162b2 neutralization. Nature. 2022 Feb;602(7898):654-6.

11.	Araf Y, Akter F, Tang YD, Fatemi R, Parvez MSA, Zheng C, et al. Omicron variant of SARS-CoV-2: Genomics, transmissibility, and responses to current COVID-19 vaccines. J Med Virol. 2022 May;94(5):1825-32.

12.	Voloch CM, da Silva Francisco R, Jr., de Almeida LGP, Cardoso CC, Brustolini OJ, Gerber AL, et al. Genomic characterization of a novel SARS-CoV-2 lineage from Rio de Janeiro, Brazil. J Virol. 2021 Mar 1.

13.	Markov PV, Katzourakis A, Stilianakis NI. Antigenic evolution will lead to new SARS-CoV-2 variants with unpredictable severity. Nat Rev Microbiol. 2022 Mar 14.

14.	Burki T. The origin of SARS-CoV-2 variants of concern. Lancet Infect Dis. 2022 Feb;22(2):174-5.

15.	Meekins DA, Gaudreault NN, Richt JA. Natural and Experimental SARS-CoV-2 Infection in Domestic and Wild Animals. Viruses. 2021 Oct 4;13(10).

16.	Cele S, Karim F, Lustig G, San JE, Hermanus T, Tegally H, et al. SARS-CoV-2 prolonged infection during advanced HIV disease evolves extensive immune escape. Cell Host Microbe. 2022 Feb 9;30(2):154-62 e5.

17.	Weigang S, Fuchs J, Zimmer G, Schnepf D, Kern L, Beer J, et al. Within-host evolution of SARS-CoV-2 in an immunosuppressed COVID-19 patient as a source of immune escape variants. Nat Commun. 2021 Nov 4;12(1):6405.

18.	Nussenblatt V, Roder AE, Das S, de Wit E, Youn JH, Banakis S, et al. Year-long COVID-19 infection reveals within-host evolution of SARS-CoV-2 in a patient with B cell depletion. J Infect Dis. 2021 Dec 23.

19.	Organization WH. [cited 08.12.2021]; Available from: https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/

20.	Ulrich L, Halwe NJ, Taddeo A, Ebert N, Schon J, Devisme C, et al. Enhanced fitness of SARS-CoV-2 variant of concern Alpha but not Beta. Nature. 2021 Dec 22.

21.	Hui KPY, Ho JCW, Cheung MC, Ng KC, Ching RHH, Lai KL, et al. SARS-CoV-2 Omicron variant replication in human bronchus and lung ex vivo. Nature. 2022 Mar;603(7902):715-20.

22.	Shuai H, Chan JF, Hu B, Chai Y, Yuen TT, Yin F, et al. Attenuated replication and pathogenicity of SARS-CoV-2 B.1.1.529 Omicron. Nature. 2022 Mar;603(7902):693-9.

23.	Han Y, Yang L, Lacko LA, Chen S. Human organoid models to study SARS-CoV-2 infection. Nat Methods. 2022 Apr;19(4):418-28.

24.   Eckerle I, Corman VM, Muller MA, Lenk M, Ulrich RG, Drosten C. Replicative Capacity of MERS
Coronavirus in Livestock Cell Lines. Emerg Infect Dis. 2014 Feb;20(2):276-9.
25.   Eckerle I, Ehlen L, Kallies R, Wollny R, Corman VM, Cottontail VM, et al. Bat airway epithelial
cells: a novel tool for the study of zoonotic viruses. PLoS One. 2014;9(1):e84679.
26.   Eckerle I, Lenk M, Ulrich RG. More novel hantaviruses and diversifying reservoir hosts--time for
development of reservoir-derived cell culture models? Viruses. 2014 Feb 26;6(3):951-67.
27.   Ehlen L, Todtmann J, Specht S, Kallies R, Papies J, Muller MA, et al. Epithelial cell lines of the
cotton rat (Sigmodon hispidus) are highly susceptible in vitro models to zoonotic Bunya-, Rhabdo-, and
Flaviviruses. Virol J. 2016 May 4;13:74.
28.   Planas D, Saunders N, Maes P, Guivel-Benhassine F, Planchais C, Buchrieser J, et al.
Considerable escape of SARS-CoV-2 Omicron to antibody neutralization. Nature. 2022
Feb;602(7898):671-5.
29.   Bekliz M, Adea K, Vetter P, Eberhardt CS, Hosszu-Fellous K, Vu DL, et al. Neutralization capacity
of antibodies elicited through homologous or heterologous infection or vaccination against SARS-CoV-
2 VOCs. Nat Commun. 2022 Jul 4;13(1):3840.
30.   Puhach O, Adea K, Hulo N, Sattonnet P, Genecand C, Iten A, et al. Infectious viral load in
unvaccinated and vaccinated individuals infected with ancestral, Delta or Omicron SARS-CoV-2. Nat
Med. 2022 Apr 8.
31.   Suzuki R, Yamasoba D, Kimura I, Wang L, Kishimoto M, Ito J, et al. Attenuated fusogenicity and
pathogenicity of SARS-CoV-2 Omicron variant. Nature. 2022 Mar;603(7902):700-5.
32.   Peacock TP, Brown JC, Zhou J, Thakur N, Sukhova K, Newman J, et al. The altered entry pathway
and antigenic distance of the SARS-CoV-2 Omicron variant map to separate domains of spike protein.
bioRxiv. 2022:2021.12.31.474653.
33.   V'Kovski P, Gultom M, Kelly JN, Steiner S, Russeil J, Mangeat B, et al. Disparate temperature-
dependent virus-host dynamics for SARS-CoV-2 and SARS-CoV in the human respiratory epithelium.
PLoS Biol. 2021 Mar;19(3):e3001158.
34.   Chan JF, Siu GK, Yuan S, Ip JD, Cai JP, Chu AW, et al. Probable Animal-to-Human Transmission
of SARS-CoV-2 Delta Variant AY.127 Causing a Pet Shop-Related COVID-19 Outbreak in Hong Kong. Clin
Infect Dis. 2022 Mar 2.
35.   Abdel-Moneim AS, Abdelwhab EM. Evidence for SARS-CoV-2 Infection of Animal Hosts.
Pathogens. 2020 Jun 30;9(7).
36.   Hale VL, Dennis PM, McBride DS, Nolting JM, Madden C, Huey D, et al. SARS-CoV-2 infection in
free-ranging white-tailed deer. Nature. 2022 Feb;602(7897):481-6.
37.   Damas J, Hughes GM, Keough KC, Painter CA, Persky NS, Corbo M, et al. Broad host range of
SARS-CoV-2 predicted by comparative and structural analysis of ACE2 in vertebrates. Proc Natl Acad
Sci U S A. 2020 Sep 8;117(36):22311-22.
38.   Luan J, Lu Y, Jin X, Zhang L. Spike protein recognition of mammalian ACE2 predicts the host
range and an optimized ACE2 for SARS-CoV-2 infection. Biochem Biophys Res Commun. 2020 May
21;526(1):165-9.
39.   Lam SD, Bordin N, Waman VP, Scholes HM, Ashford P, Sen N, et al. SARS-CoV-2 spike protein
predicted to form complexes with host receptor protein orthologues from a broad range of mammals.
Sci Rep. 2020 Oct 5;10(1):16471.
40.   Montagutelli X, van der Werf S, Rey FA, Simon-Loriere E. SARS-CoV-2 Omicron emergence
urges for reinforced One-Health surveillance. EMBO Mol Med. 2022 Mar 7;14(3):e15558.
41.   Menachery VD, Yount BL, Jr., Debbink K, Agnihothram S, Gralinski LE, Plante JA, et al. A SARS-
like cluster of circulating bat coronaviruses shows potential for human emergence. Nat Med. 2015
Dec;21(12):1508-13.
42.   Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak associated
with a new coronavirus of probable bat origin. Nature. 2020 Mar;579(7798):270-3.
43.   Lau SKP, Wong ACP, Luk HKH, Li KSM, Fung J, He Z, et al. Differential Tropism of SARS-CoV and
SARS-CoV-2 in Bat Cells. Emerg Infect Dis. 2020 Dec;26(12):2961-5.

44.   Auerswald H, Low DHW, Siegers JY, Ou T, Kol S, In S, et al. A Look inside the Replication
Dynamics of SARS-CoV-2 in Blyth's Horseshoe Bat (Rhinolophus lepidus) Kidney Cells. Microbiol Spectr.
2022 Jun 29;10(3):e0044922.
45.   Aicher S-M, Streicher F, Chazal M, Planas D, Luo D, Buchrieser J, et al. Species-Specific
Molecular Barriers to SARS-CoV-2 Replication in Bat Cells. Journal of Virology.0(0):e00608-22.
46.   Muth D, Corman VM, Roth H, Binger T, Dijkman R, Gottula LT, et al. Attenuation of replication
by a 29 nucleotide deletion in SARS-coronavirus acquired during the early stages of human-to-human
transmission. Sci Rep. 2018 Oct 11;8(1):15177.
47.   Mykytyn AZ, Lamers MM, Okba NMA, Breugem TI, Schipper D, van den Doel PB, et al.
Susceptibility of rabbits to SARS-CoV-2. Emerg Microbes Infect. 2021 Dec;10(1):1-7.
48.   Oreshkova N, Molenaar RJ, Vreman S, Harders F, Oude Munnink BB, Hakze-van der Honing RW,
et al. SARS-CoV-2 infection in farmed minks, the Netherlands, April and May 2020. Euro Surveill. 2020
Jun;25(23).
49.   Yan H, Jiao H, Liu Q, Zhang Z, Xiong Q, Wang BJ, et al. ACE2 receptor usage reveals variation in
susceptibility to SARS-CoV and SARS-CoV-2 infection among bat species. Nat Ecol Evol. 2021
May;5(5):600-8.
50.   Peacock TP, Brown JC, Zhou J, Thakur N, Newman J, Kugathasan R, et al. The SARS-CoV-2
variant, Omicron, shows rapid replication in human primary nasal epithelial cultures and efficiently
uses the endosomal route of entry. bioRxiv. 2022:2021.12.31.474653.
51.   Kim Y, Gaudreault NN, Meekins DA, Perera KD, Bold D, Trujillo JD, et al. Effects of Spike
Mutations in SARS-CoV-2 Variants of Concern on Human or Animal ACE2-Mediated Virus Entry and
Neutralization. Microbiol Spectr. 2022 Jun 29;10(3):e0178921.
52.   Gultom M, Licheri M, Laloli L, Wider M, Strässle M, V'kovski P, et al. Susceptibility of Well-
Differentiated Airway Epithelial Cell Cultures from Domestic and Wild Animals to Severe Acute
Respiratory Syndrome Coronavirus 2. Emerging Infectious Diseases. 2021;27(7):1811-20.
53.   Guo Q, Li M, Wang C, Guo J, Jiang X, Tan J, et al. Predicting hosts based on early SARS-CoV-2
samples and analyzing the 2020 pandemic. Sci Rep. 2021 Aug 31;11(1):17422.
54.   El-Masry I, Dobschuetz Sv, Plee L, Larfaoui F, Yang Z, Song J, et al. Exposure of humans or
animals to SARS-CoV-2 from wild, livestock, companion and aquatic animals. Qualitative exposure
assessment. FAO animal production and health. 2020;Paper 181.
55.   Mollentze N, Keen D, Munkhbayar U, Biek R, Streicker DG. Variation in the ACE2 receptor has
limited utility for SARS-CoV-2 host prediction. bioRxiv. 2022:2022.05.16.492068.
56.   Huang C, Jiang Y, Yan J. Comparative analyses of ACE2 and TMPRSS2 gene: Implications for the
risk to which vertebrate animals are susceptible to SARS-CoV-2. J Med Virol. 2021 Sep;93(9):5487-504.
57.   Essaidi-Laziosi MP, Alvarez CM, Puhach OP, Sattonnet-Roche P, Torriani GP, Tapparel CP, et al.
Sequential infections with rhinovirus and influenza modulate the replicative capacity of SARS-CoV-2 in
the upper respiratory tract. Emerg Microbes Infect. 2021 Dec 21:1-26.
58.   Alcaide M, Rico C, Ruiz S, Soriguer R, Muñoz J, Figuerola J. Disentangling vector-borne
transmission networks: a universal DNA barcoding method to identify vertebrate hosts from arthropod
bloodmeals. PLoS One. 2009;4(9):e7092.
59.   Ivanova NV, Clare EL, Borisenko AV. DNA Barcoding in Mammals. In: Kress WJ, Erickson DL,
editors. DNA Barcodes: Methods and Protocols. Totowa, NJ: Humana Press; 2012. p. 153-82.
60.   Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DK, et al. Detection of 2019 novel
coronavirus (2019-nCoV) by real-time RT-PCR. Eurosurveillance. 2020;25(3):2000045.
61.   Wolfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Muller MA, et al. Virological
assessment of hospitalized patients with COVID-2019. Nature. 2020 May;581(7809):465-9.
62.   Konczal M, Koteja P, Stuglik MT, Radwan J, Babik W. Accuracy of allele frequency estimation
using pooled RNA-Seq. Mol Ecol Resour. 2014 Mar;14(2):381-92.
63.   Fang S, Zheng R, Lei C, Wang J, Zheng R, Li M. Key residues influencing binding affinities of
2019-nCoV with ACE2 in different species. Brief Bioinform. 2021 Mar 22;22(2):963-75.

64.   Guo H, Hu B-J, Yang X-L, Zeng L-P, Li B, Ouyang S, et al. Evolutionary arms race between virus
and host drives genetic diversity in bat severe acute respiratory syndrome-related coronavirus spike
genes. Journal of virology. 2020;94(20):e00902-20.

Tables

Designation	Species of Origin	Species name (latin)*	Mammalian order	Geographic range	Organ of Origin
RF Kidney	Greater horseshoe bat	*Rhinolophus ferrumequinum*	*Chiroptera*	Europe	Kidney
RE Kidney	Mediterranean horseshoe bat	*Rhinolophus euryale*	*Chiroptera*	Europe	Kidney
RE Trachea	Mediterranean horseshoe bat	*Rhinolophus euryale*	*Chiroptera*	Europe	Trachea
RCL.B Trachea	Geoffroy's horseshoe bat	*Rhinolophus clivosus*	*Chiroptera*	Africa	Trachea
Pip Trachea	Common pipistrelle	*Pipistrellus pipistrellus*	*Chiroptera*	Europe	Trachea
NE-R	American mink	*Mustela vison*	*Carnivora*	America, introduced in other areas of the world	Fœtus
MV1-Lu	American mink	*Mustela vison*	*Carnivora*	America, introduced in other areas of the world	Lung
FUFE-R Fbr	Red fox	*Vulpes vulpes*	*Carnivora*	Europe	Embryo
Crocidura	Lesser white-toothed shrew	*Crocidura suaveolens*	*Soricomorpha*	Europe	Trachea
Igel 2 REC.B	European hedgehog	*Erinaceus europaeus*	*Erinaceomorpha*	Europe	Kidney
Mygla AEC.B	Bank vole	*Myodes glareolus*	*Rodentia*	Europe	Trachea
MPFLU-1-R	Domestic ferret	*Mustela putorius furo*	*Carnivora*	Europe	Lung
MPFN-1-R	Domestic ferret	*Mustela putorius furo*	*Carnivora*	Europe	Kidney
KAN-2-R	Rabbit	*Oryctolagus cuniculus*	*Lagomorpha*	Europe	Lung
RK-13	Rabbit	*Oryctolagus cuniculus*	*Lagomorpha*	Europe	Kidney
ZP	Rabbit	*Oryctolagus cuniculus*	*Lagomorpha*	Europe	Embryo
R9ab	Rabbit	*Oryctolagus cuniculus*	*Lagomorpha*	Europe	Kidney
A549	Human	*Homo sapiens*	*Primates*	Global	Lung
VERO-E6	African green monkey	*Chlorocebus aethiops*	*Primates*	Africa	Kidney

519 **Table 1:** List of cell lines used in the study. *confirmed by the sequencing of the cytochrome C subunit oxidase I gene.

Species	Accession	% ACE-2 Identity**	Key residues (KR) for binding to spike RBD																									% KR Identity*
			19	24	27	28	30	31	34	35	37	38	41	42	45	53	79	82	83	90	322	329	330	353	354	355	357	393	
Homo sapiens	BAB40370	-	S	Q	T	F	D	K	H	E	E	D	Y	Q	L	N	L	M	Y	N	N	E	N	K	G	D	R	R	-
Chlorocebus aethiops	AAY57872	94.53	S	Q	T	F	D	K	H	E	E	D	Y	Q	L	N	L	M	Y	N	N	E	N	K	G	D	R	R	100.00
Rhinolophus ferrumequinum	ADN93470	80.37	S	L	K	F	D	D	S	E	E	N	H	Q	L	N	L	N	F	N	N	N	N	K	G	D	R	R	65.38
Rhinolophus euryale	OM256535*	80.62	S	L	T	F	D	N	T	E	E	N	H	Q	L	N	L	N	F	N	K	N	N	K	G	D	R	R	65.38
Rhinolophus clivosus	OM256534*	80.87	S	L	K	F	D	D	S	E	E	N	H	Q	L	N	L	N	F	N	N	N	N	K	G	D	R	R	65.38
Pipistrellus pipistrellus	n.a.	#	S	E	D	F	D	K	S	E	E	N	H	E	L	N	R	A	F	N	S	N	N	E	D	D	R	R	50.00
Mustela vison	QPL12211	82.86	S	L	T	F	E	K	Y	E	E	E	Y	Q	L	N	H	T	Y	D	N	Q	N	K	H	D	R	R	65.38
Mustela putorius furo	BAE53380	82.48	S	L	T	F	E	K	Y	E	E	E	Y	Q	L	N	H	T	Y	D	N	Q	N	K	R	D	R	R	65.38
Vulpes vulpes	XP_025842512	83.35	S	L	T	F	E	K	Y	E	E	E	Y	Q	L	N	L	T	Y	D	N	E	N	K	G	D	R	R	76.92
Crocidura suaveolens	OM256536*	#	L	N	K	F	E	N	R	E	E	N	Y	E	L	N	I	N	F	D	N	K	N	K	G	D	R	R	50.00
Erinaceus europaeus	XP_007538670	78.88	T	E	K	F	D	D	R	Q	E	N	Y	E	L	N	T	N	Y	N	S	N	N	N	G	D	R	R	50.00
Myodes glareolus	n.a.	83.23	S	D	K	F	D	K	Q	E	E	D	Y	Q	L	N	I	S	Y	D	H	E	N	K	D	D	R	R	69.23
Oryctolagus cuniculus	QHX39726	84.85	S	L	T	F	E	K	Q	E	E	D	Y	Q	L	N	L	T	Y	N	S	E	N	K	R	D	R	R	76.92

Table 2: Comparison of ACE-2 key residues for SARS-CoV-2 binding in animal cells. Full length and key residues of ACE-2 protein sequences were compared between the different animal species including human. Eight sequences were already available. Sequences from *Rhinolophus euryale*, *Rhinolophus clivosus* and *Crocidura suaveolens* were newly added in this work (cf. accession codes). *sequence obtained in this study **percentage of identity of ACE-2 full sequence or key residues (KR) *versus H. sapiens*. #: partial sequences. n.a.: no accession number available. Differences compared to the human sequence are highlighted in light grey.

Viral RNA

Viral titer

A

in HAE at 37°C

in HAE at 37°C

B

in HAE at 33°C

in HAE at 33°C

Viral RNA

A

in Calu3 at 37°C

in Calu3 37°C

Viral titer

B

in Calu3 33°C

in Calu3 33°C

Legend (both panels):
- B1 (ancestral)
- Alpha
- Beta
- Gamma
- Zeta
- Delta
- Omicron

A

B1 (ancestral SARS-CoV-2)

B

Delta variant

C

Omicron variant

Bats   Minks and foxes   Rodents and insectivores   Ferret   Rabbit

A B1 (ancestral SARS-CoV-2) **** ****

B Delta variant

C Omicron variant

Bats Minks and foxes Rodents and insectivores Ferret Rabbit