Patterns of siderophore production and utilization at Station ALOHA from the surface to mesopelagic waters

Randelle M. Bundy[1,*] and Lauren E. Manck[2,*], Daniel J. Repeta[3], Matthew J. Church[2], Nicholas J. Hawco[4], Rene M. Boiteau[5], Jiwoon Park[1], Edward F. DeLong[4], Mak A. Saito[3]

[1]University of Washington, School of Oceanography, Seattle, WA

[2]University of Montana, Flathead Lake Biological Station, Polson, MT

[3]Woods Hole Oceanographic Institution, Marine Chemistry and Geochemistry, Woods Hole, MA

[4]University of Hawaii at Manoa, Center for Microbial Oceanography Research and Education, Honolulu, HI

[5]Oregon State University, College of Earth Ocean and Atmospheric Sciences, Corvallis, OR

[*]co-first authors and corresponding authors rbundy@uw.edu and lauren.manck@flbs.umt.edu

Running head (40 characters): Siderophore biosynthesis at Station ALOHA

Scientific Significance Statement

Microbial community production in the subtropical oligotrophic North Pacific is limited by

macronutrients such as nitrogen. However, dissolved iron is another important micronutrient that

has seasonal inputs from dust and passing eddies, keeping the availability of iron low and

episodic. Little attention has been paid to the microbial strategies for dealing with low iron to

support primary production in the oligotrophic ocean, or how limited iron availability impacts

the processing of sinking particulate organic carbon in this region. In this study, we explore iron

cycling including siderophore production and uptake by the microbial community throughout the

water column at Station ALOHA to examine how the microbial community adapts and responds

to changing iron and carbon availability on seasonal timescales.

Author contribution statement: RBundy and LEM wrote the manuscript and completed the

analyses. RBoiteau and JP contributed data and edited the manuscript. EFD, MJC, MAS and

DJR helped design the study and edit the manuscript.

Abstract

The North Pacific subtropical gyre is a globally important contributor to carbon uptake and an

oligotrophic ecosystem primarily limited by nitrogen. The microbial community is also

seasonally exposed to low iron due to biological consumption and seasonally variable iron

delivery. In this study, we examined changes in iron uptake rates, dissolved siderophore

concentrations, and siderophore biosynthesis at Station ALOHA across time (2013-2016) and

depth (surface to 500 m) to observe changes in iron acquisition and internal cycling by the

microbial community. The genetic potential for siderophore biosynthesis was widespread

throughout the upper water column, and biosynthetic gene clusters peaked in spring and summer

along with siderophore concentrations, suggesting changes in nutrient delivery, primary

production, and carbon export impact iron acquisition over the seasonal cycle. Dissolved iron

turnover times, calculated from iron-amended experiments conducted using surface (15 m) and

mesopelagic (300 m) waters, ranged from 9-252 days. The shortest average turnover times at

both depths were associated with inorganic iron additions (14 ± 9 days) and the longest with iron

bound to strong siderophores (148 ± 225 days). Uptake rates of siderophore-bound iron were

faster in the mesopelagic waters than in the surface, leading to high Fe:C uptake ratios of

heterotrophic bacteria in the upper mesopelagic. The rapid cycling and high demand for Fe at

300 m suggests differences in microbial metabolism and iron acquisition in the mesopelagic

compared to surface waters. Together, changes in siderophore production and consumption over

the seasonal cycle suggest organic carbon availability impacts iron cycling at Station ALOHA.

Keywords: siderophore, dissolved iron, siderophore biosynthesis, Station ALOHA

Temporal range: 2013-2016

Frequency or sampling interval: seasonal (monthly)

Spatial scale: site-based

1. Introduction

Oligotrophic subtropical gyres comprise the world's largest ocean biome (Longhurst 2010). Microbial communities inhabiting the oligotrophic ocean are exposed to low concentrations of nitrogen, phosphorous (Karl et al. 1997; Fitzsimmons and Boyle 2014; Letelier et al. 2019), and dissolved iron (Fitzsimmons et al. 2015). Long-term studies by the Hawaii Ocean Time-series (HOT) at Station ALOHA in the North Pacific subtropical gyre demonstrate how seasonal to interannual-scale changes in nutrient supply can impact microbial communities and carbon export (Karl and Lukas 1996; Letelier et al. 2019). The resulting time series captures both the relatively stable conditions of the oligotrophic ocean and the importance of episodic and seasonal-scale delivery of nutrients to the upper ocean from storms, eddies, dust inputs, and the shoaling and deepening of the pycnocline (DiTullio and Laws 1991; Fitzsimmons et al. 2015; Hayes et al. 2015; Pinedo-González et al. 2020; Hawco et al. 2022). Although this ecosystem is characterized by perennial nitrogen limitation, the microbial community of the euphotic zone is also consistently exposed to low dissolved iron (dFe) concentrations, particularly in the deep chlorophyll maximum (DCM; (Hogle et al. 2022, Hawco et al. 2022)), and on sub-decadal timescales, control of primary production as supported by nitrogen (N_2) fixation is thought to oscillate between phosphate and iron availability (Letelier et al. 2019).

In seawater, a majority of dFe is complexed by a pool of organic ligands (FeL) that have a high diversity of chemical structures. Complexation by organic ligands allows dFe to accumulate to concentrations above that set by its inorganic solubility (Gledhill and Buck 2012) and may help to retain Fe within the euphotic zone where it is needed to support primary production (Tortell et

al. 1999). However, ligands significantly alter the bioavailability of the dFe pool by sequestering

Fe in organic complexes that require specific cellular transport systems in order to be accessed

(Sutak et al. 2020). Changes in the external supply of Fe to the upper ocean at Station ALOHA

have been shown to trigger organic ligand production on the timescale of days (Fitzsimmons et

al. 2015).

Marine microorganisms, including both photoautotrophic and heterotrophic bacteria, have

significant cellular Fe requirements and thus are equipped with molecular strategies for acquiring

Fe from the Fe-scarce marine environment (Hopkinson et al. 2005; Sutak et al. 2020). A class of

the strongest organic Fe-binding ligands, called siderophores, are secreted by marine bacteria and

fungi in response to Fe deficiency as a high-affinity uptake strategy to solubilize and stabilize Fe

in a molecular form that can be taken up by dedicated membrane transporters (Sandy and Butler

2009; Vraspir and Butler 2009). Many structurally diverse siderophores have been identified, and

they are biosynthesized by two classes of enzymatic pathways; non-ribosomal peptide synthetase

(NRPS) pathways and/or NRPS-independent siderophore (NIS) synthetase pathways (Sandy and

Butler 2009; Hider and Kong 2010). Siderophore biosynthesis is regulated by Fe concentrations,

where elevated Fe suppresses synthesis (Sandy and Butler 2009). Some microbes can both

synthesize and internalize siderophores, while others only possess outer membrane receptors for

uptake of exogenous siderophores that they cannot produce (Cordero et al. 2012; Kramer et al.

2020). Internalizing siderophore-bound Fe requires either specific transport pathways or the

ability to extracellularly reduce Fe(III) to Fe(II), releasing it from the siderophore complex; thus

Fe complexed to siderophores has low bioavailability to the microbial community as whole

(Shaked and Lis 2012), but is readily bioavailable to microbes that possess the relevant transport

systems.

Dissolved siderophores represent 2-10% of the total dissolved Fe-binding ligand pool present at

Station ALOHA, and they likely play an important role in the competition for Fe and in

solubilization of Fe from particles (Bundy et al. 2018). North of Station ALOHA in the North

Pacific transition zone, where Fe inputs from dust in the spring are elevated (Pinedo-González et

al. 2020), siderophore concentrations and the transcription of siderophore biosynthesis and

uptake genes are also elevated, further connecting the production of siderophores as a

mechanism for solubilizing Fe from particles (Park et al. 2023). The metabolic potential for

siderophore production and uptake is prevalent in the North Pacific subtropical gyre, transition

zone, and the subpolar high nutrient low chlorophyll region (Hogle et al. 2022; Park et al. 2023).

Although the importance of siderophore production as a microbial Fe acquisition strategy in the

marine environment has been recognized for some time (Mawji et al. 2008; Boiteau et al. 2016,

2019; Bundy et al. 2018; Moore et al. 2021; Manck et al. 2022; Park et al. 2023), little is known

about their seasonal variability or turnover times. Understanding the biosynthesis and turnover

times of siderophore-bound Fe will be important for interpreting their bioavailability and

influence on Fe recycling and retention in the upper ocean. To address these gaps in our

understanding of siderophore biogeochemistry in the marine environment, we explored changes

in siderophore distributions, biosynthesis potential and turnover times at Station ALOHA across

time and depth. By placing this study in the rich context of biogeochemical measurements made

by the HOT program, we aim to better connect siderophore production and uptake to

environmental conditions, in an effort to understand the impact of siderophores on microbial Fe

acquisition and biogeochemical cycling in this important and well-studied oligotrophic

ecosystem.

2. Methods

2.1 Radioactive iron uptake and siderophore bioavailability experiments

Fe uptake experiments were performed at Station ALOHA (Fig. 1A) on two expeditions

(HOT278 in November 2015 and KM1605 in March 2016) at 15 m and 300 m. Four treatments

were used in all experiments: $FeCl_3$ (denoted as "FeL"), ferrioxamine B, ferrioxamine E, and Fe-

amphibactin. Experiments at 15 m were performed both in the light and dark, and experiments at

300 m were performed only in the dark. Fe uptake was measured in the >2.0 µm and 0.2-2.0 µm

size fraction after 12 hours. 500 nmol L^{-1} of nitrate was added to each of the 15 m experiments to

alleviate possible nitrogen limitation, which might have restricted Fe uptake (Table S1).

The isolation of amphibactin ligands used in this experiment has been described in detail

elsewhere (Bundy et al. 2018), and the other ligand standards were commercially available.

Radioactive Fe ligand stocks were prepared using a 1 mCi ^{55}Fe stock (Perkin Elmer; Lot

#091514) with a specific activity of 44.66 mCi mg^{-1} and an activity of 35.55 mCi mL^{-1}. Before

making the ligand stocks, the concentration of the ^{55}Fe stock solution was calculated by

converting the initial activity on the stock date to the current activity using the half-life of ^{55}Fe (λ

= 2.73 years) and the number of days since the stock date. A primary ^{55}Fe stock of 55 µmol L^{-1}

was used to prepare the ^{55}Fe-ligand and $^{55}FeCl_3$ stocks. The $FeCl_3$ stock (FeL treatment) was

Figure 1. (A) Map of the study region including the location of Station ALOHA (red dot). **(B)** Dissolved iron uptake rates (pmol L^{-1} day^{-1}) in the surface (15 m) and mesopelagic (300 m) at Station ALOHA for naturally complexed iron (FeL) and iron bound to the siderophores amphibactin (Amp), ferrioxamine E (DFE), and ferrioxamine B (DFB). Dark and light measurements are displayed for both the 0.2-2.0 µm and >2.0 µm size fraction. Total rates in the bottom panels represent the sum of both size fractions for each treatment. Mean values from all measurements for a given treatment are displayed and error bars represent the standard error of the mean. (B) Turnover time (days) of iron in each treatment at 15 m and 300 m based on rates of total iron uptake in the dark. Box plots display the interquartile range with median values plotted as a solid horizontal line. The y-axis is on a log_{10} scale. Turnover times at 300 m were statistically different from 15 m only in the DFB treatment (pairwise t-test, $p < 0.05$).

prepared in pH 2 (Optima HCl, Fisher Scientific) type I deionized water (MilliQ) while the

siderophore stocks were prepared in type I deionized water. Ligand stocks were equilibrated for

at least 24 hours, but up to 5 days, before use, and ligands were present in three times excess of

^{55}Fe. Duplicate ligand stocks were made with the stable isotope ^{57}Fe (Cambridge Isotopes) to set

up replicate treatments for cell counts that did not contain radioactivity.

Prior to beginning experiments at sea, quench curves were made to account for physical and

chemical quench. Physical quench was accounted for by placing a blank filter in 10 mL of

scintillation cocktail in triplicate and counting for 10 minutes on the liquid scintillation counter.

Physical quench was also assessed by filtering a range of seawater volumes (10-500 mL) onto

separate 0.2 µm filters containing no ^{55}Fe, and then placing each filter into 10 mL of scintillation

cocktail and counting for 10 minutes. These physical quench controls account for whether the

presence of cells or the filter itself impacts the counting efficiency of the liquid scintillation

counter. The physical quench was found to be extremely small (< 1 % of blanks), and for all

further analyses, blank filters were used for quench correction. Chemical quench was assessed by

setting up 10 separate vials of scintillation cocktail, each with the same activity of ^{55}Fe and

increasing acetonitrile additions (0.1-10 nmol L^{-1}). The chemical quench was also found to be

very small (< 5% of the blank) and was incorporated into the counting efficiency of the method

for all samples.

To set up the incubation experiments, trace metal clean seawater collected from X-Niskin bottles

(Ocean Test Equipment) from 15 m and 300 m was dispensed into polycarbonate bottles. Four of

the bottles were spiked with 1 nmol L^{-1} of the ^{57}Fe ligand stocks and the rest were spiked in

duplicate with 1 nmol L^{-1} of the ^{55}Fe ligand stocks. A final concentration of 0.01%

glutaraldehyde was added to four of the bottles to serve as "dead controls" and were incubated

for 30 minutes before the experiment began to account for non-specific ^{55}Fe adsorption. All

bottles were incubated in surface seawater flow-through incubators and dark treatments were

double-bagged in black polyethylene bags. After 12 hours, a flow cytometry sample was taken

from each of the ^{57}Fe treatments and was preserved with 0.01% glutaraldehyde for 30 minutes

before being flash-frozen in liquid nitrogen and stored at -80°C. At the end of the 12 hour

experiment, each bottle was sequentially filtered onto acid washed 2.0 μm and 0.2 μm filters

using low pressure vacuum filtration. Both filters were washed three times with an oxalate wash

to remove extracellular Fe (Tovar-Sanchez 2003).

A key assumption of these experiments is that all Fe remains complexed to the respective added

siderophore throughout the duration of the experiment (12 hours). To confirm this was the case,

the remainder of the treatments spiked with the ^{57}Fe-siderophore stocks were filtered with a 0.2

μm polycarbonate filter, and the filtrate was pre-concentrated onto a solid phase extraction

column (Bond Elut ENV, Agilent Technologies) and were treated and analyzed in the same

manner as the dissolved siderophore samples from the water column as described below. ^{57}Fe

peaks from LC-ICPMS analyses were examined to determine whether any free ^{57}Fe not

associated with each of the added siderophores was present. No significant free ^{57}Fe was

observed in these treatments, confirming the full complexation of each ligand in the siderophore

treatments (data not shown).

Radioactive decay on each filter and in each treatment was determined using a liquid scintillation

counter (Beckman-Coulter LS6500). ^{55}Fe disintegrations per minute (dpm) were background and

dead control corrected and converted to a concentration of Fe uptake using a standard curve. Fe

uptake rates were then calculated based on the total amount of Fe that was incorporated into

biomass onto each filter size fraction and in each treatment. Total Fe uptake over the 12-hour

incubation was converted to total Fe uptake per day (over 24 hours). We then estimated Fe

uptake rates for heterotrophic bacteria specifically, which we assumed to be the dominant

community responsible for any observed Fe uptake in the dark (Table S1). The Fe uptake per

heterotrophic bacteria cell per day (amol Fe cell^{-1} day^{-1}) was estimated using flow cytometry cell

counts of heterotrophic bacteria (in the ^{57}Fe treatments) and the total Fe uptake in the dark (Table

S1). Fe:C ratios (μmol:mol) for heterotrophic bacteria were calculated using a carbon quota of

12.4 fg C cell^{-1} (Strzepek et al. 2005; Boyd et al. 2015) and assuming a growth rate of 1 day^{-1}

(Jones et al. 1996, (Table S1)). As an additional constraint on Fe:C per cell, Fe uptake rates were

compared to average rates of bacterial carbon production measured at or near Station ALOHA –

23±8 nmol C L^{-1} day^{-1} and 0.9±1.3 nmol C L^{-1} day^{-1} at 5 m and 300 m, respectively. The rate of

bacterial production at 5 m is an average value of those measured at Station ALOHA (Viviani

and Church 2017). The mean rate at 300 m was calculated from measurements conducted on

three cruises near Station ALOHA in the North Pacific subtropical gyre – KOK1507, KM1605,

and KM1709. All data are publicly available on the Simons Collaborative Marine Atlas Project.

In each case, leucine incorporation was measured using the methods described by Smith and

Azam (1992) and a conversion factor of 1.5 kg C mol^{-1} leucine was used to convert leucine

incorporation to bacterial carbon production.

In addition to Fe uptake rates, the turnover rate of Fe in each treatment was calculated based on,

(1) $$turnover\ rate = {total\ Fe\ uptake\ rate}/{total\ Fe}$$

where the total Fe uptake rate (pmol L^{-1} day^{-1}) is the sum of the radioactive Fe uptake in both

size fractions and the total Fe is the radioactive Fe added (1 nmol L^{-1}) plus the *in situ* dFe

concentration (Bundy et al. 2018). The turnover time of Fe (in days) for each treatment was

calculated as 1/turnover rate.

2.2 Environmental parameters

Data for concentrations of nitrate+nitrite (NO_3^-+NO_2^-), chlorophyll *a* (chl *a*), rates of primary

production, and carbon flux at Station ALOHA were accessed for the study period, January 1,

2013-December 31, 2016. All data is publicly available via the HOT-Data Organization and

Graphical System. Primary production is routinely estimated based on assimilation of ^{14}C by the

HOT program at depths of 5, 25, 45, 75, 100, and 125 m (Karl et al. 2021). Particulate carbon

flux is measured from sediment traps at 150 m. Additional details on the sampling strategies used

by the HOT program can be found elsewhere (Karl and Lukas 1996) and the analytical methods

used for HOT measurements can be found online

(https://hahana.soest.hawaii.edu/hot/methods/results.html). NO_3^-+NO_2^- concentrations were

determined using the high-sensitivity chemiluminescent method (Dore and Karl 1996; Foreman

et al. 2016). Depth integrated (0-150 m) fluxes and stocks were calculated using trapezoidal

approximations, assuming homogenous mixing between 0-5 m. To integrate rates of primary

production to 150 m, monthly mean rates measured at 150 m during the first 12 years of the

HOT program (Karl et al. 2021) were added to the 0-125 m integrated value calculated during

this study period. The *e*-ratio for each HOT cruise during the study period was calculated as the

rate of particulate carbon flux measured at 150 m divided by the depth integrated (0-150 m) rate

of primary production.

Because samples for dFe (< 0.2 µm) concentrations are not routinely measured by the HOT

program, measurements made from previous studies at Station ALOHA (Fitzsimmons et al.

2015; Bundy et al. 2018) have been reproduced here. These measurements largely fall within the

spring and summer months. Therefore, to visualize potential seasonal changes in dFe

concentrations, 3-day average dFe concentrations at Station ALOHA during the study period

were obtained from the MIT Darwin model (Dutkiewicz et al. 2015) via the Simons

Collaborative Marine Atlas Project (CMAP):

https://simonscmap.com/catalog/datasets/Darwin_Nutrient. Seasons were defined in this study

as – Winter: December, January, February; Spring: March, April, May; Summer: June, July

August; Fall: September, October, November. Additionally, vertical regions were defined here

as: upper euphotic zone: ≤ 75 m, lower euphotic zone: > 75 m to ≤ 150 m, upper mesopelagic

zone: > 150 m to ≤ 300 m, mid-mesopelagic zone: > 300 m to ≤ 500 m.

2.3 Siderophore collection and analyses

Dissolved siderophore samples were collected on six cruises from 2013-2016, each within 100

km of Station ALOHA (Table S2). Samples were collected via Teflon diaphragm pump or 8 L

X-Niskin bottles on a trace metal rosette using a non-metallic line. Samples were filtered (0.2 µm

Acropak 200 or Pall capsule) into acid-cleaned fluorinated high density polyethene 200 L barrels

(expeditions from 2013-2014; Boiteau et al. 2013) or 20 L carboys (all other expeditions; Bundy

et al. 2018). For each sample collected in 2013 and 2014, 20-800 L of filtered seawater was

preconcentrated, and for all other samples 15-20 L of filtered seawater was collected. Filtered

samples were preconcentrated at a flow rate of 15-18 mL min^{-1} onto 1000 mg solid phase

extraction columns containing (Bond Elut ENV, Agilent Technologies). After preconcentration,

each column was flushed with three column volumes of trace metal clean type I deionized water

until dry and then was stored at -20ºC until analysis.

Solid phase extraction columns were slowly thawed in the dark and then eluted with three

column volumes of distilled or Optima grade methanol (Fisher Scientific). Methanol extracts

were dried to approximately 0.5-1 mL using a SpeedVac (Thermo Fisher). Either 25 or 50 µL of

the extract was first injected onto a PEEK-lined C8 column and compounds were separated using

liquid chromatography (LC) on a Dionex 3000 high pressure liquid chromatography (HPLC)

system. Details on the chromatography are presented in Boiteau et al. (2016) and Bundy et al.

(2018). Briefly, compounds were separated using a 20 minute gradient of 95% solvent A/5%

solvent B to 10% solvent A/90% solvent B (solvent A = 5 mM ammonium formate in type I

deionized water, solvent B = 5 mM ammonium formate in distilled methanol), followed by an

isocratic step from 90% to 95% solvent B (Boiteau et al. 2016; Bundy et al. 2018). Fe-bound

compounds eluted from the column were monitored by inductively-coupled plasma mass

spectrometry (ICP-MS) on an iCap-Q. The sample was introduced at a flow rate of

approximately 50 µl min^{-1} and oxygen add gas was introduced at a flow rate of approximately 25

mL min^{-1} to minimize the deposition of reduced carbon on the cones. The ICP-MS was equipped

with a perfluoroalkoxy micronebulizer (PFA-ST; Elemental Scientific), platinum sampler and

skimmer cones, and a cyclonic spray chamber cooled to 0°C. Counts per second of ^{56}Fe, ^{57}Fe,

^{54}Fe and ^{59}Co were monitored in kinetic energy discrimination mode (KED) and distinct peaks in

^{56}Fe and ^{57}Fe were integrated using in-house R scripts. Peak areas were converted to

concentration using a four-point standard curve with ferrioxamine E, and were corrected for

sensitivity changes throughout the run using the peak area of ^{59}Co from a cyanocobalamin

internal standard. All distinct quantifiable Fe peaks were categorized as putative siderophores

and the concentrations of these peaks were summed for each sample to give a total siderophore

concentration (Table S2).

To identify the putative siderophores, samples were then analyzed using LC coupled to

electrospray ionization mass spectrometry (ESI-MS) on an Orbitrap Fusion (Thermo Scientific).

The same chromatography was used, and the ICP-MS and ESI-MS data were aligned based on

the retention time of the ^{59}Co peak from cyanocobalamin in the ICP-MS trace and the presence

of the cyanocobalamin m/z in the ESI-MS data ($m/z = 678$). Putative siderophores were identified

using a targeted search of the MS^1 m/z based on a database of known siderophores (Baars et al.

2014). Fragmentation data (MS^2) was then examined and compared to literature values or in-

silico fragmentation prediction (CFM-ID 3.0), and siderophores that had a matching MS^2

fragment to literature values or in-silico prediction were assigned a structural identification

(Table S2).

2.4 Siderophore biosynthetic gene cluster analysis

Details on the sampling, processing and data analyses of metagenomes collected at Station

ALOHA are presented elsewhere (Mende et al. 2017; Luo et al. 2020). The ALOHA gene

catalog consists of ~8.9 million non-redundant genes detected from concentrated plankton

biomass (0.22-1.6 μm) collected between the surface and 1,000 m over ~1.5 years at near

monthly frequency (Mende et al. 2017). To identify putative siderophore biosynthetic gene

clusters (BGCs) within the ALOHA gene catalog, biomarker genes for both NRPS pathways and

NIS pathways were first identified. To identify NIS pathways in the ALOHA gene catalog,

hidden Markov models (HMMs) for the conserved domains pfam04183 (IucA/IucC siderophore

synthases) and pfam06276 (FhuF-like reductases) were searched for in the ALOHA gene catalog

using HMMER (Eddy 2011) which resulted in the detection of 20 NIS homologs (Table S3). NIS

pathways are specific to siderophore biosynthesis, hence the 20 identified NIS homologs could

be confidently attributed to siderophore biosynthesis. For NRPS biosynthetic pathways, we

searched for the conserved domain pfam00501, an AMP-binding domain inclusive of the

adenylation domain of NRPS modules. The resulting 28,000+ genes were then screened with

antiSMASH (Medema et al. 2011) to determine the presence of a complete or near-complete

NRPS module with adenylation, peptidyl carrier protein (pfam00550), and condensation

(pfam00668) domains. This resulted in the detection of 132 NRPS homologs. Because NRPS

biosynthetic pathways are not unique to siderophore biosynthesis, further examination of the

identified NRPS homologs was required. Each NRPS homolog, along with neighboring genes

from the ALOHA gene catalog contig assembly, were analyzed using the predictive tools within

the antiSMASH pipeline with default parameters for the relaxed detection strictness setting. In

addition to the default features, MIBiG cluster comparison and Cluster pfam analysis were also

performed. This enabled predictions of the amino acid specificity for each adenylation domain

within the NRPS modules, detection of additional conserved domains, and BLAST comparisons

to characterized biosynthetic gene clusters. This resulted in the putative identification of 18

NRPS pathways for the biosynthesis of siderophores from the ALOHA gene catalog (Table S3).

Combined, a total of 38 putative siderophore BGCs consisting of 101 unique genes were

identified and used for downstream analysis (Table S3). The antiSMASH pipeline was utilized to

assign putative structural characterizations to these siderophore BGCs based on comparisons to

known biosynthetic pathways. We note that while this is a conservative approach to identifying

siderophore BGCs that may exclude siderophores without well characterized biosynthetic

pathways, it has resulted in a list of genes with high percent identity to known siderophore

biosynthesis genes, giving us confidence in the functional annotation of genes used for

downstream analysis.

To examine the distribution of siderophore BGCs with depth and across time at Station ALOHA,

metagenomic sequence reads from plankton biomass (> 0.2 µm) collected on HOT cruises

between May 2015 and April 2016 (NCBI BioProject PRJNA352737) were mapped to open

reading frames (ORFs) of the ALOHA gene catalog. In total, this consisted of 132 samples from

5-500 m, spanning 11 months. Forward and reverse reads from each sample were quality

trimmed with Trimmomatic (Bolger et al. 2014) and mapped to the ALOHA gene catalog using

BWA (Li and Durbin 2009) with a 95% nucleotide identity threshold and minimum alignment

length of 45 bp. Reads aligned as proper pairs were then counted with FeatureCounts (Liao et al.

2014) and multiple mappers were fractionally distributed. Finally, read counts were normalized

to reads per kilobase per million (RPKM) for further analysis. Reads from the 2015-2016 dataset

recruited to 24 of the 38 siderophore BGCs identified in the ALOHA gene catalog (Table S3).

3. Results

3.1 Uptake and turnover time of siderophore-bound iron by the microbial community

Uptake rates of inorganic Fe were the highest across all treatments (Fig. 1B, Table S1). The

inorganic Fe treatments were denoted as FeL, since the added $FeCl_3$ likely associated with

unbound natural ligands present in excess of the in situ dFe in the experiment (Bundy et al.

2018). Uptake rates of FeL at both 15 m and 300 m were higher on average in the 0.2-2.0 μm

size fraction compared to the >2.0 μm size fraction. In the >2.0 μm size fraction, uptake rates of

FeL at 15 m were comparable in the light and the dark (~36 pmol L^{-1} day^{-1}). In contrast, FeL

uptake was higher (on average) in the light (85 pmol L^{-1} day^{-1}) compared to the dark (50 pmol L^{-1}

day^{-1}) in the 0.2-2.0 μm size fraction (Fig. 1B).

Siderophore-bound Fe was readily accessible by the microbial community at 15 m and 300 m at

Station ALOHA (Fig. 1B, Table S1). The relative uptake rate varied for the different Fe-

siderophore treatments. At both depths, the Fe-amphibactin amendment was taken up the fastest

(54 pmol L^{-1} day^{-1}), followed by ferrioxamine E (19 pmol L^{-1} day^{-1}) and ferrioxamine B (9 pmol

L^{-1} day^{-1}). In contrast to FeL, the uptake rate of siderophore-bound Fe was generally higher in the

dark (33 pmol L^{-1} day^{-1}) compared to the light (14 pmol L^{-1} day^{-1}). Similar to the FeL treatment,

Fe uptake was higher on average in the 0.2-2.0 μm fraction compared to the >2.0 μm fraction

across all siderophore treatments (Fig. 1B).

Surprisingly, the uptake rate of siderophore-bound Fe was higher at 300 m compared to 15 m

leading to shorter Fe turnover times (Fig. 1C). In general, at both 15 m and 300 m the turnover

times were shortest in FeL treatments, followed by Fe bound to amphibactins, ferrioxamine E

and ferrioxamine B. The average turnover time of Fe at 15 m ranged from 15±8 days in the FeL

treatment to 252±376 and 215±165 days in the ferrioxamine E and ferrioxamine B treatments,

respectively. Mean turnover times at 300 m ranged from 9±4 days in the FeL treatment to 65±4

days in the ferrioxamine B treatment (Fig. 1C). Across all treatments, turnover times of Fe were

consistently shorter at 300 m compared to 15 m (Fig. 1C), though this difference was only

significant in the ferrioxamine B treatment (pairwise t-test, $p < 0.05$).

The Fe uptake rate per cell (amol Fe day^{-1} cell^{-1}) was calculated from the total Fe uptake rate.

The Fe uptake rates in the dark from the 0.2-2.0 and >2.0 µm size fractions were summed for a

given Fe substrate (Table S1) and normalized to the abundance of heterotrophic bacteria

measured via flow cytometry, assuming that the Fe uptake in the dark treatments was dominated

by these organisms. This is likely a valid assumption for the experiments done at 300 m, but for

experiments done at 15 m there could be additional Fe uptake in the dark by phytoplankton. The

resulting Fe uptake per cell at 15 m ranged from 0.01-0.14 amol Fe cell^{-1} day^{-1}, increasing to

0.18-1.46 amol Fe cell^{-1} day^{-1} at 300 m given the higher total rates of Fe uptake and much lower

cell abundances (Table S1).

Cellular Fe:C ratios were calculated in two different ways (see section 2.2). The first, determined

an Fe quota per cell (pmol Fe cell^{-1}) at the end of each experiment assuming a bacterial growth

rate of 1 day^{-1} (Jones et al. 1996), such that the Fe uptake rate per cell is equal to the cellular Fe

quota. These cellular Fe quotas were then converted to Fe:C ratios using a fixed carbon quota of

12.4 fg C cell^{-1} (Strzepek et al. 2005; Boyd et al. 2015). The second method compared total Fe

uptake rates (pmol Fe L^{-1} day^{-1}) to average rates of bacterial carbon production measured at or

near Station ALOHA (23 nmol C L^{-1} day^{-1} and 0.9 nmol C L^{-1} day^{-1} at 15 m and 300 m,

respectively). This represents the ratio at which Fe and C are actively being acquired by bacterial

cells and does not require assumptions of growth rates or cellular carbon quotas, but does

presume a constant leucine:carbon ratio within biomass. At 15 m, the Fe:C per cell calculated

using a fixed carbon quota ranged from 6.8±3.5 µmol:mol (ferrioxamine B treatment) to 68±43

µmol:mol (FeL treatment; Table S1). In contrast, the Fe:C calculated using average rates of

bacterial production ranged from 380±192 µmol:mol to 3775±2386 µmol:mol (Table S1). We

note that assuming a lower heterotrophic bacterial growth rate of 0.05 day^{-1} in surface waters

(Kirchman 2016) with a fixed carbon quota results in higher Fe:C ratios determined by the first

method, and would reduce the discrepancy between these two methods. We however chose to

keep the estimates using a growth rate of 1 day^{-1} as our lower bound since these estimates are

comparable to previous Fe:C values reported previously (Tortell et al. 1999). Irrespective of

which approach was used, the Fe:C per cell was higher at 300 m than 15 m. The Fe:C per cell

calculated from a fixed carbon ratio at 300 m ranged from 88±6 µmol:mol (ferrioxamine B

treatment) to 711±305 µmol:mol (FeL) treatment, while Fe:C per cell based on bacterial

production ranged from 17,044±1131 to 137,500±58,941 µmol:mol in the ferrioxamine B and

FeL treatments, respectively (Table S1). These high Fe uptake rates were observed even after

oxalate washing to remove adsorbed Fe and subtraction of a dead control, indicating that they do

reflect internalization and uptake of Fe and not, for example, authigenic Fe precipitation.

3.2 Seasonal changes in nutrients and productivity at Station ALOHA during the study period

Station ALOHA is a well-studied oligotrophic time-series station in the North Pacific that is

sampled by the HOT program on a nearly monthly basis (Fig. 1A). This region is characterized

by low surface nutrient concentrations and a persistent deep chlorophyll maximum (DCM)

Figure 2. Depth profiles of **(A)** chlorophyll *a* (Chl *a*) concentrations (μg L^{-1}), **(B)** nitrate+nitrite (NO_3^-+NO_2^-) concentrations (nmol L^{-1}), **(C)** dissolved iron (dFe) concentrations (nmol L^{-1}), and **(D)** rates of primary production (mg C m^{-3} day^{-1}) during the 2013-2016 study period at Station ALOHA. Depth profiles in A, B, and C display the mean concentrations at each depth for a given season and error bars represent the standard deviation from the mean. Depth profiles in C display the mean dFe concentrations at each depth across the entire study period for available in situ measurements (open squares) and model output from the MIT Darwin model (closed squares) and error bars represent the standard deviation from the mean.

centered around 100 m (Fig. 2A). During the study period, monthly mean nitrate+nitrite (NO_3^-+NO_2^-) concentrations in the upper euphotic zone ranged from 1.4±0.4 nmol L^{-1} in August to 5.7±0.8 nmol L^{-1} in November (Fig. 2B, S1A). Below 100 m, NO_3^-+NO_2^- concentrations increased rapidly and were more variable than the upper euphotic zone. Between 100-150 m, monthly mean NO_3^-+NO_2^- concentrations during the study period reached minimum values in May (269.8±274.6 nmol L^{-1}) and maximum values in August (1967.3±1858.3 nmol L^{-1}, (Fig. 2B,

S1B)). The unusually high mean $NO_3^-+NO_2$ concentrations observed in August during the study

period, along with the high degree of variability during this month, are likely the result of

strongly negative sea level anomalies present at Station ALOHA during August 2015, increasing

nutrient concentrations in the lower euphotic zone as a result of isopycnal uplift (Barone et al.

2019). Mean monthly depth integrated stocks of $NO_3^-+NO_2^-$ (mmol N m^{-2}) in the upper and

lower euphotic zone followed similar monthly patterns to that of average $NO_3^-+NO_2^-$

concentrations (Fig. S2).

The dFe concentrations are persistently low in surface waters at Station ALOHA. Since limited

dFe data was available during the study period, 3-day averaged dFe concentrations generated

from the MIT-Darwin model were used as a potential means to explore seasonal variation in dFe

concentrations, despite the known limitations of modeling dFe. Model output predicted

minimum concentrations of dFe in the upper euphotic zone with low seasonal variability (Fig.

2C). However, dFe measurements collected during spring and summer months between 2012-

2015 at Station ALOHA suggested dFe concentrations of the upper euphotic zone varied as

much as 10-fold (0.08 to 0.87 nmol L^{-1}), variability that is not captured in the model output (Fig.

2C). This high degree of variability in the upper euphotic zone is likely a result of episodic

atmospheric dust deposition which can impact dFe concentrations in the upper euphotic zone

(Fitzsimmons et al. 2015). Given the lack of agreement between model output and observations

during the spring and summer in the upper euphotic zone, dFe availability in this region of the

water column during fall and winter remains unconstrained without further measurements. Both

model and observational data suggested consistently low (≤ 0.1 nmol L^{-1}) dFe concentrations

between 100-125 m, corresponding with the typical depth of the DCM. Below 125 m, dFe

concentrations began to increase. However, observational data suggest there is a deeper ferricline

than that indicated by model output.

During the study period, rates of primary production ranged from 0.2 – 11.8 mg C m^{-3} day^{-1}

throughout the euphotic zone, with the lowest values observed in winter (Fig. 2D, S1A). Mean

monthly rates in the upper euphotic zone peaked in October (7.8±2.4 mg C m^{-3} day^{-1}) and were

the lowest in February (4.8±1.9 mg C m^{-3} day^{-1}; Fig. S1A). Between 100-150 m, mean monthly

rates of primary production peaked in July (4.2±0.8 mg C m^{-3} day^{-1}) with minimum values

observed in December (1.1±0.3 mg C m^{-3} day^{-1}; Fig, S1B). During the study period, depth

integrated rates of primary production (0-150 m) increased from winter to summer before

decreasing in fall (Fig. 3A). Maximum monthly mean values were observed in June (728.8±87.6

mg C m^{-2} day^{-1}) and minimum values in December (453.4±23.0 mg C m^{-2} day^{-1}). The particulate

carbon flux at 150 m during the same time period showed similar monthly patterns to primary

production (Fig. 3B), but with relatively higher fluxes observed during January and February. As

a result, relatively small changes in the monthly mean *e*-ratio (calculated as the fraction of the

total depth integrated rate of primary production from 0-150 m that is captured in the particulate

carbon flux at 150 m) were detected, but followed similar patterns to those of particulate carbon

flux (Fig. 3C). The highest monthly mean values were observed in February (0.062±0.031) and

August (0.058±0.021) with the lowest in December (0.030±0.009). The average C:N ratio of the

particulate flux at 150 m increased over the summer months from 6.8±0.3 in April to 8.4±1.2 in

August, indicating that carbon-enriched particulate matter contributed to periods of high and

relatively efficient export. However, the overall low *e*-ratios observed across the study period are

consistent with intense remineralization of organic matter within the euphotic zone. Seasonal

patterns in productivity observed during the 2013-2016 study period discussed here are

consistent with those observed across the entire 30 year period of the HOT program (Karl et al.

2021).

Figure 3. Monthly and seasonal mean values of **(A)** depth-integrated rates of primary production
(mg C m^{-2} day^{-1}) from 0-150 m, **(B)** particulate carbon flux (mg C m^{-2} day^{-1}) at 150 m, **(C)** the *e*-
ratio defined as the proportion of particulate carbon flux at 150 m compared to the 0-150 m
depth-integrated rate of primary production, and **(D)** the C:N ratio of sinking particulate matter
collected at 150 m at Station ALOHA during the 2013-2016 study period. Error bars represent
the standard deviation from the mean.

3.3 Seasonal distribution of siderophores and siderophore biosynthesis genes at Station ALOHA

Dissolved siderophore concentrations and the abundance of siderophore BGCs changed

seasonally at Station ALOHA as well as with depth (Fig. 4 and 5). Total dissolved siderophore

concentrations varied from undetectable to 95.3 pmol L^{-1} during the entire study period. Note

that in the fall, siderophore concentrations were only measured at 15 m, limiting assessment of

seasonality in siderophore concentrations in the mesopelagic waters during this season. Average

siderophore concentrations throughout the whole water column were highest in the spring

(21.8±41.5 pmol L^{-1}), fall (8.4±8.1 pmol L^{-1}), and summer (3.0±4.2 pmol L^{-1}) and lower in

winter (0.4±0.6 pmol L^{-1}, Fig. 5). However, high variability was observed within each season. In

the upper euphotic zone where the most samples were collected, siderophore concentrations were

highest on average in the spring (47.8±67.2 pmol L^{-1}), fall (8.4±8.1 pmol L^{-1}), and summer

(2.4±0.1 pmol L^{-1}), followed by winter (0.4±0.3 pmol L^{-1}; Fig. 4, Table S2).

Figure 4. Seasonality in dissolved siderophore concentrations (pmol L^{-1}) at Station ALOHA
between 2013-2016. Bars display the mean values of measurements collected within the same
depth range for a given season and error bars, when present, represent the standard error of the
mean. When no data are present for a given depth range and season, no samples were collected
during that time. Depth ranges are as follows: upper euphotic 0-75 m, lower euphotic > 75 m ≤
150 m, upper mesopelagic > 150 m ≤ 300 m, mid-mesopelagic > 300 m ≤ 500 m.
510

511   The capacity for siderophore production as indicated by the presence of ORFs from siderophore

512   BGCs was detected at Station ALOHA across all seasons and at all depths sampled during the

513   study period (Fig. 5A, Table S3). Similar to dissolved siderophore concentrations, ORFs from

A

B

Figure 5. Seasonal abundance of ORFs (RPKM) from siderophore BGCs detected at Station ALOHA between May 2015-April 2016. **(A)** Box plots display the interquartile range of ORF abundances detected within a given depth range and season. Median values are plotted as solid vertical lines, and outliers are plotted as distinct points. Depth ranges are as follows: upper euphotic 0-75 m, lower euphotic > 75 m ≤ 150 m, upper mesopelagic > 150 m ≤ 300 m, mid-mesopelagic > 300 m ≤ 500 m. **(B)** Box plots display the interquartile range of ORF abundances detected within a given month or season in the euphotic (0-150 m) and mesopelagic (150-500 m) zones. Median values are displayed as solid horizontal lines and outliers are plotted as distinct points. No data available for September.

siderophore BGCs also showed low and relatively uniform median abundances (< 0.2 RPKM) in

the winter in both the euphotic and mesopelagic zones, increasing in the euphotic zone in the

spring (Fig. 5B). Monthly median ORF abundances in the euphotic zone peaked during the

summer (July, 0.30 RPKM), decreasing throughout the fall in the upper euphotic zone, but

remaining elevated in the lower euphotic zone. Lowest monthly median ORF abundances in the

euphotic zone were observed in January and February (0.04 RPKM). Mesopelagic ORF

abundances were less seasonally variable (Fig. 5B), with median ORF abundances ranging 0.05-

0.11 RPKM. Monthly median ORF abundances were maximal in April (0.15 RPKM) and ORFs

from siderophore BGCs were not detected in October in the mesopelagic.

Table 1. Siderophores measured in the water column and siderophore biosynthetic gene clusters
(BGCs) identified at Station ALOHA. Siderophores identified in the water column are grouped
according to type and are shown in the table based on that grouping. For example, multiple
different forms of amphibactins were detected (Table S2) but are all grouped here under
"amphibactin." Specific siderophore IDs can be found in Supplementary Table 2. The '-' denotes
that the specific siderophore was not found either in the water column or in the Station ALOHA
gene catalog.

Siderophore	Biosynthetic Pathway	Depth(s)	Season	BGC in gene catalog
aerobactin	NIS	-	-	yes
amphibactins	NRPS	300-400	summer	-
anguibactin	NRPS	-	-	yes
enterobactin	NIS	-	-	yes
ferrioxamaine	NIS	15-400	spring, summer, winter	yes
myxochelin	NRPS	-	-	yes
petrobactin	NIS	15	winter	yes
piscibactin	NRPS	15	winter	yes
(pseudo)alterobactin	NRPS	15	winter	yes
synechobactin	NRPS	15, 150	spring, summer, winter	-
thalassosamide	NIS	-	-	yes
vanchrobactin	NRPS	-	-	yes
vibrioferrin	NIS	15	winter	yes
vibriobactin	NRPS	-	-	yes

Several different siderophores were detected both as dissolved siderophores in the water column

and as a BGC in the Station ALOHA gene catalog (Table 1). Of those identified in the water

column, primarily hydroxamate siderophores were detected, with ferrioxamines and

amphibactins being the most common (Table 1). The detected siderophore BGCs largely

belonged to catecholate and carboxylate-type siderophores (Table 1). Two ferrioxamine BGCs

were identified in the ALOHA gene catalog, however, these BGCs were not detected in the

2015-2016 metagenomic dataset. Several siderophores were found both in the water column and

in the metagenomes, including petrobactin, pseudoalterobactin, piscibactin and vibrioferrin

(Table 1). The biosynthesis genes were primarily homologues of known genes from *Vibrio*,

Photobacterium, *Alteromonas*, and *Pseudoalteromonas* (Table S3).

4. Discussion

4.1 Turnover times of inorganic iron and iron-siderophore complexes reflect bioavailability to

the microbial community

Siderophore mediated acquisition of dFe involves secretion, uptake, and recycling of these

compounds, and the turnover time of each Fe source reflects its bioavailability. Siderophore-

bound Fe was taken up less rapidly than the inorganic Fe additions, which were likely complexed

by unbound ambient ligands. Siderophore Fe uptake had average turnover times ranging from

15-250 days (Fig. 1C) and an average turnover time of 148 days from all depths and siderophore

treatments, while FeL turned over more rapidly averaging 14 days (Fig. 1C). The different

turnover times reflects differences in the bioavailability of inorganic versus organically-bound Fe

(Lis et al. 2015). The average turnover times of FeL observed in this study were very similar to

turnover times of 14-18 days in the mixed layer during June 2019, measured at or near

background dFe concentrations at Station ALOHA (Hawco et al. 2022). The shorter turnover of

FeL likely reflects weaker binding of inorganic Fe associated with ligands naturally present in

seawater, which are more accessible to phytoplankton using ferric reductases as an uptake

mechanism (Maldonado and Price 2001; Salmon et al. 2006; Lis et al. 2015; Coale et al. 2019).

Moreover, we found rates of FeL uptake were higher in the light than in the dark, suggesting

significant uptake by photoautotrophs. Uptake rates of siderophore-bound Fe, in contrast, were

always higher in the dark than the light, suggesting that siderophore Fe consumption was likely

controlled by heterotrophic plankton and that photochemical impacts on the uptake of

siderophore-bound Fe were minimal (Barbeau 2006). The turnover time of Fe bound to

siderophores was also much longer, on the order of weeks to months, suggesting that Fe bound to

siderophores is longer-lived than the inorganic and/or weakly bound Fe pool, and may help to

retain dFe in both euphotic and upper mesopelagic waters.

Turnover times of siderophore-bound Fe varied depending on the siderophore. Fe associated

with amphibactins was taken up more quickly than ferrioxamine E or B (Fig. 1B). Amphibactins

have a lower binding affinity to dFe (log of the conditional stability constants = 12.0-12.5;

Bundy et al. 2018) compared to ferrioxamines (log of the conditional stability constants = 14.0-

14.4; (Bundy et al. 2018)), and therefore Fe could be more easily reducible from amphibactins.

Alternatively, more members of the microbial community may possess direct Fe-amphibactin

uptake pathways, but these uptake pathways are not well constrained. The relative turnover times

and bioavailability of the different siderophores therefore, appeared to be governed by their

binding strengths in this case. In our experiments, the highest uptake rates of siderophore-bound

Fe were also found in the 0.2-2.0 μm size fraction and in the dark, suggesting Fe bound to

siderophores may be more accessible to heterotrophic bacteria rather than larger

photoautotrophs.

4.2 Rates of iron acquisition by the microbial community are elevated in the mesopelagic waters

of Station ALOHA relative to the surface

An unexpected result of this work was the faster uptake rates and shorter turnover times of Fe

observed at 300 m compared to 15 m, especially in the case of siderophore-bound Fe, resulting in

higher uptake rates per cell and higher Fe:C ratios at depth. There are several experimental

details that should be considered when interpreting these observations. While these factors were

controlled for whenever feasible, the possibility remains that they have influenced the results

presented here. For example, macronutrients are elevated at 300 m relative to 15 m, so it is

possible Fe uptake rates were enhanced due to relief from macronutrient limitation. However,

our surface water treatments included modest NO_3^- amendments that should have alleviated

nitrogen limitation. The experiments at 300 m were also incubated at the same temperature as the

15 m experiments, likely enhancing the uptake rate of Fe relative to *in situ* rates due to increased

metabolic activity at higher temperatures. However, it is not clear why this would stimulate

uptake rates above those measured in surface waters. An additional possibility is that the ^{55}Fe

adsorbed to the outside of cells or detritus more readily at 300 m, causing apparent high Fe

uptake compared to 15 m. If this was the case, we would have expected the dead controls from

300 m to have significantly higher activities than those from 15 m, and this was not observed.

Furthermore, oxalate rinsing was applied to all samples, which should remove most (if not all)

adsorbed Fe. Finally, in terms of the observed Fe:C ratios, an additional source of uncertainty

may come from the conversion of Fe uptake to Fe:C using a fixed carbon quota. Fe:C ratios

based on a fixed carbon quota overlap with published values for surface waters, but resulted in

values for 300 m that were significantly higher (Table S1). If the actual carbon quota per cell is

significantly higher in heterotrophic bacteria at 300 m, then the calculated Fe:C ratios presented

here could be artificially elevated. Carbon biomass estimates are not available for any cultured

marine bacteria from mesopelagic waters, so we are not able to determine if this fixed carbon

quota is too low.

Accounting for these possible experimental artifacts, it is likely that the fast uptake rates and

short turnover times of Fe observed at 300 m are related to ecologically relevant differences in

Fe acquisition by the microbial community at this depth. The Fe requirements of heterotrophic

bacteria are not well known, but evidence suggests they are elevated relative to large

phytoplankton (Fe:C ~ 2.0 μmol: mol) or cyanobacteria (Fe:C ~ 19 μmol: mol) (Strzepek et al.

2005). Published estimates for heterotrophic bacteria Fe:C ratios range from 7.5 (Tortell et al.

1996) to 83 μmol: mol (Mazzotta et al. 2020), and the observations at 300 m in this study expand

that range to upwards of 700 μmol: mol. The range of Fe requirements observed across these

studies likely varies according to experimental growth conditions and the lifestyle strategies of

specific groups of bacteria under consideration. In the current study, the 1 nmol L^{-1} dFe additions

and slightly higher in situ dFe concentrations at 300 m may have supported luxury Fe uptake and

storage by the *in situ* bacteria community. This could lead to higher Fe uptake at these depths,

and much higher Fe:C quotas than for surface bacteria communities. Previous work has

identified multiple copies of the Fe-storage protein bacterioferritin in a cultured marine

heterotroph (Mazzotta et al. 2020) and modeling work suggests that bacteria competitively

consume Fe and that Fe storage plays a large role (Ratnarajah et al. 2021). Alternatively, it is

possible that the consumption of refractory dissolved organic matter at these depths has a higher

Fe requirement for marine bacteria, though this has not been explored to our knowledge. The

energy starved microbial community in the upper mesopelagic could have elevated metabolic

requirements for Fe (Tortell et al. 1996) relative to the surface ocean community, potentially

resulting from differences in the lability or oxidation state of organic matter. Additionally,

nitrifying bacteria and archaea, whose cellular abundances significantly increase in mesopelagic

waters, appear to have high Fe requirements (Shafiee et al. 2019; Saito et al. 2020). In addition to

the high uptake rates and Fe:C quotas observed here, other lines of evidence are starting to point

to an increased demand for Fe in the mesopelagic. For example, high concentrations of

siderophores have been observed in the upper mesopelagic ocean, suggesting active Fe uptake at

these depths (Bundy et al. 2018; Park et al. 2023; Li et al. 2024). Ultimately, understanding the

Fe requirements of mesopelagic bacteria could have important implications for carbon and Fe

cycling in the twilight zone.

4.3 Siderophore production at Station ALOHA reflects seasonal variation in primary production

and organic matter availability

Measurements of dissolved siderophores highlight seasonal differences at Station ALOHA, with

siderophore concentrations greatest in the spring, summer and fall but almost entirely absent or

low in winter (Fig. 4). There was high variability in total dissolved siderophore concentrations

and the identity of siderophores observed over the study period. Some of this variability may

have been caused by factors related to method developments over time. For example,

siderophore concentrations were relatively high during our only fall sampling period (2013), but

we were not able to identify specific siderophores with high confidence in these samples (Table

S2). These samples were collected during the development of our LC-ICP-MS and LC-ESI-MS

methods, and since that sample collection, continued improvements in detection limits and

ionization parameters have enhanced our ability to detect siderophores with high confidence.

Collecting profile samples during this time was not possible, because of the high volumes of

water that were needed (800 L). However, improvements to methodological sensitivity prior to

sample collection in 2014 allowed for the high confidence detection of several siderophores from

the winter cruise in this year despite the relatively low concentrations that were observed (Table

S2). The dissolved siderophore distributions were notably higher in the spring, summer, and fall,

even when only considering surface waters (Fig. 4). The elevated concentrations of dissolved

siderophores in spring and summer were matched by a higher potential for the microbial

community to biosynthesize siderophores during these seasons, particularly in the euphotic zone

(Fig. 5). Of the 14 different types of siderophores identified using mass spectrometry or from

gene pathways in the metagenomes, 5 were found in both (Table 1). The remaining compounds

were either only found in the water column and not in the metagenomes (2 of the 14), or

conversely, the biosynthesis genes were found (7 of the 14) but the compound was not identified

in the water column samples. A wide structural diversity of siderophores has been detected using

the LC-MS techniques employed here (Boiteau et al. 2019), making it unlikely that specific

siderophores escaped detection in the water column due to methodological bias. Therefore, the

limited overlap in the dominant siderophores measured and the BGCs identified suggests that

either some compounds may be underestimated from metagenomic approaches, or that some

siderophores cycle too rapidly to be observed in measurable quantities. Further uptake studies

examining the bioavailability of the siderophore groups dominant within the metagenomes, such

as catecholate-type siderophores, will help to further explain these differences.

Seasonality in nutrient supply, productivity, and export have all been documented in the North

Pacific subtropical gyre and Station ALOHA (Fitzsimmons et al. 2015; Hawco et al. 2021; Karl

et al. 2021) and all impact microbial community structure and thus siderophore production. In

the spring, primary production begins to increase at Station ALOHA (Fig. 2 and 3). This is

thought to be a result of increasing solar irradiance (Karl et al. 2021) and these changes are

especially evident in the lower euphotic zone ((Letelier et al. 2004, 2017); Fig. S1 and S2). This

springtime increase in productivity generally coincides with the peak in Fe delivery to this region

(Fitzsimmons et al. 2015). During this time, Fe is delivered primarily via atmospheric dust

deposition which introduces lithogenic Fe particles to the surface ocean. In this study, the highest

concentrations of dissolved siderophores were observed in surface waters during the spring and

coincided with increasing biosynthetic capacity in the euphotic zone (Fig. 4 and 5). It is likely

that siderophore secretion by heterotrophic bacteria at Station ALOHA during this time may help

solubilize particulate Fe (Bundy et al. 2018) and to compete with other microorganisms for Fe

(Boiteau et al. 2016) as the Fe demand of the microbial community begins to increase during

these productive months. For example, ferrioxamines have been shown to be produced during

the degradation of sinking (Velasquez et al. 2016) or suspended particles (Bundy et al. 2018),

and high siderophore concentrations were observed in the North Pacific Transition Zone in a

region of high dust inputs (Park et al. 2023).

As primary production increases throughout the summer months at Station ALOHA, essential

nutrients such as NO_3^- and dFe are consumed. During late summer, when the physical delivery of

NO_3^- to the surface ocean is low, the nitrogen fueling these higher rates of production is thought

to be supplied by N₂ fixation (Karl et al. 2021). Rates of N_2 fixation peak in the late summer and

early fall at Station ALOHA when warm, stratified waters select for the growth of N_2-fixing

cyanobacteria (Böttjer et al. 2017). This late summer production is also associated with a large

pulse of particulate carbon export that reaches the sea floor and is thought to be the result of

diatom growth supported by N_2-fixing endosymbionts (Karl et al. 2012). As a result of increasing

primary production and temperature, rates of heterotrophic bacterial production are also elevated

by late summer at Station ALOHA (Viviani and Church 2017). However, during this period, dust

delivery to the surface ocean decreases (Fitzsimmons et al. 2015). Thus, an increased Fe demand

of the microbial community resulting from high rates of primary production, N_2 fixation, and

bacterial production coincides with low rates of Fe delivery and likely intensifies the competition

for Fe throughout the euphotic zone as summer progresses. This has the potential to set up

conditions favoring the production of siderophores by bacteria as a means to competitively

acquire Fe or to retain or recycle Fe in the euphotic zone over longer timescales (Boyd et al.

2015; Hayes et al. 2015; Hawco et al. 2022).

In addition to dissolved siderophore concentrations, the genetic potential for siderophore

biosynthesis also peaked during the summer months. Furthermore, a majority of the genera

found to have siderophore BGCs in this study, such as *Vibrio* and *Alteromonas*, contain known

copiotrophic bacteria, suggesting that siderophore production may be common to heterotrophs

responding to episodic inputs of organic matter during periods of elevated production (Fontanez

et al. 2015; Pelve et al. 2017; Church et al. 2021; Poff et al. 2021; Leu et al. 2022). These genera

have been found to be dominant in sediment trap material from Station ALOHA and in

association with sinking eukaryotes (Fontanez et al. 2015). Thus in many cases, siderophore

production might largely be due to the Fe demand of heterotrophic bacteria as they consume

carbon from sinking particles (Hopkinson and Barbeau 2012). In this study, seasonal patterns in

the genetic potential for siderophore production within the euphotic zone closely followed those

of the abundance of both suspended and sinking particles, further suggesting that siderophore

biosynthesis may be directly tied to the dynamics of particle production. While the genetic

potential for siderophore production did not vary seasonally in the mesopelagic zone, elevated

dissolved siderophore concentrations were observed in the summer in this region of the water

column, potentially in response to the elevated particle flux escaping the euphotic zone during

this time. Finally, in winter, siderophore concentrations and the abundance of siderophore BGCs

were reduced significantly throughout the water column, likely reflecting the decrease in

metabolic activity or Fe demand of heterotrophic bacteria as productivity and the availability of

organic substrates diminish, or increased mixing that dilutes siderophores.

Together, the higher concentrations of siderophores from spring to fall and the dominance of

copiotrophic bacteria identified as potential siderophore producers in the metagenomes, predict

that siderophore production is heightened during periods of enhanced productivity at Station

ALOHA and provides compelling evidence that ligand production by the microbial community

plays an important role in the seasonal cycling and retention of Fe at Station ALOHA. Uptake

rates of Fe in the euphotic zone agree with previous measurements at Station ALOHA and imply

that Fe is cycled rapidly by the microbial community in this region. While less bioavailable than

inorganic Fe additions, siderophore-bound Fe proved to be a bioavailable source of Fe to the

microbial community in both the euphotic and mesopelagic zones. Importantly, we found the

resulting bioavailability depended on the specific siderophore structure, warranting additional

studies focused on elucidating specific compounds in the Fe-binding ligand pool and their uptake kinetics in natural communities. Due to longer turnover times of Fe bound to siderophores, the production of strong siderophores has the potential to transform dFe into a form that persists in the water column on longer time scales. The finding that Fe uptake rates and turnover times were faster in the mesopelagic waters than in the surface ocean was surprising and merits further investigation. Such results suggest Fe availability is impacting microbial dynamics and Fe cycling at depth.

5. Acknowledgements

Special thanks to the Captain and crew of the R/V *Kilo Moana* and R/V *Ka'imikai-O-Kanaloa* for help with sample collection, to the Hawaii Ocean Time-series program for the numerous biogeochemical measurements and to Kendra A. Turk-Kubo for additional flow cytometry analyses. We would also like to thank Travis Mellett for helpful comments on the manuscript. RMB was funded by a Life Sciences-Simons Early Career Investigator in Marine Microbial Ecology and Evolution Awards (Award #618401) and a Woods Hole Oceanographic Institution postdoctoral fellowship. LEM was funded by a Simons Postdoctoral Fellowship in Marine Microbial Ecology (Award #729162). MJC was funded by the Simons Collaboration on Ocean Processes (Award #721221). NJH was supported by Life Sciences-Simons Early Career Investigator in Marine Microbial Ecology and Evolution Awards (Award #924096). The authors have no conflict of interest to declare.

References

Baars, O., F. M. M. Morel, and D. H. Perlman. 2014. ChelomEx: Isotope-assisted discovery of

metal chelates in complex media using high-resolution LC-MS. Anal. Chem. **86**: 11298–

11305. doi:10.1021/ac503000e

Barbeau, K. 2006. Photochemistry of organic iron(III) complexing ligands in oceanic systems.

Photochem. Photobiol. **82**: 1505–1516. doi:10.1562/2006-06-16-ir-935

Boiteau, R. M., J. N. Fitzsimmons, D. J. Repeta, and E. A. Boyle. 2013. Detection of Iron

Ligands in Seawater and Marine Cyanobacteria Cultures by High-Performance Liquid

Chromatography-Inductively Coupled Plasma-Mass Spectrometry. Anal. Chem. **85**: 4357–

4362. doi:10.1021/ac3034568

Boiteau, R. M., D. R. Mende, N. J. Hawco, and others. 2016. Siderophore-based microbial

adaptations to iron scarcity across the eastern Pacific Ocean. Proc. Natl. Acad. Sci. **113**:

14237–14242.

Boiteau, R. M., C. P. Till, T. H. Coale, J. N. Fitzsimmons, K. W. Bruland, and D. J. Repeta.

2019. Patterns of iron and siderophore distributions across the California Current System.

Limnol. Oceanogr. **64**: 376–389. doi:10.1002/lno.11046

Bolger, A. M., M. Lohse, and B. Usadel. 2014. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics **30**: 2114–2120.

Böttjer, D., J. E. Dore, D. M. Karl, R. M. Letelier, C. Mahaffey, S. T. Wilson, J. Zehr, and M. J.

Church. 2017. Temporal variability of nitrogen fixation and particulate nitrogen export at

Station ALOHA. Limnol. Oceanogr. **62**: 200–216. doi:https://doi.org/10.1002/lno.10386

Boyd, P. W., R. F. Strzepek, M. J. Ellwood, D. A. Hutchins, S. D. Nodder, B. S. Twining, and S.

W. Wilhelm. 2015. Why are biotic iron pools uniform across high- and low-iron pelagic

790  ecosystems? Global Biogeochem. Cycles **29**: 1028–1043. doi:10.1002/2014gb005014

791 Bundy, R. M., R. M. Boiteau, C. McLean, K. A. Turk-Kubo, M. R. McIlvin, M. A. Saito, B. A.

792  S. Van Mooy, and D. J. Repeta. 2018. Distinct Siderophores Contribute to Iron Cycling in

793  the Mesopelagic at Station ALOHA. Front. Mar. Sci. **5**: 1–15.

794  doi:10.3389/fmars.2018.00061

795 Church, M. J., E. Kyi, R. O. Hall Jr, D. M. Karl, M. Lindh, A. Nelson, and E. K. Wear. 2021.

796  Production and diversity of microorganisms associated with sinking particles in the

797  subtropical North Pacific Ocean. Limnol. Oceanogr. **66**: 3255–3270.

798 Coale, T. H., M. Moosburner, A. Horák, M. Oborník, K. A. Barbeau, and A. E. Allen. 2019.

799  Reduction-dependent siderophore assimilation in a model pennate diatom. Proc. Natl. Acad.

800  Sci. U. S. A. **116**: 23609–23617. doi:10.1073/pnas.1907234116

801 Cordero, O. X., L.-A. Ventouras, E. F. DeLong, and M. F. Polz. 2012. Public good dynamics

802  drive evolution of iron acquisition strategies in natural bacterioplankton populations. Proc.

803  Natl. Acad. Sci. **109**: 20059–20064. doi:10.1073/pnas.1213344109

804 DiTullio, G. R., and E. A. Laws. 1991. Impact of an atmospheric-oceanic disturbance on

805  phytoplankton community dynamics in the North Pacific Central Gyre. Deep Sea Res. Part

806  A. Oceanogr. Res. Pap. **38**: 1305–1329.

807 Dore, J. E., and D. M. Karl. 1996. Nitrite distributions and dynamics at Station ALOHA. Deep

808  Sea Res. Part II Top. Stud. Oceanogr. **43**: 385–402.

809 Dutkiewicz, S., A. E. Hickman, O. Jahn, W. W. Gregg, C. B. Mouw, and M. J. Follows. 2015.

810  Capturing optically important constituents and properties in a marine biogeochemical and

811  ecosystem model. Biogeosciences **12**: 4447–4481. doi:10.5194/bg-12-4447-2015

812 Eddy, S. R. 2011. Accelerated profile HMM searches. PLoS Comput. Biol. **7**: e1002195.

Fitzsimmons, J. N., and E. A. Boyle. 2014. Assessment and comparison of Anopore and cross

flow filtration methods for the determination of dissolved iron size fractionation into soluble

and colloidal phases in seawater. Limnol. Oceanogr. **12**: 246–263.

doi:10.4319/lom.2014.12.246

Fitzsimmons, J. N., C. T. Hayes, S. N. Al-Subiai, R. Zhang, P. L. Morton, R. E. Weisend, F.

Ascani, and E. A. Boyle. 2015. Daily to decadal variability of size-fractionated iron and

iron-binding ligands at the Hawaii Ocean Time-series Station ALOHA. Geochim.

Cosmochim. Acta **171**: 303–324.

Fontanez, K. M., J. M. Eppley, T. J. Samo, D. M. Karl, and E. F. DeLong. 2015. Microbial

community structure and function on sinking particles in the North Pacific Subtropical

Gyre. Front. Microbiol. **6**: 469.

Foreman, R. K., M. Segura-Noguera, and D. M. Karl. 2016. Validation of Ti (III) as a reducing

agent in the chemiluminescent determination of nitrate and nitrite in seawater. Mar. Chem.

186: 83–89.

Gledhill, M., and K. N. Buck. 2012. The organic complexation of iron in the marine

environment: A review. Front. Microbiol. **3**: 69. doi:10.3389/fmicb.2012.00069

Hawco, N. J., B. Barone, M. J. Church, and others. 2021. Iron depletion in the deep chlorophyll

maximum: mesoscale eddies as natural iron fertilization experiments. Global Biogeochem.

Cycles **35**: e2021GB007112.

Hawco, N. J., S.-C. Yang, P. Pinedo-González, E. E. Black, J. Kenyon, S. Ferrón, X. Bian, and

S. G. John. 2022. Recycling of dissolved iron in the North Pacific Subtropical Gyre.

Limnol. Oceanogr. doi:https://doi.org/10.1002/lno.12212

Hayes, C. T., J. N. Fitzsimmons, E. A. Boyle, D. McGee, R. F. Anderson, R. Weisend, and P. L.

Morton. 2015. Thorium isotopes tracing the iron cycle at the Hawaii Ocean Time-series

Station ALOHA. Geochim. Cosmochim. Acta **169**: 1–16.

Hider, R. C., and X. Kong. 2010. Chemistry and biology of siderophores. Nat. Prod. Rep. **27**:

637–657. doi:10.1039/b906679a

Hogle, S. L., T. Hackl, R. M. Bundy, and others. 2022. Siderophores as an iron source for

picocyanobacteria in deep chlorophyll maximum layers of the oligotrophic ocean. ISME J.

1–11.

Hopkinson, B. M., and K. A. Barbeau. 2012. Iron transporters in marine prokaryotic genomes

and metagenomes. Environ. Microbiol. **14**: 114–128. doi:10.1111/j.1462-

2920.2011.02539.x

Hopkinson, B., J. Nunnery, and K. Barbeau. 2005. Bioavailability and chemistry of iron(III)-

porphyrin complexes. Abstr. Pap. Am. Chem. Soc. **229**: 075-GEOC.

Jones, D. R., D. M. Karl, and E. A. Laws. 1996. Growth rates and production of heterotrophic

bacteria and phytoplankton in the North Pacific subtropical gyre. Deep Sea Res. Part I

Oceanogr. Res. Pap. **43**: 1567–1580.

Karl, D., R. Letelier, L. Tupas, and J. Dore. 1997. The role of nitrogen fixation in

biogeochemical cycling in the subtropical North Pacific Ocean. Nature **388**: 533.

Karl, D. M., M. J. Church, J. E. Dore, R. M. Letelier, and C. Mahaffey. 2012. Predictable and

efficient carbon sequestration in the North Pacific Ocean supported by symbiotic nitrogen

fixation. Proc. Natl. Acad. Sci. **109**: 1842–1849.

Karl, D. M., R. M. Letelier, R. R. Bidigare, K. M. Björkman, M. J. Church, J. E. Dore, and A. E.

White. 2021. Seasonal-to-decadal scale variability in primary production and particulate

matter export at Station ALOHA. Prog. Oceanogr. **195**: 102563.

859 Karl, D. M., and R. Lukas. 1996. The Hawaii Ocean Time-series (HOT) program: Background,

860   rationale and field implementation. Deep Sea Res. Part II Top. Stud. Oceanogr. **43**: 129–

861   156.

862 Kirchman, D. L. 2016. Growth rates of microbes in the oceans. Ann. Rev. Mar. Sci. **8**: 285–309.

863 Kramer, J., Ö. Özkaya, and R. Kümmerli. 2020. Bacterial siderophores in community and host

864   interactions. Nat. Rev. Microbiol. **18**: 152–163.

865 Letelier, R. M., K. M. Björkman, M. J. Church, and others. 2019. Climate-driven oscillation of

866   phosphorus and iron limitation in the North Pacific Subtropical Gyre. Proc. Natl. Acad. Sci.

867   U. S. A. **116**: 12720–12728. doi:10.1073/pnas.1900789116

868 Letelier, R. M., D. M. Karl, M. R. Abbott, and R. R. Bidigare. 2004. Light driven seasonal

869   patterns of chlorophyll and nitrate in the lower euphotic zone of the North Pacific

870   Subtropical Gyre. Limnol. Oceanogr. **49**: 508–519.

871 Letelier, R. M., A. E. White, R. R. Bidigare, B. Barone, M. J. Church, and D. M. Karl. 2017.

872   Light absorption by phytoplankton in the North Pacific Subtropical Gyre. Limnol.

873   Oceanogr. **62**: 1526–1540.

874 Leu, A. O., J. M. Eppley, A. Burger, and E. F. DeLong. 2022. Diverse Genomic Traits

875   Differentiate Sinking-Particle-Associated versus Free-Living Microbes throughout the

876   Oligotrophic Open Ocean Water Column. MBio **13**: e0156922. doi:10.1128/mbio.01569-22

877 Li, H., and R. Durbin. 2009. Fast and accurate short read alignment with Burrows--Wheeler

878   transform. bioinformatics **25**: 1754–1760.

879 Li, J., L. Babcock-Adams, R. Boiteau, and others. 2024. Microbial iron limitation in the ocean's

880   twilight zone.

881 Liao, Y., G. K. Smyth, and W. Shi. 2014. featureCounts: an efficient general purpose program

for assigning sequence reads to genomic features. Bioinformatics **30**: 923–930.

Lis, H., Y. Shaked, C. Kranzler, N. Keren, and F. M. M. Morel. 2015. Iron bioavailability to

phytoplankton: an empirical approach. ISME J. **9**: 1003–1013. doi:10.1038/ismej.2014.199

Longhurst, A. R. 2010. Ecological geography of the sea, Elsevier.

Luo, E., J. M. Eppley, A. E. Romano, D. R. Mende, and E. F. DeLong. 2020. Double-stranded

DNA virioplankton dynamics and reproductive strategies in the oligotrophic open ocean

water column. ISME J. **14**: 1304–1315. doi:10.1038/s41396-020-0604-8

Maldonado, M. T., and N. M. Price. 2001. Reduction and transport of organically bound iron by

Thalassiosira oceanica (Bacillariophyceae). J. Phycol. **37**: 298–309. doi:10.1046/j.1529-

8817.2001.037002298.x

Manck, L. E., J. Park, B. J. Tully, A. M. Poire, R. M. Bundy, C. L. Dupont, and K. A. Barbeau.

2022. Petrobactin, a siderophore produced by Alteromonas, mediates community iron

acquisition in the global ocean. ISME J. **16**: 358–369.

Mawji, E., M. Gledhill, J. A. Milton, and others. 2008. Hydroxamate Siderophores: Occurance

and Importance in the Atlantic Ocean. Environ. Sci. Technol. **42**: 8675–8680.

Mazzotta, M. G., M. R. McIlvin, and M. A. Saito. 2020. Characterization of the Fe

metalloproteome of a ubiquitous marine heterotroph, Pseudoalteromonas (BB2-AT2):

multiple bacterioferritin copies enable significant Fe storage†. Metallomics **12**: 654–667.

doi:10.1039/d0mt00034e

Medema, M. H., K. Blin, P. Cimermancic, and others. 2011. antiSMASH: rapid identification,

annotation and analysis of secondary metabolite biosynthesis gene clusters in bacterial and

fungal genome sequences. Nucleic Acids Res. **39**: W339--W346.

Mende, D. R., J. A. Bryant, F. O. Aylward, J. M. Eppley, T. Nielsen, D. M. Karl, and E. F.

DeLong. 2017. Environmental drivers of a microbial genomic transition zone in the ocean's

interior. Nat. Microbiol. **2**: 1367–1373. doi:10.1038/s41564-017-0008-3

Moore, L. E., M. I. Heller, K. A. Barbeau, J. W. Moffett, and R. M. Bundy. 2021. Organic

complexation of iron by strong ligands and siderophores in the eastern tropical North

Pacific oxygen deficient zone. Mar. Chem. **236**: 104021.

Park, J., B. P. Durham, R. S. Key, and others. 2023. Siderophore production and utilization by

marine bacteria in the North Pacific Ocean. Limnol. Oceanogr. **68**: 1636–1653.

Pelve, E. A., K. M. Fontanez, and E. F. DeLong. 2017. Bacterial succession on sinking particles

in the ocean's interior. Front. Microbiol. **8**: 2269.

Pinedo-González, P., N. J. Hawco, R. M. Bundy, and others. 2020. Anthropogenic Asian

aerosols provide Fe to the North Pacific Ocean. Proc. Natl. Acad. Sci. U. S. A. **117**: 27862–

27868. doi:10.1073/pnas.2010315117

Poff, K. E., A. O. Leu, J. M. Eppley, D. M. Karl, and E. F. DeLong. 2021. Microbial dynamics

of elevated carbon flux in the open ocean's abyss. Proc. Natl. Acad. Sci. **118**: e2018269118.

Ratnarajah, L., S. Blain, P. W. Boyd, M. Fourquez, I. Obernosterer, and A. Tagliabue. 2021.

Resource Colimitation Drives Competition Between Phytoplankton and Bacteria in the

Southern Ocean. Geophys. Res. Lett. **48**: 1–11. doi:10.1029/2020GL088369

Saito, M. A., M. R. McIlvin, D. M. Moran, and others. 2020. Abundant nitrite-oxidizing

metalloenzymes in the mesopelagic zone of the tropical Pacific Ocean. Nat. Geosci. **13**:

355–362.

Salmon, T. P., A. L. Rose, B. A. Neilan, and T. D. Waite. 2006. The FeL model of iron

acquisition: Nondissociative reduction of ferric complexes in the marine environment.

Limnol. Oceanogr. **51**: 1744–1754.

Sandy, M., and A. Butler. 2009. Microbial iron acquisition: marine and terrestrial siderophores.

Chem. Rev. **109**: 4580–4595.

Shafiee, R. T., J. T. Snow, Q. Zhang, and R. E. M. Rickaby. 2019. Iron requirements and uptake

strategies of the globally abundant marine ammonia-oxidising archaeon, Nitrosopumilus

maritimus SCM1. ISME J. **13**: 2295–2305.

Shaked, Y., and H. Lis. 2012. Disassembling iron availability to phytoplankton. Front.

Microbiol. **3**: 123. doi:10.3389/fmicb.2012.00123

Smith, D. C., and F. Azam. 1992. A simple, economical method for measuring bacterial protein

synthesis rates in seawater using 3H-leucine. Mar. Microb. food webs **6**: 107–114.

Strzepek, R. F., M. T. Maldonado, J. L. Higgins, J. Hall, K. Safi, S. W. Wilhelm, and P. W.

Boyd. 2005. Spinning the "ferrous wheel": The importance of the microbial community in

an iron budget during the FeCycle experiment. Global Biogeochem. Cycles **19**.

doi:10.1029/2005GB002490

Sutak, R., J.-M. Camadro, and E. Lesuisse. 2020. Iron Uptake Mechanisms in Marine

Phytoplankton. Front. Microbiol. **11**. doi:10.3389/fmicb.2020.566691

Tortell, P. D., M. T. Maldonado, J. Granger, and N. M. Priece. 1999. Marine bacteria and

geochemical cycling of iron in the oceans. FEMS Microbiol. Ecol. **29**: 1–11.

doi:10.1111/j.1574-6941.1999.tb00593.x

Tortell, P. D., M. T. Maldonado, and N. M. Price. 1996. The role of heterotrophic bacteria in

iron-limited ocean ecosystems. Lett. to Nat. **383**: 330–332.

Tovar-Sanchez, a. 2003. A trace metal clean reagent to remove surface-bound iron from marine

phytoplankton. Mar. Chem. **82**: 91–99. doi:10.1016/S0304-4203(03)00054-9

Velasquez, I. B., E. Ibisanmi, E. W. Maas, P. W. Boyd, S. Nodder, and S. G. Sander. 2016.

Ferrioxamine siderophores detected amongst iron binding ligands produced during the

remineralization of marine particles. Front. Mar. Sci. **3**: 172.

Viviani, D. A., and M. J. Church. 2017. Decoupling between bacterial production and primary

production over multiple time scales in the North Pacific Subtropical Gyre. Deep Sea Res.

Part I Oceanogr. Res. Pap. **121**: 132–142.

Vraspir, J. M., and A. Butler. 2009. Chemistry of Marine Ligands and Siderophores. Ann. Rev.

Mar. Sci. **1**: 43–63. doi:10.1146/annurev.marine.010908.163712

Data Availability Statement: Iron uptake data are available as supplementary files and on

Zenodo (DOI: 10.5281/zenodo.7062571). Sequence data are available from the NCBI short read

archive (SRA) under Bioproject no. PRJNA352737.

Supplementary Information for

Patterns of siderophore production and utilization at Station ALOHA from the surface to mesopelagic waters

Randelle M. Bundy[1,*] and Lauren E. Manck[2,*], Daniel J. Repeta[3], Matthew J. Church[2], Nicholas J. Hawco[4], Rene M. Boiteau[5], Jiwoon Park[1], Edward F. DeLong[4], Mak A. Saito[3]

[1]University of Washington, School of Oceanography, Seattle, WA

[2]University of Montana, Flathead Lake Biological Station, Polson, MT

[3]Woods Hole Oceanographic Institution, Marine Chemistry and Geochemistry, Woods Hole, MA

[4]University of Hawaii at Manoa, Center for Microbial Oceanography Research and Education, Honolulu, HI

[5]Oregon State University, College of Earth Ocean and Atmospheric Sciences, Corvallis, OR

[*]co-first authors and corresponding authors rbundy@uw.edu and lauren.manck@flbs.umt.edu

Running head (40 characters): Siderophore biosynthesis at Station ALOHA

Table S1. All ^{55}Fe uptake data from each experiment. The treatments are ^{55}Fe-amphibactins (Amp), ^{55}Fe-ferrioxamine B (DFB), ^{55}Fe-ferrioxamine E (DFE) and ^{55}FeCl$_3$ (FeL).The Fe:C$^+$ of heterotrophic bacteria was calculated by assuming all dark treatments were dominated by heterotrophic bacteria Fe uptake, so the total uptake (sum of 2.0 + 0.2 µm uptake) in the 250 mL bottle over the course of 12 hours was converted to total moles of Fe and then divided by the total bacteria abundance at the end of the experiment determined by flow cytometry. A bacteria carbon per cell of 12.4 fg C cell^{-1} (Boyd et al. 2015) was used to calculate bacteria carbon from each treatment. The additional column of Fe:C* were calculated based on the comparison of iron uptake rates (pmol Fe L^{-1} day^{-1}) to rates of bacterial carbon productivity (nmol C L^{-1} day^{-1}) at Station ALOHA at 15 m and 300 m depth.

Table S2. Dissolved siderophore concentrations (pM) and identifications from all six cruises in different seasons. Only siderophores that were found complexed to iron are presented. The '-' means that a high confidence (confirmed with MS^2) siderophore identification was not possible in that sample.

Table S3. The complete list of siderophore biosynthetic genes identified in the ALOHA gene catalog with functional and taxonomic identifications given where possible. Genes specifically identified in the 2015-2016 metagenomic dataset collected at Station ALOHA are noted as such.

Figure S1: Monthly mean nitrate+nitrite ($NO_3^-+NO_2^-$) concentrations and rates of primary production in the **(A)** upper euphotic zone ($\leq$ 75 m) and **(B)** lower euphotic zone (> 75 $\leq$ 150 m) at Station ALOHA during the 2013-2016 study period.

Figure S2: Monthly and seasonal mean depth-integrated concentrations of nitrate+nitrite (NO_3^- $+NO_2^-$) and rates of primary production in the **(A)** upper euphotic zone ($\leq$ 75 m) and **(B)** lower euphotic zone (> 75 $\leq$ 150 m) at Station ALOHA during the 2013-2016 study period.