Seasonal siderophore uptake and biosynthesis associated with carbon flux at Station ALOHA

Randelle M. Bundy[1,*] and Lauren E. Manck[2,*], Rene M. Boiteau[3], Jiwoon Park[1], Edward F. DeLong[4], Nicholas J. Hawco[4], Matthew J. Church[2], Mak A. Saito[5], Daniel J. Repeta[5]

[1]University of Washington, School of Oceanography, Seattle, WA
[2]University of Montana, Flathead Lake Biological Station, Polson, MT
[3]Oregon State University, College of Earth Ocean and Atmospheric Sciences, Corvallis, OR
[4]University of Hawaii at Manoa, Center for Microbial Oceanography Research and Education, Honolulu, HI
[5]Woods Hole Oceanographic Institution, Marine Chemistry and Geochemistry, Woods Hole, MA

[*]co-first authors and corresponding authors rbundy@uw.edu and lauren.manck@flbs.umt.edu

Running head (40 characters): Siderophore biosynthesis at Station ALOHA

Author contribution statement: RBundy and LEM wrote the manuscript and completed the analyses. RBoiteau and JP contributed data and edited the manuscript. EFD, MJC, MAS and DJR helped design the study and edit the manuscript.

Scientific Significance Statement (125 words; including a setup, challenge, and resolution-
currently 114 words**)**
Microbial community production in the subtropical oligotrophic North Pacific is limited by
macronutrients such as nitrogen and phosphorous. However, dissolved iron is another important
micronutrient that has seasonal inputs from dust and passing eddies, keeping the availability of
iron low and episodic. Little attention has been paid to the microbial strategies for dealing with
low iron to support primary production in the oligotrophic ocean, or how limited iron availability
impacts the processing of sinking particulate organic carbon in this region. In this study, we
explore siderophore production and uptake throughout the water column at Station ALOHA to
examine how the microbial community adapts and responds to changing iron and carbon
availability on seasonal timescales.

Abstract (**150 words**- currently 150 words)
Dissolved iron concentrations in the North Pacific subtropical oligotrophic gyre are low, in part
due to biological consumption and seasonally restricted iron delivery to these waters. In this
study, we examined seasonal changes in dissolved siderophore concentrations, uptake rates and
biosynthesis at Station ALOHA to observe changes in iron acquisition by the microbial
community. Genetic potential for siderophore biosynthesis was widespread throughout the upper
water column (0-500 m) and peaked in spring and summer along with siderophore
concentrations. Uptake rates were faster and residence times of siderophore-bound iron were
shorter at 300 m compared to 15 m, suggesting differences in iron acquisition in the
mesopelagic. Changes in siderophore production in the water column suggest that variations in
primary production and carbon export impact iron acquisition of the microbial community over
the seasonal cycle and provides initial evidence that iron demand may change in association with
carbon export at Station ALOHA.

Keywords: siderophore, dissolved iron, siderophore biosynthesis, Station ALOHA

Data Availability Statement: Iron uptake data are available as supplementary files and on
Zenodo (DOI: 10.5281/zenodo.7062571). Sequence data are available from the NCBI short read
archive (SRA) under Bioproject no. PRJNA352737.

32 **(Main body- 3,000 words-** currently 3,598 words including references; 3,345 without
33 references**)**
34
35 **1. Introduction**
36
Oligotrophic subtropical gyres comprise the world's largest ocean biome (Longhurst 2010).
Microbial communities inhabiting the oligotrophic ocean are exposed to low concentrations of
nitrogen and phosphorous (Karl et al. 1997; Fitzsimmons and Boyle 2014; Letelier et al. 2019),
and dissolved iron (Fitzsimmons et al. 2015). The Hawaii Ocean Time-series (HOT) at Station
ALOHA in the North Pacific subtropical gyre is one of the longest running open ocean time-
series programs, and observations from the HOT program study site have demonstrated how
seasonal changes in nutrient supply impact the microbial community and carbon export (Karl
and Lukas 1996). The resulting time series captures both the relatively stable conditions of the
oligotrophic ocean and the importance of episodic and seasonal-scale delivery of nutrients to the
upper ocean from passing eddies, seasonal dust inputs, and the shoaling and deepening of the
pycnocline (Fitzsimmons et al. 2015; Hayes et al. 2015; Pinedo-González et al. 2020).
48
Seasonal changes in the external supply of iron to Station ALOHA impacts microbial community
composition and can trigger organic ligand production on the timescale of days (Fitzsimmons et
al. 2015; Hayes et al. 2015). Some of the strongest organic iron-binding ligands called
siderophores are produced by marine bacteria and fungi and can alter the bioavailability and
recycling of dissolved iron (Sandy and Butler 2009; Vraspir and Butler 2009). Siderophores are
synthesized by either a non-ribosomal peptide synthetase (NRPS) pathway or an NRPS-
independent siderophore (NIS) synthetase pathway and are secreted into the marine environment
to facilitate the uptake of iron (Sandy and Butler 2009; Hider and Kong 2010). Siderophore
biosynthesis is regulated by iron concentrations, where elevated concentrations of iron suppress
synthesis (Sandy and Butler 2009). Some microbes possess the genetic potential for both
synthesizing and internalizing siderophores, while others "cheat" and only possess outer
membrane receptors for exogenous siderophore uptake (Cordero et al. 2012; Kramer et al. 2020).
Internalizing siderophore-bound iron requires either specific transport pathways or the ability to
reduce Fe(III) to Fe(II) from the siderophore complex; thus iron complexed to siderophores has
low bioavailability to the microbial community as whole (Shaked and Lis 2012). Although
dissolved siderophores represent from 2-10% of the total dissolved iron-binding ligand pool at
Station ALOHA, they likely play an important role in competition and recycling of iron, and
facilitate the solubilization of iron from particles (Bundy et al. 2018).
67
The potential importance of siderophore production as a microbial iron acquisition strategy has
been recognized for some time, but measurements of the distributions of siderophores in
seawater are scarce (Mawji et al. 2008a; Boiteau et al. 2016, 2019; Bundy et al. 2018; Moore et
al. 2021; Manck et al. 2022) and little is known about the factors that influence their seasonal
variability. In this study, we explore seasonal changes in water column siderophore distributions,
biosynthesis potential and residence times at Station ALOHA in the context of the HOT program
to explore the conditions that impact microbial iron acquisition in this important oligotrophic
ecosystem.
76
2. Methods

2.1 Siderophore sample collection and analyses
Dissolved siderophore samples were collected from six cruises ($n = 2$ winter, $n = 2$ spring, $n = 1$
summer, $n = 1$ fall) from 2013-2016. Additional contextual data from three of the six cruises are
presented elsewhere (Table S1). Samples were collected via Teflon diaphragm pump or 8 L X-
Niskin bottles on a trace metal rosette using a non-metallic line. Samples were filtered (0.2 µm
Acropak 200 or Pall capsule filter) into either acid-cleaned fluorinated high density polyethene
200 L barrels (Boiteau et al. 2013) or acid-cleaned 20 L carboys (Bundy et al. 2018). Filtered
samples were pumped onto solid phase extraction columns (ENV resin, Agilent Technologies),
and stored at -20°C until analysis. Details on sample analysis can be found in Bundy et al.
(2018). Putative siderophores were detected using liquid chromatography inductively coupled
plasma mass spectrometry (LC-ICP-MS), where chromatographically resolved peaks were
quantified. Only iron-bound siderophores were quantified, and putative siderophores were
identified based on the alignment of an MS[1] peak with an ^{56}Fe peak in the LC-ICP-MS data and
corresponding MS[2] fragments that matched either published literature or *in silico* fragmentation
(Supplementary Methods). A siderophore assignment was not possible for all putative
siderophores, and in those cases only the total concentration is reported (Table S1). Average
dissolved siderophore concentrations were binned by depth and season to match the metagenome
sampling resolution.
96
2.2 Iron uptake and siderophore bioavailability experiments
Iron uptake experiments were performed on two expeditions (HOT278 in November 2015 and
KM1605 in March 2016). During the HOT278 expedition, experiments were performed in
surface waters (15 m; $n = 3$) and during KM1605 experiments were performed at 15 m and 300
m ($n = 2$). Four treatments were used in all experiments: $FeCl_3$ (denoted as "FeL"), ferrioxamine
B, ferrioxamine E, and Fe-amphibactin. Experiments at 15 m were performed both in the light
and dark, and experiments at 300 m were only performed in the dark. Iron uptake was measured
in the 2.0 µm and 0.2 µm size fraction after 12 hours and multiplied by 2 to calculate an iron
uptake rate per day. 500 nmol L^{-1} of nitrate was added to each of the 15 m experiments to control
for nitrogen limitation preventing the uptake of iron.
107
The isolation of the amphibactins used in this experiment have been described in detail
elsewhere (Bundy et al. 2018), and other ligand standards were commercially available.
Radioactive ^{55}Fe (Perkin Elmer) was added to the amphibactin, desferrioxamine B, and
desferrioxamine E ligand stocks and was equilibrated for at least 24 hours before use. Non-
radioactive ligand stocks were made with ^{57}Fe (Cambridge Isotopes) to set-up replicate
treatments for cell counts without radioactive iron. Trace metal clean unfiltered seawater from 15
m and 300 m was dispensed into 250 mL acid-cleaned polycarbonate bottles. Four of the bottles
were spiked with 1 nmol L^{-1} of the ^{57}Fe stocks (one for each treatment) and the rest were spiked
in duplicate with 1 nmol L^{-1} of the ^{55}Fe ligand stocks. A final concentration of 0.01%
glutaraldehyde was added to four of the bottles as dead controls and were incubated for 30
minutes before the experiment began to account for ^{55}Fe adsorption to particles.
119
120   Bottles were incubated in surface seawater flow-through incubators and dark treatments were
121   double-bagged in black polyethylene bags. After 12 hours, each bottle was filtered onto acid
122   washed 2.0 µm and 0.2 µm filters using low pressure vacuum filtration. Both filters were washed
123   three times with an oxalate wash to remove extracellular iron (Tovar-Sanchez 2003).

124
Radioactive decay for each sample was determined using a liquid scintillation counter
(Beckman-Coulter LS6500). ^{55}Fe disintegrations per minute (dpm) were background and dead
control corrected and converted to iron uptake using a standard curve. Cell count samples were
taken from ^{57}Fe treatments at the beginning and end of each experiment (Supplementary
Methods). Iron uptake per cell was estimated based on flow cytometry cell counts of
cyanobacteria, photosynthetic picoeukaryotes and heterotrophic bacteria (Table S2), and Fe:C
ratios for heterotrophic bacteria were determined only in the dark treatments using heterotrophic
bacteria abundances and an estimated carbon per cell of 12.4 fg C cell^{-1} (Boyd et al. 2015). The
turnover time of iron in each treatment was calculated by,
134
135 (1) $$turnover\ time = {total\ iron\ uptake}/{total\ iron}$$
136
137 where the total iron uptake (pmol L^{-1} day^{-1}) is the sum of the radioactive iron uptake in the 2.0
138 and 0.2 μm fraction and the total iron is the radioactive iron added (1 nmol L^{-1}) plus the *in situ*
139 dissolved iron. Residence times of iron (days) was calculated as 1/turnover time.
140
141 *2.3 Genomic sample collection and analyses*
142
Details on the sampling, processing and data analyses of metagenomes collected at Station
ALOHA are presented elsewhere (Mende et al. 2017; Luo et al. 2020). The ALOHA gene
catalog consists of ~8.9 million non-redundant genes identified at Station ALOHA between the
surface and 1,000 m over the course of ~1.5 years collected on 0.2 μm filters (Mende et al.
2017). The ALOHA gene catalog was probed for both NRPS and NIS biosynthetic gene clusters
(BGCs) by searching for the conserved domains present within biomarker genes. For NIS
pathways, hidden Markov models (HMMs) for the conserved domains pfam04183 (IucA/IucC
siderophore synthase) and pfam06276 (FhuF-like reductase) were searched using HMMER
(Eddy 2011). For NRPS biosynthetic pathways, the HMM for the conserved domain pfam00501,
an AMP-binding domain inclusive of the adenylation domain of NRPS modules, was searched.
The resulting 28,000+ genes were then screened with antiSMASH (Medema et al. 2011) to
determine the presence of a complete or near- complete NRPS module. The resulting NRPS
BGCs were then further examined to determine those specific to siderophore biosynthesis
(Supplementary Methods). In total, 38 putative siderophore BGCs were identified and used for
downstream analysis (Table S3).
158
To examine the distribution of siderophore BGCs with depth and time at Station ALOHA,
metagenomic reads from samples collected on HOT cruises between May 2015 and April 2016
(n = 132, 5-500 m) (Luo et al. 2020) were mapped to open reading frames (ORFs) of the
ALOHA gene catalog using BWA (Li and Durbin 2009), counted with FeatureCounts (Liao et al.
2014), and normalized to reads per kilobase per million (RPKM). Reads from the 2015-2016
dataset recruited to 24 of the 38 siderophore BGCs identified in the ALOHA gene catalog (Table
S3).
166
167 **3. Results**
168 *3.1 Seasonal distribution of siderophores and siderophore biosynthesis genes at Station ALOHA*
169

Dissolved siderophore concentrations and the presence of siderophore BGCs changed seasonally
at Station ALOHA (Fig. 1A). Average siderophore concentrations were highest in the spring,
summer and fall, and were lower in winter, though variability was high (Fig. 1A). Similar
patterns were observed in siderophore BGCs from metagenomes (Fig. 1B). The normalized
abundance of siderophore BGCs peaked in the spring and summer and were lowest in winter,
similar to the concentrations of total dissolved siderophores (Fig. 1C). In the spring, peaks in
siderophore BGCs were largely associated with surface waters (Fig. 2A), though subsurface
maxima were also observed below the deep chlorophyll maximum (DCM) and the euphotic zone
(250 m). Summer maxima in siderophore BGCs were almost exclusively associated with the
DCM (Fig. 2B). A similar pattern was observed in the fall, where siderophore BGCs were
maximal in the DCM and 175 m (Fig. 2C). The winter showed lower genetic potential for
siderophore biosynthesis overall, with median ORF abundances less than 0.2 RPKM at all depths
(Fig. 2D). Overall, the highest concentrations of both dissolved siderophores and normalized
read counts of siderophore BGCs were observed in the upper euphotic zone (< 75 m), the DCM
(75-125 m), and upper mesopelagic (225-300 m).
185
Several different types of siderophores were characterized in the water column and are capable
of being synthesized by the *in situ* microbial community (Table 1). Of those identified in the
water column, primarily hydroxamate siderophores were detected, with ferrioxamines and
amphibactins being the most common classes (Table 1). The siderophore BGCs identified
largely belonged to catecholate and carboxylate-type siderophores (Table 1). Two ferrioxamine
BGCs were identified in the gene catalog; however, these BGCs were not detected in the 2015-
2016 metagenomic dataset. Amphibactin BGCs, as determined by the presence of a N^5-acyl-N^5-
hydroxyornithine NRPS adenylation domain, were not detected. Several siderophores were
found both in the water column and as a BGC in the metagenomes, including petrobactin,
pseudoalterobactin, piscibactin and vibrioferrin (Table 1). The biosynthesis genes were primarily
from the genera *Vibrio*, *Photobacterium*, *Alteromonas*, and *Pseudoalteromonas* (Table S3).
197
198 *3.4 Uptake and residence time of siderophore-bound iron by the in situ microbial community*
199
Siderophore-bound iron was readily accessible by the microbial community in both the surface
(15 m) and upper mesopelagic (300 m) waters at Station ALOHA. Uptake rates of inorganic
iron, which was likely associated with unknown natural organic iron-binding ligands (denoted
"FeL"), were higher than for siderophore-bound iron (Fig. 3A). The uptake rate of FeL was also
higher when the experiment was performed in the light compared to the dark (Fig. 3A). The
uptake rate varied for the different iron-siderophore complexes. The iron-amphibactin was taken
up the fastest, followed by ferrioxamine E and ferrioxamine B. In contrast to FeL, the uptake rate
of siderophore-bound iron was always higher in the dark compared to the light (Fig. 3A).
Overall, the total iron uptake was also higher in the 0.2 μm fraction compared to the 2.0 μm
fraction in all experiments and treatments (Fig. 3A). Surprisingly, the uptake rate of siderophore-
bound iron was higher at 300 m compared to 15 m despite the 3-fold lower cell abundances,
leading to high iron uptake per cell (Table S2) and shorter residence times of iron (Fig. 3B).
Assuming the total iron uptake in the dark was due to heterotrophic bacteria, the estimated Fe:C
(μmol:mol) of the heterotrophic bacteria community in surface waters ranged from 6.8-68
μmol:mol while at 300 m they ranged from 88-710 μmol:mol (Table S2). The average residence
time of iron was shorter at 300 m compared to 15 m (Fig. 3B), though this difference was only

significant in the DFB treatment (pairwise t-test, $p < 0.05$). The average residence time of iron at
15 m ranged from 15 days in the FeL treatment to 252 and 215 days in the DFE and DFB
treatments, respectively. The average residence times at 300 m ranged from 9 days in the FeL
treatment to 65 days in the DFB treatment.
220
221 **4. Discussion**
222
223 *4.1 Siderophore production and uptake at Station ALOHA*
224
Siderophore BGCs and measurements of dissolved siderophores highlight seasonal differences in
microbial iron acquisition at Station ALOHA (Fig. 1 and 2). Both siderophore concentrations
and biosynthesis potential were greatest in the spring, summer and fall and were almost entirely
absent in winter. Although our data do not capture instantaneous siderophore production by
bacteria, it is likely that siderophore secretion is being used to solubilize particulate iron (Bundy
et al. 2018) and to compete with other organisms for iron (Boiteau et al. 2016). Several genera
found to have siderophore BGCs, such as *Vibrio* and *Alteromonas*, contain known copiotrophic
bacteria, suggesting that siderophore production may be common to heterotrophs that are
opportunistic and respond to episodic inputs of nutrients or carbon during the spring and summer
(Fontanez et al. 2015; Pelve et al. 2017; Church et al. 2021; Poff et al. 2021; Leu et al. 2022).
Interestingly, there was little overlap in the dominant siderophores measured and the BGCs
identified (Table 1). The siderophore BGCs were primarily catecholates, while those measured in
the water column were primarily hydroxamates. This implies that catecholate siderophores could
be below our detection limits, or rapidly cycled in the water column. Conversely, the structural
discrepancy between siderophores detected in BGCs and those dominating water column
measurements at Station ALOHA and in other regions (Mawji et al. 2008b; Boiteau et al. 2016;
Bundy et al. 2018), suggests that the importance of these compounds may be underestimated by
metagenomic studies.
243
Microbial iron recycling via the production and consumption of siderophore-bound iron likely
involves rapid turnover of these compounds. The iron uptake experiments conducted here
provided estimates of the residence time of weakly bound (FeL) versus siderophore-bound iron.
Siderophore-bound iron was taken up less rapidly by the microbial community and had average
residence times ranging from 15-250 days (Fig. 3A), while FeL was highly labile and had
average residence times of 1-2 weeks (Fig. 3B). The FeL treatment likely had a short residence
time since inorganic iron may associate with weaker ligands when added to seawater, which are
more accessible to phytoplankton using ferric reductases as an uptake mechanism (Maldonado
and Price 2001; Salmon et al. 2006; Lis et al. 2015; Coale et al. 2019). Iron uptake rates in FeL
treatments were higher in the light than in the dark, suggesting significant contributions by
photoautotrophs. Uptake rates of siderophore-bound iron in contrast were always higher in the
dark than the light, suggesting no significant photochemical effect on the uptake rate of
siderophores (Barbeau 2006), and that siderophore iron consumption was likely not controlled by
photosynthetic plankton.
258
The siderophore treatments had slower uptake rates and longer residence times than FeL. Iron
associated with amphibactins was taken up more quickly than ferrioxamine E or B (Fig. 3A).
Amphibactins have a lower binding affinity to dissolved iron (stability constants = 12.0-12.5)

compared to ferrioxamines (14.0-14.4; (Bundy et al. 2018)), and iron is likely more easily
reducible from amphibactins. Alternatively, more members of the microbial community may
possess Fe-amphibactin uptake pathways. Siderophore-bound iron also appeared to be more
rapidly internalized by heterotrophic bacteria compared to large photoautotrophs at Station
ALOHA. Iron uptake in the 0.2 μm size fraction was assumed to be dominated by free-living
bacteria, and the total iron uptake in this fraction was always greater than the 2.0 μm (Fig. 3A).
An unexpected result was the faster uptake rates and shorter residence times of siderophore-
bound iron at 300 m compared to 15 m. Macronutrient concentrations are elevated at 300 m
relative to 15 m, so it is possible iron uptake rates were enhanced due to relief from
macronutrient limitation. However, nitrogen amendments (500 nmol L^{-1}) to the surface
community should have alleviated nitrogen limitation. The experiments at 300 m were also
incubated at the same temperature as the 15 m experiments, likely enhancing the uptake rate of
iron relative to *in situ* rates, however it is not clear why this would stimulate uptake rates above
those measured in surface waters.
276
Fast uptake rates and short residence times of iron at 300 m are likely related to differences in
iron acquisition by the microbial community at these depths. The iron requirements of
heterotrophic bacteria are not well known, but evidence suggests they are elevated relative to
large phytoplankton (Fe:C ~ 2.0 μmol: mol) or cyanobacteria (Fe:C ~ 19 μmol: mol) (Strzepek et
al. 2005). Published estimates for heterotrophic bacteria Fe:C range from 7.5 μmol: mol (Tortell
et al. 1996) to 83 μmol: mol (Mazzotta et al. 2020). Our range in Fe:C in surface waters
encompass these values, but exceed this range at 300 m (Table S2). One explanation is the
carbon per cell estimates for heterotrophic bacteria are not representative of upper mesopelagic
bacteria, causing the Fe:C to be high. Nitrifiers for example, have both high carbon (Watson and
Mandel 1971) and iron content (Saito et al. 2020). Alternatively, the microbial community at
these depths could have elevated iron requirements, as the electron transport systems used for
respiration is iron-intensive (Tortell et al. 1996) and changes in the lability or oxidation state of
organic matter might impact iron demand. The community at 300 m may also be exposed to
elevated iron in the form of particulate iron throughout the season, and may engage in luxury
uptake and storage (Mazzotta et al. 2020). The high iron uptake rates per cell led to shorter
residence times at 300 m, particularly for siderophore-bound iron (Fig. 3B) consistent with the
shorter residence times observed in the DCM in other studies (Hawco et al. 2022). Siderophore
turnover may be particularly fast in the mesopelagic as bacteria recycle and regenerate iron from
sinking particles (Bundy et al. 2018). The reason for the enhanced iron uptake at 300 m is
unknown, but understanding the iron demands of mesopelagic bacteria could have important
implications for carbon and iron cycling in the mesopelagic.
298
299   *4.2 Environmental drivers of siderophore production and consumption*
300
Seasonality in productivity, particle accumulation, iron supply and export have all been
documented at Station ALOHA (Fitzsimmons et al. 2015; Hawco et al. 2021; Karl et al. 2021)
and such changes likely impact microbial siderophore production. As productivity increases in
the late spring in the gyre, dissolved iron concentrations are drawn down and competition for
iron intensifies (Fitzsimmons et al. 2015), setting up conditions that likely favor the production
of siderophores to recycle and retain iron in the euphotic zone (Boyd et al. 2015; Hawco et al.
2022). In the summer as carbon export fluxes increase, dissolved siderophore concentrations

increased deeper in the water column and the siderophore BGCs peaked closer to the DCM (Fig
1A, 2B). This could reflect further competition and recycling of iron later in the growing season,
including greater overall iron demand by the heterotrophic bacteria community to process
particulate organic matter. There could also be a shift in the microbial community at depth to
reflect more siderophore-producing microbes. Similarly, observations during the fall reveal a
peak in the abundance of siderophore BGCs at and below the DCM (Fig. 2C). This may reflect
intensified iron recycling or the high demand for iron by the heterotrophic bacteria community as
they respire sinking organic matter associated with the export pulse at Station ALOHA. The
short residence times of iron at 300 m also suggest intense competition for iron at these depths.
Finally, in the winter, siderophore concentrations and the presence of siderophore BGCs reduces
significantly, likely reflecting the decrease in iron demand of heterotrophic bacteria as
productivity and organic matter substrates diminish, or increased mixing that dilutes siderophore
concentrations. This dataset provides compelling evidence that ligand production by the
microbial community plays an important role in the seasonal cycling and retention of iron at
Station ALOHA.
323

5. Acknowledgements
Special thanks to the Captain and crew of the R/V *Kilo Moana* and R/V *Ka'imikai-O-Kanaloa*
for help with sample collection, and to Kendra A. Turk-Kubo for flow cytometry analyses. We
would also like to thank Travis Mellett for helpful comments on the manuscript. RMB was
funded by a Life Sciences-Simons Early Career Investigator in Marine Microbial Ecology and
Evolution Awards (Award# 618401) and a Woods Hole Oceanographic Institution postdoctoral
fellowship. LEM was funded by a Simons Postdoctoral Fellowship in Marine Microbial Ecology
(Award #729162). MJC was funded by the Simons Collaboration on Ocean Processes (Award
#721221).

Table 1. Siderophores measured in the water column and siderophore biosynthetic gene clusters (BGCs) identified at Station
ALOHA. Siderophores identified in the water column are grouped according to type and are shown in the table based on that
grouping. For example, multiple different forms of amphibactins were detected (Table S1) but are all grouped here under
"amphibactins." Specific siderophore IDs can be found in Supplementary Table 1. The '+' denotes that the specific siderophore BGC
was found in the 2015-2016 metagenomic dataset, while '-' indicates it was not detected.

Siderophore	Type	Biosynthetic Pathway	Depth(s)	Season	BGC in gene catalog
aerobactin	mixed	NIS	-	-	+
amphibactins	hydroxamate	NRPS	300-400	summer	-
anguibactin	catecholate	NRPS	-	-	+
enterobactin	catecholate	NRPS	-	-	+
ferrioxamaines	hydroxamate	NIS	15-400	spring, summer, winter	-
petrobactin	mixed	NIS/NRPS hybrid	15	winter	+
piscibactin	catecholate	NRPS	15	winter	+
(pseudo)alterobactin	catecholate	NRPS	15	winter	+
synechobactins	mixed	NRPS	15, 150	spring, summer, winter	-
vibrioferrin	carboxylate	NIS	15	winter	+
vibriobactin	catecholate	NRPS	-	-	+

Figure 1. (A) Seasonal average dissolved siderophore concentrations collected at Station ALOHA. Error bars, when present, represent the standard error of samples collected in the same depth range in the same season. When no data is present for a given depth range and season, no samples were collected during that time. (B) Distribution across depth of the abundance of siderophore BGCs detected at Station ALOHA between May 2015 – April 2016. Blue points track the median ORF abundances at each depth and boxplots display quartile ranges of the total dataset at each depth with outliers plotted as distinct points. No data present between 250 – 500 m. (C) Monthly distributions of the abundance of siderophore BGCs detected at Station ALOHA between May 2015 – April 2016. Blue points track the median ORF abundances for each month and boxplots display quartile ranges of the total dataset for each month with outliers plotted as distinct points.

Figure 2. Distribution across seasons in spring (A), summer (B), fall (C) and winter (D) and with depth of siderophore BGCs detected at Station ALOHA between May 2015 – April 2016. Blue points track the median ORF abundances and boxplots display quartile ranges of the total dataset for each season at each depth with outliers plotted as distinct points. No data present between 250 – 500 m.

Figure 3. (A) Dissolved iron uptake rates in the surface (15 m) and upper mesopelagic (300 m) in the 2.0 μm and 0.2 μm size fraction (hatched bars) in the dark (grey shaded) and in the light. (B) Residence times of iron in the inorganic iron treatment (FeL), Fe-amphibactin treatment (Amp), ferrioxamine E treatment (DFE), and ferrioxamine B treatment (DFB) at 15 m (yellow) and 300 m (blue). Residence times at 300 m were statistically different from 15 m only in the DFB treatment (pairwise t-test, $p < 0.05$)

References
Barbeau, K. 2006. Photochemistry of organic iron(III) complexing ligands in oceanic systems.
Photochem. Photobiol. **82**: 1505–1516. doi:10.1562/2006-06-16-ir-935
Boiteau, R. M., J. N. Fitzsimmons, D. J. Repeta, and E. A. Boyle. 2013. Detection of Iron
Ligands in Seawater and Marine Cyanobacteria Cultures by High-Performance Liquid
Chromatography-Inductively Coupled Plasma-Mass Spectrometry. Anal. Chem. **85**: 4357–
4362. doi:10.1021/ac3034568
Boiteau, R. M., D. R. Mende, N. J. Hawco, and others. 2016. Siderophore-based microbial
adaptations to iron scarcity across the eastern Pacific Ocean. Proc. Natl. Acad. Sci. **113**:
14237–14242.
Boiteau, R. M., C. P. Till, T. H. Coale, J. N. Fitzsimmons, K. W. Bruland, and D. J. Repeta.
2019. Patterns of iron and siderophore distributions across the California Current System.
Limnol. Oceanogr. **64**: 376–389. doi:10.1002/lno.11046
Boyd, P. W., R. F. Strzepek, M. J. Ellwood, D. A. Hutchins, S. D. Nodder, B. S. Twining, and S.
W. Wilhelm. 2015. Why are biotic iron pools uniform across high- and low-iron pelagic
ecosystems? Global Biogeochem. Cycles **29**: 1028–1043. doi:10.1002/2014gb005014
Bundy, R. M., R. M. Boiteau, C. McLean, K. A. Turk-Kubo, M. R. McIlvin, M. A. Saito, B. A.
S. Van Mooy, and D. J. Repeta. 2018. Distinct Siderophores Contribute to Iron Cycling in
the Mesopelagic at Station ALOHA. Front. Mar. Sci. **5**: 1–15.
doi:10.3389/fmars.2018.00061
Church, M. J., E. Kyi, R. O. Hall Jr, D. M. Karl, M. Lindh, A. Nelson, and E. K. Wear. 2021.
Production and diversity of microorganisms associated with sinking particles in the
subtropical North Pacific Ocean. Limnol. Oceanogr. **66**: 3255–3270.
Coale, T. H., M. Moosburner, A. Horák, M. Oborník, K. A. Barbeau, and A. E. Allen. 2019.
Reduction-dependent siderophore assimilation in a model pennate diatom. Proc. Natl. Acad.
Sci. U. S. A. **116**: 23609–23617. doi:10.1073/pnas.1907234116
Cordero, O. X., L.-A. Ventouras, E. F. DeLong, and M. F. Polz. 2012. Public good dynamics
drive evolution of iron acquisition strategies in natural bacterioplankton populations. Proc.
Natl. Acad. Sci. **109**: 20059–20064. doi:10.1073/pnas.1213344109
Eddy, S. R. 2011. Accelerated profile HMM searches. PLoS Comput. Biol. **7**: e1002195.
Fitzsimmons, J. N., and E. A. Boyle. 2014. Assessment and comparison of Anopore and cross
flow filtration methods for the determination of dissolved iron size fractionation into soluble
and colloidal phases in seawater. Limnol. Oceanogr. **12**: 246–263.
doi:10.4319/lom.2014.12.246
Fitzsimmons, J. N., C. T. Hayes, S. N. Al-Subiai, R. Zhang, P. L. Morton, R. E. Weisend, F.
Ascani, and E. A. Boyle. 2015. Daily to decadal variability of size-fractionated iron and
iron-binding ligands at the Hawaii Ocean Time-series Station ALOHA. Geochim.
Cosmochim. Acta **171**: 303–324.
Fontanez, K. M., J. M. Eppley, T. J. Samo, D. M. Karl, and E. F. DeLong. 2015. Microbial
community structure and function on sinking particles in the North Pacific Subtropical
Gyre. Front. Microbiol. **6**: 469.
Hawco, N. J., B. Barone, M. J. Church, and others. 2021. Iron depletion in the deep chlorophyll
maximum: mesoscale eddies as natural iron fertilization experiments. Global Biogeochem.
Cycles **35**: e2021GB007112.
Hawco, N. J., S.-C. Yang, P. Pinedo-González, E. E. Black, J. Kenyon, S. Ferrón, X. Bian, and
S. G. John. 2022. Recycling of dissolved iron in the North Pacific Subtropical Gyre.

Limnol. Oceanogr. doi:https://doi.org/10.1002/lno.12212

Hayes, C. T., J. N. Fitzsimmons, E. A. Boyle, D. McGee, R. F. Anderson, R. Weisend, and P. L. Morton. 2015. Thorium isotopes tracing the iron cycle at the Hawaii Ocean Time-series Station ALOHA. Geochim. Cosmochim. Acta **169**: 1–16.

Hider, R. C., and X. Kong. 2010. Chemistry and biology of siderophores. Nat. Prod. Rep. **27**: 637–657. doi:10.1039/b906679a

Karl, D., R. Letelier, L. Tupas, and J. Dore. 1997. The role of nitrogen fixation in biogeochemical cycling in the subtropical North Pacific Ocean. Nature **388**: 533.

Karl, D. M., R. M. Letelier, R. R. Bidigare, K. M. Björkman, M. J. Church, J. E. Dore, and A. E. White. 2021. Seasonal-to-decadal scale variability in primary production and particulate matter export at Station ALOHA. Prog. Oceanogr. **195**: 102563.

Karl, D. M., and R. Lukas. 1996. The Hawaii Ocean Time-series (HOT) program: Background, rationale and field implementation. Deep Sea Res. Part II Top. Stud. Oceanogr. **43**: 129–156.

Kramer, J., Ö. Özkaya, and R. Kümmerli. 2020. Bacterial siderophores in community and host interactions. Nat. Rev. Microbiol. **18**: 152–163.

Letelier, R. M., K. M. Björkman, M. J. Church, and others. 2019. Climate-driven oscillation of phosphorus and iron limitation in the North Pacific Subtropical Gyre. Proc. Natl. Acad. Sci. U. S. A. **116**: 12720–12728. doi:10.1073/pnas.1900789116

Leu, A. O., J. M. Eppley, A. Burger, and E. F. DeLong. 2022. Diverse Genomic Traits Differentiate Sinking-Particle-Associated versus Free-Living Microbes throughout the Oligotrophic Open Ocean Water Column. MBio **13**: e0156922. doi:10.1128/mbio.01569-22

Li, H., and R. Durbin. 2009. Fast and accurate short read alignment with Burrows--Wheeler transform. bioinformatics **25**: 1754–1760.

Liao, Y., G. K. Smyth, and W. Shi. 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics **30**: 923–930.

Lis, H., Y. Shaked, C. Kranzler, N. Keren, and F. M. M. Morel. 2015. Iron bioavailability to phytoplankton: an empirical approach. ISME J. **9**: 1003–1013. doi:10.1038/ismej.2014.199

Longhurst, A. R. 2010. Ecological geography of the sea, Elsevier.

Luo, E., J. M. Eppley, A. E. Romano, D. R. Mende, and E. F. DeLong. 2020. Double-stranded DNA virioplankton dynamics and reproductive strategies in the oligotrophic open ocean water column. ISME J. **14**: 1304–1315. doi:10.1038/s41396-020-0604-8

Maldonado, M. T., and N. M. Price. 2001. Reduction and transport of organically bound iron by Thalassiosira oceanica (Bacillariophyceae). J. Phycol. **37**: 298–309. doi:10.1046/j.1529-8817.2001.037002298.x

Manck, L. E., J. Park, B. J. Tully, A. M. Poire, R. M. Bundy, C. L. Dupont, and K. A. Barbeau. 2022. Petrobactin, a siderophore produced by Alteromonas, mediates community iron acquisition in the global ocean. ISME J. **16**: 358–369.

Mawji, E., M. Gledhill, J. A. Milton, and others. 2008a. Hydroxamate Siderophores: Occurance and Importance in the Atlantic Ocean. Environ. Sci. Technol. **42**: 8675–8680.

Mawji, E., M. Gledhill, J. A. Milton, and others. 2008b. Hydroxamate Siderophores: Occurrence and Importance in the Atlantic Ocean. Environ. Sci. Technol. **42**: 8675–8680. doi:10.1021/es801884r

Mazzotta, M. G., M. R. McIlvin, and M. A. Saito. 2020. Characterization of the Fe metalloproteome of a ubiquitous marine heterotroph, Pseudoalteromonas (BB2-AT2): multiple bacterioferritin copies enable significant Fe storage†. Metallomics **12**: 654–667.

453   doi:10.1039/d0mt00034e

454 Medema, M. H., K. Blin, P. Cimermancic, and others. 2011. antiSMASH: rapid identification,
455   annotation and analysis of secondary metabolite biosynthesis gene clusters in bacterial and
456   fungal genome sequences. Nucleic Acids Res. **39**: W339--W346.

457 Mende, D. R., J. A. Bryant, F. O. Aylward, J. M. Eppley, T. Nielsen, D. M. Karl, and E. F.
458   DeLong. 2017. Environmental drivers of a microbial genomic transition zone in the ocean's
459   interior. Nat. Microbiol. **2**: 1367–1373. doi:10.1038/s41564-017-0008-3

460 Moore, L. E., M. I. Heller, K. A. Barbeau, J. W. Moffett, and R. M. Bundy. 2021. Organic
461   complexation of iron by strong ligands and siderophores in the eastern tropical North
462   Pacific oxygen deficient zone. Mar. Chem. **236**: 104021.

463 Pelve, E. A., K. M. Fontanez, and E. F. DeLong. 2017. Bacterial succession on sinking particles
464   in the ocean's interior. Front. Microbiol. **8**: 2269.

465 Pinedo-González, P., N. J. Hawco, R. M. Bundy, and others. 2020. Anthropogenic Asian
466   aerosols provide Fe to the North Pacific Ocean. Proc. Natl. Acad. Sci. U. S. A. **117**: 27862–
467   27868. doi:10.1073/pnas.2010315117

468 Poff, K. E., A. O. Leu, J. M. Eppley, D. M. Karl, and E. F. DeLong. 2021. Microbial dynamics
469   of elevated carbon flux in the open ocean's abyss. Proc. Natl. Acad. Sci. **118**: e2018269118.

470 Saito, M. A., M. R. McIlvin, D. M. Moran, and others. 2020. Abundant nitrite-oxidizing
471   metalloenzymes in the mesopelagic zone of the tropical Pacific Ocean. Nat. Geosci. **13**:
472   355–362.

473 Salmon, T. P., A. L. Rose, B. A. Neilan, and T. D. Waite. 2006. The FeL model of iron
474   acquisition: Nondissociative reduction of ferric complexes in the marine environment.
475   Limnol. Oceanogr. **51**: 1744–1754.

476 Sandy, M., and A. Butler. 2009. Microbial iron acquisition: marine and terrestrial siderophores.
477   Chem. Rev. **109**: 4580–4595.

478 Shaked, Y., and H. Lis. 2012. Disassembling iron availability to phytoplankton. Front.
479   Microbiol. **3**: 123. doi:10.3389/fmicb.2012.00123

480 Strzepek, R. F., M. T. Maldonado, J. L. Higgins, J. Hall, K. Safi, S. W. Wilhelm, and P. W.
481   Boyd. 2005. Spinning the "ferrous wheel": The importance of the microbial community in
482   an iron budget during the FeCycle experiment. Global Biogeochem. Cycles **19**.
483   doi:10.1029/2005GB002490

484 Tortell, P. D., M. T. Maldonado, and N. M. Price. 1996. The role of heterotrophic bacteria in
485   iron-limited ocean ecosystems. Lett. to Nat. **383**: 330–332.

486 Tovar-Sanchez, a. 2003. A trace metal clean reagent to remove surface-bound iron from marine
487   phytoplankton. Mar. Chem. **82**: 91–99. doi:10.1016/S0304-4203(03)00054-9

488 Vraspir, J. M., and A. Butler. 2009. Chemistry of Marine Ligands and Siderophores. Ann. Rev.
489   Mar. Sci. **1**: 43–63. doi:10.1146/annurev.marine.010908.163712

490 Watson, S. W., and M. Mandel. 1971. Comparison of the morphology and deoxyribonucleic acid
491   composition of 27 strains of nitrifying bacteria. J. Bacteriol. **107**: 563–569.

492