Comparative genomics and transcriptomic response to root exudates of six rice root-associated *Burkholderia sensu lato* species

Adrian Wallner[1£], Agnieszka Klonowska[1], Ludivine Guigard[1], Isabelle Rimbault[1], Eddy Ngonkeu[2], Phuong Nguyen[3] Gilles Béna[1], Lionel Moulin[1*]

[1]PHIM Plant Health Institute, Univ Montpellier, IRD, CIRAD, INRAE, Institut Agro, Montpellier, France

[2]Institute of Agronomic Research for Development (IRAD), PO Box 2123, Yaoundé, Cameroon

[3]University of Science and Technology of Hanoi (USTH); 18 Hoang Quoc Viet, Cau Giay District, Hanoi, Vietnam

* Corresponding author: Lionel Moulin, lionel.moulin@ird.fr

£ Present address: Unité Résistance induite et Bio-protection des plantes, Université de Reims Champagne-Ardenne, Reims, France

Keywords: RNAseq, root exudates, PGPR, rhizobacteria, symbiosis, Oryza sativa

Abstract

The rhizosphere can be described as the portion of soil under influence of plant roots through deposition and exudation of various molecules. It is a nutrient rich environment that is favorable for a multitude of soil-dwelling bacteria. Beyond being a reliable nutrient provider, some bacteria will perceive the plant as a potential host and undertake root colonization leading to mutualistic or parasitic interactions. Bacteria of the *Burkholderia* and *Paraburkholderia* genera are frequently found in the rhizosphere of rice. While the latter are often described as plant growth promoting species, *Burkholderia* are mostly studied for their human opportunistic traits. Here, we used root exudate stimulation on three *Burkholderia* and three *Paraburkholderia* strains isolated from rice roots to characterize their preliminary adaptation to the rice host at the transcriptomic level. Instead of the awaited genus-dependent adaptation, we observed a strongly species-specific response for all tested strains. While all bacteria originate from the rice environment, there are great disparities in their levels of adaptation following the sensing of root exudates. We further report the shared major functions that were differentially regulated in this early step of bacterial adaptation to plant

colonization, including amino acids and putrescine metabolism, the Entner-Doudoroff (ED)

pathway as well as cyclic diguanylate monophosphate (c-di-GMP) cycling.

Background

Diverse species of *Burkholderia* and *Paraburkholderia* have been repeatedly isolated from

healthy rice roots (Govindarajan et al., 2008; Mattos et al., 2008; Wang et al., 2016a; Kang et

al., 2017; Arriel-Elias et al., 2019; Song et al., 2019; Hameed et al., 2020). Their central role

in rice ecology is further suggested by several population analyses placing *Burkholderia* and

Paraburkholderia as being among the predominant taxa found in the rice rhizosphere with

some specimens acting as microbiome structuring hubs (Ikeda et al., 2014; Hardoim, 2015; de

Souza et al., 2015; Yu et al., 2018).

Both *Burkholderia* and *Paraburkholderia* are described as efficient rhizosphere colonizers

and can achieve plant growth promotion through diverse strategies including nutrient

solubilization, phytohormone production and interference, siderophore production as well as

enhancing the plants tolerance to abiotic stresses. Several *Paraburkholderia* species are free

living nitrogen fixers and some induce nitrogen-fixing nodules on legume plants (Dall'Agnol

et al., 2017; Estrada-de los Santos et al., 2018). Further *Paraburkholderia* species are root

endophytes and enhance the growth of a number of major crops such as corn, rice and

sugarcane (Perin et al., 2006; Govindarajan et al., 2008; Mattos et al., 2008; Mitter et al.,

2013). While several *Burkholderia* species e.g., *B. glumae, B. gladioli, B. plantari*, are

responsible for serious diseases in rice crops (Kim et al., 2021), others seem to have an

intrinsic ability to colonize roots which is independent of plant genotype and do not

necessarily cause disease (Vidal-Quist et al., 2014). *B. cenocepacia* or *B. vietnamiensis* for

instance have been observed to cause plant tissue water-soaking in onion but have otherwise

been isolated from healthy plants (Engledow et al., 2004; Jacobs et al., 2008). While

Burkholderia strains are mostly reported for their biocontrol efficiency, some achieve plant

growth promotion through direct hormonal stimulation and nutrition (Trân Van et al., 2000;

Araújo et al., 2016; Batista et al., 2016; Wang et al., 2016b; Mullins et al., 2019; Chávez-

Ramírez et al., 2020). Genome mining approaches underlined the importance of secondary

metabolites in *Burkholderia* species, including an important diversity of antimicrobial

metabolites (Navarro et al., 2019). However, in contrast to *Paraburkholderia*, several

Burkholderia species were found to colonize animal hosts in addition to plants, causing

opportunistic infections and severe pathologies in humans (Galyov et al., 2010; Sousa et al.,

2011).

Here, we investigated the early strategies directing plant colonization at the onset of the interaction between rhizobacteria and their host which is mediated by the perception of root exudates (RE). Through root exudation, plants can use up to 20-40% of their photosynthetically fixed carbon to actively shape their microbiome using chemical attractants, selective carbon sources and diverse restrictive metabolites (Badri and Vivanco, 2009). Primarily genes involved in chemotaxis and motility, but also nutrient transport and metabolism are required for an efficient response to exudates. Still, as exemplified by several past studies, a bacterium's response to exudates is highly host dependent. For instance, *Pseudomonas aeruginosa* was shown to have a varying transcriptomic reaction to RE originating from two different sugar beet cultivars (Mark et al., 2005). The perception of RE can induce major metabolic adaptation in bacteria. Rice RE were found to be responsible for the differential expression of 4,4% genes in the genome of *Azoarcus* sp. BH72 (Shidore et al., 2012). More significant, a legume nodulating strain of *Paraburkholderia phymatum* differentially regulated 19.2% of its genome in response to RE of *Mimosa pudica* (Klonowska et al., 2018). Upon inoculation with maize RE, *Bacillus amyloliquefaciens* displays a two-stage response. At 24h post-treatment, the bacteria directs its metabolism towards cellular proliferation while at 48h post-treatment, most differentially expressed (DE) genes are down-regulated and production of extracellular matrix is favored (Fan et al., 2012; Zhang et al., 2015). Typically, the bacterial reaction to RE involves metabolic genes for sugar and amino-acid import and catabolism. However, functions involved at later steps of the interaction such as type 3 secretion are modulated as well upon RE perception (Hassan and Mathesius, 2012; Klonowska et al., 2018).

The present work focuses on six rice root-isolated strains evenly distributed between *Burkholderia* and *Paraburkholderia* genera. Beside the two model bacteria for plant growth promotion *B. vietnamiensis* LMG 10929 (BvLMG10929) and *P. kururiensis* M130 (PkM130), we analyzed four new rice-isolated bacterial strains, for which we provide whole-genome sequences, as well as an assessment of their rice root colonization capacity. We hypothesized that the genetic frontier between the two bacterial taxa would influence and differentiate their adaptation strategies towards plant colonization in a genus specific manner. With this in mind, we aimed at revealing each strain's adaptation signatures to the plant environment by exploring their gene expression profiles when sensing rice root exudates using a transcriptomics approach. Considering the results presented here, no distinction can be made between bacteria according to their genus as each strain displayed a specific reaction

upon plant root exudate sensing, with no significant pattern clustering con-generic bacteria,

while the regulation of several metabolic functions was shared among all bacteria.

Results

Genomic analyses of new *Burkholderia* and *Paraburkholderia* species

The newly isolated strains described in this study (Table 1) were compared to genomic

databases of *Burkholderia sensu lato* (*s.l.*) and the genomic distance to their closest relative

was computed (Figure 1). Whole genome comparisons using average nucleotide identity

(ANI) estimations showed a 98.52% score (80.14% sequence coverage) when comparing

strain ABIP441 to the proposed species *Burkholderia reimsis* sp. nov. BE51[T] (Esmaeel et al.,

2018), lying above the species threshold (> 95% identity on more than 70% cover). We thus

classified ABIP441 as *B. reimsis* (hereafter BrABIP441). We had previously proposed that

ABIP444 belongs to a novel species, *B. servocepacia* sp. nov., closely related to *Burkholderia*

cenocepacia (Wallner et al., 2019). A recent publication (Morales-Ruíz et al., 2022) updated

our proposal to validate this separate species as *B. orbicola,* with TatI-371 as the type strain.

This latter species name was thus retained in this article for ABIP444 (hereafter BoABIP444).

Strain ABIP659 displays no determining similarity to any described *Burkholderia s.l.* species.

ANI analyses suggest that it is most closely related to *Paraburkholderia tropica* without

exceeding the required threshold to infer conspecificity (91.86% identity and 76.0% sequence

coverage to *P. tropica* LMG 22274[T]). We will thus refer to it as *Paraburkholderia* sp.

(hereafter PspABIP659). At the genome level, ABIP630 shares 98.5% ANI (>70% aligned)

with *P. sabiae* LMG24235[T] and can thus be considered as belonging to this species (hereafter

PsABIP630).

All isolates demonstrate robust rice colonization capacity

As all strains were isolated from the rice rhizosphere (Table 1), we validated their ability to

colonize this environment *de novo* (Figure 2). Compared to the well investigated

BvLMG10929 and PkM130, all strains show successful root attachment and colonization.

Most bacteria reach their highest density at 7 dpi and the ratio of bacterial cells to root mass

then decreases. The three *Burkholderia spp.* all have very similar colonization efficiencies and

development profiles over time. *Paraburkholderia spp.* in turn, have more individual

colonization approaches. PsABIP630 displays the strongest abundance on roots while

PspABIP659 is the overall weakest colonizer but has the greatest dynamic in colonizing

populations. Indeed, PspABIP659 shows a striking 6-fold burst in population from 3 dpi to 7

dpi. For all strains, we evaluated the impact of inoculation on *Oryza sativa* cv. Nipponbare

growth (see Methods). No impact was observed on root and leaf weight at 60 dpi compared to
the mock treatment (Supplementary Figure S1). This is consistent with results from similar
treatments using PkM130 and BvLMG10929 (King et al., 2019).

Paraburkholderia and *Burkholderia* species share 2443 genes

An orthology search was performed to infer levels of similarity between the three
Paraburkholderia strains and the three *Burkholderia* strains (Figure 3). While all genomes are
of relatively large size compared to their Proteobacteria congeners (Table 1), there are some
discrepancies between them. The two rice-growth promoting model endophytes PkM130 and
BvLMG10929 have genomes of comparable size which are however the smallest among the
cohort analyzed here. The four remaining species have all been isolated from the rice
rhizosphere, and their genomes are considerably larger (15-20%) than those of the endophytic
PkM130 and BvLMG10929.

Altogether, 2443 genes are conserved among the six strains, representing 34% to 44% of their
total gene sets (Figure 3). Still, given the present strain selection, PspABIP659 and
PsABIP630 are best defined by their specific genome which is larger than any other
intersection. The remaining strains also have a considerable number of specific genes
although smaller than the core-genome shared between the six strains. Taken together, the
Burkholderia specific genome is 48% larger than its counterpart in *Paraburkholderia* species.

The global transcriptomic response to root exudates is strain-specific

In order to unravel bacterial genes involved in the preliminary steps of plant interaction, we
stimulated the six *Burkholderia s.l.* strains with Nipponbare rice RE and compared their
transcriptomic response to that of bacteria grown in a rich mineral medium (see Methods). A
preliminary validation of the sequencing procedure confirmed that ribodepletion efficiency
was > 70% for every strain and condition except for BvLMG10929 which only attained ~50%
efficiency (Supplementary Table S1). Nonetheless, high enough read numbers mapping
against a database of CDSs were obtained for robust bioinformatic analyses (Supplementary
Table S1). Furthermore, we verified that the exposition to RE significantly and consistently
affected the bacteria's transcriptomic responses throughout the replicates (Supplementary
Figure S2).

At a quantitative level, the general response of bacteria to RE was strain-specific with large
variations between strains. Besides BoABIP444, all strains display a majority of negatively
regulated genes upon root exudate stimulation (Figure 4). For PspABIP659, the amount of

down- and up-regulated genes is close to identical and represents the strongest response in
terms of number of genes involved which is 8-fold stronger than for the weakest responder
BrABIP441.

We further undertook an analysis of the conserved transcriptomic response across strains.
Despite the level of genetic conservation between the six strains (Figure 3), the core-genome
is not used in the primary steps of plant sensing as only two genes are DE in all six bacteria
(Figure 5). One codes for an outer membrane porin (Table 2 & Supplementary Table S2)
without further characterization. It is down-regulated in PsABIP630 and up-regulated in all
remaining strains. The second encodes a TonB protein (Table 2 & Supplementary Table S2)
which is associated with outer membrane receptors ExbB and ExbD. The differential
expression of e*xbBD* is conserved in all strains besides BrABIP441 and follows the trend of
the respective *tonB* regulation in each strain. It is up-regulated in PspABIP659 and
BvLMG10929 and down-regulated in PsABIP630 and PkM130. If we exclude BrABIP441
from the analysis, a final gene is detected to be up-regulated in every strain, the Entner-
Doudoroff pathway enzyme coding gene *edd* (Table 2 & Supplementary Table S2).

Beside the intersection between PspABIP659 and PkM130, the levels of co-regulation are
often higher in inter-generic than in intra-generic duos (Figure 5). The levels of conserved
regulation within *Burkholderia* and *Paraburkholderia* trios is noticeably low and dominantly
differential in *Paraburkholderia* (Figure 5).

The transcriptomic response is mainly nutrient directed

180
The classification according to COG categories of the DE genes comforts the hypothesis that
each strain has a dominantly individual response towards RE (Figure 6). Still, some trends can
be inferred. In each strain, most DE genes involved in carbohydrate transport and metabolism
are upregulated, indicating that the strains are taking advantage of the increased source of
carbon. The reaction to other nutrients such as amino acids, lipids and ions is more disparate
from strain to strain. Still, it appears that *Paraburkholderia* strains are globally more sensitive
to the presence of lipids contained in RE.

Our experimental approach used RE that were recovered from rice seedlings having grown for
5 days in a 6-fold diluted Hoagland nutritive medium (Supplementary Table S3). This
experimental setting has been used before to assess the plant's transcriptomic response
elicited upon bacterial inoculation (King et al., 2019). However, diluted nutrients contained in
this medium were assimilated to the recovered RE fraction. Thus, the bacterial response

concerning elements contained in the hydroponic medium must be considered with care. As it
was designed for plant nutrition the hydroponic medium does not contain any source of
carbon. It does however hold macro-nutrients such as nitrogen, phosphorus and potassium.
Additional elements eliciting a bacterial response are iron and sulfur. Still, the transcriptomic
response to these nutrients varies depending on the bacterial strain (Table 2 & Supplementary
Table S2).

Overall, we detected elements of the sulfite reductase complex (*cys*), the alkanesulfonate
transport system (*ssuBCD*) and the taurine transporter *tauABC* which can be linked to sulfure
nutrition and are DE only in *Paraburkholderia* strains. Elements linked to both assimilatory
and respiratory nitrate transport and reduction were DE in several strains (Table 2 &
Supplementary Table S2). Concerning iron nutrition, whole NRPS clusters, siderophore
receptors and bacterioferritins are dominantly negatively regulated in most strains.

While nitrogen is a main constituent of the hydroponic medium, it is present as ammonium-
conjugates and not nitrates. An increased nitrate metabolism was reported upon stimulation of
PkM130 with rice macerate (Coutinho et al., 2015). The same study further reported the
upregulation of a transporter complex involved in sulfate acquisition (ANSKv1_70577-81)
which is also upregulated in the present study. Furthermore, the iron uptake complex is
comparably downregulated in response to rice macerate (Coutinho et al., 2015) and what we
observe with RE. These observations attest that the supposedly medium-induced regulations
described above could in fact result from the composition of RE.

Amino acid metabolism is activated upon exudate sensing
Several strains respond to the presence of amino acids from the exudates by downregulating
the synthesis of dedicated transporters consistent with the strong adaptation of amino acid
metabolism in response to RE which was repeatedly reported (Mark et al., 2005; Fan et al.,
2012; Yi et al., 2017).. PkM130, BvLMG10929 and PspABIP659 all reduce the transcription
of glutamate, aspartate and branched chain amino acids transporters. PspABIP659 further
reacts in a similar fashion to the presence of arginine, lysine and histidine (Table 2 &
Supplementary Table S2). The latter is also eliciting a response in PkM130. Upregulation of
amino acid transporters was only observed for methionine in PkM130 and glycine in
BoABIP444 (BCEN4_v1_340003).

Interestingly, an operon coding for a sarcosine oxidase (*soxBDAG*), a serine deaminase (*sdaB*)
and an aldolase (*folB*) is up-regulated in all *Paraburkholderia* strains and in BoABIP444 but

not in the remaining *Burkholderia* despite them bearing homologues of the concerned genes
(Table 2 & Supplementary Table S2). Sarcosine is an intermediate in glycine synthesis which
is generated by the sarcosine oxidase with hydrogen peroxide as a byproduct. The serine
deaminase converts serine but also threonine into pyruvate and thus participates in
gluconeogenesis. The FolB aldolase is an enzyme involved in one step of tetrahydrofolate
synthesis, a cofactor used by the sarcosine oxidase.

Exudates have a limited impact on chemotaxis and motility in our experimental design
Flagellar synthesis was not significantly impacted in our experiment. The chemotaxis
pathway transducing elements *cheA*, *cheY* and *cheB* were equally devoid of regulation in all
strains except *Psp*ABIP659 where *cheA*, *cheB* but also *cheW*, *cheD* and *motB* are negatively
regulated (Table 2 & Supplementary Table S2). In other strains, chemotaxis elements present
a more scattered regulation. The steadiness of genes involved in cell motility might be
explained by their already high base level expression. The base mean of the flagellin
component *fliC* reaches up to hundreds of thousands of reads depending on the strain. While,
motility is usually activated upon root exudate sensing (Klonowska et al., 2018), in the
present setting bacteria may have had already reached their induction maxima.

Root exudate perception induces a shift in glycolysis strategies
The strains display numerous and specific reactions to the presence of carbohydrates from
RE. Several strains positively regulate their dedicated metabolism to the presence of xylulose,
ribose and glucarate (Table 2 & Supplementary Table S2).

A more conserved metabolic shift was however observed for glycolysis processes. In all
strains except BrABIP441, genes involved in gluconate import and metabolism are up
regulated. These enzymes are part of the Entner-Doudoroff (ED) pathway, an alternative to
EMP glycolysis with similar costs and outcomes (Figure 7A). We previously demonstrated
that ED pathway mutants of PkM130 and BvLMG10929 were significantly affected in their
capacity to colonize rice roots (Wallner et al., 2022). The upregulation of the ED pathway by
PkM130 and BvLMG10929 in the present study underlines its involvement in plant
colonization as one of the early metabolic adaptations. Genes, downstream of the Entner-
Doudoroff pathway, coding for a sugar ABC transporter are up-regulated in every strain
except *Psp*ABIP659 where they are repressed.

The perception of putrescine induces strain-specific adaptations

We paid particular attention to genetic elements that could indicate the onset of biofilm production as these structures are often involved in surface colonization. We noticed that RE were impacting the expression of genetic pathway involved in putrescine metabolism and cyclic diguanylate monophosphate (c-di-GMP) cycling. Upon stimulation with RE, we observe three different types of reactions to putrescine. PkM130, PspABIP659 and BvLMG10929 have a conserved response consisting in the downregulation of the previously mentioned genes. In PkM130 and PspABIP659, genes involved in putrescine formation from the precursor agmatine are also downregulated. Finally, BoABIP444 displays a fragmented regulation of the putrescine metabolism pathway but with systematic up-regulation of the respective genes (Table 2 & Figure 7B). The remaining strains, while they possess the according genes, do not adapt the pathways regulating the levels of intracellular putrescine.

Cyclic di-GMP signaling is only activated in *Paraburkholderia* species

Intracellular c-di-GMP levels are regulated by diguanylate cyclases (DGC) for synthesis and diguanylate phosphodiesterases (PDE) for degradation. *Burkholderia* strains do not differentially express any of their genes involved in c-di-GMP signaling. This is coherent with what was previously observed through a Tn-seq approach in BvLMG10929 where no gene involved in c-di-GMP cycling was required for rice colonization (Wallner et al., 2022). For *Paraburkholderia* strains, PkM130 shows the most moderate response with the downregulation of a single DGC. PsABIP630 downregulates one DGC and upregulates one PDE. The strongest response is observed for PspABIP659 which differentially expresses eleven genes involved in c-di-GMP cycling. The majority of DGC are upregulated and the only two regulated PDE are repressed. Hence, there are different strategies at play for *Paraburkholderia* strains upon sensing of RE concerning the adaptation of their c-di-GMP signaling. PkM130 and PspABIP659 elevate the intra-cellular levels of the secondary messenger while PsABIP630 decreases them. Again, these results correlate with previous observations for PkM130 where a mutant involved in c-di-GMP cycling had a severe defect in rice colonization (Wallner et al., 2022).

Discussion

In this study we analyzed the transcriptomic response to RE of six strains belonging to *Paraburkholderia* and *Burkholderia*. Given the ecologic background of both taxa we hypothesized that genera-dependent adaptation patterns could appear in the transcriptomic regulation upon sensing of rice RE. *Burkholderia* and *Paraburkholderia* strains share a close

taxonomic proximity and thus a substantial level of orthologous genes. However, their strain-
specific genomes are also of considerable size which is a premonitory sign of the disparate
transcriptomic response displayed by the different strains upon chemical stimulation with RE.
No categorizing trend could be inferred by the amount and intensity of regulated genes or by
the function of regulated genes. Each strain displays a dominantly individual response and the
genetic proximity between strains within a genus is not reflected in their transcriptomic
response. Thus, in the present cases, the adaption to the habitat is independent from the
genetic background.

Care was taken to expose the different bacteria to RE for a duration that was relative to their
respective metabolic rate. The purpose was to obtain a synchronized response for reliable
comparison. It is however conceivable that the six strains differently prioritize their responses
which could result in an overall similar strategy that cannot be detected if sampled at a single
time point. To avoid such biases induced by temporal shifts, one would have to produce a
transcriptomic kinetic. However, given the extent of differences observed here, we feel
confident that there is indeed a strain-specific response to RE.

Putrescine represents a long-overlooked metabolite in plant-bacteria interactions. While it has
been identified as a virulence factor in *Ralstonia solanacearum* (Shi et al., 2019) it is also
involved in the signalization between plant beneficial bacteria and their host (Liu et al., 2018).
Putrescine is recognized as an important signaling molecule in plants where it is involved in
various situations from organ development to biotic stress resilience (Tiburcio et al., 2014).
Its presence in RE has been reported for tomato and rice (Kuiper et al., 2001; Suzuki et al.,
2009) but has otherwise rarely been investigated. A study on the fitness determinants of
Pseudomonas sp. WCS365 in the rhizosphere of wild type *Arabidopsis thaliana* and
immunity impaired mutants demonstrated that the absence of putrescine catabolism genes
(*spuC, pauC* and *gabT*) were highly increasing the bacteria's fitness in the rhizosphere of WT
but not immunocompromised plants (Liu et al., 2018). BvLMG10929 transposon mutants
affected in putrescine metabolism and uptake were also detected to be less competitive during
rice root colonization (Wallner et al., 2022). It is suggested that putrescine might act as a
signaling molecule, informing the bacteria of the presence of a eukaryotic host. Responsive
bacteria in turn promote biofilm formation which favors attachment to the root. This
regulatory system must be extremely fine-tuned as to not trigger plant defenses by an excess
production of biofilms and other physiological changes subsequent to putrescine signalization.
In the transposon sequencing assay on *Pseudomonas* sp. WCS365, genes involved in

putrescine uptake and synthesis were found to be negative fitness regulators (Liu et al., 2018).
Consistently, PkM130, PspABIP659 and BvLMG10929 downregulate the according gene
homologues upon exudate sensing. Thus, we could hypothesize that these strains take
advantage of putrescine to regulate their metabolism and physiology towards a controlled
surface colonization of roots.

C-di-GMP is another signal molecule positively regulating biofilm formation when
accumulated (Romling et al., 2013). The regulation of enzymes involved in its metabolism
can therefore be key for a successful colonization of plant roots and evasion of host defenses.
We previously demonstrated the involvement of a c-di-GMP cycling enzyme in the root
colonization process of rice by PkM130 (Wallner et al., 2022). Inversely, in BvLMG10929
several genes involved in c-di-GMP cycling were identified as being obstructive to a
successful root colonization (Wallner et al., 2022). Recently, the importance of c-di-GMP in
biofilm production and subsequent virulence on rice was demonstrated for *Burkholderia*
glumae BGR1 (Kwak et al., 2020). The human opportunist *B. cenocepacia*, which is closely
related to *B. orbicola,* received much attention for the role played by c-di-GMP signalization
in virulence by regulation of biofilm formation and motility (Richter et al., 2019). Similarly,
the absence of the PDE-receptor coding gene *rpfR* in *Burkholderia lata* SK875, reduces the
strain's virulence and increase its biofilm production (Jung et al., 2017). RpfR is a receptor
for the *Burkholderia* specific quorum sensing signal BDSF which implies a connection
between c-di-GMP and cell-to-cell signaling (Schmid et al., 2017). Further studies linked the
production of biofilms with the sensing of nitrate and temperature which are relying on a
subsequent c-di-GMP dependent signaling pathway in *B. pseudomallei* (Mangalea et al.,
2017; Plumley et al., 2017)*.* The maize endophyte *Burkholderia gladioli* 3A12 displays c-di-
GMP dependent signaling for biocontrol purposes against fungal pathogens (Shehata and
Raizada, 2017). Although root exudate stimulation had no direct impact on the regulation of
flagellum-related genes, the regulation of genes involved in putrescine uptake and metabolism
as well as those involved in c-di-GMP signaling indicate an indirect and intricate regulation of
motility (Liu et al., 2018; Richter et al., 2019). This is coherent with the supposition that
bacteria evade host defenses by reducing the production of the common flagellin MAMPs.
This behavior was described upon exudate stimulation in *Pseudomonas aeruginosa* PA01,
Azoarcus sp. BH72 and *P. kururiensis* M130 (Mark et al., 2005; Shidore et al., 2012;
Coutinho et al., 2015).

Until recently, the ED pathway had never been reported for its activation at any stage of plant
colonization. The difference with the more classic EMP-glycolysis pathway is the generation
of NAD(P)H cofactor during ED-glycolysis. This cofactor can be essential for enzymes that
play a decisive role in rhizospheric adaptation as it is the case for thioredoxins, involved in
oxidative stress tolerance (Cumming et al., 2004; Chavarría et al., 2013). The *edd* gene,
coding for the NAD(P)H-generating enzyme, is amongst the only genes that are differentially
regulated in all six strains. In a separate study, we identified *edd* as being critical for a
successful interaction of BvLMG10929 and PkM130 with rice roots as its absence led to a
significant decrease in the bacteria's colonization capacities (Wallner et al., 2022).

Using six different strains from the *Burkholderia* and *Paraburkholderia* genera, we could find
no discerning pattern in their early response to the perception of plants through RE,
suggesting that there is a strong intra-generic diversity. Thus, instead of analyzing two genera,
we rather have six distinct strains that independently underline the importance of certain
pathways in plant-bacteria interactions. While some, such as c-di-GMP signaling were already
recognized, the implication of putrescine and the ED-pathway should receive an increased
attention and especially in later steps of plant colonization.

Methods

Plant cultivation and exudate collection

Oryza sativa ssp. *japonica* cv. Nipponbare seeds were sterilized, germinated and cultivated in
a sterile, hydroponic system as described before (King et al., 2019). Five seedlings were
planted per system and transferred to a growth chamber (28°C; 16 h light; 8 h dark). After 3
days of growth, the hydroponic medium was collected and filtered using sterile Millipore
membranes to remove residual perlite sand. The resulting medium was frozen at -80°C and
lyophilized. Dry RE were resuspended in sterile distilled water at an 80x concentration and
stored at -20°C until use.

For plant growth measurements, rice seedlings were sterilized and germinated as before and
transferred to pots containing 400 mL of a 7/2/1 mix of sterile sand (0.6-1.6 mm)/
perlite/vermiculite. Five pots containing four rice seedlings were used per condition.

Bacterial strains and cultivation

Burkholderia reimsis ABIP441, *Paraburkholderia* sp. ABIP659 and *Paraburkholderia sabiae*
ABIP630 were isolated from healthy *Oryza sativa* subsp. *japonica* roots. ABIP441 and
ABIP630 were isolated from rice roots collected from a field in Cameroon and ABIP659 from

a field in Vietnam. Additionally, we used the well characterized rice growth promoting
species *Burkholderia vietnamiensis* LMG10929 and *Paraburkholderia kururiensis* M130 as
well as the recently proposed *Burkholderia orbicola* sp. nov. (previously named *B.*
servocepacia) ABIP444 (Wallner et al., 2019; Morales-Ruíz et al., 2022).

All bacterial strains were cultured as follows: Glycerol stocks (20%) of bacterial cells
conserved at -80°C were plated on Luria's low salt LB (LBm; Sigma-Aldrich) agar plates and
incubated for 72 h at 28°C. Single colonies were used to inoculate 10 mL of LBm broth in 50
mL Falcon tubes and incubated for various amounts of time allowing the different strains to
reach an $OD_{600}=1.0$. For DNA extraction purposes, cells were used at this stage.

Plant inoculation assays

For colonization assays, bacterial strains were grown overnight in liquid LBm as described
above. Bacterial density was then adjusted to 1.10^8 cfu.mL^{-1} with regard to the relative
amount of cells at $OD_{600}=1.0$ for each strain. In a sterile setting, 5-day old rice seedlings,
prepared as described above, were inoculated with 1 mL of the bacterial solution at the base
of each plant. Two systems (10 plants total) were prepared for each condition and grown for
14 days (28°C; 16 h light; 8 h dark). Five plants were randomly harvested per condition as
biological replicates. The entire root system was sampled and transferred to a screw cap tube
containing 1 mL of sterile water and a sterile ceramic bead. Roots were weighted and then
pulverized using a Fastprep-24 (MPbio) at 6 m.s^{-1} for 40 seconds. A serial dilution of the
resulting solution was spotted out on square LBm plates and incubated at 28°C for 48 hours.

For plant growth measurements overnight broth bacterial cultures in LBm were pelleted by
centrifugation, resuspended in distilled water and adjusted to $OD_{600}=1.0$. CFU measurements
at this OD are reported in the legend of supplementary Figure S1. Plants grown in a
sand/perlite/vermiculite mix were inoculated with 1 ml of bacterial suspension per plant, 5
days after seedling transfer to pots. Plants were grown in a growth chamber (80% humidity,
28°C; 16 h light; 8 h dark) for 50 days before sampling, and fertilized with a 50% Hoagland
solution, twice a week. The length of the longest leaf as well as the root's and leave's dry
weights were measured for each plant. Statistics were performed in R (packages rstatix,
multcompView, ggplot2) and differences were estimated with a pairwise Wilcoxon non-
parametric test.

Exudate stimulation of bacterial cells

For RNA extraction, cells were prepared the as detailed above but using VSG (Vincent Succinate Glutamate) growth medium (Vincent, 1970). The pre-culture was then used to inoculate 50 mL VSG in 250 mL Erlenmeyer flasks at a final OD_{600}=0.075. After 1 h of agitation at 150 rpm and 28°C, 800 µL of concentrated root exudate solution (62-fold) was added to obtain a final exudate concentration that was equivalent to that found in the hydroponic growth system. Cells were then grown until they reached mid-exponential phase. The volume required for 10^9 cells was collected and immediately transferred to a 15mL Falcon tube containing 95% ethanol and 5% phenol. Collected cells were centrifuged at 4000 rpm for 15 minutes. The supernatant was discarded and the pellet was used to extract total RNA using the RiboPure RNA purification kit for bacteria (ThermoFisher). After extraction, the sample were subjected to a TURBO DNase (ThermoFisher) treatment to remove any residual DNA. Extraction efficiency was qualified using the 2100 Bioanalyzer system with an RNA Nano Chip (Agilent Technologies).

Ribodepletion and RNA sequencing

Ribosomic RNA (rRNA) was removed from total RNA extracts following the guidelines of Roux et al., 2018. Briefly, we generated probes targeting 5S, 16S and 23S rRNA incorporating a biotin complex in 5' and locked nucleic acids (LNAs) for improved specificity towards rRNA of the burkholderia group (Supplementary Table S4). An *in silico* approach validated that these probes are universal to all *Burkholderia s.l.* species and conserve specificity in the neighboring taxa *Ralstonia* and *Cupriavidus.* 2 µL of probes (50 µM) were used in combination with reagents from the RiboMinus Transcriptome Isolation Kit, bacteria (ThermoFisher). We used 2 µg of total RNA and 71 µL of magnetic beads. Prior to use, magnetic beads were washed twice in nuclease free water and once in 2x SSC buffer and resuspended in a final volume of 31 µL hybridization buffer. Ribodepletion efficiency was validated using the 2100 Bioanalyzer system (Agilent Technologies) with an RNA Nano Chip (Table S1).

RNA sequencing was performed at the high throughout sequencing platform at I2BC (CNRS, Gif-sur-Yvette, France) using Nextseq500 (Illumina) technology and 75 sequencing cycles (single read, 75 bp). De-multiplexing, adapter trimming and quality controls were performed by the platform using bcl2fastq2-2.18.12, Cutadapt 1.15 and FastQC v0.11.5 respectively. Raw reads and biosamples are accessible under Bioproject PRJEB42574 / ERP126457 at the European Nucleotide Archive (https://www.ebi.ac.uk/ena).

448 Reads were mapped to the reference genomes using the CLC workbench and differential gene
449 expression was inferred using the DESeq2 package. Low-count genes were removed at a
450 threshold of 5 reads and an independent hypothesis weighting was applied. One replicate had
451 to be removed from the analysis for *B. reimsis* ABIP441 as the sequencing depth was
452 insufficient to allow robust analyses (Figure S1).

DNA extraction and genome sequencing

The genomic DNA of ABIP441, ABIP630 and ABIP659 was extracted using the standard
operating procedure of the JGI as described before (Wallner et al., 2019). A genomic library
with average insert size of 350 bp was prepared for sequencing using a TruSeq Nano DNA
library preparation kit (Illumina, Inc., San Diego, CA, USA). Paired-end 2 x 125 nt
sequencing was performed by the MGX platform (CNRS, Montpellier, France) using a HiSeq
2500 (Illumina), generating 19,127,584 raw read pairs for ABIP441, 12,198,848 for ABIP630
and 16,208,618 for ABIP659. Reads with overlapping sequence were assembled using CLC
Genomics Workbench version 7.04 resulting in 311 contigs ranging from 506 bp to 27.4 kbp
for ABIP441, 84 contigs ranging from 811 bp to 424.5 kbp for ABIP630 and 122 contigs
ranging from 1162 bp to 173.7 kbp for ABIP659.

The genome quality was assessed using BUSCO v3 for the detection of mandatory single-
copy orthologues (Simão et al., 2015). Genome completeness was estimated at 97.0%, 99.0%
and 97.8% for ABIP441, ABIP630 and ABIP659 respectively. Genbank accession numbers
for the three new genomes are GCA_914492715 (ABIP441), GCA_914484815 (ABIP630),
GCA_914484835 (ABIP659), under Bioproject PRJEB42536.

Comparative genomics procedures

The genomic similarity was assessed using the Genome clustering tool of the Microscope
platform https://mage.genoscope.cns.fr/microscope/ (Vallenet et al., 2020). It is based on
Mash, a software that computes distances between two genomes. From all the pairwise
distances of the genomes set, a tree is constructed dynamically using the neighbor-joining
javascript package. The clustering annotations are computed from all-pairs distances ≤ 0.06
(or 94% ANI), corresponding to the ANI standard to define a species group. The clustering
has been computed using the Louvain Community Detection. Average Nucleotide identities
were inferred using the ANI calculator tool (https://www.ezbiocloud.net/tools/ani).

Gene of interest presence/absence was inferred by homology screening using BLASTp. For
macromolecular systems, the automatic annotation tool of the MicroScope microbial genome

annotation and analysis platform (Vallenet et al., 2020) based on the MacSyFinder v.1.0.2 software was used. These predictions were curated for T3SS using advanced comparative genomics analyses (Wallner et al., 2021). For the T6SS, we used studies on BvLMG10929 and PkM130 (Nguyen et al., 2018; Dias et al., 2019) and inferred the diversity in the remaining strains through homology screening using BLASTp with core components ClpV, TssA, TssB and TssK as queries. The same procedure was adopted for the T4SS and the VirB4, VirB6 and the VirB11 ATPase core components.

Core genome compositions were calculated using the Phyloprofile exploration tool implemented in the MicroScope microbial genome annotation and analysis platform (Vallenet et al., 2020). Homology constraints were set at minLrap $\geq$ 0.8 and identity $\geq$ 50%. To establish a list of specific genes for comparative genomics purposes, exclusive comparisons were performed for the 63 possible combinations between the 6 strains where the genomes not used in the comparison were systematically use as negative filter at the same homology constraints as above. To establish a list of co-regulated genes between the six species, non-exclusive comparisons were performed for each pairwise combination as well as for the *Burkholderia* and *Paraburkholderia* trios and the combination of all six species.

Funding

The authors gratefully acknowledge financial support from the CGIAR research program (CRP) RICE as well as from the French national research agency (ANR) funding the BURKADAPT project (ANR-19-CE20-0007). AW and LG were supported by fellowships from the French Ministry of Higher Education, Research and Innovation.

References

Araújo WL, Creason AL, Mano ET, Camargo-Neves AA, Minami SN, Chang JH, Loper JE (2016) Genome Sequencing and Transposon Mutagenesis of Burkholderia seminalis TC3.4.2R3 Identify Genes Contributing to Suppression of Orchid Necrosis Caused by B. gladioli. Mol Plant Microbe Interact **29**: 435–46

Arriel-Elias MT, de Carvalho Barros Côrtes MV, de Sousa TP, Chaibub AA, de Filippi MCC (2019) Induction of resistance in rice plants using bioproducts produced from Burkholderia pyrrocinia BRM 32113. Environ Sci Pollut Res **26**: 19705–19718

Badri D V., Vivanco JM (2009) Regulation and function of root exudates. Plant, Cell Environ **32**: 666–681

Batista BD, Taniguti LM, Monteiro-Vitorello CB, Azevedo JL, Quecine MC (2016) Draft genome sequence of Burkholderia ambifaria RZ2MS16, a plant growth-promoting rhizobacterium isolated from guarana, a tropical plant. Genome Announc. doi: 10.1128/genomeA.00125-16

Chavarría M, Nikel PI, Pérez-Pantoja D, De Lorenzo V (2013) The Entner-Doudoroff pathway empowers Pseudomonas putidaKT2440 with a high tolerance to oxidative stress. Environ Microbiol **15**: 1772–1785

Chávez-Ramírez B, Kerber-Díaz JC, Acoltzi-Conde MC, Ibarra JA, Vásquez-Murrieta M-S, Estrada-de los Santos P (2020) Inhibition of Rhizoctonia solani RhCh-14 and Pythium ultimum PyFr-14 by Paenibacillus polymyxa NMA1017 and Burkholderia cenocepacia CACua-24: A proposal for biocontrol of phytopathogenic fungi. Microbiol Res **230**: 126347

Coutinho BG, Licastro D, Mendonça-Previato L, Cámara M, Venturi V (2015) Plant-Influenced gene expression in the rice endophyte Burkholderia kururiensis M130. Mol Plant-Microbe Interact **28**: 10–21

Cumming RC, Andon NL, Haynes PA, Park M, Fischer WH, Schubert D (2004) Protein disulfide bond formation in the cytoplasm during oxidative stress. J Biol Chem **279**: 21749–21758

Dall'Agnol RF, Bournaud C, de Faria ŚM, Béna G, Moulin L, Hungria M (2017) Genetic diversity of symbiotic Paraburkholderia species isolated from nodules of Mimosa pudica (L.) and Phaseolus vulgaris (L.) grown in soils of the Brazilian Atlantic Forest (Mata Atlântica). FEMS Microbiol Ecol. doi: 10.1093/femsec/fix027

Dias GM, de Sousa Pires A, Grilo VS, Castro MR, de Figueiredo Vilela L, Neves BC (2019) Comparative genomics of *Paraburkholderia kururiensis* and its potential in bioremediation, biofertilization, and biocontrol of plant pathogens. Microbiologyopen **8**: e00801

Engledow AS, Medrano EG, Mahenthiralingam E, LiPuma JJ, Gonzalez CF (2004) Involvement of a plasmid-encoded type IV secretion system in the plant tissue watersoaking phenotype of Burkholderia cenocepacia. J Bacteriol **186**: 6015–6024

Esmaeel Q, Issa A, Sanchez L, Clément C, Jacquard C, Barka EA (2018) Draft Genome Sequence of Burkholderia reimsis BE51, a Plant-Associated Bacterium Isolated from

Agricultural Rhizosphere . Microbiol Resour Announc. doi: 10.1128/mra.00978-18

Estrada-de los Santos P, Palmer M, Chávez-Ramírez B, Beukes C, Steenkamp E,
Briscoe L, Khan N, Maluk M, Lafos M, Humm E, et al (2018) Whole Genome
Analyses Suggests that Burkholderia sensu lato Contains Two Additional Novel Genera
(Mycetohabitans gen. nov., and Trinickia gen. nov.): Implications for the Evolution of
Diazotrophy and Nodulation in the Burkholderiaceae. Genes (Basel) **9**: 389

Fan B, Carvalhais LC, Becker A, Fedoseyenko D, Von Wirén N, Borriss R (2012)
Transcriptomic profiling of Bacillus amyloliquefaciens FZB42 in response to maize root
exudates. BMC Microbiol **12**: 116

Galyov EE, Brett PJ, DeShazer D (2010) Molecular Insights into Burkholderia
pseudomallei and Burkholderia mallei Pathogenesis . Annu Rev Microbiol **64**: 495–517

Govindarajan M, Balandreau J, Kwon SW, Weon HY, Lakshminarasimhan C (2008)
Effects of the inoculation of Burkholderia vietnamensis and related endophytic
diazotrophic bacteria on grain yield of rice. Microb Ecol **55**: 21–37

Hameed A, Shahina M, Lai WA, Stothard P, Young L Sen, Lin SY, Young CC (2020)
Draft genome sequence reveals co-occurrence of multiple antimicrobial resistance and
plant probiotic traits in rice root endophytic strain Burkholderia sp. LS-044 affiliated to
Burkholderia cepacia complex. J Glob Antimicrob Resist **20**: 28–30

Hardoim PR (2015) Heading to the Origins – Rice Microbiome as Functional Extension of
the Host. doi: 10.4172/2375-4338.1000133

Hassan S, Mathesius U (2012) The role of flavonoids in root-rhizosphere signalling:
opportunities and challenges for improving plant-microbe interactions. J Exp Bot **63**:
3429–3444

Ikeda S, Sasaki K, Okubo T, Yamashita A, Terasawa K, Bao Z, Liu D, Watanabe T,
Murase J, Asakawa S, et al (2014) Low nitrogen fertilization adapts rice root
Microbiome to low nutrient environment by changing biogeochemical functions.
Microbes Environ **29**: 50–59

Jacobs JL, Fasi AC, Ramette A, Smith JJ, Hammerschmidt R, Sundin GW (2008)
Identification and onion pathogenicity of Burkholderia cepacia complex isolates from the
onion rhizosphere and onion field soil. Appl Environ Microbiol **74**: 3121–3129

Jung HI, Kim YJ, Lee YJ, Lee HS, Lee JK, Kim SK (2017) Mutation of the cyclic di-GMP phosphodiesterase gene in Burkholderia lata SK875 attenuates virulence and enhances biofilm formation. J Microbiol **55**: 800–808

Kang S-M, Waqas M, Shahzad R, You Y-H, Asaf S, Khan MA, Lee K-E, Joo G-J, Kim S-J, Lee I-J (2017) Isolation and characterization of a novel silicate-solubilizing bacterial strain *Burkholderia eburnea* CS4-2 that promotes growth of japonica rice (*Oryza sativa* L. cv. Dongjin). Soil Sci Plant Nutr **63**: 1–9

Kim N, Mannaa M, Kim J, Ra JE, Kim SM, Lee C, Lee HH, Seo YS (2021) The in vitro and in planta interspecies interactions among rice-pathogenic Burkholderia species. Plant Dis **105**: 134–143

King E, Wallner A, Rimbault I, Barrachina C, Klonowska A, Moulin L, Czernic P (2019) Monitoring of rice transcriptional responses to contrasted colonizing patterns of phytobeneficial Burkholderia s.l. reveals a temporal shift in JA systemic response. Front Plant Sci **10**: 1141

Klonowska A, Melkonian R, Miché L, Tisseyre P, Moulin L (2018) Transcriptomic profiling of Burkholderia phymatum STM815, Cupriavidus taiwanensis LMG19424 and Rhizobium mesoamericanum STM3625 in response to Mimosa pudica root exudates illuminates the molecular basis of their nodulation competitiveness and symbiotic evolutionary history. BMC Genomics **19**: 1–22

Kuiper I, Bloemberg G V, Noreen S, Thomas-Oates JE, Lugtenberg BJJ (2001) Increased Uptake of Putrescine in the Rhizosphere Inhibits Competitive Root Colonization by Pseudomonas fluorescens Strain WCS365.

Kwak GY, Choi O, Goo E, Kang Y, Kim J, Hwang I (2020) Quorum Sensing-Independent Cellulase-Sensitive Pellicle Formation Is Critical for Colonization of Burkholderia glumae in Rice Plants. Front Microbiol. doi: 10.3389/fmicb.2019.03090

Liu Z, Beskrovnaya P, Melnyk RA, Hossain SS, Khorasani S, O'Sullivan LR, Wiesmann CL, Bush J, Richard JD, Haney CH (2018) A Genome-Wide Screen Identifies Genes in Rhizosphere-Associated Pseudomonas Required to Evade Plant Defenses. MBio. doi: 10.1128/mBio.00433-18

Mangalea MR, Plumley BA, Borlee BR (2017) Nitrate sensing and metabolism inhibit biofilm formation in the opportunistic pathogen Burkholderia pseudomallei by reducing

the intracellular concentration of c-di-GMP. Front Microbiol **8**: 1353

Mark GL, Dow JM, Kiely PD, Higgins H, Haynes J, Baysse C, Abbas A, Foley T, Franks
A, Morrissey J, et al (2005) Transcriptome profiling of bacterial responses to root
exudates identifies genes involved in microbe-plant interactions. Proc Natl Acad Sci U S
A **102**: 17454–17459

Mattos KA, Pádua VLM, Romeiro A, Hallack LF, Neves BC, Ulisses TMU, Barros CF,
Todeschini AR, Previato JO, Mendonça-Previato L (2008) Endophytic colonization
of rice (Oryza sativa L.) by the diazotrophic bacterium Burkholderia kururiensis and its
ability to enhance plant growth. An Acad Bras Cienc **80**: 477–493

Mitter B, Petric A, Shin MW, Chain PSG, Hauberg-Lotte L, Reinhold-Hurek B, Nowak
J, Sessitsch A (2013) Comparative genome analysis of Burkholderia phytofirmans PsJN
reveals a wide spectrum of endophytic lifestyles based on interaction strategies with host
plants. Front Plant Sci **4**: 120

Morales-Ruíz LM, Rodríguez-Cisneros M, Kerber-Díaz JC, Rojas-Rojas FU, Ibarra JA,
Estrada-de los Santos P (2022) Burkholderia orbicola sp. nov., a novel species within
the Burkholderia cepacia complex. Arch Microbiol **204**: 1–9

Mullins AJ, Murray JAH, Bull MJ, Jenner M, Jones C, Webster G, Green AE, Neill DR,
Connor TR, Parkhill J, et al (2019) Genome mining identifies cepacin as a plant-
protective metabolite of the biopesticidal bacterium Burkholderia ambifaria. Nat
Microbiol **4**: 996–1005

Navarro MOP, Piva ACM, Simionato AS, Spago FR, Modolon F, Emiliano J, Azul AM,
Chryssafidis AL, Andrade G (2019) Bioactive Compounds Produced by Biocontrol
Agents Driving Plant Health. Microbiome Plant Heal. Dis. Springer Singapore,
Singapore, pp 337–374

Nguyen TT, Lee HH, Park I, Seo YS (2018) Genome-wide analysis of type VI system
clusters and effectors in burkholderia species. Plant Pathol J **34**: 11–22

Perin L, Martínez-Aguilar L, Castro-González R, Estrada-de Los Santos P, Cabellos-
Avelar T, Guedes H V, Reis VM, Caballero-Mellado J (2006) Diazotrophic
burkholderia species associated with field-grown maize and sugarcane. Appl Environ
Microbiol **72**: 3103–10

Plumley BA, Martin KH, Borlee GI, Marlenee NL, Burtnick MN, Brett PJ, AuCoin DP,
Bowen RA, Schweizer HP, Borlee BR (2017) Thermoregulation of biofilm formation
in burkholderia pseudomallei is disrupted by mutation of a putative diguanylate cyclase.
J Bacteriol. doi: 10.1128/JB.00780-16

Richter AM, Fazli M, Schmid N, Shilling R, Suppiger A, Givskov M, Eberl L, Tolker-
Nielsen T (2019) Key players and individualists of Cyclic-di-GMP signaling in
burkholderia cenocepacia. Front Microbiol **10**: 3286

Romling U, Galperin MY, Gomelsky M (2013) Cyclic di-GMP: the First 25 Years of a
Universal Bacterial Second Messenger. Microbiol Mol Biol Rev **77**: 1–52

Schmid N, Suppiger A, Steiner E, Pessi G, Kaever V, Fazli M, Tolker-Nielsen T, Jenal
U, Eberl L (2017) High intracellular c-di-GMP levels antagonize quorum sensing and
virulence gene expression in Burkholderia cenocepacia H111. Microbiol (United
Kingdom) **163**: 754–764

Shehata HR, Raizada MN (2017) A Burkholderia endophyte of the ancient maize landrace
Chapalote utilizes c-di-GMP-dependent and independent signaling to suppress diverse
plant fungal pathogen targets. FEMS Microbiol Lett. doi: 10.1093/femsle/fnx138

Shi Z, Wang Q, Li Y, Liang Z, Xu L, Zhou J, Cui Z, Zhang LH (2019) Putrescine is an
intraspecies and interkingdom cell-cell communication signal modulating the virulence
of dickeya zeae. Front Microbiol. doi: 10.3389/fmicb.2019.01950

Shidore T, Dinse T, Öhrlein J, Becker A, Reinhold-Hurek B (2012) Transcriptomic
analysis of responses to exudates reveal genes required for rhizosphere competence of
the endophyte Azoarcus sp. strain BH72. Environ Microbiol **14**: 2775–2787

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva E V., Zdobnov EM (2015)
BUSCO: Assessing genome assembly and annotation completeness with single-copy
orthologs. Bioinformatics **31**: 3210–3212

Song D, Chen G, Liu S, Khaskheli MA, Wu L (2019) Complete genome sequence of
Burkholderia sp. JP2-270, a rhizosphere isolate of rice with antifungal activity against
Rhizoctonia solani. Microb Pathog **127**: 1–6

Sousa SA, Ramos CG, Leitão JH (2011) Burkholderia cepacia Complex: Emerging
Multihost Pathogens Equipped with a Wide Range of Virulence Factors and

Determinants. Int J Microbiol. doi: 10.1155/2011/607575

de Souza R, Meyer J, Schoenfeld R, da Costa PB, Passaglia LMP (2015) Characterization
of plant growth-promoting bacteria associated with rice cropped in iron-stressed soils.
Ann Microbiol **65**: 951–964

Suzuki K, Okazaki K, Tawaraya K, Osaki M, Shinano T (2009) Gas chromatography–
mass spectrometry associated global analysis of rice root exudates under aseptical
conditions. Soil Sci Plant Nutr **55**: 505–513

Tiburcio AF, Altabella T, Bitrián M, Alcázar R (2014) The roles of polyamines during the
lifespan of plants: From development to stress. Planta **240**: 1–18

Trân Van V, Berge O, Ngô Kê S, Balandreau J, Heulin T (2000) Repeated beneficial
effects of rice inoculation with a strain of Burkholderia vietnamiensison early and late
yield components in low fertility sulphate acid soils of Vietnam. Plant Soil **218/2**: 273–
284

Vallenet D, Calteau A, Dubois M, Amours P, Bazin A, Beuvin M, Burlot L, Bussell X,
Fouteau S, Gautreau G, et al (2020) MicroScope: An integrated platform for the
annotation and exploration of microbial gene functions through genomic, pangenomic
and metabolic comparative analysis. Nucleic Acids Res **48**: D579–D589

Vidal-Quist JC, O'Sullivan LA, Desert A, Fivian-Hughes AS, Millet C, Jones TH,
Weightman AJ, Rogers HJ, Berry C, Mahenthiralingam E (2014) Arabidopsis
thaliana and Pisum sativum models demonstrate that root colonization is an intrinsic trait
of Burkholderia cepacia complex bacteria. Microbiology **160**: 373–384

Vincent JM (1970) A manual for the practical study of the root-nodule bacteria. A Man.
Pract. study root-nodule Bact.

Wallner A, Busset N, Lachat J, Guigard L, King E, Rimbault I, Mergaert P, Béna G,
Moulin L (2022) Differential Genetic Strategies of Burkholderia vietnamiensis and
Paraburkholderia kururiensis for Root Colonization of Oryza sativa subsp. japonica and
O. sativa subsp. indica, as Revealed by Transposon Mutagenesis Sequencing. Appl
Environ Microbiol. doi: 10.1128/AEM.00642-22

Wallner A, King E, Ngonkeu ELM, Moulin L, Béna G (2019) Genomic analyses of
Burkholderia cenocepacia reveal multiple species with differential host-Adaptation to

plants and humans. BMC Genomics **20**: 803

Wallner A, Moulin L, Busset N, Rimbault I, Béna G (2021) Genetic Diversity of Type 3
Secretion System in Burkholderia s.l. and Links With Plant Host Adaptation. Front
Microbiol **0**: 3178

Wang M, Tachibana S, Murai Y, Li L, Lau SYL, Cao M, Zhu G, Hashimoto M,
Hashidoko Y (2016a) Indole-3-Acetic Acid Produced by Burkholderia heleia Acts as a
Phenylacetic Acid Antagonist to Disrupt Tropolone Biosynthesis in Burkholderia
plantarii. Sci Rep. doi: 10.1038/srep22596

Wang XQ, Liu AX, Guerrero A, Liu J, Yu XQ, Deng P, Ma L, Baird SM, Smith L, Li
XD, et al (2016b) Occidiofungin is an important component responsible for the
antifungal activity of Burkholderia pyrrocinia strain Lyc2. J Appl Microbiol **120**: 607–18

Yi Y, de Jong A, Frenzel E, Kuipers OP (2017) Comparative Transcriptomics of Bacillus
mycoides Strains in Response to Potato-Root Exudates Reveals Different Genetic
Adaptation of Endophytic and Soil Isolates. Front Microbiol **8**: 1487

Yu Y, Zhang J, Petropoulos E, Baluja MQ, Zhu C, Zhu J, Lin X, Feng Y (2018)
Divergent Responses of the Diazotrophic Microbiome to Elevated CO2 in Two Rice
Cultivars. Front Microbiol **9**: 1139

Zhang N, Yang D, Wang D, Miao Y, Shao J, Zhou X, Xu Z, Li Q, Feng H, Li S, et al
(2015) Whole transcriptomic analysis of the plant-beneficial rhizobacterium Bacillus
amyloliquefaciens SQR9 during enhanced biofilm formation regulated by maize root
exudates. BMC Genomics **16**: 685

Table 1. **List of the six strains used in this study & their genomic characteristics.**

	B. reimsis ABIP441	*B. orbicola* ABIP444	*B. vietnamiensis* LMG10929	*P. sabiae* ABIP630	*P. sp.* ABIP659	*P. kururiensis* M130
Genbank	GCA_914492715	GCA_905184115	GCA_000959445	GCA_914484815	GCA_914484835	GCA_000341045
Genome size (bp)	7845349	7409039	6930496	8354439	8573543	7129657
Contigs	311	292	3 Chr, 1 Pl	84	122	9
Genomic objects	8330	7822	6908	8239	8395	6524
CDSs	7881	7402	6416	7779	7934	6116
RNA genes	101	96	87	93	99	88
Phage-associated genes	61	79	10	14	52	33
GC%	66.59	66.72	65.07	62.48	64.95	65.02
Isolation						
Country	Cameroon	Cameroon	Vietnam	Cameroon	Vietnam	Brazil
Compartment	Rhizosphere	Rhizosphere	Rhizosphere	Rhizosphere	Rhizosphere	Endosphere

716   Abbreviations: Chr: chromosome, Pl: plasmid.

718 Table 2. **Differentially regulated gene categories in the six *Burkholderia s.l.* strains.**

Metabolic pathways	*P. kururiensis* M130		*P. sabiae* ABIP630		*P. sp.* ABIP659		*B. vietnamiensis* LMG10929		*B. reimsis* ABIP441		*B. orbicola* ABIP444	
	down	up	down	up	down	up	down	up	down	up	down	up
Sulfite reduction	-	6	1	-	-	1	-	-	-	-	3	-
Sulfate transport	-	5	-	-	-	2	-	-	1	-	3	-
Alkanesulfonate trans.	-	4	1	-	-	3	-	2	-	-	1	-
Taurine transport	-	2	-	-	3	-	-	-	1	-	1	-
Assimilatory nitrate red.	-	8	-	-	-	-	-	4	-	-	-	-
Respiratory nitrate red.	-	6	-	-	-	6	-	-	-	-	-	-
NRPS pathways	4	-	3	-	-	-	-	2	2	-	3	-
Iron uptake complex	5	-	5	-	-	2	-	1	5	-	-	-
Xylulose catabolism	-	2	-	-	-	2	-	-	1	-	-	-
Ribose import and metab.	-	-	-	-	5	-	-	-	-	3	-	3
Putative gluconate trans.	-	4	-	4	4	-	-	4	-	4	-	1
Glucarate metabolism	-	-	-	-	-	3	-	-	-	-	1	-
ED pathway	-	3	-	2	1	2	-	5	-	-	-	3
Glycine metabolism	-	4	-	4	-	5	1	-	-	-	-	5
Myo-inositol	-	-	-	-	-	9	-	1	-	-	1	-
Urease	1	-	-	-	3	-	1	1	-	-	-	1
CO dehydrogenase	2	3	-	-	-	5	-	1	-	-	-	-
Putrescine import	4	-	-	-	4	-	4	-	-	-	-	3
Putrescine metabolism	9	-	2	-	7	1	8	-	1	-	-	3
Pyochelin synthesis	-	-	-	-	-	-	-	-	5	-	5	-
TonB-dependent transport	3	-	3	-	-	3	-	3	1	-	3	-
T6SS	1	-	-	-	-	-	-	-	-	-	-	-
Tad pilus	1	-	1	1	-	-	8	-	-	2	-	1
c-di-GMP cycling	1	-	1	1	5	6	-	-	2	1	1	2
Chemotaxis	2	3	2	-	15	1	-	-	2	1	1	2
Quorum sensing	-	1	1	-	-	-	1	-	-	-	-	-
Outer membrane porin	-	1	-	1	1	-	-	1	-	1	-	1

Figure legends:

Figure 1. Genome clustering of studied strains (in bold) with their closest species neighbors. The tree was constructed using the Genome clustering tool of the Microscope plateform (https://mage.genoscope.cns.fr/microscope/). Genbank genome accession numbers of strains under study are indicated in Table 1, and others are: *B. reimsis* BE51 (GCA_003294055) ; *B. cepacia* ATCC25416 (GCA_006094315) ; *B. cenocepacia* J2315 (GCA_000009485) ; *P. sabiae* LMG24235 (GCA_904848645) ; *P. phymatum* STM815 (GCA_000020045), *P. tropica* LMG22274 (GCA_902833865), *Ralstonia solanacearum* GMI1000 (GCA_000009125).

Figure 2. Plant colonization efficiencies for six *Burkholderia* and *Paraburkholderia* strains. Nipponbare rice seedlings were inoculated at 5 days post-germination with 1.10^8 bacterial cells and grown in a hydroponic medium. Total roots were sampled at different time points (5 replicates each) and the attached bacterial population was enumerated. The trendlines (red) were inferred using the LOESS method.

Figure 3. UpSet plot of gene conservation between six Burkholderia s.l. strains. Single dots are representative of the specific genome of each strain. Linked dots represent the conserved genes among the connected strains. All combinations totaling less than 50 conserved genes are not shown. Gene duplicates were excluded from this analysis.

Figure 4. Transcriptomic response of *Burkholderia* and *Paraburkholderia* strains to RE. Genes that underwent a 1.5-fold and higher positive or negative regulation at a confidence level of adj. pval ≤ 0.05 were considered to be significantly regulated upon stimulation with RE. The number of genes passing that threshold (A) and their regulation intensity on a log2 scale (B) are represented.

Figure 5. Specific and conserved regulation between strains upon RE stimulation. We calculated the conserved genome between different groups of strains to determine which genes are commonly regulated (either positively (red), negatively (blue) or differentially (grey) between each pair of strains, the *Burkholderia* and *Paraburkholderia* trios, all 6 strains and all 6 except BrABIP441. Single dots represent the regulation of strain-specific genes. For the PspABIP659 specific response, the histogram was reduced to fit with the figure (total regulated genes = 561, with a similar number of up and down genes)

Figure 6. COG classification of the transcriptomic response of *Burkholderia* and *Paraburkholderia* strains to RE. The number of genes falling in each COG category is expresses as a percentage of the total amount of genes regulated in the corresponding strains (Figure 4A).

Figure 7. **Regulation of Entner-doudoroff and putrescine pathways upon root exudates stimulation.** For each strain and pathway, the fold-change value is depicted next to the corresponding gene. (A) The Entner-Doudoroff pathway uses glucose as starting product and branches into the glycolysis pathway with glyceraldehyde-3P which is ultimately converted to pyruvate. No significant regulation was found in *B. reimsis* ABIP441. (B) Putrescine synthesis, catabolism and uptake pathways in *Burkholderia s.l.* strains. No significant regulation was found in *B. reimsis* ABIP441 and *P. sabiae* ABIP630.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

A

B

Fig. 5

Fig. 6

Fig. 7

A. Entner-doudoroff pathway

B. Putrescine pathway