Intracellular FUS protein accumulation leads to cytoskeletal, organelle and cellular homeostasis perturbations

Chyi Wei Chung[a], Alexandra J. Zhou[b,c], Ioanna Mela[a], Amberley D. Stephens[a], Akinori Miyashita[d,1], Peter H. St George-Hyslop[d,e,f], Clemens F. Kaminski[a], Tuomas P. J. Knowles[b,c], Gabriele S. Kaminski Schierle[a*]

[a] Department of Chemical Engineering and Biotechnology, University of Cambridge, CB3 0AS, United Kingdom.
[b] Centre for Misfolding Diseases, Yusuf Hamied Department of Chemistry, University of Cambridge, CB2 1EW, United Kingdom.
[c] Cavendish Laboratory, Department of Physics, University of Cambridge, CB3 0HE, United Kingdom.
[d] Department of Medicine (Neurology), Temerty Faculty of Medicine, University of Toronto, and University Health Network, 60 Leonard Avenue, Toronto, ON M5T 0S8, Canada.
[e] Cambridge Institute for Medical Research and Department of Clinical Neurosciences, School of Clinical Medicine, Addenbrookes Biomedical Campus, University of Cambridge, CB2 0XY, United Kingdom.
[f] Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Department of Neurology, Columbia University Irvine Medical Center, 630 West 168th Street, New York, NY, USA 10032.

[1] Present address: Department of Molecular Genetics, Brain Research Institute, Niigata University, Niigata, Japan.

*Corresponding author: gsk20@cam.ac.uk

Summary

The molecular mechanisms that connect the formation of aberrant cytoplasmic FUS condensates to biological malfunction are incompletely understood. Here, we develop an approach to determine the intracellular FUS viscosity in live mammalian cells and find that ALS-related mutant P525L-FUS forms the most viscous condensates and has impaired cytoskeletal mechanoproperties and increased euchromatin formation. We further show that some of the main cellular organelles, e.g., actin/tubulin, lysosomes, mitochondria, the endoplasmic reticulum, are significantly functionally/structurally impaired in the presence of FUS. These may be related to defects in the tubulin network, as the latter facilitates transport, formation, fusion and fission of organelles. We observe significant increases in lysosomal biogenesis, size and pH; moreover, intracellular FUS accumulation significantly promotes cytoplasmic-to-nuclear translocation of TFEB, i.e., the master gene for inducing autophagy. However, despite these, increased autophagy needed for protein aggregate clearance is not observed to occur. Our study reveals that the formation of highly viscous FUS condensates significantly impacts cytoskeletal/organelle function and cellular homeostasis, which are closely associated with cell ageing. This raises the intriguing question as to whether mutant FUS activates similar cell processes as those during cellular senescence.

Keywords: fused-in sarcoma, amyotrophic lateral sclerosis, frontotemporal dementia, lysosomal de-acidification, TFEB, cytoskeletal defects, organelle dysfunction, autophagy-lysosomal pathway.

Introduction

The aberrant condensation of fused-in sarcoma (FUS) is a hallmark of both amyotrophic lateral sclerosis (ALS) and frontotemporal dementia (FTD). Under physiological conditions, FUS undergoes liquid-liquid phase separation (LLPS) and exists in a liquid and/or a dense condensate state predominantly within cell nuclei. Prominent cytoplasmic mislocalisation and the formation of more viscous, and in many cases, gelled condensates or even fibrillar aggregates, are observed upon missense and truncation mutations. Many, but not all of these ALS-associated mutations occur within the FUS nuclear localisation signal (NLS) (Bosco et al., 2010; Conte et al., 2012; Waibel et al., 2010). Pathological phase separation and the formation of gelled condensates have also been linked to arginine hypomethylation of FUS, as commonly observed in cases of sporadic FTD, which leads to disruptions in cytoplasmic ribonucleoprotein (RNP) granule function (e.g., stress granules (SGs) and neuronal transport granules) (Hofweber et al., 2018; Murakami et al., 2015; Qamar et al., 2018).

Whilst much attention has been given to the pathological role of FUS in the formation of cytoplasmic SGs, an understanding of how FUS condensation leads to disturbances in cellular homeostasis and function would be invaluable. This knowledge could also yield avenues for therapeutic intervention in diseases associated with the aberrant assembly of these proteins, such as FUS-associated ALS (fALS-FUS) and frontotemporal lobar degeneration (FTLD-FUS).

A potentially powerful clue to the latter may come from a detailed analysis of different fALS-FUS and FTLD-FUS models. We thus first established different FUS expressing cell lines and stress models in HEK293T cells before comparing the changes in intracellular FUS viscosity of the various FUS models. Combining single particle tracking (SPT-) and fluorescence lifetime imaging microscopy (FLIM), we develop a method to quantify intracellular FUS viscosity without the need for an external sensor/tracer, and that is higher-throughput than conventional fluorescence recovery after photobleaching (FRAP). To understand how FUS condensation leads to disturbances in cellular homeostasis, we then focused our analysis on the role of FUS in the nucleus and the cytoplasm,

including its effect on the cytoskeleton and on organelle positioning and function. We show that P525L-

FUS cells form the most viscous condensates, which affect the level of euchromatin formation,

cytoskeletal proteins and organelle positioning and function, including lysosomes, mitochondria and

the endoplasmic reticulum (ER). We show that the latter is linked to the observed lysosomal de-

acidification, which acts as a trigger for the nuclear to cytoplasmic translocation of transcription factor

EB (TFEB), i.e., the master modulator of autophagic genes. Instead of the expected upregulation of

macroautophagy (henceforth referred to as autophagy), however, autophagy is blocked at an early

stage, which promotes cellular malfunction and disease progression (Aman et al., 2021; Martini-Stoica

et al., 2016).

Results

Fluorescence lifetime imaging and single particle tracking form a tool to measure FUS condensate viscosity in live cells

Studying phase separation in live cells, and thus characterising intracellular FUS viscosity is challenging (Kuimova, 2012). We have previously shown, however, that protein phase separation and aggregation can be measured *in vitro* and in live cells using FLIM (Chen et al., 2017; Esbjörner et al., 2014). We thus first investigated whether SPT, which permits exact viscosity measurements *in vitro*, can be correlated with FLIM. We generated LLPS condensates by mixing equal amounts of recombinant GFP- and unlabelled WT-FUS protein. We then added the protein mixture into a solution containing 10% polyethylene glycol 35 (PEG-35) and 150 mM potassium chloride (KCl) to mimic the molecular crowding and ionic environments of the nuclear and cytoplasmic intracellular compartments. As we were interested in aggregation kinetics and changes in the micro-rheology of FUS with time, we performed ageing experiments involving SPT and FLIM over time, giving us a means to quantify both parameters of interest (**Figure 1A**). To probe the viscosity within FUS condensates, we introduced 40 nm fluorescent nanoparticles into the condensate mixture and tracked their trajectories. Because fluctuations seen in tracked nanoparticle trajectories are predominantly due to Brownian motion (Crocker and Grier, 1996); hence, we were able to calculate intra-condensate FUS viscosity from their mean square displacement (MSD) profiles. Between the 0 to 1h time points, FUS in the condensates becomes more viscous, with viscosity values increasing from 0.11±0.01 Pas to 0.44±0.08 Pas (**Figure 1B**, blue points). We observe a correlation between FLIM and SPT measurements (**Figure 1B**). During the same time period during which increased viscosity is observed, the GFP-tagged FUS molecules in the condensates cluster more closely, as indicated from the decrease in their fluorescence lifetimes from 2.23±0.01 ns to 2.08±0.01 ns (**Figure 1B**, red points; **Figure 1C**, Condensates). These values are in turn lower than that of dispersed WT-FUS before LLPS at 2.53 ns (**Figure 1B,** green point; **Figure 1C**, dispersed), which has a viscosity of 0.004 Pas from SPT measurements. To induce the fibrillation of FUS, we removed PEG-35 from the

condensate mixture described above to yield solid fibrils with a fluorescence lifetime in the range of 2.01±0.01 ns (**Figure 1B**, magenta point; **Figure 1C**, Fibrils). As the fluorescence lifetime of GFP is independent of the viscosity of its surrounding environment (Davidson et al., 2020), we note that FLIM and SPT yield independent and complementary parameters indicative of the level of FUS compaction and viscosity, respectively. Our data further show that beyond an hour, there are no further significant changes in either fluorescence lifetime or viscosity.

To observe both physiological and aberrant condensate formation of FUS in a cell model, we created HEK293T cells expressing GFP-labelled FUS (**Figure S1A**). To ensure that our observations were not due to the accumulation of GFP, we included cells expressing a GFP only construct (i.e., Control) in our experiments. We studied WT-FUS, and two ALS-related mutants, (i.e., P525L and R495X with a point site mutation at and truncation of the NLS, respectively), thereby leading to cytoplasmic mislocalisation of FUS(Gonzalez et al., 2021). Pathological phase separation of FUS is also promoted upon hypomethylation of its arginine-rich domain, yielding irreversible FUS hydrogels that are prevalent in FUS-related forms of sporadic FTD (Hofweber et al., 2018; Murakami et al., 2015). In order to induce FUS hypomethylation, we treated the cells with adenosine periodate oxidase (AdOx), an arginine methyltransferase inhibitor, which modulates the methylation state of the RNA-binding, arginine-rich domain of FUS (Fujii et al., 2016; Qamar et al., 2018). This promotes gelation and results in the formation of clusters of hypomethylated (HYPO) intranuclear aggregates (**Figure S1B**). P525L-FUS cells lose cytoplasmic mislocalisation, whereas R495X-FUS cells (which have a truncated NLS) still retain cytoplasmic condensates after AdOx treatment. Furthermore, we induced the formation of SGs using sodium arsenite (NaAsO$_2$) (**Figure S1C**). Existing as membraneless LLPS structures of cytoplasmic RNPs, SGs are transient structures of translationally stalled mRNA complexes and proteins (Sama et al., 2013). However, it has been suggested that in neurodegeneration models, they become persistent and are the site of aberrant aggregation (Wolozin and Ivanov, 2019). We validated the formation of SGs in our cell models by immunostaining of Ras-GTPase-activating protein binding protein (G3BP), a protein marker for the assembly and dynamics of SGs (**Figure S2**) (Yang et al., 2020). Agreeing with previous

reports (Baron et al., 2013; Bosco et al., 2010), we observe colocalisation of FUS and G3BP only in the

case of mutant variants of FUS (**Figure S2B**), despite the formation of SGs in all cell models.

ALS-associated FUS mutants form the most viscous aggregates in live cells

Using FLIM, we could quantify the aggregation propensity and extent of FUS in our cell models (**Figure**

1D&E). We observe that the NLS-mutants have more aggregated FUS (cell-averaged values of 2.48±0.12

ns and 2.49±0.12 ns for P525L- and R495X-FUS, respectively), in comparison to nuclear-localised WT-

FUS (2.62±0.07 ns) with fluorescence lifetimes that do not differ significantly from control cells at

2.64±0.06 ns. As FUS in SGs and hypomethylated intranuclear aggregates have higher fluorescence

intensities than other compartments of the cell (**Figure S1B&C**), we could segment them by setting a

fluorescence intensity threshold (**Figure 1D,** AdOx & NaAsO$_2$). In comparison to R495X-, P525L-FUS has

a lower fluorescence lifetime than hypomethylated intranuclear aggregates (1.78±0.05 ns *cf.* 2.01±0.04

ns in R495X) and SGs (2.35±0.06 ns *cf.* 2.46±0.02 ns in R495X). To calculate the intracellular viscosity of

FUS, we created a fluorescence lifetime to viscosity calibration based on the *in vitro* FUS condensate

system (**Figure S3A**), noting that GFP fluorescence lifetimes of recombinant FUS and the HEK293T FUS

models fall within the same range of 1.6—2.6 ns (**Figure 1E**). Hence, we perform linear fitting (**Figure**

S3A, black line) on SPT-FLIM data of the *in vitro* protein condensate system (**Figure S3A**, grey circles),

to map fluorescence lifetime values of FUS imaged in live HEK293T cells to their corresponding viscosity

values (**Figure 1E**, secondary axis). We note that before both AdOx and NaAsO$_2$ treatment, the two NLS

mutants of FUS are more viscous at ~0.10 Pas than WT-FUS at 0.05 Pas. Moreover, we estimate that

hypomethylated aggregates differ in their viscosity to SGs by a factor of 5—10, i.e., 0.69±0.05 (AdOx –

P525L) to 0.14±0.06 Pas (NaAsO$_2$ – P525L) and 0.47±0.04 (AdOx – R495X) to 0.04±0.02 Pas (NaAsO$_2$ –

R495X.

We observe a similar trend upon calculating the diffusion coefficient of these FUS macro-

assemblies inside cells (**Figure S3B**). For the latter, we extracted MSD profiles of the FUS macro-

assemblies' trajectories over time, which we imaged in a similar manner to SPT measurements for the

in vitro condensates. For both, AdOx and NaAsO$_2$ treatments, P525L-FUS cells yield HYPO nuclear

aggregates and SGs of lowest mobility, with respective diffusion coefficients of 0.036±0.004 nm^2 s^{-1}

(compared to 0.044±0.004 nm^2 s^{-1} in WT-FUS cells) and 0.011±0.002 nm^2 s^{-1} (compared to 0.025±0.001

nm^2 s^{-1} in R495X-FUS cells). As aforementioned, R495X-FUS is the only model that retains the

cytoplasmic distribution of FUS upon AdOx treatment due to its truncated NLS (**Figure S1B**). We observe

that the hypomethylated aggregates it forms are the most mobile with a diffusion coefficient

0.076±0.007 nm^2 s^{-1} (i.e., two-fold that of P525L- and WT-FUS). Hence, using both, our fluorescence

lifetime to viscosity calibration as well as diffusion coefficient measurements, we show that there are

differences in the way different FUS variants interact with their nuclear and cytoplasmic environment,

which affect their aggregation propensity.

Figure 1: Fluorescence lifetime to viscosity calibration reveals that ALS-associated P525L-FUS forms the most viscous condensates. (A) FLIM and SPT measurements were performed on *in vitro* FUS condensates, giving a measure of condensation state and viscosity, respectively. For SPT, 40 nm nanoparticles (denoted by white arrowhead) were mixed into the condensates and the mean square displacement (MSD) profiles of their trajectory were analysed. (B) Combined SPT-FLIM data show that there is a trend between GFP fluorescence lifetimes (τ_{GFP}, FLIM in red) and viscosity (μ, SPT in blue), where more aged FUS condensates are also more viscous. Values plateau ~1 hour and show no further change upon subsequent ageing. (C) Fluorescence lifetime maps for liquid-to-solid transition of recombinant FUS. A decrease in τ_{GFP} is used to quantify increasing condensation as dispersed FUS undergoes LLPS to form condensates, which are aged over a 4-hour period. Upon irreversible solid transition (i.e., formation of fibrils), fluorescence lifetime values are at the lowest values at 2.01±0.01 ns. Scale bars, 10 μm. (D) Fluorescence lifetime maps for FUS in the GFP HEK293T model, which show that fluorescence lifetimes fall in the same 1.5—2.6 ns range as the *in vitro* system in (C). The formation of hypomethylated (HYPO) intranuclear aggregates (first row, AdOx) and SGs (second row, NaAsO₂) were induced by the addition of different stressors. These macro-assemblies can

 be easily segmented by fluorescence intensity thresholding (Segmented FUS). Scale bars, 10 µm. (E) Cell averaged-

fluorescence lifetimes values (black) alongside viscosity predicted from correlative FLIM-SPT calibration on in vitro

condensates (B). ALS-related FUS mutant cells have lower τ_{GFP} (i.e., are at a more condensed state) in comparison to WT-FUS

and control cells. Fluorescence lifetimes of segmented hypomethylated nuclear aggregates (AdOx, red) and SGs (NaAsO$_2$,

orange) are also given, with P525L- forming more viscous macroassemblies compared to R495X-FUS. *In vitro* and HEK293T

FUS cell measurements were based on 3 individual protein preparations and 3 biological repeats respectively. One-way ANOVA

test (with Holm-Sidak's multiple comparison), where n.s. is not significant, *** p<0.001 and **** p<0.0001.

P525L-FUS mutant impairs cellular mechanoproperties and enhances euchromatin formation

We have thus far established that there are differences in the condensation states of FUS, with mutant

P525L- and R495X-FUS being in a more condensed state than WT-FUS, even if the latter has been

exposed to stressors such as AdOx and NaAsO$_2$. It has been suggested that loss of FUS functionality and

its cytoplasmic mislocalisation can cause loss or gain of function defects by affecting essential

cytoskeletal proteins (Theunissen et al., 2021). To further characterise the latter in our HEK293T cell

models, we visualise filamentous (f-)actin and microtubules (i.e., the 2 main proteins of the

cytoskeleton) using far-red fluorogenic Silicon Rhodamine (SiR) dyes (Lukinavičius et al., 2014) (**Figure**

2). As the SiR-dyes are of high specificity and only become fluorescent upon binding, their fluorescence

intensity can be used as to measure f-actin and microtubule polymerisation levels. Our data reveal that

only P525L- and not R495X-FUS has a significant impact on the cytoskeleton, as shown by reduction of

f-actin, as shown by a loss in SiR-Actin intensity. A similar effect on SiR-Actin intensity can only be

induced by stressors, such as AdOx and NaAsO$_2$, which cause hypomethylation or SG formation,

respectively (**Figure 2A&B**). However, in the case of P525L-FUS, the above stressors do not cause a

further defect on the cytoskeleton as P525L-FUS alone does. We next used SiR-Tubulin and performed

similar experiments as described above to see whether FUS condensates would also affect the

microtubule protein, tubulin. It is observed that P525L-FUS, followed by R495X-FUS, cells have the

lowest SiR-Tubulin intensity levels prior to drug treatments. Interestingly though, AdOx treatment,

which promotes nuclear translocation of FUS and the formation of hypomethylated clusters, reduces the burden on microtubules mainly present in the cytoplasmic region, the effect being strongest in P525L-FUS cells, where the largest drop in SiR-Tubulin intensity has occurred prior to the addition of AdOx (**Figure 2C&D**). The opposite effect is seen when cytoplasmic SGs are formed using NaAsO$_2$, which results in significantly reduced SiR-Tubulin intensities in all models. Additionally, we quantified the amounts of f-actin and β-tubulin (a sub-unit of microtubules) in the cell samples based on a Western Blot (**Figure S4**), and we find they do not differ significantly between FUS variants. Hence, this may indicate that depolymerisation of actin and microtubules has occurred, instead of lower concentrations of either being expressed.

quantification for SiR-Actin ($I_{SiR-Actin}$). f-actin levels in P525L-FUS cells are low even without the addition of any stressors. AdOx has a greater

effect in comparison to NaAsO$_2$ on reducing f-actin levels. (C) Fluorescence intensity composite images for GFP(-FUS) (green) and SiR-Tubulin

(magenta). Scale bars, 10 μm. (D) Cell-averaged fluorescence intensity quantification for SiR-Tubulin ($I_{SiR-Tubulin}$). Tubulin levels in P525L- and

R495X-FUS cells are impacted even without the addition of any stressors. The formation of HYPO aggregates improves microtubule levels, in

contrast to SGs which impair microtubules. HEK293T measurements were based on 3 biological repeats. One-way ANOVA test (with Holm-

Sidak's multiple comparison), where * for $p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$.

To confirm that FUS condensates indeed affect cellular mechanoproperties by interfering with

the two most important cytoskeletal proteins, actin and tubulin, we perform force-displacement (FD)-

atomic force microscopy (AFM) measurements on the cytoplasmic regions of live cells (**Figure S5**). In

comparison to WT-FUS cells, we have seen that mutant P525L-FUS cells display the lowest levels of

polymerisation for both cytoskeletal proteins investigated; moreover, R495X-FUS cells have lower levels

of SiR-Tubulin detected (**Figure 2**). Corroborating this, we find that P525L-FUS cells have the softest

cytoplasm with an average apparent Young's modulus (E_{App}) of 12.3±2.2 kPa, followed by R495X at

13.7±4.0 kPa, in comparison to WT-FUS at 25.5±2.2 kPa (**Figure S5A**); these measurements correspond

to intensity quantifications of SiR-dyes (**Figure 2**). Hence, we believe that the softening we observe is

due to destabilisation of the cytoskeleton i.e., a decrease in E_{App} corresponds to lower intensities of the

SiR-dyes (**Figure S5B**). It is interesting to note that the expression of a cytoplasmic protein, such as GFP

in the control cells, also has an effect on the cell mechanoproperties as measured by FD-AFM. However,

in contrast to FUS cells, which mainly leads to accumulation in the nucleus for WT-FUS and some

cytoplasmic condensates in P525L- and R495X-FUS cells, no effect on the cytoskeletal proteins is

observed in GFP control cells.

We have shown above that AdOx treatment has a significant effect on FUS condensate

formation, but AdOx is also known as a drug that reduces DNA methylation. We therefore investigated

the effect FUS condensates might have on the level of euchromatin formation, which is strongly

increased by hypomethylation. We have recently shown that the dye SiR-DNA (or SiR-Hoechst) can be

used to determine the level of eu/heterochromatin formation in live cells (Hockings et al., 2020; Novo

et al., 2022). As SiR-DNA intercalates with DNA structures, its fluorescence lifetime is significantly

affected by the level of chromatin condensation. Since euchromatin formation is associated with

significant DNA decondensation, an increase in the fluorescence lifetime of SiR-DNA can be associated

with an increase in euchromatin formation (Hockings et al., 2020; Novo et al., 2022). We observe that

both FUS mutant cells contain more euchromatin (i.e., more decondensed DNA), with P525L-FUS cells

(3.44±0.03 ns) experiencing a greater extent of DNA decondensation than R495X-FUS cells (3.43±0.04

ns) and in comparison to the GFP control (3.40±0.03 ns) and WT-FUS cells (3.41±0.03 ns) (**Figure 3C&D**).

To further control our results, we applied two different drugs known to hinder DNA compaction, i.e.,

AdOx (due its effect on DNA hypomethylation (Schwerk and Schulze-Osthoff, 2005)) and Trichostatin A

(TSA, which induces histone diacetylase (HDAC) inhibition (Moreira et al., 2003; Vigushin et al., 2001))

to control cells, which also raised the fluorescence lifetime of SiR-DNA to 3.44±0.04 and 3.43±0.03 ns,

respectively. It is interesting to note that the effect aberrant FUS condensates has on euchromatin

formation is as severe as the one induced by drugs, such as AdOx and TSA. Taken together, cytoskeletal

defects as well as changes in euchromatin state are most prevalent in mutant variants of FUS, especially

P525L-FUS.

Figure 3: ALS-related FUS mutants lead to an increase in euchromatin formation. (A) Fluorescence lifetime of SiR-DNA is used as a measure of DNA decondensation, which is related to heterochromatin formation. Mutant FUS cells exhibit lower level of heterochromatin formation (i.e., higher $\tau_{SiR-DNA}$). GFP control cells treated with AdOx and TSA which cause DNA hypomethylation and HDAC inhibition, were included. (B) Fluorescence lifetime maps. Scale bars, 10 μm. (C) Cartoon illustration of impaired mechanoproperties of P525L-FUS cells due to weakened cytoskeleton and DNA damage in comparison to WT-FUS cells. Created on Biorender.com. HEK293T measurements were based on 3 biological repeats. One-way ANOVA test (with Holm-Sidak's multiple comparison), where * for p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001.

Aberrant FUS aggregation impacts normal organelle function

The above studies have shown that mutant FUS has a significant effect on the cytoskeleton, which, in particular for tubulin. We thus investigated whether this effect on the microtubules might interfere with organelles, such as lysosomes, mitochondria, and ER, as the microtubules are important for the positioning, formation, fission, and fusion of the latter (Mattenberger et al., 2003; Pu et al., 2016). We applied super-resolution 2-colour structured illumination microscopy (SIM) with a resolution of 100 nm (i.e., below the diffraction limit) (Young et al., 2016) in COS7 cells, which have a flat morphology ideal

for SIM, as the technique is sensitive to out-of-focus glare associated with thicker samples (Ma et al.,

2021). We show that clustering of lysosomes and mitochondria at the perinuclear area of cells is greater

in P525L-, compared to WT-FUS and control cells (**Figure S6**). We have recently found that lysosomes

are responsible for the maintenance of the tubular ER network in the periphery of cells (Lu et al., 2022,

2020). Thus, clustering of lysosomes in the perinuclear area may significantly impair the tubular ER

network in the cell periphery. Hence, to investigate if a collapse in the tubular ER structure also follows

upon intracellular FUS accumulation, we co-expressed an ER marker, mEmerald-sec61β (Nixon-Abell et

al., 2016) in mCherry (mChe) versions of our control and FUS cells. Synthetic, hypomethylation-

mimicking variants which have increased numbers of arginine (i.e., 9R and 16R) were additionally

included as further controls (Qamar et al., 2018). We observe that there is a significant impact on the

tubular ER network as indicated by a decrease in the ER tubular to sheet ratio compared to the control

upon FUS accumulation (**Figure S7A&B**). Furthermore, we also observe a decline in mitochondrial

eccentricity (i.e., more rounded structures) in the presence of FUS accumulation, and particularly in

P525L-FUS when compared to the control (**Figure S7D&E**), which may indicate defects in mitochondrial

fission and increased mitochondrial stress.

Since lysosomes also play an important role not only the formation of the ER tubular network

but also in protein homeostasis, we analysed their structure and function in the different FUS cells in

more detail. To visualise the nanoscale structures of lysosomes in live cells, we again use SIM in mChe(-

FUS) COS7 cells (**Figure 4A**, green in Composite). We additionally labelled lysosomes with LysoTracker

Deep Red (**Figure 4A,** magenta in Composite), and segmented individual lysosomes from SIM images

using a custom-written script (**Figure 4A,** where each lysosome is represented by a different colour). As

a positive control, we treated mChe only control cells with chloroquine diphosphate (CQ). As an

autophagy inhibitor, CQ promotes the formation and accumulation of autophagosomes, as it disrupts

autophagy by preventing lysosome-autophagosome fusion(Mauthe et al., 2018). We observe that

R495X (0.28±0.10 μm^2 and 71±12 counted lysosomes) and P525L-FUS (0.29±0.12 μm^2 and 62±9

counted lysosomes) have larger and more numerous lysosomes in comparison to WT- (0.12±0.04 μm^2

& 46±5 counted lysosomes), 9R- (0.15±0.05 μm^2 & 40±6 counted lysosomes) and 16R-FUS (0.14±0.03

μm^2 & 46±7 counted lysosomes), which in turn are larger than in the control (0.09±0.03 μm^2 and 23±3

counted lysosomes) (**Figure 4B&C**). As expected, the effect is most prominent in the CQ-added control

sample (0.61±0.17 μm^2 and 78±14 counted lysosomes).

To test, whether these lysosomes are still able to maintain their low physiological pH, the latter

of which is essential to maintain protein homeostasis, we employ Dextran 10,000 MW tagged to Oregon

Green 488 (Dex-OG488), which has been used as a pH sensor in a fluorescence lifetime-based assay

(Burdikova et al., 2015). Dex-OG488 accumulates in acidic compartments including lysosomes within

cells (**Figure 4D**, Composite), hence can be used as an *in situ,* non-invasive lysosomal pH sensor. We

measured Dex-OG488 fluorescence lifetime ($\tau_{Dex-OG488}$) using FLIM and analysed resulting data using fit-

free phasor plot analysis. We see modest increases in $\tau_{Dex-OG488}$ for the nuclear-localised FUS variants,

i.e., 3.92±0.99%, 3.67±0.95% (9R) and 4.31±0.72% (16R), in comparison to control cells, which indicates

a rise in lysosomal pH. The differences become more significant for the NLS-mutants, i.e., 9.61±0.84%

(R495X) and 9.66±0.67% (P525L), as well as CQ-treated control and P525L-FUS (48.4±2.4% and

52.4±1.44%, respectively) (**Figure 4E&F**). This is unsurprising as CQ is known to accumulate within

lysosomes as a deprotonated weak base, and thereby increases lysosomal pH (Chen et al., 2011). In

summary, we see that the dilated lysosomes in ALS-associated P525L- and R495X-FUS have lost their

functionality due to greater lysosomal de-acidification.

Figure 4: ALS-associated FUS mutants increase lysosomal size and biogenesis, and impact lysosomal pH. (A—C) SIM measurements to visualise lysosomal structure using LysoTracker Deep Red staining. (A) Composite SIM images showing mChe(-FUS) in green and LysoTracker Deep Red in magenta (first column, Composite). Inset corresponds to region within the red box in the Composite image, where individually segmented lysosomes are represented by a different colour. Scale bar, 10 μm. (B) Averaged lysosome area and (C) averaged lysosome count, which show that NLS-mutants, P525L and R495X have more, and Control+CQ has significantly more dilated lysosomes, in comparison to the control and nuclear-localised FUS variants. (D—F) FLIM measurements on Dex-OG488 lysosomal pH sensor. (D) Composite fluorescence intensity images showing mChe(-FUS) in green and Dex-OG488 in magenta (first column, Composite). The latter is used as a non-invasive lysosomal pH sensor with fluorescence lifetime-based readouts shown as falsely coloured images (second column, $\tau_{Dex-OG488}$).

The presence of aberrant FUS aggregation promotes the nuclear translocation of TFEB but not autophagy

We have observed that greater lysosomal de-acidification leads to increased lysosomal biogenesis, both of which are enhanced in mutant P525L- and R495X-FUS. De-acidification of lysosomes may trigger autophagy via the nuclear translocation of TFEB, which is a transcriptional modulator for the autophagy-lysosomal pathway (ALP). The latter has been identified as a therapeutic target for diseases involving lysosomal dysfunction (Tan et al., 2021; Willett et al., 2017). Under physiological conditions, TFEB is phosphorylated and remains in the cytoplasm. Nuclear translocation and activation of the TFEB pathway are usually triggered under conditions of starvation and lysosomal stress, leading to autophagosome formation, and increased lysosomal biogenesis that upregulates autophagy (Settembre et al., 2011). Highly expressed in the central nervous system, the dysfunction of TFEB has been implicated in the pathogenesis of numerous neurodegenerative diseases including ALS/FTD (Chen et al., 2015). We thus addressed the role of the TFEB when it translocates from the cytoplasm to the nucleus. Upon co-transfection with GFP-TFEB into mChe(-FUS) HEK293T cells, we observe that there is greater nuclear translocation of TFEB when FUS is co-expressed, which in turn is amplified in the NLS-FUS mutants, P525L and R495X (**Figure 5A&B**). We quantify this as the ratio of nuclear to cytoplasmic fluorescence of GFP-TFEB from widefield images (**Figure 5C**, I_{Nuc}/I_{Cyto}), and validate that the observed effect is predominantly due to FUS expression, as all FUS-expressing cells have I_{Nuc}/I_{Cyto} (e.g., 1.3±0.23 to 1.73±0.27 for WT- and P525L-FUS, respectively) that are significantly increased compared to the control cells (0.79±0.17). Control cells were also treated with CQ, which is a known stressor for lysosomal dysfunction, hence result in the induction of nuclear translocation of TFEB (Roczniak-

Ferguson et al., 2012) As expected, Control+CQ gives the greatest I_{Nuc}/I_{Cyto} of 2.35±0.31 with 98% of

cells showing nuclear translocation (i.e., I_{Nuc}/I_{Cyto} >1); the latter is similar to both R495X and P525L at

86% and 90%, respectively.

Figure 5: Nuclear translocation of TFEB is promoted by FUS accumulation, and in particular for ALS-associated mutants. (A)

Widefield fluorescence intensity images showing mChe(-FUS) in green, GFP-TFEB in magenta and their composite image for

all samples tested. Scale bar, 10 μm. (B) Images showing cytoplasmic TFEB expression in WT-FUS, and nuclear translocation in

ALS-associated R495X-FUS. (C) Quantification of nuclear to cytoplasmic GFP-TFEB intensities (violin plot, primary y-axis), and

their corresponding value of nuclear translocation (bar chart, secondary y-axis), based on images in (A). Mean and S.E.M.

values of I_{Nuc}/I_{Cyto}: 0.79±0.17 (Control), 1.31±0.23 (WT), 1.68±0.23 (9R), 1.72±0.26 (16R), 1.70±0.23 (R495X), 1.73±0.27

(P525L) and 2.35± 0.31(Control+CQ). Analysis is based on 62—200 cells over three biological repeats. One-way ANOVA (with

Holm-Sidak's multiple comparison test), where n.s. is not significant, ** is p<0.005, *** is p<0.001 and **** is p<0.0001.

We have shown that the presence of FUS accumulation in cells could act as a stressor leading

to a loss in acidification of lysosomes, the latter of which trigger the cytoplasmic to nuclear translocation

of TFEB, as indicated by the increase in lysosomal biogenesis observed. To address if autophagy is

initiated beyond lysosomal biogenesis, we performed Western Blot analysis for autophagy markers, i.e.,

p62 and LC3 (**Figure S8A&B**). With FUS accumulation, we observe a slight upregulation of p62, but not

of LC3-II, both of which indicate that autophagy is inhibited at this early stage. The most significant

difference occurs upon addition of CQ to the control, which drives up normalised LC3-II and p62 by 2-

and 5-fold, respectively (**Figure S8B**). Furthermore, a FLIM-based aggregation assay was performed on

both GFP-labelled control and P525L-FUS cells, treated with CQ, and rapamycin (i.e., an autophagy

inducer). Of the latter, a relatively low concentration of 20 nM introduced in serum-starved media was

used, as higher concentrations led to cell death. Fluorescence lifetime of the GFP tag remains the same

for P525L-FUS, indicating that there are no changes in the aggregation state of FUS within the cells

(**Figure S8C&D**). We thus show that autophagy, and therefore the clearance and degradation of FUS

(and other proteins) are inhibited, despite activation of the TFEB-lysosomal pathway. Moreover, at this

stage, rapamycin is either toxic to cells at higher doses or not efficacious in reducing FUS aggregation

at lower concentrations.

Discussion

The experimental data described here reveal that intracellular FUS accumulation leads to i) cytoskeletal and organelle dysfunction and ii) perturbations in cellular homeostasis. To study the latter, we not only analysed different cellular FUS and stress-related models with unprecedented detail but also developed an approach with which we could determine the level of viscosity in live cells.

From a micro-rheological standpoint, the liquid condensed state of FUS is a viscoelastic Maxwell fluid, possessing both viscous liquid and elastic solid behaviour (Jawerth et al., 2020). As part of this study, we have developed an approach, based on correlative SPT and FLIM to measure intracellular FUS viscosity. This provides a higher throughput method than FRAP, commonly used in this field, as several condensates can be imaged simultaneously. Moreover, this approach allows us to probe condensates which are smaller than those suitable for more conventional FRAP analysis. It provides a robust route to quantifying this intracellular viscosity, for which there is no existing tool that is widely applicable in the field. We find that NLS-mutants of FUS, P525L- and R495X-FUS, contain more viscous aggregates in comparison to WT-FUS cells. Of the two FUS mutant variants studied, P525L-FUS has the greater aggregation propensity, leading to higher accumulation of drug-induced intranuclear aggregates and SGs. We further show that these mutant variants are also associated with a loss in cytoskeletal mechanoproperties in the cytoplasm, as seen by significant impairment of their actin and tubulin networks. As part of our study, we used SiR-dyes which permit this effect to be captured in live cells, as current studies typically rely on immunofluorescence staining of fixed samples. We further corroborated this using non-fluorescence-based, physical FD-AFM measurements, which show that P525L has half the apparent Young's modulus value of WT-FUS. Similar observations in neuronal models have linked axonal cytoskeletal integrity to the proteinopathy of FUS (Castellanos-Montiel et al., 2020; Giampetruzzi et al., 2019; Theunissen et al., 2021) and other FTD/ALS-implicated proteins, e.g., TAR DNA-binding protein (TDP43) (Baskaran et al., 2018; Briese et al., 2020). This includes a related study in *Xenopus laevis* retinal ganglion cells that demonstrated P525L-FUS reduces actin density specifically in

growth cones, resulting in localised softening of the axon(van Tartwijk et al., 2022). Furthermore, several ALS-linked genes that directly impair cytoskeletal dynamics (Castellanos-Montiel et al., 2020; Garone et al., 2020). We further observe that NLS-FUS mutant cells undergo DNA decondensation. At physiological conditions, nuclear-localised FUS regulates DNA repair through its interactions with HDAC1, which becomes dramatically reduced in the case of mutant FUS (Naumann et al., 2018; Wang et al., 2013; Yang et al., 2014) and can in turn increase euchromatin formation (Dinant et al., 2008; Lukas et al., 2011; Miller et al., 2010). Since epigenetic and chromatin regulatory factors are also phase separating in the nucleus (Gibson et al., 2019; Larson et al., 2017; Sanulli et al., 2019; Strom et al., 2017), it is conceivable that mutant, and thus more viscous FUS in the nucleus may also affect epigenetic and chromatin regulatory factors and thereby further enhance euchromatin formation.

We, among others, have recently shown that the dysfunction of common biochemical pathways which attenuate organelle function (e.g., lysosomes, ER, mitochondria), are prevalent to multiple neurodegenerative disease(Fowler et al., 2019; Koh et al., 2019; Lu et al., 2020). In combination, they lead to a loss in cellular integrity, and eventually culminate in neuronal dysfunction. To address organelle distribution and function in various cellular FUS and stress-induced models, we applied several different imaging techniques and assays. With ALS-associated mutant P525L-FUS, we observe that increased lysosomal biogenesis leads to lysosomal clustering, as well as mitochondria clustering, at the perinuclear region of cells. As a result, we see a decline in tubular ER due to the loss in lysosomal distribution throughout the cell cytoplasm. Tubular ER is predominantly found in axons, where it modulates synaptic calcium for neurotransmitter release (de Juan-Sanz et al., 2017), and supplies membranes and proteins for synaptic function(Gómez-Suaga et al., 2019). Hence any disruption to its function may have significant consequences for neurons. Additionally, the tubular ER also regulates mitochondrial fission (Merkwirth and Langer, 2008), and thus the mitochondria clustering seen in FUS-expressing cells may influence the latter. P525L-FUS has the largest effect on mitochondria shape, as indicated by the significant loss of eccentricity, i.e., the mitochondria become more circular from a tubular structure, which may stem from the effect of FUS on mitochondrial fission.

Many of the above results point towards a dysregulation of the ALP which is supported by

previous studies showing aberrant FUS aggregation disrupts the latter (Baskoylu et al., 2022; Soo et al.,

2015). We observe cytoplasmic to nuclear translocation of TFEB in the presence of FUS accumulation,

which we relate to the higher intralysosomal pH measured in FUS cells. In association, we observe

increased lysosomal biogenesis in the absence of changes to autophagy levels. These results support

the key role TFEB plays in autophagy regulation, and hence the adverse effects its dysfunction yields.

Despite using non-neuronal HEK293T and COS7 cells, we believe the underlying mechanisms observed

should hold true for a more physiologically relevant model. In support of this, it was recently shown

that there is impaired autophagy and neuronal dysfunction in a P525L-FUS knock in *C. elegans* model

(Baskoylu et al., 2022). Moreover, our findings align with the fact that autophagy dysfunction occurs

early on in disease pathology, which acts as an accelerator for neurodegeneration (Cortes and la Spada,

2019). Autophagy becomes even more vital in the case of aged, i.e., non-/slow-dividing neurones. It has

been shown that autophagy-deficient mice are more likely to accumulate aggregation-prone proteins,

increasing the risk of neurodegeneration (Hara et al., 2006; Komatsu et al., 2006). Proper function of

TFEB and lysosomes have been associated with ageing and longevity (Chang et al., 2017; Lapierre et al.,

2015). A recent study has shown that promotion of lysosomal function through the overexpression of

Pep4 (i.e., Cathepsin D homologue) in a yeast cell model leads to extended lifespans (Carmona-

Gutiérrez et al., 2011). Interestingly, our and the above findings contrast with the interactions of TDP43

and TFEB, where it has been shown that the nuclear translocation of TFEB occurs with a deficiency in

the first and leads to increased lysosome/autophagosome biogenesis (Xia et al., 2016). However, the

loss in TDP43 also leads to downstream impairment of lysosome-autophagosome fusion, leading to an

accumulation of autophagic vesicles, and thus both TDP43 and FUS models display autophagy

dysfunction. Further studies will be required to pinpoint the exact mechanisms of TFEB-lysosomal

dysfunction as well as its impact on organelles, in the presence of aberrant FUS aggregation.

Interestingly, many of the defects we have observed as part of our study pinpoint towards an induction

of early senescence, such as loss of cellular mechanoproperties, increased euchromatin formation, and

loss of protein homeostasis. Aberrant phase transition of FUS had been hypothesised to drive cellular

ageing by others in the field (Alberti and Hyman, 2016). It thus remains to be determined whether

neuronal cells indeed undergo early senescence as part of a cellular stress induced by aberrant FUS and

whether the latter precedes the formation of FUS condensates.

There is no current therapeutic strategy for ALS/FTD and many other neurodegenerative

diseases. Autophagy inducers, e.g., rapamycin, have been proposed as treatments to alleviate aberrant

FUS aggregation (Zhang et al., 2011). However, in our cell models, we did not observe any alleviation in

FUS aggregation upon rapamycin treatment. We have previously proposed that perinuclear lysosomal

clustering leading to the collapse of the ER network could be a hallmark mechanism across

neurodegenerative disease (Lu et al., 2020). Moreover, this has a further impact on cellular integrity

and function. For FUS, this is additionally seen in the depolymerisation of cytoskeletal proteins. Hence,

potential therapeutic strategies may want to be targeted towards alleviating cytoskeletal defects,

organelle dysfunction and perturbations in cellular homeostasis, rather than solely towards aberrant

protein aggregation.

Our study paints a picture of the effect of aberrant FUS condensates which is highly complex,

and impacts many key cellular structures/functions, such as the cytoskeleton, organelles, and cellular

homeostasis, which in turn, cannot be alleviated by targeting only one of these defects. Interestingly,

many of the above defects mimic signs of early senescence. Indeed, similar features resembling cellular

senescence have also been described in other neurodegenerative (Martínez-Cué and Rueda, 2020;

Sahu et al., 2022). It remains to be determined whether aberrant proteins first trigger a stress response

that affects major cellular functions, such as related to cellular senescence, before or after the

formation of aberrant condensates/aggregates. Understanding the latter, will have a major impact on

the understanding common factors underlying neurodegenerative diseases and for the development

of new therapeutic approaches.

Acknowledgements

CWC is funded by the Cambridge Trust and Wolfson College for her PhD. GSKS acknowledges funding from the Wellcome Trust (065807/Z/01/Z) (203249/Z/16/Z), the UK Medical Research Council (MRC) (MR/K02292X/1), Alzheimer Research UK (ARUK) (ARUK-PG013-14), Michael J Fox Foundation (16238) and Infinitus China Ltd. PHStG-H acknowledges funding from the Canadian Institutes of Health Research (406915 Foundation Grant and Canadian Consortium on Neurodegeneration in Aging Grant), Wellcome Trust Collaborative Award (203249/Z/16/Z), US Alzheimer Society Zenith Grant (ZEN-18-529769), Alzheimer Society of Ontario Chair in Alzheimer's Disease Research and National Institute of Aging (U01AG072572; R01AG070864).

Contributions

Study conceptualisation and design: GSKS; Cell culture and plasmids: CWC, ADS & AM; Data collection: CWC, AJZ & IM; Data analysis: CWC & AJZ; Feedback: PHStG-H, CFK & TJPK; Draft manuscript preparation: CWC & GSKS; Manuscript editing: all authors. All authors have given approval of the final manuscript.

Declaration of interests

The authors declare no competing interests.

Supporting information

Methodology and supplementary figures (PDF)

Data availability

Raw data is available through Cambridge University Repository Apollo (DOI: 10.17863/CAM.83295).

Analysis scripts are available upon request.

References

Alberti S, Hyman AA. 2016. Are aberrant phase transitions a driver of cellular aging? *BioEssays* **38**:959–968. doi:10.1002/BIES.201600042

Aman Y, Schmauck-Medina T, Hansen M, Morimoto RI, Simon AK, Bjedov I, Palikaras K, Simonsen A, Johansen T, Tavernarakis N, Rubinsztein DC, Partridge L, Kroemer G, Labbadia J, Fang EF. 2021. Autophagy in healthy aging and disease. *Nat Aging* **1**:634–650. doi:10.1038/s43587-021-00098-4

Baron DM, Kaushansky LJ, Ward CL, Sama RRK, Chian R-J, Boggio KJ, Quaresma AJC, Nickerson JA, Bosco DA. 2013. Amyotrophic lateral sclerosis-linked FUS/TLS alters stress granule assembly and dynamics. *Mol Neurodegener* **8**:30. doi:10.1186/1750-1326-8-30

Baskaran P, Shaw C, Guthrie S. 2018. TDP-43 causes neurotoxicity and cytoskeletal dysfunction in primary cortical neurons. *PLoS One* **13**. doi:10.1371/JOURNAL.PONE.0196528

Baskoylu SN, Chapkis N, Unsal B, Lins J, Schuch K, Simon J, Correspondence ACH, Hart AC. 2022. Disrupted autophagy and neuronal dysfunction in C. elegans knockin models of FUS amyotrophic lateral sclerosis. *Cell Rep* **38**:110195. doi:10.1016/j.celrep.2021.110195

Bosco DA, Lemay N, Ko HK, Zhou H, Burke C, Kwiatkowski TJ, Sapp P, Mckenna-Yasek D, Brown RH, Hayward LJ. 2010. Mutant FUS proteins that cause amyotrophic lateral sclerosis incorporate into stress granules. *Hum Mol Genet* **19**:4160–4175. doi:10.1093/HMG/DDQ335

Briese M, Saal-Bauernschubert L, Lüningschrör P, Moradi M, Dombert B, Surrey V, Appenzeller S, Deng C, Jablonka S, Sendtner M. 2020. Loss of Tdp-43 disrupts the axonal transcriptome of motoneurons accompanied by impaired axonal translation and mitochondria function. *Acta Neuropathol Commun* **8**:1–16. doi:10.1186/S40478-020-00987-6

Burdikova Z, Svindrych Z, Pala J, Hickey CD, Wilkinson MG, Panek J, Auty MAE, Periasamy A, Sheehan JJ. 2015. Measurement of pH micro-heterogeneity in natural cheese matrices by fluorescence lifetime imaging. *Front Microbiol* **6**:183. doi:10.3389/FMICB.2015.00183

Carmona-Gutiérrez D, Bauer MA, Ring J, Knaue HR, Eisenberg T, Büttner S, Ruckenstuhl C, Reisenbichler A, Magnes C, Rechberger GN, Birner-Gruenberger R, Jungwirth H, Fröhlich KU, Sinner F, Kroemer G, Madeo F. 2011. The propeptide of yeast cathepsin D inhibits programmed necrosis. *Cell Death Dis* **2**:e161. doi:10.1038/CDDIS.2011.43

Castellanos-Montiel MJ, Chaineau M, Durcan TM. 2020. The Neglected Genes of ALS: Cytoskeletal Dynamics Impact Synaptic Degeneration in ALS. *Front Clinic Neurosci* **14**:380. doi:10.3389/FNCEL.2020.594975/BIBTEX

Chang JT, Kumsta C, Hellman AB, Adams LM, Hansen M. 2017. Spatiotemporal regulation of autophagy during Caenorhabditis elegans aging. *Elife* **6**:e18459. doi:10.7554/ELIFE.18459

Chen PM, Gombart ZJ, Chen JW. 2011. Chloroquine treatment of ARPE-19 cells leads to lysosome dilation and intracellular lipid accumulation: Possible implications of lysosomal dysfunction in macular degeneration. *Cell Biosci* **1**:1–10. doi:10.1186/2045-3701-1-10/FIGURES/9

Chen W, Young LJ, Lu M, Zaccone A, Strohl F, Yu N, Schierle GSK, Kaminski CF. 2017. Fluorescence self-quenching from reporter dyes informs on the structural properties of amyloid clusters formed in vitro and in cells. *Nano Lett* **17**:143–149. doi:10.1021/acs.nanolett.6b03686

Chen Y, Liu H, Guan Y, Wang Q, Zhou F, Jie L, Ju J, Pu L, Du H, Wang X. 2015. The altered autophagy mediated by
TFEB in animal and cell models of amyotrophic lateral sclerosis. *Am J Transl Res* **7**:1574–1587.

Conte A, Lattante S, Zollino M, Marangi G, Luigetti M, del Grande A, Servidei S, Trombetta F, Sabatelli M. 2012.
P525L FUS mutation is consistently associated with a severe form of juvenile Amyotrophic Lateral
Sclerosis. *Neuromuscul Disord* **22**:73–75. doi:10.1016/J.NMD.2011.08.003

Cortes CJ, la Spada AR. 2019. TFEB dysregulation as a driver of autophagy dysfunction in neurodegenerative
disease: Molecular mechanisms, cellular processes, and emerging therapeutic opportunities. *Neurobiol*
Disease **122**:83–93. doi:10.1016/J.NBD.2018.05.012

Crocker JC, Grier DG. 1996. Methods of Digital Video Microscopy for Colloidal Studies. *J Colloid Interface Sci*
179:298–310. doi:10.1006/JCIS.1996.0217

Davidson NM, Gallimore PJ, Bateman B, Ward AD, Botchway SW, Kalberer M, Kuimova MK, Pope FD. 2020.
Measurement of the fluorescence lifetime of GFP in high refractive index levitated droplets using FLIM.
Phys Chem Chem Phys **22**:14704–14711. doi:10.1039/C9CP06395A

de Juan-Sanz J, Holt GT, Schreiter ER, de Juan F, Kim DS, Ryan TA. 2017. Axonal Endoplasmic Reticulum Ca2+
Content Controls Release Probability in CNS Nerve Terminals. *Neuron* **93**:867-881.e6.
doi:10.1016/j.neuron.2017.01.010

Dinant C, Houtsmuller AB, Vermeulen W. 2008. Chromatin structure and DNA damage repair. *Epigenetics*
Chromatin **1**:9. doi:10.1186/1756-8935-1-9

Esbjörner EK, Chan F, Rees E, Erdelyi M, Luheshi LM, Bertoncini CW, Kaminski CF, Dobson CM, Kaminski Schierle
GS. 2014. Direct observations of amyloid β Self-assembly in live cells provide insights into differences in
the kinetics of Aβ(1-40) and Aβ(1-42) aggregation. *Chem Biol* **21**:732–742.
doi:10.1016/j.chembiol.2014.03.014

Fowler PC, Garcia-Pardo ME, Simpson JC, O'Sullivan NC. 2019. NeurodegenERation: The Central Role for ER
Contacts in Neuronal Function and Axonopathy, Lessons From Hereditary Spastic Paraplegias and Related
Diseases. *Front Neurosci* **13**:1051. doi:10.3389/FNINS.2019.01051

Fujii S, Takanashi K, Kitajo K, Yamaguchi A. 2016. Treatment with a Global Methyltransferase Inhibitor Induces
the Intranuclear Aggregation of ALS-Linked FUS Mutant In Vitro. *Neurochem Res* **41**:826–835.
doi:10.1007/s11064-015-1758-z

Garone MG, Alfano V, Salvatori B, Braccia C, Peruzzi G, Colantoni A, Bozzoni I, Armirotti A, Rosa A. 2020.
Proteomics analysis of FUS mutant human motoneurons reveals altered regulation of cytoskeleton and
other ALS-linked proteins via 3'UTR binding. *Sci Rep* **10**:1–8. doi:10.1038/s41598-020-68794-6

Giampetruzzi A, Danielson EW, Gumina V, Jeon M, Boopathy S, Brown RH, Ratti A, Landers JE, Fallini C. 2019.
Modulation of actin polymerization affects nucleocytoplasmic transport in multiple forms of amyotrophic
lateral sclerosis. *Nat Commun* **10**:1–15. doi:10.1038/s41467-019-11837-y

Gibson BA, Doolittle LK, Schneider MWG, Jensen LE, Gamarra N, Henry L, Gerlich DW, Redding S, Rosen MK.
2019. Organization of Chromatin by Intrinsic and Regulated Phase Separation. *Cell* **179**:484.
doi:10.1016/J.CELL.2019.08.037

Gómez-Suaga P, Pérez-Nievas BG, Glennon EB, Lau DHW, Paillusson S, Mórotz GM, Calì T, Pizzo P, Noble W,
Miller CCJ. 2019. The VAPB-PTPIP51 endoplasmic reticulum-mitochondria tethering proteins are present

in neuronal synapses and regulate synaptic activity. *Acta Neuropathol Commun* **7**:35. doi:10.1186/s40478-
019-0688-4

Gonzalez A, Mannen T, Çağatay T, Fujiwara A, Matsumura H, Niesman AB, Brautigam CA, Chook YM, Yoshizawa
T. 2021. Mechanism of karyopherin-β2 binding and nuclear import of ALS variants FUS(P525L) and
FUS(R495X). *Sci Rep* **11**:3754. doi:10.1038/s41598-021-83196-y

Hara T, Nakamura K, Matsui M, Yamamoto A, Nakahara Y, Suzuki-Migishima R, Yokoyama M, Mishima K, Saito I,
Okano H, Mizushima N. 2006. Suppression of basal autophagy in neural cells causes neurodegenerative
disease in mice. *Nature* **441**:885–889. doi:10.1038/nature04724

Hockings C, Poudel C, Feeney KA, Novo CL, Hamouda MS, Mela I, Fernandez-Antoran D, Vallejo-Ramirez PP,
Rugg-Gunn PJ, Chalut K, Kaminski CF, Kaminski-Schierle G. 2020. Illuminating chromatin compaction in live
cells and fixed tissues using SiR-DNA fluorescence lifetime. *bioRxiv* 2020.05.02.073536.
doi:10.1101/2020.05.02.073536

Hofweber M, Hutten S, Bourgeois B, Niessing D, Madl T, Dormann Correspondence D. 2018. Phase Separation of
FUS Is Suppressed by Its Nuclear Import Receptor and Arginine Methylation. *Cell* **173**:706-713.e13.
doi:10.1016/j.cell.2018.03.004

Jawerth L, Fischer-Friedrich E, Saha Suropriya, Wang J, Franzmann T, Zhang X, Sachweh J, Ruer M, Ijavi M, Saha
Shambaditya, Mahamid J, Hyman AA, Jülicher F. 2020. Protein condensates as aging Maxwell fluids.
Science (1979) **370**:1317–1323. doi:10.1126/science.aaw4951

Koh JY, Kim HN, Hwang JJ, Kim YH, Park SE. 2019. Lysosomal dysfunction in proteinopathic neurodegenerative
disorders: possible therapeutic roles of cAMP and zinc. *Mol Brain* **12**. doi:10.1186/S13041-019-0439-2

Komatsu M, Waguri S, Chiba T, Murata S, Iwata JI, Tanida I, Ueno T, Koike M, Uchiyama Y, Kominami E, Tanaka K.
2006. Loss of autophagy in the central nervous system causes neurodegeneration in mice. *Nature*
441:880–884. doi:10.1038/nature04723

Kuimova MK. 2012. Mapping viscosity in cells using molecular rotors. *Phys Chem Chem Phys* **14**:12671–12686.
doi:10.1039/C2CP41674C

Lapierre LR, Kumsta C, Sandri M, Ballabio A, Hansen M. 2015. Transcriptional and epigenetic regulation of
autophagy in aging. *Autophagy* **11**:867–880. doi:10.1080/15548627.2015.1034410

Larson AG, Elnatan D, Keenen MM, Trnka MJ, Johnston JB, Burlingame AL, Agard DA, Redding S, Narlikar GJ.
2017. Liquid droplet formation by HP1α suggests a role for phase separation in heterochromatin. *Nature*
547:236–240. doi:10.1038/nature22822

Lu M, Christensen CN, Weber JM, Konno T, Läubli NF, Scherer KM, Avezov E, Lio P, Lapkin AA, Schierle GSK,
Kaminski CF. 2022. ERnet: a tool for the semantic segmentation and quantitative analysis of endoplasmic
reticulum topology for video-rate super-resolution imaging. *bioRxiv* 2022.05.17.492189.
doi:10.1101/2022.05.17.492189

Lu M, van Tartwijk FW, Lin JQ, Nijenhuis W, Parutto P, Fantham M, Christensen CN, Avezov E, Holt CE,
Tunnacliffe A, Holcman D, Kapitein L, Kaminski Schierle GS, Kaminski CF. 2020. The structure and global
distribution of the endoplasmic reticulum network are actively regulated by lysosomes. *Sci Adv* **6**:7209–
7225. doi:10.1126/SCIADV.ABC7209

Lukas J, Lukas C, Bartek J. 2011. More than just a focus: The chromatin response to DNA damage and its role in
genome integrity maintenance. *Nat Cell Biol* **13**:1161–1169. doi:10.1038/NCB2344

Lukinavičius G, Reymond L, D'Este E, Masharina A, Göttfert F, Ta H, Güther A, Fournier M, Rizzo S, Waldmann H,
Blaukopf C, Sommer C, Gerlich DW, Arndt HD, Hell SW, Johnsson K. 2014. Fluorogenic probes for live-cell
imaging of the cytoskeleton. *Nat Meth* **11**:731–733. doi:10.1038/nmeth.2972

Ma Y, Wen K, Liu M, Zheng J, Chu K, Smith ZJ, Liu L, Gao P. 2021. Recent advances in structured illumination
microscopy. *J Phys Photonics* **3**:24009. doi:10.1088/2515-7647/abdb04

Martínez-Cué C, Rueda N. 2020. Cellular senescence in neurodegenerative diseases. *Front Cell Neurosci* **14**:16.
doi:10.3389/FNCEL.2020.00016/BIBTEX

Martini-Stoica H, Xu Y, Ballabio A, Zheng H. 2016. The Autophagy–Lysosomal Pathway in Neurodegeneration: A
TFEB Perspective. *Trends Neurosci* **39**:221–234. doi:10.1016/J.TINS.2016.02.002

Mattenberger Y, James DI, Martinou JC. 2003. Fusion of mitochondria in mammalian cells is dependent on the
mitochondrial inner membrane potential and independent of microtubules or actin. *FEBS Lett* **538**:53–59.
doi:10.1016/S0014-5793(03)00124-8

Mauthe M, Orhon I, Rocchi C, Zhou X, Luhr M, Hijlkema KJ, Coppes RP, Engedal N, Mari M, Reggiori F. 2018.
Chloroquine inhibits autophagic flux by decreasing autophagosome-lysosome fusion. *Autophagy* **14**:1435–
1455. doi:10.1080/15548627.2018.1474314

Merkwirth C, Langer T. 2008. Mitofusin 2 Builds a Bridge between ER and Mitochondria. *Cell* **135**:1165–1167.
doi:10.1016/j.cell.2008.12.005

Miller KM, Tjeertes J v., Coates J, Legube G, Polo SE, Britton S, Jackson SP. 2010. Human HDAC1 and HDAC2
function in the DNA-damage response to promote DNA nonhomologous end-joining. *Nat Struc Mol Biol*
17:1144–1151. doi:10.1038/NSMB.1899

Moreira JMA, Scheipers P, Sørensen P. 2003. The histone deacetylase inhibitor Trichostatin A modulates CD4+ T
cell responses. *BMC Cancer* **3**:1–18. doi:10.1186/1471-2407-3-30/FIGURES/7

Murakami T, Qamar S, Lin JQ, Schierle GSK, Rees E, Miyashita A, Costa AR, Dodd RB, Chan FTS, Michel CH,
Kronenberg-Versteeg D, Li Y, Yang SP, Wakutani Y, Meadows W, Ferry RR, Dong L, Tartaglia GG, Favrin G,
Lin WL, Dickson DW, Zhen M, Ron D, Schmitt-Ulms G, Fraser PE, Shneider NA, Holt C, Vendruscolo M,
Kaminski CF, St George-Hyslop P. 2015. ALS/FTD Mutation-Induced Phase Transition of FUS Liquid
Droplets and Reversible Hydrogels into Irreversible Hydrogels Impairs RNP Granule Function. *Neuron*
88:678–690. doi:10.1016/j.neuron.2015.10.030

Naumann M, Pal A, Goswami A, Lojewski X, Japtok J, Vehlow A, Naujock M, Günther R, Jin M, Stanslowsky N,
Reinhardt P, Sterneckert J, Frickenhaus M, Pan-Montojo F, Storkebaum E, Poser I, Freischmidt A,
Weishaupt JH, Holzmann K, Troost D, Ludolph AC, Boeckers TM, Liebau S, Petri S, Cordes N, Hyman AA,
Wegner F, Grill SW, Weis J, Storch A, Hermann A. 2018. Impaired DNA damage response signaling by FUS-
NLS mutations leads to neurodegeneration and FUS aggregate formation. *Nat Commun* **9**:1–17.
doi:10.1038/s41467-017-02299-1

Nixon-Abell J, Obara CJ, Weigel A v., Li D, Legant WR, Xu CS, Pasolli HA, Harvey K, Hess HF, Betzig E, Blackstone
C, Lippincott-Schwartz J. 2016. Increased spatiotemporal resolution reveals highly dynamic dense tubular
matrices in the peripheral ER. *Science (1979)* **354**. doi:10.1126/SCIENCE.AAF3928

Novo CL, Wong E v., Hockings C, Poudel C, Sheekey E, Wiese M, Okkenhaug H, Boulton SJ, Basu S, Walker S,
Kaminski Schierle GS, Narlikar GJ, Rugg-Gunn PJ. 2022. Satellite repeat transcripts modulate
heterochromatin condensates and safeguard chromosome stability in mouse embryonic stem cells. *Nat*
Commun **13**:1–16. doi:10.1038/s41467-022-31198-3

Pu J, Guardia CM, Keren-Kaplan T, Bonifacino JS. 2016. Mechanisms and functions of lysosome positioning. *J Cell*
Sci **129**:4329. doi:10.1242/JCS.196287

Qamar S, Wang GZ, Randle SJ, Ruggeri FS, Varela JA, Lin JQ, Phillips EC, Miyashita A, Williams D, Ströhl F,
Meadows W, Ferry R, Dardov VJ, Tartaglia GG, Farrer LA, Kaminski Schierle GS, Kaminski CF, Holt CE,
Fraser PE, Schmitt-Ulms G, Klenerman D, Knowles T, Vendruscolo M, St George-Hyslop P. 2018. FUS Phase
Separation Is Modulated by a Molecular Chaperone and Methylation of Arginine Cation-π Interactions.
Cell **173**:720-734.e15.

Roczniak-Ferguson A, Petit CS, Froehlich F, Qian S, Ky J, Angarola B, Walther TC, Ferguson SM. 2012. The
Transcription Factor TFEB Links mTORC1 Signaling to Transcriptional Control of Lysosome Homeostasis. *Sci*
Signal **5**:ra42. doi:10.1126/SCISIGNAL.2002790

Sahu MR, Rani L, Subba R, Mondal AC. 2022. Cellular senescence in the aging brain: A promising target for
neurodegenerative diseases. *Mech Ageing Dev* **204**:111675. doi:10.1016/J.MAD.2022.111675

Sama RRK, Ward CL, Kaushansky LJ, Lemay N, Ishigaki S, Urano F, Bosco DA. 2013. FUS/TLS assembles into stress
granules and is a prosurvival factor during hyperosmolar stress. *J Cell Physiol* **228**:2222.
doi:10.1002/JCP.24395

Sanulli S, Trnka MJ, Dharmarajan V, Tibble RW, Pascal BD, Burlingame AL, Griffin PR, Gross JD, Narlikar GJ. 2019.
HP1 reshapes nucleosome core to promote phase separation of heterochromatin. *Nature* **575**:390–394.
doi:10.1038/s41586-019-1669-2

Schwerk C, Schulze-Osthoff K. 2005. Methyltransferase inhibition induces p53-dependent apoptosis and a novel
form of cell death. *Oncogene* **24**:7002–7011. doi:10.1038/sj.onc.1208855

Settembre C, di Malta C, Polito VA, Arencibia MG, Vetrini F, Erdin S, Erdin SU, Huynh T, Medina D, Colella P,
Sardiello M, Rubinsztein DC, Ballabio A. 2011. TFEB links autophagy to lysosomal biogenesis. *Science*
(1979) **332**:1429–1433. doi:10.1126/SCIENCE.1204592

Soo KY, Sultana J, King AE, Atkinson R, Warraich ST, Sundaramoorthy V, Blair I, Farg MA, Atkin JD. 2015. ALS-
associated mutant FUS inhibits macroautophagy which is restored by overexpression of Rab1. *Cell Death*
Disc **1**:15030. doi:10.1038/cddiscovery.2015.30

Strom AR, Emelyanov A v., Mir M, Fyodorov D v., Darzacq X, Karpen GH. 2017. Phase separation drives
heterochromatin domain formation. *Nature* **547**:241–245. doi:10.1038/nature22989

Tan A, Prasad R, Jho E hoon. 2021. TFEB regulates pluripotency transcriptional network in mouse embryonic
stem cells independent of autophagy–lysosomal biogenesis. *Cell Death Dis* **12**:1–14. doi:10.1038/s41419-
021-03632-9

Theunissen F, West PK, Brennan S, Petrović B, Hooshmand K, Akkari PA, Keon M, Guennewig B. 2021. New
perspectives on cytoskeletal dysregulation and mitochondrial mislocalization in amyotrophic lateral
sclerosis. *Transl Neurodegener* **10**:1–16. doi:10.1186/S40035-021-00272-Z

van Tartwijk FW, Wunderlich LCS, Mela I, Makarchuk S, Jakobs M, Qamar S, Franze K, Kaminski Schierle GS, St
George-Hyslop P, Lin JQ, Holt CE, Kaminski CF. 2022. Mutation of the ALS/FTD-associated RNA-binding
protein FUS alters axonal cytoskeletal organisation . *bioRxiv*.

Vigushin DM, Ali S, Pace PE, Mirsaidi N, Ito K, Adcock I, Charles Coombes R. 2001. Trichostatin A Is a Histone
Deacetylase Inhibitor with Potent Antitumor Activity against Breast Cancer in Vivo 1. *Clin Cancer Res*
7:971–976.

Waibel S, Neumann M, Rabe M, Meyer T, Ludolph AC. 2010. Novel missense and truncating mutations in
FUS/TLS in familial ALS. *Neurol* **75**:815–817. doi:10.1212/WNL.0B013E3181F07E26

Wang WY, Pan L, Su SC, Quinn EJ, Sasaki M, Jimenez JC, MacKenzie IRA, Huang EJ, Tsai LH. 2013. Interaction of
FUS and HDAC1 regulates DNA damage response and repair in neurons. *Nat Neurosci* **16**:1383–1391.
doi:10.1038/NN.3514

Willett R, Martina JA, Zewe JP, Wills R, Hammond GRV, Puertollano R. 2017. TFEB regulates lysosomal
positioning by modulating TMEM55B expression and JIP4 recruitment to lysosomes. *Nat Commun* **8**:1–17.
doi:10.1038/s41467-017-01871-z

Wolozin B, Ivanov P. 2019. Stress granules and neurodegeneration. *Nat Rev Neurosci* **20**:649–666.
doi:10.1038/s41583-019-0222-5

Xia Q, Wang H, Hao Z, Fu C, Hu Q, Gao F, Ren H, Chen D, Han J, Ying Z, Wang G. 2016. TDP-43 loss of function
increases TFEB activity and blocks autophagosome–lysosome fusion. *EMBO J* **35**:121–142.
doi:10.15252/EMBJ.201591998

Yang L, Gal J, Chen J, Zhu H. 2014. Self-assembled FUS binds active chromatin and regulates gene transcription.
Proc Natl Acad Sci USA **111**:17809–17814. doi:10.1073/PNAS.1414004111

Yang P, Mathieu C, Kolaitis RM, Zhang P, Messing J, Yurtsever U, Yang Z, Wu J, Li Y, Pan Q, Yu J, Martin EW,
Mittag T, Kim HJ, Taylor JP. 2020. G3BP1 Is a Tunable Switch that Triggers Phase Separation to Assemble
Stress Granules. *Cell* **181**:325-345.e28. doi:10.1016/J.CELL.2020.03.046

Young LJ, Ströhl F, Kaminski CF. 2016. A Guide to Structured Illumination TIRF Microscopy at High Speed with
Multiple Colors. *J Vis Exp* **111**:e53988. doi:10.3791/53988

Zhang Xiaojie, Li L, Chen S, Yang D, Wang Y, Zhang Xin, Wang Z, Le W. 2011. Rapamycin treatment augments
motor neuron degeneration in SOD1G93A mouse model of amyotrophic lateral sclerosis. *Autophagy*
7:412–425. doi:10.4161/AUTO.7.4.14541