Does prefrontal connectivity during task switching help or hinder children's performance?

Sina A. Schwarze[1*], Corinna Laube[1,2], Neda Khosravani[1], Ulman Lindenberger[1,3], Silvia A. Bunge[4], Yana Fandakova[1]

[1] Center for Lifespan Psychology, Max Planck Institute for Human Development, Berlin
[2] Fresenius University of Applied Sciences, Berlin, Germany
[3] Max Planck UCL Centre for Computational Psychiatry and Ageing Research, Berlin, Germany, and London, UK
[4] Department of Psychology and Helen Wills Neuroscience Institute, University of California at Berkeley

* Corresponding author: schwarze@mpib-berlin.mpg.de
Max Planck Institute for Human Development, Lentzeallee 94, 14197 Berlin, Germany

Keywords: task switching, cognitive control, frontoparietal network, functional connectivity, children, individual differences

The authors declare no conflict of interests.

Acknowledgements: Support for this research was provided by the DFG Priority Program SPP 1772 "Human performance under multiple cognitive task requirements: From basic mechanisms to optimized task scheduling" (Grant No. FA 1196/2-1 to Y.F.). S.A.S. and N.K. are doctoral fellows of the International Max Planck Research School on the Life Course (LIFE; http://www.imprs-life.mpg.de). The authors thank Julia Delius for editorial assistance and helpful comments.

Abstract

The ability to flexibly switch between tasks is key for goal-directed behavior and continues to improve in late childhood. Children's task switching difficulties are thought to reflect less efficient engagement of sustained and transient control processes, resulting in lower performance on blocks that intermix tasks (sustained demand) and trials that require a task switch (transient demand). Sustained and transient control processes are associated with frontoparietal regions, which develop throughout childhood and may contribute to task switching development. We examined age differences in the modulation of frontoparietal regions by sustained and transient control demands in children (8–11 years) and adults. Children showed greater performance costs than adults, especially under sustained demand, along with less upregulation of sustained and transient control activation in a set of frontoparietal regions. Compared to adults, children showed greater increases in connectivity between the inferior frontal junction (IFJ) and lateral prefrontal cortex (lPFC) from single to mixed blocks. For children whose sustained activation was less adult-like, increased IFJ-lPFC connectivity was associated with better performance. Children with more adult-like sustained activation showed the inverse effect. Taken together, these results suggest that the configuration of regions that control task switching effectively undergoes developmental changes in later childhood.

Introduction

The ability to switch flexibly between tasks enables individuals to adapt their behavior to changing environments and is a key component of cognitive control (Miyake et al., 2000; Miyake and Friedman, 2012). However, this flexibility comes at a cost. When faced with the demand to switch rapidly between tasks, individuals exhibit two types of performance decline. First, they show greater error rates and longer response times (RTs) on blocks of intermixed tasks (*mixed blocks*) compared to *single blocks* in which they perform only one task. These *mixing costs* are assumed to reflect sustained control demands, including the requirement to maintain and monitor multiple sets of rules associated with each task, so-called task sets (Braver et al., 2003; Pettigrew and Martin, 2016; Rubin and Meiran, 2005). Additionally, within the mixed blocks, performance declines further on trials that require a different task than the previous trial (i.e., *switch trials*) compared to trials repeating the previous task (i.e., *repeat trials*). These *switch costs* are thought to reflect transient control processes of inhibiting the no-longer-relevant task set (Allport et al., 1994; Meiran, 1996; Wylie and Allport, 2000) and retrieving and updating the newly relevant task set (Mayr and Kliegl, 2000; Rogers and Monsell, 1995).

Meta-analyses of functional magnetic resonance imaging (fMRI) data have repeatedly highlighted the left inferior frontal junction (IFJ) and the left superior parietal lobe (SPL) as key regions involved in task switching (Derrfuss et al., 2005; Kim et al., 2012; Niendam et al., 2012; Richter and Yeung, 2014; Worringer et al., 2019; Zhang et al., 2021). Sustained control processes, including task-set maintenance and monitoring, have been associated with the anterior and dorsolateral prefrontal cortex (aPFC and dlPFC, respectively; Badre, 2008; Braver et al., 2003; Gold et al., 2010) and the SPL (Brass and von Cramon, 2004; Bunge et al., 2003). Transient processes, such as task-set updating and attentional shifts, have been associated with the IFJ and the SPL (Brass et al., 2005; Brass and von Cramon, 2004; Braver

et al., 2003; Derrfuss et al., 2005; Gurd et al., 2002). Analyses based on functional connectivity have associated a frontoparietal network (FPN) of similar regions (i.e., the IFJ, SPL, dlPFC, and the precuneus) with transient control processes, while sustained control has been associated with a cingulo-opercular network including the anterior insula/frontal operculum (aI/fO), the dorsal anterior cingulate cortex (dACC), and the aPFC (Dosenbach et al., 2008, 2007, 2006).

Flexibly switching between tasks entails larger costs in children than adults (Cragg and Chevalier, 2012; Gupta et al., 2009; Huizinga et al., 2006; Huizinga and van der Molen, 2007). While overall task switching ability improves during middle childhood, mixing and switch costs show different developmental trajectories. Until around the age of 9–11 years, children show higher switch costs than adults, with less pronounced improvements later on. In contrast, mixing costs only approach adult levels during adolescence (13–15 years; Cepeda et al., 2001; Crone et al., 2006a, 2004; Reimers and Maylor, 2005). Together, the age differences reported by previous studies hint at performance improvements related to the development of both sustained and transient control. They also suggest that sustained and transient control follow different developmental paths, with a more protracted development for sustained control, presumably reflecting the later maturation of relevant frontal brain regions and their connectivity.

Sustained control requires that task sets are kept active in working memory (Crone et al., 2004). Hence, the notion of more enduring age differences in sustained control is in line with the protracted development of working memory, which reaches adult levels only late in adolescence (Crone et al., 2006c; Luciana et al., 2005). Sustained control is also essential for the ability to prepare task sets in advance and apply control proactively (Braver, 2012). Accordingly, previous studies have reported a shift from reactive to proactive control during childhood (e.g., Munakata et al., 2012). Monitoring processes, which entail the ability to

recognize the need to change or adapt behavior or cognition represents another essential

aspect of sustained control and also continues to develop into adolescence (Fandakova et al.,

2017; Tamnes et al., 2013). All these observations converge on the proposition that children

reach mature levels of sustained control, operationally defined as task-set maintenance and

monitoring, at a later age than mature levels of transient control, defined as task-set retrieval

and updating.

The postulated differences in developmental trajectories between sustained and

transient control raise questions about concomitant changes in the brain networks

implementing either form of control. The majority of developmental neuroimaging studies of

task switching to date support a view of quantitative age differences, with children recruiting

the same set of regions as adults, albeit in a less specific manner (Bunge and Wright, 2007;

Engelhardt et al., 2019; Velanova et al., 2008; Wendelken et al., 2012; Zhang et al., 2021; but

see Crone et al., 2006b; Morton et al., 2009). According to this view, the neural

underpinnings of task switching typically observed in adulthood are in place by middle

childhood and are subsequently fine-tuned, showing more differential activation between task

conditions in adults than children (Durston et al., 2006; Luna et al., 2015; Satterthwaite et al.,

2013).

Similarly, by the age of around 10 years, children show network organizations that are

similar to those of adults, but with continuing developmental changes in connectivity

strengths within and between networks (Engelhardt et al., 2019; Grayson and Fair, 2017;

Marek et al., 2015). To date, only one study has examined age differences in network

connectivity during task switching (Ezekiel et al., 2013). While children showed lower

connectivity within the FPN than adults, they also showed greater integration of the aPFC

into the FPN as reflected in ICA-based voxel-wise factor loadings. This pattern of results

suggests that children might configure brain networks in different ways than adults when

dealing with increased demands on cognitive control during task switching. However, the study did not directly examine differences between children and adults in how connectivity patterns dynamically change with increases in control demand. Thus, it is unclear whether developmental changes in connectivity patterns that effectively respond to increased cognitive control demands consist in the gradual evolution of the adult pattern, or whether one can observe a shift from a child pattern to an adult pattern. Conceptually, it is worth noting that the brain typically provides more than one pathway to implement a task, both within and across individuals (Edelman, 1987; Lautrey, 2003; Li and Lindenberger, 2002). In the course of brain maturation, one way of implementing cognitive control may gradually become more effective than another, such that children eventually shift from one configuration to another (e.g., Van der Maas and Molenaar, 1992).

Of note, children differ considerably in the development of task-switching proficiency (Dauvier et al., 2012; Fields et al., 2021) and the pace of brain maturation (e.g., Mills et al., 2021). For example, Church et al. (2017) reported that children who showed more adult-like (i.e., greater) frontoparietal activation during the preparatory period of task switching showed less activation in other regions when perceiving the target stimulus. Thus, age changes in brain-behavior mappings during task switching may not be uniform but follow different routes in different groups of children (e.g., Lautrey, 2003), creating an additional source of heterogeneity in the neural implementation of task-switching demands among children.

We examined the neural correlates of sustained and transient control during task switching in children aged 8 to 11 years and adults aged 20 to 30 years. Based on prior studies, we expected greater mixing and switch costs in children, reflecting the continued development of task-switching ability in middle and late childhood. We predicted that children would recruit similar frontoparietal brain regions as adults but that they would show less upregulation of activation in these regions with increased control demands. Further, we

predicted that age differences in task-related activation would be more pronounced for sustained activation than for transient activation. Compared to adults, we expected children to show less pronounced increases in IFJ connectivity, a key node of the frontoparietal network, with greater sustained control demands. Alternatively, we also examined whether children would involve additional brain regions in their control network to manage increasing control demands (cf. Ezekiel et al., 2013). Finally, we explored whether individual differences in sustained activation and connectivity are associated with individual differences in task-switching performance. In particular, we were interested in the degree to which the link between connectivity patterns and task-switching performance was modulated by the maturational status of individual children. We hypothesized that children with less mature patterns of sustained activation in the core task-switching network might show heightened connectivity with other brain regions, as a vicarious neural implementation of task-switching behavior. To test this hypothesis, we computed a deviation score for each child indicating how much their pattern of sustained activation diverged from the adult average. We then tested whether these deviation scores influenced the association between connectivity and task-switching performance.

Methods

Research Participants

Children between 8 and 11 years of age (n = 117; mean age = 10.0, SD = 0.71; 59 girls) and adults between 20 and 30 years of age (n = 53; mean age = 24.7, SD = 2.6; 28 women) were recruited from the internal participant database of the Max Planck Institute for Human Development. Participants were screened for MRI suitability, had no history of psychological or neurological diseases, and German was their primary language. All participants were right-handed. Adult participants and the participating children's parents provided informed consent; the participating children additionally provided written assent.

All participants were reimbursed with 10 € per hour spent at the MRI laboratory. The study

was approved by the ethics committee of the Freie Universität Berlin and conducted in line

with the Declaration of Helsinki.

Participants were included in analyses if they performed the task in accordance with

the task rules and had good fMRI data quality. Accuracy below 50% in the run of single

blocks (run 1, see below for more details on the paradigm) or accuracy below 35% in either

of the two runs of mixed blocks (run 2 and 3) was defined as poor performance. We defined a

threshold well above chance (33%) in the single runs to make sure participants knew and

could successfully apply the rules when no additional demand on switching among them was

present. fMRI volumes with framewise displacement (Power et al., 2012) above 0.4 mm were

marked as low-quality (see Dosenbach et al., 2017). If any of the fMRI runs exceeded 50% of

low-quality volumes, the participant was excluded from further analysis. Twenty children

were excluded because of excessive motion, four children due to poor performance, and four

children due to both excessive motion and poor performance. Thus, a total of 89 children

(mean age = 10.06, SD = 0.7, 50 girls) and 53 adults were retained in the analyses reported

below.

Experimental Design

Participants performed a task-switching paradigm consisting of three different

categorization rules (Figure 1A). In every trial, a face, a scene, and an object were presented,

and participants were cued by the shape of the background (diamond, circle, or hexagon) as

to whether to perform the face task, scene task, or object task. If the stimuli were presented in

a diamond-shaped background, the face image had to be classified according to the age of the

face (child vs. young adult vs. older adult). If the stimuli were presented in a circle, the image

of the scene had to be classified according to the depicted scenery (forest vs. desert vs. sea).

For stimuli presented in a hexagon, the image of the object had to be classified according to

its color (yellow vs. red vs. purple). All stimuli and the task cue appeared at the same time.

The arrangement of images on the screen varied randomly independent of the categorization

rule. For each of the three tasks, participants had to respond via button press using their right

index, middle, and ring finger. The assignment of finger to response option was held constant

across participants. There were two different stimuli for each of the three levels of each of the

three categorization rules, e.g., two faces of children, two yellow bags, two forest scenes, etc.,

resulting in a total of 18 different stimuli.

Participants performed three runs of 99 trials each (Figure 1B). Each trial lasted 2000

ms, followed by a fixation cross for a jittered time period (1000–6000 ms). After every 33

trials, there was an extended fixation period (20 s), resulting in three blocks of 33 trials per

run. In the first run (i.e., *single blocks*), each block contained trials of a single task (i.e., the

first 33 trials were of the face task, followed by 33 trials of the scene task, and 33 trials of the

object task). Across runs two and three (i.e., *mixed blocks*), the three tasks were intermixed,

so that participants needed to repeat a task in 50% of the trials (*repeat trial*) and switch to a

different task in the other 50% of the trials (*switch trial*). The switches between task rules

were not predictable, such that participants did not know in advance which task had to be

performed next. In run two, 50 trials required a task switch and 48 a task repetition; in run

three, 48 trials were switch trials and 50 were repeat trials. As the first trial of each run could

not be categorized as a switch or repeat trial, it was excluded from all analyses.

The data analyzed here are part of a larger training study, during which participants

were scanned multiple times. The present data were acquired during the second session of the

training study. The first session of the study included a computerized assessment of general

cognitive functioning not analyzed here, as well as a shortened version of the MRI task to

ensure that participants were familiar with the rules. Prior to scanning, participants were

shown the instructions again and practiced the task in a mock MRI scanner. The mock

scanner looked identical to a regular MRI scanner, with MR scanning sounds being mimicked

via speakers so that participants could get accustomed to the scanning environment. The three

runs of the MRI task were performed in the scanner after an initial T1-weighted scan during

which participants watched a muted cartoon.

Figure 1. Experimental Design. (A) The task-switching paradigm consisted of three tasks, the face task, the scene task, and the object task. Participants had to perform the task indicated by the shape of the background. Depending on the stimulus presented, one of three buttons had to be pressed in response (here indicated by the green button). (B) Participants performed one single run, where the three tasks were presented in blocks of 33 trials and two mixed runs, where the tasks were intermixed. Each run consisted of 99 trials presented for 2 s with a jittered inter-trial interval (ITI). For a full description of the task, see text. Note: per the regulation of *bioRxiv*, the face stimuli have been removed from this figure, participants saw colored photographs of faces with neutral facial expression.

Behavioral Analysis

Trials with response times (RT) below 200 ms and above 3000 ms were excluded from all analyses. Median RTs and mean accuracies were analyzed for effects of task rule condition (single block trials vs. repeat trials in mixed blocks vs. switch trials in mixed blocks), effects of age group (adults vs. children), and their interaction using separate linear mixed-effects models with a random intercept for subject. Only correct trials were considered for the RT analysis. Accuracy was calculated as the percentage of correct responses across all responses for a given condition. Outliers were defined as mean accuracy or median RTs that deviated by 3.5 standard deviations or more from the age-group mean, and were removed from analyses of accuracy and RT separately.

Main effects of condition were followed up by pairwise comparisons, with p-values FDR-corrected for multiple comparisons. Analyses were performed in R 4.0.3 (R Core Team, 2018) with the tidyverse (Wickham et al., 2019), ggpubr (Kassambara, 2020a), rstatix (Kassambara, 2020b), rmisc (Hope, 2013), sjPlot (Lüdecke et al., 2021), lme4 (Bates et al., 2022), lmerTest (Kuznetsova et al., 2017), and emmeans (Lenth et al., 2022) packages.

fMRI Data Acquisition and Preprocessing

Anatomical and functional MR images were collected on a 3-Tesla Siemens Tim Trio MRI scanner with a 32-channel head coil at the Max Planck Institute for Human Development. A high-resolution T1-weighted structural image was acquired (220 slices; 1 mm isotropic voxels; TE = 2.35 ms; FoV = 160 x 198 x 220). Functional runs consisted of 230 whole-brain echo-planar images of 36 interleaved slices (TR = 2000 ms; TE = 30 ms; flip angle = 80°; slice thickness = 3 mm; in-plane resolution (matrix) = 3 x 3 mm (72 x 72); FoV = 216 x 216 x 129.6). During the acquisition of functional images, the task was projected on a screen behind the participant's head that they could see via a mirror mounted on the head

coil. During task execution participant motion was monitored using Framewise Integrated Real-time MRI Monitoring (FIRMM; Dosenbach et al., 2017).

Preprocessing was performed using fMRIprep (Version 20.2.0; Esteban et al., 2019). For a detailed description, see the fMRIprep documentation (link). Briefly, BOLD images were co-registered to individual anatomical templates using FreeSurfer, which implements boundary-based registration (Greve and Fischl, 2009). Additionally, they were slice-time corrected (using AFNI; Cox and Hyde, 1997), and realigned (using FSL 5.0.9; Jenkinson et al., 2002). Finally, images were resampled into MNI152NLin6Asym standard space with a voxel size of 2 mm x 2 mm x 2 mm and spatially smoothed with an 8 mm FWHM isotropic Gaussian kernel using SPM12 (Functional Imaging Laboratory, UCL, UK).

fMRI Data Analysis

General Linear Model

GLM analyses were performed using SPM12 software. For each subject, we estimated two general linear models (GLM). The first GLM modeled sustained control demands in a blocked design (*sustained* GLM). Condition regressors included *single task*, consisting of three blocks of 127 s each within the first run, and *mixed task*, consisting of six blocks of 127 s each, evenly split over the second and third run. Data were high-pass filtered at 172 s.

A second GLM modeled transient control demands in an event-related design (*transient* GLM). Here, each stimulus presentation was coded as an event with zero duration, and convolved with a canonical hemodynamic response function (HRF). Separate regressors were included for correct *switch trials* and *repeat trials*. Incorrect trials, trials with extreme RTs (below 200 ms or above 3000 ms), trials with missed responses, and the first trial of each run were modeled as nuisance regressors. Data were high-pass filtered at 128 s.

To minimize head motion artifacts, both GLMs included the amount of frame displacement per volume, in mm (Power et al., 2012), realignment parameters (three translation and three rotation parameters), and the first six anatomical CompCor components (as provided by fMRIprep; Behzadi et al., 2007) as regressors of no interest. The first five volumes of each run were discarded to allow for stabilization of the magnetic field and temporal autocorrelations were estimated using first-order autoregression.

Sustained control activation was defined as higher activation in mixed compared to single blocks (mixed > single) in the sustained GLM. Transient control activation was defined as higher activation on switch trials compared to repeat trials (switch > repeat) within the mixed runs of the transient GLM. The sustained GLM modeled control in a continuous manner, such that mixed blocks included both switch and repeat trials (e.g., Braver et al., 2003). With these separate models, it is not possible to directly compare single and repeat trials; thus, we also constructed an event-related GLM that included correct single, repeat, and switch trials as separate regressors of interest (*control* GLM). All nuisance regressors and additional parameters were identical to the transient GLM.

Region-of-Interest Definition and Analysis

First, we identified regions of interest (ROIs) involved in sustained and transient control over all subjects (whole-brain analyses; $p < .05$ family-wise-error (FWE) corrected). For sustained control, we selected voxels that showed greater activation in mixed blocks than in single blocks in the sustained GLM. We identified regions involved in transient control by selecting voxels showing greater activation on switch than on repeat trials in the transient GLM. Second, to identify ROIs involved in both sustained and transient control, which we referred to as *overlap* ROIs, we applied inclusive masking which identified common voxels for sustained and transient activation. Finally, to define regions *exclusively* involved in sustained or transient control, we applied the map of the overlap as an exclusive mask to the

sustained control contrast—which we termed *sustained* ROIs—and the transient control

contrast, termed *transient* ROIs. Only clusters of at least 50 voxels were considered for ROI

definition. If clusters were stretched across multiple anatomical areas, they were additionally

masked using anatomical regions of the Harvard-Oxford atlas (Makris et al., 2006)

thresholded at 30%. No anatomical mask is available for the IFJ; therefore this ROI was

created by splitting the cluster based on a meta-analysis (Derrfuss et al., 2005). Hemisphere

masks were applied to restrict clusters that extended beyond one hemisphere. Finally, ROIs

were binarized and used as masks for beta-parameter extraction using MarsBar (Brett et al.,

2002).

     With these ROI analyses, we sought to test whether children and adults differed in the

extent of upregulation of sustained and transient control activation with increased control

demand. Upregulation of sustained control was computed as the difference between

parameters of mixed and single blocks, and upregulation of transient control was computed as

the difference between parameters of switch and repeat trials. We then compared the extent

of upregulation between age groups for each ROI using t-test in R, reporting Cohen's d as

effect size (Cohen, 1988) and p-values FDR-corrected for multiple comparisons.

     To investigate whether children or adults recruited additional brain regions beyond

the ones we identified over both age groups, we computed exploratory whole-brain age

comparisons of the mixed > single and switch > repeat contrasts in the sustained and transient

GLM, respectively.

Psychophysiological Interaction Analysis

     Generalized psychophysiological interactions (gPPI; McLaren et al., 2012) were

analyzed using the CONN toolbox (Version 20b; Whitfield-Gabrieli and Nieto-Castanon,

2012). Unlike correlational measures of functional connectivity, PPI models how the strength of coupling between a seed region and a voxel elsewhere in the brain differs across conditions. To this end, a GLM predicting the time series of the target voxel by the interaction regressor is constructed. The interaction regressor is created by multiplying the time series of the seed region with the task regressor for each condition (i.e., the condition onset times convolved with the HRF). The PPI analysis focused on effects of sustained control, as we had predicted greater age differences here.

The main effects of the two sustained control conditions (single and mixed blocks) and the effect of the nuisance regressors previously applied to the sustained GLM were regressed from the BOLD signal time series before estimating the interaction factor. Importantly, voxel-level time series were estimated using the smoothed data, while ROI level (i.e., seed) time series were estimated using unsmoothed data to prevent a "spillage" of the BOLD signal of voxels outside the ROI into the ROI time series. Interaction parameters were estimated separately for each condition, i.e., single and mixed blocks (McLaren et al., 2012).

Based on the prominent role of the left IFJ in task switching (Derrfuss et al., 2005; Kim et al., 2012; Richter and Yeung, 2014) and the fact that it showed activation in our univariate analysis, we selected it as a seed region for the PPI analysis. More specifically, a sphere of 6 mm radius was placed around the group peak coordinates of the mixed > single contrast of the sustained GLM.

We compared the difference in seed-to-whole brain connectivity for the IFJ for mixed vs. single blocks between children and adults. Analyses were performed on a gray-matter mask of the whole brain and corrected for multiple comparisons at the cluster level (p < .05 FDR-corrected, voxel threshold at p < .001 uncorrected). Beta estimates were extracted from clusters showing age differences in connectivity for the mixed > single contrast to visualize the results and to relate connectivity patterns to performance in the task. Additionally, the

large cluster in the prefrontal cortex (PFC) was split into a medial and lateral part, as these

have been associated with different aspects during task switching (Koechlin et al., 2000).

More specifically, the medial PFC (mPFC) was defined by the superior frontal and

paracingulate gyrus and the lateral PFC (lPFC) by the frontal pole and middle frontal gyrus of

the Harvard-Oxford Atlas (Makris et al., 2006), thresholded at a probability of 30%.

Functional Activation Deviation Scores

To take individual differences among children in sustained activation into

consideration, we computed functional activation deviation scores (cf. Düzel et al., 2011;

Fandakova et al., 2015). The deviation scores reflect the similarity of task-related activation

of an individual child to the adult mean. To this end, we created a mask including all clusters

that showed sustained activation across adults (p < .05, FWE-corrected). The deviation score

for each child was then calculated as the difference between the average T-value of the

sustained control contrast (mixed > single) of all voxels inside this adult mask and outside it.

Thus, a negative score represented more adult-like activation patterns, and a more positive

score represented less adult-like patterns.

To examine relations between deviation scores, connectivity, and performance, we

used the adjusted accuracy mixing costs. Adjusted mixing costs were calculated by dividing

the difference between single- and mixed-block accuracy by the single-block accuracy, which

represents the average accuracy across trials in single blocks and thus allowed us to adjust

mixing costs to differences in overall performance. As a measure of connectivity, we used the

difference in PPI beta parameters between mixed and single blocks of the connections

between the IFJ and the lateral and medial PFC clusters separately, to investigate whether

they showed different influences on performance.

<div align="center">Results</div>

Greater Costs of Task Switching in Children

Accuracy results are shown in Figure 2A. We fit a linear mixed-effects model with accuracy as the dependent variable and the factors age group (children vs. adults) and condition (single vs. repeat vs. switch), the interaction between these two factors and a random intercept for subject using log-likelihood optimization. To enable comparisons of mixing costs (i.e., performance differences between repeat and single trials) and switch costs (i.e., performance differences between switch and repeat trials), the condition repeat was used as the reference condition in the model. The model revealed a significant effect of age group ($\beta = -0.068$, $p < .001$) and the switch condition (vs. repeat; $\beta = -0.031$, $p = .01$). The effect for the single condition (vs. repeat) was at trend level ($\beta = -0.024$; $p = .06$). Thus, children showed overall lower accuracy ($M = 0.81$, $SD = 0.16$) than adults ($M = 0.96$, $SD = 0.04$) and across age groups that accuracy was highest on single trials ($M = 0.94$, $SD = 0.08$), followed by repeat trials ($M = 0.87$, $SD = 0.13$), and then switch trials ($M = 0.80$, $SD = 0.17$). There were significant interactions of age group with both conditions (single: $\beta = -0.089$, $p < .001$; switch: $\beta = -0.054$, $p < .001$) indicating that children showed greater differences between the conditions than adults, especially with respect to the difference between single and repeat conditions. Taken together, while both age groups demonstrated reliable mixing and switch costs, the age differences for mixing costs (i.e., the interaction of age group and single condition) were more pronounced than the age differences for switch cost (i.e., the interaction of age group and switch condition).

RT results are shown in Figure 2B. The mixed-effects model of RT, set up with the same fixed and random effects as the model of accuracy, revealed significant effects of age group ($\beta = 0.28$, $p < .001$) and both conditions (single: $\beta = -0.27$, $p < .001$; switch: $\beta = 0.24$, $p < .001$; both compared to repeat as reference). There was no significant age-group-by-condition interaction, neither with the single condition ($\beta = 0.0047$, $p = .83$) nor with the

switch condition (β = 0.013, p = .54). Overall, children (M = 1.45 s, SD = 0.26) responded

more slowly than adults (M = 1.17 s, SD = 0.26). Across both groups, responses to single

trials (M = 1.08 s, SD = 0.19) were fastest, followed by repeat (M = 1.35 s, SD = 0.20) and

switch trials (M = 1.60 s, SD = 0.22).

Taken together, children and adults demonstrated reliable declines in performance

both in terms of lower accuracy and slower RTs going from single blocks to mixed blocks,

and from repeat trials to switch trials. Critically, children showed greater declines in accuracy

than adults with increasing task switching demands, with age differences being particularly

pronounced for mixing costs.

Figure 2: Behavioral results. (A) Proportion of correct responses for single, repeat, and switch trials. Across age groups accuracy was highest for single trials, followed by repeat and switch trials. Children showed greater mixing (single vs. mixed trials) and switch (repeat vs. switch trials) costs, while age differences were more pronounced for mixing costs. (B) Response time for single, repeat, and switch trials. Across age groups, responses to single trials were fastest, followed by repeat and switch trials. Mixing and switch costs in response times (RT) were similar across age groups.

Frontoparietal Brain Regions Associated with Sustained and Transient Control

Analyses of whole-brain activation associated with sustained control (mixed > single blocks) across children and adults indicated enhanced activation in multiple bilateral prefrontal and parietal regions (see Supplementary Table 1), including the IFJ, SPL, dACC, and dlPFC (see Figure 3, green color). A whole-brain analysis of transient control (switch > repeat trials) across children and adults revealed enhanced activation in left lateral and medial parietal and prefrontal regions as well as bilateral occipital regions, including the left IFJ, left SPL, left dACC, and the bilateral precuneus (Figure 3, red color; Supplementary Table 1).

Activation patterns of sustained and transient control partially overlapped. Inclusive masking revealed several regions activated by both sustained and transient control demands, including the left IFJ, left SPL, left dACC, left inferior lateral occipital cortex (iLOC), left anterior insular (aI), and left middle frontal gyrus (MFG). On the other hand, regions exclusively showing sustained control activation included the bilateral dlPFC, right IFJ, right SPL, right aI, and right MFG. The bilateral precuneus and a region in the bilateral premotor cortex (preMC) were exclusively engaged during transient control.

Thus, across age groups, we found frontoparietal regions to be associated with sustained and/or transient control, including regions—the IFJ, SPL, dACC, dlPFC, and precuneus—that have been associated with task switching in previous research in adults (Kim et al., 2012; Richter and Yeung, 2014). Of these regions, the left IFJ, SPL, and dACC showed activation for both sustained and transient control, bilateral dlPFC exclusively for sustained control, and bilateral precuneus exclusively for transient control.

To test whether sustained control effects were driven by switch trials during mixed blocks or were also present when comparing single and repeat trials only, we performed a control GLM that included single, repeat, and switch trials in an event-related design. Univariate results of the repeat > single contrast of this control GLM were comparable to the mixed > single contrast of the sustained GLM (see Supplementary Figure 1), suggesting that

sustained control effects were jointly driven by switch and repeat trials within the mixed

blocks.

Figure 3: Univariate activation across age groups. Transient control activation (switch > repeat trials) depicted in red, sustained control activation (mixed > single blocks) in green, and overlap (voxels showing transient and sustained activation) in blue. N = 142 (53 adults, 89 children), p < .05 FWE-corrected, k = 50.

Less Upregulation of Sustained and Transient Control Activation in Children

Having identified regions across the whole sample that are involved in one or both

types of control across both age groups, we next sought to examine age differences in the

modulation of activation by sustained and transient control demands within these regions. We

tested for age differences in the upregulation of sustained control activation by performing t-

tests on the difference between mixed and single parameter estimates. As shown in Figure

4A, the overlap ROIs, including left IFJ, SPL, and dACC, showed a significant age difference

in the degree of increase in activation from single to mixed blocks (IFJ: $t(118.66) = -3.86$,

$p_{corrected} < .001$, Cohen's $d = 0.66$; SPL: $t(125.71) = -4.02$, $p_{corrected} < .001$, $d = 0.68$; dACC:

$t(121.59) = -5.57$, $p_{corrected} < .001$, $d = 0.72$; FDR-corrected), such that, across all ROIs

examined, children showed less upregulation of activation from single to mixed blocks.

Similarly, we found significant age differences in the extent of upregulation of

activation by sustained control demands in the bilateral dlPFC, right IFJ, and right SPL, ROIs

that were exclusively associated with sustained control (dlPFC: $t(127.31) = -4.49$, $p_{corrected} <$

.001, $d = 0.76$; IFJ: $t(107.1) = -3.12$, $p_{corrected} = .003$, $d = 0.54$; SPL: $t(122.58) = -2.96$,

$p_{corrected}$ = .004, d = 0.5, FDR-corrected, see Figure 4A and Supplementary Figure 2),

indicating that children showed less upregulation of task-related activation in the mixed

blocks than adults. Taken together, these results suggest that even though children

upregulated brain activity in regions that have been identified as key task-switching regions

across both age groups with increasing demands on sustained control, they did so to a lesser

extent than adults, both for overlap and sustained ROIs.

Next, we tested for age differences in the upregulation of transient control activation

by performing t-tests on the difference between switch and repeat parameter estimates (see

Figure 4B). The overlap ROIs in left IFJ, SPL, and dACC showed significant age differences

in upregulation of transient control activation (IFJ: t(130.14) = –2.25, $p_{corrected}$ = .045, d =

0.38; SPL: t(138.61) = –2.30, $p_{corrected}$ = .045, d = 0.37, dACC: t(135.53) = –3.74, $p_{corrected}$ =

.002, d = 0.62, FDR-corrected). The bilateral precuneus, exclusively associated with transient

control, also showed greater upregulation of transient control activation in adults than in

children (t(134.18) = –2.32, $p_{corrected}$ = .045, d = 0.39, FDR-corrected). Taken together,

children upregulated activation due to transient control demand to a lesser extent than adults

did across the key regions implicated in task-switching.

The preceding ROI-based analyses revealed age differences in activated regions

common to adults and children. However, it is possible that children or adults showed

different activation in additional regions. Thus, we also conducted an exploratory analysis

investigating age differences in sustained and transient activation on the whole-brain level.

Consistent with the results of the ROI analysis, the whole-brain analysis of sustained control

showed greater upregulation of activation (mixed > single) for adults than children in the left

SPL, left IFJ/dlPFC, and left superior lateral occipital cortex (p < .05 FWE-corrected;

Supplementary Figure 3). No regions showed greater upregulation of sustained control

activation in children. Further, no regions showed age differences in upregulation of

activation due to transient control demand (switch > repeat); for further details, see the

Supplementary Material.

A Sustained Control Activation

B Transient Control Activation

Figure 4: Age differences in activation due to sustained and transient control demands by ROI. (A) Extracted parameter estimates for single blocks (blue) and for mixed blocks (green). Children showed less upregulation of activation due to sustained control demand in the left IFJ, SPL, and dACC, and the bilateral dlPFC. Neither age group showed upregulation of sustained control activation in the Precuneus. (B) Extracted parameter estimates for repeat trials (yellow) and for switch trials (orange). Children showed less upregulation of activation due to transient control demands in the left IFJ, SPL, dACC, and the bilateral precuneus. Neither age group showed upregulation of transient control activation in the dlPFC.

Greater Increases in Connectivity Under Increased Sustained Control Demand in Children

Based on previous research (e.g., Cepeda et al., 2001) showing that differences between adults and children in the age range investigated here are more pronounced during sustained control, and the fact that our results showed more pronounced age differences for mixing costs, we focused on connectivity associated with sustained control. More specifically, we were interested in the regions showing age differences in connectivity with the IFJ under increased sustained control demands. The left IFJ was selected as a seed, as it has been identified as a key brain region during task switching (Derrfuss et al., 2005; Kim et al., 2012; Richter and Yeung, 2014).

A whole-brain PPI analysis revealed that children showed a more pronounced increase in connectivity with the IFJ from single to mixed blocks in the bilateral angular gyrus (AG), and the medial and lateral prefrontal cortex (mPFC and lPFC, respectively) ($p < .001$, FDR cluster corrected $p < .05$). Neither of these regions overlapped substantially with the regions showing greater sustained control activation across age groups (see Supplementary Figure 5). The PFC cluster was split to account for possible differences between lateral and medial regions (Koechlin et al., 2000). The lPFC cluster was localized anteriorly adjacent to the bilateral dlPFC that showed sustained activation, and the mPFC cluster was localized anterior to the dACC showing transient and sustained activation. Figure 5 shows extracted connectivity parameters of the clusters that showed age differences in the

comparison of mixed and single blocks. Specifically, age differences were driven by the fact

that children showed higher IFJ–mPFC, IFJ–lPFC, and IFJ–AG connectivity in mixed than in

single blocks, whereas adults did not show differences in connectivity for these regions

across conditions. Additionally, Figure 5 illustrates the variation in the increase of

connectivity from single to mixed blocks among children, such that some showed very steep

increases, while others showed rather shallow increases or even decreases. Thus, these data

suggest that many, but not all children showed a greater control demand-related increase in

connectivity to broad swaths of prefrontal and parietal cortices outside the core areas

implicated in task-switching. Note, however, that while these age group differences were

identified in the aggregated group comparison, the individual estimates presented in Figure 5

suggested large interindividual differences, such that some children showed more pronounced

increases in connectivity, while others showed smaller increases.

Figure 5: Age differences in connectivity increases under greater sustained control demands.
PPI parameter estimates for clusters showing greater connectivity with the IFJ during mixed
blocks (green) than during single blocks (blue) in children compared to adults identified in a
whole-brain analysis with the left IFJ as a seed.

Connectivity Effects on Performance Depend on Adult-like Activation in Frontoparietal

Regions

Children showed less upregulation of activation in sustained control regions, but a

greater increase in IFJ connectivity with additional regions as a function of increased

demands on sustained control, including regions in lateral and medial PFC that have been

implicated in cognitive control processes (e.g. Braver et al., 2003). This pattern of results

raises the important question whether the broader connectivity pattern observed in children

represents an alternative, presumably developmentally earlier way to support task

performance specifically in those children whose frontoparietal activation patterns differ

more markedly from those observed among adults. If so, the relation of connectivity

increases and accuracy may vary across children, and may depend on the extent to which an

individual child showed a mature pattern of sustained control activation.

To test this hypothesis, we computed deviation scores indicating how similar each

child's sustained control activation was to that of the average adult. For every individual

child, this score was calculated by subtracting the average T-value of the mixed > single

contrast of all voxels *inside* the clusters that showed sustained control activation in adults

from the average T-value of the mixed > single contrast of all voxels *outside* of the clusters

that showed such activation in adults (cf. Düzel et al., 2011; Fandakova et al., 2015). Thus, a

more negative score indicates less deviation from the average adult activation pattern, while a

more positive score indicates greater deviation and therefore a less adult-like activation

pattern. This approach allowed us to take two previously observed aspects of age differences

in brain activation into account. First, age differences have been found in the strength of task-

related activation within the same regions, such that children show overall weaker activation

than adults (e.g., Wendelken et al., 2012). Second, children have been shown to have less

specific task-related activation than adults (e.g., Durston et al., 2006). Both of these aspects

would result in more positive scores, indicating a greater difference to the average adult

pattern.

We performed a linear regression on the accuracy mixing costs with deviation scores,

increases in IFJ–PFC connectivity (in response to increased sustained control demands), and

the deviation score by connectivity interaction as predictors. The goal of these analyses was

to examine whether the association between connectivity and performance depended on the

maturity of activation under sustained control demands. Regressions were performed separately for connectivity with the lPFC and mPFC clusters.

We found significant main effects of deviation score (β = 0.07, p = .009) and of increased IFJ–lPFC connectivity (β = –0.18, p = .026) on mixing costs. However, these effects were qualified by a significant interaction between deviation score and connectivity (β = –0.22, p = .03). Thus, the association between connectivity increases and performance was modulated by the extent to which children showed an adult-like pattern of sustained control activation (Figure 6). Specifically, for children with a more adult-like activation pattern (i.e., more negative deviation scores), a greater increase in connectivity for mixed blocks was associated with greater mixing costs, or lower performance. Thus, for children with more adult-like activation patterns, the inclusion of the lPFC in the control network due to greater sustained control demands was associated with poorer performance.

In contrast, for children who showed less adult-like sustained control activation (i.e., more positive deviation scores), accuracy mixing costs and connectivity increases were *negatively* correlated. Thus, for children who deviated more from the expected sustained activation pattern, stronger incorporation of the lPFC into the control network—as evidenced by heightened connectivity with IFJ in response to an increase in the need for sustained control—was related to better performance. Similar results were obtained regarding IFJ–mPFC connectivity, albeit with the interaction of deviation scores and connectivity increases at a trend level (β = –0.15, p = .1). Neither the connectivity between the IFJ and the left AG nor the right AG showed such a relationship with performance.

Taken together, these results suggest that greater lPFC involvement represents an adaptive alternative to implement task-switching affordances among children whose FPN activation patterns deviate more from the adult pattern.

Figure 6: Interaction effect of deviation score and connectivity on accuracy mixing costs in children. (A) Sustained control activation (mixed > single) across adults (p < .05, FWE-corrected) used as reference for estimating how similar an individual child's activation pattern was to this adult pattern. An example of a child showing more adult-like activation for the contrast shown in blue (p < .001, uncorrected). An example of a child showing less adult-like activation for the contrast shown in red (p < .001, uncorrected). (B) Interaction effect. Children who showed less adult-like activation patterns (red line) showed a negative association between connectivity increases from single to mixed block and mixing costs, such that greater increases in connectivity were associated with lower costs (i.e., better performance). Children showing more adult-like activation patterns (blue line) showed a positive association between connectivity increases and mixing costs, such that greater increases in connectivity were associated with greater mixing costs (i.e., worse performance).

Discussion

In the present study, we sought to shed light on the development of sustained and transient control processes by taking a closer look at (a) age group differences between children and adults in performance and neural network activations and connectivity related to transient and sustained task-switching demands; and (b) individual differences in the association between task-switching proficiency and neural network activations and connectivity among children aged 8 to 11 years. Our results indicate that children show greater mixing and switch costs than adults, accompanied by less upregulation of activation in relevant brain areas during sustained and transient increases in processing demands. Age differences were more pronounced for mixing costs and sustained control activation, in line with the notion that brain areas and circuits supporting sustained switching demands mature more slowly than those supporting transient switching demands. Task-based connectivity analyses revealed that connectivity between the IFJ and the lPFC increased with greater sustained control demands in children. Critically, the association between task-related connectivity and task-switching performance was linked to how dissimilar the activation pattern of an individual child was to the typical adult pattern: Children who showed less adult-like activation in response to sustained control demands showed better performance with greater IFJ–lPFC connectivity increases, whereas children with a more adult-like activation pattern showed the opposite association, that is, lower performance with greater IFJ–lPFC connectivity increases. This finding suggests that children whose FPN is less mature make adaptive use of an alternative neural implementation of cognitive control to cope with increases in sustained switching demands.

In general, however, children in the present study recruited the same frontoparietal regions during task switching as adults (i.e., the IFJ, SPL, dACC, dlPFC, and precuneus), but they did so in a less specific manner. These results support previous findings on the

development of task switching (Bunge and Wright, 2007; Cepeda et al., 2001; Crone et al., 2006b, 2004; Engelhardt et al., 2019; Reimers and Maylor, 2005; Velanova et al., 2008; Wendelken et al., 2012; Zhang et al., 2021). However, it is worth noting that previous findings regarding age differences in switch costs are inconsistent, with some studies reporting comparable switch costs in children and adults (Anderson et al., 2001; Luca et al., 2003; Reimers and Maylor, 2005). One potential reason for these inconsistencies is the number of task rules participants had to switch among. The majority of previous studies implemented paradigms that required switches between two task rules, whereas the present study intermixed three task rules, rendering the paradigm more complex. Our results extend previous research by showing that transient processes required during task switching, such as inhibiting the no-longer-relevant task set, and updating the new task sets (Allport et al., 1994; Mayr and Kliegl, 2000; Meiran, 2000) may still show an age-related improvement when the demands on switching are high. Future studies manipulating the number and complexity of rules during task-switching should be particularly informative with respect to the factors that contribute to individual differences in the development of transient control. Previous research in this direction has shown that age differences in working memory performance were not dependent on the ability to manage multiple rules in working memory per se, but rather on improvements in applying hierarchical rules (Unger et al., 2016), underlining the importance of considering multiple facets of control demands. Additionally, while children and adults have been shown to exhibit similar levels of inhibitory involvement during task switching, as measured with $n - 2$ repetition costs, drift diffusion models suggest that these costs were associated with different processes in children and adults (Schuch and Konrad, 2017). Specifically, while adults showed slower response selection on trials of a previously inhibited task, children showed longer task preparation on such trials, further suggesting that transient control processes continue to develop past age 11.

In our task, age differences were more pronounced in sustained control processes than in transient control processes, at both behavioral and neural levels. These results suggest that sustained control processes such as task-set maintenance and monitoring (Braver et al., 2003; Pettigrew and Martin, 2016; Rubin and Meiran, 2005) follow an extended maturational path, potentially continuing into adolescence (Cepeda et al., 2001; Crone et al., 2006a, 2004; Huizinga and van der Molen, 2007; Reimers and Maylor, 2005). One potential explanation for greater age differences in sustained than in transient control are concomitant differences in the developmental pace of underlying brain regions. In our task, several frontoparietal regions (i.e., the IFJ, SPL, and dACC) were recruited for both sustained and transient control; however, the dlPFC was exclusively activated under sustained control demands. Previous research has shown that the dlPFC is among the latest brain areas to mature during childhood and adolescence (Sowell et al., 1999; Sydnor et al., 2021; but see Fuhrmann et al., 2022). Critically, an entire network's efficiency can be hindered by a specific node, such that connectivity within a network is lower if a node of the network is lesioned, while networks that do not include this lesioned node are not affected in their efficiency (Nomura et al., 2010). Thus, while the regions associated with both transient and sustained control processes may be relatively more functional during transient control, they might be more strongly limited by the insufficient activation in the co-recruited dlPFC region during sustained control.

Additionally, connections within and between control networks continue to develop into adolescence (Fair et al., 2009, 2007; Luna et al., 2015; Marek et al., 2015), suggesting that the protracted development of sustained control processes might depend on the connectivity among regions associated with it. Age differences in structural connections between frontal and parietal regions have been shown to be relevant for task switching (Mackiewicz Seghete et al., 2013) and working memory (Østby et al., 2011; Vestergaard et

al., 2011; but see Krogsrud et al., 2018) during development. For example, task switching performance in children (10–11 years) but not adolescents (12–16 years) has been associated with greater white-matter integrity (i.e., fractional anisotropy) of the anterior corona radiata, a white-matter tract containing projections between and to many prefrontal regions (Mackiewicz Seghete et al., 2013). Thus, investigating gray-matter changes and white-matter structural connectivity differences of regions specifically involved in sustained and transient control during task switching might help distinguish how the maturation of connections within and between frontoparietal networks supports the development of these control processes.

We expected children to show smaller increases in connectivity with greater sustained control demands, reflecting the ongoing development of these task-related networks. We also found that children showed increased connectivity of the IFJ to regions that did not show reliable activation increases across children and adults in our task. One of these regions was a large PFC cluster with a lateral portion of the PFC, including the frontal pole/aPFC and middle frontal gyrus, and a medial portion, including the superior frontal gyrus and paracingulate gyrus. These results are consistent with a previous developmental connectivity study that also showed greater integration of the aPFC in a frontoparietal control network during task switching in children compared to adults (Ezekiel et al., 2013).

Interestingly, the lPFC and cognitive control functions associated with it have been suggested to follow a hierarchical organization, such that more posterior lPFC regions are associated with sensory-motor control, dorsolateral regions with contextual control, and the anterior regions with temporal control (Badre, 2008; Badre and D'Esposito, 2007; Badre and Nee, 2018; Nee and D'Esposito, 2016). The present connectivity results showing increased connectivity between the IFJ and a cluster stretching from posterior to anterior lPFC raises the question whether control hierarchies in the lPFC are less clearly established in children

than in adults (Bunge and Zelazo, 2006; Unger et al., 2016), such that children might recruit higher-order control regions for lower-order demands (cf. Crone and Steinbeis, 2017). This idea is consistent with the protracted differentiation of cognitive control process in development (e.g., Akshoomoff et al., 2018; Best and Miller, 2010; Brydges et al., 2014; Shing et al., 2010).

The lPFC in general, and the aPFC in particular, appears to support a variety of high-level cognitive processes that require the use of specific task rules (Bunge, 2004; Bunge and Zelazo, 2006; Burgess and Wu, 2012). According to these accounts of lPFC function, increased connectivity between the IFJ and the lPFC is likely to result from the increased demand to manage currently irrelevant task sets during mixed blocks. However, we found these increases only in children, not in adults. This raises the question whether children rely on increased connectivity to this region as an alternative strategy in situations with high sustained control demands. If so, proficient task-switching performance would depend on the IFJ–lPFC connectivity. Additionally, we examined whether this relationship is inversely related to the degree to which children succeed in upregulating activation in the frontoparietal regions associated with sustained control, as an indication of whether the alternative strategy is adaptive.

Our results revealed that the maturity of children's pattern of activation in response to sustained control demands moderated the relation between IFJ–lPFC connectivity and performance. More precisely, in children showing less adult-like activation patterns, increased IFJ–lPFC connectivity had a positive effect on performance, suggesting that upregulating lPFC involvement with increased sustained control demands functioned as a performance-enhancing alternate neural strategy in this group of children. These results are in line with findings on the role of the lPFC in rule maintenance and manipulation (e.g. Badre, 2012; Braver et al., 2003; Bunge, 2004), as lPFC activation and connectivity might support

the management (Lara and Wallis, 2015; Miller et al., 2018; Sala and Courtney, 2007) and

selection (Badre, 2012; Chatham and Badre, 2015; D'Ardenne et al., 2012) of multiple task-

set representations during mixed blocks, in particular when core task switching regions are

not yet sufficiently mature to become flexibly activated. The lPFC has been suggested to

allocate control resources based on superordinate goals (e.g., Eppinger et al., 2021; Ruel et

al., 2021), and to manage attention between internal information (e.g., maintained task-set

representations) and external (stimulus) information (Burgess et al., 2007). These processes

might be particularly important in task switching when internal task-set representations have

to be combined with information from the perceived stimuli to perform the task successfully.

This demand is especially enhanced during mixed blocks where the relevant task set changes

frequently and depends on external information, while the response rules have to be

maintained internally. This management process might be helpful to update task sets when

they are less clear or not available, as we would expect in children showing less adult-like

activation of frontoparietal regions.

In contrast, in children showing more adult-like activation in response to sustained

control demands, increased IFJ–lPFC connectivity in mixed blocks was negatively associated

with task performance. Interestingly, adults showed a similar brain-behavior relationship

(albeit not statistically significant), such that greater increases in connectivity were associated

with greater mixing costs. We surmise that once the core regions involved in task switching

are relatively more mature, involving additional peripheral regions may make its operation

less efficient. Indeed, under certain circumstances, increased control and lPFC involvement

can hinder learning and task performance, in particular during stimulus-driven responses and

sequence learning (Galea et al., 2010; Kruschke, 2003; Thompson-Schill et al., 2009).

Cognitive control follows a trajectory of increasing differentiation during childhood

development (Best and Miller, 2010; Brydges et al., 2014; Huizinga et al., 2006; Lee et al.,

2013; Shing et al., 2010). Similarly, throughout development the recruitment of the lPFC becomes increasingly specific (Chevalier et al., 2019; Dreher and Berman, 2002; Zhang et al., 2021), while the networks formed by different regions in the lPFC (i.e., the dlPFC, aPFC, dACC, IFJ) become increasingly segregated and specialized (Fair et al., 2009, 2007). Critically, these developmental changes in brain activation and connectivity might show non-linear and possibly non-monotonic relations to behavior that differ between and vary within individuals (Lautrey, 2003; Li and Lindenberger, 2002; Wendelken et al., 2017), further contributing to the complexity of developmental transitions in the neural machinery supporting task switching behavior. Longitudinal studies are needed to understand how brain structure, activation, and connectivity develop interactively with behavior to enable efficient task switching in the course of development. Such studies need to investigate how neural activation at one timepoint is related to connectivity at another, and importantly, how these longitudinal lead-lag relations, and individual differences therein, correlate with, and possibly predict changes in behavior.

Taken together, the presented data suggest that sustained and transient control processes during task switching follow different developmental trajectories, such that transient processes approximate adult levels relatively earlier than sustained processes do. A potential mechanism giving rise to children's task-switching difficulties might be insufficient upregulation of activation in response to increased control demands, especially for sustained control. However, increased connectivity between the IFJ and the lPFC with greater sustained control demands might offer an alternative, and possibly developmentally earlier mechanism to manage these demands, at least in some children. These findings point to a complex pattern of brain-behavior relationships during task switching in which children can cope with increased sustained control demands in at least two ways, either by upregulating activation in brain regions engaged by adults or by enhancing connectivity with brain areas involved in the

maintenance and management of multiple task sets. The longitudinal study of these patterns will reveal how these strategies shift in the course of development and how they are influenced by experience.

References

Akshoomoff, N., Brown, T.T., Bakeman, R., Hagler Jr., D.J., 2018. Developmental differentiation of executive functions on the NIH Toolbox Cognition Battery. Neuropsychology 32, 777–783. https://doi.org/10.1037/neu0000476

Allport, D.A., Styles, E.A., Hsieh, S., 1994. Shifting intentional set: Exploring the dynamic control of tasks, in: Umiltà, C., Moscovitch, M. (Eds.), Attention and Performance 15: Conscious and Nonconscious Information Processing. MIT Press, pp. 421–452.

Anderson, V.A., Anderson, P., Northam, E., Jacobs, R., Catroppa, C., 2001. Development of Executive Functions Through Late Childhood and Adolescence in an Australian Sample. Developmental Neuropsychology 20, 385–406. https://doi.org/10.1207/S15326942DN2001_5

Badre, D., 2012. Opening the gate to working memory. PNAS 109, 19878–19879. https://doi.org/10.1073/pnas.1216902109

Badre, D., 2008. Cognitive control, hierarchy, and the rostro–caudal organization of the frontal lobes. Trends in Cognitive Sciences 12, 193–200. https://doi.org/10.1016/j.tics.2008.02.004

Badre, D., D'Esposito, M., 2007. Functional Magnetic Resonance Imaging Evidence for a Hierarchical Organization of the Prefrontal Cortex. Journal of Cognitive Neuroscience 19, 2082–2099. https://doi.org/10.1162/jocn.2007.19.12.2082

Badre, D., Nee, D.E., 2018. Frontal Cortex and the Hierarchical Control of Behavior. Trends in Cognitive Sciences 22, 170–188. https://doi.org/10.1016/j.tics.2017.11.005

Bates, D., Maechler, M., Bolker, B., Walker, S., Christensen, R.H.B., Singmann, H., Dai, B., Scheipl, F., Grothendieck, G., Green, P., Fox, J., Bauer, A., Krivitsky, P.N., 2022. lme4: Linear Mixed-Effects Models using "Eigen" and S4. https://CRAN.R-project.org/package=lme4

Behzadi, Y., Restom, K., Liau, J., Liu, T.T., 2007. A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. NeuroImage 37, 90–101. https://doi.org/10.1016/j.neuroimage.2007.04.042

Best, J.R., Miller, P.H., 2010. A Developmental Perspective on Executive Function. Child Development 81, 1641–1660. https://doi.org/10.1111/j.1467-8624.2010.01499.x

Brass, M., Derrfuss, J., Forstmann, B., von Cramon, D.Y., 2005. The role of the inferior frontal junction area in cognitive control. Trends in Cognitive Sciences 9, 314–316. https://doi.org/10.1016/j.tics.2005.05.001

Brass, M., von Cramon, D.Y., 2004. Decomposing Components of Task Preparation with Functional Magnetic Resonance Imaging. Journal of Cognitive Neuroscience 16, 609–620. https://doi.org/10.1162/089892904323057335

Braver, T.S., 2012. The variable nature of cognitive control: a dual mechanisms framework. Trends in Cognitive Sciences 16, 106–113. https://doi.org/10.1016/j.tics.2011.12.010

Braver, T.S., Reynolds, J.R., Donaldson, D.I., 2003. Neural Mechanisms of Transient and Sustained Cognitive Control during Task Switching. Neuron 39, 713–726. https://doi.org/10.1016/S0896-6273(03)00466-5

Brett, M., Anton, J.-L., Valabregue, R., Poline, J.-B., 2002. Region of interest analysis using an SPM toolbox. Presented at the 8th International Conference on Functional Mapping of the Human Brain, June 2-6, 2002, Sendai, Japan. Available on CD-ROM in NeuroImage, Vol 16, No 2.

Brydges, C.R., Fox, A.M., Reid, C.L., Anderson, M., 2014. The differentiation of executive functions in middle and late childhood: A longitudinal latent-variable analysis. Intelligence 47, 34–43. https://doi.org/10.1016/j.intell.2014.08.010

Bunge, S.A., 2004. How we use rules to select actions: A review of evidence from cognitive neuroscience. Cognitive, Affective, & Behavioral Neuroscience 4, 564–579. https://doi.org/10.3758/CABN.4.4.564

Bunge, S.A., Kahn, I., Wallis, J.D., Miller, E.K., Wagner, A.D., 2003. Neural Circuits Subserving the Retrieval and Maintenance of Abstract Rules. Journal of Neurophysiology 90, 3419–3428. https://doi.org/10.1152/jn.00910.2002

Bunge, S.A., Wright, S.B., 2007. Neurodevelopmental changes in working memory and cognitive control. Current Opinion in Neurobiology, Cognitive neuroscience 17, 243–250. https://doi.org/10.1016/j.conb.2007.02.005

Bunge, S.A., Zelazo, P.D., 2006. A Brain-Based Account of the Development of Rule Use in Childhood. Curr Dir Psychol Sci 15, 118–121. https://doi.org/10.1111/j.0963-7214.2006.00419.x

Burgess, P.W., Dumontheil, I., Gilbert, S.J., 2007. The gateway hypothesis of rostral prefrontal cortex (area 10) function. Trends in Cognitive Sciences 11, 290–298. https://doi.org/10.1016/j.tics.2007.05.004

Burgess, P.W., Wu, H., 2012. Rostral prefrontal cortex (Brodmann area 10): metacognition in the brain, in: Stuss, D.T., Knight, R.T. (Eds.), Principles of Frontal Lobe Function. Oxford University Press, New York, pp. 524–544.

Cepeda, N.J., Kramer, A.F., Gonzalez de Sather, J.C.M., 2001. Changes in executive control across the life span: Examination of task-switching performance. Developmental Psychology 37, 715–730. https://doi.org/10.1037/0012-1649.37.5.715

Chatham, C.H., Badre, D., 2015. Multiple gates on working memory. Current Opinion in Behavioral Sciences 1, 23–31. https://doi.org/10.1016/j.cobeha.2014.08.001

Chevalier, N., Jackson, J., Revueltas Roux, A., Moriguchi, Y., Auyeung, B., 2019. Differentiation in prefrontal cortex recruitment during childhood: Evidence from cognitive control demands and social contexts. Developmental Cognitive Neuroscience 36, 100629. https://doi.org/10.1016/j.dcn.2019.100629

Church, J.A., Bunge, S.A., Petersen, S.E., Schlaggar, B.L., 2017. Preparatory Engagement of Cognitive Control Networks Increases Late in Childhood. Cereb Cortex 27, 2139–2153. https://doi.org/10.1093/cercor/bhw046

Cohen, J., 1988. Statistical power analysis for the behavioral sciences, 2nd ed. Lawrence Earlbaum Associates, Hillsdale, NJ.

Cox, R.W., Hyde, J.S., 1997. Software tools for analysis and visualization of fMRI data. NMR in Biomedicine 10, 171–178. https://doi.org/10.1002/(SICI)1099-1492(199706/08)10:4/5<171::AID-NBM453>3.0.CO;2-L

Cragg, L., Chevalier, N., 2012. The processes underlying flexibility in childhood. Quarterly Journal of Experimental Psychology 65, 209–232. https://doi.org/10.1080/17470210903204618

Crone, E.A., Bunge, S.A., Molen, M.W.V.D., Ridderinkhof, K.R., 2006a. Switching between tasks and responses: a developmental study. Developmental Science 9, 278–287. https://doi.org/10.1111/j.1467-7687.2006.00490.x

Crone, E.A., Donohue, S.E., Honomichl, R., Wendelken, C., Bunge, S.A., 2006b. Brain Regions Mediating Flexible Rule Use during Development. J. Neurosci. 26, 11239–11247. https://doi.org/10.1523/JNEUROSCI.2165-06.2006

Crone, E.A., Ridderinkhof, K.R., Worm, M., Somsen, R.J.M., van der Molen, M.W., 2004. Switching between spatial stimulus–response mappings: a developmental study of cognitive flexibility. Developmental Science 7, 443–455. https://doi.org/10.1111/j.1467-7687.2004.00365.x

Crone, E.A., Steinbeis, N., 2017. Neural Perspectives on Cognitive Control Development during Childhood and Adolescence. Trends in Cognitive Sciences 21, 205–215. https://doi.org/10.1016/j.tics.2017.01.003

Crone, E.A., Wendelken, C., Donohue, S., Leijenhorst, L. van, Bunge, S.A., 2006c. Neurocognitive development of the ability to manipulate information in working memory. PNAS 103, 9315–9320. https://doi.org/10.1073/pnas.0510088103

D'Ardenne, K., Eshel, N., Luka, J., Lenartowicz, A., Nystrom, L.E., Cohen, J.D., 2012. Role of prefrontal cortex and the midbrain dopamine system in working memory updating. PNAS 109, 19900–19909. https://doi.org/10.1073/pnas.1116727109

Dauvier, B., Chevalier, N., Blaye, A., 2012. Using finite mixture of GLMs to explore variability in children's flexibility in a task-switching paradigm. Cognitive Development 27, 440–454. https://doi.org/10.1016/j.cogdev.2012.07.004

Derrfuss, J., Brass, M., Neumann, J., von Cramon, D.Y., 2005. Involvement of the inferior frontal junction in cognitive control: Meta-analyses of switching and Stroop studies. Human Brain Mapping 25, 22–34. https://doi.org/10.1002/hbm.20127

Dosenbach, N.U.F., Fair, D.A., Cohen, A.L., Schlaggar, B.L., Petersen, S.E., 2008. A dual-networks architecture of top-down control. Trends in Cognitive Sciences 12, 99–105. https://doi.org/10.1016/j.tics.2008.01.001

Dosenbach, N.U.F., Fair, D.A., Miezin, F.M., Cohen, A.L., Wenger, K.K., Dosenbach, R.A.T., Fox, M.D., Snyder, A.Z., Vincent, J.L., Raichle, M.E., Schlaggar, B.L., Petersen, S.E., 2007. Distinct brain networks for adaptive and stable task control in humans. PNAS 104, 11073–11078. https://doi.org/10.1073/pnas.0704320104

Dosenbach, N.U.F., Koller, J.M., Earl, E.A., Miranda-Dominguez, O., Klein, R.L., Van, A.N., Snyder, A.Z., Nagel, B.J., Nigg, J.T., Nguyen, A.L., Wesevich, V., Greene, D.J., Fair, D.A., 2017. Real-time motion analytics during brain MRI improve data quality and reduce costs. NeuroImage 161, 80–93. https://doi.org/10.1016/j.neuroimage.2017.08.025

Dosenbach, N.U.F., Visscher, K.M., Palmer, E.D., Miezin, F.M., Wenger, K.K., Kang, H.C., Burgund, E.D., Grimes, A.L., Schlaggar, B.L., Petersen, S.E., 2006. A Core System for the Implementation of Task Sets. Neuron 50, 799–812. https://doi.org/10.1016/j.neuron.2006.04.031

Dreher, J.-C., Berman, K.F., 2002. Fractionating the neural substrate of cognitive control processes. PNAS 99, 14595–14600. https://doi.org/10.1073/pnas.222193299

Durston, S., Davidson, M.C., Tottenham, N., Galvan, A., Spicer, J., Fossella, J.A., Casey, B.J., 2006. A shift from diffuse to focal cortical activity with development. Developmental Science 9, 1–8. https://doi.org/10.1111/j.1467-7687.2005.00454.x

Düzel, E., Schütze, H., Yonelinas, A.P., Heinze, H.-J., 2011. Functional phenotyping of successful aging in long-term memory: Preserved performance in the absence of neural compensation. Hippocampus 21, 803–814. https://doi.org/10.1002/hipo.20834

Edelman, G.M., 1987. Neural Darwinism: The theory of neuronal group selection. Basic Books, New York, NY.

Engelhardt, L.E., Harden, K.P., Tucker-Drob, E.M., Church, J.A., 2019. The neural architecture of executive functions is established by middle childhood. NeuroImage 185, 479–489. https://doi.org/10.1016/j.neuroimage.2018.10.024

Eppinger, B., Goschke, T., Musslick, S., 2021. Meta-control: From psychology to computational neuroscience. Cogn Affect Behav Neurosci 21, 447–452. https://doi.org/10.3758/s13415-021-00919-4

Esteban, O., Markiewicz, C.J., Blair, R.W., Moodie, C.A., Isik, A.I., Erramuzpe, A., Kent, J.D., Goncalves, M., DuPre, E., Snyder, M., Oya, H., Ghosh, S.S., Wright, J., Durnez, J., Poldrack, R.A., Gorgolewski, K.J., 2019. fMRIPrep: a robust preprocessing pipeline for functional MRI. Nature Methods 16, 111–116. https://doi.org/10.1038/s41592-018-0235-4

Ezekiel, F., Bosma, R., Morton, J.B., 2013. Dimensional Change Card Sort performance associated with age-related differences in functional connectivity of lateral prefrontal cortex. Developmental Cognitive Neuroscience 5, 40–50. https://doi.org/10.1016/j.dcn.2012.12.001

Fair, D.A., Cohen, A.L., Power, J.D., Dosenbach, N.U.F., Church, J.A., Miezin, F.M., Schlaggar, B.L., Petersen, S.E., 2009. Functional Brain Networks Develop from a "Local to Distributed" Organization. PLOS Computational Biology 5, e1000381. https://doi.org/10.1371/journal.pcbi.1000381

Fair, D.A., Dosenbach, N.U.F., Church, J.A., Cohen, A.L., Brahmbhatt, S., Miezin, F.M., Barch, D.M., Raichle, M.E., Petersen, S.E., Schlaggar, B.L., 2007. Development of distinct control networks through segregation and integration. PNAS 104, 13507–13512. https://doi.org/10.1073/pnas.0705843104

Fandakova, Y., Lindenberger, U., Shing, Y.L., 2015. Maintenance of youth-like processing protects against false memory in later adulthood. Neurobiology of Aging 36, 933–941. https://doi.org/10.1016/j.neurobiolaging.2014.10.022

Fandakova, Y., Selmeczy, D., Leckey, S., Grimm, K.J., Wendelken, C., Bunge, S.A., Ghetti, S., 2017. Changes in ventromedial prefrontal and insular cortex support the development of metamemory from childhood into adolescence. PNAS 114, 7582–7587. https://doi.org/10.1073/pnas.1703079114

Fields, A., Bloom, P.A., VanTieghem, M., Harmon, C., Choy, T., Camacho, N.L., Gibson, L., Umbach, R., Heleniak, C., Tottenham, N., 2021. Adaptation in the face of adversity:

Decrements and enhancements in children's cognitive control behavior following early caregiving instability. Developmental Science 24, e13133. https://doi.org/10.1111/desc.13133

Fuhrmann, D., Madsen, K.S., Johansen, L.B., Baaré, W.F.C., Kievit, R.A., 2022. The midpoint of cortical thinning between late childhood and early adulthood differs between individuals and brain regions: Evidence from longitudinal modelling in a 12-wave neuroimaging sample. NeuroImage 261, 119507. https://doi.org/10.1016/j.neuroimage.2022.119507

Galea, J.M., Albert, N.B., Ditye, T., Miall, R.C., 2010. Disruption of the Dorsolateral Prefrontal Cortex Facilitates the Consolidation of Procedural Skills. Journal of Cognitive Neuroscience 22, 1158–1164. https://doi.org/10.1162/jocn.2009.21259

Gold, B.T., Powell, D.K., Xuan, L., Jicha, G.A., Smith, C.D., 2010. Age-related slowing of task switching is associated with decreased integrity of frontoparietal white matter. Neurobiology of Aging 31, 512–522. https://doi.org/10.1016/j.neurobiolaging.2008.04.005

Grayson, D.S., Fair, D.A., 2017. Development of large-scale functional networks from birth to adulthood: A guide to the neuroimaging literature. NeuroImage, Functional Architecture of the Brain 160, 15–31. https://doi.org/10.1016/j.neuroimage.2017.01.079

Greve, D.N., Fischl, B., 2009. Accurate and robust brain image alignment using boundary-based registration. NeuroImage 48, 63–72. https://doi.org/10.1016/j.neuroimage.2009.06.060

Gupta, R., Kar, B.R., Srinivasan, N., 2009. Development of task switching and post-error-slowing in children. Behavioral and Brain Functions 5, 38. https://doi.org/10.1186/1744-9081-5-38

Gurd, J.M., Amunts, K., Weiss, P.H., Zafiris, O., Zilles, K., Marshall, J.C., Fink, G.R., 2002. Posterior parietal cortex is implicated in continuous switching between verbal fluency tasks: an fMRI study with clinical implications. Brain 125, 1024–1038. https://doi.org/10.1093/brain/awf093

Hope, R.M., 2013. Rmisc: Ryan Miscellaneous. https://CRAN.R-project.org/package=Rmisc

Huizinga, M., Dolan, C.V., van der Molen, M.W., 2006. Age-related change in executive function: Developmental trends and a latent variable analysis. Neuropsychologia, Advances in Developmental Cognitive Neuroscience 44, 2017–2036. https://doi.org/10.1016/j.neuropsychologia.2006.01.010

Huizinga, M., van der Molen, M.W., 2007. Age-Group Differences in Set-Switching and Set-Maintenance on the Wisconsin Card Sorting Task. Developmental Neuropsychology 31, 193–215. https://doi.org/10.1080/87565640701190817

Jenkinson, M., Bannister, P., Brady, M., Smith, S., 2002. Improved Optimization for the Robust and Accurate Linear Registration and Motion Correction of Brain Images. NeuroImage 17, 825–841. https://doi.org/10.1006/nimg.2002.1132

Kassambara, A., 2020a. ggpubr: "ggplot2" Based Publication Ready Plots. R package version 0.4.0. https://CRAN.R-project.org/package=ggpubr

Kassambara, A., 2020b. rstatix: Pipe-Friendly Framework for Basic Statistical Tests. R package version 0.6.0. https://CRAN.R-project.org/package=rstatix

Kim, C., Cilles, S.E., Johnson, N.F., Gold, B.T., 2012. Domain general and domain preferential brain regions associated with different types of task switching: A Meta-Analysis. Human Brain Mapping 33, 130–142. https://doi.org/10.1002/hbm.21199

Koechlin, E., Corrado, G., Pietrini, P., Grafman, J., 2000. Dissociating the role of the medial and lateral anterior prefrontal cortex in human planning. PNAS 97, 7651–7656. https://doi.org/10.1073/pnas.130177397

Krogsrud, S.K., Fjell, A.M., Tamnes, C.K., Grydeland, H., Due-Tønnessen, P., Bjørnerud, A., Sampaio-Baptista, C., Andersson, J., Johansen-Berg, H., Walhovd, K.B., 2018. Development of white matter microstructure in relation to verbal and visuospatial working memory—A longitudinal study. PLOS ONE 13, e0195540. https://doi.org/10.1371/journal.pone.0195540

Kruschke, J.K., 2003. Attention in Learning. Curr Dir Psychol Sci 12, 171–175. https://doi.org/10.1111/1467-8721.01254

Kuznetsova, A., Brockhoff, P.B., Christensen, R.H.B., 2017. lmerTest Package: Tests in Linear Mixed Effects Models. J. Stat. Soft. 82. https://doi.org/10.18637/jss.v082.i13

Lara, A.H., Wallis, J.D., 2015. The Role of Prefrontal Cortex in Working Memory: A Mini Review. Frontiers in Systems Neuroscience 9. https://doi.org/ 10.3389/fnsys.2015.00173

Lautrey, Jacques, 2003. A pluralistic approach to cognitive differenciation and development, in: Sternberg, R.J., Lautrey, J., Lubart, T.I. (Eds.), Models of Intelligence: International Perspectives. American Psychological Association, Washington, DC, pp. 117–131.

Lee, K., Bull, R., Ho, R.M.H., 2013. Developmental Changes in Executive Functioning. Child Development 84, 1933–1953. https://doi.org/10.1111/cdev.12096

Lenth, R.V., Buerkner, P., Herve, M., Jung, M., Love, J., Miguez, F., Riebl, H., Singmann, H., 2022. emmeans: Estimated Marginal Means, aka Least-Squares Means. https://CRAN.R-project.org/package=emmeans

Li, S.-C., Lindenberger, U., 2002. Coconstructed functionality instead of functional normality. Behav Brain Sci 25, 761–762. https://doi.org/10.1017/S0140525X02330139

Luca, C.R.D., Wood, S.J., Anderson, V., Buchanan, J.-A., Proffitt, T.M., Mahony, K., Pantelis, C., 2003. Normative Data From the Cantab. I: Development of Executive Function Over the Lifespan. Journal of Clinical and Experimental Neuropsychology 25, 242–254. https://doi.org/10.1076/jcen.25.2.242.13639

Luciana, M., Conklin, H.M., Hooper, C.J., Yarger, R.S., 2005. The Development of Nonverbal Working Memory and Executive Control Processes in Adolescents. Child Development 76, 697–712. https://doi.org/10.1111/j.1467-8624.2005.00872.x

Lüdecke, D., Bartel, A., Schwemmer, C., Powell, C., Djalovski, A., Titz, J., 2021. sjPlot: Data Visualization for Statistics in Social Science. https://CRAN.R-project.org/package=sjPlot

Luna, B., Marek, S., Larsen, B., Tervo-Clemmens, B., Chahal, R., 2015. An Integrative Model of the Maturation of Cognitive Control. Annual Review of Neuroscience 38, 151–170. https://doi.org/10.1146/annurev-neuro-071714-034054

Mackiewicz Seghete, K.L., Herting, M.M., Nagel, B.J., 2013. White matter microstructure correlates of inhibition and task-switching in adolescents. Brain Research 1527, 15–28. https://doi.org/10.1016/j.brainres.2013.06.003

Makris, N., Goldstein, J.M., Kennedy, D., Hodge, S.M., Caviness, V.S., Faraone, S.V., Tsuang, M.T., Seidman, L.J., 2006. Decreased volume of left and total anterior insular lobule in schizophrenia. Schizophrenia Research 83, 155–171. https://doi.org/10.1016/j.schres.2005.11.020

Marek, S., Hwang, K., Foran, W., Hallquist, M.N., Luna, B., 2015. The Contribution of Network Organization and Integration to the Development of Cognitive Control. PLOS Biology 13, e1002328. https://doi.org/10.1371/journal.pbio.1002328

Mayr, U., Kliegl, R., 2000. Task-set switching and long-term memory retrieval. J Exp Psychol Learn Mem Cogn 26, 1124–1140. https://doi.org/10.1037//0278-7393.26.5.1124

McLaren, D.G., Ries, M.L., Xu, G., Johnson, S.C., 2012. A generalized form of context-dependent psychophysiological interactions (gPPI): A comparison to standard approaches. NeuroImage 61, 1277–1286. https://doi.org/10.1016/j.neuroimage.2012.03.068

Meiran, N., 2000. Reconfiguration of Stimulus Task Sets and Response Task Sets during Task Switching, in: Monsell, S., Driver, J. (Eds.), Control of Cognitive Processes: Attention and Performance XVIII. MIT Press, pp. 377–399.

Meiran, N., 1996. Reconfiguration of processing mode prior to task performance. Journal of Experimental Psychology: Learning, Memory, and Cognition 22, 1423–1442. https://doi.org/10.1037/0278-7393.22.6.1423

Miller, E.K., Lundqvist, M., Bastos, A.M., 2018. Working Memory 2.0. Neuron 100, 463–475. https://doi.org/10.1016/j.neuron.2018.09.023

Mills, K.L., Siegmund, K.D., Tamnes, C.K., Ferschmann, L., Wierenga, L.M., Bos, M.G.N., Luna, B., Li, C., Herting, M.M., 2021. Inter-individual variability in structural brain development from late childhood to young adulthood. NeuroImage 242, 118450. https://doi.org/10.1016/j.neuroimage.2021.118450

Miyake, A., Friedman, N.P., 2012. The Nature and Organization of Individual Differences in Executive Functions: Four General Conclusions. Current Directions in Psychological Science 21, 8–14. https://doi.org/10.1177/0963721411429458

Miyake, A., Friedman, N.P., Emerson, M.J., Witzki, A.H., Howerter, A., Wager, T.D., 2000. The Unity and Diversity of Executive Functions and Their Contributions to Complex "Frontal

Lobe" Tasks: A Latent Variable Analysis. Cognitive Psychology 41, 49–100. https://doi.org/10.1006/cogp.1999.0734

Morton, J.B., Bosma, R., Ansari, D., 2009. Age-related changes in brain activation associated with dimensional shifts of attention: An fMRI study. NeuroImage 46, 249–256. https://doi.org/10.1016/j.neuroimage.2009.01.037

Munakata, Y., Snyder, H.R., Chatham, C.H., 2012. Developing Cognitive Control: Three Key Transitions. Curr Dir Psychol Sci 21, 71–77. https://doi.org/10.1177/0963721412436807

Nee, D.E., D'Esposito, M., 2016. The hierarchical organization of the lateral prefrontal cortex. eLife 5, e12112. https://doi.org/10.7554/eLife.12112

Niendam, T.A., Laird, A.R., Ray, K.L., Dean, Y.M., Glahn, D.C., Carter, C.S., 2012. Meta-analytic evidence for a superordinate cognitive control network subserving diverse executive functions. Cogn Affect Behav Neurosci 12, 241–268. https://doi.org/10.3758/s13415-011-0083-5

Nomura, E.M., Gratton, C., Visser, R.M., Kayser, A., Perez, F., D'Esposito, M., 2010. Double dissociation of two cognitive control networks in patients with focal brain lesions. PNAS 107, 12017–12022. https://doi.org/10.1073/pnas.1002431107

Østby, Y., Tamnes, C.K., Fjell, A.M., Walhovd, K.B., 2011. Morphometry and connectivity of the fronto-parietal verbal working memory network in development. Neuropsychologia 49, 3854–3862. https://doi.org/10.1016/j.neuropsychologia.2011.10.001

Pettigrew, C., Martin, R.C., 2016. The role of working memory capacity and interference resolution mechanisms in task switching. Quarterly Journal of Experimental Psychology 69, 2431–2451. https://doi.org/10.1080/17470218.2015.1121282

Power, J.D., Barnes, K.A., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., 2012. Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. NeuroImage 59, 2142–2154. https://doi.org/10.1016/j.neuroimage.2011.10.018

R Core Team, 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing.

Reimers, S., Maylor, E.A., 2005. Task Switching Across the Life Span: Effects of Age on General and Specific Switch Costs. Developmental Psychology 41, 661–671. https://doi.org/10.1037/0012-1649.41.4.661

Richter, F., Yeung, N., 2014. Neuroimaging studies of task switching, in: Grange, J., Houghton, G. (Eds.), Task Switching and Cognitive Control. Oxford University Press, pp. 237–271.

Rogers, R.D., Monsell, S., 1995. The costs of a predictable switch between simple cognitive tasks. Journal of Experimental Psychology: General 124, 207–231. https://doi.org/10.1037/0096-3445.124.2.207

Rubin, O., Meiran, N., 2005. On the Origins of the Task Mixing Cost in the Cuing Task-Switching Paradigm. Journal of Experimental Psychology: Learning, Memory, and Cognition 31, 1477–1491. https://doi.org/10.1037/0278-7393.31.6.1477

Ruel, A., Devine, S., Eppinger, B., 2021. Resource-rational approach to meta-control problems across the lifespan. WIREs Cognitive Science 12, e1556. https://doi.org/10.1002/wcs.1556

Sala, J.B., Courtney, S.M., 2007. Binding of What and Where During Working Memory Maintenance. Cortex 43, 5–21. https://doi.org/10.1016/S0010-9452(08)70442-8

Satterthwaite, T.D., Wolf, D.H., Erus, G., Ruparel, K., Elliott, M.A., Gennatas, E.D., Hopson, R., Jackson, C., Prabhakaran, K., Bilker, W.B., Calkins, M.E., Loughead, J., Smith, A., Roalf, D.R., Hakonarson, H., Verma, R., Davatzikos, C., Gur, R.C., Gur, R.E., 2013. Functional Maturation of the Executive System during Adolescence. J. Neurosci. 33, 16249–16261. https://doi.org/10.1523/JNEUROSCI.2345-13.2013

Schuch, S., Konrad, K., 2017. Investigating task inhibition in children versus adults: A diffusion model analysis. Journal of Experimental Child Psychology 156, 143–167. https://doi.org/10.1016/j.jecp.2016.11.012

Shing, Y.L., Lindenberger, U., Diamond, A., Li, S.-C., Davidson, M.C., 2010. Memory maintenance and inhibitory control differentiate from early childhood to adolescence. Dev Neuropsychol 35, 679–697. https://doi.org/10.1080/87565641.2010.508546

Sowell, E.R., Thompson, P.M., Holmes, C.J., Jernigan, T.L., Toga, A.W., 1999. In vivo evidence for post-adolescent brain maturation in frontal and striatal regions. Nat Neurosci 2, 859–861. https://doi.org/10.1038/13154

Sydnor, V.J., Larsen, B., Bassett, D.S., Alexander-Bloch, A., Fair, D.A., Liston, C., Mackey, A.P., Milham, M.P., Pines, A., Roalf, D.R., Seidlitz, J., Xu, T., Raznahan, A., Satterthwaite, T.D., 2021. Neurodevelopment of the association cortices: Patterns, mechanisms, and implications for psychopathology. Neuron 109, 2820–2846. https://doi.org/10.1016/j.neuron.2021.06.016

Tamnes, C.K., Walhovd, K.B., Torstveit, M., Sells, V.T., Fjell, A.M., 2013. Performance monitoring in children and adolescents: A review of developmental changes in the error-related negativity and brain maturation. Developmental Cognitive Neuroscience 6, 1–13. https://doi.org/10.1016/j.dcn.2013.05.001

Thompson-Schill, S.L., Ramscar, M., Chrysikou, E.G., 2009. Cognition Without Control: When a Little Frontal Lobe Goes a Long Way. Curr Dir Psychol Sci 18, 259–263. https://doi.org/10.1111/j.1467-8721.2009.01648.x

Unger, K., Ackerman, L., Chatham, C.H., Amso, D., Badre, D., 2016. Working memory gating mechanisms explain developmental change in rule-guided behavior. Cognition 155, 8–22. https://doi.org/10.1016/j.cognition.2016.05.020

Van der Maas, H.L., Molenaar, P.C., 1992. Stagewise cognitive development: An application of catastrophe theory. Psychological Review 99, 395–417. https://doi.org/10.1037/0033-295X.99.3.395

Velanova, K., Wheeler, M.E., Luna, B., 2008. Maturational Changes in Anterior Cingulate and Frontoparietal Recruitment Support the Development of Error Processing and Inhibitory Control. Cereb Cortex 18, 2505–2522. https://doi.org/10.1093/cercor/bhn012

Vestergaard, M., Madsen, K.S., Baaré, W.F.C., Skimminge, A., Ejersbo, L.R., Ramsøy, T.Z., Gerlach, C., Åkeson, P., Paulson, O.B., Jernigan, T.L., 2011. White Matter Microstructure in Superior Longitudinal Fasciculus Associated with Spatial Working Memory Performance in Children. Journal of Cognitive Neuroscience 23, 2135–2146. https://doi.org/10.1162/jocn.2010.21592

Wendelken, C., Ferrer, E., Ghetti, S., Bailey, S.K., Cutting, L., Bunge, S.A., 2017. Frontoparietal Structural Connectivity in Childhood Predicts Development of Functional Connectivity and Reasoning Ability: A Large-Scale Longitudinal Investigation. J. Neurosci. 37, 8549–8558. https://doi.org/10.1523/JNEUROSCI.3726-16.2017

Wendelken, C., Munakata, Y., Baym, C., Souza, M., Bunge, S.A., 2012. Flexible rule use: Common neural substrates in children and adults. Developmental Cognitive Neuroscience 2, 329–339. https://doi.org/10.1016/j.dcn.2012.02.001

Whitfield-Gabrieli, S., Nieto-Castanon, A., 2012. Conn : A Functional Connectivity Toolbox for Correlated and Anticorrelated Brain Networks. Brain Connectivity 2, 125–41. https://doi.org/10.1089/brain.2012.0073

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L.D., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T.L., Miller, E., Bache, S.M., Müller, K., Ooms, J., Robinson, D., Seidel, D.P., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo, K., Yutani, H., 2019. Welcome to the Tidyverse. Journal of Open Source Software 4, 1686. https://doi.org/10.21105/joss.01686

Worringer, B., Langner, R., Koch, I., Eickhoff, S.B., Eickhoff, C.R., Binkofski, F.C., 2019. Common and distinct neural correlates of dual-tasking and task-switching: a meta-analytic review and a neuro-cognitive processing model of human multitasking. Brain Struct Funct 224, 1845–1869. https://doi.org/10.1007/s00429-019-01870-4

Wylie, G., Allport, A., 2000. Task switching and the measurement of "switch costs." Psychological Research 63, 212–233. https://doi.org/10.1007/s004269900003

Zhang, Z., Peng, P., Eickhoff, S.B., Lin, X., Zhang, D., Wang, Y., 2021. Neural substrates of the executive function construct, age-related changes, and task materials in adolescents and adults: ALE meta-analyses of 408 fMRI studies. Developmental Science 24, e13111. https://doi.org/10.1111/desc.13111