Slow neural oscillations explain temporal fluctuations in distractibility

Troby Ka-Yan Lui[1,2], Jonas Obleser[1,2], & Malte Wöstmann[1,2]

1. Department of Psychology, University of Lübeck, Ratzeburger Allee 160, 23562 Lübeck, Germany
2. Center of Brain, Behavior and Metabolism, University of Lübeck, Ratzeburger Allee 160, 23562 Lübeck, Germany

Corresponding author

Troby Ka-Yan Lui (kayan.lui@uni-luebeck.de)

Competing interests

The authors declare no conflict of interest.

Data availability

Raw data are available from the corresponding authors upon request. Pre-processed behavioural and neural responses as a function of distractor onset time will be made available online upon publication.

Author contributions

Conceptualization, T.K.L., J.O., and M.W.; methodology, T.K.L., J.O., and M.W.; investigation, T.K.L. ; formal analysis, T.K.L., J.O., and M.W. ; writing – original draft, T.K.L. and M.W.; writing – review & editing T.K.L., J.O., and M.W.; funding acquisition, M.W.

Funding

This work was supported by Deutsche Forschungsgemeinschaft [grant number WO 2371/1-1].

Keywords

attention; auditory; distractibility; distraction; rhythmic cognition; theta oscillations

Abstract

Human environments comprise various sources of distraction, which often occur unexpectedly in time. The proneness to distraction (i.e., distractibility) is posited to be independent of attentional sampling of targets, but its temporal dynamics and neurobiological basis are largely unknown. Brain oscillations in the theta band (3–8 Hz) have been associated with fluctuating neural excitability, which is hypothesised here to explain rhythmic modulation of distractibility. In a pitch discrimination task (N = 30) with unexpected auditory distractors, we show that distractor-evoked neural responses in the electroencephalogram and perceptual susceptibility to distraction were co-modulated and cycled approximately 3–5 times per second. Pre-distractor neural phase in left inferior frontal and insular cortex regions explained fluctuating distractibility. Thus, human distractibility is not constant but fluctuates on a subsecond timescale. Furthermore, slow neural oscillations subserve the behavioural consequences of a hitherto largely unexplained but ever-increasing phenomenon in modern environments – distraction by unexpected sound.

Introduction

Selective attention enables humans to focus on relevant information at the expense of distraction. The brain prioritizes representations of relevant events while filtering out task-irrelevant distractors (Desimone & Duncan, 1995; Picton et al., 1971). Recent research posited that distractor processing is not merely collateral to attentional sampling of targets but may follow its own dynamics (Schneider et al., 2018; Wöstmann et al., 2019, 2020). The behavioural detriments induced by different kinds of distractors (i.e., *distraction*) and the neuro-cognitive mechanisms that counteract distraction (i.e., *suppression*) have been studied in some detail (Bonnefond & Jensen, 2012; Geng & DiQuattro, 2010; van Moorselaar et al., 2020; Weisz et al., 2020; Wöstmann et al., 2019). However, the temporal dynamics and the neurobiological basis of the proneness to distraction (i.e., *distractibility*) are largely unknown.

Distractibility has long been neglected in the theoretical formulation of rhythmic attention. Originally assumed to be static (Posner et al., 1980), the attentional spotlight was proposed to be blinking at a subsecond time scale in a theta-like rhythm (i.e., 3–8 Hz) (Buschman & Kastner, 2015; Fiebelkorn & Kastner, 2019). Behaviourally, it is manifested via the waxing and waning of behavioural performance in target selection (Fiebelkorn et al., 2013; Ho et al., 2017; Kubetschek & Kayser, 2021; Landau & Fries, 2012) or working memory (Schmid et al., 2022; ter Wal et al., 2021) performance at similar frequencies. However, the temporal dynamics outside of the attentional spotlight are not well understood. While previous research studied how distractibility unfolds on relatively long temporal scales of minutes (i.e., during an experimental session (Forster & Lavie, 2014)) or years (i.e., across stages of development (Campbell et al., 2012; Kannass et al., 2006)), we found preliminary evidence for fluctuating distractibility on shorter timescales following rhythmic presentation of auditory targets (Wöstmann et al., 2020). To isolate distractibility dynamics from rhythmic entrainment or preparatory suppression, we here employ a design that uses non-rhythmic stimuli and distractors that occur unexpectedly.

A central prediction of rhythmic attention is that the phase of slow neural oscillations explains fluctuations in behaviour (VanRullen, 2016). The prediction is based on the notion that rhythmic attention arises from the periodic excitability of the attention-related brain network (Fiebelkorn & Kastner, 2019; VanRullen, 2016). In the human brain, theta neural phase (3–8 Hz) is assumed to reflect moment-to-moment changes in neural excitability (Lakatos et al., 2005). Theta phase in brain regions beyond sensory cortices, such as fronto-

parietal regions and the hippocampus, has been associated with fluctuations in target detection (Helfrich et al., 2018) and working memory encoding (Rutishauser et al., 2010; Siegel et al., 2009), respectively. While previous research has related distractibility to supra-modal regions in frontal (Chao & Knight, 1995; Wais et al., 2012) or parietal (Kanai et al., 2011) cortex, it is unclear whether and in which networks the momentary neural dynamics may subserve the waxing and waning of distractibility.

Here, we ask if the brain spontaneously alternates between states of higher and lower distractibility and whether such fluctuations have the potency to explain behavioural consequences of distraction. If so, we would expect to observe a brain-behaviour relation between the pre-distractor brain state and the distractor-induced detriment in task performance. To this end, we employed a pitch discrimination task wherein an auditory distractor could occur at variable and unexpected times in-between two target tones. A total of 17,280 behavioural and neural responses in the electroencephalogram (EEG) in N=30 participants revealed that behavioural sensitivity and distractor-evoked neural responses fluctuated in sync across distractor onset times in ~3–5 cycles per second. Critically, pre-distractor theta phase in left inferior frontal and insular cortex regions explained behavioural performance fluctuations. These effects were absent in trials without distractors, reinforcing their specificity to distractor-related neural processing.

                                Material and Methods

    Participants

Thirty participants (20 females, 10 males; mean age = 23.67, SD = 3.56) took part in the EEG

experiment. They provided written informed consent and were compensated by either €10/hour

or course credit. Participants were right-handed according to the Edinburgh Handedness

Inventory (Oldfield, 1971) (mean score = 92), with self-reported normal hearing, normal or

corrected-to-normal vision, and no psychological or neurological disorders. All procedures of

the current study were approved by the ethics committee of the University of Lübeck.

Stimuli and Procedure

Participants performed a pitch discrimination task wherein they decided whether the first (tone

1) and the second (tone 2) target tones in a trial were the same or different in pitch. Prior to the

experiment, they were instructed to answer as accurately and as fast as possible. The target

tones were 75 ms long pure tones with 5 ms rise and fall periods. In each trial, the frequencies

of tone 1 were randomly selected between musical note A#3 (233 Hz) and G#5 (830.6 Hz),

while that of tone 2 was either the same (50%) or different (higher or lower, 25% each) in

frequency compared to tone 1.

The pitch difference between tone 1 and tone 2 was titrated for each participant with an

adaptive task (see below). The offset-to-onset interval between tone 1 and tone 2 was 1550 ms.

Each distractor stimulus comprised 10 consecutive pure tones with 40 ms duration (400 ms in

total). The frequencies of the pure tones in each distractor stimulus were randomly selected

among the 12 tones between A#3 and G#5 with whole tone steps (A#3, C4, D4, E4, F#4, G#4,

A#4, C5, D5, E5, F#5, and G#5), with the constraint that there would be no repetition between

consecutive tones. Each of the 12 tone frequencies appeared at each of the 10 positions with

equal probability across trials.

In-between the two target tones, a distractor was presented in 50% of trials (distractor-

present condition) and no distractor was presented in the remaining trials (distractor-absent

condition). In the distractor-present condition, the distractor was presented at one of 24

distractor onset times (0 ms to 1150 ms, 50-ms steps, relative to the offset of tone 1), which

was selected at random on each trial. After the offset of target tone 2, participants had a 2000

ms response time window. To avoid potential temporal predictability effects of the onset of the

next trial, the inter-trial intervals were randomly selected from a truncated exponential

distribution (mean = 1460 ms), ranging between 730 and 3270 ms.

The trial order was pseudo-randomized with no repetition in probe tone frequency and

distractor onset for any two consecutive trials. In total, there were 12 trials for each unique

condition (distractor-present/absent x distractor onset x same/different target pitch) and 1152

trials for the whole experiment. All auditory materials were presented via Sennheiser

headphones (HD 25-1 II). Responses were made using a response box (The Black Box Toolkit).

The assignment of buttons to the response options ("same" or "different") was counterbalanced

across participants. Stimuli were presented via Matlab (MathWorks, Inc., Natick, USA) and

Psychtoolbox(Brainard, 1997). The auditory stimuli were presented at approximately 70 dB

SPL.

Adaptive Staircase Procedure

Prior to the main experiment, each participant's threshold for the pitch discrimination task was

titrated using an adaptive staircase procedure, implemented in the Palamedes toolbox (Prins &

Kingdom, 2018) for Matlab. For the initial 11 participants, the threshold was titrated to an

approximate accuracy of 70.7%. As the overall accuracy was relatively high even after the

adaptive staircase procedure for these 11 participants (mean = 79.59%, SD = 10.43%), the final

16 participants performed an adaptive procedure altered to yield approximately 65% accuracy

instead. Due to technical issues, performance of the remaining three participants was tracked

at 35% accuracy. As all relevant statistical analyses in the present study are within-subject, and

as paired t-tests (2-tailed) comparing the behavioural performance between distractor-absent

and distractor-present conditions were significant with (t_{29} = 8.11, p < .001) and without (t_{26} =

9.41, p < .001) these participants, their data were included in the final analysis.

Each participant went through the adaptive staircase procedure two to three times,

depending on the stability of the tracked threshold. There were in total 30 trials for each run of

the adaptive staircase procedure with an initial pitch difference of 100 cents (i.e. 1 semitone)

between tone 1 and 2. The minimum and maximum pitch difference possible in the task was 2

cents and 2000 cents, respectively. For the procedure which tracked performance at ~70.7%, a

two-down one-up procedure was used. Specifically, the pitch differences would decrease in

steps of 10 cents if participant responded correctly (i.e., different), or increase in steps of 10

cents if participant responded incorrectly (i.e., same) for 2 consecutive trials. For the procedure

which tracked performance at ~65% procedure, the pitch differences would decrease in steps of 7 cents if participant answered correctly or increase in steps of 13 cents if they answered incorrectly. The pitch difference used in the main experiment was calculated by averaging the final 10 trials in the tracking run which converged to the most stable threshold, determined by visual inspection, in the ~70.7% procedure. The same procedure was used to average the final 6 trials in the ~65% procedure. Overall accuracy averaged across all participants in the actual experiment was 73.58% (SD = 12.12%).

Behavioural Data Analysis

To understand how distractors affect pitch discrimination performance in the framework of signal detection theory, we calculated sensitivity (d') and criterion (c) separately for distractor-present and -absent conditions, using the Palamedes toolbox (Prins & Kingdom, 2018) and the following formulas:

(Formula 1)   Sensitivity = z (Hit rate) – z (False alarm rate)

(Formula 2)   Criterion = –0.5 * (z (Hit rate) + z (False alarm rate))

Hit rate was defined as the "different" response when the two tones were different in pitch, and false alarm rate the "different" response when the two tones were the same in pitch. Extreme values (0 or 1) of Hit rate or False alarm rate were adjusted (Macmillan & Kaplan, 1985): A rate of 0 was adjusted by dividing 1 by the number of trials multiplied by 2; while a value of 1 was adjusted by subtracting the same value from 1. Paired samples t-tests (2-tailed) were used to compare sensitivity and criterion in distractor-present versus -absent conditions.

To study the modulation of distractor onset times on behavioural measures in the distractor-present condition, sensitivity for each distractor onset time was calculated, resulting in a behavioural time course as a function of distractor onset time for each individual participant (see Fig. S1 & S2).

EEG Recording and Pre-processing

The experiment was conducted in an electrically shielded sound-attenuated room. A modified 10-20 international system with 64 Ag/Ag-Cl electrodes was used to record the EEG with a sampling rate of 1000 Hz (actiCHamp, Brain Products, München, Germany). The EEG recordings were band-pass filtered online from direct current (DC) to 280 Hz. TP9 was used

as the online reference and FPz as the ground electrode. Impedances were kept below 20 kOhm for all but one participant.

Matlab R2018a (MathWorks, Inc., Natick, USA) and the Fieldtrip toolbox (Oostenveld et al., 2011) were used to pre-process and analyse EEG data. The continuous EEG data were filtered (high-pass, 1 Hz; low-pass, 100 Hz) before they were segmented into epochs (-2 to 2.5s) time-locked to tone 1 onset. Independent component analysis (ICA) was used to identify and reject components corresponding to artefacts such as eye blinks, eye movements, and muscle activity (average percentage of components removed = 26.46%, SD = 8.89%). Afterwards, EEG data were re-referenced to the average of all electrodes. Epochs with amplitude changes >160 microvolts were rejected (average percentage of epochs removed = 1.35%, SD = 2%).

To obtain distractor-evoked neural responses, data were re-epoched to the onset of the distractor (-1 to 1 s) with a 200ms baseline period. Epochs belonging to the same conditions (distractor-present/absent) and distractor onset time (0 – 1150ms, 50-ms steps) were then averaged into ERP waveforms. The spectral amplitude of distractor-evoked responses at 25 Hz, which corresponds to the temporal structure of the distractor, was extracted using FFT on the ERP waveform in the time window from 0 to 520ms after distractor onset. Spectral amplitude was averaged across electrodes F1, Fz, F2, FC1, FCz, and FC2. For each participant, the 24 spectral amplitudes, corresponding to the 24 distractor onset times, resulted in a neural time course of distractor processing as a function of distractor onset time (see Fig. S1 & S2).

Distractor-evoked inter-trial phase coherence (ITPC) was also calculated across frequencies (1 – 10 Hz, 1-Hz steps) and time windows (-0.2 – 0.7 s, 0.05-s steps) for each electrode. First, Fourier coefficients were calculated (using windows with a fixed length of 0.5 s; hanning taper). Then, the complex Fourier coefficients were divided by their magnitude and averaged across trials. ITPC was calculated by taking the absolute value (i.e., magnitude) of the average complex coefficient.

Modulation of neural and behavioural measures by distractor onset time

To test whether and how distractor onset time modulates neural and behavioural measures, we used linear mixed-effect models with sine- and cosine-transformed distractor onset time, similar to Wöstmann et al. 2020 (Wöstmann et al., 2020). For time courses of sensitivity and spectral amplitude of the distractor-evoked ERP at 25 Hz separately, we first subtracted the

individually fitted quadratic trend (computed with the polyfit function in Matlab) from the original time course for each participant (see Fig. S1 & S2) as the quadratic trend was not of interest in the current study (Huang et al., 2015).

Then, we designed sine- and cosine- transformed distractor onset time vectors using the following formulas,

(Formula 3)   Sine predictor = sin $(2 * \pi * f *$ distractor onset time$)$

(Formula 4)   Cosine predictor = cos $(2 * \pi * f *$ distractor onset time$)$

Where f denotes the frequency of interest (0.5 – 8 Hz, 0.5-Hz steps). Next, we regressed the detrended sensitivity and spectral amplitude of ERP time courses on sine and cosine predictors using linear mixed models (using the fitlme function in Matlab) for each frequency of interest using the following formulas:

(Formula 5)   z(sensitivity) ~ z(sine predictor) + z(cosine predictor) + (1|participant)

(Formula 6)   z(25-Hz ERP) ~ z(sine predictor) + z(cosine predictor) + (1|participant)

Spectral magnitude for each frequency was computed by taking the square root of the sum of squared beta coefficients of sine and cosine predictors:

(Formula 7)   Spectral magnitude = sqrt (sine coef^2 + cosine coef^2)

Statistical significance of the spectral magnitude was determined by comparing the spectral magnitude of the empirical data with the 95[th] percentile of a permutation distribution, which was generated by shuffling the original behavioural/neural time course and performing the same analysis for 5,000 times.

To test whether sensitivity and spectral amplitude of the distractor-evoked ERP at 25 Hz are co-modulated, for each participant, cross-correlation coefficients across time lags of the two signals were obtained (using the "xcorr" function on z-scored time courses in Matlab). Again, we ran a similar linear mixed model as explained above, but this time with sine- and cosine- transformed time lags as predictors and used the correlation coefficients from the cross-correlation as the outcome measure. Spectral magnitude was obtained using formula 7 and statistical significance with the same permutation method mentioned above.

Phasic modulation of behavioural sensitivity

To explore the role of pre-distractor neural dynamics on the pitch discrimination performance, we examined whether pre-distractor oscillatory phase relates to behavioural sensitivity. To this end, we examined the quadratic fit of sensitivity as a function of neural phase in source space.

First, we implemented the source analysis using the Fieldtrip toolbox. First, a standard volume conduction model and standard electrode locations were used to calculate the leadfield matrix with 10-mm resolution. We applied the linearly constrained minimum variance (Van Veen et al., 1997) (LCMV) beamformer approach on the 10 Hz lowpass filtered data centred around distractor onset (-1 to 1s). We calculated a common filter including all trials by calculating the covariance matrix estimates. There were in total 2,015 source locations inside the brain.

Second, a quadratic fit analysis resolved by frequency and time probed the spectral and temporal specificity of the phasic modulation of perceptual sensitivity. To obtain trial-wise phase values for each source location, the following procedure was implemented for each trial in each source location: First, the single-trial EEG time course was projected into the source space using the common filter. Then, a sliding window (0.4s duration; moving in 50-ms steps from –0.3 to +0.3s relative to distractor onset) was employed to transform the data into the frequency domain (using FFT). Note that the time point of the sliding window refers to the mid-point of each time window. For instance, the time window centred at -0.3 included data from -0.5 to -0.1 s. The respective phase value of each frequency (2.5 – 8 Hz in 0.5-Hz steps) was then calculated using the *angle* function in MATLAB. The phase values of all trials were binned into 9 bins of equal size, ranging from -pi to pi, followed by a calculation of sensitivity for each bin. The quadratic fit of sensitivity across phase bins was estimated using the *polyfit* function (order = 2) in MATLAB. As a result, we obtained a quadratic fit index for each source location, frequency, and time of interest.

We used a source-level cluster-based permutation test (Maris & Oostenveld, 2007) to find significant clusters in voxel-frequency-time space that would exhibit phasic modulation of sensitivity. Dependent-samples t-tests were used to contrast quadratic fit coefficients against zero, followed by clustering of adjacent bins with significant effects in voxel-frequency-time space. To derive cluster p-values, summed t-values in observed clusters were tested against 5,000 permutations with shuffled condition labels (two-tailed).

To demonstrate that the significant cluster found in the above analysis does not primarily originate from auditory cortex, we localised, for comparison, the distractor-evoked inter-trial phase coherence (ITPC) at 3 – 7 Hz, strongly assumed to emerge at least to large degrees from the supratemporal plane and auditory cortex (Koerner & Zhang, 2015; Mayhew et al., 2010; Oya et al., 2018), with the following procedure for each voxel: For each trial, we projected the time series EEG data into source space using the same common filter as in the analysis on the phasic relationship with behaviour. Then, we transformed the source-projected data (0 – 300 ms after distractor onset) to the frequency domain using FFT. The same calculation as on the sensor level was used to calculate the ITPC for each frequency. ITPC across frequencies 3 – 7 Hz were then averaged to obtain one distractor-evoked ITPC value for each   voxel.

                                                Results

In the current electroencephalography (EEG) and behavioural study, we aimed at (1)

uncovering the temporal fluctuations in distraction, and (2) exploring the relationship between

such fluctuations and momentary neural phase at similar frequencies. To this end, we varied

the onset time of an auditory distractor that was presented in-between two to-be-compared

tones in a variant of a pitch discrimination task.

We probed this research question in the auditory modality as temporal information is

especially important to auditory attentional selection (Shamma et al., 2011). During the task,

participants (N = 30) had to identify whether the two target tones were the same or different in

pitch (Fig 1A). The distractor was a fast-varying, 25-Hz modulated sequence of tones that

differed in pitch, which allowed us to extract its induced 25-Hz neural response (Ding & Simon,

2009).

Distractors were present in half of the trials and absent in the remaining trials. In

distractor-present trials, the distractor onset was uniformly distributed across 24 onset times (0

– 1.15 s, in 0.05 s steps, after tone 1 offset). In distractor-absent trials, no distractor was

presented between the two tones. The inclusion of distractor-absent trials serves two purposes.

First, we could verify that the distractors had the potency to distract by comparing behavioural

performance for distractor-present versus distractor-absent trials (Wöstmann et al., 2022).

Second, participants could not anticipate whether or when a distractor would occur in a given

trial, which eliminated potential effects of such anticipation on behavioural performance

(Grabenhorst et al., 2021) or pre-stimulus neural activity (Dürschmid et al., n.d.; Herbst et al.,

2022; Stefanics et al., 2010).

Distractors interfere with pitch discrimination performance

To examine the potency of the distractors to distract, we compared participants' sensitivity and

criterion (response bias) of the pitch discrimination task between distractor-present and -absent

trials. Participants were less sensitive to the pitch difference (t_{29} = -8.11, p = <.001, Cohen's d

= -1.48), and had a more conservative response criterion (i.e., more "same pitch" responses; t_{29}

= 2.83, p = .008, Cohen's d = 0.52) on distractor-present trials (Fig 1B).

Fig 1. Experimental design and behavioural results.

A) Schematic of a distractor-present trial. Participants were instructed to indicate whether the two target tones (grey) were
the same (probability = 50%) or different (higher, probability = 25%; or lower, probability = 25%) in pitch. A 10-tone-pip
distractor sequence (white) with a 25-Hz temporal structure (i.e., 40-ms tone-pip duration; total duration 400 ms) was presented
at one of the 24 distractor onset times (dashed lines). In distractor-absent trials, no distractor was presented. B) Behavioural
results comparing distractor-present and -absent conditions. Coloured circles indicate single-subject data. Insets show bar
graphs of perceptual sensitivity (left panel) and criterion (right panel) for distractor-present (solid bar) and distractor-absent
(gradient bar) conditions, respectively. Error bars show ±1 SEM. ** p < .01. *** p < .001.

Behavioural and neural measures of distraction co-fluctuate across time

Does the impact of distraction on neural activity and goal-directed behaviour exhibit fluctuations across time? To test this, we varied distractor onset time and examined whether behavioural and neural measures of distraction would show modulations at frequencies up to 8 Hz. Behaviourally, perceptual sensitivity was calculated as an indirect measure of distraction: The more distracted, the lower the sensitivity in pitch discrimination should be (Fig 2A, yellow, see S1 Fig for individual participants' time courses). Neurally, we calculated the distractor-evoked event-related potential (ERP; Fig 2B) for each distractor onset time and used a fast Fourier transform (FFT) to extract its amplitude at 25 Hz, which corresponded to the modulation rate of the frequency-modulated distractor tone sequence (Fig 2A, blue, see S1 Fig for individual participants' time courses).

To examine temporal fluctuations of distraction, we used linear mixed-effects models with sine- and cosine- transformed distractor onset time as predictors to model behavioural (i.e., perceptual sensitivity) and neural (i.e., distractor-evoked ERP) time courses as the outcome measures. This method outperforms other methods for studying the phasic modulation of behavioural and neural responses (Zoefel et al., 2019) and has also been used previously (Wöstmann et al., 2020) to extract temporal fluctuations in the vulnerability of working memory to distraction. A quadratic trend was observed in the behavioural time course in Fig 2A as the earliest and latest distractors were most distracting due to their temporal proximity to the target tones. Before running linear mixed models, we removed the quadratic trends in the two measures as they were not of interest in the current study (Huang et al., 2015).

Fig 2E and F show the spectral magnitude (0.5–8 Hz, 0.5-Hz steps) resulting from linear mixed models on detrended perceptual sensitivity (Fig 2C, yellow) and detrended ERP amplitude (Fig 2C, blue), respectively. Statistical significance was derived by testing empirical spectral magnitude against the 95th percentile of a permutation distribution, which was derived from shuffling the behavioural and neural time courses, respectively, 5,000 times (see Methods for details).

At the behavioural level, distractor onset time modulated sensitivity below 5 Hz. At the neural level, distractor onset time modulated the distractor-evoked ERP at 4 and 5 Hz. Similar results were obtained in a control analysis, where temporal fluctuations in sensitivity in distractor-present trials were compared against distractor-absent trials (instead of permuted distractor-present trials; S3 Fig).

If these periodic neural dynamics serve as the basis for the apparent behavioural
fluctuations, we should observe the synchronization of the behavioural and neural time courses
by a common rhythm. To test this, we also examined the co-modulation of sensitivity and
distractor-evoked ERP by distractor onset time. We first calculated the cross-correlation
coefficients of the behavioural and neural time courses for individual participants (Fig 2D). We
then ran a linear mixed model with the cross-correlation coefficient as the outcome measure
and sine- and cosine-transformed time lag as predictors.

Fig 2G shows that sensitivity and distractor-evoked ERP are co-modulated at 3.5 and 5
Hz. At lag 0, there was a negative correlation between sensitivity and the distractor-evoked
ERP, consistent with the notion that stronger distractor encoding (i.e., larger distractor-evoked
ERP) corresponds to worse task performance (i.e., lower sensitivity). T-tests against zero on
the (Fisher-z transformed) correlation coefficients across participants show that this correlation
at time lag 0 was close to statistical significance (Pearson's r: $t_{29} = -1.85$, $p = 0.08$, mean
Pearson's $r = -0.08$; Spearman's r: $t_{29} = -2.13$, $p = 0.04$, mean Spearman's $r = -0.10$).

As a control analysis, the same analysis pipeline was run on the data in the distractor-
absent condition by randomly assigning a "distractor onset" for each distractor-absent trial,
which did not reveal any significant co-modulation (S4 Fig): Neither time courses of sensitivity
nor distractor-evoked ERP were modulated by distractor onset time; time lags did not modulate
the cross-correlation of these two at any frequency. The temporal co-fluctuations of
behavioural and neural measures of distraction at 3 – 5 Hz in distractor-present trials may be a
manifestation of an underlying distractibility rhythm, which we probed into next.

378 Fig 2. The analyses of behavioural and neural time courses by distractor onset time.

A) Average sensitivity (yellow solid line) and 25-Hz amplitude of the distractor-evoked event-related potential (ERP; blue

solid line) across distractor onset times. Shaded areas show ±1 SEM across participants. Dashed lines show respective

quadratic trends. B) Top panel: Distractor-evoked ERP waveform averaged across all distractor onset times at electrode Fz

(20 – 30 Hz bandpass filtered for visualization purpose). Shaded grey area marks the time window used to extract the 25-Hz

amplitude of the distractor-evoked ERP. Inset shows the scalp map of the 25-Hz amplitude of the distractor-evoked ERP

(derived via an FFT on the distractor-evoked ERP waveform). Bottom panel: Distractor-evoked inter-trial phase coherence

(ITPC) from 1 – 10 Hz and from -0.2 s – 0.6 s at Fz. Brain surface shows the ITPC values (frequencies: 3 – 7 Hz; time window:

0 – 0.3 s) in source space, which reflects the auditory response to the distractor. White outline indicates top 1% voxels with

largest ITPC values. C) Detrended time courses of behavioural and neural outcome measures. Shaded areas show ±1 SEM

across participants. D) Solid line shows average correlation coefficients, derived by averaging single-subject cross-correlations

of sensitivity and distractor-evoked ERP time courses, as a function of temporal lags. Shaded area shows ±1 SEM across

participants. E-G) Spectral magnitude across frequencies (0.5 – 8 Hz, 0.5-Hz step) for (E) detrended sensitivity, (F) distractor-

evoked ERP, and (G) the cross-correlation between the two. Shaded areas show the 95th percentile of the permutation

distribution generated from 5,000 permutations. * p < .05. ** p < .01. (uncorrected)

Pre-distractor neural phase in inferior frontal/insular cortex explains distraction

If the human brain hosts an endogenous rhythm that underlies distractibility dynamics, the

neural state prior to distractor onset should explain the participant's momentary vulnerability

to interference by a distractor. To test this, we studied how pre-distractor neural phase relates

to our previously established proxy of distraction, that is, behavioural sensitivity. We asked

when in time and in which brain network(s) such an endogenous rhythm underlying

distractibility would show up.

We employed source-projected EEG time courses to extract the quadratic relationship

between the binned pre-distractor neural phase and perceptual sensitivity. For each trial (Fig

3A), we first transformed a source-projected EEG data segment (0.4 s; sliding window) into

the frequency domain using FFT. We then extracted neural phase for a given frequency (Fig

3B). To calculate sensitivity sorted by phase bin, we first sorted the trials according to their

phase values into 9 phase bins of equal size, followed by calculation of perceptual sensitivity

for each bin (see S5 Fig for individual participants' sensitivity by phase bin). The same

procedure was repeated for a range of frequencies (2.5 – 8 Hz, 0.5-Hz steps) and time windows

(-0.3 – 0.3 s around distractor onset, 0.05-s steps). A cluster-based permutation test with the

dimensions time, frequency, and voxels, wherein the quadratic fit was tested against zero,

revealed a positive significant cluster (Fig 3C; the same analyses with 7, 8, and 10 phase bins

yielded comparable clusters across all dimensions and comparable statistical significance). The

quadratic modulation of sensitivity by neural phase at 2.5 – 7.5 Hz was most prominent in the

left insular and the inferior frontal cortices in the time window spanning ~300 ms before

distractor onset (cluster p-value = .026, two-tailed; see S6 Fig for brain surface plots from other

viewing angles).

To test whether the significant cluster overlaps with sources of auditory-evoked activity

in auditory cortex regions, we compared its source with the source of distractor-evoked inter-

trial phase coherence (ITPC) at 3 – 7 Hz (shown also in Fig 2B, bottom panel). Importantly,

although the two effects were localized in proximal cortical regions (Fig 3C, bottom panel),
their core regions were mostly non-overlapping.

For control, we conducted the same analysis on the distractor-absent trials, which
revealed no significant cluster (S7 Fig). We also tested the relationship between the pre-
distractor neural phase and the post-distractor neural measure of distraction (i.e., 25-Hz
amplitude of the distractor-evoked ERP), which did not reveal a significant effect (S8 Fig).

Fig 3. Cluster-based permutation test results on the relationship between neural phase and behavioural fluctuations.

A-B) Illustration of the source-level analysis. A) Example of a single-trial source-projected EEG time course. The moving
window (grey) was used to transform segments of the data into the frequency domain using FFT. The first grey window
corresponds to the first time window used in the time-resolved analysis (i.e., -0.5 to -0.1 s). B) Spectral representation of the
data segment in (A). Phase values across frequencies were extracted and trials were binned according to their phase values
into 9 phase bins for each frequency, time window, and source location. Bar graph shows exemplary sensitivity values
calculated from the trials sorted by phase bin. A quadratic trend was fitted to the sensitivity values across phase bins (purple
solid line). C) Results of a cluster-based permutation test, which tested quadratic fits in time-frequency-source space against
zero. Top panel shows the t-values (df = 29) across frequencies and time windows, averaged across all the voxels belonging
to the significant positive cluster. The black contour indicates the positive significant cluster. Right column shows individual
participants' quadratic coefficients for each frequency, collapsed across the time windows included in the significant cluster.
Bottom row shows individual participants' quadratic coefficients across time windows, collapsed across frequencies and
voxels included in the significant cluster. Bottom left panel shows the cluster peak effect (3 Hz; -0.2 s), which resides mainly
in left inferior frontal cortex and insular cortex. Only the t-values of the positive significant cluster are shown. The black
contour indicates the regions with the top 1% t-values across the whole brain. The t-values were interpolated and projected

onto MNI coordinates for visualization purposes. The white contour indicates distractor-evoked neural activity, quantified as the top 1% inter-trial phase coherence (ITPC) in the post-distractor time window (i.e., 0 – 0.3 s) at 3 – 7 Hz (shown also in Figure 2B). Bottom right panel shows centred perceptual sensitivity sorted by phase bins in the positive cluster at 3 Hz averaged across participants. Grey thin lines show individual centred perceptual sensitivity.

Discussion

The current study aimed to unravel the temporal dynamics of distractibility, using a pitch discrimination task with auditory distractors. The eventual degree of distraction and the neural processing of distractors were respectively quantified by distractor-evoked performance detriments and neural responses in the human electroencephalogram (EEG). We made a series of interesting observations.

First, the ~3 – 5 Hz fluctuations of behavioural sensitivity across distractor onset time urged for the question whether the same fluctuations are observed in the human brain's response to distractors. Consistently, we found that the distractor-evoked neural response covaries with behavioural sensitivity at similar frequencies. Second, while behavioural sensitivity and the distractor-evoked neural response might partly reflect post-perceptual processes (such as distractor suppression), we asked whether the brain hosts an endogenous oscillation that shapes the momentary state of distractibility. Confirming this, we found that pre-distractor neural phase in left inferior frontal/insular cortex explained rhythmic fluctuations in the momentary degree of distraction.

These major findings support the notion that temporal fluctuations in distractibility on a subsecond time scale can be explained by slow neural oscillatory dynamics in a cortical network beyond the auditory cortex.

The proneness to distraction is inherently dynamic

The current study sheds light on the dynamics of distractibility, which is an important factor often neglected in previous attention research on distraction and suppression. The ultimate degree of detriment that a distractor will cause depends on two endogenous factors: the momentary proneness to distraction (i.e., distractibility) and the ability to suppress a distractor (i.e., distractor suppression). On the one hand, research on distractor suppression often did not disentangle the active suppression of distractors (Schneider et al., 2021) from variations in distractibility. On the other hand, research on distractibility rather treated it as an individual characteristic that, if at all, only changes on a slow temporal scale such as within an experimental session (Forster & Lavie, 2014) or across developmental stages (Kannass et al., 2006). The temporal trajectory of distractibility on a faster, subsecond, time scale had hitherto been left unknown.

With distractor-evoked behavioural and neural measures, we were able to encapsulate

the temporal trajectory of distraction, which fluctuates on a subsecond temporal scale

consistent with the rate of rhythmic sampling in attention (Fiebelkorn et al., 2013; Ho et al.,

2017; Kubetschek & Kayser, 2021; Landau & Fries, 2012) and working memory (Cruzat et al.,

2021; Schmid et al., 2022; ter Wal et al., 2021). With analysis of pre-distractor neural

oscillatory phase, we were able to trace this distractibility back to a slow neural oscillatory

fluctuation in inferior frontal and insular cortex (see below for an in-depth discussion).

Participants could not anticipate whether or when the distractor would occur, thereby not being

able to engage in preparatory suppression of the upcoming distractor (Geng, 2014). The

combined analysis of pre-distractor neural phase and of post-distractor neural and behavioural

measures complementarily elucidates how the brain alternates between states of higher and

lower distractibility. These insights are essential for the inclusion of an explicit account of

distraction in models of attention in psychology and neuroscience.

Fluctuations of distractibility at $3 - 5$ Hz in the current study unveil the dynamic nature

of attention, which was underappreciated in the static spotlight metaphor of attention (Posner

et al., 1980). The attentional sampling of to-be-attended external stimuli (Fiebelkorn et al.,

2013; Ho et al., 2017; Kubetschek & Kayser, 2021) or internal memory representation (Cruzat

et al., 2021; Schmid et al., 2022; ter Wal et al., 2021) has been shown to exhibit temporal

fluctuations at similar frequencies. The waxing and waning of attentional sampling may index

inter-areal coordination between the attentional network and the sensory areas of the brain

(Dugué & VanRullen, 2017), which is associated with the alternation between stronger and

weaker attentional sampling over time (Fiebelkorn & Kastner, 2019). With much evidence on

the temporally dynamic nature of the attentional spotlight, however, there is a lack of

theoretical foundation for the inherent dynamics of cognition outside of this spotlight (Lui &

Wöstmann, 2022). With the observed fluctuations of distractibility in the theta frequency range,

an extension of the existing theory of dynamic attentional sampling to temporally dynamic

distraction is warranted.

While our results demonstrate that distractibility exhibits temporal fluctuations, they do

not reveal whether such fluctuations are independent of the fluctuations found in the attentional

sampling of memory content. Participants in the current study had to maintain the memory

representation of the pitch of tone 1 during a trial. The theta fluctuations found in the current

study thus may represent the sampling of the internal representation of tone 1, with higher

distractibility hypothetically occurring during the phase of reduced sampling of the memory

representation. Alternatively, observed theta fluctuations may represent independent
fluctuations in the proneness to distraction. Previous neuroimaging studies found that the
suppression of distracting inputs may be independent of the sampling of attended inputs
(Noonan et al., 2016; Schneider et al., 2018; Wöstmann et al., 2019). Future investigations may
manipulate both the target and distractor onset time to examine the relationship between the
temporal fluctuations underlying attentional sampling and distractibility.

Of note, as the main analysis approach used here (comparing empirical time courses to
time courses that were shuffled in time) does not distinguish between periodic and aperiodic
temporal structure (Brookshire, 2022), we are careful to conclude from the respective results
alone that distractibility is rhythmic. However, it does not negate the possibility that there is a
periodic temporal structure in distractibility. The premise of rhythmic cognition is that the
apparent fluctuations of performance reflect the periodic orchestration between brain regions
(Fiebelkorn & Kastner, 2019). In addition to fluctuations in behavioural performance, neural
evidence is therefore essential to elucidate the rhythmicity of cognition (Fiebelkorn, 2022;
Wöstmann, 2022). The current study shows a correspondence between slow neural oscillatory
phase and behaviour (using an analysis approach that does not employ shuffling-in-time),
consistent with the notion that distractibility is rhythmic. Future advancements in the analysis
approach to directly test the periodicity in cognition will further strengthen our understanding
of the distractibility dynamics.

Neural dynamics of distractibility originate in inferior frontal/insular cortices

The localisation of neural phase effects underlying distractibility dynamics beyond auditory
cortex regions might suggest that the proneness to distraction is supra-modal. In research on
visual distraction, brain regions in frontal and parietal cortices have been associated with
distractor interference in lesions (Chao & Knight, 1995) or transcranial magnetic stimulation
(Kanai et al., 2011; Wais et al., 2012) studies. The functional connectivity between the left
inferior frontal cortex and hippocampus is associated with the disruptive influence of task-
irrelevant visual distraction on working memory (Wais et al., 2010). While the current study
examined distractibility in the auditory modality, the neural origins found here overlap with
previous research on distraction in the visual modality.

The observed relationship between perceptual sensitivity and the inferior frontal/insular
theta phase suggests that fluctuations in distractibility may be related to the cognitive control

of working memory. The left inferior frontal cortex is assumed to be critical to the resolution

of competition between the maintenance of goal-relevant information and the interference from

external distraction (Irlbacher et al., 2014; Tops & Boksem, 2011; Wais et al., 2012). The

anterior insula is theorised as a gatekeeper to the brain regions responsible for goal-related

cognitive control (Molnar-Szakacs & Uddin, 2022), and is part of the ventral attention system

(Eckert et al., 2009). Specifically, the insular cortex may support the switching between

networks important to internally directed and externally directed cognition, respectively

(Uddin, 2015). The frontal theta rhythm is associated with cognitive control (Berger et al., 2019;

Cavanagh & Frank, 2014; Kamarajan et al., 2004) and the prioritization of relevant memory

representation (Riddle et al., 2020). Taken together, theta oscillations in the inferior frontal and

insular cortices may reflect the orchestration of the cognitive control system to maintain the

internal memory representation and suppress potentially distracting external inputs.

Against what might have been expected, the pre-distractor neural phase did not predict

fluctuations in the distractor-evoked neural response (S8 Fig). However, this null result might

rest on the distractor-evoked ERP being a rather unspecific proxy of distraction. Components

of the distractor-evoked ERP have been shown to reflect cognitive operations other than

distraction, such as reactive suppression (Feldmann-Wüstefeld & Vogel, 2019; Hickey et al.,

2009; Wang et al., 2019) or stimulus prediction (Volosin & Horváth, 2014). Distractibility

dynamics may only account for a small amount of variance in the distractor-evoked ERP.

Conclusions

The present study demonstrates that human proneness to distraction is not uniformly distributed

across time but fluctuates on a subsecond timescale in cycles of ~3 – 5 Hz. In the brain, time

windows of higher distractibility are coined by stronger neural responses to distractors.

Furthermore, slow neural phase in left inferior frontal/insular cortex regions explains

fluctuations in distractibility. These results unravel the temporal dynamics of distractibility and

thereby help explain human processing of an abundant kind of stimulus in increasingly

complex environments, that is, irrelevant and distracting input.

Acknowledgments

We thank Frauke Kraus and Malte Naujokat for their help in data collection.

References

Berger, B., Griesmayr, B., Minarik, T., Biel, A. L., Pinal, D., Sterr, A., & Sauseng, P. (2019). Dynamic

regulation of interregional cortical communication by slow brain oscillations during working memory.

Nature Communications, *10*(1), 4242. https://doi.org/10.1038/s41467-019-12057-0

Bonnefond, M., & Jensen, O. (2012). Alpha oscillations serve to protect working memory maintenance against

anticipated distracters. *Current Biology*, *22*(20), 1969–1974. https://doi.org/10.1016/j.cub.2012.08.029

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

https://doi.org/10.1163/156856897X00357

Brookshire, G. (2022). Putative rhythms in attentional switching can be explained by aperiodic temporal

structure. *Nature Human Behaviour*. https://doi.org/10.1038/s41562-022-01364-0

Buschman, T. J., & Kastner, S. (2015). From Behavior to Neural Dynamics: An Integrated Theory of Attention.

Neuron, *88*(1), 127–144. https://doi.org/10.1016/j.neuron.2015.09.017

Campbell, K. L., Grady, C. L., Ng, C., & Hasher, L. (2012). Age differences in the frontoparietal cognitive

control network: Implications for distractibility. *Neuropsychologia*, *50*(9), 2212–2223.

https://doi.org/10.1016/j.neuropsychologia.2012.05.025

Cavanagh, J. F., & Frank, M. J. (2014). Frontal theta as a mechanism for cognitive control. *Trends in Cognitive*

Sciences, *18*(8), 414–421. https://doi.org/10.1016/j.tics.2014.04.012

Chao, L., & Knight, R. T. (1995). Human prefrontal lesions increase distractibility to irrelevant sensory inputs.

Neuroreport: An International Journal for the Rapid Communication of Research in Neuroscience,

6(12), 1605–1610. https://doi.org/10.1097/00001756-199508000-00005

Cruzat, J., Torralba, M., Ruzzoli, M., Fernández, A., Deco, G., & Soto-Faraco, S. (2021). The phase of Theta

oscillations modulates successful memory formation at encoding. *Neuropsychologia*, *154*, 107775.

https://doi.org/10.1016/j.neuropsychologia.2021.107775

Desimone, R., & Duncan, J. (1995). Neural mechanisms of selective visual attention. *Annual Review of*

Neuroscience, *18*(1), 193–222.

Ding, N., & Simon, J. Z. (2009). Neural representations of complex temporal modulations in the human

auditory cortex. *Journal of Neurophysiology*, *102*(5), 2731–2743.

https://doi.org/10.1152/jn.00523.2009

Dugué, L., & VanRullen, R. (2017). Transcranial magnetic stimulation reveals intrinsic perceptual and

attentional rhythms. *Frontiers in Neuroscience*, *11*. https://doi.org/10.3389/fnins.2017.00154

Dürschmid, S., Reichert, C., Hinrichs, H., Heinze, H.-J., Kirsch, H. E., Knight, R. T., & Deouell, L. Y. (n.d.).

Direct evidence for prediction signals in frontal cortex independent of prediction error. *Cerebral

Cortex*, 1–9. https://doi.org/10.1093/cercor/bhy331

Eckert, M. A., Menon, V., Walczak, A., Ahlstrom, J., Denslow, S., Horwitz, A., & Dubno, J. R. (2009). At the

heart of the ventral attention system: The right anterior insula. *Human Brain Mapping*, *30*(8), 2530–

2541. https://doi.org/10.1002/hbm.20688

Feldmann-Wüstefeld, T., & Vogel, E. K. (2019). Neural Evidence for the Contribution of Active Suppression

During Working Memory Filtering. *Cerebral Cortex*, *29*(2), 529–543.

https://doi.org/10.1093/cercor/bhx336

Fiebelkorn, I. C. (2022). Detecting attention-related rhythms: When is behavior not enough? (Commentary on

van der Werf et al. 2021). *European Journal of Neuroscience*, *55*(11–12), 3117–3120.

https://doi.org/10.1111/ejn.15322

Fiebelkorn, I. C., & Kastner, S. (2019). A Rhythmic Theory of Attention. *Trends in Cognitive Sciences*, *23*(2),

87–101. https://doi.org/10.1016/j.tics.2018.11.009

Fiebelkorn, I. C., Saalmann, Y. B., & Kastner, S. (2013). Rhythmic Sampling within and between objects

despite sustained attention at a cued location. *Current Biology*, *23*(24), 2553–2558.

https://doi.org/10.1016/j.cub.2013.10.063

Forster, S., & Lavie, N. (2014). Distracted by your mind? Individual differences in distractibility predict mind

wandering. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *40*(1), 251–260.

https://doi.org/10.1037/a0034108

Geng, J. J. (2014). Attentional mechanisms of distractor suppression. *Current Directions in Psychological

Science*, *23*(2), 147–153. https://doi.org/10.1177/0963721414525780

Geng, J. J., & DiQuattro, N. E. (2010). Attentional capture by a perceptually salient non-target facilitates target

processing through inhibition and rapid rejection. *Journal of Vision*, *10*(6), 1–12.

https://doi.org/10.1167/10.6.5

Grabenhorst, M., Maloney, L. T., Poeppel, D., & Michalareas, G. (2021). Two sources of uncertainty

independently modulate temporal expectancy. *Proceedings of the National Academy of Sciences*,

118(16), e2019342118. https://doi.org/10.1073/pnas.2019342118

Helfrich, R. F., Fiebelkorn, I. C., Szczepanski, S. M., Lin, J. J., Parvizi, J., Knight, R. T., & Kastner, S. (2018).

Neural mechanisms of sustained attention are rhythmic. *Neuron*, *99*(4), 854-865.e5.

https://doi.org/10.1016/j.neuron.2018.07.032

Herbst, S. K., Stefanics, G., & Obleser, J. (2022). Endogenous modulation of delta phase by expectation–A

replication of Stefanics et al., 2010. *Cortex*, *149*, 226–245. https://doi.org/10.1016/j.cortex.2022.02.001

Hickey, C., Di Lollo, V., & McDonald, J. J. (2009). Electrophysiological Indices of Target and Distractor

Processing in Visual Search. *Journal of Cognitive Neuroscience*, *21*(4), 760–775.

https://doi.org/10.1162/jocn.2009.21039

Ho, H. T., Leung, J., Burr, D. C., Alais, D., & Morrone, M. C. (2017). Auditory sensitivity and decision criteria

oscillate at different frequencies separately for the two ears. *Current Biology*, *27*(23), 3643-3649.e3.

https://doi.org/10.1016/j.cub.2017.10.017

Huang, Y., Chen, L., & Luo, H. (2015). Behavioral oscillation in priming: Competing perceptual predictions

conveyed in alternating theta-band rhythms. *Journal of Neuroscience*, *35*(6), 2830–2837.

https://doi.org/10.1523/JNEUROSCI.4294-14.2015

Irlbacher, K., Kraft, A., Kehrer, S., & Brandt, S. A. (2014). Mechanisms and neuronal networks involved in

reactive and proactive cognitive control of interference in working memory. *Neuroscience &*

Biobehavioral Reviews, *46*, 58–70. https://doi.org/10.1016/j.neubiorev.2014.06.014

Kamarajan, C., Porjesz, B., Jones, K. A., Choi, K., Chorlian, D. B., Padmanabhapillai, A., Rangaswamy, M.,

Stimus, A. T., & Begleiter, H. (2004). The role of brain oscillations as functional correlates of

cognitive systems: A study of frontal inhibitory control in alcoholism. *International Journal of*

Psychophysiology, *51*(2), 155–180. https://doi.org/10.1016/j.ijpsycho.2003.09.004

Kanai, R., Dong, M. Y., Bahrami, B., & Rees, G. (2011). Distractibility in daily life Is reflected in the structure

and function of human parietal cortex. *Journal of Neuroscience*, *31*(18), 6620–6626.

https://doi.org/10.1523/JNEUROSCI.5864-10.2011

Kannass, K. N., Oakes, L. M., & Shaddy, D. J. (2006). A longitudinal investigation of the development of

attention and distractibility. *Journal of Cognition and Development*, *7*(3), 381–409.

https://doi.org/10.1207/s15327647jcd0703_8

Koerner, T. K., & Zhang, Y. (2015). Effects of background noise on inter-trial phase coherence and auditory

N1–P2 responses to speech stimuli. *Hearing Research*, *328*, 113–119.

https://doi.org/10.1016/j.heares.2015.08.002

Kubetschek, C., & Kayser, C. (2021). Delta/Theta band EEG activity shapes the rhythmic perceptual sampling

of auditory scenes. *Scientific Reports*, *11*(2370), 2370. https://doi.org/10.1101/2020.07.21.213694

Lakatos, P., Shah, A. S., Knuth, K. H., Ulbert, I., Karmos, G., & Schroeder, C. E. (2005). An Oscillatory

Hierarchy Controlling Neuronal Excitability and Stimulus Processing in the Auditory Cortex. *Journal*

of Neurophysiology, *94*(3), 1904–1911. https://doi.org/10.1152/jn.00263.2005

Landau, A. N., & Fries, P. (2012). Attention samples stimuli rhythmically. *Current Biology*, *22*(11), 1000–1004.

https://doi.org/10.1016/j.cub.2012.03.054

Lui, T. K.-Y., & Wöstmann, M. (2022). Effects of temporally regular versus irregular distractors on goal-

directed cognition and behavior. *Scientific Reports*, *12*, 10020. https://doi.org/10.1038/s41598-022-

13211-3

Macmillan, N. A., & Kaplan, H. L. (1985). Detection theory analysis of group data: Estimating sensitivity from

average hit and false-alarm rates. *Psychological Bulletin*, *98*(1), 185–199. https://doi.org/10.1037/0033-

2909.98.1.185

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. *Journal of*

Neuroscience Methods, *164*(1), 177–190. https://doi.org/10.1016/j.jneumeth.2007.03.024

Mayhew, S. D., Dirckx, S. G., Niazy, R. K., Iannetti, G. D., & Wise, R. G. (2010). EEG signatures of auditory

activity correlate with simultaneously recorded fMRI responses in humans. *NeuroImage*, *49*(1), 849–

864. https://doi.org/10.1016/j.neuroimage.2009.06.080

Molnar-Szakacs, I., & Uddin, L. Q. (2022). Anterior insula as a gatekeeper of executive control. *Neuroscience*

& Biobehavioral Reviews, *139*, 104736. https://doi.org/10.1016/j.neubiorev.2022.104736

Noonan, M. P., Adamian, N., Pike, A., Printzlau, F., Crittenden, B. M., & Stokes, M. G. (2016). Distinct

mechanisms for distractor suppression and target facilitation. *The Journal of Neuroscience*, *36*(6),

1797–1807. https://doi.org/10.1523/JNEUROSCI.2133-15.2016

Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory.

Neuropsychologia, *9*(1), 97–113. https://doi.org/10.1016/0028-3932(71)90067-4

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open source software for advanced

analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and*

Neuroscience, *2011*, 156869. https://doi.org/10.1155/2011/156869

Oya, H., Gander, P. E., Petkov, C. I., Adolphs, R., Nourski, K. V., Kawasaki, H., Howard, M. A., & Griffiths, T.

D. (2018). Neural phase locking predicts BOLD response in human auditory cortex. *NeuroImage*, *169*,

286–301. https://doi.org/10.1016/j.neuroimage.2017.12.051

Picton, T. W., Hillyard, S. A., Galambos, R., & Schiff, M. (1971). Human auditory attention: A central or

peripheral process? *Science*, *173*(3994), 351–353.

Posner, M. I., Snyder, C. R., & Davidson, B. J. (1980). Attention and the detection of signals. *Journal of*

Experimental Psychology: General, *109*(2), 160–174. https://doi.org/10.1037/0096-3445.109.2.160

Prins, N., & Kingdom, F. A. A. (2018). Applying the model-comparison approach to test specific research

hypotheses in psychophysical research using the palamedes toolbox. *Frontiers in Psychology*, *9*, 1250.

https://doi.org/10.3389/fpsyg.2018.01250

Riddle, J., Scimeca, J. M., Cellier, D., Dhanani, S., & D'Esposito, M. (2020). Causal evidence for a role of theta

and alpha oscillations in the control of working memory. *Current Biology*, *30*(9), 1748-1754.e4.

https://doi.org/10.1016/j.cub.2020.02.065

Rutishauser, U., Ross, I. B., Mamelak, A. N., & Schuman, E. M. (2010). Human memory strength is predicted

by theta-frequency phase-locking of single neurons. *Nature*, *464*(7290), 903–907.

https://doi.org/10.1038/nature08860

Schmid, R. R., Pomper, U., & Ansorge, U. (2022). Cyclic reactivation of distinct feature dimensions in human

visual working memory. *Acta Psychologica*, *226*, 103561.

https://doi.org/10.1016/j.actpsy.2022.103561

Schneider, D., Göddertz, A., Haase, H., Hickey, C., & Wascher, E. (2018). Hemispheric asymmetries in EEG

alpha oscillations indicate active inhibition during attentional orienting within working memory.

Behavioural Brain Research, *359*, 38–46. https://doi.org/10.1016/j.bbr.2018.10.020

Schneider, D., Herbst, S. K., Klatt, L., & Wöstmann, M. (2021). Target enhancement or distractor suppression?

Functionally distinct alpha oscillations form the basis of attention. *European Journal of Neuroscience*,

ejn.15309. https://doi.org/10.1111/ejn.15309

Shamma, S. A., Elhilali, M., & Micheyl, C. (2011). Temporal coherence and attention in auditory scene

analysis. *Trends in Neurosciences*, *34*(3), 114–123. https://doi.org/10.1016/j.tins.2010.11.002

Siegel, M., Warden, M. R., & Miller, E. K. (2009). Phase-dependent neuronal coding of objects in short-term

memory. *Proceedings of the National Academy of Sciences*, *106*(50), 21341–21346.

https://doi.org/10.1073/pnas.0908193106

Stefanics, G., Hangya, B., Hernadi, I., Winkler, I., Lakatos, P., & Ulbert, I. (2010). Phase entrainment of human

delta oscillations can mediate the effects of expectation on reaction speed. *Journal of Neuroscience*,

30(41), 13578–13585. https://doi.org/10.1523/JNEUROSCI.0703-10.2010

ter Wal, M., Linde-Domingo, J., Lifanov, J., Roux, F., Kolibius, L. D., Gollwitzer, S., Lang, J., Hamer, H.,

Rollings, D., Sawlani, V., Chelvarajah, R., Staresina, B., Hanslmayr, S., & Wimber, M. (2021). Theta

rhythmicity governs human behavior and hippocampal signals during memory-dependent tasks. *Nature

Communications*, *12*(1), 7048. https://doi.org/10.1038/s41467-021-27323-3

Tops, M., & Boksem, M. A. S. (2011). A potential role of the inferior frontal gyrus and anterior insula in

cognitive control, brain rhythms, and event-related potentials. *Frontiers in Psychology*, *2*.

https://doi.org/10.3389/fpsyg.2011.00330

Uddin, L. Q. (2015). Salience processing and insular cortical function and dysfunction. *Nature Reviews

Neuroscience*, *16*(1), 55–61. https://doi.org/10.1038/nrn3857

van Moorselaar, D., Lampers, E., Cordesius, E., & Slagter, H. A. (2020). Neural mechanisms underlying

expectation-dependent inhibition of distracting information. *ELife*, *9*, e61048.

https://doi.org/10.7554/eLife.61048

Van Veen, B. D., Van Drongelen, W., Yuchtman, M., & Suzuki, A. (1997). Localization of brain electrical

activity via linearly constrained minimum variance spatial filtering. *IEEE Transactions on Biomedical*

Engineering, *44*(9), 867–880. https://doi.org/10.1109/10.623056

VanRullen, R. (2016). Perceptual Cycles. *Trends in Cognitive Sciences*, *20*(10), 723–735.

https://doi.org/10.1016/j.tics.2016.07.006

Volosin, M., & Horváth, J. (2014). Knowledge of sequence structure prevents auditory distraction: An ERP

study. *International Journal of Psychophysiology*, *92*(3), 93–98.

https://doi.org/10.1016/j.ijpsycho.2014.03.003

Wais, P. E., Kim, O. Y., & Gazzaley, A. (2012). Distractibility during episodic retrieval is exacerbated by

perturbation of left ventrolateral prefrontal cortex. *Cerebral Cortex*, *22*(3), 717–724.

https://doi.org/10.1093/cercor/bhr160

Wais, P. E., Rubens, M. T., Boccanfuso, J., & Gazzaley, A. (2010). Neural mechanisms underlying the impact

of visual distraction on retrieval of long-term memory. *Journal of Neuroscience*, *30*(25), 8541–8550.

https://doi.org/10.1523/JNEUROSCI.1478-10.2010

Wang, B., van Driel, J., Ort, E., & Theeuwes, J. (2019). Anticipatory Distractor Suppression Elicited by

Statistical Regularities in Visual Search. *Journal of Cognitive Neuroscience*, *31*(10), 1535–1548.

https://doi.org/10.1162/jocn_a_01433

Weisz, N., Kraft, N. G., & Demarchi, G. (2020). Auditory cortical alpha/beta desynchronization prioritizes the

representation of memory items during a retention period. *ELife*, *9*, e55508.

https://doi.org/10.7554/eLife.55508

Wöstmann, M. (2022). Does attention follow a rhythm? *Nature Human Behaviour*.

https://doi.org/10.1038/s41562-022-01365-z

Wöstmann, M., Alavash, M., & Obleser, J. (2019). Alpha oscillations in the human brain implement distractor

suppression independent of target selection. *The Journal of Neuroscience*, *39*(49), 9797–9805.

https://doi.org/10.1523/JNEUROSCI.1954-19.2019

Wöstmann, M., Lui, T. K.-Y., Friese, K.-H., Kreitewolf, J., Naujokat, M., & Obleser, J. (2020). The

vulnerability of working memory to distraction is rhythmic. *Neuropsychologia*, *146*, 107505.

https://doi.org/10.1016/j.neuropsychologia.2020.107505

Wöstmann, M., Störmer, V. S., Obleser, J., Addleman, D. A., Andersen, S. K., Gaspelin, N., Geng, J. J., Luck,

S. J., Noonan, M. P., Slagter, H. A., & Theeuwes, J. (2022). Ten simple rules to study distractor

suppression. *Progress in Neurobiology*, *213*, 102269. https://doi.org/10.1016/j.pneurobio.2022.102269

Zoefel, B., Davis, M. H., Valente, G., & Riecke, L. (2019). How to test for phasic modulation of neural and

behavioural responses. *NeuroImage*, *202*, 116175. https://doi.org/10.1016/j.neuroimage.2019.116175

Fig. S1. Individual time courses of raw sensitivity (yellow) and distractor-evoked ERP (25-Hz amplitude of distractor-evoked ERP; blue).

Fig. S2. Detrended (quadratic trend removed) sensitivity (yellow) and distractor-evoked ERP (blue) individual time courses.

Fig. S3. Comparison of the spectral magnitude between distractor-present and distractor-absent conditions. Left panel shows the sensitivity time courses for distractor-present (solid) and distractor-absent (dashed) conditions. Right panel shows the averaged frequency spectra, derived from FFT on single-subject time courses. Shaded areas show ±1 SEM across individual participants. Asterisks show statistical significance when comparing the spectral magnitude of distractor-present versus -absent conditions (using uncorrected dependent-samples t-tests). * $p < .05$ ** $p < .01$

781 **Fig. S4. The same analysis pipeline as shown in Fig 2 applied to the distractor-absent condition.**

Fig. S5. Individual sensitivity (grey line) and quadratic fit (purple line) across phase bins of the significant positive
cluster at 3 Hz.

Left   Right

Lateral

Medial

786 **Fig. S6. Brain surface plots of the cluster peak effect.** Brain surface plots of the cluster peak effect (3 Hz; -0.2 s) from left
787 lateral (top left), right lateral (top right), left medial (bottom left), and right medial (bottom right) views show t-values for the
788 comparison of the quadratic fit of the sensitivity sorted by phase bins against zero. The t-values were interpolated and projected
789 onto MNI coordinates for visualization purposes.

Fig. S7. Cluster-based permutation test in the distractor-absent condition. Results of the cluster-based permutation test, across time windows and frequencies, on the quadratic relationship between neural phase and sensitivity for the distractor-absent condition. Figure shows t-values (df = 29) averaged across all the voxels belonging to the significant positive cluster in the distractor-present condition shown in Figure 3. No significant cluster was found in the distractor-absent condition.

Fig. S8. Cluster-based permutation test on distractor-evoked ERP amplitude. Results of the cluster-based permutation
test, across time windows and frequencies, on the quadratic relationship between neural phase and distractor-evoked ERP
amplitude at 25 Hz. Figure shows t-values (df = 29) averaged across all the voxels belonging to the significant positive cluster
testing the quadratic relationship between neural phase and behavioural sensitivity shown in Figure 3. As expected, and not of
main interest in the current study, a significant cluster in the post-stimulus time window (i.e., >0s) was found. More importantly,
no significant cluster was found in the pre-distractor time window (i.e., <0 s).