Neurogenomic landscape associated with status-dependent cooperative behavior

Peri E. Bolton[1,2*], T. Brandt Ryder[3,4], Roslyn Dakin[3,5], Jennifer L. Houtz[6], Ignacio T. Moore[7],

Christopher N. Balakrishnan[1†], Brent M. Horton[6*†]

*Corresponding Author: Peri Bolton peri.bolton@gmail.com

† CNB and BMH contributed equally to this work.

[1]Department of Biology, East Carolina University, Greenville, NC, USA

[2]Department of Vertebrate Zoology, National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

[3]Migratory Bird Center, Smithsonian National Zoological Park, Washington DC, USA

[4]Bird Conservancy of the Rockies, Fort Collins, CO, USA

[5]Department of Biology, Carleton University, Ottawa, ON, Canada

[6]Department of Biology, Millersville University, Millersville, PA, USA

[7]Department of Biological Sciences, Virginia Tech, Blacksburg, VA, USA

Abstract

The neurogenomic mechanisms mediating cooperative behaviors are understudied. We leveraged extensive transcriptomic and behavioral data on a neotropical bird (*Pipra filicauda*) that perform cooperative courtship displays to better understand the underlying mechanisms. Cooperative display is modulated by testosterone, it promotes cooperation in non-territorial birds, but suppresses cooperation in territory holders. We sought to understand the neurogenomic underpinnings of these three related traits: social status, cooperative behavior, and testosterone phenotype. To do this, we profiled gene expression in 10 brain nuclei spanning the social decision-making network (SDMN), and two key endocrine tissues that regulate social behavior. We associated gene expression with each bird's behavioral and endocrine profile derived from 3 years of repeated measures taken from free-living birds in the Ecuadorian Amazon. We found

distinct landscapes of constitutive gene expression were associated with social status,

testosterone phenotype, and cooperation, reflecting the modular organization and engagement of

neuroendocrine tissues. Sex-steroid and neuropeptide signaling appeared to be important in

mediating status-specific relationships between testosterone and cooperation, suggesting shared

regulatory mechanisms with male aggressive and sexual behaviors. We also identified

differentially regulated genes involved in cellular activity and synaptic potentiation, suggesting

multiple mechanisms underpin these genomic states. Finally, we identified SDMN-wide gene

expression differences between territorial and floater males that could form the basis of "status-

specific" neurophysiological phenotypes, potentially mediated by testosterone and growth

hormone. Overall, our findings provide new, systems-level insights into the mechanisms of

cooperative behavior, and suggest that differences in neurogenomic state are the basis for

individual differences in social behavior.

Introduction

41
Cooperation is the synchronized behavior(s) of two or more individuals to produce a net

fitness benefit for one or more individuals (Taborsky and Taborsky 2015). An individual's

decisions about cooperation are based on the immediate social stimulus and individual condition,

which is based on recent and distant past experiences, developmental effects, and underlying

genetics. These effects can ultimately produce repeatable, among-individual variation in

behaviors – a behavioral phenotype – where some individuals tend to be more cooperative than

others. To understand the mechanisms of cooperation, one needs to understand the

neurophysiological signal processing of the stimulus through to the behavioral response, as well

as the genetic and neurophysiological components that underpin behavioral phenotypes. Indeed,

understanding these mechanisms enables us to link how genetic and neurophysiological

architecture can constrain and facilitate the evolution of cooperation (Soares et al. 2010;

Hofmann et al. 2014; Taborsky and Taborsky 2015). Differences in cooperative behavioral

phenotype are in part achieved by the pleiotropic yet modular effects of hormone signaling

systems (Ketterson et al. 2009; Cox 2020). Hormonal pleiotropy mediates phenotypic

integration, whereby multiple traits covary with a hormonal signal (Ketterson et al. 2009; Cox

2020), sometimes in opposite directions. For example, elevated testosterone typically promotes

aggression but inhibits parental care in males (Wingfield et al. 2001). In contrast, through

variation in hormone receptor expression, tissues across the body are often modular in their

response to hormone signals, which enables diverse and flexible responses to hormonal and

social signals and trait independence (Ketterson et al. 2009; Lipshutz et al. 2019).

Cooperation, like other social behaviors, appears to be mediated by differences in

circulating hormones, such as sex steroids, and aspects of hormone-signaling pathways,

including the distribution and abundance of hormone receptors in neural tissues (Soares et al.

2010; Kasper et al. 2017). The neural substrates modulating social behaviors are conserved

across vertebrates (O'Connell and Hofmann 2011). Known as the social decision making

network (SDMN), these reciprocally interconnected brain nuclei are rich in steroid receptors,

steroidogenic enzymes, and steroid-modulated neuropeptide systems (Newman 1999; Goodson

2005; O'Connell and Hofmann 2011). The SDMN is both modular and dynamic, whereby

different combinations of brain regions are activated in response to different social contexts

(Newman 1999; Goodson 2005). This model means the common practice of measuring the

neurological response in a single nucleus is not reflective of all systems engaged in modulating a

given social behavior. Variation among individuals in the expression of neuroendocrine genes

across the SDMN likely underlies repeatable individual differences in social behavior and sets

conditions for behavioral phenotypes that respond differently to circulating hormones and social

stimuli. For example, the up or down-regulation of genes involved in sex-steroid signaling

pathways across nuclei of the SDMN can alter the sensitivity of the male brain to testosterone

and manifest as differential behavioral responses to a given hormone signal. Emerging research

is showing that behavioral phenotypes are correlated with differences in constitutive gene

expression or neurogenomic state among individuals characterized by suites of genes co-

expressed across behaviorally relevant brain regions (Antunes et al. 2021; Kabelik et al. 2021;

Lattin et al. 2021). However, given the diversity of behavioral components associated with

cooperation (Díaz-Muñoz et al. 2014; Kasper et al. 2017), questions remain on how

neuroendocrine mechanisms operate to produce the variety of social and cooperative behaviors

among individuals. Indeed, our understanding of the neuroendocrine mechanisms underlying

seemingly paradoxical male-male reproductive cooperative behaviors is only emerging (DuVal

and Goymann 2011; Díaz-Muñoz et al. 2014; Ryder et al. 2020; Vernasco et al. 2020; Jones and

DuVal 2021; Loveland et al. 2021). For the first time, we characterize gene expression across the

neuroendocrine system in relation status-specific androgenic modulation of a male-male

cooperative behavior.

The wire-tailed manakin (*Pipra filicauda*) is a neotropical lek breeding bird species in which

unrelated males perform cooperative displays to attract females (Figure 1A). Long-term display

partnerships are established between territorial and subordinate non-territorial (floater) males and

form the basis of complex social networks (Figure 1A, B) (Ryder et al. 2008; Ryder, Blake, et al.

2011). In this bird, territoriality is a pre-requisite to siring offspring, and floater males follow an

age-graded queue for territorial status (Ryder et al. 2008; Ryder et al. 2009). Cooperation is

directly linked to fitness, whereby territory holders with more display partners sire more

offspring, and floaters with more partners are more likely to ascend to territorial status (Ryder et

al. 2009). Individuals have a repeatable 'testosterone phenotype' in which territorial males tend

to have higher testosterone than floaters (Figure 1D) (Ryder, Horton, et al. 2011; Ryder et al.

2020). Remarkably, circulating testosterone has a status-specific effect on the cooperative

display behavior: experimental and observational evidence show that higher testosterone levels

are antagonistic to cooperation in territorial males but promote cooperation in floater males

(Ryder et al. 2020; Vernasco et al. 2020) (Figure 1C).

To explore the neurogenomic basis of the status-specific androgenic modulation of

cooperative behavior in these birds, we quantify transcriptome-level gene expression in the

SDMN and pituitary and testes of free-living male wire-tailed manakins sampled in the

Ecuadorian Amazon. Captive studies can obscure ecologically and evolutionary relevant

variation in environmental and behavioral responses (Griffith et al. 2017; Kasper et al. 2017), so

we demonstrate the exciting applicability of neuro-transcriptional methods in the challenging

conditions of the rainforest. In this study, we aim to characterize the genes and neuroendocrine

pathways that modulate individual variation in cooperative behavior, as well as those related

social status and individual variation in testosterone phenotype as a modifier of both social status

and cooperative behavior. We characterize the constitutive gene expression for each of these

traits within and between nodes of the SDMN and the endocrine tissues. Given the androgenic

regulation of cooperative display (Ryder et al. 2020; Vernasco et al. 2020), we hypothesize that

differences in gene regulation will manifest in SDNM nuclei with high density of sex-steroid

receptors and pituitary and gonadal tissues (Newman 1999; Goodson 2005). We further reason

that androgenic regulation will relate to differential expression of genes relating to sex-steroid

signaling and metabolism, as well as three key neuropeptide systems involved in social behavior

(Goodson and Thompson 2010; Kingsbury and Wilson 2016). Given the modular nature of the

SDNM, we predict that many gene expression patterns will be brain region-specific (Figure 1G).

However, given the pleiotropic effects of testosterone (Ketterson et al. 2009; Cox 2020), we also

predict that there will be signatures of testosterone's systemic influence on gene expression

patterns across the brain. We discuss our results in the context of modes of phenotypic evolution

and similarities with mechanisms of other social behaviors.

Materials and Methods

All methods described here were approved by Smithsonian Animal Care and Use Committee

(protocols 12-12,14-25,17-11), as well as the Ecuadorian Ministry of the Environment (MAE-

DNB-CM-2015-0008), and samples were transported with export permit (006-016-EXP-1C-

FAU-DNB/MA) and import permit (USDA APHIS 126133). Detailed methods can be found in

the supplementary material, and full code from analyses in R v 4.0.2 are available on the GitHub

site (https://periperipatus.github.io/PIFI_brain_transcriptome/) and on FigShare

(10.25573/data.22186516).

Field Methods

For RNA sequencing we sampled 16 wild territorial and floater wire-tailed manakin males from

Tiputini Biodiversity Station in the lowland Amazon rainforest of Ecuador (Table S1.1). Males

were categorized by social status as either territorial or floater. The territorial males sampled

(n=9) were males in definitive plumage that held a consistent display territory over the study

period, while floaters (n=7) had either definitive or pre-definitive plumage and did not occupy

consistent territories within leks (Ryder et al. 2008). On 12 of these 16 males, we used repeated

testosterone sampling (Figure 1D, Supplement 1) to characterize each male's testosterone

phenotype (mean testosterone). Previous research has shown that territorial males have, on

average, higher circulating testosterone than floaters (Ryder et al., 2020; Ryder, Horton, et al.,

2011), and the males sampled for this study reflect that pattern (Figure 1D). A male's

testosterone phenotype is represented by mean of the residuals of a linear regression from

repeated sampling of testosterone concentration (Supplement 1 for more detail, see also (Ryder

et al. 2020)). Mean testosterone as a measure of testosterone phenotype was shown to be

repeatable within individuals (Ryder et al. 2020). In the broader population, the relationship

between testosterone phenotype and social network measures used as a proxy for cooperation

was status-specific, with testosterone suppressing cooperation in territorial males but increasing

cooperation in floaters (Ryder et al. 2020) (Figure 1C).

Social network structure and individual variation in behavior were characterized using

autonomous radio-telemetry and tag proximity detections within territories on a total population

of 180 male manakins over multiple years (Ryder et al. 2012; Dakin and Ryder 2018; Ryder et

al. 2020). We used this tagging data to calculate social network strength for the same 12 males

with repeated testosterone measures. Social network strength is a proxy for cooperative tendency

and is the time spent interacting with other males averaged over the study period (Figure 1E),

and was the most repeatable within individuals (Ryder et al. 2020). This social network proxy for

cooperation was most strongly status specific relation to an individual's testosterone phenotype

(Ryder et al. 2020) (Figure 1C), but the overall frequency of cooperative interactions did not

differ between territorial and floater males (Figure 1E).

Microdissection and RNAseq
Methods for brain extraction, tissue preservation, cryosectioning, microdissection, and RNA
extraction followed those published previously (Horton, Ryder, et al. 2020). Briefly, birds were
euthanized by decapitation near the site of capture immediately after blood sampling for
testosterone. Whole brains were extracted within 4-6 minutes of euthanasia and immediately
frozen in powdered dry ice. To capture the full Hypothalamic-Pituitary-Gonadal axis critical to
sex-steroid regulation, we additionally sampled the pituitary gland and testes, which were
sampled immediately after the brain. Brains and testes were kept frozen on dry ice in the field (<
4 hours) until transferred into a 21-day liquid nitrogen-charged Arctic Express™ dry shipper
(Thermo Scientific) where they were stored at cryogenic temperatures until imported into the
United States. Pituitaries were preserved in RNAlater™ (Invitrogen) at ambient temperatures
until imported into the United States.

The brains were cryosectioned, and frozen sections were microdissected to excise tissue from 10
different nuclei involved in social behavior – nine of which are from the SDMN. The SDMN
reflects the interconnectivity of two networks involved in regulating social behavior (O'Connell
and Hofmann 2011), constituting nuclei from both the Social Behavior Network (SBN)
(Newman 1999; Goodson 2005) and the Mesolimbic Reward System (MRS) (Olds and Milner
1954; O'Connell and Hofmann 2011). Figure 1F indicates the nomenclature used and the
relationships of these nuclei to the SBN and MRS. To clarify bird specific nomenclature: the
arcopallium intermedium (AI) is a region that may be, in part, homologous the basolateral
amygdala of mammals, a node of the MRS (O'Connell and Hofmann 2011), which is
hypothesized to play a role in androgen-dependent manakin display behavior (Fusani et al.
2014); and the nucleus taenia (TnA) is homologous to the mammalian medial amygdala
(O'Connell and Hofmann 2011). Finally, we sampled the paraventricular nucleus (PVN) for a set
of 10 nuclei sampled. This is a region that is not yet formally recognized as part of the SDMN,
despite its neuropeptide projections to nodes of the SDMN and established role as a major
regulator of vertebrate social behavior (Goodson and Thompson 2010; Goodson and Kingsbury
2013). For a graphical representation of the tissues sampled see Figure 1F, with the full list of
individuals collected presented in Table S1.1 and tissues in Table S1.2.

Microdissected tissues from both hemispheres were combined for sequencing. RNA was
extracted and sequenced according to previously described methods (Horton, Ryder, et al. 2020).

RNA libraries for the various tissues and field seasons were randomized amongst sequencing

batches (Supplement 1, Table S1.1 & S1.2). All samples analyzed are associated with Bioproject

PRJNA437157, and SRR12660169-198, SRR19521260-271, SRR19521432-575.

Gene expression analyses

Reads were mapped to the annotated *Pipra filicauda* genome (GCA_003945595.1) using

splice-aware mapper STAR v.2.7.5 (Dobin et al. 2013), and transcripts were counted against

exons using featureCounts v.2.0.1 (Liao et al. 2014). For subsequent analyses we used a total of

170 libraries, and quality filtering of individuals and gene-counts was performed before the

analysis of each tissue (Supplement 1 & 3). To further explore tissue and brain nucleus function

in this species (Horton, Ryder, et al. 2020), we characterized differences in gene expression

among tissues and brain nuclei using principal components analysis in PCAtool*s* and Weighted

Gene Co-Expression Analysis (WGCNA) (Langfelder and Horvath 2008; Blighe and Lun 2020)

(Supplement 2). The number of individuals used in each tissue varies owing to quality filtering

and variable sampling of individuals across all brain regions (Table S1.1 & S1.2).

To describe the role of tissues and brain nuclei in relation to these hormonal and

behavioral phenotypes we conducted filtering (supplement 1), differential expression

(Supplement 3) and co-expression (Supplement 5) analysis in each tissue and nucleus separately.

For a subset of tissues, we had an extra four individuals characterized for social status from a

previous study (Horton, Ryder, et al. 2020), for which behavioral and hormone sampling data

were unavailable. Owing to this, and to not overparameterize the models that account for batch

effects we identified differentially expressed genes (DEGs) associated with each of our traits of

interest (social status, testosterone phenotype and social network strength) separately. These

were run as separate negative binomial generalized linear models for each trait that accounted for

batch effects where necessary using DESeq2 (Love et al. 2014) (For detailed description see

Supplement 3) in R v 4.0.2 (R Core Team 2020). Mean testosterone and social network strength

were input to DESeq2 as continuous variables. In these cases, significant DEGs were interpreted

to be correlated with the predictor variable, and Log_2 Fold Change (LFC) value represents the

LFC of gene expression per unit of the predictor variable. In all cases, significant DEGs were

identified using the Wald Test and the default FDR corrected p-value of 0.1 ($q<0.1$). Further, for

each tissue and nucleus we identified gene co-expression networks using WGCNA. The identity

of genes of interest identified by WGCNA were similar to those in DESeq2. For brevity and

simplicity we focus mostly on the differential expression results, but the full results are presented
in Supplement 5 (WGCNA) and Supplement 3 (differential expression).

Due to the key role testosterone plays in modulating behavior in this species (Ryder et al.
2020), we conducted further detailed analysis on nine candidate genes involved in sex-hormone
metabolism and signaling (Figure S4.1), and a further nine steroid sensitive neuropeptides and
their receptors (Figure S4.2). Any candidate gene identified by differential gene expression
(DGE) analysis with an uncorrected $p < 0.05$ was considered "significant" and examined further.
In addition to analyzing models of status, mean testosterone and social network strength
separately, we also explored whether candidate genes were involved in mediating status specific
relationships between mean testosterone and social behavior (Ryder et al. 2020) by extracting
"significant" candidate genes from a final DGE interaction analysis (model form: expression ~
status x mean testosterone). Owing to the overfitting concerns discussed above, we primarily
considered this model as a method for identifying which of our candidate genes showed status-
specific expression with respect to mean testosterone, rather than identifying new DEGs. The list
of putative relationships between genes and traits were plotted (Supplement 4), and any
relationship with highly influential observations, or with $R^2 \leq 0.2$, was excluded from our final
set. For candidates derived from the interaction effect models, and for genes that were associated
with multiple traits, we decided the most convincing relationship based on a stepdown AIC
procedure (Supplement 4). Given the substantially smaller set of candidate genes we considered
uncorrected p-values to be significant ($p<0.05$), but report the FDR corrected value as well.

To explore whether neuroanatomy constrains the types of genes expressed among brain
regions, we characterized overlap in gene expression patterns among brain nuclei by comparison
of ranked gene-lists (Supplement 6). The gene-lists were ranked by $-\log_{10}$ transforming the p-
values and multiplying by the direction of expression. Then these gene lists were compared for
similarity between each pair of tissues by calculating a) Pearson correlation and b) rank-rank
hypergeometric overlap (RRHO) test (Cahill et al. 2018).

Testosterone is pleiotropic in that it can affect multiple traits by influencing gene
expression in similar ways in many different target tissues (Ketterson et al. 2009; Cox 2020), and
it is crucial to social status-specific regulation of behavior in the wire-tailed manakin (Ryder et
al. 2020). We reasoned that these effects would be reflected in shared signatures of differential

expression across brain nuclei. To explore systemic gene expression patterns (i.e., consistent
patterns across the entire brain), we used the transformed p-values from the aforementioned
analysis and calculated the median p-value across all brain nuclei. Genes that had a median p-
value < 0.05 (transformed value > |1.3|) were considered consistently differentially expressed
across the brain. Further, the eigenvalues for each principal component (PC) from the whole-
brain PCA analysis (Figure 1G inset) were Pearson correlated with our variables of interest using
pcatools::eigencorplot() (Blighe and Lun 2020) (Supplement 1). Correlations observed with
social status were further validated by a linear mixed model in package 'nlme' (Pinheiro et al.
2016). The model form included individual as a random effect and incorporated batch variables
that were also correlated with the same PC (Supplement 1).

We used *clusterProfiler* v3.16.1(Yu et al. 2012) to determine patterns of functionality in
DEGs and WGCNA modules with Gene Ontology enrichment analyses. If the total number of
FDR corrected DEGs was less than 40, we used gene set enrichment analysis instead which
leverages the rank order of p-values and direction of expression (Supplement 3).

Results

Landscape of gene expression among neural and endocrine tissues

Our analyses revealed substantial differences in gene expression among tissues and brain
regions (Figure 1G; Figure S2.7). As expected, the pituitary was associated with high expression
levels of genes key to hormonal regulation (e.g., PRL, GH1, TSHB, POMC; Figure S2.3), while
the testes showed high expression levels of genes associated with germ cells and flagellar
motility (e.g., PKDREJ, BOLL, DRC7; Figure S2.3). Within the brain, the majority of gene
expression variation was explained by differences between brain regions (Figure 1G inset) and
was reflected in brain region-specific gene co-expression networks (Figure S2.7). These
expression patterns were more similar between adjacent nuclei, forming major brain-region
groupings of the amygdala, hypothalamus, and midbrain (Figure 1G). Our 18 candidate
neuroendocrine genes (Table S1.3) also showed tissue and brain region-specific expression
(Table 2.2, Figures S4.1-4.2).

Social status and testosterone phenotype are associated with tissue and brain region-specific
patterns of gene expression

We observed a landscape of gene expression in association with testosterone phenotype
(Figure 2A) and social status (Figure 2B), that was characterized by a high number of DEGs (or
co-expression modules) identified in some tissues, but not others. Similarly, across brain regions
we identified relationships between many of our 18 candidate genes and status, testosterone
phenotype or their interaction (Figure 2C). Mean testosterone was a continuous predictor in
DESeq2, as such DEGs showed up-or downregulation with a higher testosterone phenotype
(Figure 2A), while DEGs identified in social status indicated up or down regulation in territorial
males (Figure 2B). The number of trait-associated co-expression modules were similar to our
results from differential gene expression, so we primarily focus on the tissues with large numbers
of DEGs and significant candidate genes for the main manuscript, but see Supplements 3 and 5
for all results. The number of DEGs in each tissue was more similar between testosterone
phenotype and status than any other comparison of traits examined (R = 0.5, p = 0.08, Figure
S3.7), and this pattern appeared to hold across co-expression modules (Supplement 5) and
candidate gene analysis (Figure 2C & Figure 3B). This similarity was expected because
testosterone phenotype and status themselves are correlated (Figure 1D) (Ryder, Horton, et al.
2011; Ryder et al. 2020).

The testes and TnA each had over 1,000 genes differentially expressed (at uncorrected p-
value < 0.05) in association with both testosterone phenotype and status (144 and 252 genes after
FDR correction q<0.01 in association with mean T; 2 and 13 after FDR correction in association
with social status). DEGs identified in association with testosterone phenotype in the testes were
enriched for genes involved with response to corticosterone (GO:0051412) and synaptic
transmission and organization (e.g., GO:0007268, GO:0050808) among others (Figure S3.20).
Notable DEGs included steroidogenic acute regulatory protein (STAR, LFC = -1.1, p = 2×10^{-7}, q
= 0.0003, R^2 = 0.6) and candidate gene AVPR1A (LFC = 0.8, p = 5×10^{-5}, q = 0.01, R^2 = 0.7;
Figure S4.4). Note that Log_2 Fold Change (LFC) represents change in gene expression per unit
of the predictor variable. However, these transcriptional differences do not correspond to
significant differences in testis size in relation to testosterone phenotype (Table S3.2), nor social
status (Table S3.1 & S3.2).

Among the 252 DEGs associated with testosterone phenotype in TnA, there was non-
significant enrichment for genes involved in neural crest migration (GO:0001755), learning and
memory (GO:0007613, GO:0007612), as well as long-term synaptic potentiation (e.g.,
GO:0060291) (Figure S3.28). Among the DEGs associated with testosterone phenotype in TnA,
candidate gene estrogen receptor 2 (ESR2; aka Erβ) was upregulated in TnA of males with
higher testosterone (Figure 2C; LFC = 0.8, p = 0.001, q = 0.08, R^2 = 0.4). Other notable DEGs in
TnA included upregulation of immediate early gene JUN, also a 'coral1' hub gene (module
membership (MM) = 0.93; LFC = 0.61, p = 0.0004, q = 0.05) and corticotropin releasing
hormone receptor 2 (CRHR2, LFC = 1, q = 0.006, R^2 = 0.5).

Not all nuclei showed similar patterns between status and testosterone phenotype. For
example, ICo of the midbrain showed 161 DEGs associated with testosterone phenotype after
FDR correction, but did not stand out in association with social status (3 DEGs). Most of the
significantly enriched GO terms were biological processes relating to motor proteins and cilia
(e.g., GO:0003341, Figure S3.26). Intriguingly, although the POM did not reveal a high number
of DEGs in association with testosterone phenotype (45), our analysis revealed differential
expression for compelling gene ontology categories and candidate neuroendocrine genes (Figure
S3.25, Figure S4.10, Figure S5.7). Notably, males with higher testosterone exhibited
upregulation of candidate gene androgen receptor (AR) in POM (Figure 2D). Indeed, the
'floralwhite' module comprised many of the upregulated DEGs and was significantly associated
with both testosterone phenotype and to a lesser extent social status (Figure S5.6). This module
showed enrichment for biological processes relating to G protein-coupled receptor signaling
pathways (GO:0007186) and neuropeptide signaling pathway (GO:0007218), with statistically
non-significant enrichment for steroid receptor signaling (GO:0030518), learning and memory
(GO:0007611), and nervous system development (GO:0007399) (Figure S3.7). Indeed, many of
the module hub genes were also DEGs associated with testosterone phenotype, and interact with
candidate sex hormone pathways, prolactin releasing hormone (PRLH, MM = 0.91, LFC = 2.4, q
= 5×10^{-5}, R^2 = 0.6, Figure S3.25A) and growth regulating estrogen receptor binding 1 (GREB1,
MM = 0.86, LFC = 0.6, q = 3×10^{-5}, R^2 = 0.7, Figure S3.25A), gonadotropin releasing hormone 1
(GNRH1, MM = 0.76, LFC = 0.4, p = 0.006, q = 0.2, R^2 = 0.4). Other notable 'floralwhite' hub
genes include the candidate gene progesterone receptor (PGR, MM = 0.81) which was also
associated with social status (Figure 2C; LFC = 0.4 p = 0.006, q = 0.7) and serotonin receptor 5A

(HTR5A, MM = 0.87, LFC = 0.4, p = 0.04, q = 0.6). Candidate gene aromatase

(LOC113993669/CYP19A1) was also upregulated in males with higher testosterone phenotype

(Figure 2C, LFC = 0.6, p = 0.02, q = 0.4, R^2 = 0.5), but was found in the 'lightcoral' co-

expression module (MM= 0.86).

Despite overall similarity in gene expression associations with testosterone phenotype

and status, some brain regions were more strongly associated with status or status × testosterone

phenotype interaction (Figure 2B). Likewise, expression in many of the 18 candidate

neuroendocrine genes were associated with status or the interaction with testosterone in different

nuclei (Figure 2C). PVN of the hypothalamus showed the most DEGs associated with social

status after FDR correction (42). Within these, there was statisticaly significant enrichment for

cytoskeletal movement genes (Figure S3.12B) involved in intracellular transport and cilia (e.g.,

GO:0003341, GO:0061640), and non-significant enrichment for cholesterol biosynthesis

(GO:0006695, GO:0045540). Candidate gene expression also revealed status-specific

associations with testosterone phenotype in the PVN, that is a statistical interaction between

status and testosterone phenotype (Figure 2C). In particular, AR expression was positively

correlated with testosterone phenotype in floater males, but not in territorial males (Figure 2F;

LFC = -0.7, p = 0.008, q = 0.9). The PVN of territorial males also showed higher VIPR1 (Figure

S4.8; LFC = 0.3, p = 0.02, q = 0.4) and lower ESR2 expression (LFC = -0.6, p = 0.006, q = 0.2).

In BSTm, a region that bridges the SDMN and MRS, there were more than 1000 putative DEGs,

but only nine after FDR correction. Among these, AR expression was higher in territorial males

than in floater males (Figure 2E; LFC = 0.6, p = 0.0007, q =0.2). Further, testosterone

metabolism gene 5-alpha reductase (SRD5A2) was upregulated in the LS of territorial males

(Figure S4.20; LFC = 0.7, p = 0.0005, q = 0.4). The VIP system is also implicated in status-

specific responses to testosterone in the TnA, where VIPR1 was upregulated in association with

testosterone in territorial, but not floater males (Figure S4.17; LFC = 1, p = 0.04 q = 0.7).

GCt was a hotbed for gene expression associated with social status and status ×

testosterone phenotype interaction (Figure 1B, Figure S3.6). In association with social status,

gene set enrichment analysis revealed there was a significant tendency for downregulation of

genes relating to synapse assembly (GO:0051965), and response estradiol (GO:0032355), and

upregulation of genes relating to RNA processing (e.g., GO:0000184, GO:0006396) and

translation (e.g., GO:0006413, GO:0006412) (Figure S3.15). While, candidate neuroendocrine

genes showed status specific expression in association with testosterone phenotype in this region

including progesterone receptor (PGR, Figure S4.15; LFC = 1.9, p= 0.003, q = 0.1), estrogen

receptor 1 (ESR1; aka Erα; LFC = 1.1 p = 0.04, q = 0.4), prolactin receptor (PRLR; LFC = 0.7, p

= 0.01, q = 0.2), and vasoactive intestinal peptide (VIP; Figure S4.15; LFC = 1.6, q = 0.005, q =

0.1). Further, candidate gene aromatase was downregulated with higher testosterone phenotype

in the GCt (LFC = -1.3, p = 0.002, q = 0.3, R^2 = 0.4).

Cooperative behavior is associated with a distinct gene expression landscape

Network strength (the number of cooperative interactions between individuals) varies

widely among territorial and floater males (Figure 1E), representing a key axis of variation in

social behavior in this species. We predicted that the associations between gene expression and

cooperative behavior would be different from those observed for social status and testosterone

phenotype. Indeed, we found a distinctive landscape of constitutive gene expression across

tissues relating to cooperative behavior (Figure 3A, Figure S3.5), as well as among candidate

genes (Figure 3B). These results are based on results from a continuous predictor and can be

interpreted as genes with significant up-or downregulation with increasing social network

strength. AH had the largest number of DEGs associated with a male's social network strength.

In this nucleus there were no significantly enriched GO terms, but there was non-significant

enrichment for genes involved in RNA metabolism (e.g., GO:0043928) and regulation of

translation (e.g., GO:0017148), development (e.g., GO:0061061, GO:0042472) and synaptic

potentiation (e.g., GO:0007274) (Figure S3.35). The pituitary also stood out in the abundance of

differentially expressed genes associated with male's social network strength (109 genes after

FDR correction q < 0.1). There were no significantly enriched GO categories after FDR

correction (Figure S3.33B), but among the non-significant terms were actin filament

organization (GO:0007015), neural differentiation (e.g., GO:0031103, GO:0045664) and the cell

cycle (e.g., GO:0043065, GO:0071158.) Among the DEGs in the pituitary, candidate genes

prolactin (PRL) (Figure 3C; LFC = -0.6, p=5x10^{-5} q = 0.03, R^2 = 0.7), ESR2 (Figure S4.5; LFC

= -0.7, p = 0.04, q = 0.5, R^2 = 0.5) and VIPR2 (Figure S4.5; LFC = -0.4, p = 0.008, q = 0.3, R^2 =

0.4) were downregulated in highly cooperative males. Intriguingly, the PVN showed a weak

correlation of one of our candidate genes: oxytocin (OXT) expression decreased with a male's

social network strength (Figure 3D; LFC = -0.3, p = 0.04, q = 0.6, R^2 = 0.2). Further, although

there was less representation of sex-steroid related pathways in the candidate gene analysis
(Figure 2B), PGR was downregulated in males with higher social network strength in BSTm
(Figure 2E; LFC = -0.3, p = 5x10^{-4}, q = 0.2, R^2 = 0.5).

Many genes in the LS were differentially expressed in association with an individual's
social network strength (98 genes after FDR correction q < 0.1). There were no GO categories
significantly enriched after FDR correction (Figure S3.42), but among the top non-significant
terms were those broadly involved in development (e.g., GO:0048538, GO:0048593), and neural
organization (e.g., GO:0099175, GO:0050771). However, the expression of our candidate genes
in the LS was not linked to social network strength. Although relatively few genes were
differentially expressed in BSTm with respect to social network strength, the LS and BSTm
shared many genes with similar patterns of expression in association with strength as shown by
RRHO analysis (Cahill et al. 2018) (Figure 3G, F). In contrast, the LS and BSTm shared few
genes with similar expression in association with testosterone phenotype (Figure 4H, I) or social
status (Figure S6.1A), suggesting that these regions are uniquely engaged in modulating
cooperative behavior. Indeed, pairwise correlations of gene expression associated with
testosterone phenotype broadly recapitulated the neuroanatomical similarity of tissues sampled
(Figure 3H) and perhaps reflects similarities in pathways related to sex-steroid signaling (Figure
2C).

Testosterone phenotype and social status are associated with systemic gene expression
We reasoned that the pleiotropic effects of testosterone and its role in social status in the
manakins (Ryder et al. 2020) would also produce systemic effects on gene expression, or similar
patterns of expression across multiple regions. Social status and testosterone phenotype were
linked to the highest number of consistently differentially expressed genes (60 and 66 genes with
median uncorrected p < 0.05 respectively), while only 10 genes were consistently differentially
expressed in association with social network strength (Supplement 6). Further, PC5 of the whole
brain analysis was significantly associated with social status in a linear mixed model that
corrected for batch confounds and repeated sampling of individuals (Figure S2.5). There was
overlap between the genes and GO categories between the consistently differentially regulated
genes with social status and testosterone phenotype and genes identified as associated with PC5.
But hereon we focus on genes related to social status in the main manuscript for simplicity.

The top two most consistently differentially regulated genes were the upregulation in the

brains of territorial males of an immune system gene CLEC2B (Figure 4A) and down regulation

of development gene ROBO4 (Figure 4B). The functional categories of these genes were

reflected in the patterns of GO biological process enrichment, though none of these terms were

significant after FDR correction (Figure 4C). There was also evidence for global differential

regulation of telomere maintenance genes (Figure 4C), with downregulation of POLD-1-like

(LOC114000574) and upregulation of RAD51 in territorial males. Further, there was evidence

for global regulation of cholesterol biosynthesis (Figure 4C), and glutathione transferases (Figure

S2.6, Table S6.2), both of which are relevant in steroid hormone biosynthesis. These findings of

global expression differences among status classes could be partly caused by the upregulation of

growth hormone (GH1) in the pituitary of territorial males (Figure 4D; $LFC = 0.8$, $p = 3x10^{-5}$, q

$= 0.01$). This result holds in the model that accounts for the interaction between social status and

mean T ($LFC = 0.9$, $p = 6x10^{-5}$, $q = 0.05$).

Discussion

Using the most comprehensive transcriptome-level sampling of the SDMN to-date

(Lopes and König 2020; Antunes et al. 2021; Bentz, Niederhuth, et al. 2021; Bentz, George, et

al. 2021; Kabelik et al. 2021), our findings highlight both modular and integrative neurogenomic

organization underlying individual variation in cooperative behavior. The relative roles of each

are central to the development and expression of any complex trait (Ketterson et al. 2009;

Lipshutz et al. 2019; Cox 2020). We discuss our findings in relation to each theme in turn.

Different landscapes of gene expression with each trait reflect modular organization of the
neuroendocrine system

Modularity is reflected in how different brain nuclei, endocrine tissues, and genetic

pathways within individuals are associated with the expression of different traits. Such

neurogenomic landscapes, characterized by the upregulation of neuroendocrine genes in some

brain regions and downregulation in others, would allow differential sensitivity to testosterone

and other signaling molecules among SDMN nuclei. These neurogenomic states can be likened

to the model of immediate early gene (IEG) activation across the SDMN after a social stimulus

(Newman 1999; Goodson 2005). In that 'landscape' model, it is the pattern of IEG expression

across multiple brain nuclei that correlates with behavioral responses, not the up or

downregulation of particular genes within a single nucleus. This original model considered this

landscape of IEG expression (and later neuropeptides) in the context of proximate responses to

social stimuli. Our findings extend this model in two ways: by considering transcriptomic

variation and how long-term neurogenomic states underlie individual variation in social

behavior.

In the landscape of constitutive gene expression in male wire-tailed manakins, total

number of DEGs and candidate gene expression implicate constitutive differences in most brain

regions, but the brain regions with higher numbers of DEGs or candidate gene expression may be

more important for mediating certain traits. Some brain regions showed associations between

gene expression and both social status and testosterone phenotype (e.g., POM) or only

testosterone phenotype (e.g., TnA, testes). This may be because these centers are involved in

testosterone mediated traits irrespective of social status (O'Connell and Hofmann 2012).

Meanwhile, others appeared to be more related to social status and status specific relationships

with testosterone phenotype (e.g., PVN, GCt), or an individual's strength of cooperative

behavior (e.g., PIT, LS). Our results also suggest that the identity and function of these genes

vary across tissues and brain regions (Figure 3GH). However, most analyses revealed DEGs

relating to neural development, synaptic potentiation or organization and genes relating to

cellular metabolism, or transcription/translation regulation. Thus, individual variation is

mediated by neurogenomic states that encode synaptic modifications and connectivity

(mechanism of memory formation), larger scale shifts in cellular activity, and neural structure

(Clayton et al. 2019; George et al. 2020)). These mechanisms may be supported or feedback on

the variation in elements of sex-steroid signaling we investigated in detail (*sensu*

"neuroendocrine action potential" (Clayton et al. 2019)).

We found ample evidence for modularity in steroid related gene expression, particularly

in association with social status, testosterone phenotype or their interaction. Unsurprisingly, we

found evidence that status-specific regulation of behavior could be modulated directly by

testosterone through AR in multiple nuclei (e.g., POM, PVN, BSTm). Higher AR expression is

linked to display performance in golden-collared manakins (Schlinger et al. 2013). Further, the

conversion of testosterone to estradiol by the enzyme aromatase and subsequent binding to

estrogen receptors is a key pathway in the regulation of testosterone-dependent male behaviors

more broadly (Schlinger and Balthazart 2013), and we found evidence for the role of aromatase

and estrogen signalling role in multiple nuclei. Our results also highlight the importance of

neural VIP expression in modulating multiple aspects of social behavior (Kingsbury and Wilson

2016; Horton, Michael, et al. 2020), as it was the only neuropeptide system with widespread

patterns of differential expression with respect to all our traits of interest.

Notably, VIP and other steroid receptors may play a role in status-specific responses to

testosterone in GCt. Although tyrosine hydroxylase (TH) neurons in the GCt have been

associated with courtship behaviors (Goodson et al. 2009; Ben-Tov et al. 2021), our finding of

association with VIP and courtship behaviors broadly speaking in GCt is novel (Kingsbury and

Wilson 2016). GCt coordinates social stimuli and downstream motor processes that control

behavior. For example, in the songbird brain, GCt links POM to the song control system and

thereby mediates androgen-dependent singing behavior (Haakenson et al. 2020). Given that

cooperative display in wire-tailed manakins requires coordinated physical (and vocal displays in

cooperative *Chiroxiphia* manakins (Trainer et al. 2002)), the status-specific differential

expression of genes for steroid receptors and steroid-sensitive neuropeptides in GCt suggest that

this SDMN region may play a unique role in mediating status-specific relationships between

testosterone and cooperative behavior.

The pattern of co-upregulation of steroid related genes in the POM of males with high

testosterone phenotype (and territorial males) point to this region's role as a potent steroid

regulator. The POM is involved in regulating male aggression, sexual behavior and parental care

(O'Connell and Hofmann 2011), and stimulates gonadal testosterone production through

gonadotropin releasing hormone (*GnRH*). Indeed, the observed increase in GNRH1 transcription

may relate to transcriptional changes indicating enhanced steroid production in the testes. In our

results, the co-upregulation of AR and GNRH1 would seem contradictory due to the negative

regulatory effects of androgens and AR on *GnRH* (Brayman et al. 2012), though these may not

interact at the cell population level or may show different temporal patterns of translation.

Regardless, activation and steroid receptor concentration in POM is associated with social status

differences and social status ascension cichlid fish (Maruska et al. 2013), and mice (Lee et al.

2022). In the same cichlid species, different preoptic cell types, including homologs for PVN,

showed higher activation in one of the two partners in cooperative territorial defense

(Weitekamp and Hofmann 2017). Indeed, testosterone can link the POM to the Hypothalamic

Pituitary Adrenal (HPA) axis via the PVN (Williamson et al. 2010), which in our study also
showed evidence of status specific gene expression, including AR. Aforementioned previous
work, and the close linkages between the PVN and POM (Williamson et al. 2010), suggest these
nuclei in wire-tailed manakins could be involved in multiple elements of behavior depending on
cell types, social contexts and timescales investigated.

The number of differentially expressed genes observed in TnA implicates this region as a
hub for neurogenomic pathways involved in regulation of testosterone mediated traits. The avian
homolog of the medial amygdala (O'Connell and Hofmann 2011), TnA integrates sensory
information and has been associated with the regulation of diverse social behaviors (Raam and
Hong 2021). Indeed, our results may reflect persistent activation of this area, based on the
upregulation of IEGs such as JUN and SYT7 in males with high testosterone phenotype, and
differential regulation of other genes involved in long-term synaptic potentiation (Clayton 2000;
Marrone et al. 2008; O'Connell et al. 2012; Clayton et al. 2019). Further, our results implicate
potential estrogenic regulation in this region via increased ESR2 in males with higher
testosterone phenotype. Indeed, estrogenic regulation through ESR1 has been shown to regulate
status-specific aggression in birds (Horton et al. 2014; Merritt et al. 2020) and prosocial behavior
in prairie voles (Cushing et al. 2008). Although ESR2 is not well characterized in birds, it may
function in the medial amygdala of mammals to regulate social recognition (Lymer et al. 2018),
and sexual regulation (Nakata et al. 2016), and is associated with higher social rank in the
forebrains of cichlid fish (Burmeister et al. 2007). Further, the differential expression of stress
steroid receptor (CRHR2) hints again at the previously unexplored role of the HPA axis in
modulating display behavior and social status in this species. Indeed, social ranks often differ in
their glucocorticoid profiles (Goymann and Wingfield 2004) or regulatory responses to
glucocorticoids (Snyder-Mackler et al. 2019). The related lance-tailed manakins (*Chiroxiphia*
lanceolata), which also display cooperatively, vary in testosterone as well as corticosterone
profiles according to social status (DuVal and Goymann 2011; Jones and DuVal 2021). Finally,
the fish homolog of TnA appears mediate aggression and cooperation in cooperative territory
defense in an actor-specific manner (Weitekamp et al. 2017; Weitekamp and Hofmann 2017),
supporting a role for modulating cooperation in wire-tailed manakins as well.

The time spent interacting with other males (social network strength), our proxy for
cooperative behavior, is associated with a unique pattern of gene expression among tissues and
brain regions. Some of these signatures are difficult to parse, for example many genes are
differentially expressed in AH, yet lack clear GO enrichment. Prior to our study AH has
associated with regulation of aggression rather than cooperation in males, and acting via AVP
receptors and VIP (O'Connell and Hofmann 2011; Horton, Michael, et al. 2020). Although we
do not find any evidence for the role of these genes in the AH, our findings may suggest that
cooperative displays are derived from displays of territorial aggression, and/or gene expression is
inhibiting aggression in these displays. Indeed, cooperative displays may act as a way of
maintaining dominance hierarchies in wire-tailed manakins.

LS showed many differentially expressed genes associated with cooperation, and highly
cooperative individuals may show differences in neural structure based on the enriched GO
terms. Patterns of expression in BSTm were similar, but weaker based on fewer statistically
significant DEGs. BSTm has neural projections into LS, and vasopressin (AVP) and VIP
neurons in the BSTm and receptors for these neuropeptides in the LS are associated with
affiliative behavior and gregariousness in birds (Kelly et al. 2011; Kelly and Goodson 2013;
Kingsbury and Wilson 2016). Those prior studies suggest a role for these brain nuclei in
processing social information acutely, whereas our study suggests differences among individuals
in cooperative tendency is related to constitutive differences in gene expression and neural
structure, perhaps relating to excitability of these regions in response to social information. Thus,
cooperative display is regulated by some of the same conserved brain regions as other types of
affiliative behavior. That we only observed a large number of DEGs in the LS and not in other
nuclei may suggest that other unsampled nuclei in the mesolimbic reward system are important
for modulation of cooperative tendency (e.g., (Paula et al. 2015; Weitekamp et al. 2017; de
Abreu et al. 2020; Lopes and König 2020).

We did not find convincing evidence for any variation in sensitivity to nonapeptides
(vasopressin and oxytocin) in the brain (AVPR2 in hypothalamic nuclei showed transcript counts
<30). However, the localization of these receptors in particular brain regions may be more
important for modulating behavior than a change in their abundance (Goodson and Thompson
2010). We did find evidence for differential sensitivity to AVP in the pituitary, again implicating

stress response pathways in mediating social status and cooperation in this species. Further, we
did find a weak negative relationship between oxytocin expression and social network strength in
the PVN. Although in apparent contrast to the widely held belief that oxytocin promotes sociality
(as in humans (Feldman 2012)), this among our other observations may indicate that cooperative
displays are a derivative of aggressive displays. However, nonapeptides have complex and
sometimes species-specific relationships with affiliative behavior (Goodson and Thompson
2010; Maruska et al. 2022), and can also be associated with social status (Lee et al. 2022).
Further, the ratio of peptide to mRNA can be an important factor indicating higher turnover
(usage) in certain contexts (Maruska et al. 2022). Therefore, we do not have sufficient evidence
that oxytocin inhibits cooperation in this species, and further experimental work is needed.

Our results establish a variety of mechanisms that could underpin individual variation in
cooperative behavior, and its status specific modulation by testosterone. We find evidence for
multiple forms of neuroplasticity, through steroid receptor variation, synaptic potentiation and
neural development, as well as cellular activity (Clayton et al. 2019). Indeed, testosterone
manipulations are known to induce changes in neuroanatomy in behaviorally relevant brain
nuclei (Kabelik et al. 2008), suggesting a direct role for testosterone-mediated neural growth and
flexibility in modulating male-male cooperation (Soares et al. 2010). Our results provide
promising targets for further study on the mechanisms underlying cooperative behavior,
including examination of neuroanatomical features and the use of hormonal or other
pharmaceutical manipulations and RNAi technologies to examine causal relationships between
neural gene expression and behavior (e.g., (Merritt et al. 2020)).

Similarities in gene expression reflect the role of testosterone as a phenotypic
integrator.

Similarities in gene expression among tissues are predicted as a result of the pleiotropic
effects of testosterone and support the role of the hormone as a mediator of phenotypic
integration. The extent of gene expression similarities and differences among tissues may reflect
the degree of integration and independence (or modularity), respectively, that can either
constrain or facilitate adaptive evolution (Ketterson et al. 2009; Lipshutz et al. 2019). We
identified similarities in testosterone associated gene expression between brain regions that
broadly recapitulated tissue ontology. We also identified consistent, brain-wide (systemic) gene

expression differences associated predominantly with social status and testosterone phenotype. However, systemic results did not explain as much variation in overall neural gene expression as with some axes of personality (Kabelik et al. 2021; Lattin et al. 2021), suggesting that the brain region-specific patterns may be more important in modulating male wire-tailed manakin behavior. Moreover, the broader associations of social status with metabolic, immune, and telomere gene ontology categories suggests differences in allostatic load among status classes in male-male cooperative systems (Goymann and Wingfield 2004; Jones and DuVal 2021; Vernasco et al. 2021). Further, differential global regulation of telomeres may explain the relative stability of telomere length in territorial wire-tailed manakin males (Vernasco et al. 2021), and collectively these results may represent metabolic adaptations for territoriality.

Our results highlight two additional hormones that may have a system-wide role in the status-specific modulation of cooperative display behavior. We found differences in the expression of genes for important behavioral and metabolic hormones (prolactin and growth hormone) in the pituitary, and may reflect organism wide metabolic adaptations. Indeed, previous work on this species showed that more social males had shorter telomeres (Vernasco et al. 2021), and social bond strength has been associated with signatures of energy metabolism and stress (Anderson et al. 2022; Simons et al. 2022). The anterior pituitary is the primary production site for both hormones, and thus gene expression patterns here are likely good predictors for circulating levels of those hormones and thus signal strength. PRL transcript was downregulated in more cooperative males. Prolactin is best known for its role in promoting parental care behaviors, including in cooperative breeders (Schoech et al. 1996), but emerging studies suggest it could play a role in inhibiting aggression in cooperative breeders (Medger et al. 2019; Gilbert et al. 2022) and promotion of nest defense behavior (Mohamed et al. 2016). However, the negative relationship observed appears contradictory to a general model of increased prolactin promoting broadly affiliative social interactions, and again may reflect the origin of these cooperative displays from aggressive or competitive interactions (Díaz-Muñoz et al. 2014). The other compelling phenotypic integrator for status-dependent behaviors in the wire-tailed manakin comes from the gene encoding growth hormone (GH1). Ascension in social rank in the cooperatively displaying manakins is associated with age (McDonald 1989; DuVal 2007; Ryder et al. 2008). It was therefore surprising that territorial males in our study showed increased expression of GH1 in the pituitary as compared to floater males. However, in anolis lizards,

stimulation of the growth hormone axis is associated with increased testosterone that enables

phenotypic divergence between males and females (Cox et al. 2017). Without gross

morphological differences in wire-tailed manakin status-classes, growth-hormone could have

subtle effects on multiple systems enabling the status-specific regulation of behaviors. In

particular, status differences in GH1 expression may explain repeated signatures of neural

development and protein synthesis in our results (Sonksen 2006), and may reflect the increased

metabolic demands on displaying territorial males through muscle performance and regulation of

oxidative stress (Hu et al. 2019). The neuroethological role of growth hormone are not well

known, but its release may be linked to stress (Greenwood and Landon 1966; McCormick et al.

1998). Growth hormone administration promotes increased aggression in mice (Matte 1981), and

fish (Jönsson et al. 1998; Johansson et al. 2004). Indeed, growth hormone inhibitor somatostatin

and its receptors have also been implicated in regulating aggression and social status in cichlid

fish independent of androgens (Hofmann and Fernald 2000; Trainor and Hofmann 2006) whilst

natal androgen treatment increased aggression and somatostatin receptor expression in zebra

finches (Bentz, Niederhuth, et al. 2021). We did some evidence for negative relationship

between SSTR1 and SSTR2 with testosterone phenotype in the testes, and positive association

between SSTR1 and testosterone phenotype in the POM but these were not significant after FDR

correction. Thus, our results suggest that the growth hormone axis may play a role in the status-

specific regulation of wire-tailed manakin cooperative displays, and increased GH in territorial

males may be an important inhibitor of cooperation by promoting aggression. Future work on

social behavior should consider the role of growth hormone and somatostatin in conjunction with

sex steroids and glucocorticoids to gain novel insights into its function in behavioral regulation.

Conclusion

Across the SDMN and HPG of the male wire-tailed manakin we observed correlations with

social status, cooperation and testosterone phenotype in patterns of gene regulation and a

plurality of gene functions, including sex-steroid and neuropeptide signaling. This landscape of

gene expression speaks to the modular nature of behavioral mechanisms (Newman 1999;

Goodson 2005; Lipshutz et al. 2019) and highlights the continuum between complete phenotypic

integration and modularity (Ketterson et al. 2009). Much of the previous work on the

mechanisms of social and cooperative behaviors focus on relatively few candidate genes in

broad-scale brain divisions (e.g., hypothalamus) or in one or a few particular nuclei. Our work
presented here is one of the most comprehensive examinations to date of region-specific gene
expression in all nuclei of the SDMN, albeit correlative only. Our correlative findings help us to
better understand cooperative behavior (Soares et al. 2010) by linking neurogenomics with
endocrine mechanisms, and generate further hypotheses to test in further experiments to
implicate causality. We further demonstrate and emerging idea that constitutive landscapes in
gene expression across the brain are associated with repeatable differences in behavioral
phenotype (Antunes et al. 2021; Kabelik et al. 2021; Lattin et al. 2021). We also provide
tantalizing glimpses into nuclei and genomic pathways involved in the status-specific regulation
of this cooperative display behavior. Although questions remain on the acute mediators of
cooperative display behavior in the wire-tailed manakin, we demonstrate the power and
flexibility of a system-wide approach. We propose that landscapes of constitutive gene
expression for multiple genes and across many tissues and brain regions, like those observed
here, form the foundation of consistent inter-individual differences in any behavior, upon which
flexible and acute responses to social stimuli are built.

Acknowledgements
We want to acknowledge long hours of fieldwork in Ecuador by Ben Vernasco and Camilo
Alfonso-Cuta that made the behavioral foundation of this work. Thank you to them, and the staff
at Tiputini Biodiversity Station. Thank you to Eric Shuppe and to Matthew Fuxjager for use of
laboratory resources. Thank you to David Clayton, Julie George, Sarah London, and Barney
Schlinger for helpful discussions and manuscript reviewing, and Vanessa González for advice on
data analysis. **Funding** for the reference genome came from a private donor at Millersville
University. Funding for the behavioral and RNAseq study came from the National Science
Foundation (IOS 1353085; DBI 1457541), the Smithsonian Migratory Bird Center, the Global
Change Center at Virginia Tech, Millersville University, and the Tiputini Biodiversity Station of
the Universidad San Francisco de Quito.

Data and materials availability: Raw read data are available associated with BioProject
PRJNA437157, and all code used to conduct analyses is available here:
https://github.com/periperipatus/PIFI_brain_transcriptome

References

de Abreu MS, Maximino C, Cardoso SC, Marques CI, Pimentel AFN, Mece E, Winberg S, Barcellos LJG, Soares MC. 2020. Dopamine and serotonin mediate the impact of stress on cleaner fish cooperative behavior. *Horm. Behav.* 125.

Anderson JA, Lea AJ, Voyles TN, Akinyi MY, Nyakundi R, Ochola L, Omondi M, Nyundo F, Zhang Y, Campos FA, et al. 2022. Distinct gene regulatory signatures of dominance rank and social bond strength in wild baboons. *Philos. Trans. R. Soc. B* [Internet] 377. Available from: https://royalsocietypublishing.org/doi/10.1098/rstb.2020.0441

Antunes DF, Teles MC, Zuelling M, Friesen CN, Oliveira RF, Aubin-Horth N, Taborsky B. 2021. Early social deprivation shapes neuronal programming of the social decision-making network in a cooperatively breeding fish. *Mol. Ecol.* [Internet] 30:4118–4132. Available from: https://onlinelibrary.wiley.com/doi/full/10.1111/mec.16019

Ben-Tov M, Duarte F, Mooney R. 2021. A neural hub that coordinates learned and innate courtship behaviors. *bioRxiv* [Internet]:2021.09.09.459618. Available from: https://www.biorxiv.org/content/10.1101/2021.09.09.459618v1

Bentz AB, George EM, Wolf SE, Rusch DB, Podicheti R. 2021. Experimental competition induces immediate and lasting effects on the neurogenome in free-living female birds. *Proc. Natl. Acad. Sci. USA* 118:e2016154118.

Bentz AB, Niederhuth CE, Carruth LL, Navara KJ. 2021. Prenatal testosterone triggers long-term behavioral changes in male zebra finches: unravelling the neurogenomic mechanisms. *BMC Genomics* 22:158.

Blighe K, Lun A. 2020. PCAtools: Everything Principal Components Analysis.

Brayman MJ, Pepa PA, Mellon PL. 2012. Androgen Receptor Repression of Gonadotropin-Releasing Hormone Gene Transcription via Enhancer 1. *Mol. Cell. Endocrinol.* [Internet] 363:92. Available from: /pmc/articles/PMC3447085/

Burmeister SS, Kailasanath V, Fernald RD. 2007. Social dominance regulates androen and estrogen receptor gene expression. *Bone* 51:164–170.

Cahill K, Huo Z, Tseng GC, Logan RW, Seney ML. 2018. Improved identification of concordant and discordant gene expression signatures using an updated rank-rank hypergeometric overlap approach. *Sci. Rep.* 8:9588.

Clayton DF. 2000. The genomic action potential. *Neurobiol. Learn. Mem.* 74:185–216.

Clayton DF, Anreiter I, Aristizabal M, Frankland PW, Binder EB, Citri A. 2019. The role of the genome in experience-dependent plasticity: Extending the analogy of the genomic action potential. *Proc. Natl. Acad. Sci.* 117:201820837.

Cox RM. 2020. Sex steroids as mediators of phenotypic integration, genetic correlations, and evolutionary transitions. *Mol. Cell. Endocrinol.* [Internet] 502:110668. Available from: https://doi.org/10.1016/j.mce.2019.110668

Cox RM, Cox CL, McGlothlin JW, Card DC, Andrew AL, Castoe TA. 2017. Hormonally Mediated Increases in Sex-Biased Gene Expression Accompany the Breakdown of Between-Sex Genetic Correlations in a Sexually Dimorphic Lizard. *https://doi.org/10.1086/690105* [Internet] 189:315–332. Available from: https://www.journals.uchicago.edu/doi/10.1086/690105

Cushing BS, Perry A, Musatov S, Ogawa S, Papademetriou E. 2008. Estrogen Receptors in the Medial Amygdala Inhibit the Expression of Male Prosocial Behavior. *J. Neurosci.* [Internet] 28:10399–10403. Available from: https://www.jneurosci.org/content/28/41/10399

Dakin R, Ryder TB. 2018. Dynamic network partnerships and social contagion drive cooperation. *Proc. R. Soc. B Biol. Sci.* 285:20181973.

Díaz-Muñoz SL, DuVal EH, Krakauer AH, Lacey EA. 2014. Cooperating to compete: Altruism, sexual selection and causes of male reproductive cooperation. *Anim. Behav.* [Internet] 88:67–78. Available from: http://dx.doi.org/10.1016/j.anbehav.2013.11.008

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29:15–21.

DuVal EH. 2007. Adaptive advantages of cooperative courtship for subordinate male lance-tailed manakins. *Am. Nat.* 169:423–432.

DuVal EH, Goymann W. 2011. Hormonal correlates of social status and courtship display in the cooperatively lekking lance-tailed manakin. *Horm. Behav.* [Internet] 59:44–50. Available from: http://dx.doi.org/10.1016/j.yhbeh.2010.10.004

Feldman R. 2012. Oxytocin and social affiliation in humans. *Horm. Behav.* [Internet] 61:380–391. Available from: https://pubmed.ncbi.nlm.nih.gov/22285934/

Fusani L, Donaldson Z, London SE, Fuxjager MJ, Schlinger BA. 2014. Expression of androgen receptor in the brain of a sub-oscine bird with an elaborate courtship display. *Neurosci. Lett.* [Internet] 578:61–65. Available from: http://dx.doi.org/10.1016/j.neulet.2014.06.028

George JM, Bell ZW, Condliffe D, Dohrer K, Abaurrea T, Spencer K, Leitão A, Gahr M, Hurd PJ, Clayton DF. 2020. Acute social isolation alters neurogenomic state in songbird forebrain. *Proc. Natl. Acad. Sci.* 117:201820841.

Gilbert JD, Rossiter SJ, Bennett NC, Faulkes CG. 2022. The elusive role of prolactin in the sociality of the naked mole-rat. *Horm. Behav.* 143:105196.

Goodson JL. 2005. The vertebrate social behaviour network: evolutionary themes and variations. *Horm. Behav.* 48:11–22.

Goodson JL, Kabelik D, Kelly AM, Rinaldi J, Klatt JD. 2009. Midbrain dopamine neurons reflect affiliation phenotypes in finches and are tightly coupled to courtship. *Proc. Natl. Acad. Sci.* [Internet] 106:8737–8742. Available from: https://www.pnas.org/content/106/21/8737

Goodson JL, Kingsbury MA. 2013. What's in a name? Considerations of homologies and nomenclature for vertebrate social behavior networks. *Horm. Behav.* 64:103–112.

Goodson JL, Thompson RR. 2010. Nonapeptide mechanisms of social cognition, behavior and species-specific social systems. *Curr. Opin. Neurobiol.* 20:784–794.

Goymann W, Wingfield JC. 2004. Allostatic load, social status and stress hormones: the costs of social status matter. *Anim. Behav.* 67:591–602.

Greenwood FC, Landon J. 1966. Growth hormone secretion in response to stress in man. *Nature* 210:540–541.

Griffith SC, Crino OL, Andrew SC, Nomano FY, Adkins-Regan E, Alonso-Alvarez C, Bailey IE, Bittner SS, Bolton PE, Boner W, et al. 2017. Variation in Reproductive Success Across Captive Populations: Methodological Differences, Potential Biases and Opportunities. *Ethology* [Internet] 123:1–29. Available from: http://doi.wiley.com/10.1111/eth.12576

Haakenson CM, Balthazart J, Ball GF. 2020. Cognition and Behavior Effects of Inactivation of the Periaqueductal Gray on Song Production in Testosterone-Treated Male Canaries (*Serinus canaria*). *eNeuro* [Internet] 7:ENEURO.0048-20.2020. Available from: https://doi.org/10.1523/ENEURO.0048-20.2020

Hofmann H a., Beery AK, Blumstein DT, Couzin ID, Earley RL, Hayes LD, Hurd PL, Lacey E a., Phelps SM, Solomon NG, et al. 2014. An evolutionary framework for studying mechanisms of social behavior. *Trends Ecol. Evol.* [Internet] 29:1–9. Available from: http://linkinghub.elsevier.com/retrieve/pii/S0169534714001608

Hofmann HA, Fernald RD. 2000. Social Status Controls Somatostatin Neuron Size and Growth. *J. Neurosci.* [Internet] 20:4740–4744. Available from: https://www.jneurosci.org/content/20/12/4740

Horton BM, Hudson WH, Ortlund EA, Shirk S, Thomas JW, Young ER, Zinzow-Kramer WM, Maney DL. 2014. Estrogen receptor α polymorphism in a species with alternative behavioral phenotypes. *Proc. Natl. Acad. Sci. U. S. A.* [Internet] 111:1443–1448. Available from: http://www.pnas.org/cgi/doi/10.1073/pnas.1317165111

Horton BM, Michael C, Mackenzie P, Maney DL. 2020. Vasoactive intestinal peptide as a mediator of the effects of a supergene on social behavior. *Proc. R. Soc. B Biol. Sci.* 287:20200196.

Horton BM, Ryder TB, Moore IT, Balakrishnan CN. 2020. Gene expression in the social behavior network of the wire-tailed manakin (*Pipra filicauda*) brain. *Genes, Brain Behav.*:e12560.

Hu B, Hu S, Yang M, Liao Z, Zhang D, Luo Q, Zhang X, Li H. 2019. Growth hormone receptor gene is essential for chicken mitochondrial function in vivo and in vitro. *Int. J. Mol. Sci.* 20:1–13.

Johansson V, Winberg S, Jönsson E, Hall D, Björnsson B. 2004. Peripherally administered growth hormone increases brain dopaminergic activity and swimming in rainbow trout. *Horm. Behav.* 46:436–443.

Jones B, DuVal EH. 2021. Glucocorticoids correlate with and predict social status in the cooperatively breeding lance-tailed manakin (*Chiroxiphia lanceolata*). In: Integrative and Comparative Biology. Vol. 61. p. E428-429.

Jönsson E, Johnsson J, Björnsson B. 1998. Growth hormone increases aggressive behavior in juvenile rainbow trout. *Horm. Behav.* 33:9–15.

Kabelik D, Julien AR, Ramirez D, O'Connell LA. 2021. Social boldness correlates with brain gene expression in male green anoles. *Horm. Behav.* 133:105007.

Kabelik D, Weiss SL, Moore MC. 2008. Steroid hormones alter neuroanatomy and aggression independently in the tree lizard. *Physiol. Behav.* [Internet] 93:492. Available from: /pmc/articles/PMC4286361/

Kasper C, Vierbuchen M, Ernst U, Fischer S, Radersma R, Raulo A, Cunha-Saraiva F, Wu M, Mobley KB, Taborsky B. 2017. Genetics and developmental biology of cooperation. *Mol. Ecol.* 26:4364–4377.

Kelly AM, Goodson JL. 2013. Behavioral relevance of species-specific vasotocin anatomy in gregarious finches. *Front. Neurosci.* [Internet] 7:242. Available from: /pmc/articles/PMC3865460/

Kelly AM, Kingsbury MA, Hoffbuhr K, Schrock SE, Waxman B, Kabelik D, Thompson RR, Goodson JL. 2011. Vasotocin neurons and septal V1a-like receptors potently modulate songbird flocking and responses to novelty. *Horm. Behav.* 60:12–21.

Ketterson ED, Atwell JW, McGlothlin JW. 2009. Phenotypic integration and independence: Hormones, performance, and response to environmental change. *Integr. Comp. Biol.* 49:365–379.

Kingsbury MA, Wilson LC. 2016. The Role of VIP in Social Behavior: Neural Hotspots for the Modulation of Affiliation, Aggression, and Parental Care. *Integr. Comp. Biol.* [Internet] 56:1238–1249. Available from: https://academic.oup.com/icb/article/56/6/1238/2647097

Langfelder P, Horvath S. 2008. WGCNA: An R package for weighted correlation network analysis. *BMC Bioinformatics* 9:559.

Lattin CR, Kelly TR, Kelly MW, Johnson KM. 2021. Constitutive gene expression differs in three brain regions important for cognition in neophobic and non-neophobic house sparrows (*Passer domesticus*). *bioRxiv* [Internet]:2021.07.06.451290. Available from: https://www.biorxiv.org/content/10.1101/2021.07.06.451290v1

Lee W, Dwortz MF, Milewski TM, Champagne FA, Curley JP. 2022. Social status mediated variation in hypothalamic transcriptional profiles of male mice. *Horm. Behav.* 142:105176.

Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* 30:923–930.

Lipshutz SE, George EM, Bentz AB, Rosvall KA. 2019. Evaluating testosterone as a phenotypic integrator: From tissues to individuals to species. *Mol. Cell. Endocrinol.* [Internet] 496:110531. Available from: https://doi.org/10.1016/j.mce.2019.110531

Lopes PC, König B. 2020. Wild mice with different social network sizes vary in brain gene expression. *BMC Genomics* 21:1–14.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol.* 15:550.

Loveland JL, Giraldo-Deck LM, Lank DB, Goymann W, Gahr M, Küpper C. 2021. Functional differences in the hypothalamic-pituitary-gonadal axis are associated with alternative reproductive tactics based on an inversion polymorphism. *Horm. Behav.* [Internet] 127:104877. Available from: https://doi.org/10.1016/j.yhbeh.2020.104877

Lymer JM, Sheppard PAS, Kuun T, Blackman A, Jani N, Mahbub S, Choleris E. 2018. Estrogens and their receptors in the medial amygdala rapidly facilitate social recognition in female mice. *Psychoneuroendocrinology* 89:30–38.

Marrone DF, Schaner MJ, Mcnaughton BL, Worley PF, Barnes CA. 2008. Immediate-Early Gene expression at rest recapitulates recent experience. *J. Neurosci.* 28:1030–1033.

Maruska KP, Anselmo CM, King T, Mobley RB, Ray EJ, Wayne R. 2022. Endocrine and neuroendocrine regulation of social status in cichlid fishes. *Horm. Behav.* 139:105110.

Maruska KP, Zhang A, Neboori A, Fernald RD. 2013. Social Opportunity Causes Rapid Transcriptional Changes in the Social Behaviour Network of the Brain in an African Cichlid Fish. *J. Neuroendocrinol.* 25:145–157.

Matte AC. 1981. Growth hormone and isolation-induced aggression in wild male mice. *Pharmacol. Biochem. Behav.* [Internet] 14 Suppl 1:85–87. Available from: https://pubmed.ncbi.nlm.nih.gov/7195589/

McCormick SD, Shrimpton JM, Carey JB, O'Dea MF, Sloan KE, Moriyama S, Björnsson BT. 1998. Repeated acute stress reduces growth rate of Atlantic salmon parr and alters plasma levels of growth hormone, insulin-like growth factor I and cortisol. *Aquaculture* 168:221–235.

McDonald DB. 1989. Cooperation Under Sexual Selection : Age-Graded Changes in a Lekking Bird. *Am. Nat.* 134:709–730.

Medger K, Bennett NC, Ganswindt SB, Ganswindt A, Hart DW. 2019. Changes in prolactin, cortisol and testosterone concentrations during queen succession in a colony of naked mole-rats (Heterocephalus glaber): a case study. *Naturwissenschaften* [Internet] 106. Available from: https://pubmed.ncbi.nlm.nih.gov/31089819/

Merritt JR, Grogan KE, Zinzow-kramer WM, Sun D, Ortlund EA. 2020. A supergene-linked estrogen receptor drives alternative phenotypes in a polymorphic songbird. :1–8.

Mohamed RA, Shukry M, Mousa-balabel TM, Elbassiouny AA. 2016. Assessment of Plasma Prolactin and Nest Defense Behaviour During Breeding Cycle of Pigeon (Columba livia domestica). *J. Environ. Agric. Sci.* 7:19–22.

Nakata M, Sano K, Musatov S, Yamaguchi N, Sakamoto T, Ogawa S. 2016. Effects of Prepubertal or Adult Site-Specific Knockdown of Estrogen Receptor β in the Medial Preoptic Area and Medial Amygdala on Social Behaviors in Male Mice. *eNeuro* [Internet] 3:579–588. Available from: /pmc/articles/PMC4819287/

Newman SW. 1999. The medial extended amygdala in male reproductive behavior: A node in the mammalian social behavior network. *Ann. N. Y. Acad. Sci.* 877:242–257.

O'Connell LA, Hofmann HA. 2011. The Vertebrate mesolimbic reward system and social behavior network: A comparative synthesis. *J. Comp. Neurol.* 519:3599–3639.

O'Connell LA, Hofmann HA. 2012. Social Status Predicts How Sex Steroid Receptors Regulate Complex Behavior across Levels of Biological Organization. *Endocrinology* [Internet] 153:1341–1351. Available from: https://academic.oup.com/endo/article/153/3/1341/2424042

O'Connell LA, Matthews BJ, Hofmann HA. 2012. Isotocin regulates paternal care in a monogamous cichlid fish. *Horm. Behav.* 61:725–733.

Olds J, Milner P. 1954. Positive reinforcement produced by electrical stimulation of septal area and other regions of rat brain. *J. Comp. Physiol. Psychol.* [Internet] 47:419–427. Available from: /record/1955-06866-001

Paula JR, Messias JP, Grutter AS, Bshary R, Soares MC. 2015. The role of serotonin in the modulation of cooperative behavior. *Behav. Ecol.* [Internet] 26:1005–1012. Available from: https://academic.oup.com/beheco/article/26/4/1005/209523

Pinheiro J, Bates D, DebRoy S, Sarkar D. 2016. Linear and Nonlinear Mixed Effects Models. R. package v 3.1-128.

R Core Team. 2020. R: A language and environment for statistical computing. Available from: https://www.r-project.org/

Raam T, Hong W. 2021. Organization of neural circuits underlying social behavior: a consideration of the medial amygdala. *Curr. Opin. Neurobiol.* [Internet] 68:124. Available from: /pmc/articles/PMC8243811/

Ryder TB, Blake JG, Parker PG, Loiselle BA. 2011. The composition, stability, and kinship of reproductive coalitions in a lekking bird. *Behav. Ecol.* 22:282–290.

Ryder TB, Dakin R, Vernasco BJ, Evans BS, Horton BM, Moore IT. 2020. Testosterone modulates status-specific patterns of cooperation in a social network. *Am. Nat.* 195:82–94.

Ryder TB, Horton BM, Moore IT. 2011. Understanding testosterone variation in a tropical lek-breeding bird. *Biol. Lett.* 7:506–509.

Ryder TB, Horton BM, Van Den Tillaart M, De Dios Morales J, Moore IT. 2012. Proximity data-loggers increase the quantity and quality of social network data. *Biol. Lett.* 8:917–920.

Ryder TB, McDonald DB, Blake JG, Parker PG, Loiselle BA. 2008. Social networks in the lek-mating wire-tailed manakin (*Pipra filicauda*). *Proc. R. Soc. B Biol. Sci.* 275:1367–1374.

Ryder TB, Parker PG, Blake JG, Loiselle BA. 2009. It takes two to tango: Reproductive skew and social correlates of male mating success in a lek-breeding bird. *Proc. R. Soc. B Biol. Sci.* 276:2377–2384.

Schlinger B, Balthazart J. 2013. Aromatase and Behavior: Concepts Gained from Studies of Aromatase in the Avian Brain. In: Brain Aromatase, Estrogens, and Behavior. Available from: https://psycnet.apa.org/record/2012-25571-010

Schlinger BA, Barske J, Day L, Fusani L, Fuxjager MJ. 2013. Hormones and the neuromuscular

control of courtship in the golden-collared manakin (*Manacus vitellinus*). *Front. Neuroendocrinol.* [Internet] 34:143. Available from: /pmc/articles/PMC3995001/

Schoech SJ, Mumme L R, Wingfield.ast C J. 1996. Prolactin and helping behaviour in the cooperatively breeding Florida scrub-jay, *Aphelocoma c. coerulescens*. *Anim. Behav.* 52:445–456.

Simons ND, Michopoulos V, Wilson M, Barreiro LB, Tung J. 2022. Agonism and grooming behaviour explain social status effects on physiology and gene regulation in rhesus macaques. *Philos. Trans. R. Soc. B Biol. Sci.* 377.

Snyder-Mackler N, Sanz J, Kohn JN, Voyles T, Pique-Regi R, Wilson ME, Barreiro LB, Tung J. 2019. Social status alters chromatin accessibility and the gene regulatory response to glucocorticoid stimulation in rhesus macaques. *Proc. Natl. Acad. Sci. U. S. A.* [Internet] 116:1219–1228. Available from: https://www.pnas.org/doi/abs/10.1073/pnas.1811758115

Soares MC, Bshary R, Fusani L, Goymann W, Hau M, Hirschenhauser K, Oliveira RF. 2010. Hormonal mechanisms of cooperative behaviour. *Philos. Trans. R. Soc. B Biol. Sci.* 365:2737–2750.

Sonksen P. 2006. Growth Hormone Replacement Therapy and Life Quality. Future Perspectives. In: Nyberg F, editor. The Somatotrophic Axis in Brain Function. Elsevier Inc. p. 327–333.

Taborsky M, Taborsky B. 2015. Evolution of genetic and physiological mechanisms of cooperative behaviour. *Curr. Opin. Behav. Sci.* 6:132–138.

Trainer JM, McDonald DB, Learn WA. 2002. The development of coordinated singing in cooperatively displaying long-tailed manakins. *Behav. Ecol.* 13:65–69.

Trainor BC, Hofmann HA. 2006. Somatostatin Regulates Aggressive Behavior in an African Cichlid Fish. *Endocrinology* [Internet] 147:5119–5125. Available from: https://academic.oup.com/endo/article/147/11/5119/2500141

Vernasco BJ, Dakin R, Majer AD, Haussmann MF, Ryder TB, Moore IT. 2021. Longitudinal dynamics and behavioural correlates of telomeres in male wire-tailed manakins. *Funct. Ecol.* [Internet] 35:450–462. Available from: https://onlinelibrary.wiley.com/doi/full/10.1111/1365-2435.13715

Vernasco BJ, Horton BM, Moore IT, Ryder TB. 2020. Reduced cooperative behavior as a cost of high testosterone in a lekking passerine bird. *Behav. Ecol.* [Internet] 31:401–410. Available from: https://academic.oup.com/beheco/advance-article/doi/10.1093/beheco/arz201/5673236

Weitekamp CA, Hofmann HA. 2017. Neuromolecular correlates of cooperation and conflict during territory defense in a cichlid fish. *Horm. Behav.* 89:145–156.

Weitekamp CA, Nguyen J, Hofmann HA. 2017. Neuromolecular regulation of aggression differs by social role during joint territory defense. *Integr. Comp. Biol.* 57:631–639.

Williamson M, Bingham B, Gray M, Innala L, Viau V. 2010. The Medial Preoptic Nucleus Integrates the Central Influences of Testosterone on the Paraventricular Nucleus of the Hypothalamus and Its Extended Circuitries. *J. Neurosci.* [Internet] 30:11762. Available

from: /pmc/articles/PMC6633408/

Wingfield JC, Lynn SE, Soma KK. 2001. Avoiding the 'Costs' of Testosterone: Ecological Bases of Hormone-Behavior Interactions. *Brain. Behav. Evol.* [Internet] 57:239–251. Available from: https://www.karger.com/Article/FullText/47243

Yu G, Wang L-G, Han Y, He Q-Y. 2012. clusterProfiler: an R package for comparing biological themes among gene clusters. *Omi. A J. Integr. Biol.* 16:284–287.

Fig. 1. Circulating testosterone and neuroendocrine tissues are involved in the cooperative behavior of wire-tailed manakins.

A) Males engage in cooperative displays that form the basis of B) complex social networks. The example shown is a single lek over one week, where node size indicates a male's average social network strength (measure of overall time spent interacting), and edge width represents the frequency of interactions between a pair of individuals. Node color denotes floater (red) and territorial (gray) male status, where darker colors were included in the study. C) A male's testosterone phenotype (mean testosterone (corrected)) influences his cooperative tendency differently in territorial and floater classes. D) Twelve individuals (points) were from the larger population with repeated measures of testosterone (density), and E) proximity-data logging measures of social network strength. Note the total sample size for individuals characterized for social status is N=16 is larger than for testosterone or sociality (N=12). F) RNAseq characterized gene expression from each individual in 10 testosterone-sensitive brain regions and two endocrine tissues. Structures shown are not to scale. Colors represent functional groups (O'Connell and Hofmann 2011). G & inset) principal component analysis of transcriptomic data reveals distinct gene expression profiles among brain regions and endocrine tissues (colors and shapes as F). (photo by Alice Boyle).

Fig. 2. Neurogenomic landscape of social status and hormone phenotype.
A) There is a landscape of gene expression in male wire-tailed manakins with high testosterone phenotype (mean testosterone (corrected)) relative to lower testosterone phenotype. Red bars indicate number of genes that are upregulated in males with higher testosterone, whereas blue indicates genes that are downregulated in males with higher testosterone. Darker colors and text indicate number of differentially expressed genes after FDR correction. B) The landscape of gene expression in relation to social status, where red indicates genes upregulated in territorial males. C) Summary of candidate gene analysis and filtering, where colors indicate direction of expression (as previous panels), but intensity reflects which of the models were significant, and "i" indicates there was an interaction effect, and a black box indicates result was significant after FDR correction for all genes in that region's transcriptome. D) AR was upregulated in POM of males with higher testosterone. where point colors indicate social status (red = floater; black = territorial. Fitted line from linear regression. Subheader indicates key results from DESeq2: Log$_2$ Fold Change (LFC), uncorrected p-value (p), and correction for multiple testing (q) and R^2 correlation coefficient calculated separately. E) AR was upregulated in the BSTm of territorial males. F) AR shows status-specific expression in relation to testosterone phenotype in the PVN. Grey area represents the 95% confidence interval as determined by linear regression of gene ~ Status×mean_T. Note that maximum sample size for analysis of social status was N=16, while

only 12 individuals were sampled for testosterone and behavior, but the actual sample size varies slightly in each tissue owing to quality filtering.

Fig. 3. A distinct landscape of gene expression in relation to cooperative behavior.
A) Landscape of gene expression in male more cooperative wire-tailed manakins (social network strength). Darker colors and text indicate number of genes after FDR correction. Red indicates genes that are upregulated in males with higher social network strength, whereas blue indicates genes that are downregulated in these males. B) Summary of candidate gene analysis and filtering, where colors indicate direction of expression (as previous panels) in relation to individuals with higher social network strength. A black box indicates the result remained significant after FDR correction for all genes in that region's transcriptome. C) Prolactin (PRL) in the pituitary shows decreased expression in males with higher social network strengths. D) Oxytocin (OXT) transcript expression decreases in males with higher social network strength. E) Progesterone Receptor (PGR) has lower transcript expression in males with higher social network strength. F) RRHO2 analysis reveals similar profiles of gene expression associated with cooperation in the LS and BSTm, G) but otherwise generally low similarity of gene expression among tissues in association with cooperation. H) Similarity of gene expression among tissues in relation to testosterone phenotype roughly recapitulates ontogeny of brain regions, I) and there is no correlation between testosterone related gene expression profiles in the LS and BSTm. G,H) Cell color and number are the Pearson correlation coefficients of gene expression between tissues, correlating $\log_{10}$ transformed differential expression p-values (times the direction of

differential expression). Black boxes outline significant tissue correlation clusters based on 1,000 bootstrap resamples of the correlation matrix without FDR correction.

Fig. 4: Global neural gene expression differences between social classes
A) Immune gene CLEC2B-like was consistently upregulated in the brains of territorial individuals, and B) developmental gene ROBO4 was consistently downregulated in the brains of territorial males. C) Enriched GO categories among genes that were consistently up- or downregulated between territorial and floater males, where bar color indicates uncorrected p-value. D) These broader signatures of developmental differences could be explained by upregulation of growth hormone transcript in the pituitary of territorial males.