Neurogenomic landscape associated with status-dependent cooperative behavior

Peri E. Bolton[1,2*], T. Brandt Ryder[3,4], Roslyn Dakin[3,5], Jennifer L. Houtz[6], Ignacio T. Moore[7], Christopher N. Balakrishnan[1†], Brent M. Horton[6*†]

*Corresponding Author: Peri Bolton peri.bolton@gmail.com

† CNB and BMH contributed equally to this work.

[1]Department of Biology, East Carolina University, Greenville, NC, USA

[2]Department of Vertebrate Zoology, National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

[3]Migratory Bird Center, Smithsonian National Zoological Park, Washington DC, USA

[4]Bird Conservancy of the Rockies, Fort Collins, CO, USA

[5]Department of Biology, Carleton University, Ottawa, ON, Canada

[6]Department of Biology, Millersville University, Millersville, PA, USA

[7]Department of Biological Sciences, Virginia Tech, Blacksburg, VA, USA

Abstract

The neurogenomic mechanisms mediating cooperation in animals are understudied. Here we leverage extensive transcriptomic and behavioral data on birds that perform cooperative courtship displays to better understand the mechanisms underlying cooperative behaviors. Males of the neotropical wire-tailed manakin (*Pipra filicauda*) form cooperative display coalitions and their cooperative behaviors are modulated by testosterone in a status-specific manner. That is, testosterone promotes cooperation in non-territorial birds, but suppresses it in territory holders. We sought to understand the neurogenomic underpinnings of these three related traits: social status, cooperative behavior, and testosterone phenotype. To do this, we profiled gene expression in 10 brain nuclei spanning the social decision-making network (SDMN), and two key endocrine tissues known to regulate social behavior. We associated gene expression with each bird's

detailed behavioral and endocrine profile derived from 3 years of repeated measures taken from

free-living birds in the Ecuadorian Amazon. We found distinct landscapes of constitutive gene

expression were associated with social status, testosterone phenotype, and cooperation, reflecting

the modular organization and engagement of neuroendocrine tissues. Sex-steroid and

neuropeptide signaling appear to be important in mediating status-specific relationships between

testosterone and cooperation, which suggests shared regulatory mechanisms with male

aggressive and sexual behaviors. We also identify differentially regulated genes that are involved

in cellular activity and synaptic potentiation, suggesting that multiple mechanisms underpin these

genomic states. Finally, we identify SDMN-wide gene expression differences between territorial

and floater males that could form the basis of "status-specific" neurophysiological phenotypes,

potentially mediated by testosterone and growth hormone. Overall, our findings provide new,

systems-level insights into the mechanisms of behavior, and suggest that differences in

neurogenomic state are the basis for individual differences in social behavior.

Significance Statement

Cooperative behavior is an important determinant of fitness in many animals, yet its

neurogenomic underpinnings remain poorly understood. We leveraged an unprecedented dataset

describing gene expression in multiple brain nuclei and endocrine tissues from male wire-tailed

manakins (*Pipra filicauda*) characterized for social status, behavior, and testosterone phenotype

in the Ecuadorian Amazon. In this species, testosterone influences an unusual male-male

cooperative display, but the nature of the relationship between cooperation and circulating

testosterone depends on social status. We found evidence that individual behavioral phenotypes

are determined by a complex neurogenomic landscape across interconnected brain nuclei. We

show, on a transcriptomic scale, that brain modularity and hormone pleiotropy drive status-

specific differences in gene expression resulting individual differences in cooperative behavior.

Introduction

Cooperation is the synchronized behavior(s) of two or more individuals to produce a net fitness benefit for one or more individuals (Taborsky and Taborsky 2015). An individual's decisions about cooperation are based on the immediate social stimulus and individual condition, which is based on recent and distant past experiences, developmental effects, and underlying genetics. These effects can ultimately produce repeatable, among-individual variation in behaviors – a behavioral phenotype – where some individuals tend to be more cooperative than others. To understand the mechanisms of cooperation, one needs to understand the neurophysiological signal processing of the stimulus through to the behavioral response, as well as the genetic and neurophysiological components that underpin behavioral phenotypes. Indeed, understanding these mechanisms enables us to link how genetic and neurophysiological architecture can constrain and facilitate the evolution of cooperation (Soares *et al.* 2010; Hofmann *et al.* 2014; Taborsky and Taborsky 2015). Differences in cooperative behavioral phenotype are in part achieved by the pleiotropic yet modular effects of hormone signaling systems (Ketterson *et al.* 2009; Cox 2020). Hormonal pleiotropy mediates phenotypic integration, whereby multiple traits covary with a hormonal signal (Ketterson *et al.* 2009; Cox 2020), sometimes in opposite directions. For example, elevated testosterone typically promotes aggression but inhibits parental care in males (Wingfield *et al.* 2001). In contrast, through variation in hormone receptor expression, tissues across the body are often modular in their response to hormone signals, enabling diverse and flexible responses to hormonal and social signals and trait independence (Ketterson *et al.* 2009; Lipshutz *et al.* 2019).

Cooperation, like other social behaviors, appears to be mediated by differences in circulating hormones, such as sex steroids, and aspects of hormone-signaling pathways, including the distribution and abundance of hormone receptors in neural tissues (Soares *et al.* 2010; Kasper *et al.* 2017). The neural substrates modulating social behaviors are conserved across vertebrates (O'Connell and Hofmann 2011). Known as the social decision making network (SDMN), these reciprocally interconnected brain nuclei are rich in steroid receptors, steroidogenic enzymes, and steroid-modulated neuropeptide systems (Newman 1999; Goodson 2005; O'Connell and Hofmann 2011). The SDMN is both modular and dynamic, whereby different combinations of brain regions are activated in response to different social contexts (Newman 1999; Goodson 2005). This model means the common practice of measuring the

neurological response in a single nucleus is not reflective of all systems engaged in modulating a

given social behavior. Variation among individuals in the expression of neuroendocrine genes

across the SDMN likely underlies repeatable individual differences in social behavior and sets

conditions for behavioral phenotypes that respond differently to circulating hormones and social

stimuli. For example, the up or down-regulation of genes involved in sex-steroid signaling

pathways across nuclei of the SDMN can alter the sensitivity of the male brain to testosterone

and manifest as differential behavioral responses to a given hormone signal. Emerging research

is showing that behavioral phenotypes are correlated with differences in constitutive gene

expression or neurogenomic state among individuals characterized by suites of genes co-

expressed across behaviorally relevant brain regions (Kabelik *et al.* 2021; Lattin *et al.* 2021;

Antunes *et al.* 2021). However, given the diversity of behavioral components associated with

cooperation (Díaz-Muñoz *et al.* 2014; Kasper *et al.* 2017), questions remain on how

neuroendocrine mechanisms operate to produce the variety of social and cooperative behaviors

among individuals. Indeed, our understanding of the neuroendocrine mechanisms underlying

seemingly paradoxical male-male reproductive cooperative behaviors is only emerging (DuVal

and Goymann 2011; Díaz-Muñoz *et al.* 2014; Vernasco *et al.* 2020; Ryder *et al.* 2020; Jones and

DuVal 2021; Loveland *et al.* 2021). For the first time, we characterize gene expression across the

neuroendocrine system in relation status-specific androgenic modulation of a male-male

cooperative behavior.

The wire-tailed manakin (*Pipra filicauda*) is a neotropical lek breeding bird species in which

unrelated males perform cooperative displays to attract females (Figure 1A). Long-term display

partnerships established between territorial and subordinate non-territorial (floater) males form

the basis of complex social networks (Figure 1A, B) (Ryder *et al.* 2008; Ryder *et al.* 2011a). In

this bird, territoriality is a pre-requisite to siring offspring, and floater males follow an age-

graded queue for territorial status (Ryder *et al.* 2008; Ryder *et al.* 2009). Cooperation is directly

linked to fitness, whereby territory holders with more display partners sire more offspring, and

floaters with more partners are more likely to ascend to territorial status (Ryder *et al.* 2009).

Individuals have a repeatable 'testosterone phenotype' in which territorial males tend to have

higher testosterone than floaters (Figure 1D) (Ryder *et al.* 2011b; Ryder *et al.* 2020).

Remarkably, circulating testosterone has a status-specific effect on the cooperative display

behavior: experimental and observational evidence show that higher testosterone levels are

antagonistic to cooperation in territorial males but promote cooperation in floater males (Ryder

et al. 2020; Vernasco *et al.* 2020) (Figure 1C).

To explore the neurogenomic basis of the status-specific androgenic modulation of

cooperative behavior in these birds, we quantified transcriptome-level gene expression in the

SDMN and pituitary and testes of free-living male wire-tailed manakins sampled in the

Ecuadorian Amazon. Captive studies can obscure ecologically and evolutionary relevant

variation in environmental and behavioral responses (Kasper *et al.* 2017; Griffith *et al.* 2017), so

we demonstrate the exciting applicability of neuro-transcriptional methods in the challenging

conditions of the rainforest. In this study, we aimed to characterize the genes and neuroendocrine

pathways that modulate individual variation in cooperative behavior, as well as those related

social status and individual variation in testosterone phenotype as a modifier of both social status

and cooperative behavior. We characterized the constitutive gene expression for each of these

traits within and between nodes of the SDMN and the endocrine tissues. Given the androgenic

regulation of cooperative display (Ryder *et al.* 2020; Vernasco *et al.* 2020), we hypothesized that

differences in gene regulation would manifest in SDNM nuclei with high density of sex-steroid

receptors and pituitary and gonadal tissues (Newman 1999; Goodson 2005). We further reasoned

that androgenic regulation would relate to differential expression of genes relating to sex-steroid

signaling and metabolism, as well as three key neuropeptide systems involved in social behavior

(Goodson and Thompson 2010; Kingsbury and Wilson 2016). Given the modular nature of the

SDNM, we predicted that many gene expression patterns would be brain region-specific (Figure

1G). However, given the pleiotropic effects of testosterone (Ketterson *et al.* 2009; Cox 2020), we

also predicted that there would be signatures of testosterone's systemic influence on gene

expression patterns across the brain. We discuss our results in the context of modes of

phenotypic evolution and similarities with mechanisms of other social behaviors.

Materials and Methods

All methods described here were approved by Smithsonian Animal Care and Use Committee

(protocols 12-12,14-25,17-11), as well as the Ecuadorian Ministry of the Environment (MAE-

DNB-CM-2015-0008), and samples were transported with export permit (006-016-EXP-1C-

FAU-DNB/MA) and import permit (USDA APHIS 126133). Detailed methods can be found in

the supplementary material, and full code from analyses in R v 4.0.2 are available on the GitHub

site (https://periperipatus.github.io/PIFI_brain_transcriptome/) and on FigShare
(10.25573/data.22186516).

Field Methods

For RNA sequencing we sampled 16 wild territorial and floater wire-tailed manakin males from
Tiputini Biodiversity Station in the lowland Amazon rainforest of Ecuador (Table S1.1). Males
were categorized by social status as either territorial or floater. Territorial males (n=9) are males
in definitive plumage that hold a consistent display territory over the study period, while floaters
(n=7) have either definitive or pre-definitive plumage and do not occupy consistent territories
within leks (Ryder *et al.* 2008). On 12 of these 16 males, we used repeated testosterone sampling
(Figure 1D, Supplement 1) to characterize each male's testosterone phenotype (mean
testosterone). In this population, territorial males have, on average, higher circulating
testosterone than floaters (Ryder et al., 2020; Ryder, Horton, et al., 2011), and the males sampled
for this study reflect that pattern (Figure 1D). A male's testosterone phenotype is represented by
mean of the residuals of a linear regression from repeated sampling of testosterone concentration
(Supplement 1 for more detail, see also (Ryder *et al.* 2020)). Mean testosterone as a measure of
testosterone phenotype is repeatable within individuals (Ryder *et al.* 2020). In the broader
population, the relationship between testosterone phenotype and social network measures used as
a proxy for cooperation was status-specific, with testosterone suppressing cooperation in
territorial males but increasing cooperation in floaters (Ryder *et al.* 2020) (Figure 1C).

Social network structure and individual variation in behavior were characterized using
autonomous radio-telemetry and tag proximity detections within territories on a total population
of 180 male manakins over multiple years (Ryder *et al.* 2012; Dakin and Ryder 2018; Ryder *et*
al. 2020). We used this tagging data to calculate social network strength for the same 12 males
with repeated testosterone measures. Social network strength is a proxy for cooperative tendency
and is the time spent interacting with other males averaged over the study period (Figure 1E),
and was the most repeatable within individuals (Ryder *et al.* 2020). This social network proxy
for cooperation was most strongly status specific relation to an individual's testosterone
phenotype (Ryder *et al.* 2020) (Figure 1C), but the overall frequency of cooperative interactions
does not differ between territorial and floater males (Figure 1E).

     Microdissection and RNAseq

Methods for brain extraction, tissue preservation, cryosectioning, microdissection, and RNA

extraction followed those published previously (Horton *et al.* 2020b). Briefly, birds were

euthanized by decapitation near the site of capture immediately after blood sampling for

testosterone. Whole brains were extracted within 4-6 minutes of euthanasia and immediately

frozen in powdered dry ice. To capture the full Hypothalamic-Pituitary-Gonadal axis critical to

sex-steroid regulation, we additionally sampled the pituitary gland and testes, which were

sampled immediately after the brain. Brains and testes were kept frozen on dry ice in the field (<

4 hours) until transferred into a 21-day liquid nitrogen-charged Arctic Express™ dry shipper

(Thermo Scientific) where they were stored at cryogenic temperatures until imported into the

United States. Pituitaries were preserved in RNAlater™ (Invitrogen) at ambient temperatures

until imported into the United States.

The brains were cryosectioned, and frozen sections were microdissected to excise tissue from 10

different nuclei involved in social behavior – nine of which are from the SDMN. The SDMN

reflects the interconnectivity of two networks involved in regulating social behavior (O'Connell

and Hofmann 2011), constituting nuclei from both the Social Behavior Network (SBN)

(Newman 1999; Goodson 2005) and the Mesolimbic Reward System (MRS) (Olds and Milner

1954; O'Connell and Hofmann 2011). Figure 1F indicates the nomenclature used and the

relationships of these nuclei to the SBN and MRS. To clarify bird specific nomenclature: the

arcopallium intermedium (AI) is a region that may be, in part, homologous the basolateral

amygdala of mammals, a node of the MRS (O'Connell and Hofmann 2011), which is

hypothesized to play a role in androgen-dependent manakin display behavior (Fusani *et al.*

2014); and the nucleus taenia (TnA) is homologous to the mammalian medial amygdala

(O'Connell and Hofmann 2011). Finally, we sampled the paraventricular nucleus (PVN) for a set

of 10 nuclei sampled. This is a region that is not yet formally recognized as part of the SDMN,

despite its neuropeptide projections to nodes of the SDMN and established role as a major

regulator of vertebrate social behavior (Goodson and Thompson 2010; Goodson and Kingsbury

2013). For a graphical representation of the tissues sampled see Figure 1F, with a full list of

individuals collected in Table S1.1 and tissues in Table S1.2.

RNA was extracted and sequenced according to previously described methods (Horton *et*

al. 2020b). RNA libraries for the various tissues and field seasons were randomized amongst

sequencing batches (Supplement 1, Table S1.1 & S1.2). All samples analyzed are associated

with Bioproject PRJNA437157, and SRR12660169-198, SRR19521260-271, SRR19521432-

575. The file that matches SRA numbers with our code and metadata available on the GitHub.

Gene expression analyses

Reads were mapped to the annotated *Pipra filicauda* genome (GCA_003945595.1) using

splice-aware mapper STAR v.2.7.5 (Dobin *et al.* 2013), and transcripts were counted against

exons using featureCounts v.2.0.1 (Liao *et al.* 2014). For subsequent analyses we used a total of

170 libraries, and quality filtering of individuals and gene-counts was performed before the

analysis of each tissue (Supplement 1 & 3). To further explore tissue and brain nucleus function

in this species (Horton *et al.* 2020b), we characterized differences in gene expression among

tissues and brain nuclei using principal components analysis in PCAtool*s* and Weighted Gene

Co-Expression Analysis (WGCNA) (Langfelder and Horvath 2008; Blighe and Lun 2020)

(Supplement 2). The number of individuals used in each tissue varies owing to quality filtering

and variable sampling of individuals across all brain regions (Table S1.1 & S1.2).

To describe the role of tissues and brain nuclei in relation to these hormonal and

behavioral phenotypes we conducted filtering (supplement 1), differential expression

(Supplement 3) and co-expression (Supplement 5) analysis in each tissue and nucleus separately.

For a subset of tissues, we had an extra four individuals characterized for social status from a

previous study (Horton *et al.* 2020b), for which behavioral and hormone sampling data were

unavailable. Owing to this, and to not overparameterize the models that account for batch effects

we identified differentially expressed genes (DEGs) associated with each of our traits of interest

(social status, testosterone phenotype and social network strength) separately. These were run as

separate negative binomial generalized linear models for each trait that accounted for batch

effects where necessary using DESeq2 (Love *et al.* 2014) (For detailed description see

Supplement 3) in R v 4.0.2 (R Core Team 2020). Mean testosterone and social network strength

were input to DESeq2 as continuous variables. In these cases, significant DEGs are interpreted to

be correlated with the predictor variable, and Log_2 Fold Change (LFC) value represents the LFC

of gene expression per unit of the predictor variable. In all cases, significant DEGs were

identified using the Wald Test and the default FDR corrected p-value of 0.1 (q<0.1). Further, for

each tissue and nucleus we identified gene co-expression networks using WGCNA. The identity

of genes of interest identified by WGCNA were similar to those in DESeq2. For brevity and

simplicity we focus mostly on the differential expression results, but the full results are presented

in Supplement 5.

Due to the key role testosterone plays in modulating behavior in this species (Ryder *et al.*

2020), we conducted further detailed analysis on nine candidate genes involved in sex-hormone

metabolism and signaling (Figure S4.1), and a further nine steroid sensitive neuropeptides and

their receptors (Figure S4.2). Any candidate gene identified by differential gene expression

(DGE) analysis with an uncorrected $p < 0.05$ was considered "significant" and examined further.

In addition to analyzing models of status, mean testosterone and social network strength

separately, we also explored whether candidate genes were involved in mediating status specific

relationships between mean testosterone and social behavior (Ryder *et al.* 2020) by extracting

"significant" candidate genes from a final DGE interaction analysis (model form: expression ~

status x mean testosterone). Owing to the overfitting concerns discussed above, we primarily

considered this model as a method for identifying which of our candidate genes showed status-

specific expression with respect to mean testosterone, rather than identifying new DEGs. The list

of putative relationships between genes and traits were plotted (Supplement 4), and any

relationship with highly influential observations, or with $R^2 \leq 0.2$, was excluded from our final

set. For candidates derived from the interaction effect models, and for genes that were associated

with multiple traits, we decided the most convincing relationship based on a stepdown AIC

procedure (Supplement 4). Given the substantially smaller set of candidate genes we considered

uncorrected p-values to be significant (p<0.05), but report the FDR corrected value as well.

We characterized overlap in gene expression patterns among brain nuclei by comparison

of ranked gene-lists (Supplement 6). The gene-lists were ranked by $-\log_{10}$ transforming the p-

values and multiplying by the direction of expression. These gene lists were then compared for

similarity between each pair of tissues by calculating a) Pearson correlation and b) rank-rank

hypergeometric overlap (RRHO) test (Cahill *et al.* 2018).

Testosterone is pleiotropic in that it can affect multiple traits by influencing gene

expression in in similar ways in many different target tissues (Ketterson *et al.* 2009; Cox 2020),

and it is crucial to social status-specific regulation of behavior in the wire-tailed manakin (Ryder

et al. 2020). We reasoned that these effects would be reflected in shared signatures of differential

expression across brain nuclei. To explore systemic gene expression patterns (i.e., consistent

patterns across the entire brain), we used the transformed p-values from the aforementioned

analysis and calculated the median p-value across all brain nuclei. Genes that had a median p-

value < 0.05 (transformed value > |1.3|) were considered consistently differentially expressed

across the brain. Further, the eigenvalues for each principal component (PC) from the whole-

brain PCA analysis (Figure 1G inset) were Pearson correlated with our variables of interest using

pcatools::eigencorplot() (Blighe and Lun 2020) (Supplement 1). Correlations observed with

social status were further validated by a linear mixed model in package 'nlme' (Pinheiro *et al.*

2016). The model form included individual as a random effect and incorporated batch variables

that were also correlated with the same PC (Supplement 1).

276   We used *clusterProfiler* v3.16.1(Yu *et al.* 2012). to determine patterns of functionality in

277   DEGs and WGCNA modules with Gene Ontology enrichment analyses. If the total number of

278   FDR corrected DEGs was less than 40, we used gene set enrichment analysis instead which

279   leverages the rank order of p-values and direction of expression (Supplement 3).

Results

Landscape of gene expression among neural and endocrine tissues

283   Our analyses revealed substantial differences in gene expression among tissues and brain

regions (Figure 1G; Figure S2.7). As expected, the pituitary was associated with high expression

levels of genes key to hormonal regulation (e.g., PRL, GH1, TSHB, POMC; Figure S2.3), while

the testes showed high expression levels of genes associated with germ cells and flagellar

motility (e.g., PKDREJ, BOLL, DRC7; Figure S2.3). Within the brain, the majority of gene

expression variation was explained by differences between brain regions (Figure 1G inset) and

was reflected in brain region-specific gene co-expression networks (Figure S2.7). These

expression patterns are more similar between adjacent nuclei, forming major brain-region

groupings of the amygdala, hypothalamus, and midbrain (Figure 1G). Our 18 candidate

neuroendocrine genes (Table S1.3) also showed tissue and brain region-specific expression

(Table 2.2, Figures S4.1-4.2).

Social status and testosterone phenotype are associated with tissue and brain region-specific
patterns of gene expression

We observed a landscape of gene expression in association with testosterone phenotype
(Figure 2A) and social status (Figure 2B), that is characterized by a high number of DEGs (or co-
expression modules) identified in some tissues, but not others. Similarly, across brain regions we
identified relationships between many of our 18 candidate genes and status, testosterone
phenotype or their interaction (Figure 2C). Mean testosterone was a continuous predictor in
DESeq2, as such DEGs show up-or downregulation with a higher testosterone phenotype (Figure
2A), while DEGs in social status indicate up or down regulation in territorial males (Figure 2B).
The number of trait-associated co-expression modules were similar to our results from
differential gene expression, so we primarily focus on the tissues with large numbers of DEGs
and significant candidate genes for the main manuscript, but the full suite of results can be
viewed in supplements 3 and 5. The number of DEGs in each tissue was more similar between
testosterone phenotype and status than any other comparison of traits examined (R = 0.5, p =
0.08, Figure S3.7), and this pattern appears to hold across co-expression modules (Supplement 5)
and candidate gene analysis (Figure 2C & Figure 3B). This similarity was expected because
testosterone phenotype and status themselves are correlated (Figure 1D) (Ryder *et al.* 2011b;
Ryder *et al.* 2020).

The testes and the nucleus taenia (TnA) each had over 1,000 genes differentially
expressed (at uncorrected p-value < 0.05) in association with both testosterone phenotype and
status (144 and 252 genes after FDR correction q<0.01 in association with mean T; 2 and 13
after FDR correction in association with social status). DEGs identified in association with
testosterone phenotype in the testes were enriched for genes involved with response to
corticosterone (GO:0051412) and synaptic transmission and organization (e.g., GO:0007268,
GO:0050808) among others (Figure S3.20). Notable DEGs included steroidogenic acute
regulatory protein (STAR, LFC = -1.1, p = 2×10^{-7}, q = 0.0003, R^2 = 0.6) and candidate gene
AVPR1A (LFC = 0.8, p = 5×10^{-5}, q = 0.01, R^2 = 0.7; Figure S4.4). Note that Log_2 Fold Change
(LFC) represents change in gene expression per unit of the predictor variable. However, these
transcriptional differences do not correspond to significant differences in testis size in relation to
testosterone phenotype (Table S3.2), nor social status (Table S3.1 & S3.2).

Among the 252 DEGs associated with testosterone phenotype in the TnA, there was non-
significant enrichment for genes involved in neural crest migration (GO:0001755), learning and
memory (GO:0007613, GO:0007612), as well as long-term synaptic potentiation (e.g.,
GO:0060291) (Figure S3.28). Among the DEGs associated with testosterone phenotype in the
TnA, candidate gene estrogen receptor 2 (ESR2; aka Erβ) was upregulated in the TnA of males
with higher testosterone (Figure 2C; LFC = 0.8, p = 0.001, q = 0.08, R^2 = 0.4). Other notable
DEGs in the TnA included upregulation of immediate early gene JUN, also a 'coral1' hub gene
(module membership (MM) = 0.93; LFC = 0.61, p = 0.0004, q = 0.05) and corticotropin
releasing hormone receptor 2 (CRHR2, LFC = 1, q = 0.006, R^2 = 0.5).

Not all nuclei showed similar patterns between status and testosterone phenotype. For
example, the intercollicular nucleus (ICo) of the midbrain showed 161 DEGs associated with
testosterone phenotype after FDR correction, but did not stand out in association with social
status (3 DEGs). Most of the significantly enriched GO terms were biological processes relating
to motor proteins and cilia (e.g., GO:0003341, Figure S3.26). Intriguingly, although the POM did
not reveal a high number of DEGs in association with testosterone phenotype (45), our analysis
revealed differential expression for compelling gene ontology categories and candidate
neuroendocrine genes (Figure S3.25, Figure S4.10, Figure S5.7). Notably, males with higher
testosterone exhibited upregulation of candidate gene androgen receptor (AR) in POM (Figure
2D). Indeed, the 'floralwhite' module comprised many of the upregulated DEGs and was
significantly associated with both testosterone phenotype and to a lesser extent social status
(Figure S5.6). This module showed enrichment for biological processes relating to G protein-
coupled receptor signaling pathways (GO:0007186) and neuropeptide signaling pathway
(GO:0007218), with statistically non-significant enrichment for steroid receptor signaling
(GO:0030518), learning and memory (GO:0007611), and nervous system development
(GO:0007399) (Figure S3.7). Indeed, many of the module hub genes were also DEGs in
association with testosterone phenotype, and interact with candidate sex hormone pathways,
prolactin releasing hormone (PRLH, MM = 0.91, LFC = 2.4, q = 5x10^{-5}, R^2 = 0.6, Figure
S3.25A) and growth regulating estrogen receptor binding 1 (GREB1, MM = 0.86, LFC = 0.6, q =
3x10^{-5}, R^2 = 0.7, Figure S3.25A), gonadotropin releasing hormone 1 (GNRH1, MM = 0.76, LFC
= 0.4, p = 0.006, q = 0.2, R^2 = 0.4). Other notable 'floralwhite' hub genes include the candidate
gene progesterone receptor (PGR, MM = 0.81) which was also associated with social status

(Figure 2C; LFC = 0.4 p = 0.006, q = 0.7) and serotonin receptor 5A (HTR5A, MM = 0.87, LFC

= 0.4, p = 0.04, q = 0.6). Candidate gene aromatase (LOC113993669/CYP19A1) was also

upregulated in males with higher testosterone phenotype (Figure 2C, LFC = 0.6, p = 0.02, q =

0.4, R^2 = 0.5), but was found in the 'lightcoral' co-expression module.

Despite overall similarity in gene expression associations with testosterone phenotype

and status, some brain regions were more strongly associated with status or status × testosterone

phenotype interaction (Figure 2B). Likewise, expression in many of the 18 candidate

neuroendocrine genes were associated with status or the interaction with testosterone in different

nuclei (Figure 2C). The paraventricular nucleus (PVN) of the hypothalamus showed the most

DEGs after FDR correction (42), which showed enrichment for cytoskeletal movement genes

(Figure S3.12B) involved in intracellular transport and cilia (e.g., GO:0003341, GO:0061640),

and non-significant enrichment for cholesterol biosynthesis (GO:0006695, GO:0045540).

Candidate gene expression also revealed status-specific associations with testosterone phenotype

in the PVN, that is a statistical interaction between status and testosterone phenotype (Figure

2C). In particular, AR expression was positively correlated with testosterone phenotype in floater

males, but not in territorial males (Figure 2F; LFC = -0.7, p = 0.008, q = 0.9). The PVN of

territorial males also showed higher VIPR1 (Figure S4.8; LFC = 0.3, p = 0.02, q = 0.4) and lower

ESR2 expression (LFC = -0.6, p = 0.006, q = 0.2). In the bed nucleus of the stria terminalis

(BSTm), a region that bridges the SDMN and MRS, there were more than 1000 putative DEGs,

but only nine after FDR correction. Among these, AR expression was higher in territorial males

than in floater males (Figure 2E; LFC = 0.6, p = 0.0007, q =0.2). Further, testosterone

metabolism gene 5-alpha reductase (SRD5A2) was upregulated in the lateral septum (LS) of

territorial males (Figure S4.20; LFC = 0.7, p = 0.0005, q = 0.4). The VIP system is also

implicated in status-specific responses to testosterone in the TnA, where VIPR1 was upregulated

in association with testosterone in territorial, but not floater males (Figure S4.17; LFC = 1, p =

0.04 q = 0.7).

GCt was a hotbed for gene expression associated with social status and status ×

testosterone phenotype interaction (Figure 1B, Figure S3.6). In association with social status,

gene set enrichment analysis revealed there was a significant tendency for downregulation of

genes relating to synapse assembly (GO:0051965), and response estradiol (GO:0032355), and

upregulation of genes relating to RNA processing (e.g., GO:0000184, GO:0006396) and

translation (e.g., GO:0006413, GO:0006412) (Figure S3.15). While, candidate neuroendocrine

genes showed status specific expression in association with testosterone phenotype in this region

including progesterone receptor (PGR, Figure S4.15; LFC = 1.9, p= 0.003, q = 0.1), estrogen

receptor 1 (ESR1; aka Erα; LFC = 1.1 p = 0.04, q = 0.4), prolactin receptor (PRLR; LFC = 0.7, p

= 0.01, q = 0.2), and vasoactive intestinal peptide (VIP; Figure S4.15; LFC = 1.6, q = 0.005, q =

0.1). Further, candidate gene aromatase was downregulated with higher testosterone phenotype

in the GCt (LFC = -1.3, p = 0.002, q = 0.3, $R^2 = 0.4$).

Cooperative behavior is associated with a distinct gene expression landscape

Network strength (the number of cooperative interactions between individuals) varies

widely among territorial and floater males (Figure 1E), representing a key axis of variation in

social behavior in this species. We predicted that the associations between gene expression and

cooperative behavior would be different from those observed for social status and testosterone

phenotype. Indeed, we found a distinctive landscape of constitutive gene expression across

tissues relating to cooperative behavior (Figure 3A, Figure S3.5), as well as among candidate

genes (Figure 3B). These results are based on results from a continuous predictor and can be

interpreted as genes with significant up-or downregulation with increasing social network

strength. The Anterior Hypothalamus (AH) had the largest number of DEGs associated with a

male's social network strength. In this nucleus there were no significantly enriched GO terms,

but there was non-significant enrichment for genes involved in RNA metabolism (e.g.,

GO:0043928) and regulation of translation (e.g., GO:0017148), development (e.g., GO:0061061,

GO:0042472) and synaptic potentiation (e.g., GO:0007274) (Figure S3.35). The pituitary also

stood out in the abundance of differentially expressed genes associated with male's social

network strength (109 genes after FDR correction q < 0.1). There were no significantly enriched

GO categories after FDR correction (Figure S3.33B), but among the non-significant terms were

actin filament organization (GO:0007015), neural differentiation (e.g., GO:0031103,

GO:0045664) and the cell cycle (e.g., GO:0043065, GO:0071158.) Among the DEGs in the

pituitary, candidate genes prolactin (PRL) (Figure 3C; LFC = -0.6, p=5x10^{-5} q = 0.03, $R^2 = 0.7$),

ESR2 (Figure S4.5; LFC = -0.7, p = 0.04, q = 0.5, $R^2 = 0.5$) and VIPR2 (Figure S4.5; LFC = -

0.4, p = 0.008, q = 0.3, $R^2 = 0.4$) were downregulated in highly cooperative males. Intriguingly,

the PVN showed a weak correlation of one of our candidate genes: oxytocin (OXT) expression

decreased with a male's social network strength (Figure 3D; LFC = -0.3, p = 0.04, q = 0.6, R^2 =

0.2). Further, although there was less representation of sex-steroid related pathways in the

candidate gene analysis (Figure 2B), PGR was downregulated in males with higher social

network strength in the BSTm (Figure 2E; LFC = -0.3, p = 5×10^{-4}, q = 0.2, R^2 = 0.5).

420   Gene expression in the LS was tightly associated with an individual's social network

strength (98 genes after FDR correction q < 0.1). There were no GO categories significantly

enriched after FDR correction (Figure S3.42), but among the top non-significant terms were

those broadly involved in development (e.g., GO:0048538, GO:0048593), and neural

organization (e.g., GO:0099175, GO:0050771). However, the expression of our candidate genes

in the LS was not linked to social network strength. Although relatively few genes were

differentially expressed in the medial bed nucleus of the stria terminalis (BSTm) with respect to

social network strength, the LS and BSTm shared many genes with similar patterns of expression

in association with strength as shown by RRHO analysis (Cahill *et al.* 2018) (Figure 3G, F). In

contrast, the LS and BSTm shared few genes with similar expression in association with

testosterone phenotype (Figure 4H, I) or social status (Figure S6.1A), suggesting that these

regions are uniquely engaged in modulating cooperative behavior. Indeed, pairwise correlations

of gene expression in association with testosterone phenotype broadly recapitulates similarity of

tissues sampled (Figure 3H) and perhaps reflects similarities in pathways related to sex-steroid

signaling (Figure 2C).

Testosterone phenotype and social status are associated with systemic gene expression

436   We reasoned that the pleiotropic effects of testosterone and its role in social status in the

manakins (Ryder *et al.* 2020) would also produce systemic effects on gene expression, or similar

patterns of expression across multiple regions. Social status and testosterone phenotype were

linked to the highest number of consistently differentially expressed genes (60 and 66 genes with

median uncorrected p < 0.05 respectively), while only 10 genes were consistently differentially

expressed in association with social network strength (Supplement 6). Further, PC5 of the whole

brain analysis was significantly associated with social status in a linear mixed model that

corrected for batch confounds and repeated sampling of individuals (Figure S2.5). There was

overlap between the genes and GO categories between the consistently differentially regulated

genes with social status and testosterone phenotype and those identified in PC5 (Supplement 6),

but we focus on these common themes in relation to social status in the main manuscript for

447 simplicity. The top two most consistently differentially regulated genes are the upregulation of

448 an immune system gene C-type Lectin Domain gene (CLEC2B) (Figure 4A) and down

449 regulation of development gene roundabout guidance receptor 4 (ROBO4) (Figure 4B) in

450 territorial males. These signatures are reflected in the patterns of GO biological process

451 enrichment, though none of these terms are significant after FDR correction (Figure 4C). There

452 was also evidence for global differential regulation of telomere maintenance genes (Figure 4C),

453 with downregulation of POLD-1-like (LOC114000574) and upregulation of RAD51 in territorial

454 males. Further, there is evidence for global regulation of cholesterol biosynthesis (Figure 4C),

455 and glutathione transferases (Figure S2.6, Table S6.2), both of which are relevant in steroid

456 hormone biosynthesis. These findings of global expression differences among status classes

457 could be partly caused by the upregulation of growth hormone (GH1) in the pituitary of

458 territorial males (Figure 4D; LFC = 0.8, $p = 3 \times 10^{-5}$, q = 0.01). This result holds in the model that

459 accounts for the interaction between social status and mean T (LFC = 0.9, $p = 6 \times 10^{-5}$, q = 0.05).

460 **Discussion**

461 Using the most comprehensive transcriptome-level sampling of the SDMN to-date (Lopes and

462 König 2020; Bentz *et al.* 2021b; Kabelik *et al.* 2021; Bentz *et al.* 2021a; Antunes *et al.* 2021),

463 our findings highlight both modular and integrative neurogenomic organization underlying

464 individual variation in cooperative behavior. The relative roles of each are central to the

465 development and expression of any complex trait (Ketterson *et al.* 2009; Lipshutz *et al.* 2019;

466 Cox 2020). We discuss our findings in relation to each theme in turn.

467 **Different landscapes of gene expression with each trait reflect modular organization of the**

468 **neuroendocrine system**

469  Modularity is reflected in how different brain nuclei, endocrine tissues, and genetic

470 pathways within individuals are associated with the expression of different traits. Such

471 neurogenomic landscapes, characterized by the upregulation of neuroendocrine genes in some

472 brain regions and downregulation in others, would allow differential sensitivity to testosterone

473 and other signaling molecules among SDMN nuclei. These neurogenomic states can be likened

474 to the 'landscape' model of immediate early gene (IEG) activation across the SDMN after a

475 social stimulus (Newman 1999; Goodson 2005). In that landscape model, it is the pattern of IEG

476 expression across brain nuclei that correlates with behavioral responses, not the up or

477 downregulation of particular genes within a single nucleus. This original model considered this

478   landscape of IEG expression (and later neuropeptides) in the context of proximate responses to

479   social stimuli. Our findings extend this model in two ways: by considering transcriptomic

480   variation and how long-term neurogenomic states underlie individual variation in social

481   behavior.

482   In the landscape of constitutive gene expression in male wire-tailed manakins, total

number of DEGs and candidate gene expression implicate constitutive differences in most brain

regions, but the brain regions with higher numbers of DEGs or candidate gene expression may be

more important for mediating certain traits. Some brain regions showed associations between

gene expression and both social status and testosterone phenotype (e.g., POM) or only

testosterone phenotype (e.g., TnA, testes). This may be because these centers are involved in

testosterone mediated traits irrespective of social status (O'Connell and Hofmann 2012).

Meanwhile, others appeared to be more related to social status and status specific relationships

with testosterone phenotype (e.g., PVN, GCt), or an individual's strength of cooperative

behavior (e.g., PIT, LS). Our results also suggest that the identity and function of these genes

vary across tissues and brain regions (Figure 3GH). However, most analyses revealed DEGs

relating to neural development, synaptic potentiation or organization and genes relating to

cellular metabolism, or transcription/translation regulation. Thus, individual variation modulated

by neurogenomic states that encode synaptic modifications and connectivity (mechanism of

memory formation), larger scale shifts in cellular activity, and neural structure (Clayton *et al.*

2019; George *et al.* 2020)). These mechanisms may be supported or feedback on the variation in

elements of sex-steroid signalling we investigated in detail (*sensu "*neuroendocrine action

potential" (Clayton *et al.* 2019)).

500   We found ample evidence for modularity in steroid related gene expression, particularly

in association with social status, testosterone phenotype or their interaction. Unsurprisingly, we

found evidence that status-specific regulation of behavior could be modulated directly by

testosterone through AR in multiple nuclei (e.g., POM, PVN, BSTm). Higher AR expression is

linked to display performance in golden-collared manakins (Schlinger *et al.* 2013). Further, the

conversion of testosterone to estradiol by the enzyme aromatase and subsequent binding to

estrogen receptors is a key pathway in the regulation of testosterone-dependent male behaviors

more broadly (Schlinger and Balthazart 2013), and found evidence for their role in multiple

nuclei. Our results also highlight the importance of neural VIP expression in modulating multiple
aspects of social behavior (Kingsbury and Wilson 2016; Horton *et al.* 2020a), as it was the only
neuropeptide system with widespread patterns of differential expression with respect to both
social status, testosterone phenotype and social network strength.

Notably, VIP and other steroid receptors may play a role in status-specific responses to
testosterone in GCt. Although tyrosine hydroxylase (TH) neurons in the GCt have been
associated with courtship behaviors (Goodson *et al.* 2009; Ben-Tov *et al.* 2021), our finding of
association with VIP and courtship behaviors broadly speaking in GCt is novel (Kingsbury and
Wilson 2016). GCt coordinates social stimuli and downstream motor processes that control
behavior. For example, in the songbird brain, GCt links POM to the song control system and
thereby mediates androgen-dependent singing behavior (Haakenson *et al.* 2020). Given that
cooperative display in wire-tailed manakins requires coordinated physical (and vocal displays in
cooperative *Chiroxiphia* manakins (Trainer *et al.* 2002)), the status-specific differential
expression of genes for steroid receptors and steroid-sensitive neuropeptides in GCt suggest that
this SDMN region may play a unique role in mediating status-specific relationships between
testosterone and cooperative behavior.

The pattern of co-upregulation of steroid related genes in the POM of males with high
testosterone phenotype (and territorial males) point to this region's role as a potent steroid
regulator. The POM is known to be involved in regulating male aggression, sexual behavior and
parental care (O'Connell and Hofmann 2011), and stimulates gonadal testosterone production
through gonadotropin releasing hormone (*GnRH*). Indeed, the observed increase in GNRH1
transcription may relate to transcriptional changes indicating enhanced steroid production in the
testes. Co-upregulation of AR and GNRH1 would seem contradictory due to the negative
regulatory effects of androgens and AR on *GnRH* (Brayman *et al.* 2012), though these may not
interact at the cell population level or may show different temporal patterns of translation.
Regardless, activation and steroid receptor concentration in POM is associated with social status
differences and social status ascension cichlid fish (Maruska *et al.* 2013), and mice (Lee *et al.*
2022). In the same cichlid species, different preoptic cell types, including homologs for PVN,
showed higher activation in one of the two partners in cooperative territorial defense
(Weitekamp and Hofmann 2017). Indeed, testosterone can link the POM to the Hypothalamic

Pituitary Adrenal (HPA) axis via the PVN (Williamson *et al.* 2010), which in our study also

showed evidence of status specific gene expression, including AR. This previous work, and the

close linkages between the PVN and POM (Williamson *et al.* 2010), suggest these nuclei in

wire-tailed manakins could be involved in multiple elements of behavior depending on cell

types, social contexts and timescales investigated.

543   The number of differentially expressed genes observed in TnA implicates this region as a

hub for neurogenomic pathways involved in regulation of testosterone mediated traits. The avian

homolog of the medial amygdala (O'Connell and Hofmann 2011), the TnA integrates sensory

information and has been associated with the regulation of diverse social behaviors (Raam and

Hong 2021). Indeed, our results may reflect persistent activation of this area, based on the

upregulation of immediate early genes such as JUN and SYT7 in males with high testosterone

phenotype, and differential regulation of other genes involved in long-term synaptic potentiation

(Clayton 2000; Marrone *et al.* 2008; O'Connell *et al.* 2012; Clayton *et al.* 2019). Further, our

results implicate potential estrogenic regulation in this region via increased ESR2 in males with

higher testosterone phenotype. Indeed, estrogenic regulation through ESR1 has been shown to

regulate status-specific aggression in birds (Horton *et al.* 2014; Merritt *et al.* 2020) and prosocial

behavior in prairie voles (Cushing *et al.* 2008). Although ESR2 is not well characterized in birds,

it may function in the medial amygdala of mammals to regulate social recognition (Lymer *et al.*

2018), and sexual regulation (Nakata *et al.* 2016), and is associated with higher social rank in the

forebrains of cichlid fish (Burmeister *et al.* 2007). Further, the differential expression of stress

steroid receptor (CRHR2) hints again at the previously unexplored role of the HPA axis in

modulating display behavior in this species. Indeed, social ranks often differ in their

glucocorticoid profiles (Goymann and Wingfield 2004) or regulatory responses to

glucocorticoids (Snyder-Mackler *et al.* 2019). The related cooperative displaying lance-tailed

manakins (*Chiroxiphia lanceolata*) vary in testosterone as well as corticosterone profiles

according to social status (DuVal and Goymann 2011; Jones and DuVal 2021). Finally, the fish

homolog of TnA appears mediate aggression and cooperation in cooperative territory defense in

an actor-specific manner (Weitekamp and Hofmann 2017; Weitekamp *et al.* 2017), supporting a

role for modulating cooperation in wire-tailed manakins as well.

The frequency an individual male spends displaying with other males (i.e., network
strength), our proxy for cooperative behavior, is associated with a unique pattern of gene
expression among tissues and brain regions. Some of these signatures are difficult to parse, for
example many genes are differentially expressed in the AH, yet lack clear GO enrichment. Prior
to our study AH has associated with regulation of aggression rather than cooperation in males,
and acting via AVP receptors and VIP (O'Connell and Hofmann 2011; Horton *et al.* 2020a).
Although we do not find any evidence for the role of these genes in the AH, our findings may
suggest that cooperative displays are derived from displays of territorial aggression, and/or gene
expression is inhibiting aggression in these displays. Indeed, cooperative displays may act as a
way of maintaining dominance hierarchies in wire-tailed manakins.

The variety of pathways differentially expressed in the pituitary suggests there are whole-
organismal effects of social bond strength mediated by the pituitary. Indeed, previous work on
this species showed that more social males had shorter telomeres (Vernasco *et al.* 2021), and
social bond strength has been associated with signatures of energy metabolism and stress
(Anderson *et al.* 2022; Simons *et al.* 2022). Interestingly, PRL transcript was downregulated in
more cooperative males. Prolactin is best known for its role in promoting parental care
behaviors, including in cooperative breeders (Schoech *et al.* 1996), but emerging studies suggest
it could play a role in inhibiting aggression in cooperative breeders (Medger *et al.* 2019; Gilbert
et al. 2022) and promotion of nest defense behavior (Mohamed *et al.* 2016).

The LS showed many differentially expressed genes associated with cooperation, and
highly cooperative individuals may show differences in neural structure based on the enriched
GO terms. Patterns of expression in the BSTm were similar, but weaker, as it showed very few
statistically significant DEGs. BSTm has neural projections into the LS, and vasopressin (AVP)
and VIP neurons in the BSTm and receptors for these neuropeptides in the LS are associated
with affiliative behavior and gregariousness in birds (Kelly *et al.* 2011; Kelly and Goodson 2013;
Kingsbury and Wilson 2016). Those prior studies suggest a role for these brain nuclei in
processing social information acutely, whereas our study suggests differences among individuals
in cooperative tendency is related to constitutive differences in gene expression and neural
structure, perhaps relating to excitability of these regions in response to social information. Thus,
cooperative display is regulated by some of the same conserved brain regions as other types of

affiliative behavior. That we only observed a large number of DEGs in the LS and not in other
nuclei may suggest that other unsampled nuclei in the mesolimbic reward system are important
for modulation of cooperative tendency (e.g., (Paula *et al.* 2015; Weitekamp *et al.* 2017; de
Abreu *et al.* 2020; Lopes and König 2020).

We did not find convincing evidence for any variation in sensitivity to nonapeptides
(vasopressin and oxytocin) in the brain (AVPR2 in hypothalamic nuclei showed transcript counts
<30). However, the density of these receptors in particular brain regions may be just as important
as their abundance in modulating behavior (Goodson and Thompson 2010). We did find
evidence for differential sensitivity to AVP in the pituitary, again implicating stress response
pathways in mediating social status and cooperation in this species. Further, we did find a weak
negative relationship between oxytocin expression and social network strength in the PVN.
Although in apparent contrast to the widely held belief that oxytocin promotes sociality (as in
humans (Feldman 2012)), this among our other observations may indicate that cooperative
displays are a derivative of aggressive displays. However, nonapeptides have complex and
sometimes species-specific relationships with affiliative behavior (Goodson and Thompson
2010; Maruska *et al.* 2022), and can also be associated with social status (Lee *et al.* 2022).
Further, the ratio of peptide to mRNA can be an important factor indicating higher turnover
(usage) in certain contexts (Maruska *et al.* 2022). Therefore, we do not have sufficient evidence
that oxytocin inhibits cooperation in this species, and further experimental work is needed.

Our results establish a variety of mechanisms that could underpin individual variation in
cooperative behavior, and its status specific modulation by testosterone. We find evidence for
multiple forms of neuroplasticity, through steroid receptor variation, synaptic potentiation and
neural development, as well as cellular activity (Clayton *et al.* 2019). Indeed, testosterone
manipulations are known to induce changes in neuroanatomy in behaviorally relevant brain
nuclei (Kabelik *et al.* 2008), suggesting a direct role for testosterone-mediated neural growth and
flexibility in modulating male-male cooperation (Soares *et al.* 2010). Our results provide
promising targets for further study on the mechanisms underlying cooperative behavior,
including examination of neuroanatomical features and the use of hormonal or other
pharmaceutical manipulations and RNAi technologies to examine causal relationships between
neural gene expression and behavior (e.g., (Merritt *et al.* 2020)).

Similarities in gene expression reflect the role of testosterone as a phenotypic integrator

Similarities in gene expression among tissues are predicted as a result of the pleiotropic effects of testosterone and support the role of the hormone as a mediator of phenotypic integration. The extent of gene expression similarities and differences among tissues may reflect the degree of integration and independence (or modularity), respectively, that can either constrain or facilitate adaptive evolution (Ketterson *et al.* 2009; Lipshutz *et al.* 2019). We identified similarities in testosterone associated gene expression between brain regions that broadly recapitulated tissue ontology. We also identified consistent, brain-wide (systemic) gene expression differences associated predominantly with social status and testosterone phenotype. However, systemic results did not explain as much variation in overall neural gene expression as with some axes of personality (Kabelik *et al.* 2021; Lattin *et al.* 2021), suggesting that the brain region-specific patterns may be more important in modulating male wire-tailed manakin behavior. Moreover, the broader associations of social status with metabolic, immune, and telomere gene ontology categories suggests differences in allostatic load among status classes in male-male cooperative systems (Goymann and Wingfield 2004; Vernasco *et al.* 2021; Jones and DuVal 2021). Further, differential global regulation of telomeres may explain the relative stability of telomere length in territorial wire-tailed manakin males (Vernasco *et al.* 2021), and collectively these results may represent metabolic adaptations for territoriality.

Our results highlight two additional hormones that may have a system-wide role in the status-specific modulation of cooperative display behavior. We found differences in the expression of genes for important behavioral and metabolic hormones (prolactin and growth hormone) in the pituitary. The anterior pituitary is the primary production site for both hormones, and thus gene expression patterns here are likely good predictors for circulating levels of those hormones and thus signal strength. Transcripts encoding prolactin, best known for its role in promoting parental care behaviors, were negatively correlated with cooperative behavior. While prolactin is emerging as important in other areas of social behavior, such as nest defense (Mohamed *et al.* 2016), the negative relationship observed appears contradictory to a general model of increased prolactin promoting broadly affiliative social interactions. We speculate that this relationship reflects that cooperative display behavior may have evolved from initially aggressive or competitive interactions (Díaz-Muñoz *et al.* 2014). The other compelling

phenotypic integrator for status-dependent behaviors in the wire-tailed manakin comes from the

gene encoding growth hormone (GH1). Ascension in social rank in the cooperatively displaying

manakins is associated with age (McDonald 1989; DuVal 2007; Ryder *et al.* 2008). It was

therefore surprising that territorial males in our study showed increased expression of GH1 in the

pituitary as compared to floater males. However, in anolis lizards, stimulation of the growth

hormone axis is associated with increased testosterone that enables phenotypic divergence

between males and females (Cox *et al.* 2017). Without gross morphological differences in wire-

tailed manakin status-classes, growth-hormone could have subtle effects on multiple systems

enabling the status-specific regulation of behaviors. In particular, status differences in GH1

expression may explain repeated signatures of neural development and protein synthesis in our

results (Sonksen 2006), and may reflect the increased metabolic demands on displaying

territorial males through muscle performance and regulation of oxidative stress (Hu *et al.* 2019).

The neuroethological role of growth hormone are not well known, but its release may be linked

to stress (Greenwood and Landon 1966; McCormick *et al.* 1998). Growth hormone

administration promotes increased aggression in mice (Matte 1981), and fish (Jönsson *et al.*

1998; Johansson *et al.* 2004). Indeed, growth hormone inhibitor somatostatin and its receptors

have also been implicated in regulating aggression and social status in cichlid fish independent of

androgens (Hofmann and Fernald 2000; Trainor and Hofmann 2006) whilst natal androgen

treatment increased aggression and somatostatin receptor expression in zebra finches (Bentz *et*

al. 2021b). We did some evidence for negative relationship between SSTR1 and SSTR2 with

testosterone phenotype in the testes, and positive association between SSTR1 and testosterone

phenotype in the POM but these were not significant after FDR correction. Thus, our results

suggest that the growth hormone axis may play a role in the status-specific regulation of wire-

tailed manakin cooperative displays, and increased GH in territorial males may be an important

inhibitor of cooperation by promoting aggression. Future work on social behavior should

consider the role of growth hormone and somatostatin in conjunction with sex steroids and

glucocorticoids to gain novel insights into its function in behavioral regulation.

Conclusion

Across the SDMN and HPG of the male wire-tailed manakin we observed correlations with

social status, cooperation and testosterone phenotype in patterns of gene regulation and a

plurality of gene functions, including sex-steroid and neuropeptide signaling. This landscape of gene expression speaks to the modular nature of behavioral mechanisms (Newman 1999; Goodson 2005; Lipshutz *et al.* 2019) and highlights the continuum between complete phenotypic integration and modularity (Ketterson *et al.* 2009). Much of the previous work on the mechanisms of social and cooperative behaviors focus on relatively few candidate genes in broad-scale brain divisions (e.g., hypothalamus) or in one or a few particular nuclei. Our work presented here is one of the most comprehensive examinations to date of region-specific gene expression in all nuclei of the SDMN, albeit correlative only. Our correlative findings help us to better understand cooperative behavior (Soares *et al.* 2010) by linking neurogenomics with endocrine mechanisms, and generate further hypotheses to test in further experiments to implicate causality. We further demonstrate and emerging idea that constitutive landscapes in gene expression across the brain are associated with repeatable differences in behavioral phenotype (Kabelik *et al.* 2021; Lattin *et al.* 2021; Antunes *et al.* 2021). We also provide tantalizing glimpses into nuclei and genomic pathways involved in the status-specific regulation of this cooperative display behavior. Although questions remain on the acute mediators of cooperative display behavior in the wire-tailed manakin, we demonstrate the power and flexibility of a system-wide approach. We propose that landscapes of constitutive gene expression for multiple genes and across many tissues and brain regions, like those observed here, form the foundation of consistent inter-individual differences in any behavior, upon which flexible and acute responses to social stimuli are built.

Acknowledgements

We want to acknowledge long hours of fieldwork in Ecuador by Ben Vernasco and Camilo Alfonso-Cuta that made the behavioral foundation of this work. Thank you to them, and the staff at Tiputini Biodiversity Station. Thank you to Eric Shuppe and to Matthew Fuxjager for use of laboratory resources. Thank you to David Clayton, Julie George, Sarah London, and Barney Schlinger for helpful discussions and manuscript reviewing, and Vanessa González for advice on data analysis. **Funding** for the reference genome came from a private donor at Millersville University. Funding for the behavioral and RNAseq study came from the National Science Foundation (IOS 1353085; DBI 1457541), the Smithsonian Migratory Bird Center, the Global Change Center at Virginia Tech, Millersville University, and the Tiputini Biodiversity Station of the Universidad San Francisco de Quito.

719   **Data and materials availability:** Raw read data are available associated with BioProject

720   PRJNA437157, and all code used to conduct analyses is available here:

721   https://github.com/periperipatus/PIFI_brain_transcriptome

References

de Abreu, M. S., Maximino, C., Cardoso, S. C., Marques, C. I., Pimentel, A. F. N., Mece, E., Winberg, S., Barcellos, L. J. G., and Soares, M. C. (2020). Dopamine and serotonin mediate the impact of stress on cleaner fish cooperative behavior. *Hormones and Behavior* **125**. doi:10.1016/J.YHBEH.2020.104813

Anderson, J. A., Lea, A. J., Voyles, T. N., Akinyi, M. Y., Nyakundi, R., Ochola, L., Omondi, M., Nyundo, F., Zhang, Y., Campos, F. A., Alberts, S. C., Archie, E. A., and Tung, J. (2022). Distinct gene regulatory signatures of dominance rank and social bond strength in wild baboons. *Philosophical Transactions of the Royal Society B* **377**. doi:10.1098/RSTB.2020.0441

Antunes, D. F., Teles, M. C., Zuelling, M., Friesen, C. N., Oliveira, R. F., Aubin-Horth, N., and Taborsky, B. (2021). Early social deprivation shapes neuronal programming of the social decision-making network in a cooperatively breeding fish. *Molecular Ecology* **30**, 4118–4132. doi:10.1111/MEC.16019

Ben-Tov, M., Duarte, F., and Mooney, R. (2021). A neural hub that coordinates learned and innate courtship behaviors. *bioRxiv*, 2021.09.09.459618. doi:10.1101/2021.09.09.459618

Bentz, A. B., George, E. M., Wolf, S. E., Rusch, D. B., and Podicheti, R. (2021a). Experimental competition induces immediate and lasting effects on the neurogenome in free-living female birds. *Proceedings of the National Academy of Sciences of the USA* **118**, e2016154118. doi:10.1073/pnas.2016154118/-/DCSupplemental.Published

Bentz, A. B., Niederhuth, C. E., Carruth, L. L., and Navara, K. J. (2021b). Prenatal testosterone triggers long-term behavioral changes in male zebra finches: unravelling the neurogenomic mechanisms. *BMC Genomics* **22**, 158. doi:https://doi.org/10.1186/s12864-021-07466-9

Blighe, K., and Lun, A. (2020). PCAtools: Everything Principal Components Analysis.

Brayman, M. J., Pepa, P. A., and Mellon, P. L. (2012). Androgen Receptor Repression of Gonadotropin-Releasing Hormone Gene Transcription via Enhancer 1. *Molecular and cellular endocrinology* **363**, 92. doi:10.1016/J.MCE.2012.07.012

Burmeister, S. S., Kailasanath, V., and Fernald, R. D. (2007). Social dominance regulates androen and estrogen receptor gene expression. *Bone* **51**, 164–170. doi:10.1038/jid.2014.371

Cahill, K., Huo, Z., Tseng, G. C., Logan, R. W., and Seney, M. L. (2018). Improved identification of concordant and discordant gene expression signatures using an updated rank-rank hypergeometric overlap approach. *Scientific Reports* **8**, 9588. doi:10.1038/s41598-018-27903-2

Clayton, D. F. (2000). The genomic action potential. *Neurobiology of Learning and Memory* **74**, 185–216. doi:10.1006/nlme.2000.3967

Clayton, D. F., Anreiter, I., Aristizabal, M., Frankland, P. W., Binder, E. B., and Citri, A. (2019). The role of the genome in experience-dependent plasticity: Extending the analogy of the

genomic action potential. *Proceedings of the National Academy of Sciences* **117**, 201820837. doi:10.1073/pnas.1820837116

Cox, R. M. (2020). Sex steroids as mediators of phenotypic integration, genetic correlations, and evolutionary transitions. *Molecular and Cellular Endocrinology* **502**, 110668. doi:10.1016/j.mce.2019.110668

Cox, R. M., Cox, C. L., McGlothlin, J. W., Card, D. C., Andrew, A. L., and Castoe, T. A. (2017). Hormonally Mediated Increases in Sex-Biased Gene Expression Accompany the Breakdown of Between-Sex Genetic Correlations in a Sexually Dimorphic Lizard. *https://doi.org/10.1086/690105* **189**, 315–332. doi:10.1086/690105

Cushing, B. S., Perry, A., Musatov, S., Ogawa, S., and Papademetriou, E. (2008). Estrogen Receptors in the Medial Amygdala Inhibit the Expression of Male Prosocial Behavior. *Journal of Neuroscience* **28**, 10399–10403. doi:10.1523/JNEUROSCI.1928-08.2008

Dakin, R., and Ryder, T. B. (2018). Dynamic network partnerships and social contagion drive cooperation. *Proceedings of the Royal Society B: Biological Sciences* **285**, 20181973. doi:10.1098/rspb.2018.1973

Díaz-Muñoz, S. L., DuVal, E. H., Krakauer, A. H., and Lacey, E. A. (2014). Cooperating to compete: Altruism, sexual selection and causes of male reproductive cooperation. *Animal Behaviour* **88**, 67–78. doi:10.1016/j.anbehav.2013.11.008

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21.

DuVal, E. H. (2007). Adaptive advantages of cooperative courtship for subordinate male lance-tailed manakins. *American Naturalist* **169**, 423–432. doi:10.1086/512137

DuVal, E. H., and Goymann, W. (2011). Hormonal correlates of social status and courtship display in the cooperatively lekking lance-tailed manakin. *Hormones and Behavior* **59**, 44–50. doi:10.1016/j.yhbeh.2010.10.004

Feldman, R. (2012). Oxytocin and social affiliation in humans. *Hormones and behavior* **61**, 380–391. doi:10.1016/J.YHBEH.2012.01.008

Fusani, L., Donaldson, Z., London, S. E., Fuxjager, M. J., and Schlinger, B. A. (2014). Expression of androgen receptor in the brain of a sub-oscine bird with an elaborate courtship display. *Neuroscience Letters* **578**, 61–65. doi:10.1016/j.neulet.2014.06.028

George, J. M., Bell, Z. W., Condliffe, D., Dohrer, K., Abaurrea, T., Spencer, K., Leitão, A., Gahr, M., Hurd, P. J., and Clayton, D. F. (2020). Acute social isolation alters neurogenomic state in songbird forebrain. *Proceedings of the National Academy of Sciences* **117**, 201820841. doi:10.1073/pnas.1820841116

Gilbert, J. D., Rossiter, S. J., Bennett, N. C., and Faulkes, C. G. (2022). The elusive role of prolactin in the sociality of the naked mole-rat. *Hormones and Behavior* **143**, 105196. doi:10.1016/J.YHBEH.2022.105196

Goodson, J. L. (2005). The vertebrate social behaviour network: evolutionary themes and

variations. *Hormones and Behavior* **48**, 11–22. doi:10.1038/mp.2011.182.doi

Goodson, J. L., Kabelik, D., Kelly, A. M., Rinaldi, J., and Klatt, J. D. (2009). Midbrain dopamine neurons reflect affiliation phenotypes in finches and are tightly coupled to courtship. *Proceedings of the National Academy of Sciences* **106**, 8737–8742. doi:10.1073/PNAS.0811821106

Goodson, J. L., and Kingsbury, M. A. (2013). What's in a name? Considerations of homologies and nomenclature for vertebrate social behavior networks. *Hormones and Behavior* **64**, 103–112. doi:10.1016/J.YHBEH.2013.05.006

Goodson, J. L., and Thompson, R. R. (2010). Nonapeptide mechanisms of social cognition, behavior and species-specific social systems. *Current Opinion in Neurobiology* **20**, 784–794. doi:10.1016/J.CONB.2010.08.020

Goymann, W., and Wingfield, J. C. (2004). Allostatic load, social status and stress hormones: the costs of social status matter. *Animal Behaviour* **67**, 591–602. doi:10.1016/J.ANBEHAV.2003.08.007

Greenwood, F. C., and Landon, J. (1966). Growth hormone secretion in response to stress in man. *Nature* **210**, 540–541.

Griffith, S. C., Crino, O. L., Andrew, S. C., Nomano, F. Y., Adkins-Regan, E., Alonso-Alvarez, C., Bailey, I. E., Bittner, S. S., Bolton, P. E., Boner, W., Boogert, N., Boucaud, I. C. A., Briga, M., Buchanan, K. L., Caspers, B. A., Cichoń, M., Clayton, D. F., Derégnaucourt, S., Forstmeier, W., Guillette, L. M., Hartley, I. R., Healy, S. D., Hill, D. L., Holveck, M.-J., Hurley, L. L., Ihle, M., Tobias Krause, E., Mainwaring, M. C., Marasco, V., Mariette, M. M., Martin-Wintle, M. S., McCowan, L. S. C., McMahon, M., Monaghan, P., Nager, R. G., Naguib, M., Nord, A., Potvin, D. A., Prior, N. H., Riebel, K., Romero-Haro, A. A., Royle, N. J., Rutkowska, J., Schuett, W., Swaddle, J. P., Tobler, M., Trompf, L., Varian-Ramos, C. W., Vignal, C., Villain, A. S., and Williams, T. D. (2017). Variation in Reproductive Success Across Captive Populations: Methodological Differences, Potential Biases and Opportunities. *Ethology* **123**, 1–29. doi:10.1111/eth.12576

Haakenson, C. M., Balthazart, J., and Ball, G. F. (2020). Cognition and Behavior Effects of Inactivation of the Periaqueductal Gray on Song Production in Testosterone-Treated Male Canaries (*Serinus canaria*). *eNeuro* **7**, ENEURO.0048-20.2020. doi:10.1523/ENEURO.0048-20.2020

Hofmann, H. a., Beery, A. K., Blumstein, D. T., Couzin, I. D., Earley, R. L., Hayes, L. D., Hurd, P. L., Lacey, E. a., Phelps, S. M., Solomon, N. G., Taborsky, M., Young, L. J., and Rubenstein, D. R. (2014). An evolutionary framework for studying mechanisms of social behavior. *Trends in Ecology and Evolution* **29**, 1–9. doi:10.1016/j.tree.2014.07.008

Hofmann, H. A., and Fernald, R. D. (2000). Social Status Controls Somatostatin Neuron Size and Growth. *Journal of Neuroscience* **20**, 4740–4744. doi:10.1523/JNEUROSCI.20-12-04740.2000

Horton, B. M., Hudson, W. H., Ortlund, E. A., Shirk, S., Thomas, J. W., Young, E. R., Zinzow-Kramer, W. M., and Maney, D. L. (2014). Estrogen receptor α polymorphism in a species with alternative behavioral phenotypes. *Proceedings of the National Academy of Sciences of*

the United States of America **111**, 1443–1448. doi:10.1073/pnas.1317165111

Horton, B. M., Michael, C., Mackenzie, P., and Maney, D. L. (2020a). Vasoactive intestinal peptide as a mediator of the effects of a supergene on social behavior. *Proceedings of the Royal Society B: Biological Sciences* **287**, 20200196.

Horton, B. M., Ryder, T. B., Moore, I. T., and Balakrishnan, C. N. (2020b). Gene expression in the social behavior network of the wire-tailed manakin (*Pipra filicauda*) brain. *Genes, Brain and Behavior*, e12560. doi:10.1111/gbb.12560

Hu, B., Hu, S., Yang, M., Liao, Z., Zhang, D., Luo, Q., Zhang, X., and Li, H. (2019). Growth hormone receptor gene is essential for chicken mitochondrial function in vivo and in vitro. *International Journal of Molecular Sciences* **20**, 1–13. doi:10.3390/ijms20071608

Johansson, V., Winberg, S., Jönsson, E., Hall, D., and Björnsson, B. (2004). Peripherally administered growth hormone increases brain dopaminergic activity and swimming in rainbow trout. *Hormones and Behavior* **46**, 436–443.

Jones, B., and DuVal, E. H. (2021). Glucocorticoids correlate with and predict social status in the cooperatively breeding lance-tailed manakin (*Chiroxiphia lanceolata*). In 'Integrative and Comparative Biology'. pp. E428-429

Jönsson, E., Johnsson, J., and Björnsson, B. (1998). Growth hormone increases aggressive behavior in juvenile rainbow trout. *Hormones and Behavior* **33**, 9–15.

Kabelik, D., Julien, A. R., Ramirez, D., and O'Connell, L. A. (2021). Social boldness correlates with brain gene expression in male green anoles. *Hormones and Behavior* **133**, 105007. doi:10.1016/j.yhbeh.2021.105007

Kabelik, D., Weiss, S. L., and Moore, M. C. (2008). Steroid hormones alter neuroanatomy and aggression independently in the tree lizard. *Physiology & behavior* **93**, 492. doi:10.1016/J.PHYSBEH.2007.10.008

Kasper, C., Vierbuchen, M., Ernst, U., Fischer, S., Radersma, R., Raulo, A., Cunha-Saraiva, F., Wu, M., Mobley, K. B., and Taborsky, B. (2017). Genetics and developmental biology of cooperation. *Molecular Ecology* **26**, 4364–4377. doi:10.1111/MEC.14208

Kelly, A. M., and Goodson, J. L. (2013). Behavioral relevance of species-specific vasotocin anatomy in gregarious finches. *Frontiers in Neuroscience* **7**, 242. doi:10.3389/FNINS.2013.00242

Kelly, A. M., Kingsbury, M. A., Hoffbuhr, K., Schrock, S. E., Waxman, B., Kabelik, D., Thompson, R. R., and Goodson, J. L. (2011). Vasotocin neurons and septal V1a-like receptors potently modulate songbird flocking and responses to novelty. *Hormones and Behavior* **60**, 12–21. doi:10.1016/J.YHBEH.2011.01.012

Ketterson, E. D., Atwell, J. W., and McGlothlin, J. W. (2009). Phenotypic integration and independence: Hormones, performance, and response to environmental change. *Integrative and Comparative Biology* **49**, 365–379. doi:10.1093/icb/icp057

Kingsbury, M. A., and Wilson, L. C. (2016). The Role of VIP in Social Behavior: Neural Hotspots for the Modulation of Affiliation, Aggression, and Parental Care. *Integrative and*

Comparative Biology **56**, 1238–1249. doi:10.1093/ICB/ICW122

Langfelder, P., and Horvath, S. (2008). WGCNA: An R package for weighted correlation network analysis. *BMC Bioinformatics* **9**, 559. doi:https://doi.org/10.1186/1471-2105-9-559

Lattin, C. R., Kelly, T. R., Kelly, M. W., and Johnson, K. M. (2021). Constitutive gene expression differs in three brain regions important for cognition in neophobic and non-neophobic house sparrows (*Passer domesticus*). *bioRxiv*, 2021.07.06.451290. doi:10.1101/2021.07.06.451290

Lee, W., Dwortz, M. F., Milewski, T. M., Champagne, F. A., and Curley, J. P. (2022). Social status mediated variation in hypothalamic transcriptional profiles of male mice. *Hormones and Behavior* **142**, 105176. doi:10.1016/J.YHBEH.2022.105176

Liao, Y., Smyth, G. K., and Shi, W. (2014). featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930.

Lipshutz, S. E., George, E. M., Bentz, A. B., and Rosvall, K. A. (2019). Evaluating testosterone as a phenotypic integrator: From tissues to individuals to species. *Molecular and Cellular Endocrinology* **496**, 110531. doi:10.1016/j.mce.2019.110531

Lopes, P. C., and König, B. (2020). Wild mice with different social network sizes vary in brain gene expression. *BMC Genomics* **21**, 1–14. doi:10.1186/s12864-020-06911-5

Love, M. I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**, 550.

Loveland, J. L., Giraldo-Deck, L. M., Lank, D. B., Goymann, W., Gahr, M., and Küpper, C. (2021). Functional differences in the hypothalamic-pituitary-gonadal axis are associated with alternative reproductive tactics based on an inversion polymorphism. *Hormones and Behavior* **127**, 104877. doi:10.1016/j.yhbeh.2020.104877

Lymer, J. M., Sheppard, P. A. S., Kuun, T., Blackman, A., Jani, N., Mahbub, S., and Choleris, E. (2018). Estrogens and their receptors in the medial amygdala rapidly facilitate social recognition in female mice. *Psychoneuroendocrinology* **89**, 30–38. doi:10.1016/J.PSYNEUEN.2017.12.021

Marrone, D. F., Schaner, M. J., Mcnaughton, B. L., Worley, P. F., and Barnes, C. A. (2008). Immediate-Early Gene expression at rest recapitulates recent experience. *The Journal of Neuroscience* **28**, 1030–1033. doi:10.1523/JNEUROSCI.4235-07.2008

Maruska, K. P., Anselmo, C. M., King, T., Mobley, R. B., Ray, E. J., and Wayne, R. (2022). Endocrine and neuroendocrine regulation of social status in cichlid fishes. *Hormones and Behavior* **139**, 105110. doi:10.1016/J.YHBEH.2022.105110

Maruska, K. P., Zhang, A., Neboori, A., and Fernald, R. D. (2013). Social Opportunity Causes Rapid Transcriptional Changes in the Social Behaviour Network of the Brain in an African Cichlid Fish. *Journal of Neuroendocrinology* **25**, 145–157. doi:10.1111/j.1365-2826.2012.02382.x

Matte, A. C. (1981). Growth hormone and isolation-induced aggression in wild male mice. *Pharmacology, biochemistry, and behavior* **14 Suppl 1**, 85–87. doi:10.1016/S0091-

3057(81)80014-7

McCormick, S. D., Shrimpton, J. M., Carey, J. B., O'Dea, M. F., Sloan, K. E., Moriyama, S., and Björnsson, B. T. (1998). Repeated acute stress reduces growth rate of Atlantic salmon parr and alters plasma levels of growth hormone, insulin-like growth factor I and cortisol. *Aquaculture* **168**, 221–235. doi:10.1016/S0044-8486(98)00351-2

McDonald, D. B. (1989). Cooperation Under Sexual Selection : Age-Graded Changes in a Lekking Bird. *The American Naturalist* **134**, 709–730.

Medger, K., Bennett, N. C., Ganswindt, S. B., Ganswindt, A., and Hart, D. W. (2019). Changes in prolactin, cortisol and testosterone concentrations during queen succession in a colony of naked mole-rats (Heterocephalus glaber): a case study. *Die Naturwissenschaften* **106**. doi:10.1007/S00114-019-1621-1

Merritt, J. R., Grogan, K. E., Zinzow-kramer, W. M., Sun, D., and Ortlund, E. A. (2020). A supergene-linked estrogen receptor drives alternative phenotypes in a polymorphic songbird. , 1–8. doi:10.1073/pnas.2011347117

Mohamed, R. A., Shukry, M., Mousa-balabel, T. M., and Elbassiouny, A. A. (2016). Assessment of Plasma Prolactin and Nest Defense Behaviour During Breeding Cycle of Pigeon (Columba livia domestica). *Journal of Environmental & Agricultural Sciences* **7**, 19–22.

Nakata, M., Sano, K., Musatov, S., Yamaguchi, N., Sakamoto, T., and Ogawa, S. (2016). Effects of Prepubertal or Adult Site-Specific Knockdown of Estrogen Receptor β in the Medial Preoptic Area and Medial Amygdala on Social Behaviors in Male Mice. *eNeuro* **3**, 579–588. doi:10.1523/ENEURO.0155-15.2016

Newman, S. W. (1999). The medial extended amygdala in male reproductive behavior: A node in the mammalian social behavior network. *Annals of the New York Academy of Sciences* **877**, 242–257. doi:10.1111/j.1749-6632.1999.tb09271.x

O'Connell, L. A., and Hofmann, H. A. (2012). Social Status Predicts How Sex Steroid Receptors Regulate Complex Behavior across Levels of Biological Organization. *Endocrinology* **153**, 1341–1351. doi:10.1210/EN.2011-1663

O'Connell, L. A., and Hofmann, H. A. (2011). The Vertebrate mesolimbic reward system and social behavior network: A comparative synthesis. *Journal of Comparative Neurology* **519**, 3599–3639. doi:10.1002/cne.22735

O'Connell, L. A., Matthews, B. J., and Hofmann, H. A. (2012). Isotocin regulates paternal care in a monogamous cichlid fish. *Hormones and Behavior* **61**, 725–733. doi:10.1016/J.YHBEH.2012.03.009

Olds, J., and Milner, P. (1954). Positive reinforcement produced by electrical stimulation of septal area and other regions of rat brain. *Journal of Comparative and Physiological Psychology* **47**, 419–427. doi:10.1037/H0058775

Paula, J. R., Messias, J. P., Grutter, A. S., Bshary, R., and Soares, M. C. (2015). The role of serotonin in the modulation of cooperative behavior. *Behavioral Ecology* **26**, 1005–1012. doi:10.1093/beheco/arv039

Pinheiro, J., Bates, D., DebRoy, S., and Sarkar, D. (2016). Linear and Nonlinear Mixed Effects Models. R. package v 3.1-128.

R Core Team (2020). R: A language and environment for statistical computing. Available at: https://www.r-project.org/

Raam, T., and Hong, W. (2021). Organization of neural circuits underlying social behavior: a consideration of the medial amygdala. *Current opinion in neurobiology* **68**, 124. doi:10.1016/J.CONB.2021.02.008

Ryder, T. B., Blake, J. G., Parker, P. G., and Loiselle, B. A. (2011a). The composition, stability, and kinship of reproductive coalitions in a lekking bird. *Behavioral Ecology* **22**, 282–290. doi:10.1093/beheco/arq213

Ryder, T. B., Dakin, R., Vernasco, B. J., Evans, B. S., Horton, B. M., and Moore, I. T. (2020). Testosterone modulates status-specific patterns of cooperation in a social network. *The American Naturalist* **195**, 82–94. doi:10.1086/706236

Ryder, T. B., Horton, B. M., and Moore, I. T. (2011b). Understanding testosterone variation in a tropical lek-breeding bird. *Biology Letters* **7**, 506–509. doi:10.1098/rsbl.2010.1219

Ryder, T. B., Horton, B. M., Van Den Tillaart, M., De Dios Morales, J., and Moore, I. T. (2012). Proximity data-loggers increase the quantity and quality of social network data. *Biology Letters* **8**, 917–920. doi:10.1098/rsbl.2012.0536

Ryder, T. B., McDonald, D. B., Blake, J. G., Parker, P. G., and Loiselle, B. A. (2008). Social networks in the lek-mating wire-tailed manakin (*Pipra filicauda*). *Proceedings of the Royal Society B: Biological Sciences* **275**, 1367–1374. doi:10.1098/rspb.2008.0205

Ryder, T. B., Parker, P. G., Blake, J. G., and Loiselle, B. A. (2009). It takes two to tango: Reproductive skew and social correlates of male mating success in a lek-breeding bird. *Proceedings of the Royal Society B: Biological Sciences* **276**, 2377–2384. doi:10.1098/rspb.2009.0208

Schlinger, B. A., Barske, J., Day, L., Fusani, L., and Fuxjager, M. J. (2013). Hormones and the neuromuscular control of courtship in the golden-collared manakin (*Manacus vitellinus*). *Frontiers in neuroendocrinology* **34**, 143. doi:10.1016/J.YFRNE.2013.04.001

Schlinger, B., and Balthazart, J. (2013). Aromatase and Behavior: Concepts Gained from Studies of Aromatase in the Avian Brain. In 'Brain Aromatase, Estrogens, and Behavior'. doi:10.1093/acprof:oso/9780199841196.003.0010

Schoech, S. J., Mumme L, R., and Wingfield.ast C, J. (1996). Prolactin and helping behaviour in the cooperatively breeding Florida scrub-jay, *Aphelocoma c. coerulescens*. *Animal Behaviour* **52**, 445–456. doi:10.1006/ANBE.1996.0189

Simons, N. D., Michopoulos, V., Wilson, M., Barreiro, L. B., and Tung, J. (2022). Agonism and grooming behaviour explain social status effects on physiology and gene regulation in rhesus macaques. *Philosophical Transactions of the Royal Society B: Biological Sciences* **377**. doi:10.1098/RSTB.2021.0132

Snyder-Mackler, N., Sanz, J., Kohn, J. N., Voyles, T., Pique-Regi, R., Wilson, M. E., Barreiro,

L. B., and Tung, J. (2019). Social status alters chromatin accessibility and the gene regulatory response to glucocorticoid stimulation in rhesus macaques. *Proceedings of the National Academy of Sciences of the United States of America* **116**, 1219–1228. doi:10.1073/PNAS.1811758115/SUPPL_FILE/PNAS.1811758115.SD01.XLSX

Soares, M. C., Bshary, R., Fusani, L., Goymann, W., Hau, M., Hirschenhauser, K., and Oliveira, R. F. (2010). Hormonal mechanisms of cooperative behaviour. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 2737–2750. doi:10.1098/rstb.2010.0151

Sonksen, P. (2006). Growth Hormone Replacement Therapy and Life Quality. Future Perspectives. In 'The Somatotrophic Axis in Brain Function'. (Ed F. Nyberg.) pp. 327–333. (Elsevier Inc.) doi:10.1016/B978-012088484-1/50028-8

Taborsky, M., and Taborsky, B. (2015). Evolution of genetic and physiological mechanisms of cooperative behaviour. *Current Opinion in Behavioral Sciences* **6**, 132–138. doi:10.1016/J.COBEHA.2015.11.001

Trainer, J. M., McDonald, D. B., and Learn, W. A. (2002). The development of coordinated singing in cooperatively displaying long-tailed manakins. *Behavioral Ecology* **13**, 65–69. doi:10.1093/beheco/13.1.65

Trainor, B. C., and Hofmann, H. A. (2006). Somatostatin Regulates Aggressive Behavior in an African Cichlid Fish. *Endocrinology* **147**, 5119–5125. doi:10.1210/EN.2006-0511

Vernasco, B. J., Dakin, R., Majer, A. D., Haussmann, M. F., Ryder, T. B., and Moore, I. T. (2021). Longitudinal dynamics and behavioural correlates of telomeres in male wire-tailed manakins. *Functional Ecology* **35**, 450–462. doi:10.1111/1365-2435.13715

Vernasco, B. J., Horton, B. M., Moore, I. T., and Ryder, T. B. (2020). Reduced cooperative behavior as a cost of high testosterone in a lekking passerine bird. *Behavioral Ecology* **31**, 401–410. doi:10.1093/beheco/arz201

Weitekamp, C. A., and Hofmann, H. A. (2017). Neuromolecular correlates of cooperation and conflict during territory defense in a cichlid fish. *Hormones and Behavior* **89**, 145–156. doi:10.1016/j.yhbeh.2017.01.001

Weitekamp, C. A., Nguyen, J., and Hofmann, H. A. (2017). Neuromolecular regulation of aggression differs by social role during joint territory defense. *Integrative and Comparative Biology* **57**, 631–639. doi:10.1093/icb/icx009

Williamson, M., Bingham, B., Gray, M., Innala, L., and Viau, V. (2010). The Medial Preoptic Nucleus Integrates the Central Influences of Testosterone on the Paraventricular Nucleus of the Hypothalamus and Its Extended Circuitries. *The Journal of Neuroscience* **30**, 11762. doi:10.1523/JNEUROSCI.2852-10.2010

Wingfield, J. C., Lynn, S. E., and Soma, K. K. (2001). Avoiding the 'Costs' of Testosterone: Ecological Bases of Hormone-Behavior Interactions. *Brain, Behavior and Evolution* **57**, 239–251. doi:10.1159/000047243

Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS: A Journal of Integrative Biology* **16**, 284–287. doi:10.1089/omi.2011.0118

Fig. 1. Circulating testosterone and neuroendocrine tissues are involved in the cooperative behavior of wire-tailed manakins.

A) Males engage in cooperative displays that form the basis of B) complex social networks. The example shown is a single lek over one week, where node size indicates a male's average social network strength (measure of overall time spent interacting), and edge width represents the frequency of interactions between a pair of individuals. Node color denotes floater (red) and territorial (gray) male status, where darker colors were included in the study. C) A male's testosterone phenotype (mean testosterone (corrected)) influences his cooperative tendency differently in territorial and floater classes. D) Twelve individuals (points) were from the larger population with repeated measures of testosterone (density), and E) proximity-data logging measures of social network strength. Note the total sample size for individuals characterized for social status is N=16 is larger than for testosterone or sociality (N=12). F) RNAseq characterized gene expression from each individual in 10 testosterone-sensitive brain regions and two endocrine tissues. Structures shown are not to scale. Colors represent functional groups (O'Connell and Hofmann 2011). G & inset) principal component analysis of transcriptomic data reveals distinct gene expression profiles among brain regions and endocrine tissues (colors and shapes as F). (photo by Alice Boyle).

Fig. 2. Neurogenomic landscape of social status and hormone phenotype.
A) There is a landscape of gene expression in male wire-tailed manakins with high testosterone phenotype (mean testosterone (corrected)) relative to lower testosterone phenotype. Red bars indicate number of genes that are upregulated in males with higher testosterone, whereas blue indicates genes that are downregulated in males with higher testosterone. Darker colors and text indicate number of differentially expressed genes after FDR correction. B) The landscape of gene expression in relation to social status, where red indicates genes upregulated in territorial males. C) Summary of candidate gene analysis and filtering, where colors indicate direction of expression (as previous panels), but intensity reflects which of the models were significant, and "i" indicates there was an interaction effect, and a black box indicates result was significant after FDR correction for all genes in that region's transcriptome. D) AR was upregulated in POM of males with higher testosterone. where point colors indicate social status (red = floater; black = territorial. Fitted line from linear regression. Subheader indicates key results from DESeq2: $\log_2$ Fold Change (LFC), uncorrected p-value (p), and correction for multiple testing (q) and R^2 correlation coefficient calculated separately. E) AR was upregulated in the BSTm of territorial males. F) AR shows status-specific expression in relation to testosterone phenotype in the PVN. Grey area represents the 95% confidence interval as determined by linear regression of gene ~ Status×mean_T. Note that maximum sample size for analysis of social status was N=16, while

only 12 individuals were sampled for testosterone and behavior, but the actual sample size varies slightly in each tissue owing to quality filtering.

Fig. 3. A distinct landscape of gene expression in relation to cooperative behavior.
A) Landscape of gene expression in male more cooperative wire-tailed manakins (social network strength). Darker colors and text indicate number of genes after FDR correction. Red indicates genes that are upregulated in males with higher social network strength, whereas blue indicates genes that are downregulated in these males. B) Summary of candidate gene analysis and filtering, where colors indicate direction of expression (as previous panels) in relation to individuals with higher social network strength. A black box indicates the result remained significant after FDR correction for all genes in that region's transcriptome. C) Prolactin (PRL) in the pituitary shows decreased expression in males with higher social network strengths. D) Oxytocin (OXT) transcript expression decreases in males with higher social network strength. E) Progesterone Receptor (PGR) has lower transcript expression in males with higher social network strength. F) RRHO2 analysis reveals similar profiles of gene expression associated with cooperation in the LS and BSTm, G) but otherwise generally low similarity of gene expression among tissues in association with cooperation. H) Similarity of gene expression among tissues in relation to testosterone phenotype roughly recapitulates ontogeny of brain regions, I) and there is no correlation between testosterone related gene expression profiles in the LS and BSTm. E,F) Cell color and number are the Pearson correlation coefficients of gene expression between tissues, correlating log10 transformed differential expression p-values (times the direction of

differential expression). Black boxes outline significant tissue correlation clusters based on 1,000 bootstrap resamples of the correlation matrix.

Fig. 4: Global neural gene expression differences between social classes
A) Immune gene CLEC2B-like was consistently upregulated in the brains of territorial individuals, and B) developmental gene ROBO4 was consistently downregulated in the brains of territorial males. C) Enriched GO categories among genes that were consistently up- or downregulated between territorial and floater males, where bar color indicates uncorrected p-value. D) These broader signatures of developmental differences could be explained by upregulation of growth hormone transcript in the pituitary of territorial males.