**Genetic exchange shapes ultra-small Patescibacteria metabolic capacities in**

**the terrestrial subsurface**

Authors: Emilie Gios[1], Olivia E. Mosley[1], Nobuto Takeuchi[1], Kim M. Handley[1,*]

[1]School of Biological Sciences, The University of Auckland, Auckland, New Zealand

*Corresponding author: K. M. Handley kim.handley@auckland.ac.nz

**ABSTRACT**

Bacterial genomes are highly dynamic entities, mostly due to the extent of horizontal

gene transfer (HGT) occurring in these organisms. HGT is thought to be the main

driver of genetic variation and adaptation to local environment in bacteria. However,

little is known about the modalities of HGT within natural microbial communities,

especially the implications of genetic exchange for streamlined microorganisms such

as Patescibacteria (Candidate Phyla Radiation). We searched for evidence of genetic

exchange in 125 Patescibacteria genomes recovered from aquifer environments and

detected the presence of hundreds of genomic islands, individually transferred genes

and prophage combined, with up to 29% of genome length attributed to HGT. Results

show that most individual gene transfer events occurred between Patescibacteria, but

donors were also phylogenetically diverse groundwater microorganisms. Using gene

donor-recipient information, we identified one potential host (Omnitrophota) of the

ultra-small bacteria, and confirmed this by matching relative abundance patterns

across 16 groundwater samples. A wide variety of metabolic functions were introduced

in Patescibacteria genomes by HGT including transcription, translation and DNA

replication, recombination and repair. This study illustrates the evolutionarily dynamic

nature of Patescibacteria genomes despite the constraints of streamlining, and that

HGT in these organisms is also mediated via viral infection.

**INTRODUCTION**

The horizontal transfer of genetic material (or horizontal gene transfer; HGT), involving acquisition of exogenous DNA, is thought to be a key factor in the evolution of prokaryotes, especially in bacteria (Ochman et al., 2000). HGT accounts for a significant fraction of gene diversity and genomic fluidity in these organisms, leading to a variety of genome structures and physiological diversity (Soucy et al., 2015). This evolutionary process is important for niche adaptation, and is often driven by mobile genetic elements (MGEs) such as genomic islands (GIs), transposons and phage. Many routes exist for gene transfer: uptake of environmental DNA molecules (transformation), exchange between two cellular organisms via a physical bond (conjugation), or delivery through phage infection (transduction).

Recent metagenomic studies of subsurface ecosystems reveal a large group of unusual bacteria referred to as the Patescibacteria phylum (equating to the Candidate Phyla Radiation; CPR), which has been suggested to comprise up to 26% of all currently known bacterial diversity (Parks et al., 2017; Schulz et al., 2017). The group constitutes taxa consistently harbouring small cell and genome sizes (Brown et al., 2015). Implications of reduced size include limited metabolic functions, such as the lack of known amino acid and nucleotide biosynthesis pathways (Castelle et al., 2018). These observations led to the prediction that most ultra-small Patescibacteria have a symbiotic lifestyle (Brown et al., 2015), and some were confirmed to be episymbionts based on cultivation and microscopy studies (He et al., 2021). Close physical associations with other organisms due to metabolic dependencies, along with the long residence time of groundwater (Griebler & Lueders, 2009), and the importance of the biofilms in aquifers (Flynn et al., 2008), could favour genetic exchange in subsurface ultra-small microbial communities.

Despite proximity to other groundwater bacteria, extensive gene loss in

Patescibacteria evolution could result in selective pressures against the acquisition

and retention of exogenous DNA that are stronger than in non-streamlined genomes.

Another potential barrier to HGT lies in the fact that genome reduction in obligate

intracellular symbionts has often been associated with loss of functions involved in

DNA recombination and repair (Moran & Bennett, 2014), which could limit integration

of foreign DNA. Previous studies have, however, shown not only that a significant

fraction of Patescibacteria organisms still possesses genes required for homologous

recombination (Castelle et al., 2018), but that they are naturally competent for uptake

of eDNA molecules, mediated by DNA intake pumps (via comEC and pili complexes;

Castelle et al., 2017, 2018; Nelson & Stegen, 2015).

Investigating the mobility of adaptive genes via HGT could provide important

information about the evolutionary strategy of these ultra-small organisms, including

aspects governing putative host-symbiont interactions. Implications of HGT include

altered interaction modalities with host populations, as described in *Vibrio fischeri*.

Adding a single gene to the *V. fischeri* genome is sufficient to alter its host range in

squids (Mandel et al., 2009). Additionally, acquisition of MGEs has been shown to

constitute evidence of symbiotic relationships between HGT recipient and donor, such

as extensive gene transfer from *Wolbachia* endosymbionts to their insect hosts during

genome reduction (Dunning Hotopp et al., 2007), and putative acquisition of host-

derived ankyrin repeat proteins by bacterial sponge symbionts to inhibit phagocytosis

(Nguyen et al., 2014). How potentially adaptive genes aid niche adaptation in ultra-

small prokaryotes and fine-tune metabolic interactions with their predicted 'basibiont'

prokaryotic host cells remains largely unknown.

To understand the role of HGT in the evolution of complex, natural groundwater

communities, and particularly streamlined Patescibacteria, we analyzed 396

metagenome-assembled genomes (MAGs), including 132 Patescibacteria genomes.

We investigated genomic evidence for, and the frequency of, individual horizontal

gene transfer events (defined here as HT genes) in the MAGs, and compared these

to acquisitions of larger MGEs such as GIs and prophage. Results provide evidence

for many HGT events among Patescibacteria, and exchanges with other prokaryotes

in groundwater. We report that metabolic functions horizontally acquired by

Patescibacteria taxa appear to be biased towards features distinct from the general

groundwater communities.

**MATERIALS AND METHODS**

**Sample collection**

Groundwater samples were collected from 8 wells from 2 sand/gravel aquifers in

Canterbury, New Zealand, as described by (Mosley et al., 2022). Briefly, wells were

purged (×3–5 bore volumes), then 3–90 L of groundwater was filtered per sample.

Biomass was captured on 0.22 µm mixed cellulose ester (MCE) filters, after passing

through a 1.2 µm MCE pre-filter, using a 142 mm filter holder (Merck Millipore Ltd.,

Cork, Ireland). Samples were preserved in RNAlater (ThermoFisher Scientific,

Waltham, MA, USA), transported on dry ice, and stored at -80°C.

After collecting a groundwater sample (as above), a second sample per well

was collected following low frequency sonication (2.43 kW for 2 minutes) to induce

biofilm (and sediment) detachment from the surrounding aquifer, using a custom

sonication device (Close et al., 2020). 0.5–15 L of sediment (and hence biomass)

enriched groundwater was filtered as above.

**Nucleic acid extraction**

To remove RNAlater, filters were centrifuged (2500 g for 5 min) and washed with nuclease-free phosphate-buffered saline (pH 7.4, ThermoFisher Scientific, Waltham, MA, USA). DNA was extracted using the DNeasy PowerSoil Pro Kit (Qiagen, Valencia, CA, USA) and 0.25-0.9 g of whole filter per reaction. Samples were extracted in replicate (1–47 reactions/sample). Replicates were pooled and concentrated using a sodium acetate/ethanol precipitation with glycogen (0.1 µg/µL final concentration, Roche, Basel, Switzerland). High molecular weight DNA was confirmed via gel electrophoresis. DNA was quantified using Qubit fluorometry (ThermoFisher Scientific), and quality was checked using a NanoPhotometer (Implen, Munich, Germany). Extractions yielded 70–864 ng of DNA (8.7 ng for gwj02) for metagenomics.

**Metagenome sequencing**

Whole genome shotgun sequencing was undertaken on the 16 Canterbury samples (gwj01-16). DNA libraries were prepared using the Illumina TruSeq Nano DNA Kit with ~550 bp inserts, by Otago Genomics (University of Otago, Dunedin, NZ). The low-yield gwj02 sample was prepared with the ThruPLEX DNA-Seq Kit (Takara Bio USA, Inc., Mountain View, CA, USA). Sequences (2×250 bp) were generated using the Illumina HiSeq 2500 V4 platform. Adapter sequences were removed using Cutadapt v2.10 (Martin, 2011), and reads were quality trimmed using sickle v01.33 (parameters -q 30 -l 80). Read quality was checked using FastQC v0.11.7 (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/).

**Metagenome assembly and binning**

The 16 metagenomes were individually assembled using SPAdes v3.11.1 (Bankevich

et al., 2012) (parameters: --meta, -k 43,55,77,99,121). Metagenomes from the same

well also were co-assembled (same parameters). A dereplicated set of metagenome-

assembled genomes (MAGs) (99% average nucleotide identity, ANI, dereplication

threshold) was generated as described previously (Tee et al., 2020), and refined

according to bin_detangling script (https://github.com/dwwaite/bin_detangling).

For ultra-small prokaryotes, genome completeness and contamination were

assessed based on the presence of 51 bacterial single-copy genes (SCGs) for

Dependentiae, 43 SCGs for Patescibacteria and 38 archaeal SCGs for DPANN

(Anantharaman et al., 2016). For other prokaryotes, completeness was estimated

using CheckM v1.0.13 (Parks et al., 2015). To determine coverage, reads were

mapped onto the de-replicated MAGs using Bowtie (Langmead et al., 2009). Sample-

specific genome relative abundance was calculated by normalizing based on the

highest read count between samples (Probst et al., 2018).

**Gene prediction and functional annotation**

Protein coding sequences were predicted using Prodigal v2.6.3 (Hyatt et al., 2010).

Genes were annotated using USEARCH (Edgar, 2010) against UniRef100 (Suzek et

al., 2015), Uniprot (The UniProt Consortium, 2017) and KEGG (Kanehisa et al., 2016)

databases (e-value ≤0.001 and identity ≥50%), and HMMs against PFAM

(Sonnhammer et al., 1998) and TIGRFAM (Haft et al., 2003) databases using HMMER

v3.1b2 (Potter et al., 2018) (e-value ≤0.001). Additionally, functional annotation was

carried out using eggNOG-mapper v2 (Huerta-Cepas et al., 2017) against the

eggNOG v5.0 database (Huerta-Cepas et al., 2019), using default parameters.

**Genome taxonomy**

Taxonomic classification of MAGs was performed using the Genome Taxonomy

Database Toolkit (GTDB-Tk, v0.2.1) (Chaumeil et al., 2020) with database release 89

(Parks et al., 2020). Maximum likelihood trees were constructed using 120 bacterial

and 122 archaeal concatenated marker genes with IQ-TREE v1.6.12 (L.-T. Nguyen et

al., 2015) using 100 bootstraps, and ModelFinder (Kalyaanamoorthy et al., 2017) best-

fit model LG+F+G4 for bacteria and LG+F+I+G4 for archaea. Trees were visualized

and annotated with iTOL (Letunic & Bork, 2007). MAGs (≥80% complete) were

compared to GTDB representative genomes (downloaded 20-Jan-2020) by

calculating Average Amino-acid Identities (AAI) for blastp matches sharing ≥30%

identity over ≥70% of alignment length.

**Identification of horizontal gene transfer**

Individual HT events among MAGs were identified using MetaCHIP v1.10.0 (Song et

al., 2019) at phylum, class, order, family and genus levels. False-positive HT genes

(potentially introduced via assembly contamination) were filtered out by removing HT

genes marked by MetaCHIP as "full-length match" (where the match represents a

large proportion of contig) or "end match" (HT genes located at the end of contig) as

described by Dong et al. (2021). Information about HGT events involving members of

the Patescibacteria phylum were retrieved and visualised using the Circlize package

in R (Gu et al., 2014).

For genomic island prediction, contigs were concatenated into contiguous

sequences and uploaded to IslandViewer4 web server (Bertelli et al., 2017) for

genomic island prediction. GIs that were predicted by at least one method, and that

did not overlap two concatenated contigs, were considered for the analysis. Genes

located on GIs were compared to community gene content using blastn.

Prophage integrated in the Patescibacteria genomes were identified using

VIBRANT 1.2.1 (Kieft et al., 2020). Whenever a predicted viral sequence was

overlapping a predicted GI, CheckV quality assessment information was used and only

"High-quality" genomes were retained as prophage (Nayfach et al., 2021).

**Phylogenetic distance**

Average Amino-acid Identities (AAI) between MAGs (≥80% complete) involved in HGT

events were determined via blastp matches sharing ≥30% identity over ≥50% of query

and database alignment length.

**Orthology analysis**

To investigate the Patescibacteria pangenome, we performed an orthologous

clustering of 118,635 protein-coding sequences encoded by the recovered MAGs.

Pairwise all-vs-all blastp searches were conducted using the following cut-offs: ≥30%

nucleotide identity over ≥50% of query sequence length, e-value ≤0.001. BLAST table

outputs were used as input for orthogroup clustering using Orthofinder v2.3.1 (Emms

& Kelly, 2019) with -b option and default parameters.

**RESULTS AND DISCUSSION**

**Prevalence of HGT events in Patescibacteria genomes**

Using the community-level HGT detection tool MetaCHIP (Song et al., 2019),

we investigated the potential horizontal transfer of individual genes between aquifer

microorganisms. A total of 1,487 transfers were identified among the 396 groundwater

MAGs (Table S1), including 1,407 unique HT genes. Taxonomic breakdown of

detected HT events revealed that over half of the Patescibacteria genomes

experienced HT with another genome in the dataset, regardless of their taxonomy or

estimated genome size (Fig. 1a,b). One hundred and twenty-three (9% of) horizontally

acquired genes were identified in 68 Patescibacteria organisms, representing 1.0 $\pm$

1.2 HT (average $\pm$ standard deviation) genes received per genome and 1.1 $\pm$ 1.3 HT

genes per Mbp (Fig. 1b). In comparison, genomes of other members of the community

contained on average 3.5 $\pm$ 4.8 horizontally transferred genes per genome, but a

comparable 1.4 $\pm$ 2.1 per Mbp. The comparison of HT events per Mbp indicates that

the rate of exchange of individual genes is mostly independent of genome size (i.e.

larger genomes do not receive proportionally more genes). This is supported by

significant positive correlations between HT events per genome and estimated

genome size (both donor and recipient HT events, Fig. 2), and is consistent with

previous observations of HT acquired genes in prokaryotic genomes (Garcia-Vallvé et

al., 2000), and of transfers occurring between distantly related organisms (i.e. cross-

order transfers, >85% of the total HT events detected here) (Cordero & Hogeweg,

2009).

**Figure 1. Prevalence of HGT in Patescibacteria and the wider groundwater community.** (a) Maximum likelihood phylogenetic trees of 132 Patescibacteria MAGs recovered in this study. Trees are based on 120 concatenated bacterial marker gene alignment from GTDB-Tk. Rings from the center: number of prophage; number of HT events; number of GIs; estimated genome size (Mbp). The scale bar indicates the number of substitutions per site. (b) Number of recipient HT events between all groundwater MAGs per phylum normalized by genome size. Patescibacteria phylum is indicated in teal. The center line of each boxplot represents the median; the top and bottom lines are the first and third quartiles, respectively; and the whiskers show 1.5× the interquartile range. (c) Proportion of each genome occupied by acquired GIs and prophage sequences, comparing Patescibacteria (teal) and the rest of the community (black). Significant differences were assessed for each group using Wilcoxon Signed Rank (*: p < 0.05).

When searching for large MGEs, we found 120 GIs in 63 Patescibacteria MAGs, ranging from 2.7 to 45 kbp long, with 40 GIs over 10 kb long (average length 17.4 ± 0.7 kb) (Table S2). Additionally, 24 putative prophage genomes (30% of those overall) were found to be concentrated in 5 Patescibacteria MAGs – with the proportion of prophage in Patescibacteria being comparable to the proportion of Patescibacteria MAGs in the community (33%). This frequency of prophage in ultra-small bacterial genomes is surprising, as prophage frequency has been shown to increase with genome size (Touchon et al., 2016). Accordingly, our analysis shows that, as also

observed for HT genes, the frequency of acquisition of longer complex DNA

sequences, constituting GIs, increased with the genome size of groundwater-derived

MAGs (Fig. 2b).

**Figure 2.** Correlations between the estimated genome size of groundwater-derived
MAGs and HGT. (a) Estimated genome size compared with recipient HT events (left)
and donor HT events (right). (b) Estimated genome size compared with the number of
GIs detected (left), and the proportion of genome length occupied by GI and prophage
sequences (right).

When considering the groundwater community overall, the proportion of

genome length occupied by GIs and prophages, combined, remained constant

regardless of recipient genome size (Fig. 2b), and despite GI and prophage sequences

being similar in length between ultra-small and other microorganisms (Fig. 3, Table

S2). This is consistent with observations, by Ochman et al. (2000), that the proportion

of horizontally transferred genes is independent of the amount of protein-coding

sequences in complete prokaryotic genomes. However, when comparing the two

community fractions, results here show the proportion of genome length occupied by

putative GIs and prophage was slightly, but significantly higher in Patescibacteria than

the rest of the communities – on average 3.0 ± 4.4% versus 2.0 ± 2.8%, respectively

(Fig. 1c).

The extent of gene transfer events detected here is comparable to those

detected in a variety of other phylogenetically diverse microbial communities, e.g.

90.4% of groundwater MAGs containing detectable HGT events, compared to 78.8%

in cheese-associated communities, and 89.5% in prokaryotic isolate genomes

(Bonham et al., 2017; Ochman et al., 2000). However, the extent of genetic exchange

observed in the groundwater microbial community is likely underestimated. Ancient

transfer events will escape detection due to robust integration within the recipient

genome via modification of GC content and tetranucleotide frequencies (Lawrence &

Ochman, 1997). Moreover, while MetaCHIP has proven efficient in predicting HGT

donors/recipients relationships within microbial communities, donor taxa might not

have been captured in the MAG dataset.

**Figure 3.** Length distribution of GIs and prophage in ultra-small MAGs and other
groundwater microbial MAGs. The center line of each boxplot represents the median;
the top and bottom lines are the first and third quartiles, respectively; and the whiskers
show 1.5× the interquartile range. No significant differences were detected in GI and
prophage sequence length between ultra-small and other groundwater
microorganisms (Wilcoxon Signed Rank, p>0.05).

**HGT origins in Patescibacteria taxa**

Most HT events detected occurred between members of the Patescibacteria

(Fig. 4a), with 104 of the 123 HT genes received by Patescibacteria originating from

counterparts. This observation is consistent with research, which demonstrates that

HGT is most prevalent among closely related organisms (Bonham et al., 2017), and

that intra-species HGT occurs at far greater frequency (Majewski et al., 2000). HGT

was likely also facilitated by the spatial proximity of microorganisms (Gogarten et al.,

2002), in particular by microbial cohorts in groundwater communities (Hug et al.,

2015). This includes cohorts formed by ultra-small prokaryotes, where individuals

spatially co-occur as described previously in subsurface and soil ecosystems (Geesink

et al., 2020; Herrmann et al., 2019; Nicolas et al., 2021).

295
**Figure 4. Origins of HT genes in Patescibacteria taxa.** (a) Gene flow among phyla
donating to Patescibacteria taxa. Scale represents number of HT events and inner
bands show donor/recipient relationship (coloured according to donor organism). (b)
Example of HT gene transferred from Omnitrophota nzgw326 (upper plot) to
Patescibacteria nzgw387 (lower plot). Genes encoded on the forward strand are
displayed in blue, and genes coded on the reverse strand are displayed in green. (c)
Genome relative abundance profiles of these 2 organisms across 16 groundwater
samples.

Although a smaller number of MAGs were recovered from Omnitrophota and

Bacteroidota, organisms in these phyla shared genes more frequently with

Patescibacteria (3.5x and 3x more, respectively) than, for example, Proteobacteria,

which was the second most populous phylum (Tables 1, S3). Frequent sharing

suggests potential close relationships between these organisms and ultra-small

prokaryotes (i.e. host/symbiont). The former two bacterial groups have been identified

previously as putative HGT donors to Patescibacteria in seafloor environments (Dong

et al., 2021). Moreover, Omnitrophota and Bacteroidota organisms were found to

consistently co-occur with the ultra-small bacteria in aquifers (Chaudhari et al., 2021).

**Table 1.** Frequency of HT events to Patescibacteria at phylum level. Note the ratio of
donor HGT events to number of MAGs is biases in donor phyla with small MAG sample
sizes (Methylomirabilota, Firmicutes_B, Desulfobacterota, RBG-13-66-14).

| Donor phylum | Donor HGT events | Reconstructed donor MAGs | Ratio donor HGT events / number of MAGs |
| --- | --- | --- | --- |
| Patescibacteria | 109 | 132 | 0.83 |
| Omnitrophota | 6 | 28 | 0.21 |
| Bacteroidota | 6 | 33 | 0.18 |
| Nanoarchaeota | 2 | 23 | 0.09 |
| Proteobacteria | 3 | 51 | 0.06 |
| Actinobacteriota | 1 | 24 | 0.04 |
| Methylomirabilota | 1 | 3 | 0.33 |
| Firmicutes_B | 1 | 1 | 1.00 |
| Desulfobacterota | 1 | 1 | 1.00 |
| RBG-13-66-14 | 1 | 1 | 1.00 |

We observed further evidence for recent putative horizontal transfers to

groundwater Patescibacteria from patescibacterial counterparts and other taxa based

on positive correlations between the relative abundances of HGT pairs across sites

(29 out of 131 pairs involving Patescibacteria as donor or recipient with Pearson's

correlation coefficient >0.5, $p <0.05$, Table S4). For example, Patescibacteria nzgw387

received a copy of the *lysR* gene from Omnitrophota nzgw326 (Fig. 4b), a highly

conserved transcriptional regulator involved in numerous cellular functions (Maddocks

& Oyston, 2008), and is shown to co-occur with its donor, Omnitrophota nzgw326, in

groundwater (Fig. 4c).

A BLAST analysis of GI genes against the gene content of whole groundwater

communities showed that most of the genes located on GIs could not be matched to

any organisms within the communities (91.9% of GI genes; Table S5). Nonetheless,

we found that 7% of GI genes were related to those in Patescibacteria other than the

one carrying the GI. Interestingly, the content of one 4,250 bp long GI in

Patescibacteria nzgw468 (74% genome completeness, no contamination) was highly

similar to genes in the archaeon Halobacterota nzgw24 (99% completeness, 1.3%

contaminated), indicating a potential recent transfer event between the two distantly

related organisms. None of the 11 genes located on the GI could be functionally

characterized (as for a large proportion of genes in Patescibacteria genomes), but

were between 85 and 100% identical in amino acid composition to the ones of nzgw24.

While inter-domain transfers do occur between Archaea and Bacteria, they are less

frequent than bacterial-domain confined HGT events (Nelson et al., 1999, López-

García et al., 2015). Gene exchange between Patescibacteria and their archaeal ultra-

small counterparts (DPANN) has been reported (León-Zayas et al., 2017), particularly

in the context of membrane-associated protein encoding genes potentially involved in

host-symbiont interactions (Jaffe et al., 2016). Accordingly, we detected 9 HT events

from Patescibacteria to DPANN archaea (Micrarchaeota and Nanoarchaeota; function

of transferred genes discussed below). A further, 2 genes (one ABC-transporter and

one gene related to the BRO family, of which the function remains unknown) were

transferred from Nanoarchaeota to Patescibacteria (Table S6).

**Metabolic functions encoded in horizontally acquired regions**

Patescibacteria, or CPR bacteria, have reduced genomes and limited biosynthetic capacities (Castelle et al., 2018). The metabolic function of horizontally transferred genes that are retained in Patescibacteria genomes could provide clues about the evolution of dependencies in these ultra-small prokaryotes regarding their symbiotic interactions with hosts. Although a substantial portion of genes located in horizontally acquired genomic regions could not be assigned any function (61.6% of all genes for Patescibacteria, 38.9% for other taxa), genes located in those regions in Patescibacteria genomes that were able to be annotated are involved in a wide range of metabolic pathways (Fig. 5), and include a large number of genes involved in translation and transcription. Horizontally acquired translation and transcription genes predominantly comprised ribosomal proteins (small and large subunits), elongation factor *tuf*, which plays a central role in protein synthesis, and chaperonin *groS* (Table S6). These results reflect trends identified when considering the most frequently transferred genes to other groundwater prokaryotes, which notably included the *rpsL* gene encoding ribosomal protein S12 and transcription initiation factor *infA*, alongside translation elongation factor *tuf*. Kanhere and colleagues described similar findings (Kanhere & Vingron, 2009). By inferring HGT events based on evolutionary distances within orthologous protein families (using the COG database), authors found that genes exchanged among bacteria were primarily involved in translation (Kanhere & Vingron, 2009).

**Figure 5. Metabolic functions of genes received through horizontal transfer.** Metabolic functions acquired through HGT (HT and GI genes) by Patescibacteria (124 HT genes + 120 GIs received) and other groundwater microorganisms (1,283 HT genes + 926 GIs received) from any other member of the groundwater community (top). Breakdown of metabolic functions of HT genes (middle) and genes located on GIs (bottom).

HT genes in Patescibacteria were found to be highly enriched, compared to other phyla, in genes assigned to the EggNOG "Defense mechanisms" category (Fig. 5). These included genes (e.g. *macB* and *msbA*) encoding ABC-type systems for multidrug and antimicrobial peptide transport. Membrane-located multidrug transporters have been identified previously in relatives of ultra-small bacterium *Babela massiliensis* (Dependentiae phylum, previously Candidate division TM6). Their specialization is thought to be important at the host-symbiont interface (Jaffe et al., 2016). There is growing evidence that drug transporters are required for survival of

intracellular pathogenic bacteria in host cells (Du et al., 2015). Genes assigned to

chromatin structure were also more prevalent in the HT gene pool in Patescibacteria.

Among these genes were six SWIB/MDM2 domain containing proteins, which are

chromatin-remodelling proteins involved in transcriptional activation (Tang et al.,

2010), and were almost exclusively transferred between Patescibacteria (Table S6).

DNA packaging in the form of chromatin is well documented in Eukaryotes and

Archaea, but remains poorly understood in Bacteria (Shen & Landick, 2019). Whether

ultra-small bacteria use chromatin-remodelling proteins to manipulate their own

genomic DNA structure, or for the modification of eukaryotic/archaeal symbiotic host

DNA is unclear.

In contrast, other groundwater taxa more frequently acquired genes involved in

energy production (respiratory chain) and amino acid biosynthesis, which

Patescibacteria are known to consistently lack (Castelle et al., 2018). This could

further indicate that ultra-small bacteria must scavenge resources such as protons and

amino acids from the host or surrounding environment. A higher proportion of genes

involved in secondary metabolite pathways and signal transduction mechanisms was

also detected in other taxa (426 HT and GI genes in 105 'other' MAGs versus 18 genes

in 15 ultra-small MAGs) (Table S6). Secondary metabolites are considered luxury

metabolism (Demain & Fang, 2000) and can constitute public goods (e.g.

siderophores) (Weigert & Kümmerli, 2017). As such, biosynthetic gene clusters

encoding for these metabolites are uncommon in streamlined genomes

(Konstantinidis & Tiedje, 2004; Sharrar et al., 2020), which have reduced auxiliary

functions (Giovannoni et al., 2014), and are more likely to depend on the metabolites

produced by other organisms (Morris et al., 2012). Similarly, loss of signal transduction

mechanisms, which facilitate environmental monitoring, is a characteristic feature of

streamlined bacterial genomes, including those of bacteria with non-intracellular

lifestyles (e.g. bacterioplankton) (Gabaldón & Huynen, 2007; Swan et al., 2013).

Although it has been shown that gene function impacts HGT overall, studies

assessing functional categories of horizontally transferred genes have reported

diverging results. The 'complexity hypothesis' suggests that genes involved in cellular

housekeeping are more frequently subjected to horizontal transfer, while informational

genes (involved in transcription, translation, and related processes are more resistant

to transfers (Jain et al., 1999). Nakamura et al. (2004) found that horizontally

transferred genes are biased towards cell surface, DNA binding and pathogenicity

related functions. Conversely, other studies have found that the functional category of

a gene family is an insignificant factor in determining HGT (Choi & Kim, 2007; Cohen

et al., 2011). While we found that the genes localized in GIs harbour similar metabolic

functions overall between Patescibacteria and the rest of the groundwater community,

functions of individual genes acquired by the ultra-small bacteria through horizontal

transfer tend to differ (Fig. 5). Results obtained here therefore suggest lifestyle of the

recipient impacts the traits of genes transferred and also retained.

Biases towards horizontally acquired features distinct from the general

groundwater communities by Patescibacteria may be explained by constraints on

streamlined genomes. Selection of distinct features could occur by two mechanisms:

(i) retention of horizontally transferred genes and concentration due to selective

deletion of other less desirable features, and (ii) selective retention of acquired genes

with desirable features. The caveat being that the distribution of functions among the

hypothetical coding DNA sequence (CDS) fraction in Patescibacteria is unknown.

We could question whether the length of transferred DNA is an important factor

affecting integration and retainment of transferred genes and related metabolic

functions in streamlined ultra-small and other groundwater prokaryotes. The length of

transferred DNA was 10,324 ± 9,397 bp on average for the 1,046 detected GIs (10.8

Mbp total in all groundwater taxa) and 899 ± 562 bp for the 1,501 HT genes (1.3 Mbp

overall) (Table S2). Therefore, while a third less GIs than individually transferred genes

were detected in terms of number, the total length of sequences transferred was

substantially greater with GIs, and broadly equivalent between Patescibacteria (9,689

bp per MAG) and other groundwater microorganisms (10,406 bp per MAG). Besides

sequence length, it has been shown that recombination efficiency decreases

exponentially with sequence divergence (Majewski et al., 2000; Majewski & Cohan,

1999). However, the presence of flanking regions of identity in exchangeable regions

can remove most barriers to recombination. The shortest length of sequence

homology necessary for efficient recombination can vary greatly depending on the

organism, the recombination pathway used, and other factors (Thomas & Nielsen,

2005). In *Escherichia coli*, efficient recombination has been observed with as little as

23 bp of sequence homology (Shen & Huang, 1986).

**Impact of phylogenetic distance and direction of transfer**

We assessed how phylogenetic distance between donor-recipient pairs associates

with the metabolic function of genes acquired by Patescibacteria. Genetic exchange

occurred over a wide range of phylogenetic distances based on average amino acid

identities between genomes of groundwater microorganisms (as low as 38.5% AAI,

Fig. 6a). Within the Patescibacteria phylum, genes of diverse metabolic functions were

transferred regardless of phylogenetic distance (41.0-54.3% AAI). However, direction

of HT events (between members of the Patescibacteria phylum, other groundwater

microorganisms to Patescibacteria, and Patescibacteria to other groundwater

microorganisms) appeared to have an impact on the metabolic function of acquired

genes (Fig. 6b). A number of metabolic functions were found to be exclusively

transferred between Patescibacteria (encoded by 12 out of 104 HT genes shared

between Patescibacteria), and were involved in processes such as amino acid

metabolism (serine hydroxymethyltransferase, GlyA), cell division (actin homolog,

MreB), intracellular trafficking, co-enzyme metabolism (riboflavin biosynthesis) and

inorganic ion metabolism (cation transport) (Fig. 6b, Table S6).

**Figure 6. Pangenome of Patescibacteria and effect of phylogenetic distance and**
**direction of transfer on acquired genes.** (a) Impact of phylogenetic distance of HGT
donor to Patescibacteria (calculated as AAI) on metabolic function of horizontally
acquired genes (left) and pangenome category (right). (b) Metabolic function of HT
genes according to direction of transfer. (c) Proportion of Patescibacteria total gene
pool contained in orthogroups vs. unique, and breakdown of proportion of orthogroups
in 'core', 'shared' and 'species-specific' pangenome categories (top). Proportion of
horizontally acquired genes (HT genes and GI genes) by Patescibacteria taxa in each
pangenome category (bottom).

Others were exclusively shared from Patescibacteria to other phyla (out of n=29

transfers overall, Fig. 6b), namely two genes associated with chromatin structure and

dynamics (SWIB/MDM2 domain-containing protein) and carbohydrate metabolism

(mannose-6-phosphate isomerase, COG0662). Moreover, Patescibacteria donated a

greater fraction of replication, recombination and repair genes (GIY-YIG catalytic

domain-containing protein, UvrD subfamily helicase, bacterial nucleoid DNA-binding

protein COG0776, DNA-3-methyladenine glycosylase) to other bacteria. Among the

nine HT events from Patescibacteria to DPANN archaea reported above, three were

related to a glyoxalase (Table S6), which is key metalloenzyme in the glycolytic

pathway involved in the detoxification of reactive methylglyoxal into D-lactate. It is a

common feature across domains of life, including in Archaea (Rawat & Maupin-Furlow,

2020), and was previously reported in CPR single amplified genomes (León-Zayas et

al., 2017). The reason ultra-small archaea exhibit an apparent preference for acquiring

bacterial homologs is unclear.

A very small number of HGT events (two) also occurred exclusively from other

prokaryotes to Patescibacteria (out of n=20 transfers overall, Fig 6b). This includes

the *acpP* gene encoding an acyl carrier protein, and a gene encoding a putative

lysophospholipase       (COG2267,       alpha       beta       hydrolase       superfamily).

Lysophospholipases catalyse phospholipid hydrolysis and generate fatty acids. A

recent study coupling lipidomics and metagenomics approaches suggested that

members of the Patescibacteria phylum, that lack capacities for fatty acid biosynthesis

(Castelle et al., 2018), are able to recycle lipids from other bacteria for cell membrane

biogenesis (Probst et al., 2020). This suggests that HGT could facilitate the sourcing

of fatty acids in Patescibacteria. Phospholipase genes have also been reported in the

genomes of other ultra-small CPR prokaryotes (for example, in amoeba endosymbiont

*Vampirococcus lugosii*; Moreira et al., 2021). Patescibacteria were also more likely to

receive genes from other groundwater microorganisms associated with nucleotide

metabolism (n=2), including one thymidylate synthase genes *thyA.* The enzyme

encoded catalyzes the *de novo* synthesis of dTMP (deoxythymidine monophosphate),

an essential precursor for DNA biosynthesis (Voeller et al., 1995), which is a capacity

already encoded by the minimal biosynthetic gene repertoire of groundwater

Patescibacteria. Our results indicate that this essential function can also be

horizontally acquired from other members of groundwater microbial communities.

While the number of transfer events recorded from general groundwater

microorganisms to Patescibacteria and vice versa is low, limiting robust conclusions,

these numbers are consistent with known associations between HGT and

phylogenetic distance among prokaryotes more generally (Majewski et al., 2000).

Overall, metabolic functions gained via HGT in ultra-small prokaryotes remain poorly

studied, regardless of phylogenetic origin. Events identified here tend to confer

Patescibacteria with critical functions, such as the ability to degrade lipids, and carry

out the biosynthesis of nucleotides.

**Patescibacteria pangenome and HGT**

The evolutionary fate of a HGT event is determined by fitness conferred to the recipient

genome and cell, i.e., whether the acquisition of exogenous DNA is beneficial, neutral

or deleterious. In order to evaluate the importance of the metabolic functions acquired

through HGT in Patescibacteria, we performed a pangenome analysis of the 125

reconstructed MAGs. The pangenome of all the examined Patescibacteria contained

7,732 orthogroups, comprising 88,187 genes (74.3%) out of the total 118,635

predicted genes. Genes in the pan-genome were classified into 3 groups based on

their occurrence: (1) the "core" genes, which are present in at least 2/3 of all 132

genomes, (2) the "shared" genes are found in more than one genome, but are not core

genes, and (3) the "unique" genes, comprising species-specific genes (i.e. genes

present in one genome only) and genes not assigned to an orthogroup. Both shared

and unique genes comprise the accessory genome. Results indicated the vast majority

of orthogroups (96%) were shared between the groundwater Patescibacteria, and only

3% of orthogroups (2,634 genes) comprised the core genome (Fig. 6c). This

represents the lower end of core genome size among bacterial pangenomes (Bosi et

al., 2017; Shin et al., 2016) and could be partially due to MAG incompleteness, but is

consistent with the variation found in the core protein families by Méheust and

colleagues across members of the CPR radiation (Méheust et al., 2019).

GIs brought many unique, and hence divergent, genes to the Patescibacteria

pangenome (41.2% of GI genes, Fig. 6c). These observations are consistent with

previous work suggesting that prokaryotes contain a higher proportion of novel genes

in GIs compared with the rest of their genome (Hsiao et al., 2005). HT genes acquired

by Patescibacteria species were spread across the core and the shared gene fractions

of these organisms (45.8% shared and 51.9% core, Fig. 6c), and not the unique

fraction. A high proportion of core or shared genes would be expected given predicted

transfers occurred mostly between Patescibacteria species (83.2% of HT events, Fig.

4a). However, the lack of any unique genes reflects limitations in the detection of HT

events by the MetaCHIP tool, which considers self matches to be false positive (i.e. a

donor is required) (Song et al., 2019).

Streamlined genomes are characterised by low numbers of paralogs

(Giovannoni et al., 2014). Accordingly, among all genes in Patescibacteria, we

detected a low number of paralogous orthogroups based on the pangenome analysis

results (89 genes in 36 species-specific orthogroups – i.e. 89 paralogs out of 118,635

genes). We found that no genes located on GI regions nor HT genes were paralogs

of other genes in the Patescibacteria pangenome. Gene duplicates are generally

under purifying selection (Lynch & Conery, 2000), and have been shown to evolve

faster than non-duplicated genes with a similar level of divergence (Kondrashov et al.,

2002), introducing new proteins. Nonetheless, gene duplication is usually widespread

in bacterial genomes, with an estimated 7–41% of bacterial proteins encoded by

paralogs (Gevers et al., 2004), and is a common strategy for creating novel gene

functions for niche expansion in prokaryotes (Hooper & Berg, 2003). However, it has

been quantitatively demonstrated that gene gain via HGT rather than gene duplication

is the principal factor of innovation in the evolution of prokaryotes (Treangen & Rocha,

2011). Based on a comparably high proportion of HGT into Patescibacteria as for other

phyla, results here demonstrate HGT, and unique genes conferred by GIs, are almost

exclusively used over duplication by Patescibacteria to expand niche ranges.

**GI in Patescibacteria nzgw456 encoding a full type I restriction-modification**

**system**

Streamlined bacteria (and archaea) have been hypothesized to be largely lacking

classical defense mechanisms, such as CRISPR-Cas (Clustered Regularly

Interspaced Short Palindromic Repeats – CRISPR associated), restriction-

modification (R-M) and toxin-antitoxin systems (Koonin et al., 2017). Similarly, the

analysis of hundreds of CPR/Patescibacteria genomes recovered from diverse

environments revealed that most are missing CRISPR-Cas adaptive immune systems,

lost in the process of genome streamlining, and instead rely nonspecific defence, such

as R-M systems (Castelle et al., 2018). Although first believed to constitute a barrier

to genetic exchange between organisms (Vasu & Nagaraja, 2013), it has been recently

suggested that these systems play an important role in HGT rates by increasing the

frequency of exchange in organisms possessing numerous R-M systems (Oliveira et

al., 2016).

**Figure 7. Genomic island encoding a full type I restriction-modification system in Patescibacteria nzgw456.** Left: circular genome plot with following rings (inner to outer): GIs (in blue; star indicates location of GI depicted in the right panel); GC content; contig boundaries. Right: Linear map of GI (indicated by grey box) and protein-coding genes. Manual examination of GI was undertaken to further confirm boundaries using the DarkHorse prediction tool (Podell et al., 2008).

Out of the 120 GIs detected in Patescibacteria we identified four encoding restriction modification systems or endonucleases in patescibacterial genomes. All four MAGs contained a high number of GIs compared to other patescibacterial counterparts (average 5 ± 2.8 SD vs 1.7 ± 1.0), and three acquired more HT per Mbp than the average for Patescibacteria overall (average 2.5 ± 1.3 SD vs 1.2 ± 1.5, Table S2). One GI in Patescibacteria nzgw456 (Paceibacteria class) encodes a full type I R-M system (Fig. 7). Type I R-M systems, which are the most complex of all four known R-M systems (Murray, 2000), comprise three subunits, namely HsdR and HsdM (required for methyltransferase activity), and HsdS (required for restriction). Co-located with genes encoding these subunits on the GI was one putative ATP-dependent endonuclease. Type I R-M related genes were also found in a GI present in Patescibacteria nzgw350, and other restriction endonucleases in nzgw410 and nzgw504 (Table S7). Acquisition of type I R-M genes via HGT was recently described in CPR bacterium *V. lugosii* (Moreira et al., 2021), however not as syntenic genes. It has been suggested that R-M systems could be used by putative epibionts as a way

to source of nucleotides by degrading exogenous DNA, including that derived from

viruses (Burstein et al., 2016), in addition to protecting against phage infection (Stern

& Sorek, 2011).

**CONCLUSION**

Recent phylogenetic analyses have shed light on the massive bacterial group that is

the Patescibacteria phylum/CPR (Hug et al., 2016). Due to the reductive evolution of

Patescibacteria, it is essential to consider the processes of gene gain and loss in

respect of the predicted symbiotic lifestyles of these ultra-small microorganisms.

Results here indicate that Patescibacteria are involved in dynamic genetic exchange

with their ultra-small counterparts and other members of the microbial community. We

found a total 124 HT events involving Patescibacteria, 24 prophage and over 100 GIs

in Patescibacteria genomes. While most transfers occurred among Patescibacteria,

HGT origins were diverse and widespread across both bacterial and archaeal

domains, in particular with members of the Omnitrophota and Bacteroidota phyla. A

putative host symbiont pair was identified based on HGT evidence and co-occurrence

across groundwater samples. Functions encoded by horizontally acquired regions

comprised diverse pathways, but mainly transcription, translation and DNA replication,

recombination and repair, such as ribosomal proteins and transcription factors.

Findings show that acquired genes were distributed across core and accessory

portions of Patescibacteria genomes, and that GIs generally introduced

phylogenetically distant sequences. Some of the metabolic functions horizontally

acquired by Patescibacteria are distinct from those acquired by the general

groundwater communities, suggesting unique pressures on gene gain/loss dynamics

occurring in ultra-small prokaryotes.

**ACKNOWLEDGMENTS**

Funding was provided by a MBIE Smart Ideas grant awarded to K.M.H. (project UOAX1720). We thank D. Waite and J.S. Boey for help with bioinformatics. Computational resources were provided by New Zealand eScience Infrastructure.

The authors declare no conflict of interest.

**REFERENCES**

Anantharaman, K., Brown, C. T., Hug, L. A., Sharon, I., Castelle, C. J., Probst, A. J., Thomas, B. C., Singh, A., Wilkins, M. J., Karaoz, U., Brodie, E. L., Williams, K. H., Hubbard, S. S., & Banfield, J. F. (2016). Thousands of microbial genomes shed light on interconnected biogeochemical processes in an aquifer system. *Nature Communications*, *7*, 1–11. https://doi.org/10.1038/ncomms13219

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., Lesin, V. M., Nikolenko, S. I., Pham, S., Prjibelski, A. D., Pyshkin, A. V, Sirotkin, A. V, Vyahhi, N., Tesler, G., Alekseyev, M. A., & Pevzner, P. A. (2012). SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology : A Journal of Computational Molecular Cell Biology*, *19*(5), 455–477. https://doi.org/10.1089/cmb.2012.0021

Bertelli, C., Laird, M. R., Williams, K. P., Lau, B. Y., Hoad, G., Winsor, G. L., & Brinkman, F. S. L. (2017). IslandViewer 4: Expanded prediction of genomic islands for larger-scale datasets. *Nucleic Acids Research*, *45*(W1). https://doi.org/10.1093/nar/gkx343

Bonham, K. S., Wolfe, B. E., & Dutton, R. J. (2017). Extensive horizontal gene transfer in cheese-associated bacteria. *ELife*, *6*.

https://doi.org/10.7554/eLife.22144

Bosi, E., Fondi, M., Orlandini, V., Perrin, E., Maida, I., de Pascale, D., Tutino, M. L.,

Parrilli, E., Lo Giudice, A., Filloux, A., & Fani, R. (2017). The pangenome of

(Antarctic) Pseudoalteromonas bacteria: Evolutionary and functional insights.

*BMC Genomics*, *18*(1). https://doi.org/10.1186/s12864-016-3382-y

Brown, C. T., Hug, L. A., Thomas, B. C., Sharon, I., Castelle, C. J., Singh, A.,

Wilkins, M. J., Wrighton, K. C., Williams, K. H., & Banfield, J. F. (2015). Unusual

biology across a group comprising more than 15% of domain Bacteria. *Nature*,

*523*(7559), 208–211. https://doi.org/10.1038/nature14486

Burstein, D., Sun, C. L., Brown, C. T., Sharon, I., Anantharaman, K., Probst, A. J.,

Thomas, B. C., & Banfield, J. F. (2016). Major bacterial lineages are essentially

devoid of CRISPR-Cas viral defence systems. *Nature Communications*, *7*.

https://doi.org/10.1038/ncomms10613

Castelle, C. J., Brown, C. T., Anantharaman, K., Probst, A. J., Huang, R. H., &

Banfield, J. F. (2018). Biosynthetic capacity, metabolic variety and unusual

biology in the CPR and DPANN radiations. *Nature Reviews Microbiology*,

*16*(10), 629–645. https://doi.org/10.1038/s41579-018-0076-2

Castelle, C. J., Brown, C. T., Thomas, B. C., Williams, K. H., & Banfield, J. F. (2017).

Unusual respiratory capacity and nitrogen metabolism in a Parcubacterium

(OD1) of the Candidate Phyla Radiation. *Scientific Reports*, *7*(October 2016), 1–

12. https://doi.org/10.1038/srep40101

Chaudhari, N. M., Overholt, W. A., Figueroa-Gonzalez, P. A., Taubert, M.,

Bornemann, T. L. V., Probst, A. J., Hölzer, M., Marz, M., & Küsel, K. (2021). The

economical lifestyle of CPR bacteria in groundwater allows little preference for

environmental drivers. *BioRxiv*, 2021.07.28.454184.

https://www.biorxiv.org/content/10.1101/2021.07.28.454184v1%0Ahttps://www.b

iorxiv.org/content/10.1101/2021.07.28.454184v1.abstract

Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P., & Parks, D. H. (2020). GTDB-Tk: a

toolkit to classify genomes with the Genome Taxonomy Database.

*Bioinformatics*, *36*(6), 1925–1927. https://doi.org/10.1093/bioinformatics/btz848

Choi, I. G., & Kim, S. H. (2007). Global extent of horizontal gene transfer.

*Proceedings of the National Academy of Sciences of the United States of*

*America*, *104*(11), 4489–4494. https://doi.org/10.1073/pnas.0611557104

Close, M., Abraham, P., Webber, J., Cowey, E., Humphries, B., Fenwick, G.,

Howard, S., Huynh, K., Grace, T., Dupont, P. Y., & Weaver, L. (2020). Use of

Sonication for Enhanced Sampling of Attached Microbes from Groundwater

Systems. *Groundwater*, *58*, 901–915. https://doi.org/10.1111/gwat.12984

Cohen, O., Gophna, U., & Pupko, T. (2011). The complexity hypothesis revisited:

Connectivity Rather Than function constitutes a barrier to horizontal gene

transfer. *Molecular Biology and Evolution*, *28*(4).

https://doi.org/10.1093/molbev/msq333

Cordero, O. X., & Hogeweg, P. (2009). The impact of long-distance horizontal gene

transfer on prokaryotic genome size. *Proceedings of the National Academy of*

*Sciences of the United States of America*, *106*(51), 21748–21753.

https://doi.org/10.1073/pnas.0907584106

Demain, A. L., & Fang, A. (2000). The natural functions of secondary metabolites. In

*Advances in biochemical engineering/biotechnology* (Vol. 69).

https://doi.org/10.1007/3-540-44964-7_1

Dong, X., Zhang, C., Li, W., Weng, S., Song, W., Li, J., & Wang, Y. (2021).

Functional diversity of microbial communities in inactive seafloor sulfide

deposits. *FEMS Microbiology Ecology*, *97*(8), 1–10.

https://doi.org/10.1093/femsec/fiab108

Du, D., van Veen, H. W., Murakami, S., Pos, K. M., & Luisi, B. F. (2015). Structure,

mechanism and cooperation of bacterial multidrug transporters. In *Current*

*Opinion in Structural Biology* (Vol. 33). https://doi.org/10.1016/j.sbi.2015.07.015

Dunning Hotopp, J. C., Clark, M. E., Oliveira, D. C. S. G., Foster, J. M., Fischer, P.,

Muñoz Torres, M. C., Giebel, J. D., Kumar, N., Ishmael, N., Wang, S., Ingram,

J., Nene, R. V., Shepard, J., Tomkins, J., Richards, S., Spiro, D. J., Ghedin, E.,

Slatko, B. E., Tettelin, H., & Werren, J. H. (2007). Widespread lateral gene

transfer from intracellular bacteria to multicellular eukaryotes. *Science*,

*317*(5845). https://doi.org/10.1126/science.1142490

Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST.

*Bioinformatics*, *26*, 2460–2461. https://doi.org/10.1093/bioinformatics/btq461

Emms, D. M., & Kelly, S. (2019). OrthoFinder: Phylogenetic orthology inference for

comparative genomics. *Genome Biology*, *20*(1). https://doi.org/10.1186/s13059-

019-1832-y

Flynn, T. M., Sanford, R. A., & Bethke, C. M. (2008). Attached and suspended

microbial communities in a pristine confined aquifer. *Water Resources*

*Research*, *44*(7), 1–7. https://doi.org/10.1029/2007WR006633

Gabaldón, T., & Huynen, M. A. (2007). From endosymbiont to host-controlled

organelle: The hijacking of mitochondrial protein synthesis and metabolism.

*PLoS Computational Biology*, *3*(11).

https://doi.org/10.1371/journal.pcbi.0030219

Garcia-Vallvé, S., Romeu, A., & Palau, J. (2000). Horizontal gene transfer in

bacterial and archaeal complete genomes. *Genome Research*, *10*(11).

https://doi.org/10.1101/gr.130000

Geesink, P., Wegner, C.-E., Probst, A. J., Herrmann, M., Dam, H. T., Kaster, A.-K., &

Küsel, K. (2020). Genome-inferred spatio-temporal resolution of an uncultivated

Roizmanbacterium reveals its ecological preferences in groundwater.

*Environmental Microbiology*, *22*(2), 726–737.

https://doi.org/https://doi.org/10.1111/1462-2920.14865

Gevers, D., Vandepoele, K., Simillion, C., & Van De Peer, Y. (2004). Gene

duplication and biased functional retention of paralogs in bacterial genomes. In

*Trends in Microbiology* (Vol. 12, Issue 4).

https://doi.org/10.1016/j.tim.2004.02.007

Giovannoni, S. J., Cameron Thrash, J., & Temperton, B. (2014). Implications of

streamlining theory for microbial ecology. In *ISME Journal* (Vol. 8, Issue 8).

https://doi.org/10.1038/ismej.2014.60

Gogarten, J. P., Doolittle, W. F., & Lawrence, J. G. (2002). Prokaryotic evolution in

light of gene transfer. *Molecular Biology and Evolution*, *19*(12).

https://doi.org/10.1093/oxfordjournals.molbev.a004046

Griebler, C., & Lueders, T. (2009). Microbial biodiversity in groundwater ecosystems.

*Freshwater Biology*, *54*(4), 649–677. https://doi.org/10.1111/j.1365-

2427.2008.02013.x

Gu, Z., Gu, L., Eils, R., Schlesner, M., & Brors, B. (2014). Circlize implements and

enhances circular visualization in R. *Bioinformatics*, *30*(19).

https://doi.org/10.1093/bioinformatics/btu393

Haft, D. H., Selengut, J. D., & White, O. (2003). The TIGRFAMs database of protein

families. In *Nucleic Acids Research* (Vol. 31, Issue 1).

https://doi.org/10.1093/nar/gkg128

He, C., Keren, R., Whittaker, M., Farag, I. F., Doudna, J., Cate, J. H. D., & Banfield,

J. (2021). Genome-resolved metagenomics reveals site-specific diversity of

episymbiotic CPR bacteria and DPANN archaea in groundwater ecosystems.

*Nature Microbiology*, *6*, 354–365. https://doi.org/10.1101/2020.05.14.094862

Herrmann, M., Wegner, C.-E., Taubert, M., Geesink, P., Lehmann, K., Yan, L.,

Lehmann, R., Totsche, K. U., & Küsel, K. (2019). Predominance of Cand.

Patescibacteria in Groundwater Is Caused by Their Preferential Mobilization

From Soils and Flourishing Under Oligotrophic Conditions. *Frontiers in*

*Microbiology*, *10*(June), 1–15. https://doi.org/10.3389/fmicb.2019.01407

Hooper, S. D., & Berg, O. G. (2003). Duplication is more common among laterally

transferred genes than among indigenous genes. *Genome Biology*, *4*(8).

https://doi.org/10.1186/gb-2003-4-8-r48

Hsiao, W. W. L., Ung, K., Aeschliman, D., Bryan, J., Brett Finlay, B., & Brinkman, F.

S. L. (2005). Evidence of a large novel gene pool associated with prokaryotic

genomic Islands. *PLoS Genetics*, *1*(5). https://doi.org/10.1371

/journal.pgen.0010062

Huerta-Cepas, J., Forslund, K., Coelho, L. P., Szklarczyk, D., Jensen, L. J., von

Mering, C., & Bork, P. (2017). Fast Genome-Wide Functional Annotation

through Orthology Assignment by eggNOG-Mapper. *Molecular Biology and*

*Evolution*, *34*(8), 2115–2122. https://doi.org/10.1093/molbev/msx148

Huerta-Cepas, J., Szklarczyk, D., Heller, D., Hernández-Plaza, A., Forslund, S. K.,

Cook, H., Mende, D. R., Letunic, I., Rattei, T., Jensen, L. J., von Mering, C., &

Bork, P. (2019). eggNOG 5.0: a hierarchical, functionally and phylogenetically

annotated orthology resource based on 5090 organisms and 2502 viruses.

*Nucleic Acids Research*, *47*(D1), D309–D314.

https://doi.org/10.1093/nar/gky1085

Hug, L. A., Thomas, B. C., Brown, C. T., Frischkorn, K. R., Williams, K. H., Tringe, S.

G., & Banfield, J. F. (2015). Aquifer environment selects for microbial species

cohorts in sediment and groundwater. *ISME Journal*, *9*(8), 1846–1856.

https://doi.org/10.1038/ismej.2015.2

Hyatt, D., Chen, G.-L., Locascio, P. F., Land, M. L., Larimer, F. W., & Hauser, L. J.

(2010). Prodigal: prokaryotic gene recognition and translation initiation site

identification. *BMC Bioinformatics*, *11*, 119. https://doi.org/10.1186/1471-2105-

11-119

Jaffe, A. L., Corel, E., Pathmanathan, J. S., Lopez, P., & Bapteste, E. (2016).

Bipartite graph analyses reveal interdomain LGT involving ultrasmall

prokaryotes and their divergent, membrane-related proteins. *Environmental*

*Microbiology*, *18*(12), 5072–5081. https://doi.org/10.1111/1462-2920.13477

Jain, R., Rivera, M. C., & Lake, J. A. (1999). Horizontal gene transfer among

genomes: The complexity hypothesis. *Proceedings of the National Academy of*

*Sciences of the United States of America*, *96*(7).

https://doi.org/10.1073/pnas.96.7.3801

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A., & Jermiin, L. S.

(2017). ModelFinder: fast model selection for accurate phylogenetic estimates.

*Nature Methods*, *14*(6), 587–589. https://doi.org/10.1038/nmeth.4285

Kanehisa, M., Sato, Y., Kawashima, M., Furumichi, M., & Tanabe, M. (2016). KEGG

as a reference resource for gene and protein annotation. *Nucleic Acids*

*Research*, *44*(D1), D457–D462. https://doi.org/10.1093/nar/gkv1070

Kanhere, A., & Vingron, M. (2009). Horizontal gene transfers in prokaryotes show

differential preferences for metabolic and translational genes. *BMC Evolutionary*

*Biology*, *9*(1), 1–13. https://doi.org/10.1186/1471-2148-9-9

Kieft, K., Zhou, Z., & Anantharaman, K. (2020). VIBRANT: automated recovery,

annotation and curation of microbial viruses, and evaluation of viral community

function from genomic sequences. *Microbiome*, *8*(1), 90.

https://doi.org/10.1186/s40168-020-00867-0

Kondrashov, F. A., Rogozin, I. B., Wolf, Y. I., & Koonin, E. V. (2002). Selection in the

evolution of gene duplications. *Genome Biology*, *3*(2).

https://doi.org/10.1186/gb-2002-3-2-research0008

Konstantinidis, K. T., & Tiedje, J. M. (2004). Trends between gene content and

genome size in prokaryotic species with larger genomes. *Proceedings of the*

*National Academy of Sciences*, *101*(9), 3160–3165.

https://doi.org/10.1073/pnas.0308653100

Koonin, E. V., Makarova, K. S., & Wolf, Y. I. (2017). Evolutionary Genomics of

Defense Systems in Archaea and Bacteria. *Annual Review of Microbiology*, *71*.

https://doi.org/10.1146/annurev-micro-090816-093830

Langmead, B., Trapnell, C., Pop, M., & Salzberg, S. L. (2009). Ultrafast and

memory-efficient alignment of short DNA sequences to the human genome.

*Genome Biology*, *10*(3), R25. https://doi.org/10.1186/gb-2009-10-3-r25

Lawrence, J. G., & Ochman, H. (1997). Amelioration of bacterial genomes: Rates of

change and exchange. *Journal of Molecular Evolution*, *44*(4).

https://doi.org/10.1007/PL00006158

León-Zayas, R., Peoples, L., Biddle, J. F., Podell, S., Novotny, M., Cameron, J.,

Lasken, R. S., & Bartlett, D. H. (2017). The metabolic potential of the single cell

genomes obtained from the Challenger Deep, Mariana Trench within the

candidate superphylum Parcubacteria (OD1). *Environmental Microbiology*,

*19*(7), 2769–2784. https://doi.org/10.1111/1462-2920.13789

Letunic, I., & Bork, P. (2007). Interactive Tree Of Life (iTOL): an online tool for

phylogenetic tree display and  annotation. *Bioinformatics (Oxford, England)*,

*23*(1), 127–128. https://doi.org/10.1093/bioinformatics/btl529

Lynch, M., & Conery, J. S. (2000). The evolutionary fate and consequences of

duplicate genes. *Science*, *290*(5494).

https://doi.org/10.1126/science.290.5494.1151

Maddocks, S. E., & Oyston, P. C. F. (2008). Structure and function of the LysR-type

transcriptional regulator (LTTR) family proteins. *Microbiology*, *154*(12), 3609–

3623. https://doi.org/10.1099/mic.0.2008/022772-0

Majewski, J., & Cohan, F. M. (1999). DNA sequence similarity requirements for

interspecific recombination in Bacillus. *Genetics*, *153*(4).

https://doi.org/10.1093/genetics/153.4.1525

Majewski, J., Zawadzki, P., Pickerill, P., Cohan, F. M., & Dowson, C. G. (2000).

Barriers to genetic exchange between bacterial species: Streptococcus

pneumoniae transformation. *Journal of Bacteriology*, *182*(4).

https://doi.org/10.1128/JB.182.4.1016-1023.2000

Mandel, M. J., Wollenberg, M. S., Stabb, E. V., Visick, K. L., & Ruby, E. G. (2009). A

single regulatory gene is sufficient to alter bacterial host range. *Nature*,

*458*(7235). https://doi.org/10.1038/nature07660

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput

sequencing reads. *EMBnet.Journal; Vol 17, No 1: Next Generation Sequencing*

*Data Analysis*. https://doi.org/10.14806/ej.17.1.200

Méheust, R., Burstein, D., Castelle, C. J., & Banfield, J. F. (2019). The distinction of

CPR bacteria from other bacteria based on protein family content. *Nature*

*Communications*, *10*(1), 4173. https://doi.org/10.1038/s41467-019-12171-z

Moran, N. A., & Bennett, G. M. (2014). The Tiniest Tiny Genomes. *Annual Review of*

*Microbiology*, *68*(1), 195–215. https://doi.org/10.1146/annurev-micro-091213-

112901

Morris, J. J., Lenski, R. E., & Zinser, E. R. (2012). The black queen hypothesis:

Evolution of dependencies through adaptive gene loss. *MBio*, *3*(2), 1–7.

https://doi.org/10.1128/mBio.00036-12

Mosley, O. E., Gios, E., Weaver, L., Close, M., Daughney, C., van der Raaij, R.,

Martindale, H., & Handley, K. M. (2022). Metabolic Diversity and Aero-Tolerance

in Anammox Bacteria from Geochemically Distinct Aquifers. *MSystems*, *7*(1),

2021.09.16.460709. https://doi.org/10.1128/msystems.01255-21

Murray, N. E. (2000). Type I Restriction Systems: Sophisticated Molecular Machines

(a Legacy of Bertani and Weigle). *Microbiology and Molecular Biology Reviews*,

*64*(2). https://doi.org/10.1128/mmbr.64.2.412-434.2000

Nakamura, Y., Itoh, T., Matsuda, H., & Gojobori, T. (2004). Biased biological

functions of horizontally-transferred genes in prokaryotic genomes. *Nature*

*Genetics*, *36*(7), 760–766. https://doi.org/10.1038/ng1381

Nayfach, S., Camargo, A. P., Schulz, F., Eloe-Fadrosh, E., Roux, S., & Kyrpides, N.

C. (2021). CheckV assesses the quality and completeness of metagenome-

assembled viral genomes. *Nature Biotechnology*, *39*(5).

https://doi.org/10.1038/s41587-020-00774-7

Nelson, W. C., & Stegen, J. C. (2015). The reduced genomes of Parcubacteria

(OD1) contain signatures of a symbiotic lifestyle. *Frontiers in Microbiology*,

*6*(JUN), 1–14. https://doi.org/10.3389/fmicb.2015.00713

Nguyen, L.-T., Schmidt, H. A., von Haeseler, A., & Minh, B. Q. (2015). IQ-TREE: A

Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood

Phylogenies. *Molecular Biology and Evolution*, *32*(1), 268–274.

https://doi.org/10.1093/molbev/msu300

Nguyen, M. T. H. D., Liu, M., & Thomas, T. (2014). Ankyrin-repeat proteins from

sponge symbionts modulate amoebal phagocytosis. *Molecular Ecology*, *23*(6).

https://doi.org/10.1111/mec.12384

Nicolas, A. M., Jaffe, A. L., Nuccio, E. E., Taga, M. E., Firestone, M. K., & Banfield,

J. F. (2021). Soil Candidate Phyla Radiation Bacteria Encode Components of

Aerobic Metabolism and Co-occur with Nanoarchaea in the Rare Biosphere of

Rhizosphere Grassland Communities. *MSystems*, *6*(4).

https://doi.org/10.1128/msystems.01205-20

Ochman, Howard, Lawrence, Jeffrey G., & Groisman, Eduardo A. (2000). Lateral

gene transfer and the nature of bacterial innovation. *Nature*, *405*(6784), 299–

304. http://adsabs.harvard.edu/abs/2000Natur.405..299O

Oliveira, P. H., Touchon, M., & Rocha, E. P. C. (2016). Regulation of genetic flux

between bacteria by restriction-modification systems. *Proceedings of the*

*National Academy of Sciences of the United States of America*, *113*(20).

https://doi.org/10.1073/pnas.1603257113

Parks, D. H., Chuvochina, M., Chaumeil, P. A., Rinke, C., Mussig, A. J., &

Hugenholtz, P. (2020). A complete domain-to-species taxonomy for Bacteria

and Archaea. *Nature Biotechnology*, *38*(September).

https://doi.org/10.1038/s41587-020-0501-8

Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P., & Tyson, G. W. (2015).

CheckM: assessing the quality of microbial genomes recovered from isolates,

single cells, and metagenomes. *Genome Research*, *25*(7), 1043–1055.

https://doi.org/10.1101/gr.186072.114

Parks, D. H., Rinke, C., Chuvochina, M., Chaumeil, P. A., Woodcroft, B. J., Evans, P.

N., Hugenholtz, P., & Tyson, G. W. (2017). Recovery of nearly 8,000

metagenome-assembled genomes substantially expands the tree of life. *Nature*

*Microbiology*, *2*(11). https://doi.org/10.1038/s41564-017-0012-7

Potter, S. C., Luciani, A., Eddy, S. R., Park, Y., Lopez, R., & Finn, R. D. (2018).

HMMER web server: 2018 update. *Nucleic Acids Research*, *46*(W1).

https://doi.org/10.1093/nar/gky448

Probst, A. J., Elling, F. J., Castelle, C. J., Zhu, Q., Elvert, M., Birarda, G., Holman, H.

Y. N., Lane, K. R., Ladd, B., Ryan, M. C., Woyke, T., Hinrichs, K. U., & Banfield,

J. F. (2020). Lipid analysis of CO2-rich subsurface aquifers suggests an

autotrophy-based deep biosphere with lysolipids enriched in CPR bacteria.

*ISME Journal*, *14*(6). https://doi.org/10.1038/s41396-020-0624-4

Probst, A. J., Ladd, B., Jarett, J. K., Geller-McGrath, D. E., Sieber, C. M. K.,

Emerson, J. B., Anantharaman, K., Thomas, B. C., Malmstrom, R. R.,

Stieglmeier, M., Klingl, A., Woyke, T., Ryan, M. C., & Banfield, J. F. (2018).

Differential depth distribution of microbial function and putative symbionts

through sediment-hosted aquifers in the deep terrestrial subsurface. *Nature*

*Microbiology*, *5*, 328–336. https://doi.org/10.1038/s41564-017-0098-y

Rawat, M., & Maupin-Furlow, J. A. (2020). Redox and thiols in archaea. In

*Antioxidants* (Vol. 9, Issue 5). https://doi.org/10.3390/antiox9050381

Schulz, F., Eloe-Fadrosh, E. A., Bowers, R. M., Jarett, J., Nielsen, T., Ivanova, N. N.,

Kyrpides, N. C., & Woyke, T. (2017). Towards a balanced view of the bacterial

tree of life. *Microbiome*, *5*(1). https://doi.org/10.1186/s40168-017-0360-9

Sharrar, A. M., Crits-Christoph, A., Méheust, R., Diamond, S., Starr, E. P., &

934 Banfield, J. F. (2020). Bacterial secondary metabolite biosynthetic potential in

935 soil varies with phylum, depth, and vegetation type. *MBio*, *11*(3).

936 https://doi.org/10.1128/MBIO.00416-20

937 Shen, B. A., & Landick, R. (2019). Transcription of Bacterial Chromatin. In *Journal of*

938 *Molecular Biology* (Vol. 431, Issue 20). https://doi.org/10.1016/j.jmb.2019.05.041

939 Shen, P., & Huang, H. V. (1986). Homologous recombination in Escherichia coli:

940 dependence on substrate length and homology. *Genetics*, *112*(3).

941 https://doi.org/10.1093/genetics/112.3.441

942 Shin, J., Song, Y., Jeong, Y., & Cho, B. K. (2016). Analysis of the core genome and

943 pan-genome of autotrophic acetogenic bacteria. In *Frontiers in Microbiology*

944 (Vol. 7, Issue SEP). https://doi.org/10.3389/fmicb.2016.01531

945 Song, W., Wemheuer, B., Zhang, S., Steensen, K., & Thomas, T. (2019). MetaCHIP:

946 Community-level horizontal gene transfer identification through the combination

947 of best-match and phylogenetic approaches. *Microbiome*, *7*(1).

948 https://doi.org/10.1186/s40168-019-0649-y

949 Sonnhammer, E. L. L., Eddy, S. R., Birney, E., Bateman, A., & Durbin, R. (1998).

950 Pfam: Multiple sequence alignments and HMM-profiles of protein domains.

951 *Nucleic Acids Research*, *26*(1). https://doi.org/10.1093/nar/26.1.320

952 Soucy, S. M., Huang, J., & Gogarten, J. P. (2015). Horizontal gene transfer: Building

953 the web of life. In *Nature Reviews Genetics* (Vol. 16, Issue 8).

954 https://doi.org/10.1038/nrg3962

955 Stern, A., & Sorek, R. (2011). The phage-host arms race: Shaping the evolution of

956 microbes. In *BioEssays* (Vol. 33, Issue 1).

957 https://doi.org/10.1002/bies.201000071

958 Suzek, B. E., Wang, Y., Huang, H., McGarvey, P. B., Wu, C. H., & Consortium, U.

(2015). UniRef clusters: a comprehensive and scalable alternative for improving

sequence similarity searches. *Bioinformatics (Oxford, England)*, *31*(6), 926–932.

https://doi.org/10.1093/bioinformatics/btu739

Swan, B. K., Tupper, B., Sczyrba, A., Lauro, F. M., Martinez-Garcia, M., Gonźalez, J.

M., Luo, H., Wright, J. J., Landry, Z. C., Hanson, N. W., Thompson, B. P.,

Poulton, N. J., Schwientek, P., Acinas, S. G., Giovannoni, S. J., Moran, M. A.,

Hallam, S. J., Cavicchioli, R., Woyke, T., & Stepanauskas, R. (2013). Prevalent

genome streamlining and latitudinal divergence of planktonic bacteria in the

surface ocean. *Proceedings of the National Academy of Sciences of the United*

*States of America*, *110*(28). https://doi.org/10.1073/pnas.1304246110

Tang, L., Nogales, E., & Ciferri, C. (2010). Structure and function of SWI/SNF

chromatin remodeling complexes and mechanistic implications for transcription.

In *Progress in Biophysics and Molecular Biology* (Vol. 102, Issues 2–3).

https://doi.org/10.1016/j.pbiomolbio.2010.05.001

Tee, H. S., Waite, D., Payne, L., Middleditch, M., Wood, S., & Handley, K. M. (2020).

Tools for successful proliferation: diverse strategies of nutrient acquisition by a

benthic cyanobacterium. *ISME Journal*, *14*(8), 2164–2178.

https://doi.org/10.1038/s41396-020-0676-5

The UniProt Consortium. (2017). UniProt: the universal protein knowledgebase.

*Nucleic Acids Research*, *45*(D1), D158–D169.

https://doi.org/10.1093/nar/gkw1099

Touchon, M., Bernheim, A., & Rocha, E. P. C. (2016). Genetic and life-history traits

associated with the distribution of prophages in bacteria. *ISME Journal*, *10*(11).

https://doi.org/10.1038/ismej.2016.47

Treangen, T. J., & Rocha, E. P. C. (2011). Horizontal transfer, not duplication, drives

984 the expansion of protein families in prokaryotes. *PLoS Genetics*, *7*(1).

985 https://doi.org/10.1371/journal.pgen.1001284

986 Vasu, K., & Nagaraja, V. (2013). Diverse Functions of Restriction-Modification

987 Systems in Addition to Cellular Defense. *Microbiology and Molecular Biology*

988 *Reviews*, *77*(1). https://doi.org/10.1128/mmbr.00044-12

989 Voeller, D. M., Changchien, L. ming, Maley, G. F., Maley, F., Takechi, T., Turner, R.

990 E., Montfort, W. R., Allegra, C. J., & Chu, E. (1995). Characterization of a

991 specific interaction between Escherichia coli thymidylate synthase and

992 escherichia coli thymidylate synthase mRNA. *Nucleic Acids Research*, *23*(5).

993 https://doi.org/10.1093/nar/23.5.869

994 Weigert, M., & Kümmerli, R. (2017). The physical boundaries of public goods

995 cooperation between surface-attached bacterial cells. *Proceedings of the Royal*

996 *Society B: Biological Sciences*, *284*(1858).

997 https://doi.org/10.1098/rspb.2017.0631

998