Circular RNAs arising from synaptic host genes are modulated by SFPQ RNA-binding protein and increased during human neuronal differentiation

Michelle Watts[1,2], Marika Oksanen[1,2,3], Sanna Lejerkrans[1,2], Francesca Mastropasqua[1,2], Myriam Gorospe[3] and Kristiina Tammimies[1,2,*]

[1]Center of Neurodevelopmental Disorders (KIND), Centre for Psychiatry Research, Department of Women's and Children's Health, Karolinska Institutet and Stockholm Health Care Services, Region Stockholm, Stockholm, Sweden.

[2]Astrid Lindgren Children's Hospital, Karolinska University Hospital, Region Stockholm, Stockholm, Sweden.

[3]Laboratory of Genetics and Genomics, National Institute on Aging Intramural Research Program, NIH, Baltimore, MD, USA.

[*]Correspondence:

Kristiina Tammimies, PhD

E-mail: kristiina.tammimies@ki.se

Postal address: BioClinicum J9:30, Visionsgatan 4, 171 56 Solna, Sweden

Abstract

Background

Circular RNAs (circRNAs) are emerging as a key component of the complex neural transcriptome

implicated in brain development. However, the specific expression patterns and functions of circRNAs

in human neuronal differentiation have not been explored.

Results

Using total RNA sequencing analysis, we identified expressed circRNAs during the differentiation of

human neuroepithelial stem (NES) cells into developing neurons and discovered that many circRNAs

originated from host genes associated with synaptic function. Interestingly, when assessing

population data, exons giving rise to circRNAs in our dataset had a higher frequency of genetic

variants. Additionally, screening for RNA-binding protein sites identified enrichment of Splicing Factor

Proline and Glutamine Rich (SFPQ) motifs in increased circRNAs, several of which were reduced by

SFPQ knockdown and enriched in SFPQ ribonucleoprotein complexes.

Conclusions

Our study provides an in-depth characterisation of circRNAs in a human neuronal differentiation

model and highlights SFPQ as both a regulator and binding partner of circRNAs elevated during

neuronal maturation.

Keywords

Circular RNA; SFPQ; Neuronal differentiation; Genetic variation

Background

CircRNAs represent a unique class of RNA molecules formed following non-canonical back-splicing

of exon-exon junctions (Cocquerelle *et al*, 1992; Nigro *et al*, 1991). These covalently closed,

circularised RNAs lack 5'cap and 3'poly-A structures resulting in highly stable molecules, resistant to enzymatic degradation (Jeck *et al*, 2013; Memczak *et al*, 2013). Since their initial discovery, the circRNA transcriptome has expanded rapidly with the advent of sequencing approaches and bioinformatic tools designed to detect back-spliced junctions in RNA-sequencing (RNAseq) data (Memczak *et al.*, 2013; Salzman *et al*, 2013; Salzman *et al*, 2012). Although once thought to be non-functional by-products of spurious splicing events, numerous functional roles of circRNAs have now been identified, including microRNA (miRNA) sponging (Memczak *et al.*, 2013), transcriptional regulation (Conn *et al*, 2017; Pandey *et al*, 2020), RNA-binding protein function (Panda *et al*, 2017; Tsitsipatis *et al*, 2021), and modulation of protein interactions (Barbagallo *et al*, 2019; Du *et al*, 2016). These roles highlight the vast diversity of possible functions for circRNAs, most of which are unknown.

As is often observed for mechanisms such as microRNAs or splicing factors that diversify the mammalian transcriptome, circRNA species are highly enriched in the nervous system (Memczak *et al.*, 2013; Rybak-Wolf *et al*, 2015; You *et al*, 2015). Spatio-temporal regulation of circRNA expression within the brain is also dynamic, differing across brain regions and developmental stages (Chen *et al*, 2019; Veno *et al*, 2015; You *et al.*, 2015). Brain and neuronal expressed circRNAs have also been found to arise from back-splicing of linear host genes associated with synaptic functions and display specific enrichment within synaptosomes as well as regulation in response to neuronal activation (Gokool *et al*, 2020; You *et al.*, 2015). Under various circumstances, this regulation appears independent of corresponding host genes and, in some cases, circRNA levels far exceed linear host RNA levels (Rybak-Wolf *et al.*, 2015; You *et al.*, 2015), underscoring the importance of precise modulation of circRNAs during neuronal development.

Accumulating evidence suggests that circRNAs are not only important for neuronal function but also for brain development. Differential expression of circRNAs has been found to occur during neuronal differentiation, both in mice and in cell culture models such as the human neuroblastoma cell line SH-SY5Y, with several specific circRNAs were found to have a regulatory role in this process (Hollensen *et al*, 2020; Li *et al*, 2022b; Suenkel *et al*, 2020; Yang *et al*, 2019; Yang *et al*, 2018). Additionally, a number of circRNAs have been identified as being dysregulated in Autism Spectrum

Disorder (ASD) (Chen *et al*, 2020; Gokool *et al.*, 2020), while a depletion of circRNAs has been

observed in individuals with Schizophrenia (SCZ) (Mahmoudi *et al*, 2019). Currently, however, our

understanding of circRNA regulation and function in human neurodevelopment remains largely

unexplored.

While evaluation of circRNAs in the human brain is limited to post-mortem analyses, derivation

of mature neurons from human induced pluripotent stem cells (hiPSCs) allows for modelling of

neuronal differentiation in a human genetic background (Dolmetsch & Geschwind, 2011).

Computational methods to detect back-spliced junctions in total RNAseq data also allow for the

discovery of circRNAs without the need for specialised treatment of RNA samples as well as

comparison with linear RNA expression from the same data (Hansen, 2018). We combined these

methodologies to examine the circRNA landscape of hiPSC-derived neuroepithelial stem (NES) cells

and developing neurons, revealing an increase in circRNAs during differentiation, arising from host

genes with synaptic functions. CircRNA-forming exons were also found to harbour genetic variants at

a higher rate in comparison to exons from the same host gene that were not circularized. In addition,

we investigated the potential functionality of elevated circRNAs and identified an enrichment of

binding motifs for Splicing Factor Proline and Glutamine Rich (SFPQ). Predicted circRNA targets of

SFPQ were found to be depleted during neuronal differentiation following SFPQ knockdown including

a novel circRNA from the Regulating Synaptic Membrane Exocytosis 1 (RIMS1) host gene that was

specifically enriched by SFPQ immunoprecipitation above its corresponding linear isoform.

Results

Detection of robust circRNA production during NES differentiation from total RNAseq data

To analyse circRNAs in human neuronal differentiation, hiPSC-derived NES cells were differentiated

as previously described (Becker *et al*, 2020). Total RNA was collected and sequenced from cells in

their proliferative, neural progenitor state (NES) and two time points of differentiation. First, at day five

(D5), when a neural identity has been adopted and neurite outgrowth/axon specification is underway,

and second at day 28 (D28), when extensive neural networks, but not yet mature active connections, have been established **[Fig 1a]**. To reduce spurious circRNA discovery, three detection programs (CIRCexplorer2, CIRI, MapSplice) with reported low false positive rates were utilised (Hansen, 2018). A total of 60,531 detected circRNAs were filtered for minimum expression, leaving 6,540 circRNAs (~10.8%) for downstream analysis. These were further filtered to include only circRNAs detected by at least two programs, which represented the majority of the expression filtered dataset (6,385; ~97.6%) **[Fig 1b]**. Among these, 66% were known circRNAs reported in circBase (Glazar *et al*, 2014), 34% were novel circRNAs (not yet reported in circBase), and 8.5% of the dataset were '*de novo*' circRNAs that did not annotate to known splice sites. The entire filtered dataset of 6,385 circRNAs was derived from 2,743 unique host genes, with 1,377 genes producing two or more circRNA isoforms and 50 'hotspot' genes that generated over ten circRNAs **[Additional File 2: Table S1a]**. All circRNAs analysed and reported in Additional File 2: Table S1 have been assigned an ID based on host gene name and back-spliced junction genomic locations. Where available, corresponding circBase IDs are provided. For simplicity, specific circRNAs referenced within this text are referred to only by host gene name with alphabetical indexing for referencing circRNA isoforms.

Differentially expressed circRNAs (DEcircs) in NES differentiation were detected using DESeq2 with log fold-change (LFC) shrinkage estimation for increased stringency. We observed a predominant increase of circRNA expression with pairwise comparisons finding 33 circRNAs up and 8 circRNAs down at D5 compared to NES and 660 circRNAs up and 95 down at D28 compared to NES (p-adjusted <0.05, LFC ≥|2|) **[Fig 1c; Additional File 2: Table S1b-d]**. Almost all circRNAs (32/33) whose expression was increased at the early differentiation time point (D5) remained significantly increased at D28 **[Fig 1d]**. Clustering results using the likelihood ratio test, which incorporates all time points, similarly highlighted a primary cluster of circRNAs with a pattern of increasing abundance throughout differentiation **[Fig 1e; Additional File 2: Table S1e,f]**.

To validate whether detected circRNAs occurred endogenously and are differentially expressed during differentiation, we selected 20 DEcircs and designed divergent primer pairs to selectively amplify back-spliced junctions **[Additional File 1: Fig S1a]**. PCR amplification and gel

electrophoresis found a clear, predominant band of expected size for 19/20 primer pairs in samples that were depleted of linear RNA by RNAse R treatment **[Fig 1f]**. No amplification was observed from primers designed for *de novo* circRNA *circZNF1* and primers for *circRERE_a* and *circRIMS1_b* produced bands in addition to the expected product **[Fig 1f]**. Sanger sequencing confirmed that for *circRERE_a* primers, these corresponded to either PCR artifact or to the *circRERE_b* isoform, while primers for *circRIMS1_b* produced two additional *circRIMS1* isoforms that contained either one or two additional exons not annotated in NCBI **[Additional File 1: Fig S1b,c]**. These unannotated exons (UEs) mapped to the RIMS1 intronic region between exons 26 and 27 (UE-1 = 28-b; GRCh37-chr6:73005640-73005667, UE-2 = 90-b; GRCh37-chr6:73008851-73008880). Using quantitative PCR (qPCR) analysis of replicate samples, similar differential expression of circRNAs was also observed at D28 when compared to NES, particularly for increased circRNAs **[Additional File 1: Fig S1d]** and qPCR LFC values strongly correlated with RNAseq LFC values (R = 0.8, p = 1.2e-4). Finally, secondary divergent primer pairs were designed to confirm circRNA annotation and determine internal exon structure in circRNAs with non-consecutive back-spliced exons **[Additional File 1: Fig S2a]**. Among primer pairs which amplified enough product, all except *circRIMS1_a*, produced a single band. Sanger sequencing confirmed back-spliced exon identity in all circRNAs as well as internal exon retention, where relevant **[Fig 1g and Additional File 1: Fig S2b-m]**. A second larger product of the primer pair for *circRIMS1_a* (*circRIMS1_a2*) was found to also contain the 90-b UE-2 described above. To determine if these UEs are expressed in linear RIMS1 RNA, we designed primers flanking this region and performed PCR amplification on D28 cDNA. A single product was detected that we confirmed by Sanger sequencing excluded both UEs, indicating these exons may be circRNA specific **[Additional File 1: Fig S2n]**. Finally, as an additional validation, the dataset was compared with circRNAs detected by Gokool *et al*, which represents the largest dataset of human brain circRNAs (197 post-mortem samples), and ~80% of circRNAs in our study were found to be represented. Taken together, these results indicate that our pipeline for circRNA detection from total RNAseq data was successful in identifying genuine circRNAs that are differentially expressed during neuronal differentiation.

CircRNAs increased during NES differentiation arise from synaptic host genes

To gain insight into the host genes from which circRNAs are produced during differentiation, functional

annotations associated with circRNA host genes were examined. As NES differentiation primarily

increased circRNA expression, we initially tested host gene ontology enrichment among Cluster 1

circRNAs as well as positive gene set enrichment analysis (GSEA) of D28 vs NES DEcircs. This

highlighted significantly enriched terms predominantly related to synaptic functions including synapse

assembly, synaptic vesicle cycle and ion transport in addition to other neuronal development

processes such as neuronal projection and cAMP signalling **[Fig 2a,b and Additional File 3: Table**

S2a,b]. While similar pathways were identified as positively enriched from GSEA of D5 vs NES, only

one reached significance (GO:0051962-Positive regulation of nervous system development*;* p-

adjusted <0.05). Of interest, GSEA of D28 vs D5 found only cell-cell adhesion as an enriched process,

while multiple processes related to early neuronal specification, such as cell projection and

axonogenesis, were negatively enriched **[Additional File 1: Fig S3a and Additional File 3: Table**

S2c]. GSEA of host genes of circRNAs decreased at D28 vs NES did not identify any enriched

pathways. However, among Cluster 2 circRNA hosts, terms associated with morphogenesis and

development were overrepresented **[Additional File 1: Fig S3b and Additional File 3: Table S2d]**.

Additionally, we examined the enrichment of neurodevelopmental disorder (NDD) risk genes since

circRNA expression has previously been found to be disrupted in individuals with ASD and SCZ. The

tested genes included ASD genes from the Simons Foundation Autism Research Initiative (SFARI)

database, collated SCZ risk genes **[Additional File 4: Table S3]**, and developmental disorder-related

genes covering broader phenotypes (Li *et al*, 2022a). We found significant enrichment of ASD genes

among hosts genes of circRNAs increased at D28 vs NES, and significant enrichment of SCZ risk

genes in Cluster 4 circRNA hosts **[Additional File 1: Fig S3c,d]**.

Employing the same RNAseq dataset, we performed differential expression analysis of linear

RNAs to identify regulatory patterns and gene ontology terms. The expression of linear RNAs was

found to follow a similar regulatory pattern to that of circRNAs, with a higher proportion of linear RNAs

increased at D28 vs NES (2,904 genes compared to 1,363 decreased; p-adjusted <0.05, LFC ≥|2|)

[Additional File 1: Fig S4a,b and Additional File 5: Table S4a]. GSEA also revealed enrichment of

similar pathways related to synaptic function among linear RNAs increased by differentiation

[Additional File 1: Fig S4c and Additional File 5: Table S4b]. Since multiple studies have found

discordances between the changes in circRNA levels and the changes in the levels of linear RNAs

from the same host gene, we compared changes in circRNA and linear counterparts. As shown, a

moderate but significant positive correlation was detected (R = 0.5, p <2.2e-16). Plotting of LFC

correlates highlighted this same positive linear association; however, there was an evident shift from

the linear distribution suggesting that highly increased circRNAs had higher fold-changes than their

linear counterparts [**Fig 2c**]. A direct comparison of circRNAs to linear RNA counterparts found this

same trend among the top 20 circRNAs with an average increase of circRNA expression of ~6.7 fold

higher than linear RNAs **[Additional File 1: Fig S5]**. Additionally, among circRNAs increased at D28

vs NES, a large majority (71.6%) were over two-fold higher than their linear counterparts from the

same host genes, supporting a non-linear relationship between the expression of these two RNA

forms.

To further explore the relationship between linear RNA and circRNA regulation, we analysed

differential exon usage in our dataset using DEXseq. All exon bins detected by DEXseq were analysed

as exons, including those corresponding to circRNA-forming exons. Comparing cells differentiated up

to D28 with NES cells we identified a large number of differentially used exons (19,502 exons; p-

adjusted <0.05, lfc ≥|2|). Among these, the number of exons with elevated expression was over three

times higher than exons with reduced expression (4,680 down:14,822 up), implying substantially

increased exon inclusion at D28. This corresponds with alternative splicing of 4,904 RNAs with an

average of ~31 exons. When enrichment of gene ontology terms among alternatively spliced RNAs

were examined, pathways related to the ribosome and spliceosome were most significantly enriched

[Additional File 1: Fig S4d].

To establish if alternative splicing is associated with circRNA biogenesis and/or increased

abundance of circRNAs at D28, the rate of alternate splicing (based on differential exon usage at D28

vs. NES) was compared among three groups: all linear RNAs, linear RNAs sharing host RNAs with

circRNAs, and linear RNAs sharing host RNAs with circRNAs increased at D28. While alternative

splicing was ~3-fold higher among RNAs which host circRNAs (p < 0.0001) compared to all RNAs,

there was no difference in alternate splicing rate between all host RNAs and hosts of circRNAs

increased at D28 (p = 0.153) **[Fig 2d]**. Similarly, we found no correlation between circRNA fold-

changes and estimated exon fold-changes (R = -0.018), suggesting that increased abundance of

circRNAs during NES differentiation is independent of general splicing of host precursor RNAs.

Variant frequency is increased in circRNA-forming exons

To further assess the role of circRNAs in human health and disorders, we examined genetic variant

burden in circRNA-forming exons. Exons were classified into circ and nonCirc groups based on all

exons of the longest annotated transcript containing the back-spliced exon(s). CircRNA exons were

further differentiated as exons that either form BSJs or do not form BSJs (nonBSJ exons) **[Fig 3a]**.

Genetic variants included in the Genome Aggregation Database (gnomAD) were initially utilised to

examine the frequency of circRNA exon variation in the general population. Rates of genetic variation

were found to be slightly higher in both circ (p = 2.2e-16) and BSJ (p = 1.4e-5) exons when compared

with nonCirc and nonBSJ exons, respectively **[Fig 3b]**. However, no shift was observed in the

annotation of variant impact among exon groups **[Fig 3c]** or in the proportion of missense variants or

the proportion of missense variants identified as deleterious **[Additional File 1: Fig S6a,b]**. When we

limited the analysis to splice variants, a higher rate among circRNA-forming exons in comparison to

nonCirc exons was detected (p < 2.2e-16). Furthermore, among circRNA-forming exons, the rate of

splice variants was significantly lower in BSJ exons compared with nonBSJ exons (p = 1.1e-4) **[Fig**

3d]. These observations suggest circRNA-forming and BSJ exons are subject to increased variation

while splicing variants are less frequent among back-spliced exons.

210   Next, we examined the frequency of rare *de novo* variants found in ASD from a large exome

211   sequencing study from 21,219 family-based samples, including 12,166 variants in 6,430 individuals

212   with ASD (Satterstrom *et al*, 2020). We identified similar results as for gnomAD variants in that *de*

novo ASD variants occurred at a slightly higher rate in circ and BSJ exons **[Fig 3e]**. This increase in circ exon variants was significant specifically when examining circRNAs formed from ASD-associated SFARI host genes (p = 0.0384) **[Fig 3f]**. We note however that as genetic variation overall was increased in circularised exons, that this may not be indicative of specific enrichment of ASD-associated variants in circRNAs. Similarly, a trend for increased rare *de novo* variants in BSJ above non-BSJ exons was also detected (p = 0.06017) among 2,179 unaffected sibling controls from the same cohort **[Additional File 1: Fig S6c,d]**.

Due to a lack of a comparable large resource for SCZ, we collated SCZ variants reported from several sources **[Additional File 1: Table S5]**. In total, 2,035 rare *de novo* variants identified in individuals diagnosed with either Schizophrenia or schizoaffective disorder were included. When comparing the frequency of SCZ variants, we found no differences across exon groups either among all transcripts or when comparing variants specifically in SCZ risk genes **[Additional File 1: Fig S6e,f]**.

Binding site enrichment in circRNAs differentially expressed during NES differentiation

Given that circRNA functionality remains largely unknown, differentially expressed circRNAs were screened *en masse* for potential function by searching for binding sites among circRNA sequences. The discovery that the highly abundant circRNA, *CDR1as,* functions in sequestering miR-7 through numerous seed sites (Memczak *et al.*, 2013; Piwecka *et al*, 2017) has since led to frequent reporting of predicted circRNA:miRNA:mRNA interactions (Chen *et al.*, 2020; Dell'Orco *et al*, 2021; Liu *et al*, 2021; Ma *et al*, 2019). Therefore, we first investigated miRNA sponging by exploring possible interactions between miRNAs and DEcircs. We limited our search to 125 human miRNAs previously associated with neurodevelopment or NDDs, which were most of interest to our study **[Additional File 6: Table S6a,b]**. To construct putative circRNA:miRNA:mRNA interaction axes, binding sites for these miRNAs were predicted in both DEcircs as well as 3'UTRs of differentially expressed RNAs **[Additional File 6: Table S6c,d]**. Based on canonical miRNA regulation where miRNAs repress gene expression and circRNA sponges alleviate target repression, circRNAs with predicted miRNA binding

sites were matched to mRNAs containing binding sites for the same miRNA and with the same pattern

of regulation. We found no circRNAs to have a high number of binding sites for any particular miRNA

with a maximum of three sites identified in any single circRNA. This is consistent with other reports

that enrichment of miR-7 sites in CDR1as is a unique phenomenon among circRNAs (Guo *et al*, 2014;

You *et al.*, 2015). We report identified circRNA:miRNA:mRNA interactions in **Additional File 6: Table**

S6e, however, we do not expect that these circRNAs function in miR-sponging.

Next, we examined the enrichment of RNA-binding protein (RBP) motifs among DEcirc

sequences. A number of motifs were found to be enriched among different subsets of DEcircs,

including two motifs enriched among increased circRNAs **[Additional File 7: Table S7a]**. These two

motifs were associated with two splicing factor RBPs, namely, TIA1 Cytotoxic Granule Associated

RNA Binding Protein Like 1 (TIAL1) and Splicing Factor Proline and Glutamine Rich (SFPQ) **[Fig**

4a,b]. These results highlighted a possible functional link between TIAL1 and SFPQ RBPs and the

increased abundance of circRNAs in neuronal differentiation.

SFPQ regulates expression of circRNAs in NES differentiation

To further test the relationship between increased circRNAs and enriched RBP motifs, expression of

SFPQ and TIAL1 mRNA in NES differentiation was first established. Normalised counts from RNAseq

analysis indicated that both RBPs were expressed at all time points albeit with SFPQ levels around

tenfold higher than TIAL1 **[Fig 4c,d]**. We also noted a significant reduction in expression of both SFPQ

and TIAL1 at D28, although the log-fold reduction was low, this pattern is reflective of SFPQ

expression in mice, which is highest during early neurodevelopmental stages and later diminishes

(Takeuchi *et al*, 2018). From motif analysis of circRNA sequences, a number of 'top' putative targets

for SFPQ and TIAL1 were subsequently chosen based on motif frequency, base mean expression

and LFC values **[Additional File 7: Table S7b,c]**. Elevated expression of all selected circRNA targets

during NES differentiation was confirmed using divergent primer qPCR with all targets increased ~4

fold or greater as early as day eight of differentiation (D8) **[Additional File 1: Fig S7a]**. To determine

if these RBPs affected target circRNA expression, we knocked down SFPQ and TIAL1 during NES

differentiation with cells treated with siRNA on days zero (NES) and eight (D8) and RNA collected on

D8 and D16. For quantification of protein levels, cell lysates were also collected four days after each

siRNA treatment **[Additional File 1: Fig S7b]**. Quantification of mRNA at D8 and D16 found *SFPQ*

to be robustly reduced by siRNA treatment compared to non-targeting control siRNA (siNTC) **[Fig**

4e]. SFPQ protein levels were also reduced ~50% at D4, however, protein expression was returned

to baseline levels at D12 indicating that only transient knockdown of SFPQ was achieved **[Additional**

File 1: Fig S7c,d]. TIAL1 protein expression was too low to be detected in any samples and *TIAL1*

mRNA levels also appeared unaffected by siRNA treatment **[Fig 4e and Additional File 1: Fig**

S7c,d]. TIAL1 and its predicted targets were therefore excluded from further analyses.

Following siRNA-SFPQ treatment, we quantified the relative expression of select circRNA

targets alongside their corresponding linear mRNA form. Knockdown of SFPQ resulted in depletion

of all circRNA targets at both D8 and D16 with an apparent greater decrease of circRNAs compared

to linear counterparts, an effect that was especially pronounced at D16 of differentiation **[Fig 4f]**. This

indicates that SFPQ affects expression of both circRNAs and their linear counterparts, but we expect

that changes in circRNA expression are exacerbated by their enhanced stability (Jeck *et al.*, 2013;

Memczak *et al.*, 2013), even beyond D8 when SFPQ protein was no longer suppressed. To further

explore this relationship between SFPQ and circRNA expression we examined SFPQ expression by

qPCR at additional timepoints of differentiation **[Additional File 1: Fig S8a]**. Consistent with RNAseq

data, we found SFPQ expression reduced at later timepoints of differentiation when circRNA

expression is highest. Although this appears counterintuitive for a regulator of circRNA formation,

when we further profiled selected SFPQ circRNA targets throughout differentiation up to D50 we found

circRNA abundance increases linearly during early differentiation (D8) and plateaus during later

stages **[Additional File 1: Fig S8b-f].** Next, as a qualitative analysis, two circRNAs that exhibited the

greatest decrease following SFPQ knockdown, *circERC2* and *circMYT1L* were selected for *in situ*

hybridisation. circRNA probes were designed to hybridise to the back-spliced junction along with a

control probe designed to a linear splice junction of one back-spliced exon. No specific signal was

seen with a negative scrambledCirc *in situ* probe or for *circMYT1L*, likely due to low expression as

linear MYT1L was only detected at low levels **[Additional File 1: Fig S9a,b]**. However, *circERC2* was

detected with a clear increase in expression observed following differentiation **[Fig 4g-j]**. *In situ*

detection of *circERC2* showed outspread co-localised with linear ERC2 and expression throughout

the neuronal soma **[Fig 4g-j]**. This localisation pattern also appeared unchanged by siRNA-SFPQ

treatment **[Additional File 1: Fig S9c-f].** Finally, to determine if SFPQ can bind to the predicted

circRNA targets, RNA immunoprecipitation followed by qPCR analysis (RIP-qPCR) was performed

on neurons at D28 of differentiation. We found that SFPQ immunoprecipitation enriched all circRNA

targets greater than four-fold and, in the case of *circRIMS1_d*, specific enrichment of the circRNA

exceeded that of the linear isoform **[Fig 4k]**. Collectively, this suggests a role for SFPQ not only in

the regulation of circRNAs during neuronal differentiation but also as a potential binding partner of

increased circRNAs.

CircRNAs increased during differentiation are flanked by long introns

While the SFPQ targets examined were selected based on enrichment for a singular SFPQ motif,

additional motifs for SFPQ are known and could be involved in the effects of SFPQ knockdown.

Additionally, SFPQ has previously been shown to have an important role in circRNA biogenesis

through interactions with circRNA flanking introns, specifically regulating circRNAs with long flanking

introns that contain distal inverted Alu repeat elements (IAE) (Stagsted *et al*, 2021). The observed

decrease of circRNA targets following SFPQ knockdown could be due to a more general effect of

SFPQ promoting circRNA formation through flanking intronic regions. We therefore tested flanking

introns of increased circRNAs for enrichment of RBP motifs. Although four motifs for SFPQ were

found to be enriched in flanking introns, this represented a small minority among a total of 349

enriched motifs from 76 RBPs, the majority of which were motifs corresponding to members of the

HNRNP (93) and SRSF (91) protein families **[Additional File 7: Table S7d]**. Previously, SFPQ

binding sites have been identified as enriched both in close proximity (+/- 2000 bp) to BSJ sites of

circRNAs (Stagsted *et al.*, 2021) as well at 5' ends of long introns from SFPQ-bound genes (Takeuchi

et al., 2018). The frequency of SFPQ motifs in flanking introns of circRNA subsets was consequently

examined to determine if any position-based pattern was evident. We observed that the frequency of

SFPQ motifs among flanking introns of DEcircs is highest at intron ends distal to the BSJ and lowest

in regions closest to the BSJ **[Fig 5a]**. Analysing motif position relative to the 5' site of both upstream

and downstream flanking introns indicated that this pattern was driven by an enrichment of SFPQ

binding sites at 5' intronic sites **[Fig 5b]**. While this indicates that 5' SFPQ motifs may be important

for biogenesis of circRNAs differentially expressed in this dataset, this pattern was not unique to

increased circRNAs and, in fact, circRNAs decreased during differentiation appear to have the highest

frequency of 5' SFPQ motifs in their downstream flanking introns.

We further compared both flanking intron length and the location of inverted Alu repeats in all

circRNAs and circRNAs increased at D28 as potential features of SFPQ-regulated circRNAs.

Contrasts were made only between introns flanking circRNAs as, consistent with prior studies (Ivanov

et al, 2015; Jeck *et al.*, 2013), we found flanking introns were significantly longer overall when

compared to all introns in circRNA host genes **[Additional File 1: Fig S10a-d]**. Based on grouping of

flanking introns by length and presence of proximal IAEs, the proportion of intronic types was

unchanged in the subset of increased circRNAs **[Fig 5c]**. A comparison of mean length, however,

found that flanking introns were significantly longer in increased circRNAs (p = 2.6e-11) whereas

distance to Alu elements in circRNAs that contained IAEs did not differ significantly (p = 0.1615) **[Fig**

5d-g]. When this analysis was repeated on decreased circRNAs we found both mean intron length

and Alu to be unchanged **[Additional File 1: Fig S10e-h]**. These results suggest that intronic length,

but not IAE distance, may be an important factor in the increased abundance of circRNAs during NES

differentiation.

Discussion

Despite mounting evidence highlighting an important role for circRNA regulation in normal brain

function, our understanding of circRNA generation and function during neurodevelopment remains

poorly understood. Our study provides a comprehensive profile of circRNA expression in a human

model of neuronal differentiation. Although previous reports have suggested high conservation of

circRNAs, more recent evidence indicates that circRNA exons exhibit similar conservation to

neighbouring linear exons and that the majority of circRNAs are species-specific with only ~20% of

mouse circRNAs having human homologs (Gruhl *et al*, 2021; Guo *et al.*, 2014; Santos-Rodriguez *et*

al, 2021). Exploring circRNA expression and function within a human-genetic context is therefore

critical to understanding the role of circRNAs in both disease and normal brain development.

Recent expansions of the human circRNAome also coincide with the creation of various

bioinformatic tools for the detection of circRNAs from sequencing of either total RNA or RNA enriched

for circRNA by RNase treatment (Hansen, 2018; Sharma *et al*, 2019). Sequencing of circRNA

enriched samples offers improvements in detecting lowly expressed circRNAs and the advantage of

reconstructing complete circRNA sequences (Rybak-Wolf *et al.*, 2015). By contrast, lowly expressed

circRNAs are unlikely to be captured in total RNAseq data, and detection of circRNAs, which relies

only on BSJ reads, is prone to false positives. However, the pipeline employed in our study,

demonstrates that the identity and expression levels of endogenous circRNAs can be reliably

determined from total RNAseq by intersecting results of multiple tools, at least for abundant circRNA

species that are most likely to be biologically relevant.

An overall pattern of increased circRNA expression following neuronal differentiation has

previously been observed in studies of rodent neuronal development, with elevation of circRNAs

occurring following neuronal maturation *in vitro* and within the developing mouse hippocampus

(Rybak-Wolf *et al.*, 2015; You *et al.*, 2015). Enrichment of GO terms related to neuronal and synaptic

functions among hosts of increased circRNAs in our dataset is also consistent with several reports

highlighting neuronal genes, and especially synaptic factors, as hosts for numerous circRNAs

(Ashwal-Fluss *et al*, 2014; You *et al.*, 2015). Identification of negatively enriched GO terms related to

early, but not late differentiation (cell projection/axonogenesis) at D28 vs D5, however, indicates that

circRNAs with altered expression arise from host RNAs functioning in biological pathways relevant to

the stage of neuronal maturation. We also observed greater relative increases in the levels of

circRNAs than in the levels of linear counterparts, supporting prior evidence of specific circRNA

regulation during neuronal development. Non-specific accumulation of circRNAs, however, has been

reported in non-proliferating cells and with aging in *C. elegans* (Bachmayr-Heyda *et al*, 2015; Cortes-

Lopez *et al*, 2018) while brain-specific circRNAs accumulate during aging in more complex organisms

like *Drosophila* and mouse (Gruner *et al*, 2016; Westholm *et al*, 2014). Dissecting the mechanisms of

both specific circRNA regulation as well as circRNA accumulation will therefore be crucial for

understanding circRNA functions in the developing and aging brain.

Although mechanisms of circRNA biogenesis are still poorly understood, BSJ formation has

been shown to occur co-transcriptionally and appears closely linked to linear mRNA splicing (Ashwal-

Fluss *et al.*, 2014). Differential exon usage has been reported to be in competition with exon

circularisation (Ashwal-Fluss *et al.*, 2014), whereas other studies identified circRNAs formed primarily

from exons with high inclusion rates (Gokool *et al.*, 2020). While we observed a significant increase

of alternative splicing among transcripts that host circRNAs, alternative splicing was comparable

among host RNAs of increased circRNAs, and no correlation was found between circRNA fold-

changes and fold-changes of circularised exons. Pathways enriched among differentially spliced

RNAs also diverged from those pathways enriched among host precursors giving rise to circRNAs,

the latter featuring processes associated with the ribosome and the spliceosome.

Since the majority of individual circRNAs remain unstudied, examining variants occurring in

circRNA exons could provide insight into their function and possible roles in evolutionary brain

development and disorders. Based on gnomAD data, circRNA exons had slightly higher variant

frequency while splice site variant frequency in BSJ exons was significantly lower than nonBSJ exons.

This increase in variation was reflected in a higher frequency of rare *de novo* ASD variants in

circularised exons of SFARI genes. A separate study also found an enrichment of ASD GWAS SNPs

among all circBase circRNAs as well as an enrichment of GWAS SNPs for four out of ten other traits

tested (Rajcsanyi *et al*, 2022). Additionally, numerous SNPs which effect circRNA abundance

(circQTLs) have been identified in humans with the majority found to be independent, linked only to

circRNA not mRNA expression (Ahmed *et al*, 2019). Further, over 600 independent circQTLs have

recently been associated with 389 circRNAs specifically in the ASD brain (Mai *et al*, 2022). Taken

together, this indicates that circRNA-forming exons in humans may be subject to increased genetic

variation and alterations of circRNA function and/or circRNA expression may be an important

consideration when determining effects of variants associated with ASD.

While there is mixed evidence of RBP binding site enrichment in circRNAs (Guo *et al.*, 2014;

You *et al.*, 2015), we identified significant enrichment of limited motifs among subsets of DEcircs.

SFPQ, found enriched among increased circRNAs, is associated with various aspects of gene

regulation and mRNA processing (Heyd & Lynch, 2010; Kaneko *et al*, 2007; Peng *et al*, 2002;

Rosonina *et al*, 2005). Though typically localised to nuclear paraspeckles (Passon *et al*, 2012) non-

nuclear SFPQ pools, linked to distal axonal transport of RNA granules, have been identified in

neuronal cells (Fukuda *et al*, 2021; Furukawa *et al*, 2015; Thomas-Jinu *et al*, 2017). Analogously,

SFPQ has been associated with multiple neurodegenerative disorders (Ishigaki *et al*, 2020; Younas

et al, 2020) and loss of SFPQ causes gross defects in mouse brain development (Takeuchi *et al.*,

2018). These effects are associated with defective transcription of long genes, many of which are

selectively expressed in the brain, enriched for synaptic functions, and associated with

neurodevelopmental disorders (Gabel *et al*, 2015; King *et al*, 2013; Takeuchi *et al.*, 2018). A recent

analysis in human cell lines also found that SFPQ knockdown primarily depleted circRNAs with long

flanking introns containing distal Alu elements, as opposed to short introns with proximal Alu elements

(Stagsted *et al.*, 2021). Although complementary sequences such as Alu repeats can facilitate

circRNA production (Jeck *et al.*, 2013; Westholm *et al.*, 2014), long flanking introns rather than flanking

proximal inverted repeats have been shown to be more important in biogenesis of the most abundant

and conserved human circRNAs (Stagsted *et al*, 2019; Westholm *et al.*, 2014).

We observed that SFPQ appears to be an important player in regulation of circRNAs and,

consistent with previously examined SFPQ targets (Stagsted *et al.*, 2021), introns flanking circRNAs

increased by NES differentiation were significantly longer than introns flanking all detected circRNAs,

while Alu element distance did not appear to be a defining characteristic. Depletion of circRNAs

following SFPQ knockdown during differentiation indicates that SFPQ is involved in circRNA

biogenesis in our model, possibly through previously described mechanisms such as 5' end enriched

SFPQ binding in long flanking introns. However, motif enrichment in increased circRNA flanking

introns indicates many other RBPs were enriched to a greater extent, and patterns of intronic SFPQ

motif enrichment appeared relatively uniform across all DEcircs. This finding indicates that, while

SFPQ is likely a critical factor in circRNA biogenesis, additional factors other than intronic SFPQ

binding must be involved in directing specific circRNA regulation. Given that SFPQ was uniquely

enriched within increased circRNA sequences, we speculate that these exonic binding sites may also

play a role in circRNA regulation. Previously, in addition to intronic sites, exonic sites were shown to

contribute to circRNA biogenesis from *musclebind like splicing regulator 1* (*MBLN1*) pre-mRNA

through binding of its own protein product, MBLN1, thereby regulating its own expression (Ashwal-

Fluss *et al.*, 2014). Here, we show that multiple circRNA targets were also enriched by SFPQ

immunoprecipitation, suggesting SFPQ binding sites in these circRNAs are occupied and could

function in sequestering SFPQ. Further research is needed to untangle the functional significance of

SFPQ binding in circRNAs elevated during differentiation either in regulating their own biogenesis,

altering host gene splicing and/or sequestering of SFPQ protein.

Conclusions

As increasing evidence points to a significant role for circRNAs in neurodevelopment, our data

provides an important resource for future studies. We evaluate alternative splicing and differential

abundance of linear RNAs and circRNAs and scrutinise genetic variation as well as RBP binding sites,

providing an in-depth characterisation of circRNAs in human neuronal differentiation. We also

discerned further correlates between circRNAs and neurodevelopmental disorders and identified the

splicing factor SFPQ as an important player in neuronal maturation, as both a regulator of circRNA

levels and possibly a target itself of circRNA regulation. We note, however, that these mechanisms

are not mutually exclusive and a more complex regulatory picture, encompassing splicing modulation,

circRNA biogenesis and SFPQ:circRNA binding, is likely yet to be resolved.

Methods

NES cell culture and differentiation

Established neuroepithelial stem (NES) cells, previously derived from iPSCs established from a male neurotypical donor, CTRL-9-II; RRID:CVCL_JL74 (Uhlin *et al*, 2017), were seeded on tissue culture flasks coated with 20 µg/ml poly-L-ornithine (Sigma-Aldrich P3655), and 1 µg/ml laminin (Sigma-Aldrich L2020). Cells were grown in DMEM/F12+GlutaMAX medium (ThermoFisher 31331093) supplemented with 0.05X B27 (ThermoFisher 17504044), 1X N2 (ThermoFisher 17502001), 10 ng/ml bFGF (fisher scientific CTP0261), 10 ng/ml EGF (PeproTech AF-100-15) and 10 U/ml penicillin/streptomycin (ThermoFisher 15140122). Medium was exchanged 50% daily and cells maintained in 5% CO_2 at 37°C, passaging once 100% confluent and seeding at a density of $5x10^4$ cells/cm^2. Neural differentiation was induced by growth factor withdrawal the day after plating with media B27 concentration increased to 0.5X. Media was exchanged 50% every second day up until D15, after which media was supplemented with 0.4 ug/ml laminin and exchanged 50% every three days.

RNA extraction and RNA sequencing

Cells were lysed in TRIzol reagent (ThermoFischer 15596026) before separating with chloroform and mixing the aqueous phase with isopropanol as per manufacturer directions. RNA was then isolated from the isopropanol/chloroform solution using the ReliaPrep RNA Cell Miniprep kit (Promega Z6010). For linear RNA depletion, RNA was incubated with 20 U RNase R (Lucigen RNR07250) for 30 mins at 37 °C, as described previously (Pandey *et al*, 2019), before proceeding with the same RNA extraction protocol. For RNAseq, three to five biological replicates were extracted for each cell line and time point. Libraries were prepared with Illumina Truseq Stranded total RNA RiboZero GOLD kit and sequenced on the NovaSeq6000 platform with a 2x151 setup using NovaSeqXp workflow in S4 mode flowcell. Raw reads were processed using cutadapt v3.2 to trim adaptor sequences and low-

quality base pairs and discard short reads (options: -m 20 -e 0.1 -q 20 -O 1). The GRCh37 genome

assembly was used for all alignment, annotation, and downstream analysis steps.

CircRNA detection from RNAseq

Back-spliced junctions were detected from paired end reads using three separate programs;

CIRCexplorer2 v2.3.8 (Zhang *et al*, 2016), MapSplice2 v2.1.8 (Wang *et al*, 2010) and CIRI2 v2.0.6

(Gao *et al*, 2018). For CIRCexplorer2, modules were run with default options, reads were pre-aligned

with TopHat v2.0.9 tophat_fusion (with Bowtie v1.1.2 and Samtools v0.1.19) using –fusion-min-dist

200. MapSplice was run using python v2.7.6 and options –fusion-non-canonical –min-fusion-dist 200.

CIRI2 was run using Perl v5.26.2 with default settings after aligning reads with BWA v0.7.17 (with

Samtools v1.9). The R package circRNAprofiler v1.4.2 (Aufiero *et al*, 2020) was used as described in

the manual to merge and filter back-spliced junction reads from all three detection programs as well

as for circRNA annotation and extraction of circRNA sequences.

Differential expression and gene ontology analyses

Based on clustering analysis, two outlier samples (1xNES/1xD5) were initially removed before

differential expression analysis of both circRNAs and genes using DESeq2 v1.30.1 and applying the

ashr method of LFC shrinkage (Stephens, 2017). Thresholds of LFC values ≥ |2| and p-adjusted <0.05

were used for sub-setting circRNAs and genes as differentially expressed. For analysis of expression

over all three time points, results from DESeq2 likelihood ratio test were analysed with DEGreport

v1.26.0. Analysis of differential exon usage was performed using the DEXSeq package (v1.40.0)

(Anders *et al*, 2012) using provided python scripts to generate a flattened annotation file of exon bins,

with aggregation disabled, and to count reads within exon bins. The standard analysis pipeline

described in the tutorial was followed, defining exon bins as differentially used if estimated LFC was

≥|2|, padj <0.05 and exon base mean >10 and classifying genes as differentially spliced if at least one

exon of a gene was differentially used. As some exon bins defined by DEXSeq differed from annotated

exon regions, differential exon usage data was compared with circRNA exons based on position

information and using the intersect function of bedtools software v2.29.2. For annotated exons which

contained multiple exon bins, exon fold-changes were calculated as the mean estimated exon bin

fold-change. The clusterProfiler package v3.18.1 was used for all gene ontology analyses using either

biological process terms annotated in org.Hs.eg.db v3.12.0 or user input annotations for analysis of

NDD-associated gene enrichment. As GSEA requires only one entry per gene and numerous host

genes had multiple circRNAs, data were split into negatively and positively regulated circRNAs and

the minimum and maximum circRNA LFC values were assigned to host genes with duplicate entries.

PCR, quantitative PCR and Sanger sequencing validation of circRNAs

For all experiments, cDNA was generated using the iScript cDNA synthesis kit (Bio-Rad). To validate

circRNA detection, divergent primers **[Additional File 1: Fig S1b]** were designed as previously

described (Panda & Gorospe, 2018). PCR was performed using HotStarTaq (Qiagen) and bands

visualized by gel electrophoresis. For Sanger sequencing, sequences were amplified using

HotStarTaq and circular primers **[Additional File 1: Fig S1b]**. Products were isolated using DNA

Clean and Concentrator-5 kit (Zymo Research) or, if multiple products, bands were separated using

gel electrophoresis and extracted using Zymoclean Gel DNA Recovery Kit (Zymo Research).

Samples were sequenced by Eurofins Scientific before comparing alignments to predicted

annotations with nucleotide BLAST and visualizing alignments with ApE v2.0.49.0. Divergent primers

and SYBR green (Bio-Rad) were used for quantitative PCR (qPCR) analyses. For differential

expression validation, ciRNA expression was normalised to two reference circRNAs, *circRTN4* and

circCASK1G3, found to have high and consistent expression throughout NES differentiation. For

SFPQ targets circRNAs were normalized to *U6* snRNA. For siRNA experiments all genes/linear RNAs

were normalised to *GAPDH* mRNA. SPAG9 and CPSF6 were used as reference genes for profiling

SFPQ expression across differentiation time-points. For known circRNAs, divergent BSJ primers were

designed using circInteractome (Dudekula *et al*, 2016). For novel circRNA divergent BSJ junctions

and for all circRNA primers, circRNA sequences were recreated in SnapGene Viewer v5.1.1 based

on annotations from circRNAprofiler and exon sequences in Ensembl GRCh37 and primers designed

using NCBI primer blast. All primer sequences are listed in **Additional File 1: Table S8**.

Variant analysis

For circRNAs which annotated to known splice sites, exon regions were extracted based on the

longest transcript containing the upstream and downstream back-spliced exons and sub-grouped as

in **Fig 3a**, with the assumption that no exons were spliced out of circRNAs. GnomAD GRCh37 exomes

files v2.1.1 were downloaded for each chromosome and converted into bed format. *De novo* variants

associated with ASD cases and controls were sourced from Satterstrom *et al.* (2020) and SCZ

variants were pooled from various sources listed in **Additional File 1: Table S5**. All variants were

intersected with exon subgroups using bedtools and all downstream analysis was performed in R,

normalising all exon variant frequencies by exon length. SIFT annotations for gnomAD exon variants

were predicted using Ensembl VEP v104.

miRNA and RBP binding site prediction

A list of 125 miRNAs associated with neurodevelopment or NDDs **[Additional File 6: Table S6a,b]**

was collated from literature review and pubmed searches for miRNA/microRNA with terms such as

neurodevelopment/neuronal/neuron differentiation etc. Ensembl biomart was used to extract 3'UTR

sequences for DEGs and, if multiple 3'UTR sequences were annotated to a single gene, the longest

3'UTR sequence was used. miRNA binding sites in circRNA and 3'UTR sequences were predicted

using remote RNA22 v2 with the following parameters: seedmismatchesallowed = 0; seedsize = 8;

minpairmatchs = 12; minenergy = -12; maxguwobbles = 1; threshold = 7000000. To examine

enrichment of RNA-binding motifs in circRNAs, separate fasta files of DEcircs were generated for

circRNAs decreased or increased at D28vNES / D28vD5 and D5vNES as well as all 4 LRT clusters.

RNA-binding protein motifs were downloaded from the ATtRACT database v0.99β. Motif enrichment

was tested using the AME function of MEME-suite v5.3.0 with default settings and a background

dataset of all annotated (5839/6385) circRNAs. For individual motif site detection, the MEME-suite

FIMO function was used with a threshold p-value of 0.001.

siRNA mediated silencing and protein detection

For siRNA mediated knockdown, Accell Human SMARTpool siRNAs targeting SFPQ (E-006455-00-

0010) or TIAL1 (E-011405-00-0010) were added to cell media at a final concentration of 0.3 µM.

Accell non-targeting control pool (D-001910-10-20), transfected at the same concentration was used

as a control treatment (siNTC). Protein samples were collected 96 hr after siRNA treatment. After

removing media and washing once with PBS cells collected with a cell scraper in a lysis buffer of 50

nM Tris-HCl, 100 mM 100 mM NaCl, 5 mM EDTA, 1 mM EGTA and 1X HALT protease inhibitor

cocktail (PIC; Fischer Scientific) and samples sonicated at 36% amplitude for 6X 1 sec. The Qubit

Protein Assay Kit (Thermo Scientific) was used to measure protein concentration. Protein was

detected using the Simple-Western-JESS capillary western blot system (Bio Techne) multiplexed for

total protein and chemiluminescence detection of mouse αSFPQ (1:2000; Sigma-Aldrich

WH0006421M2, RRID:AB_1843565) or mouse αTIAL1 (1:10; Santa Cruz sc-398372). The Compass

for S.W. software v5.0.1 was used to analyse data and calculate peak area normalised to total protein

area.

In situ hybridisation

Locked nucleic acid (LNA) probes for *in situ* were designed to be 45-55 bps in length with a melting

temperature of ~70-75°C. Probes were designed to be complementary to the BSJ for circRNA

detection while linear control probes were designed to target a linear splice junction, within the same

transcript, containing one of the BSJ exons. CircRNA probes plus a negative control scrambled

circRNA probe labelled with 5'FAM and linear probes labelled with 5'Cy3 were ordered from

Integrated DNA Technologies (IDT) as HPLC purified, Affinity Plus DNA oligos and resuspended in

TE buffer. Details of all probe sequences are listed in **Table S9**. Cells were plated on glass coverslips

coated with 100 µg/ml poly-L-ornithine and 2 µg/ml laminin. *In situ* hybridisation was performed as

described previously with some modifications (Zirkel & Papantonis, 2018). Cells were washed with

PBS before incubation in fresh fixative (4% PFA, 0.9% NaCl and 5% Glacial acetic acid in UltraPure

H_2O) for 17 min at room temp. Coverslips were washed for 5 min with PBS before adding ice-cold

70% ethanol and storing at least overnight or up to a few weeks at -20°C. After removing from -20°C

cells were recovered for 5 min at room temp in PBS, permeabilised for 10 min in 0.5% Triton X-

100/0.5% saponin (Sigma-Aldrich S7900) and washed in PBS. Cells were then treated with 1%

formaldehyde in PBS for 10 min and washed once in PBS for 10 min. Cells were dehydrated with

ethanol at 70, 90 and 100% for 3 min each and air-dried. For each coverslip 15 ng of labelled probe

was diluted in 50 µL of hybridisation buffer consisting of 10% Dextran sulfate (Sigma-Aldrich 265152),

0.1% Sodium dodecyl sulfate (Sigma-Aldrich 71736), 30% Ethylene carbonate (Sigma-Aldrich

E26258), 20 µg/mL Yeast tRNA (Thermo Fischer AM7119) and 2X SSC. Probes were denatured at

83°C for 5 min and hybridised overnight at 37°C in a box humidified with 2X SSC. Coverslips were

washed 3 times in 2X SSC for 10 min at 42°C, followed by two 3-min PBS washes at room temp.

Cells were blocked for 30 min in 0.1% Triton X/5% Donkey serum/PBS and counterstained with

1:1000 Hoechst. After two 3-min washes in PBS, coverslips were air-dried for 3-5 min and mounted

onto glass slides using Prolong Diamond Antifade Mountant (Thermo Fischer). Images were acquired

using an LSM 700 Zeiss Confocal Microscope and contrast was enhanced uniformly.

Formaldehyde RNA immunoprecipitation

Formaldehyde RNA immunoprecipitation (fRIP) was performed as previously described (Hendrickson

et al, 2016) with some modifications. NES cells were differentiated until D28 in a T75 flask. To collect,

cells were washed with PBS, rinsed with 6 mL Accutase for 10 sec and incubated in 4 mL TrypLE

select for 2 min/37°C before addition of 200 µL of Neuronal isolation enzyme (Thermo Scientific,

88285) and further incubating 1 min/37°C. Dissociation was stopped with 6 mL of Trypsin inhibitor

and solution collected, any remaining cells were collected by rinsing flasks with 6 mL of Wash Medium

(DMEM/F12+GlutaMAX, 5% FBS, 10 U/ml penicillin/streptomycin). Clumps were broken up by gentle

pipetting and cells spun at 400 x g for 3 min and resuspended in Wash Medium at 5×10^6 cells/mL.

Crosslinking was performed by addition of formaldehyde to a final concentration of 0.1% and

incubation for 10 min with rotation. Glycine was added at a concentration of 125 mM, incubating for 5

min with rotation to stop the reaction. Cells were pelleted, washed twice in cold PBS and once in PIC,

and stored at -80°C. For immunoprecipitation, cell pellets were resuspended and lysed by incubation

for 10 min at 4 °C with rotation in RIPA lysis buffer (50 mM TRIS [pH 8], 150 mM KCL, 0.1% SDS, 1%

Triton-X, 5 mM EDTA, 0.5% sodium deoxycholate, 0.5 mM DTT, 1X PIC, 100 U/mL RNaseOUT). The

lysate was sonicated 15 times (0.7s on, 1.3s off, Vibra-Cell VCX-600, Sonics) at an amplitude of 10%

and centrifuged at max speed for 10 min at 4°C. Supernatant was collected and diluted with an equal

volume of Binding buffer (150 mM KCl, 25 mM Tris [pH 7.5], 5 mM EDTA, 0.5 % IGEPAL) with 0.5

mM DTT, 1X PIC and 100 U/mL RNaseOUT added. Cells were incubated 2 hr at 4°C with rotation

after addition of either 6 μg mouse αSFPQ (Sigma-Aldrich WH0006421M2, RRID:AB_1843565)) or 6

μg normal mouse αIgG (Santa Cruz sc-2025, RRID:AB_737182). To each immunoprecipitation, 50

μL of Dynabeads Protein G (Thermo Fischer 10004D), pre-washed 2X in Binding buffer, was added

before incubating 1 hr/4°C with rotation. Beads were washed 2X 10 min/4°C with rotation in 1 mL

Binding buffer/PIC/RNaseOUT. To reverse cross-linking, beads were resuspended in 56 μL UltraPure

water and 33 μL reverse crosslinking buffer (PBS (without Mg or Ca), 6% N-lauroyl sarcosine, 30 mM

EDTA, 15 mM DTT), 2 mg/mL Proteinase K and 400 U/mL RNaseOUT were added to each reaction,

before incubating 1 hr/42°C followed by 1 hr/55°C. Bead/buffer mix was added to 1 mL TRIzol and

RNA extraction performed as described above. For each fRIP, 5 ng of RNA was used for cDNA

synthesis and targets were analysed by qPCR, normalizing SFPQ-fRIP to IgG-fRIP.

Intron analysis

Sequences for introns flanking circRNA back-spliced exons were extracted using BSgenome v1.58.0

and Biostrings v2.58.0 based on annotations output by circRNAprofiler. RBP binding sites were

predicted using MEME-suite as described above and SFPQ motif frequency for each subset of

circRNAs normalised by the number of circRNAs. All GRCh37 repeat elements were downloaded

from UCSC table browser RepeatMasker track and Alu elements extracted using grep. Bedtools

intersect was used to subset Alu elements that overlapped with intronic regions by at least 50% of the

repeat element length and bedtools closest tool used to find Alu elements closest to the BSJ site.

CircRNAs were classified as having inverted Alu elements when the closest Alu elements in the

downstream and upstream flanking introns were in opposite orientation (on opposite strands).

CircRNAs were classified as having proximal IAE flanking introns based on a total Alu distance from

the BSJ less than the median for all circRNAs with IAEs. Classification of circRNAs with short or long

flanking introns was also classified based on median length, totalled from both flanking introns.

Statistics and data visualisation

All statistics and data visualisation were performed in R v4.0.4. Specific statistical tests performed are

described within the relevant results section. All replicates specified within figure legends indicate

biological replicates i.e., cells cultured separately at successive passages. Various R packages were

used for visualisation of data including ggplot2 v3.3.3, EnhancedVolcano v1.8.0 and VennDiagram

v1.6.20

Declarations

Ethics approval and consent to participate

Not Applicable

Consent for publication

Not Applicable

Availability of data and materials

Generated raw sequencing data is unavailable as it contains potentially identifiable human data. All

other related data and code are available at: https://github.com/Tammimies-Lab/circRNA_neuro.git

Ethics approval and consent to participate

Not Applicable

Competing interests

The authors have no competing interests to declare.

Funding

The project was supported by the Swedish Research Council (KT), Swedish Foundation for Strategic

Research (KT), the Swedish Brain Foundation – Hjärnfonden (KT), Strategic Research Area

Neuroscience Stratneuro (KT, MW), The Swedish Foundation for International Cooperation in

Research and Higher Education STINT (KT), The Wenner Gren Foundation (MW), Sällskapet

Barnavård (MW), the Board of Research at Karolinska Institutet (KT) and the National Institute of

Aging's Intramural Research Program, National Institutes of Health (MG). Open access funding

provided by Karolinska Institutet.

Authors' contributions

MW, KT & MG designed experiments. MW, MO, SL & FM performed experiments. MW analysed data

and wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgments

The authors acknowledge support from the National Genomics Infrastructure in Stockholm funded by

Science for Life Laboratory, the Knut and Alice Wallenberg Foundation and the Swedish Research

Council, the SNIC/Uppsala Multidisciplinary Center for Advanced Computational Science for

assistance with massively parallel sequencing and access to the UPPMAX computational

infrastructure, and the iPS Core facility at Karolinska Institutet for assistance with the generation of

iPSC and NES cells.

References

Ahmed I, Karedath T, Al-Dasim FM, Malek JA (2019) Identification of human genetic variants

controlling circular RNA expression. *RNA* 25: 1765-1778

Anders S, Reyes A, Huber W (2012) Detecting differential usage of exons from RNA-seq data.

Genome Res 22: 2008-2017

Ashwal-Fluss R, Meyer M, Pamudurti NR, Ivanov A, Bartok O, Hanan M, Evantal N, Memczak S,

Rajewsky N, Kadener S (2014) circRNA biogenesis competes with pre-mRNA splicing. *Mol Cell* 56:

55-66

Aufiero S, Reckman YJ, Tijsen AJ, Pinto YM, Creemers EE (2020) circRNAprofiler: an R-based

computational framework for the downstream analysis of circular RNAs. *BMC Bioinformatics* 21: 164

Bachmayr-Heyda A, Reiner AT, Auer K, Sukhbaatar N, Aust S, Bachleitner-Hofmann T, Mesteri I,

Grunt TW, Zeillinger R, Pils D (2015) Correlation of circular RNA abundance with proliferation--

exemplified with colorectal and ovarian cancer, idiopathic lung fibrosis, and normal human tissues.

Sci Rep 5: 8057

Barbagallo D, Caponnetto A, Brex D, Mirabella F, Barbagallo C, Lauretta G, Morrone A, Certo F,

Broggi G, Caltabiano R *et al* (2019) CircSMARCA5 Regulates VEGFA mRNA Splicing and

Angiogenesis in Glioblastoma Multiforme Through the Binding of SRSF1. *Cancers (Basel)* 11

Becker M, Mastropasqua F, Reising JP, Maier S, Ho ML, Rabkina I, Li D, Neufeld J, Ballenberger L,

Myers L *et al* (2020) Presynaptic dysfunction in CASK-related neurodevelopmental disorders. *Transl*

Psychiatry 10: 312

Chen BJ, Huang S, Janitz M (2019) Changes in circular RNA expression patterns during human foetal

brain development. *Genomics* 111: 753-758

Chen YJ, Chen CY, Mai TL, Chuang CF, Chen YC, Gupta SK, Yen L, Wang YD, Chuang TJ (2020)

Genome-wide, integrative analysis of circular RNA dysregulation and the corresponding circular RNA-

microRNA-mRNA regulatory axes in autism. *Genome Res* 30: 375-391

Cocquerelle C, Daubersies P, Majerus MA, Kerckaert JP, Bailleul B (1992) Splicing with inverted order

of exons occurs proximal to large introns. *EMBO J* 11: 1095-1098

Conn VM, Hugouvieux V, Nayak A, Conos SA, Capovilla G, Cildir G, Jourdain A, Tergaonkar V,

Schmid M, Zubieta C *et al* (2017) A circRNA from SEPALLATA3 regulates splicing of its cognate

mRNA through R-loop formation. *Nat Plants* 3: 17053

Cortes-Lopez M, Gruner MR, Cooper DA, Gruner HN, Voda AI, van der Linden AM, Miura P (2018)

Global accumulation of circRNAs during aging in Caenorhabditis elegans. *BMC Genomics* 19: 8

Dell'Orco M, Elyaderani A, Vannan A, Sekar S, Powell G, Liang WS, Neisewander JL, Perrone-

Bizzozero NI (2021) HuD Regulates mRNA-circRNA-miRNA Networks in the Mouse Striatum Linked

to Neuronal Development and Drug Addiction. *Biology (Basel)* 10

Dolmetsch R, Geschwind DH (2011) The human brain in a dish: the promise of iPSC-derived neurons.

Cell 145: 831-834

Du WW, Yang W, Liu E, Yang Z, Dhaliwal P, Yang BB (2016) Foxo3 circular RNA retards cell cycle

progression via forming ternary complexes with p21 and CDK2. *Nucleic Acids Res* 44: 2846-2858

Dudekula DB, Panda AC, Grammatikakis I, De S, Abdelmohsen K, Gorospe M (2016)

CircInteractome: A web tool for exploring circular RNAs and their interacting proteins and microRNAs.

RNA Biol 13: 34-42

Fukuda Y, Pazyra-Murphy MF, Silagi ES, Tasdemir-Yilmaz OE, Li Y, Rose L, Yeoh ZC, Vangos NE,

Geffken EA, Seo HS *et al* (2021) Binding and transport of SFPQ-RNA granules by KIF5A/KLC1

motors promotes axon survival. *J Cell Biol* 220

Furukawa MT, Sakamoto H, Inoue K (2015) Interaction and colocalization of HERMES/RBPMS with

NonO, PSF, and G3BP1 in neuronal cytoplasmic RNP granules in mouse retinal line cells. *Genes*

Cells 20: 257-266

Gabel HW, Kinde B, Stroud H, Gilbert CS, Harmin DA, Kastan NR, Hemberg M, Ebert DH, Greenberg

ME (2015) Disruption of DNA-methylation-dependent long gene repression in Rett syndrome. *Nature*

522: 89-93

Gao Y, Zhang J, Zhao F (2018) Circular RNA identification based on multiple seed matching. *Brief*

Bioinform 19: 803-810

Glazar P, Papavasileiou P, Rajewsky N (2014) circBase: a database for circular RNAs. *RNA* 20: 1666-

1670

Gokool A, Anwar F, Voineagu I (2020) The Landscape of Circular RNA Expression in the Human

Brain. *Biol Psychiatry* 87: 294-304

Gruhl F, Janich P, Kaessmann H, Gatfield D (2021) Circular RNA repertoires are associated with

evolutionarily young transposable elements. *Elife* 10

Gruner H, Cortes-Lopez M, Cooper DA, Bauer M, Miura P (2016) CircRNA accumulation in the aging

mouse brain. *Sci Rep* 6: 38907

Guo JU, Agarwal V, Guo H, Bartel DP (2014) Expanded identification and characterization of

mammalian circular RNAs. *Genome Biol* 15: 409

Hansen TB (2018) Improved circRNA Identification by Combining Prediction Algorithms. *Front Cell*

Dev Biol 6: 20

Hendrickson DG, Kelley DR, Tenen D, Bernstein B, Rinn JL (2016) Widespread RNA binding by

chromatin-associated proteins. *Genome Biol* 17: 28

Heyd F, Lynch KW (2010) Phosphorylation-dependent regulation of PSF by GSK3 controls CD45

alternative splicing. *Mol Cell* 40: 126-137

Hollensen AK, Thomsen HS, Lloret-Llinares M, Kamstrup AB, Jensen JM, Luckmann M, Birkmose N,

Palmfeldt J, Jensen TH, Hansen TB *et al* (2020) circZNF827 nucleates a transcription inhibitory

complex to balance neuronal differentiation. *Elife* 9

Ishigaki S, Riku Y, Fujioka Y, Endo K, Iwade N, Kawai K, Ishibashi M, Yokoi S, Katsuno M, Watanabe

H *et al* (2020) Aberrant interaction between FUS and SFPQ in neurons in a wide range of FTLD

spectrum diseases. *Brain* 143: 2398-2405

Ivanov A, Memczak S, Wyler E, Torti F, Porath HT, Orejuela MR, Piechotta M, Levanon EY,

Landthaler M, Dieterich C *et al* (2015) Analysis of intron sequences reveals hallmarks of circular RNA

biogenesis in animals. *Cell Rep* 10: 170-177

Jeck WR, Sorrentino JA, Wang K, Slevin MK, Burd CE, Liu J, Marzluff WF, Sharpless NE (2013)

Circular RNAs are abundant, conserved, and associated with ALU repeats. *RNA* 19: 141-157

Kaneko S, Rozenblatt-Rosen O, Meyerson M, Manley JL (2007) The multifunctional protein

p54nrb/PSF recruits the exonuclease XRN2 to facilitate pre-mRNA 3' processing and transcription

termination. *Genes Dev* 21: 1779-1789

King IF, Yandava CN, Mabb AM, Hsiao JS, Huang HS, Pearson BL, Calabrese JM, Starmer J, Parker

JS, Magnuson T *et al* (2013) Topoisomerases facilitate transcription of long genes linked to autism.

Nature 501: 58-62

Li D, Choque Olsson N, Becker M, Arora A, Jiao H, Norgren N, Jonsson U, Bolte S, Tammimies K

(2022a) Rare variants in the outcome of social skills group training for autism. *Autism Res* 15: 434-

446

Li W, Shan B, Cheng X, He H, Qin J, Zhao H, Tian M, Zhang X, Jin G (2022b) circRNA Acbd6

promotes neural stem cell differentiation into cholinergic neurons via the miR-320-5p-Osbpl2 axis. *J

Biol Chem* 298: 101828

Liu Z, Wu X, Yu Z, Tang X (2021) Reconstruction of circRNA-miRNA-mRNA associated ceRNA

networks reveal functional circRNAs in intracerebral hemorrhage. *Sci Rep* 11: 11584

Ma N, Pan J, Ye X, Yu B, Zhang W, Wan J (2019) Whole-Transcriptome Analysis of APP/PS1 Mouse

Brain and Identification of circRNA-miRNA-mRNA Networks to Investigate AD Pathogenesis. *Mol

Ther Nucleic Acids* 18: 1049-1062

Mahmoudi E, Fitzsimmons C, Geaghan MP, Shannon Weickert C, Atkins JR, Wang X, Cairns MJ

(2019) Circular RNA biogenesis is decreased in postmortem cortical gray matter in schizophrenia and

may alter the bioavailability of associated miRNA. *Neuropsychopharmacology* 44: 1043-1054

Mai TL, Chen CY, Chen YC, Chiang TW, Chuang TJ (2022) Trans-genetic effects of circular RNA

expression quantitative trait loci and potential causal mechanisms in autism. *Mol Psychiatry*

Memczak S, Jens M, Elefsinioti A, Torti F, Krueger J, Rybak A, Maier L, Mackowiak SD, Gregersen

LH, Munschauer M *et al* (2013) Circular RNAs are a large class of animal RNAs with regulatory

potency. *Nature* 495: 333-338

Nigro JM, Cho KR, Fearon ER, Kern SE, Ruppert JM, Oliner JD, Kinzler KW, Vogelstein B (1991)

Scrambled exons. *Cell* 64: 607-613

Panda AC, Gorospe M (2018) Detection and Analysis of Circular RNAs by RT-PCR. *Bio Protoc* 8

Panda AC, Grammatikakis I, Munk R, Gorospe M, Abdelmohsen K (2017) Emerging roles and context

of circular RNAs. *Wiley Interdiscip Rev RNA* 8

Pandey PR, Rout PK, Das A, Gorospe M, Panda AC (2019) RPAD (RNase R treatment,

polyadenylation, and poly(A)+ RNA depletion) method to isolate highly pure circular RNA. *Methods*

155: 41-48

Pandey PR, Yang JH, Tsitsipatis D, Panda AC, Noh JH, Kim KM, Munk R, Nicholson T, Hanniford D,

Argibay D *et al* (2020) circSamd4 represses myogenic transcriptional activity of PUR proteins. *Nucleic*

Acids Res 48: 3789-3805

Passon DM, Lee M, Rackham O, Stanley WA, Sadowska A, Filipovska A, Fox AH, Bond CS (2012)

Structure of the heterodimer of human NONO and paraspeckle protein component 1 and analysis of

its role in subnuclear body formation. *Proceedings of the National Academy of Sciences of the United*

States of America 109: 4846-4850

Peng R, Dye BT, Perez I, Barnard DC, Thompson AB, Patton JG (2002) PSF and p54nrb bind a

conserved stem in U5 snRNA. *RNA* 8: 1334-1347

Piwecka M, Glazar P, Hernandez-Miranda LR, Memczak S, Wolf SA, Rybak-Wolf A, Filipchyk A,

Klironomos F, Cerda Jara CA, Fenske P *et al* (2017) Loss of a mammalian circular RNA locus causes

miRNA deregulation and affects brain function. *Science* 357

Rajcsanyi LS, Diebels I, Pastoors L, Kanber D, Peters T, Volckmar AL, Zheng Y, Grosse M, Dieterich

C, Hebebrand J *et al* (2022) Evidence for correlations between BMI-associated SNPs and circRNAs.

Sci Rep 12: 12643

Rosonina E, Ip JY, Calarco JA, Bakowski MA, Emili A, McCracken S, Tucker P, Ingles CJ, Blencowe

BJ (2005) Role for PSF in mediating transcriptional activator-dependent stimulation of pre-mRNA

processing in vivo. *Mol Cell Biol* 25: 6734-6746

Rybak-Wolf A, Stottmeister C, Glazar P, Jens M, Pino N, Giusti S, Hanan M, Behm M, Bartok O,

Ashwal-Fluss R *et al* (2015) Circular RNAs in the Mammalian Brain Are Highly Abundant, Conserved,

and Dynamically Expressed. *Mol Cell* 58: 870-885

Salzman J, Chen RE, Olsen MN, Wang PL, Brown PO (2013) Cell-type specific features of circular

RNA expression. *PLoS Genet* 9: e1003777

Salzman J, Gawad C, Wang PL, Lacayo N, Brown PO (2012) Circular RNAs are the predominant

transcript isoform from hundreds of human genes in diverse cell types. *PLoS One* 7: e30733

Santos-Rodriguez G, Voineagu I, Weatheritt RJ (2021) Evolutionary dynamics of circular RNAs in

primates. *Elife* 10

Satterstrom FK, Kosmicki JA, Wang J, Breen MS, De Rubeis S, An JY, Peng M, Collins R, Grove J,

Klei L *et al* (2020) Large-Scale Exome Sequencing Study Implicates Both Developmental and

Functional Changes in the Neurobiology of Autism. *Cell* 180: 568-584 e523

Sharma D, Sehgal P, Hariprakash J, Sivasubbu S, Scaria V (2019) Methods for Annotation and

Validation of Circular RNAs from RNAseq Data. *Methods Mol Biol* 1912: 55-76

Stagsted LV, Nielsen KM, Daugaard I, Hansen TB (2019) Noncoding AUG circRNAs constitute an

abundant and conserved subclass of circles. *Life Sci Alliance* 2

Stagsted LVW, O'Leary ET, Ebbesen KK, Hansen TB (2021) The RNA-binding protein SFPQ

preserves long-intron splicing and regulates circRNA biogenesis in mammals. *Elife* 10

Stephens M (2017) False discovery rates: a new deal. *Biostatistics* 18: 275-294

Suenkel C, Cavalli D, Massalini S, Calegari F, Rajewsky N (2020) A Highly Conserved Circular RNA

Is Required to Keep Neural Cells in a Progenitor State in the Mammalian Brain. *Cell Rep* 30: 2170-

2179 e2175

Takeuchi A, Iida K, Tsubota T, Hosokawa M, Denawa M, Brown JB, Ninomiya K, Ito M, Kimura H,

Abe T *et al* (2018) Loss of Sfpq Causes Long-Gene Transcriptopathy in the Brain. *Cell Rep* 23: 1326-

1341

Thomas-Jinu S, Gordon PM, Fielding T, Taylor R, Smith BN, Snowden V, Blanc E, Vance C, Topp S,

Wong CH *et al* (2017) Non-nuclear Pool of Splicing Factor SFPQ Regulates Axonal Transcripts

Required for Normal Motor Development. *Neuron* 94: 931

Tsitsipatis D, Grammatikakis I, Driscoll RK, Yang X, Abdelmohsen K, Harris SC, Yang JH, Herman

AB, Chang MW, Munk R *et al* (2021) AUF1 ligand circPCNX reduces cell proliferation by competing

with p21 mRNA to increase p21 production. *Nucleic Acids Res* 49: 1631-1646

Uhlin E, Ronnholm H, Day K, Kele M, Tammimies K, Bolte S, Falk A (2017) Derivation of human iPS

cell lines from monozygotic twins in defined and xeno free conditions. *Stem Cell Res* 18: 22-25

Veno MT, Hansen TB, Veno ST, Clausen BH, Grebing M, Finsen B, Holm IE, Kjems J (2015) Spatio-

temporal regulation of circular RNA expression during porcine embryonic brain development. *Genome*

Biol 16: 245

Wang K, Singh D, Zeng Z, Coleman SJ, Huang Y, Savich GL, He X, Mieczkowski P, Grimm SA, Perou

CM *et al* (2010) MapSplice: accurate mapping of RNA-seq reads for splice junction discovery. *Nucleic*

Acids Res 38: e178

Westholm JO, Miura P, Olson S, Shenker S, Joseph B, Sanfilippo P, Celniker SE, Graveley BR, Lai

EC (2014) Genome-wide analysis of drosophila circular RNAs reveals their structural and sequence

properties and age-dependent neural accumulation. *Cell Rep* 9: 1966-1980

Yang M, Xiang G, Yu D, Yang G, He W, Yang S, Zhou G, Liu A (2019) Hsa_circ_0002468 Regulates

the Neuronal Differentiation of SH-SY5Y Cells by Modulating the MiR-561/E2F8 Axis. *Med Sci Monit*

25: 2511-2519

Yang Q, Wu J, Zhao J, Xu T, Zhao Z, Song X, Han P (2018) Circular RNA expression profiles during

the differentiation of mouse neural stem cells. *BMC Syst Biol* 12: 128

You X, Vlatkovic I, Babic A, Will T, Epstein I, Tushev G, Akbalik G, Wang M, Glock C, Quedenau C

et al (2015) Neural circular RNAs are derived from synaptic genes and regulated by development and

plasticity. *Nat Neurosci* 18: 603-610

Younas N, Zafar S, Shafiq M, Noor A, Siegert A, Arora AS, Galkin A, Zafar A, Schmitz M, Stadelmann

C *et al* (2020) SFPQ and Tau: critical factors contributing to rapid progression of Alzheimer's disease.

Acta Neuropathol 140: 317-339

Zhang XO, Dong R, Zhang Y, Zhang JL, Luo Z, Zhang J, Chen LL, Yang L (2016) Diverse alternative

back-splicing and alternative splicing landscape of circular RNAs. *Genome Res* 26: 1277-1287

Zirkel A, Papantonis A (2018) Detecting Circular RNAs by RNA Fluorescence In Situ Hybridization.

Methods Mol Biol 1724: 69-75

Figure Legends

Figure 1. Detection and validation of circRNAs in human neuronal differentiation**. a.** Schematic

representation of sample and data collection. **b.** Venn diagram of circRNAs detected from three different

circRNA detection programs, numbers highlighted in bold indicate circRNAs included in downstream

analysis. **c.** Volcano plots of differentially expressed circRNAs at D5 and D28 of differentiation compared

to NES. **d.** Venn diagram of significantly dysregulated circRNAs (padj < 0.05 and LFC ≥|2|). **e.** Cluster

analysis of circRNAs differentially expressed throughout differentiation. **f.** Detection of top differentially

expressed circRNAs by PCR with divergent junction primers from RNAse R-treated samples. Samples

from D28 were used to detect 15 increased circRNAs (left) and samples from NES cells were used for

detection of 5 decreased circRNAs (right). Asterisks indicate reactions with additional bands or no product,

blue highlighting indicates the product is an isoform of the intended circRNA target, red highlighting

indicates PCR artifact. **g.** Example schematic of sanger sequencing validation of circRNA exon retention

in *circATRNL1*. Exons are shown in purple, green arrows indicate primer location, and sequencing

alignment is shown in orange.

Figure 2. CircRNAs increased during NES differentiation arise from host genes with synaptic function. **a.**

GSEA of gene ontology – biological process terms among the genes that host circRNAs increased at D28

of differentiation compared to NES. **b.** Most highly enriched gene ontology – biological process terms

among host genes of Cluster 1 circRNAs. **c.** Correlation of LFC values for circRNAs and their

corresponding linear counterparts. **d.** Alternative splicing comparison between all RNAs, RNA which host

circRNAs and RNAs that host circRNAs increased at D28 vs. NES. **** = p <0.0001, hypergeometric p-

value test.

Figure 3. Genetic variant frequency in circRNA-forming exons**. A.** Schematic depicting how exons were subdivided for comparative analysis. BSJ = back-spliced junctions. **B.** Frequency of GnomAD exon variants for circRNAs detected in this study, normalised for exon size (Kb). **C.** Proportion of GnomAD exon variants annotated as having either low, medium, or high impact. **D.** Frequency of GnomAD splice-region exon variants/Kb. **E.** Frequency of rare *de novo* ASD variants/Kb. **F.** Frequency of rare *de novo* ASD variants/Kb in exons of circRNAs arising from ASD-associated SFARI host genes. Two sample Z-test/Wilcox test, **** = p < 0.0001, *** = p < 0.001 * = p < 0.05. DNVs = *de novo* variants.

Figure 4. RNA-binding protein SFPQ binds and regulates subsets of circRNAs during neuronal differentiation**. a.** SFPQ motif enriched among circRNAs increased at D28 v D5. **b.** TIAL1 motif enriched among circRNAs increased at D28 v NES. **c.** Normalised counts of SFPQ RNA during NES differentiation. **d.** Normalised counts of TIAL1 RNA during NES differentiation. FC = LFC from DEseq2 analysis, **** = DEseq2 adjusted p-value < 0.0001. **e.** Detection of SFPQ and TIAL1 mRNA following siRNA treatment (*n* = 3). **f.** Relative expression of SFPQ targets following SFPQ knockdown (*n* = 3)**. g-j.** *In situ* detection of *circERC2* and its linear isoform in undifferentiated NES cells (**g**), and cells at D5 (**h**) and D28 (**i,j**) of differentiation. Blue = Hoechst nuclear counter stain, magenta = linear ERC2 probe, green = *circERC2* probe. Scale bar in g = 25 µm, applies to g-j. images are representative of staining from biological replicates (*n* = 3). **k.** SFPQ RIP followed by RT-qPCR analysis to identify SFPQ targets and *5S* rRNA control, normalised to IgG RIP samples from D28 differentiated neurons (*n* = 2). **** = p < 0.0001, *** = p < 0.001, ** = p < 0.01, * = p < 0.05, one-way ANOVA, Tukey post-hoc analysis, Bonferroni adjusted. Error bars represent standard error.

Figure 5. Long introns flank circRNAs increased by NES differentiation. **a.** Frequency of SFPQ motifs in circRNA flanking introns by distance relative to intron splice site distal from the BSJ. **b.** Frequency of SFPQ motifs in circRNA flanking introns by distance relative to the 5' intron end. Scale in b applies to a and b. **c.** Proportion of flanking intron types among all circRNAs compared to increased circRNAs. Intron types were defined as either, short/long introns with a proximal IAE (pIAE-short/pIAE-long) or short/long introns

without proximal IAE (Short/Long). **d.** Mean intron length of flanking introns. **e.** Mean distance to IAE from

BSJ. **f.** Density plot of intron length totalled from both flanking introns. **g.** Density plot of IAE distance

totalled from Alu element distance on both flanking introns. Error bars in d and e represent standard error,

dashed lines in g and f indicate median values. Two sample Z-test, **** = p < 0.0001.

Additional Files

Additional File 1

Format: Portable document format (.pdf)

Title: Supplementary information

Description: Figure S1. Validation of circRNA expression. Figure S2. Validation of circRNA exon retention.

Figure S3. Pathway enrichment among differentially expressed circRNAs. Figure S4. Pathway enrichment

among differentially expressed linear RNAs. Figure S5. Log2 fold-change values of top increased

circRNAs and their corresponding linear host RNAs (D28 vs NES). Figure S6. Variant analysis in circRNA

exons. Table S5. Source studies for compilation de novo variants identified in individuals with

Schizophrenia or Schizoaffective disorder. Figure S7. SFPQ and TIAL1 target detection and siRNA

knockdown. Figure S8. Extended expression profile of SFPQ and identified circRNA targets in NES

differentiation. Figure S9. In Situ detection of circRNAs during NES differentiation. Figure S10. Analysis of

introns from circRNA host genes. Table S8. Primer sequences.

Additional File 2

Format: Excel workbook (.xlsx)

Title: Table S1

Description: Supplementary Table 1. Differential expression analysis of circRNAs in NES differentiation.

a. Hotspot host genes which generated over 10 circRNAs. b. Pairwise differential expression (DE) analysis

of circRNAs at D28 of differentiation compared to NES. c. Pairwise DE analysis of circRNAs at D5 of

differentiation compared to NES. d. Pairwise DE analysis of circRNAs at D28 compared to D5 of

differentiation. e. Likelihood ratio test DE circRNA analysis of all time points. f. Clustering of LRT results.

Additional File 3

Format: Excel workbook (.xlsx)

Title: Table S2

Description: Supplementary Table 2. Gene ontology analysis of circRNAs differentially expressed during

differentiation. a. Biological processes enriched among host genes of Cluster 1 cicrRNAs. b. Positive gene

set enrichment analysis of D28 compared to NES circRNA host genes. c. Positive gene set enrichment

analysis of D5 compared to NES circRNA host genes. d. Biological processes enriched among host genes

of Cluster2 circRNAs.

Additional File 4

Format: Excel workbook (.xlsx)

Title: Table S3

Description: Supplementary Table 3. Collated list of Schizophrenia risk genes. a. Gene list. b. Source

studies.

Additional File 5

Format: Excel workbook (.xlsx)

Title: Table S4

Description: Supplementary Table 4. Analysis of linear RNA expression in NES differentiation. a.

Differential expression analysis of linear RNAs at D28 compared to NES. b. Positive gene set enrichment

analysis of linear RNAs at D28 compared to NES.

Additional File 6

Format: Excel workbook (.xlsx)

Title: Table S6

Description: Supplementary Table 6. Prediction of circRNA:miRNA:mRNA interactions. a. Collated list of

miRNAs linked to neurodevelopment. b. Reference studies for miRNA list. c. miRNA sites predicted in

DEcircs. d. miRNA sites predicted in 3'UTRs of differentially expressed mRNAs. e. circRNA:miRNA:mRNA

interactions with correlated expression during differentiation.

Additional File 7

Format: Excel workbook (.xlsx)

Title: Table S7

Description: Supplementary Table 7. Prediction of RNA binding protein (RBP) motifs in circRNAs and thier

flanking introns. a. Significantly enriched RBP motifs within sequences of differentially expressed circRNA

subsets. b. SFPQ target selection from positive circRNA sequences increased at Day 28 compared to Day

5. c. TIAL1 target selection from positive Cluster 1 circRNA sequences and circRNA sequences increased

at Day 28 compared to Day 5. d. Significantly enriched RBP motifs within flanking intron sequences of

circRNA increased at Day 28 compared to NES.

a.

	BSJ	
up-D28vNES		
up-D28vD5		
dwn-D28vNES		
dwn-D28vD5		
All circRNAs		

4Kb 8Kb 12Kb 16Kb 20Kb 16Kb 12Kb 8Kb 4Kb

b.

upstream   downstream

5'   5'

up-D28vNES		
up-D28vD5		
dwn-D28vNES		
dwn-D28vD5		
All circRNAs		

4Kb 8Kb 12Kb 16Kb 20Kb   4Kb 8Kb 12Kb 16Kb 20Kb

Motif frequency

2 4 6 8

c.

Flanking intron type (%)

- Long
- Short
- pIAE-Long
- pIAE-Short

circs   upD28_circs

d.

Mean intron length (nts)

****

circs   upD28_circs

e.

Mean Alu distance (nts)

circs   upD28_circs

f.

Density

— circRNAs
— upD28-circcs

Total flanking intron length (Log₁₀[nts])

g.

Density

— circRNAs
— upD28-circs

Total distance to Alu elements (Log10[nts])