Title:

Perineuronal net loosening associates with the remodeling of GABAergic

terminals on motor cortical layer 5 interneurons and precedes motor

coordination recovery post stroke

Short Title:

PNN remodeling post stroke

Egor Dzyubenko[1,*], Katrin I. Willig[2], Dongpei Yin[1], Maryam Sardari[1], Erdin Tokmak[1],

Patrick Labus[1], Ben Schmermund[1], and Dirk M. Hermann[1,*]

[1]Department of Neurology and Center for Translational Neuro- and Behavioral Sciences

(C-TNBS), University Hospital Essen, Hufelandstraße 55, Essen D-45122, Germany

[2]Group of Optical Nanoscopy in Neuroscience, Max Planck Institute for Multidisciplinary

Sciences, City Campus, Hermann-Rein-Str. 3, 37075 Göttingen, Germany

*Corresponding authors

Egor Dzyubenko, email: egor.dzyubenko@uk-essen.de;

Dirk M Hermann, email: dirk.hermann@uk-essen.de

Words (Abstract):   235

Words (main text and figure legends):   6896

Figures: 6

Supplementary figures: 4

Supplementary codes: 2

References: 63

Abstract

Stroke remains one of leading causes of long-term disability worldwide, and the development of effective restorative therapies is hindered by an incomplete understanding of intrinsic recovery mechanisms in the brain. Here, we explored the involvement of perineuronal nets (PNNs), the facet-like layers of extracellular matrix surrounding fast-spiking interneurons, in brain remodeling and neurological recovery after focal cerebral ischemia in mice with and without induced stroke tolerance. Due to insufficient resolution of conventional microscopy methods, the contribution of ultrastructural changes in PNNs to post stroke brain plasticity remained unknown. Using superresolution stimulated emission depletion (STED) and structured illumination (SR-SIM) microscopy, we revealed that PNN facets become larger and less dense in the post-acute stroke phase. Morphological alterations in PNNs are likely mediated by activated microglia cells, which preferentially enwrap the PNN-coated neurons. The loosening of PNNs is transient and associates with the increased number of GABAergic axonal terminals on inhibitory interneurons in the motor cortical layer 5. The coherent remodeling of PNNs and inhibitory synapses precedes the recovery of motor coordination during the late post-acute stroke phase. In the chronic stroke phase, the initial morphology of PNNs is restored, and the number of GABAergic axonal terminals on motor cortical interneurons is reduced. Our data suggests a novel mechanism of motor cortical plasticity after stroke, and we propose that prolonging PNN loosening during the post-acute period can extend the opening neuroplasticity window into the chronic stroke phase.

Keywords: extracellular matrix, cerebral ischemia, motor recovery, synaptic rewiring, parvalbumin interneurons, fluorescence nanoscopy

Introduction

Brain remodeling is essential for regaining compromised motor activity and coordination post stroke. Neurological recovery after stroke involves several neuroplasticity mechanisms including corticospinal tract rewiring [36], sprouting of interhemispheric cortico-cortical projections [40] and remodeling of local intracortical connectivity [7]. Motor cortical activity defining motor commands during skilled limb movements is selectively distributed across neurons with distinct projection patterns [47], suggesting that reorganization of both long-range and local connectivity is similarly pivotal for successful recovery after stroke. Experimental treatments promoting pyramidal tract plasticity have been proposed [50], but harnessing the plasticity of intracortical projections requires deeper understanding of local connectivity changes in the motor cortex post stroke. Cortical oscillations underlying motor learning arise from the fast-spiking activity of layer 5 (L5) interneurons [45], which are the main source of inhibition in neocortical microcircuits [43, 46]. Although motor cortical L5 interneurons are critical for controlling coordinated movements, their involvement in post stroke brain remodeling has not been systematically studied to the best of our knowledge.

Cortical L5 fast-spiking interneurons express parvalbumin (PV), potassium channels with rapid activation and deactivation kinetics (Kv3.1) and are coated with perineuronal nets (PNNs) on the extracellular side [11, 34, 42]. PNNs are condensed lattice-like layers of extracellular matrix (ECM) composed of multiple macromolecules including hyaluronic acid, chondroitin sulfate proteoglycans (CSPGs), and link proteins [10, 56]. These polymeric assemblies propagate to the extracellular space (ECS) and are anchored to neuronal surface via hyaluronic acid synthases [29, 39]. PNNs are rigid structures resistant to chemical decomposition [17] that are formed in an activity-dependent manner [18] and restrict synaptic plasticity [28, 49]. They compartmentalize neuronal surface, limit astrocyte-neuron direct membrane contacts and new synapse formation, stabilize existing connectivity and hypothetically contribute to potassium buffering (for review see [20, 26]). Thereby, PNNs maintain the fast-spiking properties of interneurons [57] and inhibitory control in neuronal networks [19].

In the adult brain, CSPGs within PNNs restrict neuronal plasticity by inhibiting

axonal sprouting [30], and experimental approaches involving non-specific ECM digestion

have been shown to promote functional recovery post stroke [31]. It is unlikely though

that the complete removal of ECM and PNNs in particular can be implemented clinically

because it induces epileptiform activity [2, 3] and impairs memory formation [32, 38].

Therefore, understanding the impact of more delicate PNN alterations on neuroplasticity

is imperative for developing novel therapies targeting brain ECM. In a previous study, we

developed a method allowing for topological quantification of PNN morphology based on

superresolution fluorescence microscopy [21]. We showed that despite their rigidity,

PNNs are subject to subtle modulation post stroke and anticipated that their transient

loosening can support brain remodeling. In this work, we have further elaborated our

method and investigated PNN morphology with unprecedented resolution and linked it to

synaptic remodeling and neurological recovery post stroke.

Materials and methods

Legal issues, animal housing and randomization

Experimental procedures were conducted in accordance with European Union

(Directive 2010/63/EU) guidelines for the care and use of laboratory animals and

approved by the local government (Bezirksregierung Düsseldorf). C57BL/6j mice were

kept in groups of 5 animals per cage, inverse 12/12 h light/dark cycle and access to food

and water *ad libitum*. All efforts were made to reduce the number of animals in the

experiments. The groups were randomly assigned using dummy names, and the

experimenters were blinded to group coding during sample preparation, data acquisition

and analysis.

Cerebral ischemia, stroke tolerance induction, and tissue collection

Wildtype male C57/Bl6 mice at the age of 2 months were randomly assigned into

three groups: stroke, preconditioning and stroke, and naive control. Each group included

7 animals. Focal cerebral ischemia was induced by transient left sided intraluminal middle

cerebral artery occlusion (tMCAO) for 30 min as described previously [22]. In brief, mice

were anesthetized with 1.5% v/v isoflurane (carrier gas was N_2O with 30% v/v O_2) and

150 µl of buprenorphine was injected subcutaneously. After exposing and ligating the
lower part of the left common carotid artery (CCA), a silicon-coated nylon monofilament
was introduced through a fine incision and advanced until the bifurcation of the middle
cerebral artery (MCA). Cessation of blood supply in the MCA territory was verified by
measuring laser Doppler flow (LDF). After 30 minutes, the filament was removed to
induce reperfusion, which was controlled by LDF recording. The tMCAO procedure
resulted in highly reproducible ischemic lesions located in striatum and adjacent cortical
areas, but not motor cortex and the produced infarcts had similar volumes as in our
previous studies [21, 52].

Stroke tolerance was induced by inflammatory preconditioning with 1 mg/kg LPS
that was injected intraperitoneally 3 days before tMCAO to trigger robust peripheral
immune response that we reported previously [52]. Experimental endpoints were defined
as 7, 14, 28, and 42 days post ischemia (DPI) to evaluate post stroke brain remodeling
during post-acute and chronic stroke phases (Fig. 1A). Upon reaching the pre-defined
endpoints, animals were anesthetized with 100 µl of ketamine-xylazine (1:3) and
sacrificed by transcardiac perfusion with 4% paraformaldehyde (PFA) in normal saline.
The brains were removed and immersed in 4% PFA solution for 12 hours at 4°C. Tissues
were cryoprotected in sucrose gradient solutions (10-30%), carefully dried, frozen, and
stored at -80 °C until further processing.

Infarct volume and brain atrophy measurement

Coronal sections of the brain (20 µm thick) were collected at 500 µm intervals
across the forebrain using a Leica CM1950 cryostat and placed onto cold microscope
slides (ThermoFisher Scientific, Cat# J1800AMNT). The sections were stained with cresyl
violet (that is, Nissl) and scanned using the AxioObserver Z1 microscope (objective Plan-
Apochromat 10×/0.45 M25; Zeiss, Jena, Germany). The infarct volume (IV) was
measured at 7 DPI as $IV = \Sigma(IA^*\Delta)$, where IA is the infarcted area on a section and Δ is
the interval between sections. Brain atrophy at 42 DPI was determined by subtracting the
areas of surviving tissue in the ipsilesional hemisphere from the area of the contralesional
hemisphere. Atrophy volume (AV) was calculated as $AV = \Sigma(AA^*\Delta)$, where AA is the
atrophy area on a section and Δ is the interval between sections.

Neurological deficits and motor performance tests

General and focal neurological deficits were analyzed using Clark's neuroscore [12] daily until 7 DPI, every 3 days until 14 DPI, and weekly until 42 DPI. Post stroke recovery of motor activity and coordination was assessed by tight rope at baseline, 7, 14, 21, 28, 35, and 42 DPI as described previously [52]. In brief, the tight rope test measures the time until the animals reach the platform from the middle of a 60-cm-long rope. Mice were pre-trained over 1–2 days before MCAO ensuring that they were able to reach the platform within 10 s.

Immunohistochemical procedures

Free-floating coronal sections (30 µm thick) were obtained at the level of bregma +0.5 to +1 mm using a Leica CM1950 cryostat and stored until use at -20 °C in 1:1 mixture of phosphate buffer saline (PBS) and ethylene glycol with 1% polyvinyl pyrrolidone. For immunohistochemistry, the sections were rinsed 0.1 M PBS and permeabilized with 0.3% w/v Triton X-100 in PBS. Non-specific antibody binding was blocked by applying a mixture of 10% v/v ChemiBLOCKER (Cat# 2170, Millipore, Burlington, MA, U.S.A.) and 5% v/v normal donkey serum in PBS for 12 hours at room temperature with gentle agitation. Sections were incubated with primary antibodies for 48 hours at 4 °C in PBS with 0.01% w/v Triton X-100. Astrocytes and microglia/macrophage cells were labeled using rat anti-GFAP (1:300; Cat# 13-0300, Thermo Fisher Scientific, Waltham, MA, U.S.A) and rabbit anti-IBA1 (1:500; Cat# 019-19741, Wako, Neuss, Germany) or guinea pig anito-IBA1 (1:300, Cat# 234308, Synaptic Systems, Goettingen, Germany) antibodies. Interneurons were detected with rabbit anti-Parvalbumin (1:500; Cat# 195002, Synaptic Systems) and rabbit anti-Kv3.1b (1:1000, Cat# APC-014, Alomone Labs, Jerusalem, Israel). PNNs were labeled with biotinylated WFA (1:100, B-1355, Vector Laboratories, Burlingame, USA). Glutamatergic and GABAergic synaptic terminals were detected using guinea pig anti-VGLUT1 (1:500, Cat# 135304, Synaptic Systems) and guinea pig anti-VGAT (1:500, Cat# 131103, Synaptic Systems) antibodies. For fluorescence detection, we used secondary antibodies conjugated to Alexa 488, 594 and 647 or streptavidin conjugated to Atto 495, Atto 590 or Abberior STAR RED dyes. Nuclei were counterstained with DAPI (1:1000, D1306, ThermoFisher). The bound antibodies were stabilized by incubating the sections

in 2% w/v PFA for 30 min at room temperature. For high-resolution imaging with confocal, SR-SIM and STED microscopy with oil immersion objectives, the refractive index of stained tissues was adjusted to 1.5 using 2,2'-thiodiethanol (TDE, Cat# 166782, Merck, Darmstadt, Germany), which is widely used for tissue clearing [13].

Quantification of glial and neuronal markers

The expression of GFAP and IBA1 markers was analyzed using a AxioObserver Z1 microscope (objective Plan-Apochromat 10x/0.45 M25; Zeiss). In the whole-section images obtained by tiling, 600x600 µm regions of interest (ROIs) were selected in the motor cortical layer 5 as shown in Fig 1E, G, and the mean pixel intensity was measured using ImageJ (National Institutes of Health, Bethesda, MD, U.S.A.). In each animal, four images obtained from two adjacent brain sections were analyzed in the ipsilesional and contralesional motor cortex.

Cell density of interneurons expressing PV, Kv3.1 and PNNs was quantified manually in 600x600x10 µm ROIs obtained in the motor cortical L5 regions using the LSM 710 confocal microscope (Zeiss, 20x Plan Apochromat objective, NA 0.8, pixel size 0.21 µm). In each animal, four image stacks obtained from two adjacent brain sections were analyzed in the ipsilesional and contralesional motor cortex.

Quantification of synaptic inputs and microglia-neuron surface contact

The number of axonal terminals perforating PNNs was quantified in 33x33x5 µm z-stacks obtained with high-resolution confocal microscopy using the LSM 710 microscope (Zeiss, 100x alpha Plan-Apochromat objective, NA 1.46, voxel size 60x60x500 nm). ROIs were positioned in the motor cortex L5 containing a single PNN coated neuron. Synaptic terminals expressing VGAT or VGLUT1 that associated with WFA labeling were counted using an automated ImageJ routine (see Supplementary Code 1).

The surface of microglia-neuron contacts was quantified in 75x75x10 µm z-stacks obtained with high-resolution confocal microscopy using the LSM 710 microscope (Zeiss, 63x alpha Plan-Apochromat objective, NA 1.4, voxel size 70x70x450 nm). ROIs were positioned in the motor cortex L5 containing a single PNN coated neuron. Surfaces

representing IBA1 (microglia) and Kv3.1 (fast-spiking interneurons) labeled cells were generated by automated thresholding with IMARIS 9.9 software (Oxford Instruments, Stockholm, Sweden) using the standard Surfaces function. Area of contact between cells was quantified as the intersection between IBA1 and Kv3.1 surfaces.

Microscopy techniques and resolution measurements

The morphology of WFA-labeled PNNs was analyzed using the four 3D imaging methods: two-photon excitation (2P), confocal, superresolution structured illumination (SR-SIM), and 3D stimulated emission depletion (STED) microscopy.

2P microscopy of PNNs labeled with Atto 495 was performed using a Leica TCS SP8 (25x HCX IRAPO L 25x water immersion objective, NA 0.95) microscope. ROIs (92x92x11.5 µm, voxel size 90x90x480 nm) were scanned as z-stacks. Excitation laser (Titanium:Sapphire Chameleon Vision II) was tuned to 940 nm, and the output power was 1.712 W. Emitted fluorescence (500 – 570 nm detection wavelength) was collected using a hybrid detector. Imaging resolution estimated by measuring full width at half maximum (FWHM) of Ø 100 nm TetraSpeck microspheres (Cat# T7279, Thermo Fisher Scientific) using the same microscope settings as for PNN imaging.

Confocal microscopy of PNNs labeled with Atto 590 was performed using a LSM 710 (100x alpha Plan-Apochromat objective, NA 1.46) microscope. ROIs (65.4x65.4x5.5 µm, voxel size 60x60x350 nm) were scanned as z-stacks. Excitation laser (561 nm DPSS) was used at 10% of maximum output power to reduce photobleaching. Confocal pinhole size (585 – 655 nm detection wavelength) was adjusted to 1 Airy unit. Imaging resolution was estimated by measuring FWHM of Ø 100 nm TetraSpeck microspheres (Cat# T7279, Thermo Fisher Scientific) using the same microscope settings as for PNN imaging.

SR-SIM microscopy of PNNs labeled with Atto 590 or Abberior Star RED was performed using a Carl Zeiss Elyra PS.1 (100x alpha Plan-Apochromat objective, NA 1.46) microscope. ROIs (49.4x49.4x6.7 µm, voxel size 20x20x100 nm) were scanned as z-stacks. Excitation lasers (561 and 642 nm OPSL) were used at 5% of maximum output power to reduce photobleaching. We used the 51 µm grid (5 rotations, 5 phases) for 585

– 655 nm and 649 – 755 nm detection wavelengths. The output superresolution images
were computed using the automatic processing mode using the Zen Black software
(Zeiss). Imaging resolution was estimated by measuring FWHM of Ø 100 nm TetraSpeck
microspheres (Cat# T7279, Thermo Fisher Scientific) using the same microscope settings
as for PNN imaging.

STED microscopy of PNNs labeled with Abberior Star RED was performed using
the custom-built setup [61] at the Max Planck Institute for Multidisciplinary Sciences in
Göttingen, Germany. We used an oil immersion objective (HCX PL APO 100x/1.40 OIL
STED, Leica Microsystems). ROIs (25x25x5 µm, voxel size 30x30x100 nm) were
scanned as z-stacks. Pixel dwell time was 10 µs. Pulsed excitation light of 650 nm was
spectrally filtered from a white light source [61] and applied with an average power of 2.7
µW at the back focal plane of the objective. The STED laser (Katana 08 HP, OneFive
GmbH, Regensdorf, Swiss), providing nanosecond pulses at 775 nm, was employed with
a power of 221 mW. The STED beam was shaped by a spatial light modulator (Abberior
Instruments) with a 2π-vortex and π phase delay for x/y- and z-depletion, respectively.
The x/y- lateral and z-axial resolution was independently adjusted and estimated by
measuring the FWHM of Ø 40 nm TransFluoSpheres (Ex/Em 633/720 nm, Cat# T8870,
Thermo Fisher Scientific) using the same microscope settings as for PNN imaging.

Reconstruction and analysis of PNN morphology

Ultrastructural organization of PNNs was visualized by superresolution microscopy
and their morphology was analyzed using our in-house quantitative approach based on
graph-based computational reconstruction [21]. The recent optimized MATLAB code is
provided here as Supplementary Code 2. In each animal, a total of 8 individual PNNs
were analyzed in the ipsilesional and contralesional motor cortex. In brief, the 3D image
stacks obtained by superstition STED or SR-SIM microscopy were imported into the
IMARIS 9.9 program (Oxford Instruments) and local fluorescence intensity maxima were
defined as net nodes representing PNN mesh vertices. The internode connections
(edges) were reconstructed using the non-redundant nearest neighbor search algorithm
in MATLAB. The resulting graphs were used to derive morphological metrics that
characterize ultrastructure of PNN facets. More specifically, we quantified average

internode distances (*L*) indicating PNN facet size and the average node degrees (*D*)

reflecting facet density.

Statistical analysis

All quantitative data was presented as box plots indicating the mean (empty

square), the median (line), 25-75 % range (borders) and SD (whiskers) of data

distribution. For all datasets, the normality of distribution was analyzed using the

Kolmogorov-Smirnov test. The differences between groups were evaluated using one-

way (infarct volume and brain atrophy) or two-way (all other readouts) analysis of variance

(ANOVA) and post-hoc pairwise t-tests. For multiple comparisons, Bonferroni corrections

were applied.

Results

Inflammatory preconditioning reduces the infarct size but not delayed brain atrophy

We investigated post stroke brain remodeling in mice with and without stroke

tolerance induced by inflammatory preconditioning. Focal cerebral ischemia was induced

by transient left sided intraluminal occlusion of middle cerebral artery (tMCAO) for 30

minutes resulting in highly reproducible ischemic lesions located in striatum and adjacent

cortical areas, but not motor cortex. Inflammatory preconditioning was performed by

injecting 1 mg/kg LPS intraperitoneally 3 days before tMCAO (Fig. 1A), which triggered

robust peripheral immune response that we reported previously [52]. In agreement with

previous studies [41, 51], the inflammatory preconditioning significantly reduced infarct

volume at 7 days post ischemia (DPI), as indicated by Nissl staining quantifications (Fig

1B, C). However, preconditioning with LPS did not attenuate the delayed brain atrophy at

42 DPI (Fig 1B, D).

Inflammatory preconditioning alters glial responses post stroke

Post-stroke reactive gliosis was evaluated using IBA1 and GFAP

immunohistochemistry detecting microglia/macrophages and reactive astrocytes,

correspondingly (Fig 1E, G). Focal cerebral ischemia triggered microglia/macrophage

activation not only in lesion-associated areas, but also in the ipsilesional motor cortex L5,

as indicated by IBA1 immunoreactivity quantification at 7 DPI (Fig. 1F). Microglia/macrophage activation persisted at 42 DPI and was also observed in the contralesional hemisphere. Inflammatory preconditioning increased IBA1 immunoreactivity at 42 DPI, but not 7 DPI. In contrast, astrocytic reactivity at 7 DPI was reduced at 7DPI in the group exposed to both preconditioning and stroke, as indicated by GFAP immunoreactivity quantification (Fig. 1H).

Interneurons lose PNNs partially and transiently after stroke

Cortical L5 fast spiking interneurons are critical for oscillatory activity in the motor cortex, pyramidal tract output activity regulation, and motor control [23]. These cells express parvalbumin, potassium channels Kv3.1, and are coated with a pattern of aggregated extracellular matrix forming PNNs [34, 54]. PNNs regulate several types of neuroplasticity [8, 49, 53, 57], but their role in post stroke brain remodeling remains under investigated.

We analyzed the expression of PNNs around motor cortical L5 interneurons (regions of interest are outlined in Fig 1E, G) using the *Wisteria floribunda* agglutinin (WFA) that binds glycan chains of extracellular proteoglycans enriched in PNNs [17, 33]. Co-labelling of parvalbumin (PV) and Kv3.1 with WFA (Fig 2A, C) indicated that stroke reduced the expression of PNNs in the ipsilesional motor cortex L5 at 7 DPI (Supplementary Fig S1A). In healthy brains, 73.4±5.5% PV[+] and 93.5±2.1% Kv3.1[+] expressed PNNs, as evidenced by PV[+]/PNN[+] (Fig 2B) and Kv3.1[+]/PNN[+] (Fig 2D) cell quantifications. In mice exposed to stroke only, on average 34.3% PV[+] and 15.3% Kv3.1[+] neurons in the ipsilesional motor cortex lost PNN coatings during the first week post stroke. PNN expression was completely restored by 42 DPI. Inflammatory preconditioning reduced the transient loss of PNNs around PV[+] neurons at 7 DP, compared with the stroke only group (Fig 2B). In all timepoints and conditions evaluated, the majority of PNN[+] cells were also PV[+] (96%) and Kv3.1[+] (97%), and stroke altered neither PV[+] nor Kv3.1[+] cell density, as shown in Supplementary Fig S1.

1
2 **Figure 1. Brain damage and reactive gliosis induced by focal cerebral ischemia.**
3 (**A**) Timeline and experimental endpoints. (**B**) Cresyl violet (Nissl) staining shows focal infarcts at
4 7 DPI, quantified in (**C**), and brain atrophy at 42 DPI, quantified in (**D**). (**E**) Representative WFA

(green) and IBA1 (red) staining at 7 and 42 DPI. (**F**) IBA1 immunoreactivity in the motor cortex
L5. (**G**) Representative WFA (green) and GFAP (red) staining at 7 and 42 DPI. (**H**) GFAP
immunoreactivity in the motor cortex L5. (**E**, **G**) White squares outline 600x600 μm regions of
interest selected for analysis. Nuclei (blue) were stained with DAPI. (**C**, **D**, **F**, **H**) Graphs are box
plots with data as dots, means as squares, medians as lines, interquartile ranges as boxes and
whiskers showing SD. Asterisks and hashes denote significant differences with the control and
stroke groups, correspondingly, as indicated by two-way ANOVA and t-tests (*,#$p < 0.05$, **,##$p <$
0.01, ***$p < 0.001$), n = 7. Scale bars, 1 mm. DPI, days post ischemia; ns, not significant.

Figure 2. PNN expression in the motor cortex L5 post stroke.
Representative immunolabeling of WFA (green) and (**A**) PV (red) or (**C**) Kv3.1 (red) shows PNN
expression around motor cortical L5 interneurons. (**B**, **D**) Percentage of PV (**B**) and Kv3.1 (**D**)
expressing neurons coated with PNNs. Graphs are bar plots showing mean±SD and data as dots.
Asterisks and hashes denote significant differences with the control and stroke groups,
correspondingly, as indicated by two-way ANOVA and t-tests (#$p < 0.05$, **$p < 0.01$, ***$p < 0.001$),
n = 7. Scale bars, 100 μm. DPI, days post ischemia; PV, parvalbumin.

PNN morphology analysis requires superresolution imaging

PNNs are exceptionally stable mesh-like structures in the extracellular space
(ECS) consisting of densely packed proteoglycans bound together by link proteins [9, 17].
Chondroitin sulfate carrying proteoglycans of PNNs repel axons [4, 60], and the
perisomatic synapses on fast-spiking interneurons are established within the PNN facets
forming synaptic pockets, which can be visualized with confocal microscopy (Fig. 3A, B).

After stroke, PNNs undergo morphological changes beyond all-or-none
degradation, and the understanding of their ultrastructural remodeling requires three-
dimensional superresolution imaging [21, 54]. We analyzed the morphology of PNNs
using our in-house quantitative approach combining 3D superresolution fluorescence
imaging and graph-based computational reconstruction. In brief, local fluorescence
intensity maxima of PNN mesh vertices were defined as net nodes, and the internode
connections (edges) were reconstructed using the non-redundant nearest neighbor
search algorithm. Graphs thus obtained reflect the morphology of PNNs and allow for the
quantitative analysis of their ultrastructure. In comparison to our previous report [21], we
significantly improved imaging resolution by adjusting the refractive index of stained
tissues (also known as tissue clearing), using fluorophores with longer emission
wavelengths, and applying additional microscopy techniques.

We compared PNN morphology (Fig. 3C) visualized by four cutting-edge 3D
imaging methods: two-photon (2P) excitation, confocal, superresolution structured
illumination (SR-SIM) and stimulated emission depletion (STED) microscopy. Lateral
(D_{xy}) and axial (D_z) imaging resolution (Fig. 3D) was estimated as full width at half-
maximum (FWHM) using sub-resolution fluorescent beads (Ø 100 nm and Ø 40 nm for
STED) embedded in the stained tissue. 2P microscopy with a small numerical aperture
(NA=0.95) water immersion objective had the lowest resolution (D_{xy} = 443±48 nm, D_z =
1114±242 nm) among the compared methods, resulting in the incomplete PNN graphs
with long internode distances (Fig. 3E, F). Confocal, SR-SIM and 3D STED imaging was
performed using oil immersion objectives with high numerical apertures (NA=1.4 or 1.46).

Figure 3. PNN morphology analysis using confocal, 2P, SIM and STED microscopy.
(**A**, **B**) Confocal microscopy of WFA-labeled neurons shows characteristic facet-like morphology
of PNNs. PNN facets form synaptic pockets (arrowheads), within which presynaptic terminals
contact neuronal surface (dash line) and establish synapses. GABAergic (**A**) and glutamatergic
(**B**) terminals are shown by vesicular GABA (VGAT, red) and glutamate (VGLUT1) transporter
labeling, correspondingly. Arrowheads and dash lines indicate putative synaptic pockets and

neuronal surfaces, correspondingly. Scale bars, 2 µm. (**C**) PNNs in the motor cortex L5 (control brains) were labeled with biotinylated WFA and streptavidin conjugated to Atto 490 (2P microscopy) or Star RED (confocal, SR-SIM and 3D STED microscopy) fluorophores. Images are maximum intensity z-projections. Scale bars, 10 µm. (**D**) Imaging resolution was estimated using sub-resolution microspheres (see Methods). Scale bars, 500 nm. (**E**) PNN morphology was reconstructed as graphs with nodes positioned at local fluorescence intensity maxima and edges generated by non-redundant nearest neighbor search algorithm. (**F**) Histograms show internode distance distributions for the single PNNs shown in (**C**) and (**E**). 2P, two-photon excitation; SR-SIM, superresolution structured illumination microscopy; STED, stimulated emission depletion; D_{xy}, lateral resolution; D_z, axial resolution; ESC, extracellular space.

Lateral resolution of confocal microscopy (D_{xy} = 217±15 nm) was very close to the diffraction limit (d = λ/2NA = 633/2.92 = 216.8 nm), but the relatively low axial resolution (D_z = 757±82 nm) led to the merging of closely spaced PNN nodes, resulting in imperfect graph reconstruction. Both SR-SIM and 3D STED methods had superior resolution that was beyond the diffraction limit in all three dimensions (Fig. 3B). While SR-SIM had slightly better D_z (350±17 versus 369±58 nm), 3D STED significantly outperformed SR-SIM in terms of D_{xy} (75±10 versus 147±12 nm). Of note, D_z and D_{xy} is individually adjustable for STED and was optimized in each sample. Both SR-SIM and 3D STED imaging allowed for precise reconstruction of PNN morphology using graphs, the mathematical constructs designed for topological analysis (Fig. 3C, D).

To compare PNN morphology resolved by SR-SIM and 3D STED, we sequentially visualized the same PNN-coated neuron with the two methods (Fig. 4A, B) and aligned the obtained 3D image stacks based on the intensity maxima positions. Both methods revealed highly similar PNN structures, and both major vertices of the meshes and putative synaptic pockets were found at the same positions (Fig. 4C, D). Regions of interest showing putative synaptic pockets are magnified in Supplementary Fig. S2.

Due to superior lateral resolution, 3D STED microscopy detected multiple smaller vertices in addition to those detected by SR-SIM (Fig. 4D). These intensity maxima were predominantly observed at a distance more than 1 µm from neuronal surface and seldom associated with synaptic pockets. We therefore concluded that these particles represent less condensed interstitial brain matrix and not PNNs. The small particles were filtered for PNN graph reconstruction, based on the cutoff defined by the Gaussian mixture model

showing bimodal distribution of WFA-labeled particle volumes measured by STED

microscopy (Supplementary Fig. S3A, B).

Stroke induces transient loosening of PNNs in both hemispheres

We quantified morphological changes in motor cortical L5 PNNs post stroke by

measuring internode distances and node degrees (that is, number of neighbors

connected to a vertex) of the graphs reconstructing the organization of WFA-labeled

meshes (Fig. 4 E-H). While the average internode distance (L) indicates PNN facet size,

the average node degree (D) reflects facet density.

In mice exposed to stroke only, the size of PNN facets measured with SR-SIM (Fig.

4E) was significantly increased in the ipsilesional motor cortex at 7 DPI (L=573±59 nm)

and 14 DPI (L=581±55 nm), and in the contralesional hemisphere at 14 DPI (L=591±45

nm), compared with the control (L=536±45 nm). At 28 and 42 DPI, PNN facet size

decreased back to control levels. Inflammatory preconditioning minimized the observed

effect, and PNN facet size was not different from the control at 7 and 42 DPI in mice

exposed to preconditioning and stroke. We observed only a minor decrease in the density

of PNN facets (Fig. 4G) at 14 DPI in the stroke group and at 7 DPI in the preconditioning

and stroke group in both ipsilesional and contralesional hemispheres.

The size of PNN facets measured with STED microscopy was significantly

increased in the ipsilesional motor cortex at 7 DPI (L=598±60 nm versus L=545±33 nm

in control) in mice exposed to stroke only (Fig. 4F). At 42 DPI, PNN facet size decreased

back to control levels. In mice exposed to preconditioning and stroke, the facet size was

not different from the control at 7 and 42 DPI, but a minor decrease in the facet density

was observed at 7 DPI in both ipsilesional and contralesional hemispheres (Fig. 4H). In

the interstitial matrix, we observed no alterations induced by stroke (Supplementary Fig.

S3C-E). Therefore, post stroke morphological changes in motor cortical PNNs revealed

by STED microscopy confirmed the results obtained using SR-SIM.

Figure 4. PNN morphology in the motor cortex L5 post stroke.

(**A**, **B**) Maximum intensity z-projections show the morphology of the same PNN sequentially visualized using (**A**) SR-SIM and (**B**) STED microscopy. Scale bars, 5 µm. (**C**, **D**) High magnification single plane images, mesh vertices, and the reconstructed graphs are shown for the same region visualized using (**C**) SR-SIM and (**D**) STED microscopy. Arrowheads and dash lines indicate putative synaptic pockets and neuronal surface, correspondingly. White outlines show PNN vertices, red outlines are interstitial particles. Scale bars, 1 µm. (**E**, **F**) PNN facet size was quantified as average internode distance. (**G**, **H**) PNN facet density was quantified as average node degree. Graphs are box plots with data as dots, means as squares, medians as lines, interquartile ranges as boxes and SD as whiskers. Asterisks and hashes denote significant differences with the control and stroke groups, correspondingly, as indicated by two-way ANOVA and t-tests (*,#p < 0.05, **p < 0.01, ***p < 0.001), (**E**, **G**) n = 7, (**F**, **H**) n = 5. DPI, days post ischemia; SR-SIM, superresolution structured illumination microscopy; STED, stimulated emission depletion; ESC, extracellular space.

Transient PNN loosening associates with bilateral synaptic remodeling post stoke

Because PNNs have a major impact on synaptic plasticity [26], we further explored whether the transient loosening of PNN facets post stroke (Fig. 5A) associates with synaptic remodeling. The density of GABA- and glutamatergic terminals perforating PNNs was quantified in 33x33x5 µm regions of interest positioned in the motor cortex L5 containing a single PNN coated neuron (Fig. 5B). We quantified the local GABAergic and glutamatergic projections targeting the PNN-coated interneurons by immunolabeling the vesicular transporters of GABA (VGAT) and glutamate (VGLUT1), correspondingly.

In healthy brains, motor cortical L5 interneurons received 75±15 (mean±SD) GABAergic inputs from their network partners, as indicated by the quantifications of VGAT terminals perforating PNNs (Fig. 5C). In mice exposed to stroke only, the number of GABAergic terminals perforating PNNs increased at 7 and 14 DPI (92±26 and 90±15 inputs per cell) in the ipsilesional motor cortex but was decreased at 42 DPI (60±21 inputs per cell). In the contralesional motor cortex, interneurons received more GABAergic inputs at 14 and 28 DPI (86±18 and 89±13 inputs per cell) but the number of VGAT terminals decreased back to control levels at 42 DPI. In mice treated with inflammatory preconditioning, the number of GABAergic terminals decreased at 42 DPI (57±24 inputs per cell) bilaterally but did not differ with the control at 7 DPI.

In healthy brains, motor cortical L5 interneurons received 104±27 (mean±SD) glutamatergic inputs from their local network partners, as indicated by the quantifications

1   of VGLUT1 terminals perforating PNNs (Fig. 5D), and their number was not significantly

2   altered post stoke.

Figure 5. Coherent remodeling of PNNs and their perforating synaptic terminals precedes motor coordination recovery after stroke. (**A**) Representative maximum intensity z-projections show transient loosening of motor cortical PNNs after stroke detected by SR-SIM. Inlets show artistic representations of PNN facets. (**B**) Single-plane confocal images show representative immunolabeling of GABAergic axonal terminals expressing VGAT (red) and glutamatergic axonal terminals expressing VGLUT1 (red). PNNs were labeled with WFA (cyan), nuclei are shown in blue. Scale bars, 5 µm. (**C**, **D**) Quantifications show the number of GABAergic (**C**) and glutamatergic (**D**) terminals perforating PNNs. Graphs are box plots with data as dots, means as squares, medians as lines, interquartile ranges as boxes and whiskers showing SD. Asterisks and hashes denote significant differences with the control and stroke groups, correspondingly, as indicated by two-way ANOVA and t-tests (*,#$p < 0.05$, **$p < 0.01$), n = 7. (**E**) Motor coordination recovery measured by tight rope test. Data are mean±SEM. n = 7. DPI, days post ischemia; ns, not significant.

Motor recovery follows motor cortical tissue remodeling post stroke

We analyzed neurological recovery after stroke using Clark's score for general and focal deficits and tight rope test for measuring motor performance. Neurological deficits (Supplementary Fig S4) manifested in the acute phase and decreased during the first week post stroke. Motor activity and coordination measured by the tight rope test (Fig 5E) were decreased at 7 DPI. Starting 28 DPI, we observed a gradual recovery of coordinated movements. Motor coordination was significantly restored by 42 DPI. We observed no significant influence of immune preconditioning on the motor performance recovery after stroke. These data indicate that the transient loosening of PNNs and synaptic remodeling in the motor cortex L5 precede the recovery of coordinated motor activity post stroke.

Microglia-neuron surface contact increases post stroke

In a healthy brain, microglia cells establish direct contact with neuronal plasma membranes [16] and can promote synaptic plasticity by remodeling extracellular matrix [44]. In the fast-spiking PNN coated interneurons, the direct surface contact with microglia should be difficult because of the inhibitory properties of incorporated proteoglycans [1, 20, 26]. However, the transient loosening of PNNs after stroke (Fig. 5A) may facilitate microglia-interneuron interaction. We explored this possibility by quantifying the surface-to-surface contacts between microglia/macrophages labeled with IBA1 and the fast-spiking neurons expressing Kv3.1 channels (Fig. 6A) in the motor cortex L5 after stroke.

Figure 6. Microglia-interneuron contacts post stroke.

(**A**) Confocal Z-projections show representative IBA1 (red), Kv3.1 (green) and WFA (white) immunolabeling in the motor cortex L5. Yellow arrowheads highlight speckles of WFA-labeled material inside IBA1-labelled cell bodies at 42 DPI. Corresponding 3D rendering of IBA1 and Kv3.1 surfaces is shown below. Scale bars, 5 µm. (**B**) Surface contact between microglia and Kv3.1[+]/PNN[+] cells. (**C**) Surface contact between microglia and Kv3.1[+] neurons with and without PNNs at 7 DPI. Graphs are box plots with data as dots, means as squares, medians as lines, interquartile ranges as boxes and whiskers showing SD. Asterisks and hashes denote significant differences with the control and 7 DPI groups, correspondingly, as indicated by two-way ANOVA and t-tests ([#]$p < 0.05$, ***$p < 0.001$, ****$p < 0.00001$). n = 5. DPI, days post ischemia.

Under all experimental conditions that we investigated, multiple microglia
processes were observed in close proximity to neuronal membranes expressing Kv3.1,
and at least two microglia cell bodies were present within 50 µm radius around the soma
of every Kv3.1$^+$ neuron. While in the healthy brains the direct contacts between microglia
and Kv3.1$^+$/PNN$^+$ neurons were point-like and not numerous, the IBA1/Kv3.1 surface
overlap strongly increased at 7 DPI in both ipsilesional and contralesional hemispheres
(Fig. 6B). At 42 DPI, microglia-interneuron contact surface decreased in comparison to 7
DPI but remained significantly larger than in control. In mice with inflammatory
preconditioning, the IBA1/Kv3.1 surface overlap was increased similar to the stroke only
group.

As demonstrated by 3D surface rendering (Fig. 6A), microglia enwrapped
significant parts of neuronal surfaces at 7 DPI. Interestingly, the extensive contacts
between Kv3.1$^+$ neuronal membranes and microglial processes were observed only on
Kv3.1$^+$/PNN$^+$ cells, indicating the high preference of microglia to contacting PNN-coated
neurons (Fig. 6C). In addition, we observed multiple WFA-labeled speckles inside IBA1-
labeled cells at 42 DPI (Fig. 6A), suggesting that microglia/macrophages can phagocyte
PNN components after stroke.

Discussion

We herein demonstrate that the increased size and reduced density of PNN facets
associate with synaptic reorganization preceding the recovery of motor coordination after
stroke. Noteworthy, morphological changes in motor PNNs revealed by SR-SIM were
confirmed using 3D STED microscopy. The coherent remodeling of PNN ultrastructure
and GABAergic axonal terminals on motor cortical L5 interneurons suggests a novel
mechanism of stroke recovery that involves ECM modulation in both ipsi- and
contralesional hemispheres. During the subacute stroke phase at 7 DPI, PNN loosening
correlates with the increasing number of perforating axonal terminals expressing VGAT,
which agrees with increased GABAergic phasic activity during the first week post stroke
[37]. In the chronic stroke phase at 42 DPI, PNN morphology returns back to norm, but
the number of GABAergic inputs received by motor cortical L5 interneurons is significantly

reduced. We hypothesize that these dynamic changes in motor cortical inhibitory
connectivity arise from the tripartite interaction between PNNs, synapses, and microglia.

PNNs cover the soma, proximal dendrites and initial axonal segments of the fast-
spiking interneurons that express calcium binding protein PV and potassium channels
Kv3.1 with fast activation and deactivation kinetics [34, 42]. By creating facet-like
structures, PNNs compartmentalize neuronal surface and restrict synapse formation to
the areas devoid of inhibitory CSPGs that repel axons [4]. The three-dimensional
organization of PNN facets resembles wells that are approximately 1 μm deep, with
neuronal plasma membrane at the bottom and opening towards the extracellular space.
Because most of these wells are occupied by perforating axonal terminals, we here
propose to call them synaptic pockets. After stroke, the increased size of synaptic pockets
allows for new synapse formation. Recent findings show that synapses continuously
wane and re-emerge *in vivo*, and that GABAergic terminals are especially dynamic with
about 60% of them retracting and returning within a few days [59]. We propose that the
loosening of PNNs creates larger permissive regions supporting GABAergic synapse
plasticity and results in the increased GABAergic input to fast-spiking interneurons at 7
and 14 DPI. These new synapses are not stabilized in the long term however, leading to
the decreased number of VGAT terminals perforating PNNs at 42 DPI.

The removal of new GABAergic synapses that are established during the post-
acute stroke phase is likely mediated by activated microglia. Recent evidence indicates
that microglia cells can selectively sculpt inhibitory connectivity [24] by eliminating
presynaptic terminals using a phagocytic mechanism known as trogocytosis [62]. Here,
we observed that every PNN-coated fast-spiking interneuron in the motor cortex L5 is
always adjacent to a few highly ramified microglia cells. Our data shows that although
increased IBA1 immunoreactivity persists in the chronic stroke phase, the surface-to-
surface contact between interneurons and microglia is more extensive at 7 DPI than at
42 DPI. Interestingly, the surface of microglia-interneuron contacts on PNN+ cells is 30
times larger than on PNN- neurons at 7DPI, indicating the high microglial preference for
enwrapping PNN-coated neurons after stroke.

At 42 DPI, we observed speckles of WFA-labeled substance within IBA1-labeled

cell bodies, suggesting that microglia activation contributes to PNN loosening at 7 DPI.

Notably, activated microglia facilitate the loss of PNNs in rodent models of Alzheimer's

[15] and Huntington's [14] diseases. We suppose that after stroke, microglia attenuate

plasticity inhibiting PNN properties in the early post-acute phase, allowing for the

formation of additional GABAergic synapses on motor cortical L5 interneurons. The

increased inhibitory input to inhibitory interneurons can switch the excitation-inhibition

balance in motor cortical microcircuits towards excitation, thereby promoting motor

coordination recovery that we observed starting 28 DPI. During the chronic stroke phase,

PNNs recover completely and restrict new synapse formation, while the non-stabilized

inputs are removed by microglia. Thus, PNNs and microglia negatively regulate inhibitory

input to inhibitory interneurons, which can limit functional recovery in the chronic stroke

phase.

Admittedly, our hypothesis is based on correlations and needs further validation

using intravital superresolution imaging. Here, we revealed the transient remodeling of

PNNs in the motor cortical L5 after stroke using SR-SIM and confirmed this effect using

STED microscopy. While SR-SIM can be applied for multi-label imaging in relatively large

volumes, it requires sophisticated computational processing that may generate artifacts

[35]. STED microscopy has superior resolution compared to SR-SIM and does not

generate any processing-related artifacts [6]. However, STED microscopy is more

challenging to perform in multi-color modes and commonly uses short-distance objectives

that make 3D imaging in large volumes limited.

In this work, we visualized and quantified synaptic terminals and microglial-neuron

contacts using high-resolution confocal microscopy. Nevertheless, our measurements

relied on immunohistochemical procedures incompatible with intravital imaging. In the

future, the tripartite interaction between PNN, presynaptic terminals and microglia post

stroke can be verified using reversibly switchable fluorescent proteins and multi-label *in*

vivo STED microscopy [63].

The superior resolution of STED microscopy and exceptional photostability of WFA

labeling allowed for the detection of interstitial ECM particles that did not associate with

PNNs. In rodent models of learning and memory formation, interstitial matrix regulates axonal sprouting and synaptic input density, while PNNs control the number of synaptic spines and receptor mobility [25]. In the stroke model we used herein, the interstitial matrix density was not affected, and the dynamic modulation of GABAergic input density on motor cortical L5 interneurons associated with ultrastructural rearrangements in PNNs. This evidence indicates that the role of cortical PNNs in post stroke recovery differs from the memory-related function of hippocampal PNNs.

In spite of the prominent changes in PNN morphology, the number of glutamatergic inputs on the fast-spiking motor cortical L5 interneurons is not affected post stroke. Noteworthy, the glutamatergic terminals that we detected here by VGLUT1 immunolabeling represent the local excitatory input within motor cortical microcircuits and not the thalamocortical afferents [27]. The higher stability of glutamatergic inputs on interneurons post stroke may involve additional mechanisms independent of PNNs [5, 58], which agrees with our recent study showing that the depletion of ECM primarily affects inhibitory and not excitatory synapses [19].

Here, we compared PNN morphology alterations, synaptic remodeling, and reactive gliosis after focal cerebral ischemia in mice with and without induced stroke tolerance. Inflammatory preconditioning via systemic injection of LPS has been shown to reduce the severity of stroke in animal models [41, 55]. Our data indicates that exposure to inflammatory stress before stroke reduces infarct volume, attenuates PNN loosening and prevents synaptic alterations during the subacute stroke phase. In the chronic phase though, GABAergic input is similarly reduced in mice with and without induced stroke tolerance. Inflammatory preconditioning does not reduce brain atrophy and even increases microglial reactivity at 42 DPI. We also observed no significant effect of inflammatory preconditioning on motor recovery in the chronic stroke phase, which calls into question the translational value of this approach for improving long-term recovery after stroke.

Our results suggest that the alternating morphology and integrity of PNNs in the motor cortex can be harnessed for promoting neurological recovery in the chronic stroke phase. While the intrinsic brain remodeling post stroke involves only transient loosening

of PNNs, prolonging the partial PNN degradation during the post-acute period can extend the opening neuroplasticity window into the chronic stroke phase. In addition, modulation of CSPG sulfation is a promising target for improving post stroke rehabilitation. The plasticity-inhibiting properties of CSPGs depend on the ratio between 4-sulfated and 6-sulphated disaccharides [30, 60]. Outgrowing axons avoid 4-sulfated CSPGs, and their selective cleavage with arylsulfatase B has been proposed for promoting nerve regeneration [48]. Conclusively, exploring the possibilities for modulating chemical composition and ultrastructure of cortical PNNs is a novel and promising target for post-stroke neuroplasticity research.

Acknowledgements

The authors sincerely appreciate the grant support by the German Research Foundation (DFG, projects 389030878 and 405358801, to DMH, project 467228103 to ED). For the maintenance of equipment and advise on microscopy applications, the authors are especially thankful to the IMCES (imaging center Essen) staff.

Author contributions

E.D. and D.M.H. designed and planned the study. D.Y. and M.S. performed animal surgeries and coded the experimental groups to blind other experimenters. D.Y. conducted animal behavior tests. E.D., E.T, P.L and B.S. performed immunocytochemistry and SR-SIM imaging. E.D. and K.W. performed STED imaging and analyzed data. E.D. performed widefield, confocal, 2P microscopy and associated analyses. E.D., K.W. and D.M.H. drafted the manuscript. All authors discussed the data and contributed to the final version of the manuscript.

References

Afshari FT, Kwok JC, White L, Fawcett JW (2010) Schwann cell migration is integrin-dependent and inhibited by astrocyte-produced aggrecan. Glia 58: 857-869 Doi 10.1002/glia.20970

Arranz AM, Perkins KL, Irie F, Lewis DP, Hrabe J, Xiao F, Itano N, Kimata K, Hrabetova S, Yamaguchi Y (2014) Hyaluronan deficiency due to Has3 knock-out causes altered neuronal activity and seizures via reduction in brain extracellular space. J Neurosci 34: 6164-6176 Doi 10.1523/JNEUROSCI.3458-13.2014

Balashova A, Pershin V, Zaborskaya O, Tkachenko N, Mironov A, Guryev E, Kurbatov L, Gainullin M, Mukhina I (2019) Enzymatic Digestion of Hyaluronan-Based Brain Extracellular Matrix in vivo Can Induce Seizures in Neonatal Mice. Frontiers in neuroscience 13: 1033 Doi 10.3389/fnins.2019.01033

Becker CG, Becker T (2002) Repellent Guidance of Regenerating Optic Axons by Chondroitin Sulfate Glycosaminoglycans in Zebrafish. The Journal of Neuroscience 22: 842-853 Doi 10.1523/jneurosci.22-03-00842.2002

Benson DL, Huntley GW (2012) Synapse adhesion: a dynamic equilibrium conferring stability and flexibility. Current Opinion in Neurobiology 22: 397-404 Doi https://doi.org/10.1016/j.conb.2011.09.011

Berning S, Willig KI, Steffens H, Dibaj P, Hell SW (2012) Nanoscopy in a living mouse brain. Science 335: 551 Doi 10.1126/science.1215369

Carmichael ST, Wei L, Rovainen CM, Woolsey TA (2001) New patterns of intracortical projections after focal cortical stroke. Neurobiol Dis 8: 910-922 Doi 10.1006/nbdi.2001.0425

Carulli D, Broersen R, de Winter F, Muir EM, Meskovic M, de Waal M, de Vries S, Boele HJ, Canto CB, De Zeeuw CIet al (2020) Cerebellar plasticity and associative memories are controlled by perineuronal nets. Proc Natl Acad Sci U S A 117: 6855-6865 Doi 10.1073/pnas.1916163117

Carulli D, Pizzorusso T, Kwok JC, Putignano E, Poli A, Forostyak S, Andrews MR, Deepa SS, Glant TT, Fawcett JW (2010) Animals lacking link protein have attenuated perineuronal nets and persistent plasticity. Brain 133: 2331-2347 Doi 10.1093/brain/awq145

Carulli D, Rhodes KE, Brown DJ, Bonnert TP, Pollack SJ, Oliver K, Strata P, Fawcett JW (2006) Composition of perineuronal nets in the adult rat cerebellum and the cellular origin of their components. J Comp Neurol 494: 559-577 Doi 10.1002/cne.20822

Chow A, Erisir A, Farb C, Nadal MS, Ozaita A, Lau D, Welker E, Rudy B (1999) K(+) channel expression distinguishes subpopulations of parvalbumin- and somatostatin-containing neocortical interneurons. J Neurosci 19: 9332-9345 Doi 10.1523/jneurosci.19-21-09332.1999

Clark W, Lessov N, Dixon M, Eckenstein F (1997) Monofilament intraluminal middle cerebral artery occlusion in the mouse. Neurological Research 19: 641-648 Doi 10.1080/01616412.1997.11740874

Costantini I, Ghobril J-P, Di Giovanna AP, Mascaro ALA, Silvestri L, Müllenbroich MC, Onofri L, Conti V, Vanzi F, Sacconi Let al (2015) A versatile clearing agent for multi-modal brain imaging. Scientific Reports 5: 9808 Doi 10.1038/srep09808

Crapser JD, Ochaba J, Soni N, Reidling JC, Thompson LM, Green KN (2019) Microglial depletion prevents extracellular matrix changes and striatal volume reduction in a model of Huntington's disease. Brain 143: 266-288 Doi 10.1093/brain/awz363

Crapser JD, Spangenberg EE, Barahona RA, Arreola MA, Hohsfield LA, Green KN (2020) Microglia facilitate loss of perineuronal nets in the Alzheimer's disease brain. EBioMedicine 58: 102919 Doi 10.1016/j.ebiom.2020.102919

Cserep C, Posfai B, Lenart N, Fekete R, Laszlo ZI, Lele Z, Orsolits B, Molnar G, Heindl S, Schwarcz ADet al (2020) Microglia monitor and protect neuronal function through specialized somatic purinergic junctions. Science 367: 528-537 Doi 10.1126/science.aax6752

Deepa SS, Carulli D, Galtrey C, Rhodes K, Fukuda J, Mikami T, Sugahara K, Fawcett JW (2006) Composition of perineuronal net extracellular matrix in rat brain: a different disaccharide composition for the net-associated proteoglycans. J Biol Chem 281: 17789-17800 Doi 10.1074/jbc.M600544200

Dityatev A, Bruckner G, Dityateva G, Grosche J, Kleene R, Schachner M (2007) Activity-dependent formation and functions of chondroitin sulfate-rich extracellular matrix of perineuronal nets. Developmental neurobiology 67: 570-588 Doi 10.1002/dneu.20361

Dzyubenko E, Fleischer M, Manrique-Castano D, Borbor M, Kleinschnitz C, Faissner A, Hermann DM (2021) Inhibitory control in neuronal networks relies on the extracellular matrix integrity. Cell Mol Life Sci 78: 5647-5663 Doi 10.1007/s00018-021-03861-3

Dzyubenko E, Gottschling C, Faissner A (2016) Neuron-Glia Interactions in Neural Plasticity: Contributions of Neural Extracellular Matrix and Perineuronal Nets. Neural Plast 2016: 5214961 Doi 10.1155/2016/5214961

Dzyubenko E, Manrique-Castano D, Kleinschnitz C, Faissner A, Hermann DM (2018) Topological remodeling of cortical perineuronal nets in focal cerebral ischemia and mild hypoperfusion. Matrix Biol 74: 121-132 Doi 10.1016/j.matbio.2018.08.001

Dzyubenko E, Manrique-Castano D, Pillath-Eilers M, Vasileiadou P, Reinhard J, Faissner A, Hermann DM (2022) Tenascin-C restricts reactive astrogliosis in the ischemic brain. Matrix Biol: Doi 10.1016/j.matbio.2022.04.003

Estebanez L, Hoffmann D, Voigt BC, Poulet JFA (2017) Parvalbumin-Expressing GABAergic Neurons in Primary Motor Cortex Signal Reaching. Cell reports 20: 308-318 Doi 10.1016/j.celrep.2017.06.044

Favuzzi E, Huang S, Saldi GA, Binan L, Ibrahim LA, Fernández-Otero M, Cao Y, Zeine A, Sefah A, Zheng Ket al (2021) GABA-receptive microglia selectively sculpt developing inhibitory circuits. Cell 184: 4048-4063.e4032 Doi 10.1016/j.cell.2021.06.018

Fawcett JW, Fyhn M, Jendelova P, Kwok JCF, Ruzicka J, Sorg BA (2022) The extracellular matrix and perineuronal nets in memory. Mol Psychiatry: Doi 10.1038/s41380-022-01634-3

Fawcett JW, Oohashi T, Pizzorusso T (2019) The roles of perineuronal nets and the perinodal extracellular matrix in neuronal function. Nat Rev Neurosci 20: 451-465 Doi 10.1038/s41583-019-0196-3

Fremeau RT, Voglmaier S, Seal RP, Edwards RH (2004) VGLUTs define subsets of excitatory neurons and suggest novel roles for glutamate. Trends in Neurosciences 27: 98-103 Doi https://doi.org/10.1016/j.tins.2003.11.005

Frischknecht R, Heine M, Perrais D, Seidenbecher CI, Choquet D, Gundelfinger ED (2009) Brain extracellular matrix affects AMPA receptor lateral mobility and short-term synaptic plasticity. Nature neuroscience 12: 897-904 Doi 10.1038/nn.2338

Frischknecht R, Seidenbecher CI (2008) The crosstalk of hyaluronan-based extracellular matrix and synapses. Neuron Glia Biol 4: 249-257 Doi 10.1017/S1740925X09990226

Galtrey CM, Fawcett JW (2007) The role of chondroitin sulfate proteoglycans in regeneration and plasticity in the central nervous system. Brain research reviews 54: 1-18 Doi 10.1016/j.brainresrev.2006.09.006

Gherardini L, Gennaro M, Pizzorusso T (2015) Perilesional treatment with chondroitinase ABC and motor training promote functional recovery after stroke in rats. Cereb Cortex 25: 202-212 Doi 10.1093/cercor/bht217

Gogolla N, Caroni P, Luthi A, Herry C (2009) Perineuronal nets protect fear memories from erasure. Science 325: 1258-1261 Doi 10.1126/science.1174146

Haji-Ghassemi O, Gilbert M, Spence J, Schur MJ, Parker MJ, Jenkins ML, Burke JE, van Faassen H, Young NM, Evans SV (2016) Molecular Basis for Recognition of the Cancer Glycobiomarker, LacdiNAc (GalNAc[β1→4]GlcNAc), by Wisteria floribunda Agglutinin. J Biol Chem 291: 24085-24095 Doi 10.1074/jbc.M116.750463

Härtig W, Derouiche A, Welt K, Brauer K, Grosche J, Mäder M, Reichenbach A, Brückner G (1999) Cortical neurons immunoreactive for the potassium channel Kv3.1b subunit are predominantly surrounded by perineuronal nets presumed as a buffering system for cations. Brain Research 842: 15-29 Doi 10.1016/s0006-8993(99)01784-9

Heintzmann R, Huser T (2017) Super-Resolution Structured Illumination Microscopy. Chemical Reviews 117: 13890-13908 Doi 10.1021/acs.chemrev.7b00218

Hermann DM, Chopp M (2012) Promoting brain remodelling and plasticity for stroke recovery: therapeutic promise and potential pitfalls of clinical translation. Lancet Neurol 11: 369-380 Doi 10.1016/s1474-4422(12)70039-x

Hiu T, Farzampour Z, Paz JT, Wang EH, Badgely C, Olson A, Micheva KD, Wang G, Lemmens R, Tran KVet al (2016) Enhanced phasic GABA inhibition during the repair phase of stroke: a novel therapeutic target. Brain 139: 468-480 Doi 10.1093/brain/awv360

Hylin MJ, Orsi SA, Moore AN, Dash PK (2013) Disruption of the perineuronal net in the hippocampus or medial prefrontal cortex impairs fear conditioning. Learn Mem 20: 267-273 Doi 10.1101/lm.030197.112

Kwok JC, Carulli D, Fawcett JW (2010) In vitro modeling of perineuronal nets: hyaluronan synthase and link protein are necessary for their formation and integrity. J Neurochem 114: 1447-1459 Doi 10.1111/j.1471-4159.2010.06878.x

Liu Z, Li Y, Zhang ZG, Cui X, Cui Y, Lu M, Savant-Bhonsale S, Chopp M (2010) Bone marrow stromal cells enhance inter- and intracortical axonal connections after ischemic stroke in adult rats. J Cereb Blood Flow Metab 30: 1288-1295 Doi 10.1038/jcbfm.2010.8

Marsh B, Stevens SL, Packard AEB, Gopalan B, Hunter B, Leung PY, Harrington CA, Stenzel-Poore MP (2009) Systemic Lipopolysaccharide Protects the Brain from Ischemic Injury by Reprogramming the Response of the Brain to Stroke: A Critical Role for IRF3. The Journal of Neuroscience 29: 9839-9849 Doi 10.1523/jneurosci.2496-09.2009

Matsuda Y-T, Miyamoto H, Joho RH, Hensch TK (2021) Kv3.1 channels regulate the rate of critical period plasticity. Neuroscience research 167: 3-10 Doi https://doi.org/10.1016/j.neures.2021.04.003

Naka A, Adesnik H (2016) Inhibitory Circuits in Cortical Layer 5. Front Neural Circuits 10: 35 Doi 10.3389/fncir.2016.00035

Nguyen PT, Dorman LC, Pan S, Vainchtein ID, Han RT, Nakao-Inoue H, Taloma SE, Barron JJ, Molofsky AB, Kheirbek MAet al (2020) Microglial Remodeling of the Extracellular Matrix Promotes Synapse Plasticity. Cell 182: 388-403 e315 Doi 10.1016/j.cell.2020.05.050

Otsuka T, Kawaguchi Y (2021) Pyramidal cell subtype-dependent cortical oscillatory activity regulates motor learning. Communications Biology 4: 495 Doi 10.1038/s42003-021-02010-7

Packer AM, Yuste R (2011) Dense, Unspecific Connectivity of Neocortical Parvalbumin-Positive Interneurons: A Canonical Microcircuit for Inhibition? The Journal of Neuroscience 31: 13260-13271 Doi 10.1523/jneurosci.3131-11.2011

Park J, Phillips JW, Guo J-Z, Martin KA, Hantman AW, Dudman JT (2022) Motor cortical output for skilled forelimb movement is selectively distributed across projection neuron classes. Science Advances 8: eabj5167 Doi doi:10.1126/sciadv.abj5167

Pearson CS, Mencio CP, Barber AC, Martin KR, Geller HM (2018) Identification of a critical sulfation in chondroitin that inhibits axonal regeneration. Elife 7: Doi 10.7554/eLife.37139

Pizzorusso T, Medini P, Berardi N, Chierzi S, Fawcett JW, Maffei L (2002) Reactivation of ocular dominance plasticity in the adult visual cortex. Science 298: 1248-1251 Doi 10.1126/science.1072699

Reitmeir R, Kilic E, Kilic U, Bacigaluppi M, ElAli A, Salani G, Pluchino S, Gassmann M, Hermann DM (2011) Post-acute delivery of erythropoietin induces stroke recovery by promoting perilesional tissue remodelling and contralesional pyramidal tract plasticity. Brain 134: 84-99 Doi 10.1093/brain/awq344

Rosenzweig HL, Lessov NS, Henshall DC, Minami M, Simon RP, Stenzel-Poore MP (2004) Endotoxin Preconditioning Prevents Cellular Inflammatory Response During Ischemic Neuroprotection in Mice. Stroke 35: 2576-2581 Doi doi:10.1161/01.STR.0000143450.04438.ae

Sardari M, Skuljec J, Yin D, Zec K, de Carvalho TS, Albers D, Wang C, Pul R, Popa-Wagner A, Doeppner TRet al (2021) Lipopolysaccharide-induced sepsis-like state compromises post-ischemic neurological recovery, brain tissue survival and remodeling via mechanisms involving microvascular thrombosis and brain T cell infiltration. Brain Behav Immun 91: 627-638 Doi 10.1016/j.bbi.2020.10.015

Shi W, Wei X, Wang X, Du S, Liu W, Song J, Wang Y (2019) Perineuronal nets protect long-term memory by limiting activity-dependent inhibition from parvalbumin interneurons. Proceedings of the National Academy of Sciences 116: 27063-27073 Doi 10.1073/pnas.1902680116

Sigal YM, Bae H, Bogart LJ, Hensch TK, Zhuang X (2019) Structural maturation of cortical perineuronal nets and their perforating synapses revealed by superresolution imaging. Proceedings of the National Academy of Sciences 116: 7071-7076 Doi 10.1073/pnas.1817222116

Stevens SL, Vartanian KB, Stenzel-Poore MP (2014) Reprogramming the Response to Stroke by Preconditioning. Stroke 45: 2527-2531 Doi doi:10.1161/STROKEAHA.114.002879

Su W, Matsumoto S, Sorg B, Sherman LS (2019) Distinct roles for hyaluronan in neural stem cell niches and perineuronal nets. Matrix Biology 78-79: 272-283 Doi https://doi.org/10.1016/j.matbio.2018.01.022

Tewari BP, Chaunsali L, Campbell SL, Patel DC, Goode AE, Sontheimer H (2018) Perineuronal nets decrease membrane capacitance of peritumoral fast spiking interneurons in a model of epilepsy. Nat Commun 9: 4724 Doi 10.1038/s41467-018-07113-0

Van Horn MR, Ruthazer ES (2019) Glial regulation of synapse maturation and stabilization in the developing nervous system. Current Opinion in Neurobiology 54: 113-119 Doi https://doi.org/10.1016/j.conb.2018.10.002

Villa KL, Berry KP, Subramanian J, Cha JW, Oh WC, Kwon HB, Kubota Y, So PT, Nedivi E (2016) Inhibitory Synapses Are Repeatedly Assembled and Removed at Persistent Sites In Vivo. Neuron 89: 756-769 Doi 10.1016/j.neuron.2016.01.010

Wang H, Katagiri Y, McCann TE, Unsworth E, Goldsmith P, Yu ZX, Tan F, Santiago L, Mills EM, Wang Yet al (2008) Chondroitin-4-sulfation negatively regulates axonal guidance and growth. J Cell Sci 121: 3083-3091 Doi 10.1242/jcs.032649

Wegner W, Ilgen P, Gregor C, van Dort J, Mott AC, Steffens H, Willig KI (2017) In vivo mouse and live cell STED microscopy of neuronal actin plasticity using far-red emitting fluorescent proteins. Scientific Reports 7: 11781 Doi 10.1038/s41598-017-11827-4

Weinhard L, di Bartolomei G, Bolasco G, Machado P, Schieber NL, Neniskyte U, Exiga M, Vadisiute A, Raggioli A, Schertel Aet al (2018) Microglia remodel synapses by presynaptic trogocytosis and spine head filopodia induction. Nature Communications 9: 1228 Doi 10.1038/s41467-018-03566-5

Willig KI, Wegner W, Müller A, Clavet-Fournier V, Steffens H (2021) Multi-label in vivo STED microscopy by parallelized switching of reversibly switchable fluorescent proteins. Cell reports 35: 109192 Doi https://doi.org/10.1016/j.celrep.2021.109192

Supplementary Figures

Figure S1. Expression of Kv3.1 and parvalbumin in the motor cortical PNN⁺ neurons.

(**A**) Cell density of neurons expressing PNNs. (**B**) Percentage of PNN⁺ neurons expressing Kv3.1.

(**C**) Percentage of PNN⁺ neurons expressing PV. (**D**, **E**) Cell density of neurons expressing PV

(**D**) and Kv3.1 (**E**). Graphs are bar plots showing mean±SD and data as dots. Asterisks and

hashes denote significant differences with the control group, as indicated by two-way ANOVA and

t-tests (*p < 0.05, **p < 0.01), n = 7. DPI, days post ischemia; PV, parvalbumin.

Figure S2. Ultrastructure of a putative synaptic pocket.
High magnification single plane images, mesh vertices, and the reconstructed graphs are shown
for the same region visualized using (**A**) SR-SIM and (**B**) STED microscopy. Scale bars, 500 nm.

Figure S3. STED microscopy detects both PNN vertices and interstitial matrix particles.
(**A**) Histograms show the distribution of WFA-labeled particle volumes measured with SR-SIM and STED microscopy in control mice (n=5). Bin size, 0.01 μm^3. (**B**) Gaussian mixture model indicates the bimodal distribution of PNN vertex volumes measured by STED microscopy. The 0.03 μm^3 cutoff was chosen to filter the small vertices not associating with PNNs. (**C**) High magnification single plane STED image, PNN vertices, interstitial matrix particles, and the reconstructed interstitial matrix graph are shown. Scale bars, 500 nm. Interparticle distance (**D**) and meshwork density (**E**) quantifications indicate no significant alterations in the interstitial matrix post stroke. Graphs are box plots with data as dots, means as squares, medians as lines, interquartile ranges as boxes and whiskers showing SD. n=5. DPI, days post ischemia.

Figure S4. Neurological deficits post stroke.

(**A**) Clark's general deficits scoring. (**B**) Clark's focal deficits scoring. Data are mean±SEM. n = 7. DPI, days post ischemia.

1 **Supplementary code 1**

```
2  //This ImageJ macros was designed for batch processing and quantification of synaptic
3  inputs in 33x33x5 micrometer image stacks
4
5  name=getTitle();
6  rename("processed");
7  run("Split Channels");

8  selectWindow("C2-processed");
9  run("Enhance Contrast...", "saturated=0.3 normalize equalize process_all use");
10 run("Gaussian Blur...", "sigma=10 stack");
11 run("Auto Threshold", "method=Otsu white stack");

12 selectWindow("C1-processed");
13 run("Gaussian Blur 3D...", "x=1 y=1 z=1");
14 run("Maximum 3D...", "x=1 y=1 z=1");
15 run("Subtract Background...", "rolling=10 stack");
16 run("Auto Threshold", "method=Triangle white stack use_stack_histogram");
17 run("Watershed", "stack");

18 imageCalculator("AND create stack", "C1-processed","C2-processed");
19 selectWindow("Result of C1-processed");
20 run("3D Objects Counter", "threshold=128 slice=5 min.=40 max.=2940179
21 exclude_objects_on_edges surfaces summary");

22 run("Concatenate...", "all_open open");
23 newname="C:/Users/Murgot/Desktop/tempres/"+"processed_"+name;
24 saveAs("Tiff",newname);
25 close();

26
```

1 **Supplementary code 2**

```matlab
% This MatLab code was created by Egor Dzyubenko to quantify the density and
% topology of perineuronal nets
% This code is the modified and improved version of the algorithm from our previous
% work (Dzyubenko et al, 2018)

% Required import variables
% OriginalImageName (filename)
% xyz (positions of PNN mesh vertices in space)
% V (volume of each node), optional, we use it for filtration purposes

[ind,name]=grp2idx(OriginalImageName);
range=1.5; % range should not be smaller than the expected distance between objects
nn=length(name);

%The part below allows to filter the dot objects depending on their size
fltrd=find(V>=0.005); %V>=0.001 & V<0.03    V>=0.03
OriginalImageName=OriginalImageName(fltrd,:);
xyz=xyz(fltrd,:);
V=V(fltrd,:);

%check the number of dots
dotNum=zeros(nn,1);
for ww=1:nn
 pp=length(find(ind==ww));
 dotNum(ww,1)=pp;
end
%find the mesh vertices and whether they should be connected
for n=1:nn % nn is length(name). can use intervals
%preassign matrices for single cells
tic
cellind=find(OriginalImageName==name(n));
nodexyz=xyz(cellind,:);
NodeN=length(cellind);
%find indexes Idx and distances D of nodes within the range
[Idx,D] = rangesearch(nodexyz, nodexyz, range);
neighborD=NaN(NodeN,2);
neighborIdx=zeros(NodeN,2);
conn=zeros(NodeN,NodeN);
dist=NaN(NodeN, NodeN);
%get two neighbors for the node if they exist. their places in the cell are
%always 2nd and 3rd
for k=1:NodeN
    if length(D{k})>1
    neighborD(k,1)=D{k}(2);
    neighborIdx(k,1)=Idx{k}(2);
    end
    if length(D{k})>2
    neighborD(k,2)=D{k}(3);
    neighborIdx(k,2)=Idx{k}(3);
    end
end
%create connections with 2 closest neibors nodes if they exist
for i=1:NodeN
        if any(neighborIdx(i,1))       %first neighbor
        first=neighborIdx(i,1);
```

```matlab
        conn(i,first)=1;
        dist(i,first)=neighborD(i,1);
        conn(first,i)=0;                %remove duplicates
        dist(first,i)=NaN;
        end

        if any(neighborIdx(i,2))        %second neighbor
        second=neighborIdx(i,2);
        conn(i,second)=1;
        dist(i,second)=neighborD(i,2);
        conn(second,i)=0;               %remove duplicate
        dist(second,i)=NaN;
        end
    end

% histogram(dist); %optional

%Construct the graph
[row,col]=find(conn==1);
EdgeTable=table([row,col],'VariableNames',{'EndNodes'});
if ~isempty(EdgeTable)
G = graph(EdgeTable);
%you can optionally plot the graph (not recommended for batch mode)
% figure
% plot(G,'EdgeColor','g','EdgeAlpha',0.4,'NodeColor','none','XData',
nodexyz(:,1),'YData', nodexyz(:,2),'ZData', nodexyz(:,3));
% You can optionally create and analyse the total surface of the PNN
% shp = alphaShape(nodexyz(:,1),nodexyz(:,2),nodexyz(:,3),radiusmax);
% plot(shp);
% Surf=surfaceArea(shp);
%Getting quantification data (feel free to add what you like!)
NodeDeg=degree(G);
NodeTotal=NodeN;
NodeConnDeg=NodeDeg(NodeDeg>1);
PercentNodesInMesh=100*length(NodeConnDeg)/NodeTotal;
MeanDegreeConnectivity=mean(NodeConnDeg);
MeanInternodeDistance=mean(dist,'all','omitnan');
%Set results display
results{n,1}=name(n);
results{n,2}=NodeTotal;
results{n,3}=PercentNodesInMesh;
results{n,4}=MeanDegreeConnectivity;
results{n,5}=MeanInternodeDistance;
toc
end

clearvars -except results V n OriginalImageName range xyz name
end

SummaryTable=array2table(results,
'VariableNames',{'name','NodeTotal','PercentNodesInMesh','MeanDegreeConnectivity','Me
anInternodeDistance',});
```