Integrative dissection of gene regulatory elements at base resolution

Zeyu Chen[1,2,3,*], Nauman Javed[1,2,3,*], Molly Moore[2], Jingyi Wu[1,2,3], Michael Vinyard[2,4,5], Luca Pinello[2,4],

Fadi J. Najm[2,#], Bradley E. Bernstein[1,2,3,#]

[1]Department of Cancer Biology, Dana-Farber Cancer Institute, Boston, MA, USA. [2]Broad Institute of MIT

and Harvard, Cambridge, MA, USA. [3]Department of Cell Biology and Pathology, Harvard Medical

School, Boston, MA, USA. [4]Department of Pathology, Massachusetts General Hospital and Harvard

Medical School. [5]Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA,

USA. *Equal contributions. #Correspondence: fadinajm@broadinstitute.org;

bradley_bernstein@dfci.harvard.edu

Abstract

Although vast numbers of putative gene regulatory elements have been cataloged, the sequence

motifs and individual bases that underlie their functions remain largely unknown. Here we

combine deep learning, epigenetic perturbations and base editing to dissect regulatory

sequences within the exemplar immune locus encoding CD69. Focusing on a differentially

accessible and acetylated upstream enhancer, we find that the complementary strategies

converge on a ~150 base interval as critical for CD69 induction in stimulated Jurkat T cells. We

pinpoint individual cytosine to thymine base edits that markedly reduce element accessibility

and acetylation, with corresponding reduction of CD69 expression. The most potent base edits

may be explained by their effect on binding competition between the transcriptional activator

GATA3 and the repressor BHLHE40. Systematic analysis of GATA and bHLH/Ebox motifs

suggests that interplay between these factors plays a general role in rapid T cell transcriptional

responses. Our study provides a framework for parsing gene regulatory elements in their

endogenous chromatin contexts and identifying operative engineered variants.

Genome-wide maps of chromatin state and transcription factor (TF) binding have nominated more than

a million cell type-specific regulatory elements (REs) in the human genome as potential context-specific

regulators of gene expression[1–4]. A critical next step is to determine their functions and sequence

determinants. Computational tools that predict functional bases and/or gene targets are rapidly evolving,

but require systematic benchmarking against perturbational data[5,6]. CRISPR interference (CRISPRi)[7–11]

with fusions between dCas9 and the KRAB repressor provides a means to suppress a regulatory

element and evaluate consequent transcriptional changes[8–11]. Traditional CRISPR-based genetic

perturbations offer increased resolution[9–13], but may incur variable sequence changes due to

heterogeneity of indels after DNA repair. Recent studies have used base editors[14–16] to evaluate protein

coding sequences at high-throughput[17,18], suggesting the potential of these tools to interrogate

noncoding REs in systematic fashion.

Here,we integrated deep learning[6], CRISPRi, dCas9[19] and base editing[14,16] to parse non-coding

regulatory sequences within the CD69 locus (**Fig 1a**). CD69 encodes a key molecule for T cell signal

transduction[20] and tissue residency[21]. CD69 expression is rapidly induced upon stimulation via T cell

receptor cross linking or exposure to PMA/ionomycin in both CD4+ T cells and the Jurkat T cell line (**Fig.**

S1a-S1b). Chromatin accessibility maps revealed several putative regulatory sites that gain accessibility

upon stimulation in primary T cells and Jurkat cells (**Fig. 1b**).   Additionally, we incorporated predictions

from a deep learning model (Enformer[6]) trained on CAGE-seq data[22] and chromatin maps[1]. Genomic

intervals corresponding to the promoter, 3' UTR and an upstream RE were predicted to impact CD69

transcriptional output (**Fig. 1b**).

We used CRISPRi to test the functional impact of these putative REs (**Fig. 1b and Table S1**). We

targeted four REs that gain accessibility upon stimulation by infecting Jurkat cells with lentiviral

constructs containing KRAB-dCas9[23] and sgRNAs. We selected positive cells, applied PMA/ionomycin

stimulation, and measured CD69 surface protein expression by flow cytometry. We found that sgRNAs

targeting RE-4, located 4 kb upstream of the TSS, had the strongest suppressive effect on CD69

expression in stimulated cells (**Fig. 1c and S1c**), while sgRNAs targeting the TSS-proximal element

RE-3 had a weaker suppressive effect (**Fig. 1c and S1c**). RE-4 corresponds to a DNase hypersensitive

site bound by multiple TFs[1] and is capable of enhancing transcription in luciferase reporter assay[24] or

CRISPR activation[25]. The deep learning model predicts that CD69 expression is primarily impacted by a

~170 bp sequence interval within RE-4 (**Fig. 1d**).

To resolve the functional sequences within these elements and test the Enformer predictions, we

designed a library of 101 sgRNAs that target sequences spanning RE-3 and RE-4 (**Fig. S2a-S2b and**

Table S2). We reasoned that dCas9 without the repressive KRAB domain would specifically occlude

TFs overlapping its target site, and thus affect a narrower interval than the KRAB-dCas9 construct[26]. We

infected Jurkat cells with a pooled lentiviral CRISPR library composed of dCas9 and the 101 sgRNAs,

selected for puromycin resistance and stimulated with PMA/ionomycin for 5 hours. We then isolated

genomic DNA from pre-sorted and sorted CD69- and CD69+ subsets (**Fig. S2c**), and amplified the

sgRNA-cassettes for sequencing. The relative effect of each sgRNA on CD69 expression was

calculated based on its enrichment/depletion in CD69+ relative to CD69- libraries. Multiple sgRNAs

spanning the ~1.7 kb tiled sequence restrained CD69 activation (**Fig. 1e and S2d**).

To assay individual functional bases in CD69 REs, we complemented the dCas9 tiling using a Cytidine

Base Editor (CBE) screen. We infected Jurkat cells with lentiviral constructs containing CBE and the

same pool of 101 sgRNAs as above (**Fig. S2a-S2b, and Table S2**). We stimulated and sorted the cells,

and then sequenced the sgRNA-cassettes from pre-sorted, CD69- and CD69+ subsets (**Fig. S2c**). This

identified multiple sgRNAs that significantly reduced CD69 activation (**Fig. 1f and S2e**). Notably, the

CBE and dCas9 perturbations both pinpointed a ~150 bp interval within RE4 centered at sg#70 as

critical for CD69 expression (**Fig. 2a**; Chr12:9764860-9765010). This experimentally identified interval

closely coincided with the region identified by the deep learning model (**Fig.1d**). The screening assays

also revealed a second high scoring interval centered at sg#48 (**Fig. 2a**; Chr12:9765200-9765310). Both

implicated intervals are over-represented for multiple TF motifs relevant to immune function, including

GATA, bHLH/Ebox, TCF, ETS and STAT (**Fig. 2b**). We also scanned the locus for annotated expression

quantitative trait loci (eQTLs). However, the implicated RE-4 interval is highly conserved evolutionarily,

devoid of natural variation in the human population, and thus invisible to eQTL analysis (**Fig 2c**)[27].

These findings highlight the importance of engineered variants for parsing highly conserved regulatory

sequences.

We next sought to validate incurred base edits and their transcriptional consequences. The top scoring

CBE screen sgRNA, sg#70, is predicted to incur C->T transitions at positions 948 and/or 952 within

RE-4 (chr12: 9,764,948 and chr12: 9,764,952). We infected Jurkat cells with a CBE vector containing

either sg#70 or a control sgRNA (sgCtrl) and measured CD69 by flow cytometry (gating strategy in **Fig.**

S3a). This confirmed that CBE-sg#70 strongly reduced CD69 induction upon stimulation (**Fig 3a and**

S3b). We next amplified and sequenced the target region from genomic DNA isolated from Jurkat cells

infected with CBE-sg#70[28]. In unsorted cells, C-948 was replaced by T on ~57.0% of alleles. The

proportion of C-948 edited alleles was higher in sorted CD69- Jurkat cells (67.0%) and lower in the

CD69+ population (53.6%), consistent with a suppressive effect on CD69 induction (**Fig. 3b**). In contrast,

edits to the other candidate site, C-952, were less frequent (14.4% at baseline, 16.6% in CD69-, 13.9%

in CD69+, **Fig. 3b**). These results confirm the initial observation that the C-948->T edit strongly impacts

transcriptional induction of CD69 in response to stimulation.

We also examined the impact of the C-948 base edit on chromatin accessibility. ATAC-seq profiles

revealed reduced RE-4 accessibility in cells harboring the CBE-sg#70 construct, relative to CBE controls

(**Fig. 3c and S3c**). The reduced accessibility was specific to RE-4, as we did not observe any other

accessibility changes in the CD69 locus or neighboring genomic regions (**Fig. S3d**), nor in the vicinity of

other activation associated genes such as CD28 and NR4A1 (**Fig. S3e**). Hence, the single base

substitution at position C-948 reduces RE-4 accessibility and suppresses CD69 induction in stimulated

Jurkat cells.

We next considered the mechanism that underlies the potent effect of this single base mutation. C-948

directly overlaps a GATA motif predicted by the Enformer model to impact CD69 expression in Jurkat

cells (**Fig. 3d and S3f**). Importantly, the C-948->T edit disrupts a critical position in the GATA motif. The

GATA motif is adjacent to a bHLH/Ebox motif that also scores in the deep learning model. We therefore

sought to identify specific TFs that are dynamically expressed and likely to bind these respective motifs

(**Fig. 3e**). GATA3 is highly expressed in Jurkat cells, up-regulated upon stimulation, and broadly

implicated in T-cell lineage commitment[29]. Among bHLH factors, BHLHE40 and BHLHE22 are both

highly expressed and strongly induced upon stimulation. BHLHE40 in particular has established roles in

T cell differentiation, inflammation and autoimmunity[30].

Whereas GATA3 is generally associated with transcriptional activation in T cells, BHLHE40 is a

transcriptional repressor[30–34]. The close juxtaposition of their cognate motifs suggests that only one

factor - either activator or repressor - could bind the implicated site at a given time. We therefore

hypothesized that the paired motifs constitute a dynamic regulatory switch that contributes to CD69

induction. The potent suppressive effect of the base edit could then be explained by its ability to displace

the GATA3 activator by disrupting its motif, while also favoring BHLHE40 repressor binding by relieving

steric hindrance with the GATA factor.

To investigate this hypothesis, we used ChIP-seq to map GATA3, BHLHE40 and the

enhancer-associated histone acetylation mark H3K27Ac in Jurkat cells. Whereas a strong GATA3

binding peak is evident over RE-4 in stimulated Jurkat cells, binding is lost in CBE-sg#70 infected cells

(**Fig. 3f**). Remarkably, GATA3 loss in the edited cells is accompanied by broader BHLHE40 binding over

RE-4, consistent with a switch in TF binding at the edited site (**Fig. 3f**). H3K27ac signal over RE-4 is also

reduced in the CBE-sg#70 edited cells, providing further support for the model that a switch from

activator to repressor binding suppresses element activity (**Fig. 3f**).

We further tested our model with GATA3 and BHLHE40 loss- and gain-of-function experiments. First,

we confirmed that GATA3 knockout suppressed CD69 induction in stimulated Jurkat cells (**Fig. S4a**).

Next, we investigated GATA3-BHLHE40 antagonism by lentiviral BHLHE40 overexpression. We found

that BHLHE40 overexpression suppressed CD69 induction in both control and CBE-sg#70 edited Jurkat

cells (**Fig. 4a**). However, the magnitude of suppression was greater in the edited cells, potentially due to

a relief from GATA factor competition. In contrast, BHLHE41, the homolog of BHLHE40, has no effect on

CD69 expression (**Fig. S4b**). We also assessed the impact of BHLHE40 overexpression on chromatin

accessibility using ATAC-seq. We found that overexpression reduced RE-4 accessibility, indicating a

direct repressive impact on the element and consistent with our proposed TF switch model (**Fig. 4b**).

Further evidence for the importance of interplay between these TFs emerged in our examination of the

second interval identified in our dCas9 and CBE screens. Remarkably, we found that the top base editor

hit in this interval (sg#48) also incurs a C->T edit that disrupts a GATA motif flanked by a bHLH/Ebox

motif (**Fig. 2a-2b**). Here again, the respective motifs are too close to permit coordinate binding. Hence,

this second hit may also be explained by its impact on competitive binding dynamics between the

GATA3 activator and BHLHE40 repressor. This prompted us to examine whether interplay between

these factors plays a more general role in T cell transcriptional responses. We collated the set of GATA3

bound sites in Jurkat cells that contain a GATA motif and a bHLH/Ebox motif within the corresponding

accessible site. We found that BHLHE40 overexpression reduced the aggregate accessibility of these

sites (**Fig. 4c-4d**), consistent with a global repressive role[35,36]. Overall, 909 (95.4%) of these GATA3

sites with proximate bHLH/Ebox motifs became less accessible upon BHLHE40 overexpression, while

just 44 (4.6%) became more accessible (FDR < 0.2). We also examined the edge-edge distance

between the GATA and bHLH/Ebox motifs (**Fig. 4e**). Sites that were repressed by BHLHE40

overexpression showed a strong enrichment for motif spacing of 0 to 3 bp, consistent with competition

between factors (FDR < 0.05). In contrast, a similar analysis of sites that were not repressed revealed a

preferential spacing of 6 to 9 bp between motifs, consistent with previously reported sites of coordinate

GATA and Ebox factor (e.g., TAL1) binding[37]. These findings suggest that the precise spacing of

competitive or collaborative TF binding motifs can have a substantial impact on regulatory element

dynamics.

Finally, we considered the influence of the interacting TFs on T cell phenotypes. GATA3 is an

established T cell regulator that promotes Th2 over Th1 differentiation[38], and is also implicated in the

maintenance of naive T cells[39]. Although BHLHE40 has also been associated with Th2 responses, we

found that BHLHE40 overexpression downregulated multiple immune gene targets involved in Th2

differentiation (IL4R, IL21R, EGR2, etc.) or naive T cell maintenance (CD248, BACH2, etc.)(**Fig. 4f**).

Gene Set Enrichment Analysis confirmed that BHLHE40 overexpression enriched Th1 response genes

and effector T cell signatures, while depleting genes associated with Th2 responses or naive T cells[40]

(**Fig. S5**). These results are consistent with a general role of BHLHE40 in restraining GATA3 mediated

activation at certain immune loci. Notably, many of the immune loci subject to opposing regulation

contain elements with closely spaced GATA and bHLH/Ebox motifs (0-3 bp), consistent with a specific

impact of competitive TF binding on T cell transcriptional programs and phenotypes (**Fig. 4g-4h**).

In conclusion, we integrated chromatin maps, deep learning, CRISPR interference and base editing to

parse the regulatory sequences that underlie the induction of an exemplar gene in stimulated Jurkat T

cells. Top scoring base edits pinpoint a narrow but critical sequence interval within a CD69 enhancer

and reveal a general role for dynamic interplay between the GATA3 activator and the BHLHE40

repressor in controlling T cell transcriptional responses. Our study and results emphasize the

importance of epigenetic perturbations and artificial sequence variants for characterizing regulatory

sequences, which tend to be highly conserved and may be invisible to methods that rely on natural

genetic variation.

Figure Legends

Figure 1. Integrative analysis of the CD69 regulatory landscape.

a) Gene regulatory landscape characterization by successive functional assays and deep learning.

b) Genomic tracks depict accessibility of the CD69 locus in primary CD4+ T cells and Jurkat cells,

without or with stimulation (PMA/ionomycin). Enformer signal track shows the predicted contribution of

underlying sequence to CD69 expression (magnitude of the model gradient at each position with respect

to CD69 promoter signal, summed over 128 bp bins) in Jurkat cells. Grey bars depict regions with

differential accessibility in stimulated Jurkat cells, relative to resting (FDR=0.2). CRISPRi sgRNA

positions are also indicated. ATAC signal corresponds to reads per genomic content (RPGC).

c) Flow cytometry of CD69 expression in Jurkat cells targeted with the indicated CRISPRi sgRNA

following a stimulation time course. Samples gated from the lentiviral transduced population

(mCherry+).

d) Expanded view of Enformer signal at single base resolution over RE-4, as denoted in panel b.

e) Enrichment/depletion plot of dCas9 sgRNAs in CD69+ Jurkat cells, relative to CD69- cells (y-axis;

Log_2 Odds Ratio of normalized sgRNA reads). sgRNAs along the x-axis according to their 5' starting

position on the positive strand. Each data point represents mean±s.e.m.

f) Enrichment/depletion plot of Cytidine Base Editor (CBE) sgRNAs in CD69+ Jurkat cells, relative to

CD69- cells (as in panel e).

For c,e,f, data represent 2-3 biological independent experiments. A 170 bp region critical for CD69

activation is denoted (d-f, light red).

Figure 2. A critical sequence interval within RE-4 influences CD69 expression.

a) Enrichment/depletion plot of sgRNAs in dCas9 and CBE tiling screens as in Fig 1e/f, limited to the

central portion of RE-4 with sgRNAs shown to scale. Expected C->T edit positions highlighted for

CBE-sgRNAs sg#70 and sg#48 (dashed grey lines).

b) Transcription factor motif locations (grouped by broad motif class) for key immune regulators shown

across the same interval as in panel a (FDR<0.05). Dark grey areas represent overlapping motifs.

c) Zoomed out view of the CD69 locus shows CBE sgRNA depletion (red boxes indicate significantly

depleted sgRNAs), common SNPs (black vertical stripes), eQTLs (blue vertical stripes)[27] and

PhastCon100 conservation score (green stripes).

Figure 3. Top scoring base edits target competitive TF binding sites.

a) Flow cytometry plots of CD69 signal for CBE-sgCtrl and CBE-sg#70 Jurkat cells under resting or

stimulated conditions. Bar plot depicts the proportion of CD69+ cells in CBE-sgCtrl (grey) and

CBE-sg#70 (red) after stimulation. P-value based on unpaired t test, **P<0.01. Data are from 4

independent experiments each with 2-3 technical replicates, mean±s.e.m.

b) Table depicts frequency of incurred base edits in CBE-sg#70 infected Jurkat cells. PCR amplicons

from unsorted, CD69- and CD69+ populations were sequenced by Illumina Nextseq500. Consensus

sequence is shown along with stacked bars that depict the proportions of cytosine and thymine bases in

the sequencing data (numbers indicate percent of alleles with C->T edit). Shaded boxes indicate the

sg#70 target sequence.

c) Chromatin accessibility shown over the CD69 locus for stimulated CBE-sg#70 (red) and CBE-sgCtrl

(grey) Jurkat cells. Bar plot depicts the mean ATAC-seq signal over RE-4 (TMM normalized counts per

million; CPM). P-value based on unpaired t test, *P<0.05. Data are from 3 replicates, mean±s.e.m.

d) Enformer signal (letter height) for the sg#70 target region indicates the predicted impact of each base

on CD69 expression. The sgRNA directly coincides with a GATA motif and a bHLH/E-box motif, and

incurs an edit that disrupts the former (vertical dashed line).

e) Volcano plot depicts gene expression fold-change (x-axis) and significance (y-axis) for TF genes in

stimulated Jurkat cells, relative to resting cells. Labels identify differential GATA (red) and bHLH/Ebox

(blue) family members.

f) Genomic tracks for the CD69 locus depict chromatin accessibility (ATAC), H3K27 acetylation

(H3K27ac), GATA3 binding and BHLHE40 binding in CBE-sgCtrl (grey) and CBE-sg#70 (red) Jurkat

cells. Y-axis represents the $-\log_{10}$(p-value) to input controls.

Jurkat cells in a, b, c and f were stimulated with PMA/ionomycin for 2 hours.

Figure 4. GATA3-BHLHE40 competition impacts global T cell transcriptional responses.

a) Flow cytometry plots of CD69 signal for stimulated Jurkat cells transduced with CBE-sg#70 and a

BHLHE40 overexpression construct (BHLHE40-OE), or with corresponding controls (sgCtrl and Ctrl-LV,

respectively). Bar plot depicts the proportion of CD69+ cells in each condition. P-value based on

unpaired t test, ****P<0.0001. Data are from 3 independent experiments with 2-3 technical replicates,

mean±s.e.m.

b) Chromatin accessibility in the CD69 locus for CBE-sg#70 Jurkat cells transduced with either

BHLHE40 overexpression lentivirus (light blue) or control (grey). Cells were stimulated with

PMA/ionomycin. P-value based on unpaired t test without multiple testing correction, *P<0.05. Bar plot

data are from 2 replicates, mean±s.e.m ATAC-seq signal over RE-4 (TMM normalized CPM).

c) Plot depicts aggregate accessibility (y-axis) for GATA3 bound sites that also harbor bHLH/E-box

motifs (centered on the motifs). Data shown for stimulated Jurkat cells transduced with either BHLHE40

overexpression lentivirus (light blue) or control (grey).

d) For the set of GATA3 bound sites with bHLH/E-box motifs in c, volcano plot depicts fold-change

(x-axis) and significance (y-axis) of chromatin accessibility in Jurkat cells transduced with BHLHE40

overexpression lentivirus, relative to control. Differentially accessible sites (FDR < 0.1) are indicated in

red.

e) For differentially accessible sites in d, histogram shows the number of sites (y-axis) with the indicated

spacing (x-axis) between GATA and bHLH/E-box motifs. Sites are stratified by whether their

accessibility is reduced (red) or increased (grey) in the BHLHE40 overexpressing cells. Sites with

significant peak differential between reduced and increased accessibility (FDR < 0.05) are denoted (*).

f) Heatmap shows differentially expressed genes with BHLHE40 binding in their REs, in BHLHE40

overexpressing Jurkat cells relative to control. Cells were stimulated with PMA/ionomycin.

g-h) Genomic views of the IL21R(f) and CD248(g) loci show ChIP-seq data for H3K27Ac, BHLHE40,

and GATA3 in stimulated Jurkat cells (signal corresponds to P-value enrichment over input).

Accessibility (ATAC) and expression (RNA-seq) are also shown for BHLHE40 overexpressing Jurkat

cells and controls. Sites with combined GATA and bHLH/Ebox motifs are indicated (pink shade).

Jurkat cells in b-f were stimulated with PMA/ionomycin for 2 hours.

Methods

Guide library design and cloning

Pooled libraries for expression of sgRNAs were generated as detailed previously (Joung et al Nature

Protocol 2017). Briefly, DNA oligos were annealed into double stranded fragments with compatible

overhangs and ligated into BsmBI sites into vectors. Vector backbones were CRISPRi+guide puro

(pXPR_066, Broad GPP), lentiCRISPR v2-dCas9 (gift of Thomas Gilmore, Addgene 112233) and

rApobec-nCas9-UGI-puro (pRDA_256, Broad GPP) . Libraries were then transformed by electroporation

into electrocompetent coli (Invitrogen) and spread onto bioassay plates. Bacterial colonies were

harvested and isolated using the Plasmid Plus Midi Kit (Qiagen). Four putative regulatory, ATACseq

accessible regions were identified near the CD69 locus. Peak proximity and acceptable on-target

efficacy scores[41] determined sgRNA selection for the CRISPRi tests. After RE3 and RE4 were identified,

all sgRNAs possible in these peak regions were selected and included for screening with dCas9 and

base editors and can be found in **Table S2**.

Cell culture and stimulation

The Jurkat cell line (ATCC, Clone E6.1, TIB152) was cultured in complete RPMI (RPMI Medium

1640,Gibco, 11875085, 1% Penicillin-Streptomycin, Gibco, 15140122, 10% Heat Inactivate Fetal Bovine

Serum, Peak Serum, 20mM HEPES, Gibco,15630080, 1% Sodium Pyruvate, Gibco, 11360070, and 1%

NEAA,Gibco, 11140050) at a maximum density of 2×10^6 cells/ml in 25 cm or 75 cm cell culture dishes.

Stimulation of Jurkat cells for 2-7 hour experiments was achieved with 50ng/ml Phorbol 12-myristate

13-acetate (PMA, Sigma-Alrich, P8139) and 500ng/ml ionomycin calcium salt from Streptomyces

conglobatus (ionomycin, Sigma-Alrich, I0634).

Cryopreserved CD4[+] T cells isolated from healthy donors were obtained from AllCells. On the day of

stimulation, cells were thawed in RPMI 1640 medium supplemented with 2mM L-glutamine and 50%

FBS, counted and resuspended in TexMACS medium (Miltenyi Biotec) supplemented with 20 IU/mL

human Interleukin-2 (IL-2) and 1% penicillin-streptomycin. Cells were seeded at 1 million cells per well in

a 48-well plate. Cells were either left untreated or stimulated with 10 µL T Cell TransAct™, human

(Miltenyi Biotec) via CD3 and CD28 for 24hrs.

Lentivirus production

293T cells approaching 70-80% confluency in 10 cm cell culture dishes were used for packaging. Cells

were pre-treated with 25 uM chloroquine diphosphate (Millipore Sigma, C6628) in 3 ml of complete

DMEM (Gibco DMEM with 1% Penicillin-Streptomycin and 10% Heat Inactivate Fetal Bovine Serum)

and incubate in the 37°C and 5% CO2 incubator for more than 30 minutes. Lipofectamine 3000

Transfection Reagent (ThermoFisher, L3000001) was used to deliver plasmids into 293T cells. Briefly,

15 ug lentiviral vector plasmid, 15 ug of psPAX2 and 5 ug pMD.G plasmid were vortexed with 40 ul

P3000 reagent in 1.5 ml OptiMEM (ThermoFisher, 31985070). Then 40 ul Lipofectamine was added to

1.5 ml OptiMEM and briefly vortexed. The two OptiMEM solutions were combined and mixed well by

vortexing for 30s and incubated at room temperature for at least 20 minutes. Carefully, the OptiMEM

mixture was added dropwise to 293T cells and incubated in a 37°C and 5% CO_2 incubator for 6 hours.

Media were aspirated and replaced with 5ml of fresh complete RPMI. Lentiviral supernatant was

harvested between 24 hours and 48 hours after transfection.

Lentivirus tranduction

Jurkat cells were resuspended in 1ml media and seeded at a density of 2-5 x 10^5 cells per well of a

12-well plate. Lentiviral supernatant was supplemented with 8 ug/ml polybrene (Sigma-Alrich) added to

the Jurkat cells. The plate was then centrifuged at 2000xg, 32°C for 60mins. Cells were then incubated

at 37°C and 5% CO_2 overnight and changed into complete RPMI on the next day. For GFP+ or

mCherry+ marked lentivirus, cells were sorted or analyzed 5 days after transfection via flow cytometry.

For blasticidin selection, 5 ug/ml of blasticidin was added to the transduced cells and selected for 14

days. For puromycin selection, 5 ug/ml of puromycin was added to the transduced cells and selected for

3 days.

Flow cytometry and sorting

Suspended cells were centrifuged down at 300xg, room temperature for 5 minutes. The cells were

stained with the antibody cocktail in the staining buffer of a 1:1 mix of PBS and Brilliant Staining Buffer

(BD, 566349), at room temperature for 20 mins or at 4°C for 30-40 mins. Cells were washed once in

PBS with 1% FBS and then resuspended in the same buffer. Flow cytometry or FACS was processed on

either BD LSRFortessa X-20 or SONY SH800 following the manufacturing instructions. Antibodies and

dyes used from BD: Brilliant Violet 510™ anti-human CD69 Antibody (310936)；APC anti-human CD69

Antibody (310910) Zombie NIR™ Fixable Viability Kit (423106).

At least 2 x 10^5 CRISPR library infected Jurkat cells were collected as a pre-sorted baseline. 2-4 x 10^6

CRISPR library infected Jurkat cells were resuspended in 2 ml of complete RPMI and stimulated with 50

ng/ml PMA and 500 ng/ml ionomycin for 5 hours, and then processed for FACS as described above.

Sorted CD69- and CD69+ populations were collected for genomic DNA isolation.

Genomic DNA isolation and sequencing

Genomic DNA (gDNA) was isolated using QIAamp DNA Micro Kit (QIAGEN, 6304) or DNeasy Blood

and Tissue Kit (QIAGEN, 69504) according to the manufacturer's protocol. The gDNA concentrations

were quantified by Qubit. For PCR amplification, at least 330 ng of gDNA was used per reaction for

greater than 500-fold library coverage. Each reaction contained 1.5 ul Titanium Taq (Takara), 10 µl of

10× Titanium Taq buffer, 8 µl deoxyribonucleotide triphosphate provided with the enzyme, 5 µl DMSO,

0.5 µl P5 stagger primer mix (stock at 100 µM concentration), 10 µl of a uniquely barcoded P7 primer

(stock at 5 µM concentration), and water up to 100ul.

P5 Primer: AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCT TCCGATCT

TTGTGGAAAGGACGAAACACCG

P7 Primer: CAAGCAGAAGACGGCATACGAGATNNNNNNNNGTGACTGGAGTTCAGAC

GTGTGCTCTTCCGATCTCCAATTCCCACTCCTTTCAAGACCT

PCR cycling conditions included: an initial 5 min at 95°C; followed by 30 s at 54°C, 30 s at 53°C, 20 s at

72°C, for 28 cycles; and a final 10-min extension at 72°C. PCR primers were synthesized at Integrated

DNA Technologies. PCR products were purified with Agencourt AMPure XP SPRI beads according to

the manufacturer's instructions (Beckman Coulter, A63880). Samples were sequenced on a MiSeq

(Illumina). Reads were counted by alignment to a reference file of all possible guide RNAs present in the

library. The read was then assigned to a condition on the basis of the 8-nt index included in the P7

primer.

Amplicon sequencing

To assess base editing frequency of the sg#70 locus, we designed primers flanking this region resulting

in a 214bp product. Forward primer: GGTGAGACGTCAGAAAGGAAGT and reverse primer:

AATTCACCCACTGAAAGGAAAA. Amplicons were next ligated with Illumina Truseq adaptors, cleaned

and size selected with AMPure XP SPRI beads, and sequenced on a MiSeq paired end run. FASTQ

files were processed with CRISPResso2 v2 with standard settings for base editor[28].

ATAC-Seq experimental processing

ATAC-Seq lysis buffer contains 10mM Tris-HCl (pH=7.4), 10mM NaCl, 3mM $MgCl_2$, 0.1% Tween-20, 0.1%

NP40, 0.1% Digitonin, 1% BSA and topped up with ddH_2O. ATAC-Seq washing buffer contains 10mM

Tris-HCl (pH=7.4), 10mM NaCl, 3mM MgCl2, 1% BSA and topped up with ddH_2O.

5 x 10^4 cells were centrifuged down with the resuspension buffer (PBS with 1%BSA) in a low-binding

eppendorf tube at 4°C, 500xg for 5 mins. Each pellet is resuspended with 50 ul of lysis buffer and

incubated on ice for 5 minutes. 50 ul of wash buffer was added to the lysis buffer containing nuclei and

centrifuged down at 4°C, 500xg for 5 minutes. The supernatant is then removed and 50ul resuspension

buffer is added to the tube without disturbing the pellet. Nucleus are then centrifuged down at 4°C,

500xg for 5minutes. Tagmentation of the genome DNA is processed using the Illumina tagmentation kit

(20034197) for 30 mins in 37°C. Fragmented products are then isolated via MinElute Reaction

Purification Kit (QIAGEN, 28003) according to the manufacturer's instructions.   Illumina Nextera XT

SetA indexes and NEBNext High-Fidelity 2X PCR Master mix (NEB, M0541) are used to amplify the

fragmented products of each sample, with 12 PCR cycles of 98°C-10s, 63°C-30s and 72°C-1min. PCR

products are then isolated via MinElute PCR Purification Kit (QIAGEN, 28004) following manufacturer's

instructions.

RNA-Seq experimental processing

Whole RNA was extracted from over 1 × 10^5 cells using the QIAGEN RNeasy Micro kit (QIAGEN,

74004) according to the manufacturer's instructions. 1ug RNA was then used to prepare the RNA-Seq

library. Poly-A+ RNA is enriched using Dynabeads mRNA Direct Kit (ThermoFisher, 610.12) according

to the manufacturer's instructions and eluted in 18ul Tris-HCl buffer(pH=7.4). Zinc fragmentation are

processed using RNA Fragmentation Reagents(ThermoFisher, AM8740), followed by Turbo DNase

(ThermoFisher, AM2238) and FastAP enzyme (EF0651) treatment. Then the fragmented RNA are

cleaned-up using Dynabeads MyOne Silane (ThermoFisher, 37002D) and eluted in 7ul of nuclease-free

water. Next, RNA-adaptors are ligased to eluted RNA using T4 RNA ligase (NEB, M0204L) at 23°C for 1

hour and adaptor-ligated RNA was cleaned-up using Dynabeads MyOne Silane (ThermoFisher, 37002D)

and eluted in 13.5ul of nuclease-free water. First strand of cDNA is synthesized using AffinityScript RT

Enzyme (Agilent, 600107) according to the manufacturer's instructions at 54°C for 1 hour. First-strand

cDNA was cleaned-up using Dynabeads MyOne Silane (ThermoFisher, 37002D) and eluted in 5.5ul of

nuclease-free water, followed by cDNA adaptor ligation. After another round of clean-up, the

adaptor-ligated cDNA was processed to library PCR amplification using Phusion Master Mix (NEB,

M0531L) with IDT adaptor indexes. The final library was cleaned-up with AMPure XP Beads (Beckman

Coulter, B23318) to a final size around 280bps.

ChIP-seq experimental processing

Jurkat cells were pelleted (2.5X10^7 per sample) and fixed using 1% formaldehyde at 37°C for 10 mins

then quenched by glycine. Samples were next washed with cold PBS+proteinase inhibitor

(ThermoFisher, 78429), resuspended in lysis buffer (1% SDS, 0.25% DOC, 50mM Tris-HCl, pH=7.4),

and incubated on ice for 10 mins. Samples were diluted up to 1ml in eppendorf using ChIP dilution buffer

(0.01% SDS, 150mM NaCl, 0.25% Triton, 50mM Tris-HCl, pH=7.4) and sonicated using a Covaris E220,

with the following settings: 24 mins with 5% duty factor, 140W max power and 200 cycles/burst. Each

sample was then split into 4 eppendorf tubes: 1) 20ul, top up to 200ul for input; 2)180ul, top up to 1ml for

H3K27Ac ChIP (2.5ul, Active Motif, 39133); 3) 400ul, top up to 1ml for GATA3 ChIP (10ul, CST-D13C9,

5852); 4) 400ul, top up to 1ml for BHLHE40 ChIP (10ul, Novus Biological, NB100-1800). The tubes were

incubated overnight at 4°C on a rotator.

On the next day, Protein G beads (ThermoFisher, 10003D) were washed and added to the

antibody-containing suspension and rotated at 4°C for 2 hours. The beads were then washed with

ice-cold RIPA wash buffer: RIPA-500, LiCl, and 10mM Tris-HCl buffer (pH=8.5). The beads were eluted

in a wash buffer (10mM Tris-HCl, pH=8.0, 0.1% SDS, 150mM NaCl, 5mM DTT) and incubated at 65°C

on a shaker for 1 hour. Samples were then treated with RNAse (Roche, 11119915001) at 37°C for 30

mins and then with proteinase K (Invitrogen, 25530-015) at 63°C for 3 hours. AMPure XP Beads

(Beckman Coulter, B23318) were used to purify the DNA fragments from the samples. Eluted fragments

were then processed for DNA end-repair (Epicenter Biotech, ER0720), Klenow A base adding (Klenow

from NEB, M0212L), adaptor ligation (Ligase from NEB, M2200S) and PCR amplification (PFU Ultra II

HS 2x master mix from Agilent, 600850-51) according to manufacturer's protocols. Index primers were

ordered from Integrative DNA Technology. PCR was set up with the following conditions: 2 mins for

95°C; 30 sec at 95°C, 30 sec at 55°C,30 sec at 72°C for 16 cycles; 1 min at 72°C. PCR products were

purified using AMPure XP Beads with a final size of around 300 bps.

Enformer predictions

The published Enformer model was downloaded from https://tfhub.dev/deepmind/enformer/1 and used

as described by the authors at

https://github.com/deepmind/deepmind-research/blob/master/enformer/enformer-usage.ipynb. Briefly,

we calculated the gradient of the model for unstimulated Jurkat T-cells with respect to the predicted

CAGE-seq signal at the CD69 promoter. This was achieved by centering a 393216 bp genomic window

within the CD69 promoter (chr12:9760820-9760903) and computing the gradient for human output head

4831 with respect to output bins 446-450.The absolute value of the gradients were then summed in

128bp bins for coarse grain resolution (**Fig 1**).

For identifying TF motifs using Enformer base importance scores, we used the TFModisco suite [42]. This

tool clusters short stretches of bases using base importance scores from deep learning algorithms to

discover motifs that can then be matched to known databases. First we centered 393216 bp genomic

windows as above at the promoter of each of 2195 genes that were differentially expressed (FDR=0.01,

see RNA-seq processing below) between resting and stimulated Jurkat cells. Then, we computed the

gradient of the model at each base within the window for output head 4831 as above with respect to the

CAGE-seq signal at the promoter, corresponding to bins 446-450.   For each window, we also computed

the model gradient on a dinucleotide shuffled version of the sequence which was averaged across all

genes in order to obtain an empirical null distribution of gradients. In order to reduce computing time, we

extracted model gradients, sequence, and null gradients for the 750 centered bp window centered at

each ATAC-seq peak detected from unstimulated Jurkat cells. Predictions were run in parallel across all

genes simultaneously using a custom WDL/Google Cloud script. Finally, hypothetical contribution

scores at each position within the 750 bp input window were computed as the model gradient

corresponding to each non-reference base. TFmodisco was then used with default settings in order to

identify putative regulatory motifs. Candidate seqlets were then matched to HOCOMOCO v11 motifs[43]

using Tomtom from the MEME-suite V5.4.1[44].

ATAC-seq data processing, differential accessibility analysis, and coverage tracks

All ATAC-seq data were aligned and processed using the ENCODE uniform ATAC-seq processing

pipeline v2.1.3 available at https://github.com/ENCODE-DCC/atac-seq-pipeline. The pipeline was

configured to use default parameters, adapter auto-detection, the bowtie2 aligner [45], and MACS2 [46] for

peak calling. GRCh38 V29 and associated mitochondrial genomes and blacklists were obtained from

https://www.encodeproject.org/references/ENCSR938RZZ/. Differential accessibility analysis was

conducted with the CSAW package v 1.28[47]. Briefly, a consensus set of peaks was obtained from the

union of peaks across all input samples/replicates. Peaks lying within blacklist regions and with low

signal(less than -3 $\log_{10}$CPM) were removed. Reads were counted in 300 bp windows genomewide and

merged to a maximum width of 5 kb. Finally, counts were TMM normalized and significant differentially

accessible peaks were identified based on a genome-wide (for global analysis) or local window (<= 1

Mb) FDR correction.

ATAC-seq tracks in all figures were computed by pooling replicates where applicable using samtools

merge [48] and creating signal tracks using DeepTools bamcoverage [49]. Signal tracks were normalized

using the reads per genomic bin normalization (RPGC) options in DeepTools with the pre-computed

effective genome size for GRCh38 in order to create coverage bigwigs. Average ATAC-seq coverage

profiles over bHLH/E-box sites were obtained by taking genome-wide motif-scans (see motif analysis

below) and filtering to keep bHLH(Ebox) motif sites overlapping the union of ATAC-seq peaks between

the BHLHE40-OE and BHLHE40-WT conditions. DeepTools plotProfile function with a 4kb window

centered at each bHLH(Ebox) motif was then used to compute the mean coverage profile for each

condition.

RNA-seq data processing, differential expression analysis, and gene-set enrichment

RNA-seq datasets were processed using a custom pipeline utilizing fastp v0.23.2([50] for automatic

adapter trimming with default settings for paired-end datasets, and the STAR aligner [51] v2.7.9a with

default GTEx [52] settings obtained from https://github.com/broadinstitute/gtex-pipeline. Gene

quantifications were obtained using Salmon v1.6 [53] and the GENCODE V38 annotation [54] with the

seqBias, gcBias, posBias, and validateMappings flags enabled.

Differential expression analysis between conditions was conducted using DESeq V2 [55] with default

settings. Log fold change values were corrected with the lfcShrink option using the apeglm method. For

BHLHE40-OE gene expression analysis, BHLHE40-OE was compared to wild-type only in the

stimulated case. Unless otherwise stated, significance is based on an FDR=0.05 cutoff.

Heatmap was constructed using the Complex heatmap package v 2.12.0 [56]. Genes were subsetted to

only keep those differentially expressed between BHLHE40-OE and BHLHE40-WT at FDR=0.05, and

those with a nearby(< 50 kb peak-promoter distance) BHLHE40 ChIP-seq peak(see ChIP-seq

processing below).

Gene-set enrichment analysis was conducted using the FGsea package v 1.22.0 [57] and the

ImmuneSigDB gene sets [40]. The product of $-\log_{10}$(p-value) and $\log_2$FoldChange was used as the ranking

metric for input to gene set enrichment. Significant pathways were collapsed using the

collapsePathways function from FGsea with default settings. Normalized enrichment scores for enriched,

down-regulated gene sets (identified as those containing the suffix _DN) were multiplied by -1.

ChIP-seq data processing

ChIP-seq read alignment, quality filtering, duplicate marking and removal, peak calling, signal

generation, and quality-control was conducted using the ENCODE ChIP-seq pipeline v2.1.6 available at

https://github.com/ENCODE-DCC/chip-seq-pipeline2.   GRCh38 V29 and blacklists were obtained from

https://www.encodeproject.org/references/ENCSR938RZZ/. In brief, reads were aligned to the GRCh38

genome using bowtie2(-X2000), filtered to remove poor quality reads(Samtools) and

de-duplicated(Picard MarkDuplicates). Both histone and TF peaks were then called using MACS2[46]. All

datasets were assessed for enrichment quality and replicate concordance using the included ENCODE

ChIP-seq quality control pipeline. H3K27ac ChIP-seq datasets passed all ENCODE QC standards and

were not further processed beyond obtaining signal tracks as described below. For GATA3 ChIP-seq in

the unedited cells (sgCtrl-P258), we noted a small number of peaks and therefore increased the MACS2

q-value cutoff to 0.05 and max number of peaks to 500000 to improve sensitivity. Consensus peak sets

were then obtained using IDR analysis between replicates with an FDR cutoff of 0.05. De-novo motif

discovery using the XSTREME [58] program from the MEME-suite[44] yielded the expected GATA motif

among the top discovered motifs. A final peak set for sgCtrl-P258 was then obtained by retaining peaks

that contained the expected GATA motif, and that overlapped open chromatin regions as defined by

ATAC-seq. For BHLHE40-P258(sgCtrl), replicates showed poor concordance and the replicate(replicate

1) with the greater number of peaks was selected for further analysis. A peak set was obtained replicate

1 was obtained by taking the intersection of peaks between pseudo-replicates for this dataset. Motif

discovery for this peak set yielded the expected e-box motif(CANNTG) among the top recovered motifs.

As with GATA3, only peaks overlapping with ATAC-seq peaks and which contained an E-box/bHLH

motif were retained.   Target genes for each factor were then nominated by computing the set of genes

with an annotated TSS within 50kb of a called peak using bedtools closest.

Motif Analysis

To avoid motif redundancy, we obtained non-redundant motif cluster definitions and the corresponding

PWMs from https://resources.altius.org/~jvierstra/projects/motif-clustering-v2.0beta/ [59]. Motif scan of the

RE-4 region corresponding to chr12: 9764556 - 9765505 was conducted by extracting the regions

genomic sequence using bedtools getfasta [60] and scanned using the MOODs motif scanner v1.9.4.1 [61]

with a p-value cutoff of 0.0001 and background base probabilities of 2.977e-01 2.023e-01 2.023e-01

2.977e-01. We filtered to keep motifs matched with MOODs score > 4, and further clustered motifs

based on whether the motif cluster name contained GATA, bHLH, TCF, ETS, NFKB, NFAT, CREB, or

STAT.

For motif spacing analysis, genome-wide motif scans were obtained from

https://www.vierstra.org/resources/motif_clustering. Each GATA3 ChIP-seq peak(see ChIP-seq analysis)

was intersected with the set of consensus ATAC-seq peaks for both the BHLHE40-OE and WT

stimulated Jurkat samples. For the remaining peaks, the highest MOODs score GATA motif located

within the middle ⅓ of the peak was chosen as the representative GATA motif. The closest bHLH/E-box

motif was then located using the bedtools closest tool with the -t all and -d options enabled. The

motif-motif distance was then computed as the edge-edge distance between the central GATA motif and

the core CANNTG E-box motif(i.e. the sequence GATAGGCACCTG would yield a distance of 2 bp).

Motif pairs were then separated based on whether the ATAC-seq peak showed reduced or increased

accessibility with BHLHE40-OE(FDR=0.2).   Motif spacing enrichment was then computed by obtaining

the number of motif-spacings at each distance from 1 to 50 bp for each group separately.Enrichment

significance was tested based on the SpaMo approach [62]. For a given motif-pair with a maximum

motif-motif distance of r, we assume that the number of sequences exhibiting a motif-motif distance $0 < x$

$< r$ is binomially distributed, with all spacings being equally likely. This means that the probability of a

given pair having a specific motif-spacing, ignoring strand and motif-orientation, is $1 / r$. We choose

restrict $r <= 50$, reasoning that motif pairs exceeding this edge-edge distance are unlikely to represent

meaningful TF interactions, and to reduce the burden of multiple testing. The probability that we will

observe a specific number of motif-pairs *n* out of a total number of motif-pairs *m* exhibiting a specific

motif-spacing *x* by chance alone is then computed as 1.0 - CDF_binomial(n,m,1.0 / r). This p-value is

then corrected for multiple testing using the BH procedure.

Common SNP, eQTL and Conservation Score Analysis

Common SNPs are adapted from Ensembl GRCh38[63], with a cut-off of more than 1% minor allele

frequency. Expression quantitative trait loci(eQTL) are adapted from eQTLGEN[27], only cis-eQTLs are

considered here with FDR< 0.05 (data from https://eqtlgen.org/cis-eqtls.html, gene locus for CD69).

Conservation score are adapted via phastCons100way score(0-1, clear to dark green)[64], sg#70 regions

are marked specifically in the plot.

Data and code accessibility

For Jurkat and CD4+ T cell ATAC-seq datasets, we adapted GSE155555[5] and GSE124867 for

accessibility analysis on the CD69 loci. For the wild-type Jurkat RNA-Seq, we adapted GSE90718. The

ATAC-Seq datasets for 1) CBE+sgCtrl and CBE+sg#70, 2) Ctrl-GFP and BHLHE40-GFP, 3)

CBE+sg#70+Ctrl-GFP and CBE+sg#70+BHLHE40-GFP are generated in this study, the datasets will be

released on GEO at publication. The RNA-Seq datasets for Ctrl-GFP and BHLHE40-GFP are generated

in this study; the datasets will be released on GEO at publication. The ChIP-Seq datasets for H3K27Ac,

GATA3 and BHLHE40 on CBE+sgCtrl and CBE+sg#70 are generated in this study, the datasets will be

released on GEO at publication.

Analysis code and custom scripts will be made available on github at publication time.

Acknowledgment

We thank S. M. Bevil, S. Battaglia, G. Rahme, K. Macias and J. Verga for technical assistance. This

work was supported by funds from the NCI/NIH Director's Fund (DP1CA216873 to B.E.B.), the Gene

Regulation Observatory and the Variant-to-Function Initiative at the Broad Institute. Z.C. is supported by

NCI-CA-234842. L.P. is supported by the National Human Genome Research Institute (NHGRI)

(R35HG010717, UM1HG012010). J.W. is supported by a postdoctoral fellowship from the

Damon-Runyon Cancer Research Foundation. M.E.V. is supported by the National Cancer Institute

(NCI) of the National Institutes of Health under the Ruth L. Kirschstein National Research Service Award

(F31CA257625). B.E.B. is the Richard and Nancy Lubin Family Endowed Chair at the Dana-Farber

Cancer Institute and an American Cancer Society Research Professor.

Author information

579   Z.C., N.J., F.J.N., and B.E.B. conceived the study. Z.C., F.J.N., and B.E.B. designed the experiments.

580   Z.C. and M.M. performed the experiments. M.E.V. and L.P. provided computational assistance. Z.C.,

581   N.J., J.W. analyzed the data. Z.C., N.J., F.J.N., and B.E.B. interpreted the data and wrote the

582   manuscript.

Competing interests

585   B.E.B. declares outside interests in Fulcrum Therapeutics, HiFiBio, Arsenal Biosciences, Cell Signaling

586   Technologies, and Chroma Medicine.

References:

1.   ENCODE Project Consortium *et al.* Expanded encyclopaedias of DNA elements in the human and

mouse genomes. *Nature* **583**, 699–710 (2020).

2.   Stunnenberg, H. G., International Human Epigenome Consortium & Hirst, M. The International

Human Epigenome Consortium: A Blueprint for Scientific Collaboration and Discovery. *Cell* **167**,

1897 (2016).

3.   Wittkopp, P. J. & Kalay, G. Cis-regulatory elements: molecular mechanisms and evolutionary

processes underlying divergence. *Nat. Rev. Genet.* **13**, 59–69 (2011).

4.   Andersson, R. & Sandelin, A. Determinants of enhancer and promoter activities of regulatory
elements. *Nat. Rev. Genet.* **21**, 71–87 (2020).
5.   Nasser, J. *et al.* Genome-wide enhancer maps link risk variants to disease genes. *Nature* **593**, 238–
243 (2021).
6.   Avsec, Ž. *et al.* Effective gene expression prediction from sequence by integrating long-range
interactions. *Nat. Methods* **18**, 1196–1203 (2021).
7.   Gilbert, L. A. *et al.* CRISPR-mediated modular RNA-guided regulation of transcription in eukaryotes.
Cell **154**, 442–451 (2013).
8.   Fulco, C. P. *et al.* Systematic mapping of functional enhancer–promoter connections with CRISPR
interference. *Science* **354**, 769–773 (2016).
9.   Canver, M. C. *et al.* BCL11A enhancer dissection by Cas9-mediated in situ saturating mutagenesis.
Nature **527**, 192–197 (2015).
10. Korkmaz, G. *et al.* Functional genetic screens for enhancer elements in the human genome using
CRISPR-Cas9. *Nat. Biotechnol.* **34**, 192–198 (2016).
11. Sanjana, N. E. *et al.* High-resolution interrogation of functional elements in the noncoding genome.
Science **353**, 1545–1549 (2016).
12. Rajagopal, N. *et al.* High-throughput mapping of regulatory DNA. *Nat. Biotechnol.* **34**, 167–174
(2016).
13. Diao, Y. *et al.* A tiling-deletion-based genetic screen for cis-regulatory element identification in
mammalian cells. *Nat. Methods* **14**, 629–635 (2017).
14. Gaudelli, N. M. *et al.* Programmable base editing of A•T to G•C in genomic DNA without DNA
cleavage. *Nature* **551**, 464–471 (2017).
15. Kim, Y. B. *et al.* Increasing the genome-targeting scope and precision of base editing with
engineered Cas9-cytidine deaminase fusions. *Nat. Biotechnol.* **35**, 371–376 (2017).
16. Komor, A. C., Kim, Y. B., Packer, M. S., Zuris, J. A. & Liu, D. R. Programmable editing of a target
base in genomic DNA without double-stranded DNA cleavage. *Nature* **533**, 420–424 (2016).
17. Cuella-Martin, R. *et al.* Functional interrogation of DNA damage response variants with base editing
screens. *Cell* **184**, 1081–1097.e19 (2021).
18. Hanna, R. E. *et al.* Massively parallel assessment of human variants with base editor screens. *Cell*
184, 1064–1080.e20 (2021).

19. Qi, L. S. *et al.* Repurposing CRISPR as an RNA-guided platform for sequence-specific control of
gene expression. *Cell* **184**, 844 (2021).

20. Cibrián, D. & Sánchez-Madrid, F. CD69: from activation marker to metabolic gatekeeper. *Eur. J.*
Immunol. **47**, 946–953 (2017).

21. Sathaliyawala, T. *et al.* Distribution and compartmentalization of human circulating and
tissue-resident memory T cell subsets. *Immunity* **38**, 187–197 (2013).

22. FANTOM Consortium and the RIKEN PMI and CLST (DGT) *et al.* A promoter-level mammalian
expression atlas. *Nature* **507**, 462–470 (2014).

23. Alerasool, N., Segal, D., Lee, H. & Taipale, M. An efficient KRAB domain for CRISPRi applications
in human cells. *Nat. Methods* **17**, 1093–1096 (2020).

24. Laguna, T. *et al.* New insights on the transcriptional regulation of CD69 gene through a potent
enhancer located in the conserved non-coding sequence 2. *Mol. Immunol.* **66**, 171–179 (2015).

25. Mumbach, M. R. *et al.* Enhancer connectome in primary human cells identifies target genes of
disease-associated DNA elements. *Nature Genetics* vol. 49 1602–1612 (2017).

26. Dominguez, A. A., Lim, W. A. & Qi, L. S. Beyond editing: repurposing CRISPR–Cas9 for precision
genome regulation and interrogation. *Nat. Rev. Mol. Cell Biol.* **17**, 5–15 (2015).

27. Võsa, U. *et al.* Large-scale cis- and trans-eQTL analyses identify thousands of genetic loci and
polygenic scores that regulate blood gene expression. *Nat. Genet.* **53**, 1300–1310 (2021).

28. Clement, K. *et al.* CRISPResso2 provides accurate and rapid genome editing sequence analysis.
Nat. Biotechnol. **37**, 224–226 (2019).

29. Ho, I.-C., Tai, T.-S. & Pai, S.-Y. GATA3 and the T-cell lineage: essential functions before and after
T-helper-2-cell differentiation. *Nat. Rev. Immunol.* **9**, 125–135 (2009).

30. Cook, M. E., Jarjour, N. N., Lin, C.-C. & Edelson, B. T. Transcription Factor Bhlhe40 in Immunity
and Autoimmunity. *Trends Immunol.* **41**, 1023–1036 (2020).

31. Huynh, J. P. *et al.* Bhlhe40 is an essential repressor of IL-10 during Mycobacterium tuberculosis
infection. *J. Exp. Med.* **215**, 1823–1838 (2018).

32. Emming, S. *et al.* A molecular network regulating the proinflammatory phenotype of human memory
T lymphocytes. *Nat. Immunol.* **21**, 388–399 (2020).

33. Asanoma, K. *et al.* Regulation of the Mechanism of TWIST1 Transcription by BHLHE40 and
BHLHE41 in Cancer Cells. *Mol. Cell. Biol.* **35**, 4096–4109 (2015).

34. Zawel, Yu & Torrance. DEC1 is a downstream target of TGF-β with sequence-specific

657   transcriptional repressor activities. *Proc. Estonian Acad. Sci. Biol. Ecol.*

658   35. Azmi, S., Sun, H., Ozog, A. & Taneja, R. mSharp-1/DEC2, a basic helix-loop-helix protein functions

659   as a transcriptional repressor of E box activity and Stra13 expression. *J. Biol. Chem.* **278**, 20098–

660   20109 (2003).

661   36. Sato, F. *et al.* Functional analysis of the basic helix-loop-helix transcription factor DEC1 in circadian

662   regulation. Interaction with BMAL1. *European Journal of Biochemistry* vol. 271 4409–4419 (2004).

663   37. Sanda, T. *et al.* Core transcriptional regulatory circuit controlled by the TAL1 complex in human T

664   cell acute lymphoblastic leukemia. *Cancer Cell* **22**, 209–221 (2012).

665   38. Wan, Y. Y. GATA3: a master of many trades in immune regulation. *Trends Immunol.* **35**, 233–242

666   (2014).

667   39. Singer, M. *et al.* A Distinct Gene Module for Dysfunction Uncoupled from Activation in

668   Tumor-Infiltrating T Cells. *Cell* **171**, 1221–1223 (2017).

669   40. Godec, J. *et al.* Compendium of Immune Signatures Identifies Conserved and Species-Specific

670   Biology in Response to Inflammation. *Immunity* **44**, 194–206 (2016).

671   41. Doench, J. G. *et al.* Optimized sgRNA design to maximize activity and minimize off-target effects of

672   CRISPR-Cas9. *Nat. Biotechnol.* **34**, 184–191 (2016).

673   42. Shrikumar, A., Tian, K. & Avsec, Ž. Technical Note on Transcription Factor Motif Discovery from

674   Importance Scores (TF-MoDISco) version 0.5.6.5. *arXiv* (2018) doi:

675   https://doi.org/10.48550/arXiv.1811.00416.

676   43. Kulakovskiy, I. V. *et al.* HOCOMOCO: towards a complete collection of transcription factor binding

677   models for human and mouse via large-scale ChIP-Seq analysis. *Nucleic Acids Res.* **46**, D252–

678   D259 (2018).

679   44. Bailey, T. L., Johnson, J., Grant, C. E. & Noble, W. S. The MEME Suite. *Nucleic Acids Res.* **43**,

680   W39–49 (2015).

681   45. Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat. Methods* **9**, 357–

682   359 (2012).

683   46. Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol.* **9**, R137 (2008).

684   47. Lun, A. T. L. & Smyth, G. K. csaw: a Bioconductor package for differential binding analysis of

685   ChIP-seq data using sliding windows. *Nucleic Acids Res.* **44**, e45 (2016).

48. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079
(2009).

49. Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data analysis.
Nucleic Acids Res. **44**, W160–5 (2016).

50. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics vol. 34 i884–i890 (2018).

51. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

52. GTEx Consortium. The GTEx Consortium atlas of genetic regulatory effects across human tissues.
Science **369**, 1318–1330 (2020).

53. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and
bias-aware quantification of transcript expression. *Nat. Methods* **14**, 417–419 (2017).

54. Frankish, A. *et al.* GENCODE 2021. *Nucleic Acids Res.* **49**, D916–D923 (2021).

55. Anders, S. & Huber, W. Differential expression analysis for sequence count data. *Genome Biol.* **11**,
R106 (2010).

56. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

57. Korotkevich, G. *et al.* Fast gene set enrichment analysis. *bioRxiv* 060012 (2021)
doi:10.1101/060012.

58. Grant, C. E. & Bailey, T. L. XSTREME: Comprehensive motif analysis of biological sequence
datasets. *bioRxiv* (2021) doi:10.1101/2021.09.02.458722.

59. Vierstra, J. *et al.* Global reference mapping of human transcription factor footprints. *Nature* **583**,
729–736 (2020).

60. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics **26**, 841–842 (2010).

61. Korhonen, J., Martinmäki, P., Pizzi, C., Rastas, P. & Ukkonen, E. MOODS: fast search for position
weight matrix matches in DNA sequences. *Bioinformatics* **25**, 3181–3182 (2009).

62. Whitington, T., Frith, M. C., Johnson, J. & Bailey, T. L. Inferring transcription factor complexes from
ChIP-seq data. *Nucleic Acids Res.* **39**, e98 (2011).

63. Cunningham, F. *et al.* Ensembl 2022. *Nucleic Acids Res.* **50**, D988–D995 (2022).

64. Siepel, A. *et al.* Evolutionarily conserved elements in vertebrate, insect, worm, and yeast genomes.

Genome Res. **15**, 1034–1050 (2005).

Figure 1. Integrative analysis of the CD69 regulatory landscape.

A) Gene regulatory landscape characterization by successive functional assays and deep learning.

B) Genomic tracks depict accessibility of the CD69 locus in primary CD4+ T cells and Jurkat cells, without or with stimulation (PMA/ionomycin). Enformer signal track shows the predicted contribution of underlying sequence to CD69 expression (magnitude of the model gradient at each position with respect to CD69 promoter signal, summed over 128 bp bins)in Jurkat cells. Grey bars depict regions with differential accessibility in stimulated Jurkat cells, relative to resting (FDR=0.2). CRISPRi sgRNA positions are also indicated. ATAC signal corresponds to reads per genomic content (RPGC).

C) Flow cytometry of CD69 expression in Jurkat cells targeted with the indicated CRISPRi sgRNA following a stimulation time course. Samples gated from the lentiviral transduced population (mCherry+).

D) Expanded view of Enformer signal at single base resolution over RE-4, as denoted in panel b.

E) Enrichment/depletion plot of dCas9 sgRNAs in CD69+ Jurkat cells, relative to CD69- cells (y-axis; Log₂ Odds Ratio of normalized sgRNA reads). sgRNAs along the x-axis according to their 5' starting position on the positive strand. Each data point represents mean±s.e.m.

F) Enrichment/depletion plot of Cytidine Base Editor (CBE) sgRNAs in CD69+ Jurkat cells, relative to CD69- cells (as in panel e).

For c,e,f, data represent 2-3 biological independent experiments. A 170 bp region critical for CD69 activation is denoted (d-f, light red).

Figure 2. A critical sequence interval within RE-4 influences CD69 expression.

A) Enrichment/depletion plot of sgRNAs in dCas9 and CBE tiling screens as in Fig 1e/f, limited to the central portion of RE-4 with sgRNAs shown to scale. Expected C->T edit positions highlighted for CBE-sgRNAs sg#70 and sg#48 (dashed grey lines).

B) Transcription factor motif locations (grouped by broad motif class) for key immune regulators shown across the same interval as in panel a (FDR<0.05). Dark grey areas represent overlapping motifs.

C) Zoomed out view of the CD69 locus shows CBE sgRNA depletion (red boxes indicate significantly depleted sgRNAs), common SNPs (black vertical stripes), eQTLs (blue vertical stripes)27 and PhastCon100 conservation score (green stripes).

Figure 3. Top scoring base edits target competitive TF binding sites.
A) Flow cytometry plots of CD69 signal for CBE-sgCtrl and CBE-sg#70 Jurkat cells under resting or stimulated conditions. Bar plot depicts the proportion of CD69+ cells in CBE-sgCtrl (grey) and CBE-sg#70 (red) after stimulation. P-value based on unpaired t test, **P<0.01. Data are from 4 independent experiments each with 2-3 technical replicates, mean±s.e.m.
B) Table depicts frequency of incurred base edits in CBE-sg#70 infected Jurkat cells. PCR amplicons from unsorted, CD69- and CD69+ populations were sequenced by Illumina Nextseq500. Consensus sequence is shown along with stacked bars that depict the proportions of cytosine and thymine bases in the sequencing data (numbers indicate percent of alleles with C->T edit). Shaded boxes indicate the sg#70 target sequence.
C) Chromatin accessibility shown over the CD69 locus for stimulated CBE-sg#70 (red) and CBE-sgCtrl (grey) Jurkat cells. Bar plot depicts the mean ATAC-seq signal over RE-4 (TMM normalized counts per million; CPM). P-value based on unpaired t test, *P<0.05. Data are from 3 replicates, mean±s.e.m.
D) Enformer signal (letter height) for the sg#70 target region indicates the predicted impact of each base on RE-4 accessibility. The sgRNA directly coincides with a GATA motif and a bHLH/E-box motif, and incurs an edit that disrupts the former (vertical dashed line).
E) Volcano plot depicts gene expression fold-change (x-axis) and significance (y-axis) for TF genes in stimulated Jurkat cells, relative to resting cells. Labels identify differential GATA (red) and bHLH/Ebox (blue) family members.
F) Genomic tracks for the CD69 locus depict chromatin accessibility (ATAC), H3K27 acetylation (H3K27ac), GATA3 binding and BHLHE40 binding in CBE-sgCtrl (grey) and CBE-sg#70 (red) Jurkat cells. Y-axis represents the -log10(p-value) to input controls.
Jurkat cells in A, B, C and F were stimulated with PMA/ionomycin for 2 hours.

Figure 4. GATA3-BHLHE40 competition impacts global T cell transcriptional responses.

A) Flow cytometry plots of CD69 signal for stimulated Jurkat cells transduced with CBE-sg#70 and a BHLHE40 overexpression construct (BHLHE40-OE), or with corresponding controls (sgCtrl and Ctrl-LV, respectively). Bar plot depicts the proportion of CD69+ cells in each condition. P-value based on unpaired t test, ****P<0.0001. Data are from 3 independent experiments with 2-3 technical replicates, mean±s.e.m.

B) Chromatin accessibility in the CD69 locus for CBE-sg#70 Jurkat cells transduced with either BHLHE40 overexpression lentivirus (light blue) or control (grey). Cells were stimulated with PMA/ionomycin. P-value based on unpaired t test without multiple testing correction, *P<0.05. Bar plot data are from 2 replicates, mean±s.e.m ATAC-seq signal over RE-4 (TMM normalized CPM).

C) Plot depicts aggregate accessibility (y-axis) for GATA3 bound sites that also harbor bHLH/E-box motifs (centered on the motifs). Data shown for stimulated Jurkat cells transduced with either BHLHE40 overexpression lentivirus (light blue) or control (grey).

D) For the set of GATA3 bound sites with bHLH/E-box motifs in c, volcano plot depicts fold-change (x-axis) and significance (y-axis) of chromatin accessibility in Jurkat cells transduced with BHLHE40 overexpression lentivirus, relative to control. Differentially accessible sites (FDR < 0.1) are indicated in red.

E) For differentially accessible sites in d, histogram shows the number of sites (y-axis) with the indicated spacing (x-axis) between GATA and bHLH/E-box motifs. Sites are stratified by whether their accessibility is reduced (red) or increased (grey) in the BHLHE40 overexpressing cells. Sites with significant peak differential between reduced and increased accessibility (FDR < 0.05) are denoted (*).

F) Heatmap shows differentially expressed genes with BHLHE40 binding in their REs, in BHLHE40 overexpressing Jurkat cells relative to control. Cells were stimulated with PMA/ionomycin.

G-H) Genomic views of the IL21R(F) and CD248(G) loci show ChIP-seq data for H3K27Ac, BHLHE40, and GATA3 in stimulated Jurkat cells (signal corresponds to P-value enrichment over input). Accessibility (ATAC) and expression (RNA-seq) are also shown for BHLHE40 overexpressing Jurkat cells and controls. Sites with combined GATA and bHLH/Ebox motifs are indicated (pink shade).

Jurkat cells in B-F were stimulated with PMA/ionomycin for 2 hours.