Ketone body metabolism declines with age in mice in a sex-dependent manner

Authors:

Brenda Eap[1,3], Mitsunori Nomura[1], Oishika Panda[1], Thelma Y Garcia[1], Christina D King[1], Jacob P Rose[1], Teresa C Leone[4], Daniel P Kelly[4], Birgit Schilling[1], John C Newman[1,2,3]

[1]Buck Institute for Research on Aging, Novato, CA

[2]Division of Geriatrics, University of California, San Francisco, San Francisco, CA

[3]Leonard Davis School of Gerontology, University of Southern California, Los Angeles CA

[4] Cardiovascular Institute, Department of Medicine, Perelman School of Medicine at the University of Pennsylvania, Philadelphia, PA

* Correspondence to: jnewman@buckinstitute.org

Abstract

Understanding how our cells maintain energy homeostasis has long been a focus of aging biology. A decline in energy metabolism is central to many age-related diseases such as Alzheimer's disease, heart failure, frailty, and delirium. Intervening on pathways involved in energy homeostasis can extend healthy lifespan. When the primary energy substrate glucose is scarce, mice and humans can partially switch cellular energetic needs to fat-derived ketone bodies (i.e., beta-hydroxybutyrate (BHB), acetoacetate, acetone). Aging is associated with glucose intolerance and insulin insensitivity, yet, surprisingly, what role ketone body metabolism might play in compensating for impaired glucose utilization in age-related diseases is understudied. Here, we investigate how endogenous ketone body production and utilization pathways are modulated by age across the lifespan of male and female C57BL/6N mice (3 mo old, 12 mo old, 22 mo old). We show how different ages have different metabolic and gene expression responses to 1-week ketogenic diet (KD). We hypothesized that there would be a compensatory ketogenic response with age but instead saw declines in plasma BHB concentrations under fasting and non-fasting conditions with strong sexual dimorphism. Under KD, both sexes increased BHB concentrations at all ages, but only males showed strong gene expression induction. We also observed tissue-specific changes with age in baseline ketone metabolism, and surprising induction of extrahepatic ketogenic genes under KD. We found significant residual blood concentrations of BHB in KD even after a knockout of liver BHB production. Overall, these findings show that older mice have impaired non-fasting ketogenesis but are capable of increasing their ketogenic capacity under stimulation (i.e., KD) to meet energetic demands in aging.

38 Therapies to augment non-fasting ketogenesis or provide exogenous ketones may be

39 useful to improve energy homeostasis in diseases and conditions of aging.

Introduction

Aging is associated with glucose intolerance and insulin insensitivity, resulting in a deregulation of the nutrient-responsive pathway, a common hallmark of aging[1–3]. While lifestyle interventions like fasting, caloric restriction and dietary restriction can extend healthy mammalian lifespan, impactful translation to improve human health requires a deeper mechanistic understanding of these interventions. For examples, IGF-1 signaling and the mammalian target of rapamycin (mTOR) pathways are promising translational targets with drugs now in clinical trials (e.g., NCT02432287, NCT03733132, NCT01649960, and NCT05414292)[4–7]. One of the commonly observed metabolic change associated with these interventions across species is the elevation of blood ketone bodies. Similar to the discoveries of rapamycin and metformin, endogenous metabolites like ketone bodies have also been shown to prevent the deregulation of the nutrient-responsive pathway[8–13].

Ketone bodies are synthesized in the liver from non-esterified fatty acids (NEFAs) mobilized from adipose tissue. Ketone bodies, of which beta-hydroxybutyrate (BHB) is most abundant, are commonly known as glucose-sparing energy molecules for extra-hepatic tissues such as brain, heart, kidney, and skeletal muscle[14]. In healthy adults, ketosis under non-fasting conditions maintains plasma BHB concentrations to a range of 50 to 250 μM[15]. However, a higher non-fasting ketosis can be achieved through fasting, exercise, and dietary interventions like a ketogenic diet, increasing concentrations above 500 μM and into the low millimolar range[16–22]. In addition to its role as an alternative oxidative fuel, BHB and acetoacetate are increasingly understood to have a variety of signaling activities also relevant to aging. BHB specifically has been

shown to play important roles in health and disease. In brain aging, two studies showed

that feeding a non-obese KD could extend the healthy lifespan of wild-type mice

including improving memory function[20,21]. In humans, ketone body uptake differs in

normal brain aging, mild cognitive impairment, and Alzheimer's Disease[23]. KD,

intermittent fasting, and ketogenic pharmacotherapies like SGLT2i have shown to

protect against diabetic kidney disease[24]. In the failing heart, it's been shown that the

heart prefers ketone bodies as more energy efficient substrates, which helps explain

why patients with heart failure have elevated concentrations of ketone bodies[25–27]. BHB

has also shown to regulate inflammation by inhibiting the NLRP3 inflammasome,

reducing proinflammatory IL-1B, IL-18, and caspase-1 activation, in heart, kidney and

the central nervous system[28,29].

This growing literature suggests that ketogenic interventions have promise for

various diseases and conditions of aging, but there are significant gaps. Much of the

work in aging was carried out in male mice only, despite clear metabolic differences

between males and females in glucose and fat metabolism[30,31]. Work to date has

focused on interventions, while how the baseline endogenous ketogenic system

changes with age is not understood. And little is known about how the response of the

endogenous system to interventions like KD changes with age. Our most recent work

showed an improvement in memory of aging male mice, leading us to further elucidate

the effects of KD across ages and sexes[20,21]. From the same work, we showed that

ketogenesis is tightly regulated by the peroxisome proliferator activated receptor alpha

(PPARα). Therefore, we sought to understand the regulation of ketone body metabolism

across ages, sexes, and diets to investigate the hypothesis that ketogenesis is

compensatory to aging and is tissue-specific to hypometabolism of glucose we

commonly see with age. In this study, we first test the different responses of male and

female C57BL/6 mice to fasting across different ages. We use a typical high-

carbohydrate diet to study non-fasting ketogenesis across "normal" aging metabolism.

We then use KD to stimulate the endogenous ketogenic system and test how the

response differs across ages and sexes. We find that non-fasting hepatic ketogenesis

declines across ages more so in males in both fasted and non-fasted states, but could

be augmented by KD in a sex-dependent manner. We also show that aging induces

organ-specific ketogenesis to compensate for age-related declines in energetic

demands.

Results

Fasting-induced ketosis declines in aging and is sex-dependent.

We first tested if the kinetics of fasting ketosis, particularly the production of BHB,

changes with age, a topic surprisingly understudied given the links between aging

biology and fasting[32]. Male and female C57BL/6N mice of three ages (young adult, 3

months old; middle-aged adult, 12 months old; and older adult, 22 months old) were

fasted for 16 hours (6am-10pm) and monitored every 2 hours with blood drawn for

glucose and BHB measurements. Both male and female blood glucose levels

decreased to approximately 100 mg/dL by the end of their fasting period **(Fig. 1a, e)**.

Blood BHB concentrations increased as expected, but with striking age- and sex-

dependent variability (**Fig. 1b, f**). The overall area under the curve (AUC) of BHB

concentration declined sharply with age in both sexes **(Fig. 1c, g)**. Interestingly, the

trajectory of plasma BHB concentrations was different in females, with an early

intermediate peak and faster late rise, compared to males. In sum, the overall BHB AUC

during a 16-hour fast was 1.3-2 times higher in females compared to males, as was the

final BHB concentration at the end of fasting **(Fig. 1d, h)**, despite the similar overall

declines with age between sexes.

Age-related changes in metabolic state after fasting is sex-dependent.

The fasting data suggest that both age and sex affect energy substrate plasticity. To

better characterize differences in energy substrate utilization during fasting, we

observed male and female C57BL/6N mice at the same three ages (3 mo, 12 mo, 22

mo) in metabolic chambers for 4.5 days with a 24-hour fasting period in between **(Fig.**

2a, e). The respiratory exchange ratio (RER) is the ratio between CO_2 emission and O_2

consumption, with values typically ranging between 0.7 (primarily fat oxidation) and 1.0

(primarily carbohydrate oxidation). The RER of male mice both before and after fasting

was around 0.94 at all ages **(Fig. 2b, d)**. In contrast, female mice at 12 and 22 months

(but not 3 months) had a RER of 0.80 before fasting, increasing to 0.94 only after

fasting with the smallest change in the oldest females (**Fig. 2f, h**). Therefore, older

female mice appeared to be more dependent on fat oxidation at baseline compared to

young female mice or male mice at all ages **(Extended Fig. 1a-c)**. During the fasting

period, RER of all mice dropped to 0.79, a large change for males but minimal change

from baseline for females (**Fig. 2c, g**). In contrast to the striking decline in fasting BHB

with age, the differences with age in overall fuel utilization before, during, and after

fasting via RER were relatively small. Fuel utilization did show important sex differences

with fat metabolism predominating in non-fasting middle-aged and old female mice. This

difference may begin to explain the consistently higher BHB concentrations in fasting

observed for female mice.

Ketogenic diet helps augment non-fasting ketosis in aging. To further understand

how non-fasting ketosis changes in aging between sexes, the same three age groups of

male and female mice were acclimated to a control diet (77% carbohydrate, 10%

protein, 13% fat) for 14 days and then fed either the same control diet or a high-fat

ketogenic diet (90% fat and 10% protein) for an additional week **(Fig. 3a, f)**. We

collected tissue and plasma on day 7 just before the start of the light cycle (6am). A

ketogenic diet is a means of stimulating ketone body production without fasting or

energy restriction, and like fasting is a potent inducer of genes related to ketone body

production and utilization[20]. We reasoned that the control-fed group would show how

the basal ketogenic system changes with age, while the ketogenic diet would show the

latent potential of the ketogenic system upon stimulation. Changes in body weights in

both sexes on the control diet were less than a gram **(Extended Fig. 2a, b)**. Circulating

plasma BHB concentrations were within the typical range (0.05-0.3 mM) of a

predominantly carbohydrate diet and were similar between males and females, but were

generally lower in older mice though this reached statistical significance only in females

(Fig. 3b). Two major regulators of ketogenesis are insulin and substrate availability in

the form of non-esterified fatty acids (NEFA). Insulin levels increased with age as

expected **(Fig. 3c)**, accompanied by a decline in NEFA **(Fig. 3d)** albeit again only

significant in females. Glucose concentrations also declined with age in these non-

obese mice **(Fig. 3e)**. Together the rise in insulin and decline in NEFA help explain the

decline in basal, non-fasting BHB concentrations. Overall, these data show that non-

fasting ketosis does decline with age, more so in females. Body weights of young males

and females fed ketogenic diet for one week, along with middle-aged females increased

by 1-1.5 grams after the 1-week diet, but body weights declined by 1.5 grams in both

old male and female mice **(Extended Fig. 2c, d)**. Plasma BHB concentrations induced

by ketogenic diet remained relatively consistent across all ages **(Fig. 3g)**, in contrast to

both our prior fasting and control diet BHB blood concentration data. Insulin levels did

not increase significantly with age on ketogenic diet **(Fig. 3h)**, and NEFA levels

remained more stable as well **(Fig. 3i)**. We observed a dramatic drop in glucose levels

in old males and females along with a respective increase in plasma BHB

concentrations, exceeding a normal KD-range of ~2.0 mM **(Fig 3j)**. Overall, these data

suggest that aging mice retain the latent capacity to induce the ketogenic system under

the strong stimulus of a ketogenic diet. In addition, similar trajectories were observed in

males and females, leading us to investigate upstream regulators of ketogenesis next.

KD activates overall ketogenic capacity via upstream regulators of ketogenesis

but declines with aging. Ketogenesis is tightly regulated by fatty acid availability and,

transcriptionally, by the key regulator of fatty acid and ketone body metabolism,

peroxisome proliferator activated receptor alpha (PPARα). To better understand how

transcriptional regulation of ketogenesis changes with age and differs between sexes,

we examined the gene expression of enzymes, transporters, and upstream regulators

involved in ketogenesis in the liver via RT-qPCR. In control-fed males, we observed

strong age-related declines in gene expression of several regulators of ketogenesis

including PPARα, cAMP responsive element binding protein 3 like 3 (*Creb3l3*), the

PPARα transcriptional target murine cytochrome P450, family 4, subfamily a,

polypeptide 10 (*Cyp4a10*)*,* phosphoenolpyruvate carboxykinase 1 (*Pck1*), a target of

gluconeogenesis and importantly, 3-Hydroxy-3-Methyl-CoA synthase 2, (*Hmgcs2),* the

rate-limiting enzyme of ketogenesis **(Fig. 4a, Extended Fig. 3a-n)**. In our higher-

powered repeat control cohort, we were able to confirm the key age-related expression

declines in *PPARα and Hmgcs2* **(Extended Fig. 4a-h)**. KD strongly induces ketogenic

gene expression in males, but for several key genes the strength of induction declines

with age, including *PPARα*, *Creb3l3*, *Cyp4a10,* murine cytochrome P450, family 4,

subfamily a, polypeptide 14 (*Cyp4a14)*, and the ketogenesis pathways enzymes acetyl-

CoA acetyltransferase (*Acat1*), 3-hydroxymethyl-3-methylglutaryl-CoA lyase (*Hmgcl*),

and *Hmgcs2* **(Fig. 4a, Extended Fig. 3a-n)**. Interestingly, KD suppression of genes

involved in fatty acid synthesis (a key feature of KD) such as fatty acid synthase (*Fasn*),

acetyl-CoA carboxylase (*Acaa*), stearoyl-CoA desaturase (*Scd*), and ATP citrate lyase

(*Acly*), was strong at all ages **(Fig. 4a)**. Mass spectrometric analysis (via data-

independent acquisitions)[33,34] comparing relative protein abundance between young KD

vs young CD and old KD vs old CD, respectively, in the same liver samples revealed

similar patterns at the protein level. We observed an increase in CYP4A10, CYP4A14,

PCK1, ACAT1, HMGCS2, BDH1, and SLC16A1 in young and old KD-fed males and a

decrease in FASN and ACYL **(Fig. 4c)**. In control-fed females, we saw weak age-

related expression patterns for ketogenic genes and relatively less overall induction with

KD **(Fig. 4b, Extended Fig. 5a-n)**. Several upstream regulators and ketogenic pathway enzymes showed peak control diet expression at middle age, but only *PPARα* and *Bdh1* retained this pattern in the replication cohort (**Extended Figure 6a-h**). There was no consistent age-related pattern of KD gene expression induction, and less KD induction overall at any age other than for males. Similarly to males, we saw strong downregulation in genes related to fatty acid synthesis by KD across all ages **(Fig. 4b)**.

For a more systematic interpretation of these gene expression changes, we calculated a normalized composite score incorporating all of the upstream regulator and ketogenesis genes for each individual mouse **(Fig. 4d)**. Overall, ketogenic gene expression was relatively consistent across ages for control-fed males. KD strongly induced gene expression at all ages in males, but most strongly in young males. By contrast, females showed a life course trajectory of modestly higher ketogenic gene expression at middle age then lower at old age, but with little induction by KD at any age. Overall, our results show that males are more sensitive to gene expression changes by KD and that young male mice in particular are able to most strongly boost their ketogenic capacity through upregulation of PPARα and its targets compared to females and older males, similar to what we showed in our long term feeding of KD in old male mice (**Fig. 4e**)[20].

Tissue-specific changes in peripheral ketogenesis and ketolytic genes with age.
Based on the changes we observed in hepatic ketogenesis, we investigated how these changes affect ketolysis in peripheral tissues like the heart and brain. In the heart, both

control-fed males and females showed the highest expression of the rate-limiting

ketolytic gene, 3-oxoacid CoA-transferase (*Oxct1*) along with the membrane

transporter, solute carrier family member 1 (*Slc16a1*), at the oldest age **(Fig. 5a-b and**

Extended Fig. 7a-j). Interestingly, *Oxct1* was relatively suppressed by KD in the heart,

while the key ketogenic gene *Hmgcs2* was strongly induced by KD in the oldest males

and females **(Fig. 5a-b and Extended Fig. 7a-j)**. Overall, this suggests a pattern of

compensatory increase in ketone body utilization with aging in the heart, along with

possible augmentation by local ketogenesis under the stimulus of KD. In whole brain,

only male mice showed an increase in ketolytic genes with age solute carrier family 16

member 7,(*Slc16a7*)*, Acat1,* and *Oxct1*), not females **(Fig. 5c-d, Extended Fig. 8a-l)**.

Both male and female brains showed a similar pattern of strong KD gene expression

induction only at the youngest age, with stronger induction in females. Interestingly, this

KD gene expression induction in young mice included the ketogenic enzyme *Hmgcs2*.

Using LC-MS, we measured (R)-BHB in whole brains and observed relatively consistent

amounts across ages and sexes under control diet feeding **(Fig. 5e-f)**, suggesting that

homeostatic mechanisms preserve brain BHB availability despite age-related changes

in plasma BHB concentrations. KD dramatically increased brain (R)-BHB in all ages and

both sexes, but least so in middle-aged males. A third organ, kidney, showed age-

related decline in ketolytic enzymes (*Oxct1* and *Bdh1*) in males but not females

(Extended Fig. 9a-j), along with strong induction by KD of *Hmgcs2* at all ages in both

males and females. Altogether, we observed organ-specific patterns in age-related

changes in the ketogenic system that likely represent the adaptive role of ketone-based

energy production specific to each organ. Surprisingly, we more consistently observed

induction of *Hmgcs2* by KD than ketolytic genes, suggesting a possible role for local

ketogenesis within these tissues either for signaling roles or to augment an age-related

decline in hepatic ketogenesis.

249   Finally, to directly test if induced extra-hepatic ketogenesis meaningfully

contributes to blood BHB concentrations under KD, we generated a liver-specific

knockout (KO) mouse of *Bdh1* (Albumin-Cre; *Bdh1*[fl/fl]) that should produce only AcAc,

not BHB, from the liver[35] **(Extended Fig. 10a)**. Unlike heart-specific *Bdh1* KO mice[36] or

whole-body *Bdh1* KO mice[37], peripheral utilization of BHB should be unaffected in the

mouse model. Body weights and blood glucose in these KOs should also remain

unaffected **(Extended Fig. 10b-c, i-j)**. Ketogenesis, or perhaps only interconversion of

AcAc to BHB, has been described in the kidney but under fasting conditions does not

contribute to circulating BHB[38]. Surprisingly, we observed about 1/3 of normal blood

BHB concentrations under KD in the liver KO **(Extended Fig. 10d,k)**, suggesting a role

for non-hepatic ketogenesis under the strong stimulus of KD in compensating for loss of

hepatic BHB production[35,39–41]. We confirmed efficacy of the knockout via RT-qPCR

showing increased induction of *Hmgcs2* with KD, emphasizing ketogenesis, but an

insignificant change in *Bdh1,* supporting the lack of AcAc to BHB conversion in livers of

both male and female liver BDH1 KOs **(Extended Fig. 10e-h, l-o)**.

Discussion

266   Studies showing lifespan extension through interventions like fasting, dietary and

caloric restriction consistently show that metabolism and aging are tightly connected.

Though these interventions hold promise in preserving healthy longevity, the role of

pathways invoked by these interventions in normal aging often remains unclear. For an

example, elevating the concentration of circulating ketone bodies in the blood via caloric

restriction or ketogenic diet is associated with increased healthy longevity in mice, but

whether ketone bodies change in aging under normal conditions has been

understudied[16,42–44]. In this study, we systematically investigated ketosis in aging in

C57BL/6N male and female mice under three conditions, a 16-hour fast, a typical high-

carbohydrate diet and high-fat KD to recapitulate non-fasting and KD-induced ketosis

respectively.

We observed three important sex differences that should critically inform mouse

(and human) studies of metabolism in aging. First, during a 16-hour fast we observed

higher concentrations of circulating plasma BHB in females than males, although both

declined with age. Second, female RER is generally lower (more fat metabolizing) at

baseline. Third, we only observed substantial gene expression induction by KD in

males, despite similar non-fasting ketogenic responses. Our data are consistent with

limited studies of sex differences in ketosis in humans. One clinical study showed

fasting and postprandial plasma triacylglycerol concentrations were significantly higher

in men, but women had higher plasma BHB[45]. Women have higher lipolytic release than

men, which may help explain sexually dimorphic differences in lipid and glucose

metabolism[4]. Several studies also revealed higher lipogenesis and lipid storage in

females, hypothesized to be for reproduction, which could explain why female RER is

lower[46,47]. A caveat to comparing mouse to human fasting studies is that mouse BHB

concentrations exceeded 1.0 mM within 16 hours while heathy human adults reach that

point only after 48-72 hours of fasting, likely due to the faster metabolic rate of mice

versus humans[15,17,48]. Most metabolic studies in mice are undertaken only in young

males, and our data show that the fasting/ketogenic response of young male mice is not

representative of all ages and sexes. The relative lack of gene expression induction by

KD in female mice was striking from a mechanistic standpoint. Both non-fasting and

mechanistic studies in mice should include both sexes and, if targeting an age-related

disease, use older mice whenever possible.

Our data also provide important clues to the heretofore obscure role of ketosis in

aging. We hypothesized that ketogenic capacity might increase with age, in part due to

reports of a compensatory role of ketone bodies in specific age-related diseases such

as heart failure and Alzheimer's disease. Mechanistic studies in mice have

demonstrated that the failing heart adaptively switches to favor ketone body

metabolism, and that provision of ketone bodies either exogenously or via KD improves

heart function in heart failure models[25,36,49]. Instead we observed an age-related decline

in BHB induction under fasting, a decline in non-fasting blood BHB concentrations, and

a decline in expression of many ketogenic genes in the liver. Our data are consistent

with a more limited but cross-sectional aging study in male mice that also showed that

glucose and BHB were lower in the serum in the old, suggesting reduced production

from the liver[50]. Adipose tissue is a critical organ for nutrient sensing and energy

storage and has been widely studied to explain changes in metabolic health. Rising

insulin levels and declining circulating NEFA possibly through decreased lipolysis in

aging may proximately explain declining blood BHB levels with age[51–53]. We also

suspect that the high content of carbohydrates in our control diet and the rising insulin

levels with age could also inhibit ketogenic capacity through mTOR activation[44,54]. Aging

also induces inflammatory activation of NLRP3 and NF-κB, releasing pro-inflammatory

markers like IL-6 and TNFα, which could also diminish compensatory ketogenesis

through PPARα inhibition[55]. Despite the many KD studies done mainly in males (mice

and humans), a study administering exogenous ketone supplements in rats showed that

age and sex both modulate circulating blood BHB. In older rats, the ketone ester

increased concentrations of blood BHB more efficiently in male rats, supporting our data

for the KD-fed males[56].

In our study, we did find gene expression evidence of age-related compensatory

utilization of ketone bodies in extrahepatic organs, despite the overall age-related

decline in hepatic ketogenesis. In age-related diseases like heart failure and

Alzheimer's Disease, tissue-specific ketone body utilization increases, and overall

utilization is determined by the circulating concentrations of ketone bodies in the blood

[36,39,57–61]. From our data, heart and brain, but not kidney, showed increased expression

of ketolytic genes with age, but more so in control-fed animals, underscoring the

importance of ketone body utilization in heart failure and the aging brain. Despite the

levels being measured in whole brain, higher expression levels of the astrocyte specific

monocarboxylate transporter, *Slc16a1 and* neuron-specific monocarboxylate

transporter, *Slc16a2* also support an age-related increase in brain ketolysis[62–64].

However, when given a strong stimuli like KD, ketogenesis through *Hmgcs2*

gene expression dramatically increases in heart and kidney across all ages, suggesting

a role for compensatory tissue-specific ketogenesis. More specifically, heart fibroblast

and kidney mesangial cells have been shown to express *Hmgcs2*[65,66]. Cortex glia in

Drosophila melanogaster have been shown to also participate in ketogenesis through

AMPK signaling[67]. Based on our measurements of brain R-BHB, we suspect that the

higher concentrations could help explain higher ketogenic activity to meet brain

energetic demands[68–71]. Our previous study also showed that old male mice fed the

same KD used in this study for 12 months had improved cognitive function, which was

absent in mice fed both the high-carbohydrate and high-fat diets[21,72]. Further, we found

that knocking out liver *Bdh1*, which prevents liver production of BHB, still left ~1/3 of

circulating BHB intact under KD, suggesting that extrahepatic ketogenesis under KD is

a meaningful contributor to total body BHB production. In addition to providing extra-

hepatic tissues with an alternative energy substrate, ketone bodies also have important

signaling properties. One paper showed that AcAc can protect the liver from high-fat-

diet-induced fibrosis through mediating metabolic plasticity in macrophages[73]. A study

also showed that ketone bodies have an important role in the gut, such as mediating

intestinal stem cell homeostasis under different responses to diets[74].

This study is the first to systematically examine ketone body metabolism across

ages and sexes. The power of our cohorts limited the detection of metabolic and gene

expression changes to some extent, and so key measures were validated in a repeat,

larger, non-fasting control-fed cohort. Although we were able to distinguish between

ketogenesis and ketolysis through gene expression analyses, ketone production and

oxidation rates are not determined solely by gene expression. Future experiments

directly measuring oxidation rates using stable isotope labeling across sexes and ages

will deepen our understanding of the changes in ketone body metabolism. We also

acknowledge that in mice, sexes, ages, and strains can all influence the effects of

dietary interventions[75,76]. While KD remains to be an effective therapeutic concept in

brain aging, the driver of the sexual dimorphic changes we observed in our study

requires further elucidation. We also have reasons to believe that sex hormones, which

were not accounted for, regulate hepatic transport of ketone bodies and could help

explain the sexual dimorphic changes in ketone body metabolism[77]. Overall, this study

identifies critical age- and sex-differences in ketone body metabolism under fasting,

non-fasting, and ketogenic diet conditions, and suggests the potential of interventions

that target the endogenous ketogenic system to treat age-related diseases.

Methods

Mouse strains, housing and husbandry

C57BL/6 mice were obtained from the National Institute on Aging's Aged Rodent

Colony. *Bdh1*[fl/fl] mice were previously published[36]. Albumin-Cre mice[78] were obtained

from the Jackson Laboratory (#003574). All mice were maintained in animal facility

under a 6:00 am to 6:00 pm light cycle. All mice were housed in groups before being

single caged for their respective diets for one week. All mice were maintained according

to the National Institutes of Health guidelines and all procedures and protocols were

approved by the Institutional Animal Care and Use Committee at Buck Institute for

Research on Aging (Novato, CA).

Mouse diets and feeding

Food was provided ad libitum at all times. Per-calorie macronutrient content for

customized diets (Envigo) is as follows: Control, 10% protein, 13% fat, and 77%

carbohydrate (TD.150345); KD 10% protein and 90% fat (TD.160153). Both diets are

matched on a per-calorie basis for micronutrient content, fiber, and preservatives. All

mice were acclimated on a control diet for 2 weeks in the groups they arrived before

they were single caged for this study.

Metabolic Cage

C57BL/6 male and female mice (3, 12, 22 months) were single-caged in metabolic

chambers (Promethion Cages from Sable Systems International) for 108 hours. Raw

data files were analyzed using CalR (https://CalRapp.org).

Plasma Beta-Hydroxybutyrate Measurements

Blood was obtained via distal tail-snip (10-40 uL) or cardiac puncture if the animal was

euthanized for tissue collection in lithium-heparin coated microvettes (Sarstedt CB 300

LH). Blood was spun at 1500 x g for 15 mi at 4C and kept in -80C until thawed for

assay. Plasma BHB levels were measured using StanBio LiquiColor Beta-

Hydroxybutyrate colorimetric test per manufacturer's protocol (2440-058).

Glucose Measurements

Glucose levels were measured using a Accu-Chek blood glucose meter. Glucose

measurements for Figure 1 was every 2 hours during the 16-hour fast (6am-10pm).

Glucose measurements for Figure 3 was done before light period (6am) following mice

euthanasia.

Insulin Measurements

Insulin levels were measured using Mercodia Ultrasensitive Mouse Insulin ELISA per

manufacturer's protocol (Mercodia Cat. 10-1249-01).

Non-esterified fatty acid (NEFA) Measurements

NEFA levels were measured using NEFA Randox ELISA per manufacturer's protocol

(Randox Cat. FA115).

Real-time quantitative PCR

Mice were euthanized before the light period (6am) at the end of the 1-week feeding

period and tissues were immediately collected and snap-frozen for -80°C storage. Snap

frozen tissues were homogenized using RiboZol RNA Extraction Reagent (VWR; N580)

and Next Advance Bullet Blender (BBY24M). RNA was extracted using the Qiagen

RNEasy Mini Kit (Qiagen Cat. 169012183) and checked for purity using Thermo

Scientific Nanodrop 2000 (Serial No. D631). cDNA synthesis was carried out with

Superscript cDNA Synthesis Kit (BioRad Cat. 1708891) and real-time quantitative PCR

was performed using iTaq Universal SYBR Green Supermix (BioRad Cat. 1725121) in

BioRad CFX96 Real-Time System. Gene expression analyses were normalized to

either *B2m* (liver) or *Rplp0* (heart, brain, kidney) housekeeping genes.

Absolute Quantification of brain (R)-BHB

Frozen whole brains (~30-50 mg) were homogenized in extraction buffer [3:1, v/v

acetonitrile and HPLC-grade H2O] with Next Advance Bullet Blender (BBY24M).

Derivatization was adapted from the following paper[79]. Extracted samples were dried

using DNA SpeedVac System (ThermoFisher Scientific Model DNA130-115) and

resuspended in 98:2 H2O:Methanol containing .1% formic acid, mixed and centrifuged

at 10,000xg for 10 minutes. Supernatant was then transferred into HPLC vials. A

sample volume of 2 µL was injected into the UPLC-MS/MS Thermo Q Exactive with

Vanquish Horizon in Full MS and PRM scan modes using positive ionization. The

analysis was performed on a Accucore Vanquish C18+ column (100 x 2.1 mm, 1.5µm

particle size; ThermoFisher Cat. #20073385). The following mobile phases were used:

A) HPLC-graded H2O containing 0.1% (v/v) formic acid and B) methanol containing

0.1% (v/v) formic acid with the gradient starting at 2% B for 0.5 min and gradually

increasing to 10.4%B until 10.0 min, 2%B until 10.1 min, and 2%B until 12.0 at 0.150

mL/min flow rate for a total run time of 12.0 minutes. Column was maintained at 40°C.

LC system was hyphenated to Thermo Q Exactive MS equipped with heated

electrospray ionization (HESI) source. The MS system was operated in Full Scan MS or

PRM modes using positive ionization. MS scan range was 50.0 to 750.0 m/z in Full MS

scan mode. The resolution was set to 140,000 with AGC target 3e6, isolation window

1.0 m/z and optimal collision energy was 50 (arbitrary units). For PRM scan mode the

isolation window was set to 0.4 m/z and resolution was set to 70,000 with AGC target

1e6. Common HESI parameters were auxiliary gas: 5, sheath gas flow: 50, sweep gas:

0, spray voltage 3 kV, capillary temperature 320°, S-lens 55.0, and auxiliary gas

temperature: 150°C. Quantification of area response ratios were processed and

acquired using Thermo Scientific Xcalibur software (OPTON-30965). Area Response

Ratios (ES/IS) from three technical replicates were averaged, and a simple regression

line was constructed. To quantify the 'samples', Area Response Ratios (ES/IS) from

three technical replicates were averaged and concentrations were calculated using the

Prism-generated calibration curve equations. Amounts were calculated as

[(100*concentration) pmol]/[mg tissue weight].

Proteomics Analysis, Protein Digestion and Desalting

Livers of wild type and mice fed a ketogenic diet both young and old, with 4 biological

replicates each, were homogenized in 800 µL of lysis buffer containing 8 M urea, 200 mM

triethylammonium bicarbonate (TEAB), pH 8, 75 mM sodium chloride, 1 µM trichostatin

A, 3 mM nicotinamide, and 1x protease/phosphatase inhibitor cocktail (Thermo Fisher

Scientific, Waltham, MA), with 2 cycles in a Bead Beater TissueLyser II (Qiagen,

Germantown, MD) at 24 Hz for 2 min each. Lysates were clarified by spinning at 15,700

x g for 15 min at 4°C, and the supernatant containing the soluble proteins was collected.

Protein concentrations were determined using a Bicinchoninic Acid Protein (BCA) Assay

(Thermo Fisher Scientific, Waltham, MA). Protein lysates (3.1 mg each) were reduced

using 20 mM dithiothreitol (DTT) in 50 mM TEAB for 30 min at 37 °C, and after cooling to

room temperature, alkylated with 40 mM iodoacetamide (IAA) for 30 min at room

temperature in the dark. Samples were diluted 4-fold with 50 mM TEAB, pH 8, and

proteins were digested overnight with a solution of sequencing-grade trypsin (Promega,

San Luis Obispo, CA) in 50 mM TEAB at a 1:50 (wt:wt) enzyme:protein ratio at 37°C. This

reaction was quenched with 1% formic acid (FA) and the sample was clarified by

centrifugation at 2,000 x g for 10 min at room temperature. Clarified peptide samples were

desalted with Oasis 30-mg Sorbent Cartridges (Waters, Milford, MA). 100 µg of each

peptide elution were aliquoted for analysis of relative protein-level changes, after which

all desalted samples were vacuum dried. The 100 µg whole protein lysate aliquots were

re-suspended in 0.2% FA in water at a final concentration of 1 µg/µL. Finally, indexed

retention time standard peptides (iRT; Biognosys, Schlieren, Switzerland)[80] were spiked

in the samples according to manufacturer's instructions.

Mass Spectrometric Acquisitions

Reverse-phase HPLC-MS/MS analyses were performed on a Dionex UltiMate 3000

system coupled online to an Orbitrap Exploris 480 mass spectrometer (Thermo Fisher

Scientific, Bremen, Germany). The solvent system consisted of 2% ACN, 0.1% FA in

water (solvent A) and 80% ACN, 0.1% FA in ACN (solvent B). Digested peptides (400 ng)

were loaded onto an Acclaim PepMap 100 C_{18} trap column (0.1 x 20 mm, 5 µm particle

size; Thermo Fisher Scientific) over 5 min at 5 µL/min with 100% solvent A. Peptides (800

ng) were eluted on an Acclaim PepMap 100 C_{18} analytical column (75 µm x 50 cm, 3 µm

particle size; Thermo Fisher Scientific) at 300 nL/min using the following gradient: linear

from 2.5% to 24.5% of solvent B in 125 min, linear from 24.5% to 39.2% of solvent B in

40 min, up to 98% of solvent B in 1 min, and back to 2.5% of solvent B in 1 min. The

column was re-equilibrated for 30 min with 2.5% of solvent B, and the total gradient length

was 210 min. Each sample was acquired in data-independent acquisition (DIA) mode[33,34].

Full MS spectra were collected at 120,000 resolution (Automatic Gain Control (AGC)

target: 3e6 ions, maximum injection time: 60 ms, 350-1,650 *m/z*), and MS2 spectra at

30,000 resolution (AGC target: 3e6 ions, maximum injection time: Auto, Normalized

[33,34]Collision Energy (NCE): 30, fixed first mass 200 *m/z*). The isolation scheme consisted

of 26 variable windows covering the 350-1,650 *m/z* range with an overlap of 1 *m/z*[81].

Data Analysis, DIA-MS Data Processing and Statistical Analysis

DIA data was processed in Spectronaut (version 16.0.220524.53000) using directDIA.

Data was searched against a database containing all *Mus musculus* entries extracted

from UniProtKB-TrEMBL (86,521 protein entries; August 24, 2021). Trypsin/P was set as

digestion enzyme and two missed cleavages were allowed. Cysteine

carbamidomethylation was set as fixed modification, and methionine oxidation, and

protein N-terminus acetylation variable modifications. Data extraction parameters were

set as dynamic. Identification was performed requiring a 1% q-value cutoff on the

precursor ion and protein level. For the protein lysate samples, quantification was based

on the peak areas of extracted ion chromatograms (XICs) of 3 – 6 MS2 fragment ions,

and local normalization was applied. iRT profiling was selected. Quantification was based

on the peak areas of XICs of 3 – 6 MS2 fragment ions, without normalization as well as

data filtering using q-value and imputation of missing values with empirical noise.

Differential expression analysis comparing ketogenic diet to wild-type mouse livers was

performed using a paired t-test, and p-values were corrected for multiple testing,

specifically applying group wise testing corrections using the Storey method[82].

Quantifiable proteins consist of protein groups with at least two unique peptides and

significantly-altered proteins have a q-value < 0.01 and absolute Log_2(fold-change) > 0.58

(Supplementary Tables).

Data Availability

Raw data and complete MS data sets have been uploaded to the Mass Spectrometry

Interactive Virtual Environment (MassIVE) repository, developed by the Center for

Computational Mass Spectrometry at the University of California San Diego, and can be

downloaded using the following link:

https://massive.ucsd.edu/ProteoSAFe/dataset.jsp?task=934f911545e84050ac0c18346d

12006c   (MassIVE ID number: MSV000090365; ProteomeXchange ID: PXD036921).

Enter the username and password in the upper right corner of the page:

Username: *MSV000090365_reviewer*

Reviewer password: *winter*

Heatmap Differential Expression Analysis

An average value was calculated for each gene across ages of both control and KD

groups of each sex. Values are to represent minimum and maximum differential

expressions for each gene. This analysis does not provide statistical significance, but

rather allowed for a better representation of KD-induction compared to our control

groups.

Ketogenic Composite Score Analysis

Composite scores were calculated to represent an overall trajectory of ketogenic genes

shown on the heatmap. The standard deviations and means of young control diet from

each respective sex were normalized and used to calculate ketogenic scores for 12 and

22 months of both control and KD groups. Each individual mouse was normalized as

the number of baseline standard deviations from the baseline mean.

Quantification and Statistical analysis

Figures are presented as mean ± SEM, P values are calculated by either one-way

ANOVA or two-way ANOVA with additional post-hoc testing for group comparisons.

Statistical calculations are carried out in Prism 9.4.1 (GraphPad Software, San Diego,

California USA, www.graphpad.com). Exact replicate numbers "N" are specified in the

figure legends for each experiment, and refer to the number of individual mice.

References

1. López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. The

hallmarks of aging. *Cell* **153**, 1194–1217 (2013).

2. Paolisso, G. *et al.* Glucose tolerance and insulin action in healthy centenarians. *Am.*

J. Physiol. **270**, E890-894 (1996).

3. Marmentini, C. *et al.* Aging Reduces Insulin Clearance in Mice. *Front. Endocrinol.*

12, 679492 (2021).

4. Lee, S.-H. & Min, K.-J. Caloric restriction and its mimetics. *BMB Rep.* **46**, 181–187

(2013).

5. Fontana, L., Partridge, L. & Longo, V. D. Extending healthy life span--from yeast to

humans. *Science* **328**, 321–326 (2010).

6. Mannick, J. B. *et al.* Targeting the biology of ageing with mTOR inhibitors to improve

immune function in older adults: phase 2b and phase 3 randomised trials. *Lancet*

Healthy Longev. **2**, e250–e262 (2021).

7. Justice, J. N. *et al.* A framework for selection of blood-based biomarkers for

geroscience-guided clinical trials: report from the TAME Biomarkers Workgroup.

GeroScience **40**, 419–436 (2018).

8. Harrison, D. E. *et al.* Rapamycin fed late in life extends lifespan in genetically

heterogeneous mice. *Nature* **460**, 392–395 (2009).

9. Wilkinson, J. E. *et al.* Rapamycin slows aging in mice. *Aging Cell* **11**, 675–682

(2012).

10. Barzilai, N., Crandall, J. P., Kritchevsky, S. B. & Espeland, M. A. Metformin as a

Tool to Target Aging. *Cell Metab.* **23**, 1060–1065 (2016).

11. Kulkarni, A. S., Gubbi, S. & Barzilai, N. Benefits of Metformin in Attenuating the

Hallmarks of Aging. *Cell Metab.* **32**, 15–30 (2020).

12. Rojas-Morales, P., Tapia, E. & Pedraza-Chaverri, J. β-Hydroxybutyrate: A signaling

metabolite in starvation response? *Cell. Signal.* **28**, 917–923 (2016).

13. Asadi Shahmirzadi, A. *et al.* Alpha-Ketoglutarate, an Endogenous Metabolite,

Extends Lifespan and Compresses Morbidity in Aging Mice. *Cell Metab.* **32**, 447-

456.e6 (2020).

14. Robinson, A. M. & Williamson, D. H. Physiological roles of ketone bodies as

substrates and signals in mammalian tissues. *Physiol. Rev.* **60**, 143–187 (1980).

15. Cahill, G. F. Fuel metabolism in starvation. *Annu. Rev. Nutr.* **26**, 1–22 (2006).

16. London, E. D. *et al.* Effects of fasting on ketone body concentrations in healthy men

of different ages. *J. Gerontol.* **41**, 599–604 (1986).

17. Cahill, G. F. *et al.* Hormone-fuel interrelationships during fasting. *J. Clin. Invest.* **45**,

1751–1769 (1966).

18. Balasse, E. O. & Féry, F. Ketone body production and disposal: Effects of fasting,

diabetes, and exercise. *Diabetes. Metab. Rev.* **5**, 247–270 (1989).

19. Koeslag, J. H., Noakes, T. D. & Sloan, A. W. Post-exercise ketosis. *J. Physiol.* **301**,

79–90 (1980).

20. Newman, J. C. *et al.* Ketogenic Diet Reduces Midlife Mortality and Improves

Memory in Aging Mice. *Cell Metab.* **26**, 547-557.e8 (2017).

21. Roberts, M. N. *et al.* A ketogenic diet extends longevity and healthspan in adult

mice. *Cell Metab.* **26**, 539-546.e5 (2017).

22. Tognini, P. *et al.* Distinct Circadian Signatures in Liver and Gut Clocks Revealed by

Ketogenic Diet. *Cell Metab.* **26**, 523-538.e5 (2017).

23. Croteau, E. *et al.* A cross-sectional comparison of brain glucose and ketone

metabolism in cognitively healthy older adults, mild cognitive impairment and early

Alzheimer's disease. *Exp. Gerontol.* **107**, 18–26 (2018).

24. Tomita, I. *et al.* SGLT2 Inhibition Mediates Protection from Diabetic Kidney Disease

by Promoting Ketone Body-Induced mTORC1 Inhibition. *Cell Metab.* **32**, 404-419.e6

(2020).

25. Aubert, G. *et al.* The Failing Heart Relies on Ketone Bodies as a Fuel. *Circulation*

133, 698–705 (2016).

26. Voros, G. *et al.* Increased Cardiac Uptake of Ketone Bodies and Free Fatty Acids in

Human Heart Failure and Hypertrophic Left Ventricular Remodeling. *Circ. Heart Fail.*

11, e004953 (2018).

27. Bedi, K. C. *et al.* Evidence for Intramyocardial Disruption of Lipid Metabolism and

Increased Myocardial Ketone Utilization in Advanced Human Heart Failure.

Circulation **133**, 706–716 (2016).

28. Youm, Y.-H. *et al.* The ketone metabolite β-hydroxybutyrate blocks NLRP3

inflammasome-mediated inflammatory disease. *Nat. Med.* **21**, 263–269 (2015).

29. Benyó, Z. *et al.* GPR109A (PUMA-G/HM74A) mediates nicotinic acid-induced

flushing. *J. Clin. Invest.* **115**, 3634–3640 (2005).

30. Jensen, M. D. Gender differences in regional fatty acid metabolism before and after

meal ingestion. *J. Clin. Invest.* **96**, 2297–2303 (1995).

31. Romanski, S. A., Nelson, R. M. & Jensen, M. D. Meal fatty acid uptake in adipose

tissue: gender effects in nonobese humans. *Am. J. Physiol.-Endocrinol. Metab.* **279**,

E455–E462 (2000).

32. Anton, S., Ezzati, A., Witt, D., McLaren, C. & Vial, P. The effects of intermittent

fasting regimens in middle-age and older adults: Current state of evidence. *Exp.*

Gerontol. **156**, 111617 (2021).

33. Gillet, L. C. *et al.* Targeted data extraction of the MS/MS spectra generated by data-

independent acquisition: a new concept for consistent and accurate proteome

analysis. *Mol. Cell. Proteomics MCP* **11**, O111.016717 (2012).

34. Collins, B. C. *et al.* Multi-laboratory assessment of reproducibility, qualitative and

quantitative performance of SWATH-mass spectrometry. *Nat. Commun.* **8**, 291

(2017).

35. Stagg, D. B. *et al.* Diminished ketone interconversion, hepatic TCA cycle flux, and

glucose production in D-β-hydroxybutyrate dehydrogenase hepatocyte-deficient

mice. *Mol. Metab.* **53**, 101269 (2021).

36. Horton, J. L. *et al.* The failing heart utilizes 3-hydroxybutyrate as a metabolic stress

defense. *JCI Insight* **4**, 124079 (2019).

37. Otsuka, H. *et al.* Deficiency of 3-hydroxybutyrate dehydrogenase (BDH1) in mice

causes low ketone body levels and fatty liver during fasting. *J. Inherit. Metab. Dis.*

43, 960–968 (2020).

38. Venable, A. H. *et al.* Fasting-induced HMGCS2 expression in the kidney does not

contribute to circulating ketones. *Am. J. Physiol. Renal Physiol.* **322**, F460–F467

(2022).

39. Cuenoud, B. *et al.* Metabolism of Exogenous D-Beta-Hydroxybutyrate, an Energy

Substrate Avidly Consumed by the Heart and Kidney. *Front. Nutr.* **7**, (2020).

40. Mccann, W. P. The oxidation of ketone bodies by mitochondria from liver and

peripheral tissues. *J. Biol. Chem.* **226**, 15–22 (1957).

41. McGarry, J. D. & Foster, D. W. Regulation of hepatic fatty acid oxidation and ketone

body production. *Annu. Rev. Biochem.* **49**, 395–420 (1980).

42. Hennebelle, M. *et al.* Preliminary evaluation of a differential effect of an α-linolenate-

rich supplement on ketogenesis and plasma ω-3 fatty acids in young and older

adults. *Nutr. Burbank Los Angel. Cty. Calif* **32**, 1211–1216 (2016).

43. Kuhla, A. *et al.* Lifelong caloric restriction reprograms hepatic fat metabolism in

mice. *J. Gerontol. A. Biol. Sci. Med. Sci.* **69**, 915–922 (2014).

44. Sengupta, S., Peterson, T. R., Laplante, M., Oh, S. & Sabatini, D. M. mTORC1

controls fasting-induced ketogenesis and its modulation by ageing. *Nature* **468**,

1100–1104 (2010).

45. Pramfalk, C. *et al.* Sex-Specific Differences in Hepatic Fat Oxidation and Synthesis

May Explain the Higher Propensity for NAFLD in Men. *J. Clin. Endocrinol. Metab.*

100, 4425–4433 (2015).

46. Della Torre, S. *et al.* Short-Term Fasting Reveals Amino Acid Metabolism as a Major

Sex-Discriminating Factor in the Liver. *Cell Metab.* **28**, 256-267.e5 (2018).

47. Sparks, L. M. *et al.* Effect of adipose tissue on the sexual dimorphism in metabolic

flexibility. *Metabolism.* **58**, 1564–1571 (2009).

48. Robinson, A. M. & Williamson, D. H. Physiological roles of ketone bodies as

substrates and signals in mammalian tissues. *Physiol. Rev.* **60**, 143–187 (1980).

49. Yurista, S. R. *et al.* Ketone Ester Treatment Improves Cardiac Function and

Reduces Pathologic Remodeling in Preclinical Models of Heart Failure. *Circ. Heart*

Fail. **14**, e007684 (2021).

50. Petr, M. A. *et al.* A cross-sectional study of functional and metabolic changes during

aging through the lifespan in male mice. *eLife* **10**, e62952 (2021).

51. Arner, P. Catecholamine-induced lipolysis in obesity. *Int. J. Obes. Relat. Metab.*

Disord. J. Int. Assoc. Study Obes. **23 Suppl 1**, 10–13 (1999).

52. Mullins, G. R. *et al.* Catecholamine-induced lipolysis causes mTOR complex

dissociation and inhibits glucose uptake in adipocytes. *Proc. Natl. Acad. Sci.* **111**,

17450–17455 (2014).

53. Catecholamine-induced lipolysis in adipose tissue of the elderly. - PMC.

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC296613/.

54. Caron, A., Richard, D. & Laplante, M. The Roles of mTOR Complexes in Lipid

Metabolism. *Annu. Rev. Nutr.* **35**, 321–348 (2015).

55. Flint, T. R. *et al.* Tumor-Induced IL-6 Reprograms Host Metabolism to Suppress

Anti-tumor Immunity. *Cell Metab.* **24**, 672 (2016).

56. Kovács, Z., Brunner, B., D'Agostino, D. P. & Ari, C. Age- and Sex-Dependent

Modulation of Exogenous Ketone Supplement-Evoked Effects on Blood Glucose

and Ketone Body Levels in Wistar Albino Glaxo Rijswijk Rats. *Front. Neurosci.* **14**,

618422 (2020).

57. Luong, T. V., Abild, C. B., Bangshaab, M., Gormsen, L. C. & Søndergaard, E.

Ketogenic Diet and Cardiac Substrate Metabolism. *Nutrients* **14**, 1322 (2022).

58. Cotter, D. G., Schugar, R. C. & Crawford, P. A. Ketone body metabolism and

cardiovascular disease. *Am. J. Physiol. Heart Circ. Physiol.* **304**, H1060-1076

(2013).

59. Cunnane, S. C. *et al.* Can ketones compensate for deteriorating brain glucose

uptake during aging? Implications for the risk and treatment of Alzheimer's disease.

Ann. N. Y. Acad. Sci. **1367**, 12–20 (2016).

60. Kashiwaya, Y. *et al.* d-β-Hydroxybutyrate protects neurons in models of Alzheimer's

and Parkinson's disease. *Proc. Natl. Acad. Sci.* **97**, 5440–5444 (2000).

61. Fortier, M. *et al.* A ketogenic drink improves brain energy and some measures of

cognition in mild cognitive impairment. *Alzheimers Dement.* **15**, 625–634 (2019).

62. Ding, F., Yao, J., Rettberg, J. R., Chen, S. & Brinton, R. D. Early decline in glucose

transport and metabolism precedes shift to ketogenic system in female aging and

Alzheimer's mouse brain: implication for bioenergetic intervention. *PloS One* **8**,

e79977 (2013).

63. Pierre, K., Pellerin, L., Debernardi, R., Riederer, B. M. & Magistretti, P. J. Cell-

specific localization of monocarboxylate transporters, MCT1 and MCT2, in the adult

mouse brain revealed by double immunohistochemical labeling and confocal

microscopy. *Neuroscience* **100**, 617–627 (2000).

64. Pierre, K. & Pellerin, L. Monocarboxylate transporters in the central nervous system:

distribution, regulation and function. *J. Neurochem.* **94**, 1–14 (2005).

65. Tabula Muris Consortium *et al.* Single-cell transcriptomics of 20 mouse organs

creates a Tabula Muris. *Nature* **562**, 367–372 (2018).

66. Schaum, N. *et al.* Ageing hallmarks exhibit organ-specific temporal signatures.

Nature **583**, 596–602 (2020).

67. Silva, B. *et al.* Glia fuel neurons with locally synthesized ketone bodies to sustain

memory under starvation. *Nat. Metab.* 1–12 (2022) doi:10.1038/s42255-022-00528-

6.

68. Pan, J. W., Rothman, T. L., Behar, K. L., Stein, D. T. & Hetherington, H. P. Human

brain beta-hydroxybutyrate and lactate increase in fasting-induced ketosis. *J. Cereb.*

Blood Flow Metab. Off. J. Int. Soc. Cereb. Blood Flow Metab. **20**, 1502–1507

(2000).

69. Zhang, Y., Kuang, Y., LaManna, J. C. & Puchowicz, M. A. Contribution of brain

glucose and ketone bodies to oxidative metabolism. *Adv. Exp. Med. Biol.* **765**, 365–

370 (2013).

70. Zhang, Y., Zhang, S., Marin-Valencia, I. & Puchowicz, M. A. Decreased carbon

shunting from glucose toward oxidative metabolism in diet-induced ketotic rat brain.

J. Neurochem. **132**, 301–312 (2015).

71. Marosi, K. *et al.* 3-Hydroxybutyrate regulates energy metabolism and induces BDNF

expression in cerebral cortical neurons. *J. Neurochem.* **139**, 769–781 (2016).

72. Newman, J. C. *et al.* Ketogenic diet reduces mid-life mortality and improves memory

in aging mice. *Cell Metab.* **26**, 547-557.e8 (2017).

73. Puchalska, P. *et al.* Hepatocyte-Macrophage Acetoacetate Shuttle Protects against

Tissue Fibrosis. *Cell Metab.* **29**, 383-398.e7 (2019).

74. Cheng, C.-W. *et al.* Ketone Body Signaling Mediates Intestinal Stem Cell

Homeostasis and Adaptation to Diet. *Cell* **178**, 1115-1131.e15 (2019).

75. Pak, H. H. *et al.* Fasting drives the metabolic, molecular and geroprotective effects

of a calorie-restricted diet in mice. *Nat. Metab.* **3**, 1327–1341 (2021).

76. Green, C. L. *et al.* Sex and genetic background define the metabolic, physiologic,

and molecular response to protein restriction. *Cell Metab.* **34**, 209-226.e5 (2022).

77. Cao, J., Ng, M. & Felmlee, M. A. Sex Hormones Regulate Rat Hepatic

Monocarboxylate Transporter Expression and Membrane Trafficking. *J. Pharm.*

Pharm. Sci. Publ. Can. Soc. Pharm. Sci. Soc. Can. Sci. Pharm. **20**, 435–444 (2017).

78. Postic, C. *et al.* Dual roles for glucokinase in glucose homeostasis as determined by

liver and pancreatic beta cell-specific gene knock-outs using Cre recombinase. *J.*

Biol. Chem. **274**, 305–315 (1999).

79. Tsutsui, H. *et al.* Simultaneous determination of DL-lactic acid and DL-3-

hydroxybutyric acid enantiomers in saliva of diabetes mellitus patients by high-

throughput LC-ESI-MS/MS. *Anal. Bioanal. Chem.* **404**, 1925–1934 (2012).

80. Escher, C. *et al.* Using iRT, a normalized retention time for more targeted

measurement of peptides. *Proteomics* **12**, 1111–1121 (2012).

81. Bruderer, R. *et al.* Optimization of Experimental Parameters in Data-Independent

Mass Spectrometry Significantly Increases Depth and Reproducibility of Results.

Mol. Cell. Proteomics MCP **16**, 2296–2309 (2017).

82. Burger, T. Gentle Introduction to the Statistical Foundations of False Discovery Rate

in Quantitative Proteomics. *J. Proteome Res.* **17**, 12–22 (2018).

Acknowledgements

This work was supported by NIH grant R01AG67333 (J.C.N.), and Buck intramural

funds (J.C.N.), NIA T32 AG052374 (B.E), NIA T32 AG000266 (M.N), and NIH

R01HL151345 (D.P.K). We thank S. Galkina from Buck Institute's Phenotyping Core for

administering the metabolic cage study, A. Holcom for her assistance in the repeat

control cohort tissue collection, and B. J. Stubbs for critical reading of the manuscript.

Some figure illustrations were created with BioRender.com.

Authorship contributions

B.E. conceived the studies, designed the experiments, analyzed experiments, and

wrote the manuscript. M.N., O.P., and T.Y.G helped perform experiments. J.P.R.,

C.D.K., and B.S. performed and analyzed proteomics experiments. T.C.L. and D.P.K

generated Bdh1$^{fl/fl}$ mice. D.P.K provided important insights and suggestions to

manuscript. J.C.N. supervised the study and co-wrote the manuscript.

Competing interests

792   **Additional Information**

793   All data needed to evaluate the conclusions in the paper are present in the paper and

794   Extended Figures. Data requests should be submitted to the corresponding author.

Figure 1

Fig. 1 | Fasting-induced ketosis declines in aging and is sex-dependent.

a, Trajectory of 16-hour fasting glucose levels in male mice (3, 12, 22 months), 6am-

10pm **b,** Trajectory of 16-hour fasting plasma BHB levels in males, 6am-10pm **c,** AUC

of plasma BHB in males **d,** values of fasting AUC and final fasting concentration in

males **e,** Trajectory of 16-hour fasting glucose levels in female mice (3, 12, 22 months)

f, Trajectory of 16-hour fasting plasma BHB levels in females **g,** AUC of plasma BHB in

females **h,** values of fasting AUC and final fasting concentration in females **a-c, e-g,**

n=5-6 per age group, all data are presented as mean ± SEM; Ordinary one-way

ANOVA, post-hoc Tukey's for multiple comparisons test between age groups,

*p<0.0332, **p<0.0021, ***p<0.0002, ****p<0.001.

Figure 2

Fig. 2 | Age-related changes in metabolic state after fasting is sex-dependent.

a, Respiratory Exchange Ratio (RER) of male mice (3, 12, 22 months) over the course

of 4.5 days **b,** Average RER for before fasting in males **c,** Average RER for 24 hour

fasting in males **d,** Average RER for after fasting in males **e,** Respiratory Exchange

Ratio (RER) of female mice over the course of 4.5 days **f,** Average RER for before

fasting in female mice (3, 12, 22 months) **g,** Average RER for 24 hour fasting in females

h, Average RER for after fasting in females. **a,e,** n=5 per age group **b-d, f-h,** n=5 per

age group; all data are presented as mean ± SEM; Ordinary one-way ANOVA, post-hoc

Tukey's for multiple comparisons test between age groups, *p<0.0332, **p<0.0021,

***p<0.0002. ****p<0.001.

Figure 3

Fig. 3 | Ketogenic diet helps augment non-fasting ketosis in aging. a, Control diet,

sex and age groups, and feeding timeline **b,** Day 7 plasma BHB concentrations of

pooled cohort **c,** Day 7 insulin levels **d,** Day 7 non-esterified fatty acids levels **e,** Day 7

blood glucose levels **f,** Ketogenic diet, sex and age groups, and feeding timeline **g,** Day

7 plasma BHB concentrations **h,** Day 7 insulin levels **i,** Day 7 non-esterified fatty acids

levels **j,** Day 7 blood glucose levels **b,** n=16-26 per sex groups **c-e, g-j,** n=6-7 per sex,

diet and age groups, all data are presented as mean ± SEM; Ordinary two-way ANOVA,

post-hoc Tukey's for multiple comparisons test between age groups, *p<0.0332,

**p<0.0021, ***p<0.0002. ****p<0.001.

Liver Gene Expression

Figure 4

Fig. 4 | KD activates overall ketogenic capacity via upstream regulators of ketogenesis but declines with aging. a, Hepatic differential gene expression in control and KD-fed males **b,** Hepatic differential gene expression in control and KD-fed females **c,** Volcano plot of mass spectrometric analysis (via data-independent acquisitions) comparing relative protein abundance between young KD vs young CD and old KD vs old CD males (n=4/age and diet groups) **d,** Composite scores of control and KD-fed mice normalized to young mice **e,** schematic of regulators ketogenesis in the liver and ketolysis in extra-hepatic tissues **a, b ,d,** n=6-7 per sex and age groups, all data are presented as mean ± SEM; Ordinary two-way ANOVA, post-hoc Tukey's for multiple comparisons test between age groups, *p<0.0332, **p<0.0021, ***p<0.0002. ****p<0.001.

Figure 5

Fig. 5 | Tissue-specific changes in peripheral ketogenesis and ketolytic genes with age. a, Heatmap of heart gene expression in control and KD-fed males **b,** Heatmap of heart gene expression in control and KD-fed females **c,** Heatmap of brain gene expression in control and KD-fed males **d,** Heatmap of brain gene expression in control and KD-fed females **e,** Concentrations of *(R)*-BHB in whole brains of males **f,** Concentrations of *(R)*-BHB in whole brains of females **e-f,** n=6-7 per sex and age groups, all data are presented as mean ± SEM; Ordinary two-way ANOVA, post-hoc Tukey's for multiple comparisons test between age groups, *p<0.0332, **p<0.0021, ***p<0.0002. ****p<0.001.

Figure 6

Fig. 6 | KD boosts the age-related declines non-fasting and fasting-induced ketosis in a sex-dependent manner. Non-fasted and fasted mice all showed a decline in circulating blood BHB with age, which was augmented by KD. The increases in blood BHB by KD were observed in both sex but was stronger in males. Changes in ketone body metabolism in extra-hepatic organs like the heart, kidney, and brain were diet-dependent. In control-fed animals, ketolysis was favored over ketogenesis whereas KD provided fats for tissue-specific ketogenesis. Overall, our work shows that hepatic ketogenesis declines in the liver with age, incapable of compensating for the hypometabolism of glucose we commonly see in aging. However, with KD, ketogenesis is promoted systemically to help compensate for this decline.