Title: Plant functional types and tissue stoichiometry explain nutrient transfer in common

arbuscular mycorrhizal networks of temperate grasslands

Authors: Hilary Rose Dawson[1,2,3*], Katherine L. Shek[1,2,4], Toby M. Maxwell[1,5], Paul B. Reed[1,2,6],

Barbara Bomfim[1,7], Scott D. Bridgham[1,2], Brendan J. M. Bohannan[1,2], Lucas C.R. Silva[1,2,8*]

Affiliations:

1. Institute of Ecology and Evolution, University of Oregon, Eugene, OR 97403 USA

2. Department of Biology, University of Oregon, Eugene, OR 97403 USA

3. Research School of Biology, Australian National University, Canberra, ACT 2601 Australia

4. Department of Natural Resources and the Environment, University of New Hampshire,

Durham NH 03824

5. Department of Biological Sciences, Boise State University, Boise, Idaho, 83725 USA

6. Institute for Applied Ecology, Corvallis, OR 97333 USA

7. Climate and Ecosystem Sciences Division, Lawrence Berkeley National Laboratory,

Berkeley, CA 94720 USA

8. Environmental Studies Program, University of Oregon, Eugene, OR 97403 USA

ORCIDs: H.R.D.: 0000-0003-4613-762X; K.L.S.: 0000-0001-5629-5782; T.M.M.: 0000-0001-

5171-0705; P.B.R.: 0000-0001-7143-7515; B.B.: 0000-0001-9510-2496; S.D.B.: 0000-0003-

0614-2678; B.J.M.B.: 0000-0003-2907-1016; L.C.R.S.: 0000-0002-4838-327X

Corresponding authors: H.R.D. hilary.dawson@anu.edu.au; L.C.R.S. lsilva7@uoregon.edu

Acknowledgements: The authors thank the Siskiyou Field Institute, The Nature Conservancy,

and Capitol Land Trust for providing sites for this experiment, Laurel Pfeifer-Meister, Bitty Roy,

Bart Johnson, Graham Bailes, Aaron Nelson, and Matthew Krna for their contributions to

experimental design, Emily Scherer for her assistance with sample analysis, Bitty Roy, Leonel

Sternberg, and Timothy A. Doane for comments on the manuscript, and numerous others for

assistance with the HOPS project. This experiment was funded by National Science Foundation

Macrosystems Biology grant #1340847, Plant Biotic Interactions grant #1758947, and

Convergence Accelerator Pilot grant #1939511.

Author contributions: L.C.R.S, S.D.B., and B.J.M.B. designed the research; H.R.D., K.L.S.,

T.M.M., B.B., and P.B.R. performed the research; H.R.D. led the data analysis and manuscript

preparation with K.L.S. assisting; all authors contributed to revising the manuscript and

approved the publication.

Conflicts of interest: The authors have no conflicts of interest to declare.

Statement of inclusion: Our study was performed in the region where all authors were located

when the research began. Several authors are long-time residents of the region while others were

based at local institutions as students or in shorter term research positions. Our study was

designed to address concerns about sustainability in the region, concerns that are held by

residents, managers, and researchers alike.

Data availability: Data used in these analyses are available online at the Data Dryad repository

(https://doi.org/10.5061/dryad.7pvmcvdxt).

Title: Plant functional types and tissue stoichiometry explain nutrient transfer in common

arbuscular mycorrhizal networks of temperate grasslands

Abstract:

Plants and mycorrhizal fungi form mutualistic relationships that affect how resources

flow between organisms and within ecosystems. Common mycorrhizal networks (CMNs) could

facilitate preferential transfer of carbon and limiting nutrients, but this remains difficult to

predict. Do CMNs favor fungal resource acquisition at the expense of plant resource demands (a

fungi-centric view), or are they passive channels through which plants regulate resource fluxes (a

plant-centric view)? We used stable isotope tracers ($^{13}CO_2$ and $^{15}NH_3$), plant traits, and

mycorrhizal DNA to quantify above- and belowground carbon and nitrogen transfer between 18

plant species along a 520-km latitudinal gradient in the Pacific Northwest, USA. Plant functional

type and tissue stoichiometry were the most important predictors of interspecific resource

transfer. Of "donor" plants, 98% were ^{13}C-enriched, but we detected transfer in only 2% of

"receiver" plants. However, all donors were ^{15}N-enriched and we detected transfer in 81% of

receivers. Nitrogen was preferentially transferred to annuals (0.26 ± 0.50 mg N per g leaf mass)

compared to perennials (0.13 ± 0.30 mg N per g leaf mass). This corresponded with tissue

stoichiometry differences. Our findings suggest that plants and fungi that are located closer

together in space and with stronger demand for resources over time are more likely to receive

larger amounts of those limiting resources.

Keywords: arbuscular mycorrhizal fungi, carbon, common mycorrhizal networks, ecological
stoichiometry, grasslands, nitrogen, plant functional type

Introduction:

Plant-mycorrhizal associations are thought to have emerged as rudimentary root systems over 400 million years ago, facilitating the expansion of terrestrial life that followed (Kenrick & Strullu-Derrien, 2014). The transformative power of early fungal symbioses is still evident today in all major plant lineages, from bryophytes to angiosperms (Heijden et al., 2015). Over 85% of all contemporary flowering plant species form symbioses with fungi, with arbuscular mycorrhizal (AM) associations being the most common (Brundrett, 2009). The relationships between plants and AM fungi dominates both managed and unmanaged landscapes and are estimated to be responsible for up to 80% of global primary productivity (Heijden et al., 2015). Fungi can form symbioses with more than one individual plant, particularly AM fungi which have low host specificity (Selosse et al., 2006). By extension, it has been widely hypothesized that the multi-plant-fungal relationships form "common mycorrhizal networks" (CMNs) which facilitate carbon and nutrient transfer between organisms, beyond the immediate plant-fungus mutualism formed by individuals. Although CMNs are traditionally defined by strict criteria that are difficult to test experimentally (Karst et al., 2008), the concept is ecologically relevant and useful in designing new experiments that may bring insight into CMNs structure and function. Here, we use 'CMN' under the proposed new definition of Rillig et al (2024) "where at least one mycorrhizal fungal genet interacts (connecting and colonizing or growing in close proximity) with the roots of a minimum of two plants of the same or different species." Although Rillig et al. (2024) claim it is unlikely that carbon and nutrient exchanges will occur without hyphal continuity, we disagree that hyphal continuity is necessary for these exchanges. Plants leak exudates into the soil that can be picked up by fungal hyphae near the root surface (Vives-Peris et al., 2020). The same fungi can be in symbiosis with multiple plants. Although this interaction

is not what is classically considered a CMN (what Rillig et al. have renamed as a CMN with

hyphal continuity, or CMN-HC), it remains in essence fungal transport of carbon and nutrients

derived from the plants involved (a CMN by Rillig et al.'s revised definition). Both CMN-HCs

and CMNs could have ecological significance. The CNM relationships are often simplified to the

mutual exchange of carbon-based photosynthates for soil nutrients that are more readily

bioavailable to fungi (Smith & Read, 2008), but exist along a continuum from parasitic to

mutualistic (Johnson et al., 1997; Karst et al., 2008; Luo et al., 2023).

The literature holds myriad and often complimentary, but sometimes contradictory,

hypotheses that could explain the CMN mutualism as a key structural and functional component

of ecosystems. For example, the "economics" hypothesis (Kiers et al., 2011) proposes that plants

and fungi engage in "trades" of nutrients mined by fungi in exchange for plant photosynthates

(Averill et al., 2019; Fellbaum et al., 2014; Werner & Dubbert, 2016). In the economics

hypothesis, the terms of trade between plant and fungi are mediated by supply and demand for

limiting resources, which could create a dynamic market emerging from interactions between

environmental, biochemical, and biophysical variables. The "Wood Wide Web" hypothesis

emerged from the analysis of isotopically labeled carbon transferred between plants, presumably

through fungal mycorrhizae. Simard et al. (1997) hypothesized that plants that allocate carbon to

sustain common fungal symbionts also benefit from shared nutrients, while plants associating

with mycorrhizal fungi outside that network cannot. Complementing the analogy, the "kinship"

hypothesis proposes that plants of the same species preferentially receive more resources in

CMNs (Pickles et al., 2017; Tedersoo et al., 2020).

The past two decades have seen extensive but inconclusive research on these hypotheses

and how they relate to empirical measurements of CMN structure and function. On the one hand,

economic analogies suggest that the reciprocally regulated exchange of resources between plants

and fungi in CMNs should favor the most beneficial cooperative partnerships (Fellbaum et al.,

2014; Kiers et al., 2011). On the other hand, reciprocal transfer is only found in a subset of

symbionts under specific conditions, while increased competition in CMNs is a more common

observation (Walder & Van Der Heijden, 2015; Weremijewicz et al., 2016). At the core of this

controversy is whether CMNs actively support fungal resource acquisition at the expense of

plant resource demands (i.e., a fungi-centric view) or function as passive channels through which

plants regulate resource fluxes (i.e., a plant-centric view). If plant-centric, we expect to find that

the structure and functioning of CMNs give rise to consistent spatiotemporal patterns of resource

allocation similar to those predicted by the kinship hypothesis. If fungi-centric, we expect to find

that spatiotemporal patterns of resource allocation reflect the composition and functioning of the

fungal community regardless of the connecting plant nodes in CMNs. Other perspectives

emphasize that CMNs are experimentally under-documented and that this is an area that warrants

further research (Henriksson et al., 2023; Karst et al., 2023; Rillig et al., 2024; Robinson et al.,

2024). Given that data exist to support multiple, sometimes opposing views (Figueiredo et al.,

2021; Silva & Lambers, 2021); we posit that CMNs are neither plant- nor fungi-centric.

In this study, we ask if interactions among biophysical and biogeochemical processes

could explain resource transfer in CMNs with more accuracy than previous plant- or fungi-

centric analogies. We use a grassland system where fungal ASVs are frequently found within the

roots of multiple plants in a small area, a system that meets the broader CNM definition given by

Rillig et al (2024). In our study, we focus on dynamics in a system that has a high probability of

connectivity. We quantify interspecific carbon and nitrogen transfer focusing on plant traits that

are known to regulate physiological performance (Dawson et al., 2022), rather than aiming to

prove that CMNs are the only explanation. By measuring plant traits and environmental variables

that affect resource-use efficiencies across different species we describe how the transfer of

carbon and nitrogen occurs in paired experiments designed to affect soil water and nutrient mass

flow. We labeled perennial plants central to each plot (hereafter, 'donors') with stable

isotopically enriched gases and monitored leaf ^{15}N and ^{13}C for the surrounding plants

('receivers'), monitored from immediately after labeling to 21 days after. We replicated our

paired experimental setting at three different locations, with study sites distributed across a 520

km latitudinal gradient. We also sequenced strain-level variation in root fungal DNA, plant

functional types, and leaf stoichiometric traits to test if relatedness (same or different species as

the donor) explained difference in resource transfer. Both plants and fungi have economic

spectra characterized by contrasting traits and nutrient strategies which together form an

interacting continuum potentially driven by resource use and availability (Ward et al., 2022). It is

unclear to what extent plant or fungi characteristics drive these plant-fungal interactions.

Therefore, we focused on quantifying how plant-fungal interactions influence the structure and

functioning of CMNs across environmental gradients and resource constraints.

Materials and Methods:

We conducted our experiment at three sites situated on a 520 km latitudinal transect that

spans three Mediterranean climates: cool, moist (northern site; Tenino, WA) to warm, moist

(central site; Eugene, OR) to warm, dry (southern site; Selma, OR). Each circular plot was 3 m in

diameter. Half of our plots were restored prairie systems (n = 10 per site) while the other half of

the plots had introduced pasture grasses established prior to restoration (n = 10 per site).

Restored prairie plots were mowed, raked, received herbicide, and seeded in 2014-2015,

followed by seeding in fall 2015, 2016, and 2017 (Reed et al., 2019). We erected rainout shelters

that excluded 40% of the rainfall on half the plots at each site (n = 10 rain exclusion, 10 control

per site; Fig. 1). Due to the climatically driven differences in communities across sites, not all

species were present at all sites (Table S1); however, all functional groups were present at all

sites and most species were present at more than one site. Our experimental design was nested in

a multi-year experiment where data loggers were used to continuously measure temperature and

moisture in all the manipulated plots.

We recognize that there are many challenges for establishing field experiments of CMN

effects, such as treatments for severed versus intact connections (Karst et al., 2023). The plants

we experimented on grow in a shared plot where nutrients could transfer via soil, water, bacteria,

other fungal guilds, or other non-CMN mechanisms. This does not exclude the possibility that

fungi play a large role in these interactions, especially in a system where all plant species have

the potential to engage in the most common form of a mycorrhizal connection (Table S1;

Heijden et al., 2015). As Rillig et al. (2024) point out, maintaining a strict definition for CMN

limits research from exploring meaningful ecological interactions. We have approached this

experiment under their proposed broader definition of CMN, an approach which allows us to

advance our understanding of how CMNs may function in natural ecosystems.

Previous work demonstrated that the rainout shelters had minimal effects on aboveground

community structure or function (Dawson et al., 2022), possibly due to the shoulder season

effect of the Mediterranean rain seasonality. This network of experimental sites was established

in 2010 and has been extensively studied since then (Brambila et al., 2023; DeMarche et al.,

2021; Reed, Bridgham, et al., 2021; Reed et al., 2019, 2023; Reed, Peterson, et al., 2021; Reed,

Pfeifer□Meister, et al., 2021), including work on mycorrhizal fungi (Vandegrift et al., 2015;

Wilson et al., 2016). Treatments had marginal effects on the soil water potential (especially

during the early growing season). Despite those differences, there were no significant changes in

the plant community composition or productivity under rain exclusion, which also did not affect

morphological and functional traits (i.e., specific leaf area, iWUE, and C:N ratios) of the

functional groups we selected for this experiment (Dawson et al., 2022; Reed, Pfeifer☐Meister,

et al., 2021).

Isotopic labelling

At each site, we selected a healthy perennial forb (*Sidalcea malviflora* ssp. *virgata* in

restored prairie plots [except in one plot where we used *Eriophyllum lanatum* due to a lack of *S.*

malviflora ssp. virgata]), or a perennial grass (*Alopecurus pratensis, Schedonorus arundinaceus,*

or *Agrostis capillaris*) in pasture plots at the center of each plot to receive the isotopic labels. On

sunny days between 11AM and 3PM, we applied isotopically enriched carbon (^{13}C) and nitrogen

(^{15}N) as a pulse of carbon dioxide (CO_2) and ammonia (NH_3) to the leaves of target "donor"

species common across experimental sites. Although gases are not the primary source of nitrogen

for most plants, applying gaseous nitrogen allowed us to limit the amount leaked into the soil

compared to applying nitrogen directly to the soil (Silva et al., 2015). Plant leaves are known to

uptake ammonia (Farquhar et al., 1980; Sutton et al., 2008). We performed the labeling

experiment using custom-made clear chambers with internal fans, following established

protocols (e.g., Earles et al., 2016; Silva et al., 2015; Sperling et al., 2017). Before performing

the experiment, we tested our approach in the field to optimize gas exposure and labeling

amounts, which included checking for leaks and contamination outside of the chamber. We

covered the donor plant with a clear plastic cylindrical chamber and injected gas in sequence at

20 minute intervals. For $^{13}CO_2$, we made three injections of 2 mL pure CO_2 (98 atm % ^{13}C) to

double the amount of CO_2 in the chamber each time. For NH_3, we made two injections of 10 mL

pure NH_3 (98 atm% ^{15}N). The dates of application were based on peak productivity estimated

from Normalized Different Vegetation Index (NDVI) at each site (see Reed et al., 2019 for

details). We sampled leaves from each donor plant immediately after labeling (time point 0) as

well as from all plants approximately 4 days (time point 1), 10 days (time point 2), and 21 days

(time point 3) post-labelling (Fig. 1, Table S2). Time points were chosen to balance the

potentially rapid transfer of nutrients through the system with the logistical difficulties of a single

team sampling along a 520 km gradient. We also collected leaves at time points 1, 2, and 3 from

up to twelve plants in each plot representing three replicates of grass/forb structural groups and

annual/perennial life history strategies (Table S1). The number of plants and groups depended on

which plants were growing in each plot.

At the end of the experiment, we harvested entire plants and the soil surrounding the

roots at time point 3 and kept them in cool conditions until processing. We separated the roots

and rhizosphere soils, and selected approximately ten ~3 cm fine root fragments per sample (i.e.,

third order or finer, where available) for DNA extraction and identification. All roots and

rhizosphere soils were stored at -80º C until processing.

Baseline and Resource Transfer Calculations

Before isotopic labeling, we collected soil, leaves, and roots from each site. We collected

soils in late spring and early summer 2019 to 20 cm depth in each plot. From these soil samples,

we removed root fragments that represented the typical roots seen in each plot. We collected

leaves for each species in each plot; however, these leaves were contaminated with ^{15}N during

transport from the highly enriched time point 0 donor samples that were transported with them.

To replace contaminated samples, we separately sampled leaves from biomass samplings

collected in late spring and early summer 2019, ensuring that annual and perennial grasses and

forbs were represented at each site.

We oven-dried all samples at 65°C to constant mass and encapsulated them for stable

isotope analysis. All stable isotope analysis was done at UC Davis Stable Isotope Facilities. We

calculated the amount of carbon and nitrogen in each plant compartment (leaves and roots) using

standard label recovery equations (Silva et al., 2015), using baseline values measured before

application of the labelled gases to capture background variations in isotopic composition of

unenriched leaves, roots, and soil samples.

We designated all samples with greater than two standard deviations above baseline

samples as "enriched" in a particular isotope. Baseline values were calculated on a site by rain

exclusion treatment basis by plant functional type basis (Table S3). Site-specific baseline soil

and root isotope ratios represent the whole community because we were unable to differentiate

which roots belonged to which plants from our soil cores.. In all cases, baseline values fell within

the expected range for our region (Fig. S1). For each enriched sample, we calculated isotope

excess as

Equation 1

$$atm\%_{excess} = atm\%_{post\ label} - atm\%_{baseline}$$

249   For enriched donor plants, we calculated % derived from label immediately following label

250   application as

251   **Equation 2**

$$\%DFL_{donor} = \frac{atm\%_{excess}}{atm\%_{labelling\ gas} - atm\%_{baseline}} * 100$$

For enriched receiver plants, we calculated % derived from label (%NDFL and %CDFL,

respectively) for each relevant point in space and time as

Equation 3

$$\%DFL_{receiver} = \frac{atm\%_{receiver\ excess}}{atm\%_{donor\ excess}} * 100$$

When we calculated %DFL in roots, we used donor leaves as the source (atm% donor excess).

We then calculated the amount derived from label on a per mass basis as

Equation 4

$$DFL\ (\text{mg N / g leaf}) = 1000 * 1g * \left(\frac{N\ or\ C\%}{100}\right) * \left(\frac{DFL\%}{100}\right)$$

We calculated intrinsic water-use efficiency following Farquhar and Richards (1984) as

detailed in (Dawson et al., 2022) using the baseline [13]C values from the original baseline

samples. The equations given in Farquhar and Richards (1984) scale δ13C of plant tissue

samples with atmospheric data and by plant physiological constants. Because of sampling

discrepancies, we did not have intrinsic water-use efficiency data for 32 plants.

We selected a subset of rhizosphere soils that represented six donor plants at each site

divided equally between restored prairie and pasture plots, and selected the three most highly

[15]N-enriched interspecific receivers in each plot across the sites. In addition, we sampled three

most highly enriched interspecific receivers at each site and restored prairie-introduced pasture

combination. We sampled the top three enriched intraspecific receivers at each site and

treatment. In total, this came to 48 post-labelling soil samples in 29 plots.

Fungal DNA analysis

We extracted DNA from roots of 450 plants harvested at time point 3 (21 days post-label)

using Qiagen DNeasy Powersoil HTP kits (Qiagen, Hilden, Germany). We only analyzed DNA

from roots, not from the soils collected from each plant's rhizosphere. We characterized each

sample's AM fungal composition with a two-step PCR protocol that amplified a ~550bp

fragment of the SSU rRNA gene (the most well-supported region for AM fungal taxonomic

resolution (Dumbrell et al., 2011). We used WANDA (5'- CAGCCGCGGTAATTCCAGCT- 3')

and AML2 (5'- GAACCCAAACACTTTGGTTTCC-3') primers (Langmead & Salzberg, 2012;

Lee et al., 2008). We used primers with unique indices so we could multiplex several projects on

a single run. We quantified successful PCR amplicons with the Quant-iT PicoGreen dsDNA

Assay Kit (Invitrogen, Waltham, MA, USA) on a SpectraMax M5E Microplate Reader

(Molecular Devices, San Jose, CA, USA) before purifying with QIAquick PCR Purification kits

(Qiagen). We sequenced the purified pools on the Illumina MiSeq platform (paired-end 300bp,

Illumina Inc., San Diego, CA, USA) at the University of Oregon Genomics and Cell

Characterization Core Facility (Eugene, OR, USA). Reads were deduplicated with UMI-tools

using unique molecular identifiers (UMIs) inserted during PCR (Smith et al., 2017).

We assigned amplicon sequence variants (ASVs) using the dada2 pipeline (version

1.18.0) with standard quality filtering and denoising parameters (Callahan et al., 2016). The

dada2 pipeline maintains strain-level diversity at the scale of individual sequence variants rather

than clustering sequences into OTUs. This fine-scale measure of fungal sequence diversity was

particularly important for our analyses to maintain the greatest chance of detecting a single AM

fungal 'individual' in multiple plant root samples. Taxonomy was assigned to ASVs using the

MaarjAM database (2019 release) (Öpik et al., 2010). We used a Bayesian mixture model in the

DESeq2 package (Love et al., 2014) to scale ASV counts within and across samples to avoid

artificial taxon abundance biases (Anders & Huber, 2010).

Replication statement

Scale of inference	Scale at which the factor of interest is applied	Number of replicates at the appropriate scale
Individuals	Individual donors and receivers	1,441 leaves, 450 root DNA samples, and 364 root stable isotope samples from 60 donor and 502 receiver plants
Species	Functional groups	18 species with at least 3 individual plant replicates each (details in Table S1) Grouped by 61 annual forb plants, 131 annual grass plants, 81 perennial forb plants, 161 perennial grass plants
Community	Pooled by site and	3 sites (northern, central, and southern latitude sites spanning ~520 km; 20 plots per site; 60 total)
	by climate and restoration treatment	Rain exclusion (10 plots per site; 30 total) vs ambient (10 control plot per site; 30 total); Restored prairie (10 plots per site; 30 total) vs pasture (10 plots per site; 30 total) treatments;

300
Data analysis

We performed all analyses in R ver. 4.0.4 (R Core Team, 2022). Graphs were made in

ggplot2 (Wickham, 2016). We removed one outlier plant with a ^{15}N atm% more than twice as

high as the next highest measurement. We also removed five mislabeled samples. To meet

statistical assumptions, we only included data from enriched plants with successful root fungal

DNA extraction in our analyses and figures. We limited receivers to those for which we also had

sufficient root fungal DNA. In total, from 1441 leaves measured for isotopic content and with

successfully recovered fungal DNA, we analyzed data from 353 unique plants: 54 donors and

353 receivers. We excluded two plots (one central rain exclusion restored plot and one central

rain exclusion pasture plot) because either no donor or no receiver leaves were recovered.

We tested the relationship between receiver leaf %DFL and plant traits (grass/forb,

annual/perennial, iWUE, C:N, degrees of connectivity, interaction term between grass/forb and

annual/perennial) and site conditions (position on latitude gradient, pasture/restored, rain

exclusion treatment, distance from donor, time from labelling) with a mixed-effect ANOVA

(plot nested within site as random effect; Table 1). We used a Tukey post-hoc test for differences

within groups shown in the following figures and tables. We constructed a phyloseq object using

the ASV table with normalized counts (McMurdie & Holmes, 2013), and used iGraph,

metagMisc, and RCy3 (Gustavsen et al., 2019; Mikryukov, 2017; Nepusz & Csardi, 2006) to

create networks for each plot. In each network, nodes represented individual plants and edges

between nodes represent plants sharing at least one fungal DNA sequence variant. The weighted

edges are based on how many fungal ASVs were shared among plants. We calculated degrees of

connectivity with tidygraph (Petersen, 2022) to examine how many plants each individual plant

was 'connected' to (by means of shared fungal ASVs) in each plot (Fig. S2). We also calculated

whether each receiver plant shared fungal ASVs with the central donor plant in each plot. We

visualized shifts in AM fungal community composition using non-metric multi-dimensional

scaling (NMDS) in the vegan package, demonstrating the AM fungal community similarity

across plants (Oksanen, et al., 2022).

Results:

All donors were ^{15}N-enriched in their leaves at time of labelling. Two donors were not

^{13}C-enriched in their leaves at time of labelling. We recovered DNA data from the roots of

88.3% of donors. We sampled 1,444 leaves from 434 receiver plants at three time points. Of

these leaves, 81.0% were ^{15}N-enriched and 2.4% were ^{13}C-enriched. We recovered DNA from

the roots of 77.9% of the receivers. At time point 3 (~21 days post-labelling), we collected roots

from 46 of the initial 60 donors and 306 of the initial 434 receivers. Of the roots of the collected

donors, 97.8% were still 15N-enriched and 23.9% were still ^{13}C-enriched at time point 3. Of the

roots of the collected receivers, 33.0% were still ^{15}N-enriched and 10.5% were still ^{13}C-enriched

at time point 3.

Assimilation of isotopic tracers was similar between labeled "donor" plants with no

significant differences on average between sites or experimental treatments within sites,

including rainfall exclusion or restored status (Fig. S3). At all sites, foliar assimilation of ^{15}N and

^{13}C by donor plants led to enrichment levels ranging from approximately 5-10 fold higher than

baselines. Foliar enrichment levels decreased consistently at all sites and treatments over the 21

day sampling period. Annual forbs had the greatest enrichment level and perennial forbs the

lowest enrichment level (Fig. 2). We found significant spatial and temporal differences in foliar

and root isotope ratios in donors and receivers resulting from interspecific transfer of carbon and

nitrogen (Fig. S4; Table 1). Receiver foliar enrichment levels did not correlate with donor foliar

enrichment levels within the same plot (Fig. S5).

Allocation and transfer varied significantly between functional groups due to their

intrinsic differences in tissue stoichiometry (Table 1). We selected 18 common annual/perennial

and grass/forb species of receiver plants, which revealed significant differences between

functional types for NDFL (Fig. 2) but little detectable CDFL relative to baseline (Fig. S4, S6).

Rain exclusion treatment, restoration treatment, and site did not affect interspecific transfer of

nitrogen (Fig. S7; ANOVA, P > 0.05, Table 1). We observed very low carbon enrichment, but of

the 2.4% of leaves that were enriched in carbon, plant functional type and site affected carbon

transfer (Table S4). We did, however, observe significant differences in nitrogen transfer by

plant functional type (Table 1), mirroring intrinsic differences in tissue stoichiometry and iWUE

(Fig. 3), despite no significant enrichment in soils collected from the rhizosphere of those same

plants (Fig. S8). We also found that C:N affected NDFL, although not in a simple linear manner

and with no apparent correlation between NDFL and iWUE (Fig. 3). We did detect a low level of

soil enrichment in 1 out of 30 receiver soil samples (0.377 atm% ^{15}N) and 5 out of 18 donor soil

samples (ranging from 0.376 to 0.479 atm% ^{15}N; Fig. S8).

Annuals had greater ^{15}N foliar enrichment compared to perennials (ANOVA, P < 0.001,

Tables 1 and 2). Foliar enrichment decreased over both time and space (ANOVA P < 0.001; Fig.

S3, Fig. S4). On average, annuals had a lower leaf nitrogen content and higher C:N than

perennials (Table 2, Fig. 3). Forbs had higher NDFL than grasses (ANOVA, P < 0.001, Table 1)

as well as a lower C:N. There was a significant interaction between annual/perennial and

grass/forb form (ANOVA, P = 0.003, Table 1). There was no correlation in ^{15}N-enrichment and

whether the donors and receivers were the same species (Table 1).

Fungal community composition demonstrated a high degree of potential connectivity

between plants of different species but no obvious pattern of connectedness that could explain

preferential nutrient transfer by plant functional groups. We found that 97.25% ± 8.01 (SD) of all

plant roots within each experimental plot shared at least one fungal DNA sequence variant

(ASV) with another plant of the same plot. Fungal community composition was similar across

plant functional groups (Fig. 4, PERMANOVA pseudo-F statistic = 2.269, R^2 = 0.005, p =

0.001). Annual plants shared fungi with more plants in the same plot (4.74 plants ± 2.62 SD)

compared to perennials (4.04 plants ± 2.49 SD ; t-test, P = 0.019; Table S5), but degrees of

connectivity did not predict nitrogen transfer (Table 1). Seventy-three percent of plants were

colonized by four or fewer fungi and shared fungi with five or fewer other plants in the plot,

making it difficult to determine if strength of connectivity altered nitrogen transfer (Fig. S9).

Discussion:

In a system where fungal DNA sequence variants were shared between ~97% of plants,

we found that the transfer of nitrogen is regulated by plant functional traits that are known to

influence resource use and allocation in plant communities. Using a recently proposed definition

of CMNs that prioritizes ecological understanding (Rillig et al., 2024), our data suggest that the

assimilation and allocation of limiting resources in CMNs was neither plant- nor fungi-centric. In

our study, the rates and direction of resource transfer in our potential CMNs, inferred from pulse

labeling and recovery of ^{15}N in leaves and rhizospheres, could be predicted from leaf C:N and

distance from donor species (Table 1). Across all sites and treatments, we observed a stronger

sink for ^{15}N in annual plants (Table 2), indicating preferential transfer of limiting resources to

that functional group of plants. We applied the ^{15}N enriched label to perennial species in all

plots, so this greater enrichment in annual plants precludes a preference for receiver plants of the

same species as would be expected under the kinship hypothesis (Table 1). Although nearly all

plants shared fungal ASVs in their roots (Fig. 4), connectivity did not predict ^{15}N transfer (Table

1). However, we did not test hyphal continuity of our network and we draw our conclusions

based on an ecological definition of CMNs, not under Rillig et al.'s (2024) definition of CMN-

HC. Our data suggest that rates and direction of resource transfer in CMNs reflect plant nutrient

requirements and spatial proximity.

We conducted repeated spatiotemporal sampling of isotopic-enrichment levels at

increasing distances from donor species, days to weeks after labelling, and in well-established

communities exposed to multiple years of experimental treatments, expecting to find evidence of

kinship (i.e., greatest resource transfer in plants of the same species), driven by CMN economics

(i.e., ^{15}N transfer rates coinciding with ^{13}C investment in root and fungal mass). Annuals

received, on average, an order of magnitude higher enrichment than perennials even though our

donor plants were perennials. We also did not find evidence of a relationship between ^{15}N in

leaves and ^{13}C in roots because we did detect ^{15}N-enrichment in leaves but no ^{13}C-enrichment in

roots. We did, however, find significant ^{13}C-enrichment in the donor plants. Therefore, our data

do not support either hypothesis, and instead suggest AM fungi form CMNs where the rates and

direction of resource transfer ultimately reflects a sink-source strength effect, consistent with

previous observations of stoichiometric source-sink manipulations of carbon and nitrogen within

plants (Cai et al., 2021; Ruiz-Vera et al., 2017; Tegeder & Masclaux☐Daubresse, 2018), but in

our case observed at the community scale.

Nitrogen enrichment levels remained high in leaves and many roots at the end of the

experiment, allowing us to measure NDFL across the community and infer the main predictors

of N transfer. However, carbon enrichment levels faded before plants were harvested

approximately 21 days post-labelling. After controlling for variation in assimilation rates by

calculating NDFL, we found that annual plants received greater ^{15}N-enrichment than perennial

plants. Plants closest to the donor were most enriched, and ^{15}N-enrichment decreased over time

(Fig. S4, Fig. S5). Although the rainout shelters had limited effect, there were major differences

across the latitudinal gradient represented by the sites in temperature and soil moisture

availability (Dawson et al., 2022; Reed et al., 2019); however, neither treatment nor site affected

our results.

The major predictors of differences in allocation of ^{13}C and ^{15}N to roots and subsequent

transfer to "receiver" species were the intrinsic difference in plant functional types and correlated

traits, including measured leaf C:N. Previous studies in northern California under environmental

conditions similar to those found in our southernmost experimental site showed rapid (days to

weeks) transfer of ^{15}N applied to the leaves of ectomycorrhizal pines to surrounding annual AM

plant receivers (He et al., 2006). Those results demonstrated "direct fungal connections are not

necessary for N transfer among plants" and, similar to our results, the "leaves of the annual

plants had greater ^{15}N derived from source and were more enriched (^{15}N at % excess and δ^{15}N

values) than perennial receivers, irrespective of the mycorrhizal type." Similarly, as proposed by

He et al. (2006), our data suggest that annual plants were a strong sink for N which could be

explained by stoichiometric gradients that affect root exudation and recapture of N-containing

materials from rhizodeposition (Høgh-Jensen & Schjoerring, 2001; Mayer et al., 2003).

Although our study included two species with symbiotic N fixation ability that could have

altered some of the baseline data, even if a plot was fully dominated by legumes that difference

would represent a minor effect relative to the pulse label application. Our data corroborate rapid

transfer among AM plants, with no detectable enrichment in root or soil ^{13}C near roots 21 days

post-labelling, but do not allow us to determine general mechanisms that are responsible for the

^{15}N transfers. Other methods of transfer–such as by fungi of other functional guilds, bacteria, or

water flow in the soil–were possible given that plants shared a common growing medium in each

plot.

We inferred a high connectivity between plants within each given treatment and site

given the highly similar fungal composition in the root systems of both perennial and annual

plants (Fig. 4). Given the constraints of ASV-identified data, we did this analysis on a strain-

level scale and it is possible that separate spores of the same ASV separately infected plants

within the same plot. However, we are reasonably confident in our use of fungal ASVs as a

proxy for connectivity given the strong overlap in our community and because individuals of one

ASV can anastomose in the soil (Mikkelsen et al., 2008). This overlap could explain the lack of

support for the kinship hypothesis in our dataset and offers further support for stoichiometric

gradients in general, and C:N gradients in particular, as a principal control of terms of trade in

CMNs (Kiers et al., 2011). Because we observed such high rates of shared ASVs (~97%) and we

observed high levels of N-enrichment in receivers at our first post-label sampling point four days

after application, we could not test whether receivers connected to donors or whether plants were

connected to the network affected ^{15}N transfer. We found unexpectedly low soil isotopic-

enrichment (fig. S8) which suggested that the labels did not remain in the fungal network. This

low level was likely driven by the fact that we did not sample soils until 21 days after labeling

and hyphal turnover for grass-associated AM fungi can be less than one month (See et al., 2022).

Because N is a major limiting nutrient in this system, according to the current paradigm of

CMNs, it would be quickly taken up and recycled or transferred rather than accumulating in the

soil.

We found that plant-soil stoichiometric gradients and functional traits were the strongest

predictors of resource sharing in a possible grassland arbuscular mycorrhizal CMN. We interpret

this finding as evidence of biochemical and biophysical sinks, in which nutrients are allocated to

plants with the greatest need for those nutrients, either through a 'passive' mycorrhizal network

as suggested by the high number of shared fungal ASVs or direct uptake from soil or water flow.

Expanding on previous studies, we propose that AM fungi facilitate spatiotemporal dynamics of

carbon and nitrogen transfer through CMNs in ways that are neither plant- nor fungi-centric.

That is, plants and fungi that are located closer together in space and with stronger demand for

resources over time are more likely to receive larger amounts of those limiting resources.

References

Anders, S., & Huber, W. (2010). Differential expression analysis for sequence count data.

Genome Biology, *11*(10), R106. https://doi.org/10.1186/gb-2010-11-10-r106

Averill, C., Bhatnagar, J. M., Dietze, M. C., Pearse, W. D., & Kivlin, S. N. (2019). Global

imprint of mycorrhizal fungi on whole-plant nutrient economics. *Proceedings of the*

National Academy of Sciences, *116*(46), 23163–23168.

https://doi.org/10.1073/pnas.1906655116

Brambila, A., Reed, P. B., Bridgham, S. D., Roy, B. A., Johnson, B. R., Pfeifer-Meister, L., &

Hallett, L. M. (2023). Disturbance: A double-edged sword for restoration in a changing

climate. *Restoration Ecology*, *31*(1), e13675. https://doi.org/10.1111/rec.13675

Brundrett, M. C. (2009). Mycorrhizal associations and other means of nutrition of vascular

plants: Understanding the global diversity of host plants by resolving conflicting

information and developing reliable means of diagnosis. *Plant and Soil*, *320*(1–2), 37–77.

https://doi.org/10.1007/s11104-008-9877-9

Cai, Z., Xie, T., & Xu, J. (2021). Source–sink manipulations differentially affect carbon and

nitrogen dynamics, fruit metabolites and yield of Sacha Inchi plants. *BMC Plant Biology*,

21(1), 160. https://doi.org/10.1186/s12870-021-02931-9

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P.

(2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature*

Methods, *13*(7), 581–583. https://doi.org/10.1038/nmeth.3869

Davison, J., García De León, D., Zobel, M., Moora, M., Bueno, C. G., Barceló, M., Gerz, M.,

León, D., Meng, Y., Pillar, V. D., Sepp, S., Soudzilovaskaia, N. A., Tedersoo, L.,

Vaessen, S., Vahter, T., Winck, B., & Öpik, M. (2020). Plant functional groups associate

with distinct arbuscular mycorrhizal fungal communities. *New Phytologist*, *226*(4), 1117–

1128. https://doi.org/10.1111/nph.16423

Dawson, H. R., Maxwell, T. M., Reed, P. B., Bridgham, S. D., & Silva, L. C. R. (2022). Leaf

traits predict water□use efficiency in U.S. Pacific Northwest grasslands under rain

exclusion treatment. *Journal of Geophysical Research: Biogeosciences*, *127*(10),

e2022JG007060. https://doi.org/10.1029/2022JG007060

DeMarche, M. L., Bailes, G., Hendricks, L. B., Pfeifer□Meister, L., Reed, P. B., Bridgham, S.

D., Johnson, B. R., Shriver, R., Waddle, E., Wroton, H., Doak, D. F., Roy, B. A., &

Morris, W. F. (2021). Latitudinal gradients in population growth do not reflect

demographic responses to climate. *Ecological Applications*, *31*(2), e2242.

https://doi.org/10.1002/eap.2242

Dumbrell, A. J., Ashton, P. D., Aziz, N., Feng, G., Nelson, M., Dytham, C., Fitter, A. H., &

Helgason, T. (2011). Distinct seasonal assemblages of arbuscular mycorrhizal fungi

revealed by massively parallel pyrosequencing. *New Phytologist*, *190*(3), 794–804.

https://doi.org/10.1111/j.1469-8137.2010.03636.x

Earles, J. M., Sperling, O., Silva, L. C. R., McElrone, A. J., Brodersen, C. R., North, M. P., &

Zwieniecki, M. A. (2016). Bark water uptake promotes localized hydraulic recovery in

coastal redwood crown. *Plant, Cell & Environment*, *39*(2).

       https://doi.org/10.1111/pce.12612

Farquhar, G. D., Firth, P. M., Wetselaar, R., & Weir, B. (1980). On the gaseous exchange of

       ammonia between leaves and the environment: Determination of the ammonia

       compensation point. *Plant Physiology*, *66*(4), 710–714.

       https://doi.org/10.1104/pp.66.4.710

Farquhar, G. D., & Richards, R. A. (1984). Isotopic composition of plant carbon correlates with

       water-use efficiency of wheat genotypes. *Functional Plant Biology*, *11*(6), 539–552.

Fellbaum, C. R., Mensah, J. A., Cloos, A. J., Strahan, G. E., Pfeffer, P. E., Kiers, E. T., &

       Bücking, H. (2014). Fungal nutrient allocation in common mycorrhizal networks is

       regulated by the carbon source strength of individual host plants. *New Phytologist*,

       203(2), 646–656. https://doi.org/10.1111/nph.12827

Figueiredo, A. F., Boy, J., & Guggenberger, G. (2021). Common mycorrhizae network: A

       review of the theories and mechanisms behind underground interactions. *Frontiers in*

       Fungal Biology, *2*, 735299. https://doi.org/10.3389/ffunb.2021.735299

Gustavsen, J. A., Pai, S., Isserlin, R., Demchak, B., & Pico, A. R. (2019). Rcy3: Network

       biology using cytoscape from within R. *F1000Research*, *8*, 1774.

       https://doi.org/10.12688/f1000research.20887.3

He, X., Bledsoe, C. S., Zasoski, R. J., Southworth, D., & Horwath, W. R. (2006). Rapid nitrogen

       transfer from ectomycorrhizal pines to adjacent ectomycorrhizal and arbuscular

       mycorrhizal plants in a California oak woodland. *New Phytologist*, *170*(1), 143–151.

       https://doi.org/10.1111/j.1469-8137.2006.01648.x

Heijden, M. G. A., Martin, F. M., Selosse, M., & Sanders, I. R. (2015). Mycorrhizal ecology and

evolution: The past, the present, and the future. *New Phytologist*, *205*(4), 1406–1423.

https://doi.org/10.1111/nph.13288

Henriksson, N., Marshall, J., Högberg, M. N., Högberg, P., Polle, A., Franklin, O., & Näsholm,

T. (2023). Re□examining the evidence for the mother tree hypothesis – resource sharing

among trees via ectomycorrhizal networks. *New Phytologist*, *239*(1), 19–28.

https://doi.org/10.1111/nph.18935

Høgh-Jensen, H., & Schjoerring, J. K. (2001). Rhizodeposition of nitrogen by red clover, white

clover and ryegrass leys. *Soil Biology and Biochemistry*, *33*(4–5), 439–448.

https://doi.org/10.1016/S0038-0717(00)00183-8

Johnson, N. C., Graham, J., & Smith, F. A. (1997). Functioning of mycorrhizal associations

along the mutualism–parasitism continuum. *New Phytologist*, *135*(4), 575–585.

https://doi.org/10.1046/j.1469-8137.1997.00729.x

Karst, J., Jones, M. D., & Hoeksema, J. D. (2023). Positive citation bias and overinterpreted

results lead to misinformation on common mycorrhizal networks in forests. *Nature*

Ecology & Evolution, *7*(4), 501–511. https://doi.org/10.1038/s41559-023-01986-1

Karst, J., Marczak, L., Jones, M. D., & Turkington, R. (2008). The Mutualism–Parasitism

Continuum in Ectomycorrhizas: A Quantitative Assessment Using Meta-Analysis.

Ecology, *89*(4), 1032–1042. https://doi.org/10.1890/07-0823.1

Kenrick, P., & Strullu-Derrien, C. (2014). The Origin and Early Evolution of Roots. *Plant*

Physiology, *166*(2), 570–580. https://doi.org/10.1104/pp.114.244517

Kiers, E. T., Duhamel, M., Beesetty, Y., Mensah, J. A., Franken, O., Verbruggen, E., Fellbaum,

C. R., Kowalchuk, G. A., Hart, M. M., Bago, A., Palmer, T. M., West, S. A.,

Vandenkoornhuyse, P., Jansa, J., & Bücking, H. (2011). Reciprocal rewards stabilize

cooperation in the mycorrhizal symbiosis. *Science*, *333*(6044), 880–882.

https://doi.org/10.1126/science.1208473

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature*

Methods, *9*(4), 357–359. https://doi.org/10.1038/nmeth.1923

Lee, J., Lee, S., & Young, J. P. W. (2008). Improved PCR primers for the detection and

identification of arbuscular mycorrhizal fungi: PCR primers for arbuscular mycorrhizal

fungi. *FEMS Microbiology Ecology*, *65*(2), 339–349. https://doi.org/10.1111/j.1574-

6941.2008.00531.x

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550.

https://doi.org/10.1186/s13059-014-0550-8

Luo, X., Liu, Y., Li, S., & He, X. (2023). Interplant carbon and nitrogen transfers mediated by

common arbuscular mycorrhizal networks: Beneficial pathways for system functionality.

Frontiers in Plant Science, *14*, 1169310. https://doi.org/10.3389/fpls.2023.1169310

Mayer, J., Buegger, F., Jensen, E. S., Schloter, M., & Heß, J. (2003). Estimating N

rhizodeposition of grain legumes using a 15N in situ stem labelling method. *Soil Biology*

and Biochemistry, *35*(1), 21–28. https://doi.org/10.1016/S0038-0717(02)00212-2

McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R package for reproducible interactive

analysis and graphics of microbiome census data. *PLoS ONE*, *8*(4), e61217.

https://doi.org/10.1371/journal.pone.0061217

Mikkelsen, B. L., Rosendahl, S., & Jakobsen, I. (2008). Underground resource allocation

between individual networks of mycorrhizal fungi. *New Phytologist*, *180*(4), 890–898.

https://doi.org/10.1111/j.1469-8137.2008.02623.x

Mikryukov, V. (2017). *vmikk/metagMisc: V.0.0.0.9000* (v.0.0.0.9000) [Computer software].

https://doi.org/10.5281/ZENODO.571403

Nepusz, T., & Csardi, G. (2006). *The igraph software package for complex network research*

[R]. https://cran.r-project.org/web/packages/igraph

Oksanen J, Simpson G, Blanchet F, Kindt R, Legendre P, Minchin P, O'Hara R, Solymos P,

Stevens M, Szoecs E, Wagner H, Barbour M, Bedward M, Bolker B, Borcard D,

Carvalho G, Chirico M, De Caceres M, Durand S, … Weedon J. (2022). *vegan:*

Community Ecology Package (2.6-2) [Computer software]. https://CRAN.R-

project.org/package=vegan

Öpik, M., Vanatoa, A., Vanatoa, E., Moora, M., Davison, J., Kalwij, J. M., Reier, Ü., & Zobel,

M. (2010). The online database Maarj *AM* reveals global and ecosystemic distribution

patterns in arbuscular mycorrhizal fungi (Glomeromycota). *New Phytologist*, *188*(1),

223–241. https://doi.org/10.1111/j.1469-8137.2010.03334.x

Petersen, T. (2022). *tidygraph: A Tidy API for Graph Manipulation* [Computer software].

https://tidygraph.data-imaginist.com/authors.html

Pickles, B. J., Wilhelm, R., Asay, A. K., Hahn, A. S., Simard, S. W., & Mohn, W. W. (2017).

Transfer of [13]C between paired Douglas☐fir seedlings reveals plant kinship effects and

uptake of exudates by ectomycorrhizas. *New Phytologist*, *214*(1), 400–411.

https://doi.org/10.1111/nph.14325

R Core Team. (2022). *R: A language and environment for statistical computing.* [R]. R

Foundation for Statistical Computing. https://www.R-project.org/

Reed, P. B., Assour, H. R., Okotie-Oyekan, A., Bailes, G. T., Johnson, B. R., Nelson, A. A.,

Pfeifer-Meister, L., Roy, B. A., & Bridgham, S. D. (2023). Climate Effects on Prairie

Productivity Partially Ameliorated by Soil Nutrients and Plant Community Responses.

Ecosystems, *26*(5), 983–999. https://doi.org/10.1007/s10021-022-00811-3

Reed, P. B., Bridgham, S. D., Pfeifer☐Meister, L. E., DeMarche, M. L., Johnson, B. R., Roy, B.

A., Bailes, G. T., Nelson, A. A., Morris, W. F., & Doak, D. F. (2021). Climate warming

threatens the persistence of a community of disturbance☐adapted native annual plants.

Ecology, *102*(10), e03464. https://doi.org/10.1002/ecy.3464

Reed, P. B., Peterson, M. L., Pfeifer☐Meister, L. E., Morris, W. F., Doak, D. F., Roy, B. A.,

Johnson, B. R., Bailes, G. T., Nelson, A. A., & Bridgham, S. D. (2021). Climate

manipulations differentially affect plant population dynamics within versus beyond

northern range limits. *Journal of Ecology*, *109*(2), 664–675. https://doi.org/10.1111/1365-

2745.13494

Reed, P. B., Pfeifer☐Meister, L. E., Roy, B. A., Johnson, B. R., Bailes, G. T., Nelson, A. A.,

Boulay, M. C., Hamman, S. T., & Bridgham, S. D. (2019). Prairie plant phenology driven

more by temperature than moisture in climate manipulations across a latitudinal gradient

in the Pacific Northwest, USA. *Ecology and Evolution*, *9*(6), 3637–3650.

https://doi.org/10.1002/ece3.4995

Reed, P. B., Pfeifer☐Meister, L. E., Roy, B. A., Johnson, B. R., Bailes, G. T., Nelson, A. A., &

Bridgham, S. D. (2021). Introduced annuals mediate climate☐driven community change

in Mediterranean prairies of the Pacific Northwest, USA. *Diversity and Distributions*,

27(12), 2584–2595. https://doi.org/10.1111/ddi.13426

Rillig, M. C., Lehmann, A., Lanfranco, L., Caruso, T., & Johnson, D. (2024). Clarifying the

definition of common mycorrhizal networks. *Functional Ecology*, *00*, 1–7.

https://doi.org/10.1111/1365-2435.14545

Robinson, D. G., Ammer, C., Polle, A., Bauhus, J., Aloni, R., Annighöfer, P., Baskin, T. I., Blatt,

M. R., Bolte, A., Bugmann, H., Cohen, J. D., Davies, P. J., Draguhn, A., Hartmann, H.,

Hasenauer, H., Hepler, P. K., Kohnle, U., Lang, F., Löf, M., … Näsholm, T. (2024).

Mother trees, altruistic fungi, and the perils of plant personification. *Trends in Plant*

Science, *29*(1), 20–31. https://doi.org/10.1016/j.tplants.2023.08.010

Ruiz-Vera, U. M., De Souza, A. P., Long, S. P., & Ort, D. R. (2017). The role of sink strength

and nitrogen availability in the down-regulation of photosynthetic capacity in field-grown

Nicotiana tabacum L. at elevated co2 concentration. *Frontiers in Plant Science*, *8*, 998.

https://doi.org/10.3389/fpls.2017.00998

See, C. R., Keller, A. B., Hobbie, S. E., Kennedy, P. G., Weber, P. K., & Pett☐Ridge, J. (2022).

Hyphae move matter and microbes to mineral microsites: Integrating the hyphosphere

into conceptual models of soil organic matter stabilization. *Global Change Biology*,

28(8), 2527–2540. https://doi.org/10.1111/gcb.16073

Selosse, M.-A., Richard, F., He, X., & Simard, S. W. (2006). Mycorrhizal networks: Des liaisons

dangereuses? *Trends in Ecology & Evolution*, *21*(11), 621–628.

https://doi.org/10.1016/j.tree.2006.07.003

Silva, L. C. R., & Lambers, H. (2021). Soil-plant-atmosphere interactions: Structure, function,

and predictive scaling for climate change mitigation. *Plant and Soil*, *461*, 5–27.

Silva, L. C. R., Salamanca-Jimenez, A., Doane, T. A., & Horwath, W. R. (2015). Carbon dioxide

level and form of soil nitrogen regulate assimilation of atmospheric ammonia in young

trees. *Scientific Reports*, *5*(1), 13141. https://doi.org/10.1038/srep13141

Simard, S. W., Perry, D. A., Jones, M. D., Myrold, D. D., Durall, D. M., & Molina, R. (1997).

Net transfer of carbon between ectomycorrhizal tree species in the field. *Nature*,

388(6642), 579–582. https://doi.org/10.1038/41557

Smith, T., Heger, A., & Sudbery, I. (2017). UMI-tools: Modeling sequencing errors in Unique

Molecular Identifiers to improve quantification accuracy. *Genome Research*, *27*(3), 491–

499. https://doi.org/10.1101/gr.209601.116

Sperling, O., Silva, L. C. R., Tixier, A., Théroux-Rancourt, G., & Zwieniecki, M. A. (2017).

Temperature gradients assist carbohydrate allocation within trees. *Scientific Reports*,

7(1), 3265. https://doi.org/10.1038/s41598-017-03608-w

Sutton, M. A., Erisman, J. W., Dentener, F., & Möller, D. (2008). Ammonia in the environment:

From ancient times to the present. *Environmental Pollution*, *156*(3), 583–604.

https://doi.org/10.1016/j.envpol.2008.03.013

Tedersoo, L., Bahram, M., & Zobel, M. (2020). How mycorrhizal associations drive plant

population and community biology. *Science*, *367*(6480), eaba1223.

https://doi.org/10.1126/science.aba1223

Tegeder, M., & Masclaux☐Daubresse, C. (2018). Source and sink mechanisms of nitrogen

transport and use. *New Phytologist*, *217*(1), 35–53. https://doi.org/10.1111/nph.14876

Vives-Peris, V., De Ollas, C., Gómez-Cadenas, A., & Pérez-Clemente, R. M. (2020). Root

exudates: From plant to rhizosphere and beyond. *Plant Cell Reports*, *39*(1), 3–17.

https://doi.org/10.1007/s00299-019-02447-5

Walder, F., & Van Der Heijden, M. G. A. (2015). Regulation of resource exchange in the

arbuscular mycorrhizal symbiosis. *Nature Plants*, *1*(11), 15159.

https://doi.org/10.1038/nplants.2015.159

Ward, E. B., Duguid, M. C., Kuebbing, S. E., Lendemer, J. C., & Bradford, M. A. (2022). The

functional role of ericoid mycorrhizal plants and fungi on carbon and nitrogen dynamics

in forests. *New Phytologist*, *235*(5), 1701–1718. https://doi.org/10.1111/nph.18307

Weremijewicz, J., Sternberg, L. D. S. L. O., & Janos, D. P. (2016). Common mycorrhizal

networks amplify competition by preferential mineral nutrient allocation to large host

plants. *New Phytologist*, *212*(2), 461–471. https://doi.org/10.1111/nph.14041

Werner, C., & Dubbert, M. (2016). Resolving rapid dynamics of soil–plant–atmosphere

interactions. *New Phytologist*, *210*(3), 767–769. https://doi.org/10.1111/nph.13936

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis* (2nd ed. 2016). Springer

International Publishing: Imprint: Springer. https://doi.org/10.1007/978-3-319-24277-4

Figure 1. Schema of experimental set up, sampling, and effects of rainout shelters. Figure

shows the three location of the sites (orange points on map) in relation to the climatic gradient

(cool to warm, wet to dry). Each plot had a central perennial plant that was labeled with

isotopically enriched gas (shown with a grey cylinder here) and we sampled plants of each

functional type at three distances from the central labelled plant. Inset shows the temporal

sampling scheme of leaves, roots and soil.

Figure 2. Annuals had greater leaf nitrogen derived from label (NDFL) scaled by mass

than perennials. Points represent enriched individual receiver plants with associated DNA data

at one time point of sampling. Y-axis is log10 scale with raw value datapoints. Boxplots show

medians and interquartile ranges, lines show the largest (or smallest) value no further than 1.5x

IQR from the hinge. Letters above the boxplots indicate significant differences.

Figure 3. Stoichiometric and functional traits compared to nitrogen derived from transfer.

A) Leaf C:N by annual/perennial. B) Percent [15]N derived from label (DFL) compared to C:N in

leaves. C) iWUE by annual/perennial as measured before labelling. D) Percent [15]NDFL

compared to iWUE in leaves at all time points. Boxplots show medians and interquartile ranges,

703   lines show the largest (or smallest) value no further than 1.5x IQR from the hinge. Letters above

704   boxplots indicate significant differences.

Figure 4. NMDS of fungal communities. Each point represents an individual plant ordinated by
the Bray Curtis dissimilarities for AM fungal composition. PERMANOVA pseudo-F statistic =
2.269, R^2 = 0.005, p = 0.001.

710　**Table 1. Mixed-effects ANOVA results effects on leaf nitrogen derived from label**

711　**(%NDFL).** Random effect is plot nested within site. Only enriched receiver leaves with

712　associated DNA data were included in the analysis (n = 1094). Leaf %CDFL results available in

713　Table S4.

	F-statistic	DF	P-value
Annual/perennial	9.81	**1**	**<0.001**
Grass/forb	11.0	**1**	**<0.001**
Same species as donor	0.47	1	0.463
Degree of connectivity	0.02	1	0.68
iWUE	0.03	1	0.81
C:N	35.51	**1**	**<0.001**
Site	0.40	2	0.71
Drought treatment	1.09	1	0.48
Restoration treatment	0.85	1	0.34
Distance from donor	9.61	**1**	**0.002**
Time from labelling	83.59	**1**	**<0.001**
Annual/perennial: Grass/forb interaction	7.75	**1**	**0.01**

716 **Table 2. Leaf nitrogen derived from label (NDFL) and leaf tissue nitrogen (N%) four days**

717 **after labeling.**

			%NDFL				NDFL (mg N/g leaf)				N (%)			
		n	*Mean*	*SD*	*Min*	*Max*	*Mean*	*SD*	*Min*	*Max*	*Mean*	*SD*	*Min*	*Max*
Annual	**Forb**	54	6.528	9.769	0.101	56.005	0.749	0.819	0.041	3.858	0.749	0.819	0.041	3.858
	Grass	92	4.847	9.032	0.098	63.884	0.34	0.463	0.011	2.79	0.340	0.463	0.011	2.790
Perennial	**Forb**	67	0.952	1.069	0.090	5.582	0.214	0.228	0.021	1.156	0.214	0.228	0.021	1.156
	Grass	148	1.632	3.066	0.088	22.434	0.252	0.484	0.013	2.948	0.252	0.484	0.013	2.948