Title: Agnostic fungi: plant traits and tissue stoichiometry explain nutrient transfer in common

arbuscular mycorrhizal networks of temperate grasslands

Authors: Hilary Rose Dawson[1], Katherine L. Shek[1,2], Toby M. Maxwell[1,3], Paul B. Reed[1,4],

Barbara Bomfim[1,5], Scott Bridgham[1], Brendan Bohannan[1], Lucas C.R. Silva[1,6]

Affiliations:

1. Department of Biology, University of Oregon, Eugene, OR 97403 USA

2. Department of Natural Resources and the Environment, University of New Hampshire,

Durham NH

3. Department of Biological Sciences, Boise State University, Boise, Idaho, 83725 USA

4. Institute for Applied Ecology, Corvallis, OR 97333 USA

5. Climate and Ecosystem Sciences Division, Lawrence Berkeley National Laboratory, Berkeley,

CA 94720 USA

6. Environmental Studies Program, University of Oregon, Eugene, OR 97403 USA

Abstract:

Plants and mycorrhizal fungi form close mutualistic relationships that affect the structure and

function of ecosystems. Recent studies show how certain plant communities can form

associations with the same mycorrhizal fungus leading to preferential transfer of carbon and

limiting nutrients, such as nitrogen, between different species through common mycorrhizal

networks. This phenomenon has been documented in a wide range of ecosystems but its

mechanisms remain poorly understood. In an experimental prairie-pasture plant system,

replicated at three sites across a 520 km latitudinal gradient in the Pacific Northwest, USA, we

identified general processes that reconcile competing hypotheses related to resource transfer in

common mycorrhizal networks. We used stable isotope tracers, paired with analyses of fungal

DNA and plant and fungal community structure and traits, to quantify above- and belowground

allocation and transfer of carbon and nitrogen between 18 different plant species. We applied

isotopically enriched carbon (^{13}C) and nitrogen (^{15}N) as a pulse of carbon dioxide (CO_2) and

ammonia (NH_3) to the leaves of target "donor" species common across experimental sites. At

each site, we performed >1360 measurements of foliar and root isotopic enrichment, which

revealed morphological traits and tissue stoichiometry as the most important predictors of

interspecific resource transfer. Assimilation of isotopic tracers was similar between labeled

"donor" plants with no significant difference between sites for labeled "donor" plants with no

significant difference in transfer between species among experimental treatments, including

rainfall exclusion or species diversity. However, allocation of carbon and nitrogen to roots and

transfer to "receiver" species varied significantly between functional groups (annual/perennial or

grass/forb) due to differences in tissue stoichiometry. Our findings points to a simple mechanistic

answer for long-standing questions regarding mutualism and transfer of resources between plants

via mycorrhizal networks, an explanation that can lead to general predictions of preferential flow

of limiting resources in other ecosystems.

Introduction

Plant-mycorrhizal associations are thought to have emerged as rudimentary root systems over 400 million years ago, facilitating the expansion of terrestrial life that followed (Kenrick & Strullu-Derrien, 2014). The transformative power of early fungal symbioses is still evident today in all major plant lineages, from bryophytes to angiosperms, all of which form mycorrhizal connections (van der Heijden et al. 2015). Over 85% of all contemporary flowering plant species form symbioses with fungi, with arbuscular mycorrhizal (AM) associations being the most common (Brundrett 2009). Today, the relationships between plants and AM fungi dominates managed and unmanaged landscapes and are estimated to be responsible for up to 80% of global primary productivity (van der Heijden et al. 2015). Fungi are gregarious and form partnerships with more than one individual plant (Selosse et al. 2006). By extension, it has been widely hypothesized that the multi-plant-fungal relationships form "common mycorrhizal networks" (CMNs) which facilitate carbon and nutrient transfer between organisms, beyond the immediate plant-fungus mutualism formed by individuals. This relationship is often simplified to the mutual exchange of carbon-based compounds created by plant photosynthesis for limiting soil nutrients mined by fungi (Smith and Read 2008). However, a synthesis of empirical research at macro- and micro-levels of ecological organization reveals a gap in understanding of structural and functional properties that could predict how carbon and nutrient are transferred in CMNs (Silva and Lambers 2021).

The literature holds myriad (often complimentary, but sometimes contradictory) hypotheses that could explain the CMN mutualism as a key structural and functional component of ecosystems. For example, the "Wood Wide Web" hypothesis emerged from the analysis of isotopically labeled carbon transferred between plants, presumably through fungal mycorrhizae (Simard et al. 1997). This observation set the foundation for economic analogies based on the idea that certain plants which allocate carbon to sustain common fungal symbionts can also benefit from shared nutrients, while plants associating with different mycorrhizal fungi cannot. The "economics" hypothesis (Kiers et al. 2011) proposes that plants and fungi engage in "trades" of nutrients mined by fungi in exchange for carbon-based photosynthates from plants (Fellbaum et al. 2014, Werner and Dubbert 2016, Averill et al. 2019). In this hypothesis, the terms of trade between plant and fungi are mediated by supply and demand for limiting resources, which could create a dynamic market emerging from interactions between environmental, biochemical, and

biophysical variables. Complementing the economic analogy, the "kinship" hypothesis proposes

that phylogenetic distance within or across species promotes functional gradients and preferential

flow of resources in CMNs (Tedersoo et al. 2020), in which transfer of carbon and nutrients are

predicted to be more frequent and abundant between related individuals (e.g., seedlings and trees

of the same species) than between unrelated individuals (Pickles et al. 2017).

The past two decades have seen extensive but inconclusive research on these hypotheses

and how they relate to empirical measurements of CMN structure and function. There is a

conspicuous ambiguity separating theory and empirical observations. On the one hand, economic

analogies suggest that the reciprocally regulated exchange of resources between plants and fungi

in CMNs should favor the most beneficial cooperative partnerships (Kiers et al. 2011, Fellbaum

et al. 2014). On the other hand, reciprocal transfer is only found in a subset of symbionts under

specific conditions, while amplified competition in CMNs is a more common observation

(Walder and van der Heijden 2015, Weremijewicz et al. 2016). At the core of this controversy is

the question of whether CMNs actively support fungal growth at the expense of plant resource

demands (i.e., a fungi-centric view) or function as passive channels through which plants

regulate resource fluxes (i.e., a plant-centric view). Data exist to support both of these opposing

views (see Silva and Lambers 2021, Figueiredo et al. 2021); therefore we posit that CMNs are

neither plant- nor fungi-centric. Perhaps more than gregarious, AM fungi might be "agnostic"

with respect to plant species partnerships, which would imply that that the rates and direction of

resource transfer in AM CMNs are ultimately a reflection of sink-source strength gradients.

To test the "agnostic fungi" hypothesis, we investigated how interactions between general

biophysical and biogeochemical processes operating at the soil-plant-atmosphere interface could

explain resource transfer in CMNs better than previous plant- or fungi-centric analogies. We

studied how soil resource-use efficiency and plant traits that regulate physiological performance

affect the transfer of carbon and nitrogen in paired experiments where we also measured fungal

DNA, plant community structure, and plant traits to test the effects of relatedness in resource

transfer. Our experimental setting includes high- and low-diversity prairie and pasture

experimental sites under ambient conditions and rain exclusion shelters, designed to affect soil

water and nutrient mass flow, replicated at three different sites across a 520 km latitudinal

gradient. Due to the latitudinal and natural climatic difference in communities across sites, not all

species were present at all sites (Table S1); however, all "donor" species and all "receiver"

functional groups were present at all sites. This allowed us to draw general inferences about the
structure and function of AM CMNs in temperate grasslands under a broad range on ecological
and environmental conditions.

Methods

We conducted our experiment at three sites situated on a 520 km latitudinal transect that
spans cool, moist (northern site; Tenino, WA) to warm, moist (central site; Eugene, OR) to
warm, dry (southern site; Selma, OR) climates. Half of our plots were manipulated to restore a
high-diversity prairie community (n = 10 per site) while the other half of the plots were the
existing introduced low-diversity pasture grass community (n = 10 per site). Manipulated high-
diversity plots were mowed, raked, received herbicide, and seeded in 2014-2015, followed by
seeding in fall 2015, 2016, and 2017 (Reed et al. 2019). We erected rainout shelters that
excluded 40% of the rainfall on half the plots at each site (n = 10 rain exclusion, 10 control per
site, except the central site where there were only 8 rain exclusion plots; Fig. 1).
Plots were either planted high diversity prairie communities or previously existing
pasture communities, and all plants in our system have the potential to form associations with
AM fungi. Half the plots were under rainout shelters that had little to no effect on aboveground
community structure or function, where significant differences exist between plant functional
groups with respect to ecophysiological performance and tissue stoichiometry (Dawson et al. *In
Press*). We labeled perennial plants central to each plot (hereafter, 'donors') and monitored leaf
^{15}N and ^{13}C for the surrounding plants ('receivers') for up to 21 days. In addition, we sampled
roots and soils at 21 days post-labelling. We analyzed fungal DNA collected from the root
samples, as well as stable isotopes in all samples. Our experimental design was nested in a multi-
year experiment where in situ data loggers were used to continuously measure variables in all the
manipulated plots. We calculated soil matric potentials to account for soil differences between
sites (Saxton and Rawls 2006). Rain exclusion only changed soil temperature at the northern site
(Fig. 1c) and soil matric potential at the central site (Fig. 1d).
This network of sites has been established since 2010 and has been extensively studied
since then (Reed et al. 2019, 2021a, 2021b, 2021c, Peterson et al. 2020) including work on
mycorrhizal fungi (Wilson et al. 2016). Previous results show that the rain exclusion treatment
significantly changed soil water potential, which should also affect nutrient uptake through mass

flow. At each site, pasture plots that have not been seeded by additional species adjoin the high
diversity plots. These are dominated by one to a few species of introduced perennial grasses
which we targeted as donors in pasture plots. For high diversity plots, we targeted native
perennial species that are common to ambient and drought treatments at all sites, and that have
shown divergent changes in productivity in response to a warming-induced decline in
mycorrhizal colonization attributed to that treatment's drying effect (Wilson et al. 2016). Despite
those differences, we did not find significant changes in the average plant community
composition or productivity under rain exclusion, which also did not affect morphological and
functional traits (e.g. specific leaf area, iWUE, and C:N ratios) of the functional groups we
selected for this experiment (Dawson in press) (Reed et al. 2021c). Therefore, this experimental
setting is ideal to test general hypotheses related to how plant-fungal relationships mediate
carbon and nutrient dynamics in CMNs under different ecological and environmental conditions.

147
148 **Figure 1. Schema of experimental set up, sampling, and effects of rainout shelters.** Effects
149 of rain exclusion treatment are shown on soil matric potential and soil temperature at each site
150 for each growing season. Growing season is defined as 1 Oct to 30 Jun and each season is
151 labeled by the spring months of the growing year (e.g. 2018 is 1 Oct 2017 to 30 Jun 2018).

Isotopic labelling

On a sunny day between 11AM and 3PM, we applied isotopically enriched carbon (^{13}C)
and nitrogen (^{15}N) as a pulse of carbon dioxide (CO_2) and ammonia (NH_3) to the leaves of target
"donor" species common across experimental sites. At each site, we selected a healthy perennial
forb (*Sidalcea malviflora* in diverse prairie plots [except in one plot where we used *Eriophyllum
lanatum* due to a lack of *S. malviflora*], *Alopecurus pratensis, Schedonorus arundinaceus,* or
Agrostis capillaris in low diversity pasture plots) at the center of each plot to receive the isotopic
labels. We performed the labeling experiment using custom-made field-deployable clear
chambers with internal fans built to allow gas mixing and fast in situ assimilation of isotopic
labels, following established protocols developed in previous isotopically labeling studies (Silva
et al. 2015, e.g., Earles et al. 2016, Sperling et al. 2017). We covered the donor plant with a clear
plastic cylinder and injected gas in sequence at 20 minute intervals. For $^{13}CO_2$, we made three
injections of 2 mL pure CO_2 (enriched at 98 atm % ^{13}C) to double the amount of CO2 in the
chamber each time. For NH3, we made two injections of 10 mL pure NH3 (98 atm% 15N). The
dates of application were based on peak NDVI at each site (Reed et al. 2019). We sampled
leaves from each donor plant immediately after labeling (time point 0) as well as from all plants
approximately 4 days (time point 1), 10 days (time point 2), and 21 days (time point 3) post-
labelling as logistics permitted (Fig 1, Table S2). By applying enriched gases only to the aerial
portions of established perennial donor plants we introduced the isotopic tracer to both the plant
and mycorrhizal fungal tissues without exposing the soil to the tracers. At the end of the
experiment we harvested entire plants and surrounding rhizospheres at time point 3 and kept
them in cool conditions until processing. We separated the roots and rhizospheres, and collected
roots from each plant, selecting approximately ten ~3cm fine root fragments per sample (i.e.
third order or finer, where available) for DNA extraction. All roots and rhizosphere soils were
stored at -80º C until processing.

Baseline and Resource Transfer Calculations

Before isotopic labeling, we collected soil, leaf, and roots from each site. We collected
soils in late spring and early summer 2019 to a 20 cm depth in each plot. From these soil
samples, we removed root fragments that represented the typical roots seen in each plot. We

collected leaves for each species in each plot; however, these leaves were contaminated with ^{15}N

during transport. To replace these, we separately sampled leaves from biomass samplings

performed at the same plots, ensuring that annual and perennial grasses and forbs were

represented at each site. The biomass samples were collected in late spring and early summer

2019. Baseline ^{13}C values were calculated from the original samples because the contamination

only occurred with ^{15}N.

We oven-dried all samples at 65°C to constant mass and encapsulated them for stable

isotope analysis. All stable isotope analysis was done at UC Davis Stable Isotope Facilities using

an Elementar Vario EL Cube or Micro Cube elemental analyzer (Elementar Analysensysteme

GmbH, Hanau, Germany) interfaced to a PDZ Europa 20-20 isotope ratio mass spectrometer

(Sercon Ltd., Cheshire, UK). We calculated the amount of carbon and nitrogen in each plant

compartment (leaves and roots) using standard label recovery equations (Silva et al 2015), using

baseline values measured before application of the labelled gases to capture background

variations in isotopic composition of unenriched leaves, roots, and soil samples

At each relevant point in space and time, we calculated % nitrogen and % carbon derived from

label (%NDFL and %CDFL, respectively) as follows (He et al. 2006, Kramer et al. 2002, Silva

et al. 2015 Nature).

Equation 1

$$\frac{atm\%_{post-label} - atm\%_{baseline}}{atm\%_{labelling\ gas}}$$

Baseline values were calculated from the site- and perenniality-specific values for each plant

material. Site-specific baseline soil and root isotope ratios represent the whole community

because of interconnected rhizospheres where is it is not possible to identify specific species. In

all cases, baseline values fell within the expected range for our region (Fig 2).

We used the baseline ^{13}C values to calculate intrinsic water-use efficiency using the method of

Dawson et al. (*in press*).

Figure 2. Natural abundance of stable isotopes in leaves, roots, and soil before labelling.
Dashed lines indicate mean value.

To better understand post-labelling soil enrichment, we selected a subset of rhizosphere soils that represented six donor plants at each site divided equally between high and low diversity plots, and selected the three most highly [15]N enriched interspecific receivers in each plot. In addition, we sampled three most highly enriched interspecific receivers at each site and diversity combination (if they had not been sampled previously). Because pasture plots were sometimes monodominant grasses, we sampled the top three enriched intraspecific receivers at each site and treatment. In total, this came to 48 post-labelling soil samples in 29 plots.

Fungal analysis

We extracted DNA from roots of 450 plants harvested at time point 3 using Qiagen DNeasy Powersoil HTP kits (Qiagen, Hilden, Germany). We only analyzed DNA from roots, not from the soils collected from each plant's rhizosphere. We characterized each sample's AM fungal composition with a two-step PCR protocol that amplified a ~550bp fragment of the SSU

rRNA gene (the most well-supported region for AM fungal taxonomic resolution (Dumbrell et

al. 2011)). We used WANDA (5'- CAGCCGCGGTAATTCCAGCT- 3') and AML2 (5'-

GAACCCAAACACTTTGGTTTCC-3') primers (Lee et al. 2008, Langmead and Salzberg

2012). We used primers with unique indices so we could multiplex several projects on a single

run. We quantified successful PCR amplicons with the Quant-iT PicoGreen dsDNA Assay Kit

(Invitrogen, Waltham, MA, USA) on a SpectraMax M5E Microplate Reader (Molecular

Devices, San Jose, CA, USA) before purifying with QIAquick PCR Purification kits (Qiagen).

We sequenced the purified pools on the Illumina MiSeq platform (paired-end 300bp, Illumina

Inc., San Diego, CA, USA) at the University of Oregon Genomics and Cell Characterization

Core Facility (Eugene, OR, USA). We used a custom, in-house bioinformatics pipeline to

demultiplex reads and filter out duplicate reads generated by PCR duplication using unique

molecular identifiers (UMIs) inserted during PCR processing.

We assigned amplicon sequence variants (ASVs) using the dada2 pipeline with standard

quality filtering and denoising parameters (Callahan et al. 2016). The dada2 pipeline maintains

strain-level diversity at the scale of individual sequence variants rather than clustering sequences

into OTUs. This fine-scale measure of fungal sequence diversity was particularly important for

our analyses to maintain the greatest chance of detecting a single AM fungal 'individual' in

multiple plant root samples. Taxonomy was assigned to ASVs using the MaarjAM database

(Öpik et al. 2010). We used a Bayesian mixture model to scale ASV counts within and across

samples to avoid artificial taxon abundance biases (Anders and Huber 2010).

Data analysis

We performed all analyses in R ver. 4.0.4 (R Core team 2015). We tested the relationship

between NDFL and plant traits and site conditions with a mixed-effect ANOVA. We constructed

a phyloseq object (McMurdie and Holmes 2013) and used iGraph, metagMisc, DESeq2, and

RCy3 (Nepusz and Csardi 2006, Love et al. 2014, Mikryukov 2017, Gustavsen et al. 2019) to

create networks for each plot. Nodes represented individual plants and edges between nodes

showed where plants shared at least one fungal partner. We weighted edges based on how many

fungal ASVs were shared between plants. We used tidygraph (Petersen 2022) to calculate

degrees of connectivity, or how many plants each individual plant was connected to in each plot

(Fig. 3). We also calculated whether each receiver shared fungal ASVs with the central donor

plant. We visualized individual plot networks in CytoScape. We created a non-metric multi-

dimensional scaling (NMDS) plot using the vegan package demonstrating the mycorrhizal fungal

community similarity between plants (Oksanen et al. 2022).

Figure 3. Example of how networks were constructed for each plot. The grey cylinder

indicates the donor plant for the plot. Numbers beneath the receivers are the distance (in

centimeters) from the donor. Degrees were calculated as how many plants each individual plant

was connected to; for example, the perennial grass at 71 cm has 7 degrees.

Results

Assimilation of isotopic tracers was similar between labeled "donor" plants with no

significant differences on average between sites or experimental treatments within sites,

including rainfall exclusion or species diversity (Fig S1). At all sites, foliar assimilation of ^{15}N

and ^{13}C by donor plants led to enrichment levels ranging from approximately 5-fold to one order

of magnitude higher than baselines, with only a few that were less than 2-fold from baseline (two

donor individuals *of S. malviflora* at the northern site and two donor individuals of *S.*

arundinaceus at the southern site). Foliar enrichment levels decreased consistently at all sites and

treatments over the twenty-one day sampling period, suggesting translocation within plants and

interspecific transfer between species. We found significant spatial and temporal differences in

foliar and root isotope ratios in donors and receivers resulting from interspecific transfer of

carbon and nitrogen (Fig 4; Table 1). Receiver foliar enrichment levels did not correlate with

donor foliar enrichment levels within the same plot (Fig S2).

Figure 4. Decreased receiver enrichment with distance from donor in leaves and roots.

Dashed lines indicate natural abundance means; grey boxes indicate range of natural abundance

variation shown in figure 2. We interpret this as no systematic enrichment of 13C; however,

there is high 15N enrichment in leaves. Y-axes are log10 scale. Note that the y-axis scales are

different between 15N leaves and 15N roots. Inset box shows distribution of 15N atm% for

enriched (greater than natural abundance) annual and perennial leaves and roots.

Table 1. ANOVA results effects on nitrogen derived from label (NDFL)

	χ^2	P-value
Fixed effects		
Perenniality	32.095	**<0.001**
Grass/forb	18.673	**<0.001**
iWUE	0.010	0.921
Degree of connectivity	1.149	0.284
C:N	46.115	**<0.001**
Site	0.360	0.548
Drought treatment	0.534	0.465
Diversity treatment	2.802	0.094
Functional diversity	1.713	0.191
Distance from donor	20.965	**<0.001**
Time from labelling	82.693	**<0.001**
Random effect		
Plot		**<0.001**

Allocation of ^{13}C and ^{15}N tracers to roots and subsequent transfer to "receiver" species

did not follow expected patterns (e.g., despite donors being perennial, annual receivers were

more enriched, indicating that relatedness did not drive the transfer, Fig. 4, Table 1) but varied

significantly between functional groups due to their intrinsic differences in physiological traits

and tissue stoichiometry (Table 1). We selected 18 representative annual/perennial and grass/forb

species of receiver plants, which revealed significant differences between functional groups for

NDFL (Fig 5) but no detectable CDFL relative to baseline (Fig. 4).

Rain exclusion treatment, plot diversity treatment, and site did not affect interspecific

transfer of carbon or nitrogen (Fig S; ANOVA, $P > 0.05$, table 1). We observed significant

differences in nitrogen transfer by functional group, mirroring intrinsic differences in tissue

stoichiometry and iWUE, despite no significant enrichment in soils collected from the

rhizosphere of those same plants (fig S4). We did, however, detect a low level of soil enrichment

in 4 out of 23 donor soil samples (ranging from 0.382 to 0.479 atm% 15N). Annuals had greater

[15]N foliar enrichment compared to perennials (ANOVA, $P < 0.001$, table 1, fig 4 insets). Foliar

enrichment decreased over both time and space (ANOVA $P < 0.001$). On average, annuals had a

lower nitrogen content and higher C:N than perennials (table 1). Annuals had an average of

0.16% ±0.23 SD NDFL at the first sampling; they ranged from 0 to 1.53%. Perennials had an

average of 0.04% ±0.03 NDFL at the first sampling; they ranged from 0.01 to 0.15%. C:N

affected NDFL (table 1), but NDFL did not correlate with either one (Fig. 5).

Figure 5. Nitrogen content by perenniality and intrinsic water-use efficiency. A) Leaf C:N

by perenniality. B) iWUE by perenniality as measured before labelling. C) Percent [15]N derived

from label (DFL) compared to C:N in leaves. D) Percent [15]N DFL compared to iWUE in leaves

at all time points. Grey line indicates 0%.

Our analysis of fungal community composition shows a high degree of connectivity

between plants of different species but no obvious pattern of connectedness that could explain

preferential nutrient transfer by functional groups. Overall, plants in all experimental plots and

sites were highly connected. We infer this from fungal communities within the roots of each

donor and receiver plant where we found that 97.25 % ± 8.01 (SD) of all plants within each

experimental plot shared at least one fungal DNA sequence variant (ASV) with another plant of

the same plot. All plants shared at least one ASV with the plot donor plant, with fungal

communities being apparently agnostic with respect to functional groups (Fig 6). Annual plants

shared fungi with more plants in the same plot (4.74 plants ± 2.62 SD) compared to perennials

(4.04 plants ± 2.49 SD ; t-test, P = 0.019; table S3), but degrees of connectivity did not predict

transfer (table 1).

Figure 6. NMDS of fungal communities. Each point is an individual plant ordinated by its

fungal community.

Discussion

We designed this experiment to be informed by previous hypotheses on how carbon and

nitrogen transfer occurs in highly connected CMNs of the Pacific Northwest. We conducted

repeated spatiotemporal sampling of isotopic enrichment levels at increasing distances from

donor species, days to weeks after labelling, and in well-established communities exposed to

multiple years of experimental treatments, expecting to find evidence of kinship (i.e., greatest

resource transfer in related plants), driven by CMN economics (i.e., ^{15}N transfer rates coinciding

with ^{13}C investment in root and fungal mass). Our data do not support either hypothesis and

instead suggest AM fungi are "agnostic" with respect to plant species partnerships, forming

CMNs where the rates and direction of resource transfer ultimately reflects a sink-source

strength in ecophysiological performance and stoichiometric gradients. We found a ^{15}N signature

in leaves that was especially strong in annual receivers, indicating preferential transfer of

limiting resources to that functional group. However, we did not find ^{13}C enrichment in most

samples, above- or belowground, and both ^{15}N and ^{13}C signatures of receiver plants were

independent of donor or soil enrichment levels.

Our dual-isotope (^{13}C and ^{15}N) labelling approach was imposed upon an existing climate

experiment to investigate in the terms of trade between carbon and nitrogen from assimilation to

translocation among plant compartments and fungal partners. Our results are consistent with

earlier experiments performed in greenhouse experiments under low stress (e.g., temperature

controlled and irrigated twice daily) (Silva et al. 2015). However, post-assimilation translocation

did not support previous data. Plants acquire CO_2 and NH_3 from the atmosphere and release

those same gases into their surroundings, exhibiting a foliar compensation point at which the

evolution of gases is equal to assimilation (Farquhar et al. 1980). This compensation point

depends on the partial pressure of CO_2 or NH_3 in the mesophyll, and therefore on its partial

pressure in the atmosphere, with an increase in leaf uptake expected for both gases as their

concentration rises. Our pulse experiment worked for both gases, causing lasting enrichment

levels in the leaves of doner species and, over time, in the leaves of receiver individuals of other

species, suggesting translocation and transfer (Fig 4 and S1). This result represents an integrated

measure of the total foliar assimilation of NH_3 and CO_2 after a single pulse of isotopically

enriched gases.

Nitrogen enrichment levels remained high in leaves and many roots at the end of the

experiment, allowing us to measure NDFL across the community and infer the main drivers of N

transfer. However, carbon enrichment levels faded before plants were harvested approximately

21 days post-labelling, either due to losses via respiration or redistribution in CMNs where it

became undetectable in all but a few cases. After controlling for variation in assimilation rates,

we found that annual plants received greater ^{15}N enrichment than perennial plants. Plants closest

to the donor were most enriched, and ^{15}N enrichment decreased over time (fig 4). We did not

detect ^{13}C enrichment in roots, shoots, or soils. Annuals had lower total nitrogen content than

perennials, mirroring intrinsic differences in stoichiometry and water-use efficiency (although

with no significant correlation found; Fig 5), indicating that kinship did not drive this transfer as

all donor plants were perennials. Although this experiment was conducted along a latitudinal

climate gradient with rainout shelters, which affected soil water content (fig 1), neither treatment

nor site affected our results.

The major drivers of differences in allocation of ^{13}C and ^{15}N to roots and subsequent

transfer to "receiver" species were the intrinsic difference in physiological traits and tissue

stoichiometry between functional groups (annual/perennial or grass/forb; table 1). This result

points to a simple mechanistic explanation to reconcile long-standing questions regarding

mutualism and hierarchical transfer of resources between plants via mycorrhizal networks, an

explanation that can lead to general predictions of preferential flow of limiting resources in other

ecosystems. For example, previous studies in northern California under environmental conditions

similar to those found in our southernmost experimental site showed rapid (days to weeks)

transfer of ^{15}N applied to the leaves of EM pines to surrounding annual AM plant receivers (He

et al. 2006). Those results were interpreted as evidence for agnostic CMNs because "direct

fungal connections are not necessary for N transfer among plants" and "leaves of the annual

plants had greater ^{15}N derived from source (NDFS) and were more enriched (^{15}N at % excess and

δ^{15}N values) than perennial receivers, irrespective of the mycorrhizal type". Similarly, as

proposed by He et al. (2006), our observation of agnostic ^{15}N transfer from donor to receivers

suggests that annual plants, with their extensive root systems, were a strong sink for N which

could be explained by stoichiometric gradients that affect root exudation and recapture of N-

containing materials from rhizodeposition. Our data corroborate rapid agnostic transfer among

AM plants, with no detectable enrichment in root or soil ^{13}C near roots 21 days post-labelling,

but do not allow us to determine which mechanism is responsible for the ^{15}N transfers.

We found a significant degree of connectivity between plants within each given treatment

and site, with a near complete overlap in their shared CMN communities (Fig 6). This overlap

could explain the lack of support for the kinship hypothesis in our dataset and offers further

support for stoichiometric gradients as the fundamental control of terms of trade in CMNs. The

marketplace hypothesis proposes that host plants provide the mycorrhizal fungus with carbon in

the form of simple sugars. The rate which fungi take up these sugars is controlled by their

affinity to the host. Different C sources from a host trigger changes in fungal monosaccharide

transporter gene expression, causing fungi to uptake N from the soil (Fellbaum et al. 2012).

However, host and fungus-specific gene expression differences would be weak in our study sites

because our DNA results demonstrate that our plants were highly connected by shared arbuscular

mycorrhizal (AM) fungal ASVs. In our case, the ^{15}N we applied to leaves could have

mineralized, been recycled at the soil/fungal interface, and converted into basic organic

compounds such as amino acids (e.g. via glutamine synthetase or argininosuccinate synthase)

which act as charge balance in the catabolic arm of the urea cycle. The fact that we supplied a

carbon source (acetate) independent from the C supply of the host could have reduced N

transport in the AM symbiosis. Apoplastic transport would then explain how some of the labeled

C ended up in plants even though most of it is expected to go to the fungal symbionts.

This mechanism also explains how N and mineral elements (e.g. P) are transferred through

CMNs from plant to plant even though they are not expected to limit fungal growth (Ward et al.

2022).

Prior work suggests that plant-to-plant transfer of nitrogen via fungi is unlikely. More

likely the nitrogen label "leaked" into the soil where fungi harvested it and transferred it to

another, connected, plant (Ingraffia et al. 2021). Regardless, greater fungal connectivity increases

nitrogen transfer (He et al. 2009, Ingraffia et al. 2021). This suggests that the nitrogen transfer

we observed in our system was mediated by fungi. However, because we had few plants that

were disconnected from the network, we cannot say with certainty that connectivity is necessary

for this transfer.

In conclusion, we found that plant-soil stoichiometric gradients and functional traits were

the strongest drivers of resource sharing in grassland CMNs. We interpret this finding as

evidence of biophysical and biophysical sinks, in which nutrients are allocated to plants with the

greatest need for those nutrients, either through a 'passive' mycorrhizal network or direct uptake

from soil, or that nutrients should be allocated through water flow. Expanding on previous

studies, we propose that agnostic fungi facilitate spatiotemporal dynamics of carbon and nitrogen

through CMNs in was that are neither plant- nor fungi-centric. That is, plants and fungi that are

located closer together in space and with stronger demand for resources over time are more

likely to receive larger amounts of those limiting resources.

Acknowledgements

The authors thank the Siskiyou Field Institute, The Nature Conservancy, and Capitol Land Trust

for providing sites for this experiment, Laurel Pfeifer-Meister, Bitty Roy, Bart Johnson, Graham

Bailes, Aaron Nelson, and Matthew Krna for their contributions to experimental design, Emily

Scherer for her assistance with sample analysis, and numerous others for assistance with the

HOPS project. This experiment was funded by National Science Foundation Macrosystems

Biology grant #1340847, Plant Biotic Interactions grant #1758947, and Convergence Accelerator

Pilot grant #1939511.

References

Anders, S., and W. Huber. 2010. Differential expression analysis for sequence count data. Genome Biology 11:1–12.

Averill, C., J. M. Bhatnagar, M. C. Dietze, W. D. Pearse, and S. N. Kivlin. 2019. Global imprint of mycorrhizal fungi on whole-plant nutrient economics. Proceedings of the National Academy of Sciences of the United States of America 116:23163–23168.

Callahan, B. J., P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson, and S. P. Holmes. 2016. DADA2: High-resolution sample inference from Illumina amplicon data. Nature Methods 2016 13:7 13:581–583.

Dumbrell, A. J., P. D. Ashton, N. Aziz, G. Feng, M. Nelson, C. Dytham, A. H. Fitter, and T. Helgason. 2011. Distinct seasonal assemblages of arbuscular mycorrhizal fungi revealed by massively parallel pyrosequencing. New Phytologist 190:794–804.

Earles, J. M., O. Sperling, L. C. R. Silva, A. J. McElrone, C. R. Brodersen, M. P. North, and M. A. Zwieniecki. 2016. Bark water uptake promotes localized hydraulic recovery in coastal redwood crown. Plant, Cell & environment 39:320–328.

Farquhar, G. D., P. M. Firth, R. Wetselaar, and B. Weir. 1980. On the Gaseous Exchange of Ammonia between Leaves and the Environment: Determination of the Ammonia Compensation Point. Plant physiology 66:710–714.

Fellbaum, C. R., E. W. Gachomo, Y. Beesetty, S. Choudhari, G. D. Strahan, P. E. Pfeffer, E. T. Kiers, and H. Bücking. 2012. Carbon availability triggers fungal nitrogen uptake and transport in arbuscular mycorrhizal symbiosis. Proceedings of the National Academy of Sciences of the United States of America 109:2666–2671.

Fellbaum, C. R., J. A. Mensah, A. J. Cloos, G. E. Strahan, P. E. Pfeffer, E. T. Kiers, and H. Bücking. 2014. Fungal nutrient allocation in common mycorrhizal networks is regulated by the carbon source strength of individual host plants. New Phytologist 203:646–656.

Figueiredo, A. F., J. Boy, and G. Guggenberger. 2021. Common Mycorrhizae Network: A Review of the Theories and Mechanisms Behind Underground Interactions. Frontiers in Fungal Biology 0:48.

Gustavsen, J. A., S. Pai, R. Isserlin, B. Demchak, and A. R. Pico. 2019. RCy3: Network biology using Cytoscape from within R. F1000Research 8:1774.

He, X., C. S. Bledsoe, R. J. Zasoski, D. Southworth, and W. R. Horwath. 2006. Rapid nitrogen transfer from ectomycorrhizal pines to adjacent ectomycorrhizal and arbuscular mycorrhizal plants in a California oak woodland. New Phytologist 170:143–151.

He, X., M. Xu, G. Y. Qiu, and J. Zhou. 2009. Use of 15N stable isotope to quantify nitrogen transfer between mycorrhizal plants. Journal of Plant Ecology 2:107–118.

van der Heijden, M. G. A., F. M. Martin, M. A. Selosse, and I. R. Sanders. 2015. Mycorrhizal ecology and evolution: the past, the present, and the future. New Phytologist 205:1406–1423.

Ingraffia, R., D. Giambalvo, A. S. Frenda, E. Roma, P. Ruisi, and G. Amato. 2021. Mycorrhizae differentially influence the transfer of nitrogen among associated plants and their competitive relationships. Applied Soil Ecology 168:104127.

Kenrick, P., and C. Strullu-Derrien. 2014. The Origin and Early Evolution of Roots 1. Plant Physiology:570–580.

Kiers, E. T., M. Duhamel, Y. Beesetty, J. A. Mensah, O. Franken, E. Verbruggen, C. R. Fellbaum, G. A. Kowalchuk, M. M. Hart, A. Bago, T. M. Palmer, S. A. West, P.

Vandenkoornhuyse, J. Jansa, and H. Bücking. 2011. Reciprocal rewards stabilize
cooperation in the mycorrhizal symbiosis. Science 333:880–882.
Langmead, B., and S. L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2. Nature
Methods 2012 9:4 9:357–359.
Lee, J., S. Lee, and J. P. W. Young. 2008. Improved PCR primers for the detection and
identification of arbuscular mycorrhizal fungi. FEMS Microbiology Ecology 65:339–349.
Love, M. I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biology 15:1–21.
McMurdie, P. J., and S. Holmes. 2013. phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. PLOS ONE 8:e61217.
Mikryukov, V. 2017, May 4. metagMisc.
Nepusz, T., and G. Csardi. 2006. The igraph software package for complex network research.
InterJournal.
Oksanen, J., G. Simpson, F. Blanchet, R. Kindt, P. Legendre, P. Minchin, R. O'Hara, P.
Solymos, M. Stevens, E. Szoecs, H. Wagner, M. Barbour, M. Bedward, B. Bolker, D.
Borcard, G. Carvalho, M. Chirico, M. de Caceres, S. Durand, H. Evangelista, R. FitzJohn,
M. Friendly, B. Furneaux, G. Hannigan, M. Hill, L. Lahti, D. McGlinn, M. Ouellette, E.
Ribeiro Cunha, T. Smith, A. Stier, C. ter Braak, and J. Weedon. 2022. vegan: Community
Ecology Package.
Öpik, M., A. Vanatoa, E. Vanatoa, M. Moora, J. Davison, J. M. Kalwij, Ü. Reier, and M. Zobel.
2010. The online database MaarjAM reveals global and ecosystemic distribution patterns in
arbuscular mycorrhizal fungi (Glomeromycota). New Phytologist 188:223–241.
Petersen, T. 2022. tidygraph: A Tidy API for Graph Manipulation.
Peterson, M. L., G. Bailes, L. B. Hendricks, L. Pfeifer☐Meister, P. B. Reed, S. D. Bridgham, B.
R. Johnson, R. Shriver, E. Waddle, H. Wroton, D. F. Doak, B. A. Roy, and W. F. Morris.
2020. Latitudinal gradients in population growth do not reflect demographic responses to
climate. Ecological Applications.
Pickles, B. J., R. Wilhelm, A. K. Asay, A. S. Hahn, S. W. Simard, and W. W. Mohn. 2017.
Transfer of ^{13}C between paired Douglas-fir seedlings reveals plant kinship effects and
uptake of exudates by ectomycorrhizas. New Phytologist 214:400–411.
R Core team. 2015. R Core Team.
Reed, P. B., S. D. Bridgham, L. E. Pfeifer-Meister, M. L. DeMarche, B. R. Johnson, B. A. Roy,
G. T. Bailes, A. A. Nelson, W. F. Morris, and D. F. Doak. 2021a. Climate warming
threatens the persistence of a community of disturbance-adapted native annual plants.
Ecology 102:e03464.
Reed, P. B., M. L. Peterson, L. E. Pfeifer☐Meister, W. F. Morris, D. F. Doak, B. A. Roy, B. R.
Johnson, G. T. Bailes, A. A. Nelson, and S. D. Bridgham. 2021b. Climate manipulations
differentially affect plant population dynamics within versus beyond northern range limits.
Journal of Ecology 109:664–675.
Reed, P. B., L. E. Pfeifer☐Meister, B. A. Roy, B. R. Johnson, G. T. Bailes, A. A. Nelson, M. C.
Boulay, S. T. Hamman, S. D. Bridgham, L. E. Pfeifer-Meister, B. A. Roy, B. R. Johnson, G.
T. Bailes, A. A. Nelson, M. C. Boulay, S. T. Hamman, and S. D. Bridgham. 2019. Prairie
plant phenology driven more by temperature than moisture in climate manipulations across
a latitudinal gradient in the Pacific Northwest, USA. Ecology and Evolution 9:3637–3650.

Reed, P. B., L. E. Pfeifer☐Meister, B. A. Roy, B. R. Johnson, G. T. Bailes, A. A. Nelson, and S.
D. Bridgham. 2021c. Introduced annuals mediate climate☐driven community change in
Mediterranean prairies of the Pacific Northwest, USA. Diversity and Distributions 00:1–12.
Saxton, K. E., and W. J. Rawls. 2006. Soil Water Characteristic Estimates by Texture and
Organic Matter for Hydrologic Solutions. Soil Science Society of America Journal
70:1569–1578.
Selosse, M. A., F. Richard, X. He, and S. W. Simard. 2006, November 1. Mycorrhizal networks:
des liaisons dangereuses? Elsevier Current Trends.
Silva, L. C. R., and H. Lambers. 2021. Soil-plant-atmosphere interactions: structure, function,
and predictive scaling for climate change mitigation. Plant and Soil:1–23.
Silva, L. C. R., A. Salamanca-Jimenez, T. A. Doane, and W. R. Horwath. 2015. Carbon dioxide
level and form of soil nitrogen regulate assimilation of atmospheric ammonia in young
trees. Scientific reports 5:13141.
Simard, S. W., D. A. Perry, M. D. Jones, D. D. Myrold, D. M. Durall, and R. Molina. 1997. Net
transfer of carbon between ectomycorrhizal tree species in the field. Nature 388:579–582.
Smith, S. E., and D. J. (David J.) Read. 2008. Mycorrhizal symbiosis. Academic Press.
Sperling, O., L. C. R. Silva, A. Tixier, G. Théroux-Rancourt, and M. A. Zwieniecki. 2017.
Temperature gradients assist carbohydrate allocation within trees. Scientific Reports
7:3265.
Tedersoo, L., M. Bahram, and M. Zobel. 2020. How mycorrhizal associations drive plant
population and community biology. Science 367.
Walder, F., and M. G. A. van der Heijden. 2015. Regulation of resource exchange in the
arbuscular mycorrhizal symbiosis. Nature Plants 1.
Ward, E. B., M. C. Duguid, S. E. Kuebbing, J. C. Lendemer, and M. A. Bradford. 2022. The
functional role of ericoid mycorrhizal plants and fungi on carbon and nitrogen dynamics in
forests. New Phytologist.
Weremijewicz, J., L. da S. L. O. Sternberg, and D. P. Janos. 2016. Common mycorrhizal
networks amplify competition by preferential mineral nutrient allocation to large host
plants. New Phytologist 212:461–471.
Werner, C., and M. Dubbert. 2016. Resolving rapid dynamics of soil-plant-atmosphere
interactions. New Phytologist 210:767–769.
Wilson, H., B. R. Johnson, B. Bohannan, L. E. Pfeifer-Meister, R. Mueller, and S. D. Bridgham.
2016. Experimental warming decreases arbuscular mycorrhizal fungal colonization in
prairie plants along a Mediterranean climate gradient. PeerJ 2016:1–22.
557