Title: Agnostic fungi: plant functional types and tissue stoichiometry explain nutrient transfer in

common arbuscular mycorrhizal networks of temperate grasslands

Authors: Hilary Rose Dawson[1,2*], Katherine L. Shek[1,2,3], Toby M. Maxwell[1,4], Paul B. Reed[1,2,5],

Barbara Bomfim[1,6], Scott D. Bridgham[1,2], Brendan J. M. Bohannan[1,2], Lucas C.R. Silva[1,2,7*]

Affiliations:

1. Institute of Ecology and Evolution, University of Oregon, Eugene, OR 97403 USA

2. Department of Biology, University of Oregon, Eugene, OR 97403 USA

3. Department of Natural Resources and the Environment, University of New Hampshire,

Durham NH 03824

4. Department of Biological Sciences, Boise State University, Boise, Idaho, 83725 USA

5. Institute for Applied Ecology, Corvallis, OR 97333 USA

6. Climate and Ecosystem Sciences Division, Lawrence Berkeley National Laboratory, Berkeley,

CA 94720 USA

7. Environmental Studies Program, University of Oregon, Eugene, OR 97403 USA

Corresponding authors: H.R.D. hilaryd@uoregon.edu L.C.R.S. lsilva7@uoregon.edu

ORCIDs:

H.R.D.: 0000-0003-4613-762X

K.L.S.: 0000-0001-5629-5782

T.M.M.: 0000-0001-5171-0705

P.B.R.: 0000-0001-7143-7515

B.B.: 0000-0001-9510-2496

S.D.B.: 0000-0003-0614-2678

B.J.M.B.: 0000-0003-2907-1016

L.C.R.S.: 0000-0002-4838-327X

Abstract:

Plants and mycorrhizal fungi form mutualistic relationships that affect how resources

flow between organisms and within ecosystems. Common mycorrhizal networks (CMNs) could

facilitate preferential transfer of carbon and limiting nutrients but this remains difficult to predict.

Do CMNs favor fungal resource acquisition at the expense of plant resource demands (a fungi-

centric view), or are they passive channels through which plants regulate resource fluxes (a

plant-centric view)? We used stable isotope tracers ($^{13}CO_2$ and $^{15}NH_3$), plant traits, and

mycorrhizal DNA to quantify above- and belowground carbon and nitrogen transfer between 18

plant species along a 520-km latitudinal gradient in the Pacific Northwest, USA. Plant functional

type and tissue stoichiometry were the most important predictors of interspecific resource

transfer. Of "donor" plants, 98% were ^{13}C-enriched, but we detected transfer in only 2% of

"receiver" plants. However, all donors were ^{15}N-enriched and we detected transfer in 81% of

receivers. Nitrogen was preferentially transferred to annuals (0.26 ± 0.50 mg N per g leaf mass)

compared to perennials (0.13 ± 0.30 mg N per g leaf mass). This corresponded with tissue

stoichiometry differences. Our findings point to a simple mechanistic answer for long-standing

questions regarding transfer of resources between plants via mycorrhizal networks.

Keywords: arbuscular mycorrhizal fungi, carbon, common mycorrhizal networks, ecological

stoichiometry, grasslands, nitrogen, plant functional type

Introduction:

Plant-mycorrhizal associations are thought to have emerged as rudimentary root systems over 400 million years ago, facilitating the expansion of terrestrial life that followed[1]. The transformative power of early fungal symbioses is still evident today in all major plant lineages, from bryophytes to angiosperms [2]. Over 85% of all contemporary flowering plant species form symbioses with fungi, with arbuscular mycorrhizal (AM) associations being the most common [3]. Today, the relationships between plants and AM fungi dominates both managed and unmanaged landscapes and are estimated to be responsible for up to 80% of global primary productivity [2]. Fungi are gregarious and form symbioses with more than one individual plant [4]. By extension, it has been widely hypothesized that the multi-plant-fungal relationships form "common mycorrhizal networks" (CMNs) which facilitate carbon and nutrient transfer between organisms, beyond the immediate plant-fungus mutualism formed by individuals. This relationship is often simplified to the mutual exchange of carbon-based photosynthates for limiting soil nutrients that are more readily bioavailable to fungi [5], but is known to exist along a continuum from parasitic to mutualistic [6–9]. A synthesis of empirical research at macro- and micro-levels of ecological organization reveals a gap in understanding of structural and functional properties that could predict how carbon and nutrients are transferred between plants through fungi in CMNs [10].

The literature holds myriad and often complimentary, but sometimes contradictory, hypotheses that could explain the CMN mutualism as a key structural and functional component of ecosystems. For example, the "economics" hypothesis [11] proposes that plants and fungi engage in "trades" of nutrients mined by fungi in exchange for plant photosynthates [12–14]. In the economics hypothesis, the terms of trade between plant and fungi are mediated by supply and

demand for limiting resources, which could create a dynamic market emerging from interactions

between environmental, biochemical, and biophysical variables. The "Wood Wide Web"

hypothesis emerged from the analysis of isotopically labeled carbon transferred between plants,

presumably through fungal mycorrhizae. Simard et al. [15] concluded that plants that allocate

carbon to sustain common fungal symbionts also benefit from shared nutrients, while plants

associating with mycorrhizal fungi outside that network cannot. Complementing the economic

analogy, the "kinship" hypothesis proposes that plants of the same species preferentially receive

more resources in CMNs [16,17].

The past two decades have seen extensive but inconclusive research on these hypotheses

and how they relate to empirical measurements of CMN structure and function. On the one hand,

economic analogies suggest that the reciprocally regulated exchange of resources between plants

and fungi in CMNs should favor the most beneficial cooperative partnerships [11,12]. On the

other hand, reciprocal transfer is only found in a subset of symbionts under specific conditions,

while increased competition in CMNs is a more common observation [18,19]. At the core of this

controversy is whether CMNs actively support fungal resource acquisition at the expense of

plant resource demands (i.e., a fungi-centric view) or function as passive channels through which

plants regulate resource fluxes (i.e., a plant-centric view). If plant-centric, we expect to find that

the structure and functioning of CMNs give rise to consistent spatiotemporal patterns of resource

allocation akin to those predicted by the kinship hypothesis. If fungi-centric, we expect to find

that spatiotemporal patterns of resource allocation reflect the composition and functioning of the

fungal community regardless of the connecting plant nodes in CMNs. Data exist to support both

opposing views [see 10,20]; therefore, we posit that CMNs are neither plant- nor fungi-centric.

Instead, perhaps rather than gregarious, AM fungi are "agnostic" with respect to plant species

composition and relatedness, and yet affected by major plant functional traits that are known to

influence resource use and allocation. There is no clear evidence to suggest that AM

fungi exhibit specificity for plant species as do host-specific pathogens [21]. Even in the systems

dominated by ectomycorrhizal (EM) associations where most previous research into CMN

structure and function was developed, fungi tend to be generalists, although plant taxa-specific

EM fungi occur at relatively high proportion, especially in early successional temperate plant

communities [22,23]. Building on previous research, our hypothesis represents a first-principles

simplification in which gradients in resource limitation, rather than species-specific plant-fungi

interactions, might allow us to better understand and predict the rates and direction of resource

transfer in CMNs.

To test the "agnostic fungi" hypothesis, we asked if interactions among biophysical and

biogeochemical processes could explain resource transfer in CMNs with more accuracy than

previous plant- or fungi-centric analogies. We studied how soil resource-use efficiency and plant

traits that are known to regulate physiological performance [24] and that are expected to affect

the transfer of carbon and nitrogen [25] in paired experiments including restored prairie and

pasture experimental sites under ambient conditions and rain exclusion shelters (designed to

affect soil water and nutrient mass flow). To test this, we applied ^{15}N-enriched NH3 to plant

leaves, which are known to uptake ammonia [26,27], and sampled surrounding plants to examine

the transfer of nitrogen using the tracer. Although gases are not the primary source of nitrogen

for most plants, applying gaseous nitrogen allowed us to limit the amount leaked into the soil

compared to applying nitrogen directly to the soil [28]. We replicated our paired experimental

setting at three different locations, with study sites distributed across a 520 km latitudinal

gradient. Due to the climatically driven differences in communities across sites, not all species

were present at all sites (Table S1); however, all functional groups were present at all sites and

most species were present at more than one site. In this experiment, we also sequenced strain-

level variation in root fungal DNA, plant functional types, and leaf stoichiometric traits to test if

relatedness (same or different species as the donor) explained difference in resource transfer.

This allowed us to generalize our inferences about the structure and function of arbuscular

CMNs in temperate grasslands. The diversity of plant growth forms included in our study also

affect CMN composition and function. For example, Davison et al. [29] found plant growth form

altered AM fungal community and functional diversity. Plant total biomass can also increase

nitrogen received through a CMN, an effect that may be compounded by whether plants are

woody or herbaceous [30]. This could be explained by larger or denser plants providing more

carbon. Both plants and fungi have economic spectra characterized by fast or slow traits and

nutrient strategies which together form an interacting continuum potentially driven by N

availability [31]. It is unclear to what extent plant or fungi characteristics drive these plant-fungal

interactions. Therefore, we designed an experiment to quantify how plant-fungal interactions

influence the structure and functioning of CMNs across broad environmental gradients and

resource constraints.

Materials and Methods:

We conducted our experiment at three sites situated on a 520 km latitudinal transect that

spans three Mediterranean climates: cool, moist (northern site; Tenino, WA) to warm, moist

(central site; Eugene, OR) to warm, dry (southern site; Selma, OR). Each circular plot was 3 m in

diameter. Half of our plots were restored prairie systems (n = 10 per site) while the other half of

the plots had introduced pasture grasses established prior to restoration (n = 10 per site).

Restored prairie plots were mowed, raked, received herbicide, and seeded in 2014-2015,

followed by seeding in fall 2015, 2016, and 2017 [32]. We erected rainout shelters that excluded

40% of the rainfall on half the plots at each site (n = 10 rain exclusion, 10 control per site; Fig.

1).

All plants in our system have the potential to form associations with AM fungi (Table S1)

[33–35]. Previous work demonstrated that the rainout shelters had minimal effects on

aboveground community structure or function [24], possibly due to the shoulder season effect of

the Mediterranean rain seasonality. This network of experimental sites was established in 2010

and has been extensively studied since then [32,36–39] including work on mycorrhizal fungi

[40,41]. Treatments had marginal effects on the soil water potential (especially during the early

growing season). Pasture plots were dominated by one to a few species of introduced perennial

grasses which we targeted as donors in these plots. We used different species of perennial grass

at each site. For restored prairie plots, we targeted the native perennial forb *Sidalcea malviflora*

ssp. *virgata* common to ambient and drought treatments at all sites. Despite those differences, we

did not find significant changes in the average plant community composition or productivity

under rain exclusion, which also did not affect morphological and functional traits (e.g., specific

leaf area, iWUE, and C:N ratios) of the functional groups we selected for this experiment

[24,36].

To test our new hypothesis, we built upon those preliminary data with a dual isotope label

approach as follows. We labeled perennial plants central to each plot (hereafter, 'donors') and

monitored leaf ^{15}N and ^{13}C for the surrounding plants ('receivers') for up to 21 days. In addition,

we sampled roots and soils at 21 days post-labelling. We characterized AM fungal DNA isolated

from the root samples, as well as stable isotopes in all leaf and root samples. Our experimental

design was nested in a multi-year experiment where data loggers were used to continuously

measure temperature and moisture in all the manipulated plots.

Figure 1. Schema of experimental set up, sampling, and effects of rainout shelters. Figure shows

the three location of the sites (orange points on map) in relation to the climatic gradient (cool to

warm, wet to dry). Each site has twenty plots which are a full factorial representation of the

restoration and rain exclusion treatments. Each plot had a central perennial plant that was labeled

with isotopically enriched gas (shown with a grey cylinder here) and we sampled plants of each

functional type at three distances from the central labelled plant. Inset shows the temporal

sampling scheme of leaves, roots and soil.

Isotopic labelling

At each site, we selected a healthy perennial forb (*Sidalcea malviflora* ssp. *virgata* in

restored prairie plots [except in one plot where we used *Eriophyllum lanatum* due to a lack of *S.*

malviflora ssp. virgata]), or a perennial grass (*Alopecurus pratensis, Schedonorus arundinaceus,*

or *Agrostis capillaris*) in pasture plots at the center of each plot to receive the isotopic labels. On

sunny days between 11AM and 3PM at peak productivity for each site, we applied isotopically

enriched carbon (^{13}C) and nitrogen (^{15}N) as a pulse of carbon dioxide (CO_2) and ammonia (NH_3)

to the leaves of target "donor" species common across experimental sites. We performed the

labeling experiment using custom-made field-deployable clear chambers with internal fans built

to allow gas mixing and fast in situ assimilation of isotopic labels, following established

protocols developed in previous isotopically labeling studies [e.g., 42–44]. Before performing

the experiment, we tested our approach in the field to optimize gas exposure and labeling

amounts, which included checking for leaks and contamination outside of the chamber. We

covered the donor plant with a clear plastic cylindrical chamber and injected gas in sequence at

20-minute intervals. For $^{13}CO_2$, we made three injections of 2 mL pure CO_2 (enriched at 98 atm

% ^{13}C) to double the amount of CO_2 in the chamber each time. For NH_3, we made two injections

of 10 mL pure NH_3 (98 atm% ^{15}N) at 20 minute intervals. The dates of application were based on

peak productivity estimated from Normalized Different Vegetation Index (NDVI) at each site

[see 32 for details]. We sampled leaves from each donor plant immediately after labeling (time

point 0) as well as from all plants approximately 4 days (time point 1), 10 days (time point 2),

and 21 days (time point 3) post-labelling as logistics permitted (Fig. 1, Table S2). We also

collected leaves at time points 1, 2, and 3 from up to twelve plants in each plot representing three

replicates of factorial grass/forb structural groups and annual/perennial life history strategies

(Table S1). The number of plants and represented groups depended on which plants were

growing in each plot. For example, pasture plots were limited to only grasses and the northern

pasture plots had only perennial grasses.

By applying enriched gases only to the aboveground portions of established perennial

donor plants, we introduced the isotopic tracer to both the plant and mycorrhizal fungal tissues

with minimal exposure of the soil to the tracers. At the end of the experiment, we harvested

entire plants and the soil surrounding the roots at time point 3 and kept them in cool conditions

until processing. We separated the roots and rhizospheres, and collected roots from each plant,

selecting approximately ten ~3 cm fine root fragments per sample (i.e., third order or finer,

where available) for DNA extraction and identification. All roots and rhizosphere soils were

stored at -80º C until processing.

Baseline and Resource Transfer Calculations

Before isotopic labeling, we collected soil, leaves, and roots from each site. We collected

soils in late spring and early summer 2019 to 20 cm depth in each plot. From these soil samples,

we removed root fragments that represented the typical roots seen in each plot. We collected

leaves for each species in each plot; however, these leaves were contaminated with ^{15}N during

transport. To replace contaminated samples, we separately sampled leaves from biomass

samplings collected in late spring and early summer 2019, ensuring that annual and perennial

grasses and forbs were represented at each site.

We oven-dried all samples at 65º C to constant mass and encapsulated them for stable

isotope analysis. All stable isotope analysis was done at UC Davis Stable Isotope Facilities using

an Elementar Vario EL Cube or Micro Cube elemental analyzer (Elementar Analysensysteme

GmbH, Hanau, Germany) interfaced to a PDZ Europa 20-20 isotope ratio mass spectrometer

(Sercon Ltd., Cheshire, UK). We calculated the amount of carbon and nitrogen in each plant

compartment (leaves and roots) using standard label recovery equations [44], using baseline

values measured before application of the labelled gases to capture background variations in

isotopic composition of unenriched leaves, roots, and soil samples.

We designated all samples with greater than two standard deviations above baseline

samples as "enriched" in a particular isotope. Baseline values were calculated on a site by rain

exclusion treatment basis by plant functional type basis (Table S3). We calculated these values

for each plant functional type because our sampling scheme was designed around these

functional types. Site-specific baseline soil and root isotope ratios represent the whole

community because of interconnected rhizospheres where is it was not possible to identify

specific species. In all cases, baseline values fell within the expected range for our region (Fig.

2). For each enriched sample, we calculated isotope excess as

Equation 1

$$atm\%_{excess} = atm\%_{post\ label} - atm\%_{baseline}$$

For enriched donor plants, we calculated % derived from label immediately following label

application as

Equation 2

$$\%DFL_{donor} = \frac{atm\%_{excess}}{atm\%_{labelling\ gas} - atm\%_{baseline}} * 100$$

For enriched receiver plants, we calculated % derived from label (%NDFL and %CDFL,

respectively) for each relevant point in space and time as [40]

Equation 3

$$\%DFL_{receiver} = \frac{atm\%_{receiver\ excess}}{atm\%_{donor\ excess}} * 100$$

When we calculated %DFL in roots, we used donor leaves as the source (atm% donor excess).

We then calculated the amount derived from label on a per mass basis as

Equation 4

$$DFL\ (\text{mg N / g leaf}) = 1000 * 1g * \left(\frac{N\ or\ C\%}{100}\right) * \left(\frac{DFL\%}{100}\right)$$

We calculated intrinsic water-use efficiency following Farquhar and Richards [45] using

the baseline ^{13}C values from the original samples because the contamination only occurred with

^{15}N. Because of sampling discrepancies, we did not have intrinsic water-use efficiency data for

32 plants.

Figure 2. Natural abundance of stable isotopes in leaves, roots, and soil before labelling.

Dashed lines indicate mean value. Exact values used to calculate derivation from label (Eq. 1 –

Eq. 4) are available in Table S3.

To better understand post-labelling soil enrichment, we selected a subset of rhizosphere

soils that represented six donor plants at each site divided equally between restored prairie and

pasture plots, and selected the three most highly [15]N-enriched interspecific receivers in each plot

across the sites. In addition, we sampled three most highly enriched interspecific receivers at

each site and restored prairie-introduced pasture combination (if they had not been sampled

previously). Because pasture plots were sometimes monodominant grasses, we sampled the top

three enriched intraspecific receivers at each site and treatment. In total, this came to 48 post-

labelling soil samples in 29 plots.

Fungal DNA analysis

We extracted DNA from roots of 450 plants harvested at time point 3 (21 days post-label)

using Qiagen DNeasy Powersoil HTP kits (Qiagen, Hilden, Germany). We only analyzed DNA

from roots, not from the soils collected from each plant's rhizosphere. We characterized each

sample's AM fungal composition with a two-step PCR protocol that amplified a ~550bp

fragment of the SSU rRNA gene (the most well-supported region for AM fungal taxonomic

resolution [46]. We used WANDA (5'- CAGCCGCGGTAATTCCAGCT- 3') and AML2 (5'-

GAACCCAAACACTTTGGTTTCC-3') primers [47,48]. We used primers with unique indices

so we could multiplex several projects on a single run. We quantified successful PCR amplicons

with the Quant-iT PicoGreen dsDNA Assay Kit (Invitrogen, Waltham, MA, USA) on a

SpectraMax M5E Microplate Reader (Molecular Devices, San Jose, CA, USA) before purifying

with QIAquick PCR Purification kits (Qiagen). We sequenced the purified pools on the Illumina

MiSeq platform (paired-end 300bp, Illumina Inc., San Diego, CA, USA) at the University of

Oregon Genomics and Cell Characterization Core Facility (Eugene, OR, USA). Reads were

deduplicated with UMI-tools using unique molecular identifiers (UMIs) inserted during PCR

[49].

We assigned amplicon sequence variants (ASVs) using the dada2 pipeline (version

1.18.0) with standard quality filtering and denoising parameters [50]. The dada2 pipeline

maintains strain-level diversity at the scale of individual sequence variants rather than clustering

sequences into OTUs. This fine-scale measure of fungal sequence diversity was particularly

important for our analyses to maintain the greatest chance of detecting a single AM fungal

'individual' in multiple plant root samples. Taxonomy was assigned to ASVs using the

MaarjAM database (2019 release) [51]. We used a Bayesian mixture model in the DESeq2

package [52] to scale ASV counts within and across samples to avoid artificial taxon abundance

biases [53].

Data analysis

We performed all analyses in R ver. 4.0.4 [54]. We removed one outlier plant with a ^{15}N

atm% more than twice as high as the next highest measurement. We also removed five

mislabeled samples. To meet statistical assumptions, we only included data from enriched plants

with successful root fungal DNA extraction in our analyses and figures. We limited receivers to

those for which we also had sufficient root fungal DNA. In total, from 1440 leaves measured for

isotopic content and with successfully recovered fungal DNA, we analyzed data from 391 unique

plants: 53 donors and 338 receivers.

We tested the relationship between receiver leaf %DFL and plant traits (grass/forb,

annual/perennial, iWUE, C:N, degrees of connectivity, interaction term between grass/forb and

annual/perennial) and site conditions (position on latitude gradient, pasture/restored, rain

exclusion treatment, distance from donor, time from labelling) with a mixed-effect ANOVA

(plot as random effect; Table 1). We constructed a phyloseq object using the ASV table with

normalized counts[55], and used iGraph, metagMisc, and RCy3 [56–58] to create networks for

each plot. In each network, nodes represented individual plants and edges between nodes

represent plants sharing at least one fungal DNA sequence variant. The weighted edges are based

on how many fungal ASVs were shared among plants. We calculated degrees of connectivity

with tidygraph [59] to examine how many plants each individual plant was 'connected' to (by

means of shared fungal ASVs) in each plot (Fig. 3). We also calculated whether each receiver

plant shared fungal ASVs with the central donor plant in each plot. We visualized individual plot

networks in CytoScape. We visualized shifts in AM fungal community composition using non-

metric multi-dimensional scaling (NMDS) in the vegan package, demonstrating the AM fungal

community similarity across plants [60].

Figure 3. Example of how networks were constructed for each plot. The grey cylinder

indicates the donor plant for the plot. Numbers beneath the receivers are the distance (in

centimeters) from the donor. Degrees were calculated as how many plants each individual plant

was connected to by shared fungal ASVs; for example, the perennial grass at 71 cm has 7

degrees.

Results:

All donors were [15]N-enriched in their leaves at time of labelling. Two donors were not

[13]C-enriched in their leaves at time of labelling. We recovered DNA data from the roots of

88.3% of donors. We sampled 1,444 leaves from 434 receiver plants at three time points. Of

these leaves, 81.0% were [15]N-enriched and 2.4% were [13]C-enriched. We recovered DNA from

the roots of 77.9% of the receivers. At time point 3 (roughly 21 days post-labelling), we

collected roots from 46 of the initial 60 donors and 306 of the initial 434 receivers. Of the roots

of the collected donors, 97.8% were still 15N-enriched and 23.9% were still [13]C-enriched at time

point 3. Of the roots of the collected receivers, 33.0% were still ^{15}N-enriched and 10.5% were

still ^{13}C-enriched at time point 3.

Assimilation of isotopic tracers was similar between labeled "donor" plants with no

significant differences on average between sites or experimental treatments within sites,

including rainfall exclusion or restored status (Fig. S1). At all sites, foliar assimilation of ^{15}N and

^{13}C by donor plants led to enrichment levels ranging from approximately 5-10 fold higher than

baselines. Foliar enrichment levels decreased consistently at all sites and treatments over the 21

day sampling period. Annual forbs had the greatest enrichment level and perennial forbs the

lowest enrichment level (Fig. 4). We found significant spatial and temporal differences in foliar

and root isotope ratios in donors and receivers resulting from interspecific transfer of carbon and

nitrogen (Fig. S2; Table 1). Receiver foliar enrichment levels did not correlate with donor foliar

enrichment levels within the same plot (Fig. S3).

Figure 4. Annuals had greater leaf nitrogen derived from label (NDFL) scaled by mass than perennials. Points represent enriched individual receiver plants with associated DNA data at one time point of sampling. All three post-enrichment time points are shown. Y-axis is log10 scale. See Fig. S2 for values over distance and Fig. S4 for NDFL in roots as well as CDFL in roots and leaves.

Table 1. ANOVA results effects on leaf nitrogen derived from label (%NDFL). Only

enriched receiver leaves with associated DNA data were included in the analysis (n = 1094).

Leaf %CDFL results available in Table S4.

	DF	χ^2	P-value
Fixed effects			
Annual/perennial	**1**	**10.33**	**<0.001**
Grass/forb	**1**	**11.45**	**<0.001**
Same species as donor	1	0.54	0.463
Degree of connectivity	1	0.17	0.68
iWUE	1	0.06	0.81
C:N	**1**	**36.78**	**<0.001**
Site	2	0.69	0.71
Drought treatment	1	0.51	0.48
Restoration treatment	1	0.91	0.34
Distance from donor	**1**	**9.72**	**0.002**
Time from labelling	**1**	**84.05**	**<0.001**
Annual/perennial: Grass/forb interaction	**1**	**7.87**	**0.01**
Random effect			
Plot			**<0.001**

Allocation of ^{13}C and ^{15}N tracers to roots and subsequent transfer to "receiver" species

did not follow expected patterns (e.g., despite donors being perennial, annual receivers were

more enriched, indicating that relatedness did not drive the transfer, Fig. 4, Table 1). Allocation

and transfer varied significantly between functional groups due to their intrinsic differences in

tissue stoichiometry (Table 1). We selected 18 common annual/perennial and grass/forb species

of receiver plants, which revealed significant differences between functional types for NDFL

(Fig. 4) but little detectable CDFL relative to baseline (Fig. S2). Rain exclusion treatment,

restoration treatment, and site did not affect interspecific transfer of nitrogen (Fig. S5; ANOVA,

$P > 0.05$, Table 1). We observed very low carbon enrichment, but of the 2.4% of leaves that were

enriched in carbon, plant functional type and site affected carbon transfer (Table S4). We did,

however, observe significant differences in nitrogen transfer by plant functional type (Table 1),

mirroring intrinsic differences in tissue stoichiometry and iWUE (Fig. 5), despite no significant

enrichment in soils collected from the rhizosphere of those same plants (Fig. S6). We also found

that C:N affected NDFL, although not in a simple linear manner and with no apparent correlation

between NDFL and iWUE (Fig. 5). We did detect a low level of soil enrichment in 1 out of 30

receiver soil samples (0.377 atm% ^{15}N) and 5 out of 18 donor soil samples (ranging from 0.376

to 0.479 atm% ^{15}N; Fig. S6).

Annuals had greater ^{15}N foliar enrichment compared to perennials (ANOVA, $P < 0.001$,

Tables 1 and 2). Foliar enrichment decreased over both time and space (ANOVA $P < 0.001$; Fig.

S1, Fig. S2). On average, annuals had a lower leaf nitrogen content and higher C:N than

perennials (Table 2, Fig. 5). Forbs had higher NDFL than grasses (ANOVA, $P < 0.001$, Table 1)

as well as a lower C:N. There was a significant interaction between annual/perennial and

grass/forb form (ANOVA, $P = 0.003$, Table 1). There was no correlation between the relatedness

of donors and receivers in 15N-enrichment (table 1).

Table 2. Leaf nitrogen derived from label (NDFL) and leaf tissue nitrogen (N%) four days

after labeling.

			%NDFL				NDFL (mg N/g leaf)				N (%)			
		n	*Mean*	*SD*	*Min*	*Max*	*Mean*	*SD*	*Min*	*Max*	*Mean*	*SD*	*Min*	*Max*
Annual	**Forb**	54	6.528	9.769	0.101	56.005	0.749	0.819	0.041	3.858	0.749	0.819	0.041	3.858
	Grass	92	4.847	9.032	0.098	63.884	0.34	0.463	0.011	2.79	0.340	0.463	0.011	2.790
Perennial	**Forb**	67	0.952	1.069	0.090	5.582	0.214	0.228	0.021	1.156	0.214	0.228	0.021	1.156
	Grass	148	1.632	3.066	0.088	22.434	0.252	0.484	0.013	2.948	0.252	0.484	0.013	2.948

Figure 5. Stoichiometric and functional traits compared to nitrogen derived from transfer.

A) Leaf C:N by annual/perennial. B) Percent [15]N derived from label (DFL) compared to C:N in

leaves. C) iWUE by annual/perennial as measured before labelling. D) Percent [15]NDFL

compared to iWUE in leaves at all time points. Letters above boxplots indicate significant

differences.

Our analysis of fungal community composition shows a high degree of connectivity

between plants of different species but no obvious pattern of connectedness that could explain

preferential nutrient transfer by plant functional groups. We found that 97.25% ± 8.01 (SD) of all

plant roots within each experimental plot shared at least one fungal DNA sequence variant

(ASV) with another plant of the same plot. Fungal community composition was similar across

plant functional groups (Fig. 6, PERMANOVA pseudo-F statistic = 2.269, R^2 = 0.005, p =

0.001). Annual plants shared fungi with more plants in the same plot (4.74 plants ± 2.62 SD)

compared to perennials (4.04 plants ± 2.49 SD ; t-test, P = 0.019; Table S5), but degrees of

connectivity did not predict nitrogen transfer (Table 1). Seventy-three percent of plants were

colonized by four or fewer fungi and shared fungi with five or fewer other plants in the plot,

making it difficult to determine if strength of connectivity altered nitrogen transfer (Fig. S7).

Figure 6. NMDS of fungal communities. Each point represents an individual plant ordinated by

the Bray Curtis dissimilarities for AM fungal composition. PERMANOVA pseudo-F statistic =

2.269, $R^2 = 0.005$, p = 0.001.

Discussion:

In our extensively replicated spatiotemporal isotopic tracer experiment, the assimilation

and allocation of limiting resources in CMNs was neither plant- nor fungi-centric. Our data show

that AM fungi are "agnostic" with respect to plant species composition and relatedness, and that

the transfer of nitrogen (an important nutrient that can limit plant growth in our system) is

regulated by major functional traits that are known to influence resource use and allocation in

plant communities. The rates and direction of resource transfer in CMNs, inferred from pulse

labeling and recovery of ^{15}N in leaves and rhizospheres, can be predicted from leaf C:N and

distance from donor species (Table 1). Across all sites and treatments, we observed a stronger

sink for ^{15}N in annual plants (Table 2), indicating preferential transfer of limiting resources to

that functional group of plants. We applied the 15N enriched label to perennial species in all

plots, so this greater enrichment in annual plants precludes a preference for receiver plants of the

same species as would be expected under the kinship hypothesis (Table 1). Although plants

shared ~97% fungal ASVs in their roots (Fig. 6), connectivity did not predict ^{15}N transfer (Table

1), and thus we found no evidence for a fungi-centric hypothesis based on which all connected

plants would have been expected to be equally enriched in ^{15}N. Our data suggest that rates and

direction of resource transfer in CMNs reflect plant nutrient requirements and spatial proximity.

We designed this experiment to be informed by previous hypotheses on how carbon and

nitrogen transfer occurs in highly connected CMNs, where a dilution of the applied ^{13}C label is

to be expected given the vast amounts of carbon in the soil and plant biomass. We conducted

repeated spatiotemporal sampling of isotopic-enrichment levels at increasing distances from

donor species, days to weeks after labelling, and in well-established communities exposed to

multiple years of experimental treatments, expecting to find evidence of kinship (i.e., greatest

resource transfer in plants of the same species), driven by CMN economics (i.e., ^{15}N transfer

rates coinciding with ^{13}C investment in root and fungal mass). Our donor plants were perennials

while annuals received, on average, an order of magnitude higher enrichment. Thus, we

discarded kinship as a possible driver of transfer in this system. We also did not find evidence of

a relationship between ^{15}N in leaves and ^{13}C in roots because we did detect ^{15}N-enrichment in

leaves but no ^{13}C-enrichment in roots. We did, however, find significant ^{13}C-enrichment in the

donor plants. Therefore, our data do not support either hypothesis, and instead suggest AM fungi

are "agnostic" with respect to plant species partnerships, forming CMNs where the rates and

direction of resource transfer ultimately reflects a sink-source strength effect, consistent with

previous observations of stoichiometric source-sink manipulations of carbon and nitrogen within

plants [61–63], but in our case observed at the community scale.

Nitrogen enrichment levels remained high in leaves and many roots at the end of the

experiment, allowing us to measure NDFL across the community and infer the main drivers of N

transfer. However, carbon enrichment levels faded before plants were harvested approximately

21 days post-labelling, either due to losses via respiration or redistribution in CMNs where it

became undetectable in all but a few cases. After controlling for variation in assimilation rates by

calculating NDFL, we found that annual plants received greater ^{15}N-enrichment than perennial

plants. Plants closest to the donor were most enriched, and ^{15}N-enrichment decreased over time

(Fig. S2, Fig. S3). Plant level of connectivity through an assumed CMN did not predict ^{15}N-

enrichment. Although the rainout shelters had limited effect, there were major differences across

the latitudinal gradient represented by the sites in temperature and soil moisture availability

[24,32]; however, neither treatment nor site affected our results.

The major drivers of differences in allocation of ^{13}C and ^{15}N to roots and subsequent

transfer to "receiver" species were the intrinsic difference in plant functional types and correlated

traits, including measured leaf C:N. Previous studies in northern California under environmental

conditions similar to those found in our southernmost experimental site showed rapid (days to

weeks) transfer of ^{15}N applied to the leaves of ectomycorrhizal pines to surrounding annual AM

plant receivers [64]. Those results were interpreted as evidence for agnostic CMNs because

"direct fungal connections are not necessary for N transfer among plants" and "leaves of the

annual plants had greater ^{15}N derived from source and were more enriched (^{15}N at % excess and

δ^{15}N values) than perennial receivers, irrespective of the mycorrhizal type." Similarly, as

proposed by He et al. [64], our observation of agnostic ^{15}N transfer from donor to receivers

suggests that annual plants were a strong sink for N which could be explained by stoichiometric

gradients that affect root exudation and recapture of N-containing materials from rhizodeposition

[65,66]. Although our study included two species with symbiotic N fixation ability could have

altered some of the baseline data but even if a plot was fully dominated by legumes that

difference would represent a minor effect relative to the pulse label application. Our data

corroborate rapid agnostic transfer among AM plants, with no detectable enrichment in root or

soil ^{13}C near roots 21 days post-labelling, but do not allow us to determine general mechanisms

that are responsible for the ^{15}N transfers.

We inferred a high connectivity between plants within each given treatment and site

given the highly similar fungal composition in the root systems of both perennial and annual

plants (Fig. 6). Given the constraints of ASV-identified data, we did this analysis on a strain-

level scale and it is possible that separate spores of the same ASV separately infected plants

within the same plot. However, we are reasonably confident in our use of fungal ASVs as a

proxy for connectivity given the strong overlap in our community and because individuals of one

ASV can anastomose in the soil [67]. This overlap could explain the lack of support for the

kinship hypothesis in our dataset and offers further support for stoichiometric gradients in

general, and C:N gradients in particular, as a principal control of terms of trade in CMNs

[11]. Because we observed such high rates of shared ASVs (~97%) and we observed high levels

of N-enrichment in receivers at our first post-label sampling point four days after application, we

could not test whether receivers connected to donors or whether plants were connected to the

network affected ^{15}N transfer. We found unexpectedly low soil isotopic-enrichment (fig. S6)

which suggested that the labels did not remain in the fungal network. This low level was likely

driven by the fact that we did not sample soils until 21 days after labeling and hyphal turnover

for grass-associated AM fungi can be less than one month [68]. Because N is a major limiting

nutrient in this system, according to the current paradigm of CMNs, it would be quickly taken up

and recycled or transferred rather than accumulating in the soil.

We found that plant-soil stoichiometric gradients and functional traits were the strongest

drivers of resource sharing in grassland CMNs. We interpret this finding as evidence of

biochemical and biophysical sinks, in which nutrients are allocated to plants with the greatest

need for those nutrients, either through a 'passive' mycorrhizal network or direct uptake from

soil, or that nutrients should be allocated through water flow. Expanding on previous studies, we

propose that agnostic AM fungi facilitate spatiotemporal dynamics of carbon and nitrogen

transfer through CMNs in ways that are neither plant- nor fungi-centric. That is, plants and fungi

that are located closer together in space and with stronger demand for resources over time are

more likely to receive larger amounts of those limiting resources.

Acknowledgements:

The authors thank the Siskiyou Field Institute, The Nature Conservancy, and Capitol

Land Trust for providing sites for this experiment, Laurel Pfeifer-Meister, Bitty Roy, Bart

Johnson, Graham Bailes, Aaron Nelson, and Matthew Krna for their contributions to

experimental design, Emily Scherer for her assistance with sample analysis, Bitty Roy, Leonel

Sternberg, Timothy A. Doane, and five anonymous reviewers for comments on the manuscript,

and numerous others for assistance with the HOPS project. This experiment was funded by

National Science Foundation Macrosystems Biology grant #1340847, Plant Biotic Interactions

grant #1758947, and Convergence Accelerator Pilot grant #1939511.

Competing interests:

The authors have no competing interests to declare.

Author contributions:

L.C.R.S, S.D.B., and B.J.M.B. designed the research; H.R.D., K.L.S., T.M.M., B.B., and

P.B.R. performed the research; H.R.D. led the data analysis and manuscript preparation with

K.L.S. assisting; all authors contributed to revising the manuscript and approved the publication.

Data availability:

Data used in these analyses will be available online at the Data Dryad repository

(https://doi.org/10.5061/dryad.7pvmcvdxt) after publication.

References:

521   **References:**

1. Kenrick P, Strullu-Derrien C. 2014 The Origin and Early Evolution of Roots 1. *Plant Physiol* , 570–580. (doi:10.1104/pp.114.244517)

2. van der Heijden MGA, Martin FM, Selosse MA, Sanders IR. 2015 Mycorrhizal ecology and evolution: the past, the present, and the future. *New Phytologist* **205**, 1406–1423. (doi:10.1111/NPH.13288)

3. Brundrett MC. 2009 Mycorrhizal associations and other means of nutrition of vascular plants: understanding the global diversity of host plants by resolving conflicting information and developing reliable means of diagnosis. *Plant Soil* **320**, 37–77. (doi:10.1007/s11104-008-9877-9)

4. Selosse MA, Richard F, He X, Simard SW. 2006 Mycorrhizal networks: des liaisons dangereuses? *Trends Ecol Evol*. **21**, 621–628. (doi:10.1016/j.tree.2006.07.003)

5. Smith SE, Read DJ (David J). 2008 *Mycorrhizal symbiosis*. Academic Press.

6. Mariotte P, Meugnier C, Johnson D, Thébault A, Spiegelberger T, Buttler A. 2013 Arbuscular mycorrhizal fungi reduce the differences in competitiveness between dominant and subordinate plant species. *Mycorrhiza* **23**, 267–277. (doi:10.1007/S00572-012-0465-8/FIGURES/4)

7. Klironomos JN. 2003 Variation in plant response to native and exotic arbuscular mycorrhizal fungi. *Ecology* **84**, 2292–2301. (doi:10.1890/02-0413)

8. Johnson NC, Graham JH, Smith FA. 1997 Functioning of mycorrhizal associations along the mutualism–parasitism continuum. *New Phytologist* **135**, 575–585. (doi:10.1046/J.1469-8137.1997.00729.X)

9.   Luo X, Liu Y, Li S, He X. 2023 Interplant carbon and nitrogen transfers mediated by

common arbuscular mycorrhizal networks: beneficial pathways for system functionality.

Front Plant Sci **14**. (doi:10.3389/fpls.2023.1169310)

10.   Silva LCR, Lambers H. 2021 Soil-plant-atmosphere interactions: structure, function, and

predictive scaling for climate change mitigation. *Plant Soil* , 1–23. (doi:10.1007/s11104-

020-04427-1)

11.   Kiers ET *et al.* 2011 Reciprocal rewards stabilize cooperation in the mycorrhizal

symbiosis. *Science (1979)* **333**, 880–882. (doi:10.1126/science.1208473)

12.   Fellbaum CR, Mensah JA, Cloos AJ, Strahan GE, Pfeffer PE, Kiers ET, Bücking H. 2014

Fungal nutrient allocation in common mycorrhizal networks is regulated by the carbon

source strength of individual host plants. *New Phytologist* **203**, 646–656.

(doi:10.1111/nph.12827)

13.   Werner C, Dubbert M. 2016 Resolving rapid dynamics of soil-plant-atmosphere

interactions. *New Phytologist* **210**, 767–769. (doi:10.1111/nph.13936)

14.   Averill C, Bhatnagar JM, Dietze MC, Pearse WD, Kivlin SN. 2019 Global imprint of

mycorrhizal fungi on whole-plant nutrient economics. *Proc Natl Acad Sci U S A* **116**,

23163–23168.

(doi:10.1073/PNAS.1906655116/SUPPL_FILE/PNAS.1906655116.SD01.CSV)

15.   Simard SW, Perry DA, Jones MD, Myrold DD, Durall DM, Molina R. 1997 Net transfer

of carbon between ectomycorrhizal tree species in the field. *Nature* **388**, 579–582.

(doi:10.1038/41557)

16.   Tedersoo L, Bahram M, Zobel M. 2020 How mycorrhizal associations drive plant

population and community biology. *Science (1979)* **367**.

(doi:10.1126/SCIENCE.ABA1223/ASSET/64588F4D-CCDD-4041-891C-

3C19CD9AEDF9/ASSETS/GRAPHIC/367_ABA1223_F3.JPEG)

17.   Pickles BJ, Wilhelm R, Asay AK, Hahn AS, Simard SW, Mohn WW. 2017 Transfer of

^{13}C between paired Douglas-fir seedlings reveals plant kinship effects and uptake of

exudates by ectomycorrhizas. *New Phytologist* **214**, 400–411. (doi:10.1111/nph.14325)

18.   Walder F, van der Heijden MGA. 2015 Regulation of resource exchange in the arbuscular

mycorrhizal symbiosis. *Nat Plants* **1**. (doi:10.1038/nplants.2015.159)

19.   Weremijewicz J, Sternberg L da SLO, Janos DP. 2016 Common mycorrhizal networks

amplify competition by preferential mineral nutrient allocation to large host plants. *New*

Phytologist **212**, 461–471. (doi:10.1111/nph.14041)

20.   Figueiredo AF, Boy J, Guggenberger G. 2021 Common Mycorrhizae Network: A Review

of the Theories and Mechanisms Behind Underground Interactions. *Frontiers in Fungal*

Biology **0**, 48. (doi:10.3389/FFUNB.2021.735299)

21.   Lee E-H, Eo J-K, Ka K-H, Eom A-H. 2013 Mycobiology diversity of arbuscular

mycorrhizal fungi and their roles in ecosystems. **41**, 121–125.

(doi:10.5941/MYCO.2013.41.3.121)

22.   Spencer MW, Roy BA, Thornton TE, Silva LCR, McGuire KL. 2023 Logging has legacy

effects on the structure of soil fungal communities several decades after cessation in

Western Cascade forest stands. *Frontiers in Forests and Global Change* **6**, 41.

(doi:10.3389/FFGC.2023.791766)

23.   McMahen K, Guichon SHA, Anglin CD, Lavkulich LM, Grayston SJ, Simard SW. 2022

Soil microbial legacies influence plant survival and growth in mine reclamation. *Ecol Evol*

12, e9473. (doi:10.1002/ECE3.9473)

24.   Dawson HR, Maxwell TM, Reed PB, Bridgham SD, Silva LCR. 2022 Leaf traits predict

water-use efficiency in U.S. Pacific Northwest grasslands under rain exclusion treatment.

J Geophys Res Biogeosci **127**, e2022JG007060. (doi:10.1029/2022JG007060)

25.   Peaucelle M *et al.* 2019 Covariations between plant functional traits emerge from

constraining parameterization of a terrestrial biosphere model. *Global Ecology and*

Biogeography **28**, 1351–1365. (doi:10.1111/GEB.12937)

26.   Farquhar GD, Firth PM, Wetselaar R, Weir B. 1980 On the Gaseous Exchange of

Ammonia between Leaves and the Environment: Determination of the Ammonia

Compensation Point. *Plant Physiol* **66**, 710–714. (doi:10.1104/pp.66.4.710)

27.   Sutton MA, Erisman JW, Dentener F, Möller D. 2008 Ammonia in the environment: From

ancient times to the present. *Environmental Pollution* **156**, 583–604.

(doi:10.1016/j.envpol.2008.03.013)

28.   Silva LCR, Salamanca-Jimenez A, Doane TA, Horwath WR. 2015 Carbon dioxide level

and form of soil nitrogen regulate assimilation of atmospheric ammonia in young trees.

Sci Rep **5**, 13141. (doi:10.1038/srep13141)

29.   Davison J *et al.* 2020 Plant functional groups associate with distinct arbuscular

mycorrhizal fungal communities. *New Phytologist* **226**, 1117–1128.

(doi:10.1111/nph.16423)

30.   He Y, Cornelissen JHC, Wang P, Dong M, Ou J. 2019 Nitrogen transfer from one plant to

another depends on plant biomass production between conspecific and heterospecific

species via a common arbuscular mycorrhizal network. *Environmental Science and*

Pollution Research **26**, 8828–8837. (doi:10.1007/S11356-019-04385-X/FIGURES/4)

31.   Ward EB, Duguid MC, Kuebbing SE, Lendemer JC, Bradford MA. 2022 The functional

role of ericoid mycorrhizal plants and fungi on carbon and nitrogen dynamics in forests.

New Phytologist (doi:10.1111/NPH.18307)

32.   Reed PB *et al.* 2019 Prairie plant phenology driven more by temperature than moisture in

climate manipulations across a latitudinal gradient in the Pacific Northwest, USA. *Ecol*

Evol **9**, 3637–3650. (doi:10.1002/ece3.4995)

33.   Soudzilovskaia NA *et al.* 2020 FungalRoot: global online database of plant mycorrhizal

associations. *New Phytologist* **227**, 955–966. (doi:10.1111/nph.16569)

34.   Chaudhary VB *et al.* 2016 Data Descriptor: MycoDB, a global database of plant response

to mycorrhizal fungi. *Nature* , 3:160028. (doi:10.1038/sdata.2016.28)

35.   Dickie IA, Martínez-García LB, Koele N, Grelet GA, Tylianakis JM, Peltzer DA,

Richardson SJ. 2013 Mycorrhizas and mycorrhizal fungal communities throughout

ecosystem development. *Plant Soil* **367**, 11–39. (doi:10.1007/s11104-013-1609-0)

36.   Reed PB, Pfeifer-Meister LE, Roy BA, Johnson BR, Bailes GT, Nelson AA, Bridgham

SD. 2021 Introduced annuals mediate climate-driven community change in Mediterranean

prairies of the Pacific Northwest, USA. *Divers Distrib* **00**, 1–12. (doi:10.1111/ddi.13426)

37.   Reed PB *et al.* 2021 Climate manipulations differentially affect plant population dynamics

within versus beyond northern range limits. *Journal of Ecology* **109**, 664–675.

(doi:10.1111/1365-2745.13494)

38.   Reed PB *et al.* 2021 Climate warming threatens the persistence of a community of

disturbance-adapted native annual plants. *Ecology* **102**, e03464. (doi:10.1002/ecy.3464)

39.   Peterson ML *et al.* 2020 Latitudinal gradients in population growth do not reflect

demographic responses to climate. *Ecological Applications* (doi:10.1002/eap.2242)

40.   Wilson H, Johnson BR, Bohannan B, Pfeifer-Meister LE, Mueller R, Bridgham SD. 2016

635   Experimental warming decreases arbuscular mycorrhizal fungal colonization in prairie

636   plants along a Mediterranean climate gradient. *PeerJ* **2016**, 1–22.

637   (doi:10.7717/peerj.2083)

638   41.   Vandegrift R, Roy BA, Pfeifer-Meister L, Johnson BR, Bridgham SD. 2015 The

639   herbaceous landlord: integrating the effects of symbiont consortia within a single host.

640   *PeerJ* **3**, e1379. (doi:10.7717/peerj.1379)

641   42.   Earles JM, Sperling O, Silva LCR, McElrone AJ, Brodersen CR, North MP, Zwieniecki

642   MA. 2016 Bark water uptake promotes localized hydraulic recovery in coastal redwood

643   crown. *Plant Cell Environ* **39**, 320–328. (doi:10.1111/pce.12612)

644   43.   Sperling O, Silva LCR, Tixier A, Théroux-Rancourt G, Zwieniecki MA. 2017

645   Temperature gradients assist carbohydrate allocation within trees. *Sci Rep* **7**, 3265.

646   (doi:10.1038/s41598-017-03608-w)

647   44.   Silva LCR, Salamanca-Jimenez A, Doane TA, Horwath WR. 2015 Carbon dioxide level

648   and form of soil nitrogen regulate assimilation of atmospheric ammonia in young trees.

649   *Sci Rep* **5**, 13141. (doi:10.1038/srep13141)

650   45.   Farquhar GD, Richards RA. 1984 Isotopic Composition of Plant Carbon Correlates With

651   Water-Use Efficiency of Wheat Genotypes. *Functional Plant Biology* **11**, 539–552.

652   46.   Dumbrell AJ, Ashton PD, Aziz N, Feng G, Nelson M, Dytham C, Fitter AH, Helgason T.

653   2011 Distinct seasonal assemblages of arbuscular mycorrhizal fungi revealed by

654   massively parallel pyrosequencing. *New Phytologist* **190**, 794–804. (doi:10.1111/J.1469-

655   8137.2010.03636.X)

656   47.   Langmead B, Salzberg SL. 2012 Fast gapped-read alignment with Bowtie 2. *Nature*

657   *Methods 2012 9:4* **9**, 357–359. (doi:10.1038/nmeth.1923)

658   48.   Lee J, Lee S, Young JPW. 2008 Improved PCR primers for the detection and

659   identification of arbuscular mycorrhizal fungi. *FEMS Microbiol Ecol* **65**, 339–349.

660   (doi:10.1111/J.1574-6941.2008.00531.X)

661   49.   Smith T, Heger A, Sudbery I. 2017 UMI-tools: modeling sequencing errors in Unique

662   Molecular Identifiers to improve quantification accuracy. *Genome Res* **27**, 491–499.

663   (doi:10.1101/gr.209601.116)

664   50.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016

665   DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*

666   *2016 13:7* **13**, 581–583. (doi:10.1038/nmeth.3869)

667   51.   Öpik M, Vanatoa A, Vanatoa E, Moora M, Davison J, Kalwij JM, Reier Ü, Zobel M. 2010

668   The online database MaarjAM reveals global and ecosystemic distribution patterns in

669   arbuscular mycorrhizal fungi (Glomeromycota). *New Phytologist* **188**, 223–241.

670   (doi:10.1111/J.1469-8137.2010.03334.X)

671   52.   Love MI, Huber W, Anders S. 2014 Moderated estimation of fold change and dispersion

672   for RNA-seq data with DESeq2. *Genome Biol* **15**, 1–21. (doi:10.1186/S13059-014-0550-

673   8/FIGURES/9)

674   53.   Anders S, Huber W. 2010 Differential expression analysis for sequence count data.

675   *Genome Biol* **11**, 1–12. (doi:10.1186/GB-2010-11-10-R106/COMMENTS)

676   54.   R Core team. 2015 R Core Team. *R: A Language and Environment for Statistical*

677   *Computing. R Foundation for Statistical Computing , Vienna, Austria. ISBN 3-900051-07-*

678   *0, URL http://www.R-project.org/.* **55**, 275–286.

55.   McMurdie PJ, Holmes S. 2013 phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data. *PLoS One* **8**, e61217.

(doi:10.1371/JOURNAL.PONE.0061217)

56.   Nepusz T, Csardi G. 2006 The igraph software package for complex network research.

57.   Mikryukov V. 2017 metagMisc. (doi:10.5281/ZENODO.571403)

58.   Gustavsen JA, Pai S, Isserlin R, Demchak B, Pico AR. 2019 RCy3: Network biology

using Cytoscape from within R. *F1000Res* **8**, 1774.

(doi:10.12688/F1000RESEARCH.20887.3)

59.   Petersen T. 2022 tidygraph: A Tidy API for Graph Manipulation.

60.   Oksanen J *et al.* 2022 vegan: Community Ecology Package.

61.   Cai Z, Xie T, Xu J. 2021 Source–sink manipulations differentially affect carbon and

nitrogen dynamics, fruit metabolites and yield of Sacha Inchi plants. *BMC Plant Biol* **21**,

1–14. (doi:10.1186/S12870-021-02931-9/FIGURES/8)

62.   Ruiz-Vera UM, De Souza AP, Long SP, Ort DR. 2017 The role of sink strength and

nitrogen availability in the down-regulation of photosynthetic capacity in field-grown

Nicotiana tabacum L. at elevated CO2 concentration. *Front Plant Sci* **8**, 998.

(doi:10.3389/FPLS.2017.00998/BIBTEX)

63.   Tegeder M, Masclaux-Daubresse C. 2018 Source and sink mechanisms of nitrogen

transport and use. *New Phytologist* **217**, 35–53. (doi:10.1111/NPH.14876)

64.   He X, Bledsoe CS, Zasoski RJ, Southworth D, Horwath WR. 2006 Rapid nitrogen transfer

from ectomycorrhizal pines to adjacent ectomycorrhizal and arbuscular mycorrhizal plants

in a California oak woodland. *New Phytologist* **170**, 143–151. (doi:10.1111/j.1469-

8137.2006.01648.x)

65.   Mayer J, Buegger F, Jensen ES, Schloter M, Heß J. 2003 Estimating N rhizodeposition of

grain legumes using a 15N in situ stem labelling method. *Soil Biol Biochem* **35**, 21–28.

(doi:10.1016/S0038-0717(02)00212-2)

66.   Høgh-Jensen H, Schjoerring JK. 2001 Rhizodeposition of nitrogen by red clover, white

clover and ryegrass leys. *Soil Biol Biochem* **33**, 439–448. (doi:10.1016/S0038-

0717(00)00183-8)

67.   Mikkelsen BL, Rosendahl S, Jakobsen I. 2008 Underground resource allocation between

individual networks of mycorrhizal fungi. *New Phytologist* **180**, 890–898.

(doi:10.1111/J.1469-8137.2008.02623.X)

68.   See CR, Keller AB, Hobbie SE, Kennedy PG, Weber PK, Pett-Ridge J. 2022 Hyphae

move matter and microbes to mineral microsites: Integrating the hyphosphere into

conceptual models of soil organic matter stabilization. *Glob Chang Biol* **28**, 2527–2540.

(doi:10.1111/gcb.16073)

Supporting information

Table S1. List of species at each site with number of leaf samples per species.

Table S2. Collection dates for each time point at each site.

Table S3. Baseline values used to calculate amount derived from label (DFL).

Table S4. ANOVA results effects on leaf carbon derived from label (%CDFL).

Table S5. List of fungal ASVs by annuals and perennials.

Figure S1. Enrichment decreases over time in labelled plants.

Figure S2. Decreased receiver enrichment with distance from donor in leaves and roots.

Figure S3. Receiver enrichment did not correlate with donor enrichment.

Figure S4: Enrichment derived from label for A) CDFL in leaves, B) NDFL in roots, and C)

CDFL in roots.

Figure S5. No trend in ^{15}N-enrichment across site and rain exclusion treatment.

Figure S6. Limited soil enrichment 21 days after labelling.

Figure S7. Histogram of connectivity metrics showing left skew.