TITLE

When traits vary across species but performance doesn't: One solution to the paradox

AUTHORS

Kristen Nolting and Kent Holsinger

ABSTRACT

Variation in plant traits related to size and architecture, water relations, and resource uptake

reflect important dimensions of plant function that may explain how different species are

adapted to their local environments. Within a local community, variation in many of these traits

may promote coexistence of competing species through niche partitioning. If differences in traits

are associated with differences in performance, and there is substantial trait variation within

local communities, we might expect competitive asymmetries that would lead to competitive

exclusion. Alternatively, if combinations of trait differences converge on similar performance trait

differences associated with niche partitioning may lead to phenotypically diverse species that

exhibit similar performance. Here we develop a model showing that when the major axes of trait

covariation do not align with the axis of performance variation, species that differ in structural

traits, i.e., species having 'alternative designs,' may achieve similar performance. We apply this

model to data from commonly co-occurring *Protea* species in the Cape Floristic Region, South

Africa and demonstrate that despite large differences in traits that are related to performance,

trait covariation leads to minimal performance differences among species.

KEYWORDS

Intraspecific trait variation, trait covariation, alternative designs, plant functional traits

INTRODUCTION

Hutchinson noted in his seminal paper – *"An Homage to Santa Rosalia"* – that there seem to be

more species on the planet than niches to hold them (Hutchinson 1959). Since then, many

investigators have sought to identify niche axes along which species differ that might allow

species to coexist, often using traits to understand the functional link between an organism's

phenotype and its fitness in a given environment (Schoener 1968; Brown and Lieberman 1973;

Cody 1986; Gillespie 2004; Treurnicht et al. 2020). In contrast, a few authors have instead

proposed that trait differences among species do not lead to competitive differences among

them (Hubbell 2001).

Global surveys of trait variation in land plants reveal large differences (Reich et al. 1997;

Wright et al. 2004; Díaz et al. 2016), and studies of trait differences among species within

diverse clades such as *Helianthus* demonstrate that the magnitude of interspecific trait variation

can rival that seen in global surveys (Mason and Donovan 2015). Even within local plant

communities and assemblages the trait variation observed among co-occurring species can

span orders of magnitude (Wright et al. 2004; Baraloto et al. 2010; Aiello-Lammens et al. 2017)

and can be greater than expected given regional species pools (Swenson et al. 2012). As these

examples suggest, it is common to find substantial trait variation at almost any scale, even

among individuals within a species growing at a single site (Messier et al. 2010; Umaña and

Swenson 2019; Nolting et al. 2021; Taseski et al. 2021). In contrast, substantial performance

differences are less commonly documented, especially among co-occurring species (Becker et

al. 1999; Worthy et al. 2020; Nolting et al. 2021). Why do we fail to see large differences in

performance when we see large differences in individual structural traits that are related to

performance?

Twenty years ago Marks and Lechowicz (2006) hinted at a solution to this paradox. They

showed that selection on multivariate trait combinations within a single environment can

produce many different functional forms ("alternative designs") with roughly equivalent

performance. This result suggests that while individual *trait* differences among species may

reflect differential resource use (and niche partitioning), **performance** differences among

species within a community are likely to be minimal, a conclusion consistent with the

expectation from coexistence theory (Chesson 2000; Clark et al. 2007), whereby coexisting

species should be roughly equivalent in performance if stabilizing differences are not

pronounced. Marks and Lechowicz (2006) argue that while trait variation in natural communities

is often assumed to reflect species' optimization to microenvironments, it could be that "…aside

from major environmental heterogeneity, alternative designs are a major, if not the main, cause

of functional diversity in nature" (Marks and Lechowicz 2006, p. 62). In spite of their provocative

suggestion, few studies have investigated this outcome in empirical systems and the way in

which trait covariation can lead to alternative designs has not been described.

Here we use an extensive collection of structural and physiological performance traits

measured on individual plants to demonstrate that multivariate combinations of structural traits

differ markedly among species but lead to similar physiological performance among co-

occurring *Protea s*pecies sampled from natural communities in the Cape Floristic Region, South

Africa (data originally published in Nolting et al. 2021). We develop a simple model to explain

these results, and we conclude that the presence of persistent 'alternative designs' (i.e., large

variation among species in individual structural traits that converge on similar performance,

(Marks and Lechowicz 2006; Marks 2007)) within natural communities could be the result of

selection that minimizes performance differences among species. This outcome may be

prevalent in natural plant and animal communities, and has important implications for

understanding the ecological and evolutionary mechanisms contributing to persistence of

species-diverse assemblages world-wide.

MATERIALS AND METHODS

EMPIRICAL DATA COLLECTION

We use data from (Nolting et al. 2021) to explore the relationship between structural trait

variation and physiological performance. Specifically, we demonstrate that (1) traits predict

physiological performance relatively well, and (2) species differ markedly in their traits, but do

not substantially differ in their physiological performance.

Study Site and Species

Nolting et al. (2021) investigated multivariate trait-performance relationships for five *Protea*

species sampled across three sites in the Cape Floristic Region, South Africa. The Cape

Floristic Region (CFR) is the smallest of the world's six floral kingdoms (area ~90,000 km^2), but

it hosts over 9,000 plant species (Linder 2003; Peter Linder 2005) making it the most diverse of

the five Mediterranean regions (Cowling et al. 1996). Like other Mediterranean biomes, the CFR

is characterized by warm, dry summers and cool, wet winters (Cowling et al. 1996), but there is

also a west-east seasonality gradient with winter rainfall in the west and relatively constant, low

rainfall in the east. *Protea* species primarily occur in fynbos habitat, which is characterized as a

shrubland with nutrient-poor soils (Allsopp, N. Colville, J.F. Verboom, G.A. 2014) and natural,

recurring fires that sweep through sites on average every 10-12 years (Van Wilgen et al. 2010).

The genus *Protea* L. (Proteaceae) represents a morphologically diverse (~112 species in

Africa, with ~60% endemic to the Cape Floristic Region; (Schnitzler et al. 2011)), iconic, and

locally abundant plant genus that typifies many fynbos communities. *Protea* exhibit a diversity of

form and function with respect to growth form (e.g., low growing, prostrate shrubs to upright

shrubby trees), response to fire (e.g., reseeding species that die with fire but release all of their

seeds during the fire event from persistent infructescences versus resprouting species that

regrow after fire in addition to releasing seeds), and vegetative traits associated with carbon and

water use strategies (e.g., leaf size and shape variation). Previous studies on *Protea* have

demonstrated that variation in structural traits (e.g., leaf mass per area and wood density) and

physiological traits (e.g., photosynthetic rates and stem hydraulic conductance) are associated

with fitness differences (Carlson et al. 2011; Prunier et al. 2012; Nolting et al. 2021). Further,

variation in these traits is associated with variation in the abiotic environment, such as

precipitation and temperature gradients across the CFR (Carlson et al. 2011; Mitchell et al.

2015).

Trait Collection

The samples included in Nolting et al. (2021) occur at three sites and include 11-20 individuals

of each species per site (a total of 8 populations and 132 individuals). On each individual plant

sampled, Nolting et al. (2021) recorded measures of plant size (e.g., height and canopy area)

and reproductive effort measured as the total number of inflorescences and infructescences

retained on a plant. The *Protea* species included in the study either regenerate from seed or

resprout after fire, meaning that adults (a) are all roughly the same age within a stand as they

either recruit as seedlings together post-fire or regrow from persistent stems, and (b) all retain

their infructescences until these 'seedheads' split and release seeds during a fire. These

characteristics make it possible to record reproductive effort for individuals (e.g., inflorescences

plus infructescences) which has been shown to covary with seed production (Carlson and

Holsinger 2013), representing a proxy for fitness (Nolting et al. 2021).

On each individual, Nolting et al. (2021) also measured area-based, light-saturated

photosynthetic rates (A_{area}), stomatal conductance (g_s), and instantaneous water-use efficiency

(WUE_{Instan}) in the field. They collected branches to measure sapwood-specific hydraulic

conductivity (K_s) following Espino and Schenk (2011), and quantified the Huber value of each

stem (the sapwood cross-sectional area of a stem relative to the total one-sided leaf area per

area, (Mencuccini and Bonosi 2001)) on which hydraulic measures were recorded, to calculate

leaf-specific conductivity (LSC, sapwood-specific conductivity divided by total leaf area of the

stem, Tyree and Zimmerman 2002) and leaf-specific photosynthetic rates (LSP, light-saturated

photosynthetic rate per area divided by Huber value). From the stems and leaves on which

physiological traits were measured, Nolting et al. (2021) measured a suite of structural traits

(e.g., traits related to organ morphology and allocation): leaf area (LA, cm^2), leaf mass per area

(LMA, g cm^{-2}), lamina density (LD, g cm^{-3}), leaf length-width ratio (LWR, unitless), stomatal pore

length (SL, mm), stomatal pore density (SD, mm^{-2}), bark thickness (BT, mm), and wood density

(WD, g cm^{-3}). These structural and physiological performance traits were chosen based on the

large literature in plant functional ecology suggesting that they are key indices of carbon uptake

and allocation and water-use strategies (see Nolting et al. 2021 for further discussion of

functional significance of each trait, and of trait complexes). Further, Nolting et al. (2021)

demonstrate that many of these structural and physiological performance traits are directly

and/or indirectly associated with individual plant size and reproductive effort.

TRAIT DIFFERENCES DO NOT LEAD TO PERFORMANCE DIFFERENCES IN PROTEA

Nolting et al. (2021) showed that the measures of physiological performance we investigate

below are associated with plant size and reproductive effort, which we take to be reasonable

proxies for fitness. Thus, we focus on investigating the relationship between eight individual

structural traits and six measures of physiological performance.

Structural Traits Predict Physiological Performance

We evaluated the relationship between structural traits and each of our physiological

performance traits using Bayesian linear mixed effects models (one model for each of the six

physiological performance traits) using the package 'brms' (Bürkner 2017, 2018, 2021) (version

2.16.3) in the R programing environment (R Core Team 2020; version 4.0.2). All performance

and structural traits were standardized to a mean of zero and a unit standard deviation prior to

analysis. Sample site, species identity, and the interaction between site and species were

included as random effects. We used independent N(0,1) priors on the intercept and regression

coefficients, and half Cauchy (0,5) priors on the variance parameters. We used four chains with

a warm-up of 2000 iterations and a sampling of 2000 iterations for a total of 8000 samples from

the posterior with delta = 0.999. All parameters had Rhat < 1.01, and no divergent transitions

were reported. In Figure 1 we plot the predicted performance values versus the observed

values, and report the mean Bayesian R^2 estimates (Gelman et al. 2017). Consistent with

Nolting et al. (2021), we demonstrate that combinations of structural traits predict performance

reasonably well (i.e., full model R^2 values range from 0.26 to 0.44; marginal R^2 values – fixed

effects only – ranging from 0.19 to 0.36).

Figure 1: For each multiple regression modeling the relationship between the eight structural traits and each of the six performance traits, we plot the predicted performance values versus observed (with traits centered with a mean of zero and scaled to unit variance). (A) Ks: Stem-specific hydraulic conductance, (B) LSC: Leaf-specific conductivity, (C) Photosynthetic rate, (D) LSP: Leaf-specific photosynthetic rate, (E) Stomatal conductance, and (F) Instantaneous water-use efficiency. The dotted gray line reflects the 1:1 line. Colors represent different species and shapes represent different sites. We report the mean Bayesian R2 value for the fixed effects for each model.

Species Differ in Structural Traits but Not in Physiological Performance Traits

We evaluated the among species variation (relative to residual variation) in each of our structural and performance traits using a simple regression model with species as a random intercept. We fit these models in 'brms' as above, using the same priors, model specification, and sampling. As Nolting et al. (2021) discuss in more detail, the proportion of variation among

190 species in structural traits (Figure 2a) is much greater than that in physiological performance

191 traits (Figure 2b). As individuals sampled were of similar size within each population, size-

192 mediated trait differences are unlikely to contribute to the observed differences. In short, there

193 are substantial differences among species in structural traits that are associated with

194 performance, but there are only minimal differences among species in physiological

195 performance traits.

Figure 2: Among species variance partitioning for the eight structural traits (A) and the six performance

traits (B). The gray histograms reflect the full posterior distribution, the black circles represent the mean of

the posterior, the thick black lines reflect the 50 percent credible intervals, and the thin black line reflects

the 95 percent credible intervals. Adapted from Nolting et al. 2021, with permission from *Annals of*

Botany.

EXPLAINING THE PARADOX WITH A SIMPLE MODEL

Our intuition tells us that the variance in physiological performance is smaller than we might

expect because many different combinations of traits, i.e., alternative designs, lead to similar

performance. Here we develop that intuition further by investigating how the variance in

performance is related to covariation among the traits that influence performance. We begin

with a simple simulation to illustrate the phenomenon.

Simulating Variance in Performance in a Two-Trait Model

Assume that we have a single performance trait that is linearly predicted by two structural traits:

217
$$y_i = \beta_0 + \beta_1 x_{i1} + \beta_2 x_{i2} + \varepsilon_i \; ,$$

219   where performance trait y and structural traits x_1 and x_2 are measured on all individuals, i, with ϵ_i

220   as the residual error. β_1 and β_2 reflect the associations between traits x_1 and x_2 with y, and β_0 is

221   the intercept. For simplicity, we will assume that x_1 and x_2 are centered and scaled to have a

222   mean of zero and standard deviation of 1. Traits x_1 and x_2 are non-independent, and their

223   correlation is given by ρ.

224   For the simulation, we assume $\rho = 0.8$, and we explore two scenarios: one in which the

225   regression coefficients (β_1 and β_2) describing the relationship between our two traits and

226   performance are both positive and of the same magnitude, and a second scenario in which the

227   regression coefficients are of the same magnitude but opposite in sign:

229   **Scenario 1:** $\beta_1 = 0.5, \beta_2 = 0.5.$

230   **Scenario 2:** $\beta_1 = 0.5, \beta_2 = -0.5.$

We'll also assume $\beta_0 = 0$ (i.e., the expected intercept is zero) in our models under each

scenario, which is a reasonable approximation if we've centered and scaled the response

variable as well as the covariates. We display the outcomes under these two scenarios, by

simulating trait data under each scenario for 1000 individuals, sampled from a multivariate

normal distribution. The performance value, y_i, for each individual in each scenario was

determined from the simulated trait data and the linear model described for each scenario,

assuming a sampling variance of 0.05. Figure 3a (scenario 1) and 3b (scenario 2) reflect the

outcome of a single realization of this process. We also report the variance in our performance

trait across 1000 realizations for each scenario, and we summarize the distribution of reported

variance across realizations in Figure 3c (scenario 1) and Figure 3d (scenario 2).

Figure 3a shows the distribution of simulated trait data from one realization in scenario 1

with the different color of each point corresponding to the performance predicted for that

individual from the linear regression. In Figure 3c, we use the same simulated trait data but the

colors correspond to the performance predicted under scenario 2. The magenta dashed line

represents the regression vector associated with performance in each scenario, and the gray

arrows are the principal components associated with the trait data.

It may not be surprising that in scenario 2, where there is a positive association of trait x_1

with performance and a negative association of trait x_2 with performance, there is less variation

across the 1000 individuals in performance than in scenario 1 in which both traits have a

positive association with performance. If we examine these figures a bit more carefully, we also

notice that the regression vector is parallel to major axis of trait variation (PC1) in scenario 1

and orthogonal to it in scenario 2. This suggests that when many individual structural traits are

254   associated with a single performance trait, there might be little variation in the performance trait

255   if the regression vector is orthogonal to the major axes of trait variation.

Figure 3: Simulating variation in performance with two traits under two different scenarios. In scenario

one (A) both trait one and trait two are positively associated with performance (e.g., β_1 and $\beta_2 = 0.5$), and

in scenario two (B) trait one is positively associated with performance and trait two is negatively

associated with performance (e.g., $\beta_1 = 0.5$ and $\beta_2 = -0.5$). Trait data was simulated from a multivariate

normal distribution with a correlation matrix described by rho = 0.8, and performance values were

sampled under each scenario with this trait data and a sampling variance of 0.05. In (A) and (B) the gray

arrows correspond to the principal components of the trait data, and the magenta line reflects the

regression vector for each scenario. The black dashed ellipses reflect the 95% confidence limits. Panels

(C) and (D) summarize the variation in the performance trait, y, across 1000 realizations under each

scenario. In (C) and (D) the solid dashed lines represent the mean of the distribution, and the dashed

black lines reflect the 95% confidence limits.

270   *Developing a General Model*

272   We specify the relationship between structural traits and performance traits using the

273   multivariate version of our two-trait model:

$$
\begin{aligned}
y_i &= \beta_0 + \sum_{k=1}^{K} \beta_k x_{ik} + \varepsilon_i \\
&= \beta_0 + x_i^T \beta
\end{aligned}
,
$$

where y_i is the observed physiological performance of individual i, β_0 is the intercept, x_{ik} is the

trait value for trait k in individual i, β_k is the regression coefficient for trait k, and ϵ_i is the

residual error (with mean 0 and variance σ^2). x_i is the vector of traits in individual i and β_k ($k =$

$1, \ldots, K$) is the regression vector. We assume that traits are drawn independently from a

multivariate distribution with mean μ and covariance Σ_W. We will take expectations with respect

to the distribution of x_{ik}.

The variance of y is

$$
\begin{aligned}
Var(y_i) &= E(y_i - \mu)^2 \\
&= E\left((\beta_0 + x_i^T \beta + \epsilon_i) - (\beta_0 + \mu^T \beta) \right)^2 \\
&= E(((x_i - \mu)^T \beta)^2 + x_i^T \beta \epsilon_i + \epsilon_i^2) \\
&= \beta^T \Sigma \beta + \sigma^2
\end{aligned}
$$

291   Using the spectral decomposition theorem (Szabo 2015) we can rewrite Σ as

$$
\Sigma = \lambda_1 u_1 u_1^T + \cdots + \lambda_K u_K u_K^T
$$

295 where the eigenvalues λ_i are numbered from the largest to the smallest, and the corresponding

296 eigenvalues have unit length. This allows us to rewrite $Var(y_i)$ as

298
$$Var(y_i) = \beta^T (\lambda_1 u_1 u_1^T + \cdots + \lambda_K u_K u_K^T)\beta + \sigma^2$$

300 Suppose that the vector of regression coefficients is parallel with the *i*th principal component. It

301 is, by definition, orthogonal to all of the remaining components, and $Var(y_i)$ reduces to

303
$$Var(y_i) = \lambda_i \beta^T u_i u_i^T \beta + \sigma^2$$

Thus if the regression vector is parallel to one of the minor axes of trait variation, the variation in

performance will be much smaller than if it is parallel to one of the major axes of trait variation.

More generally, if the regression vector is mostly orthogonal to the major axes of trait variation,

there may be little variation in performance.

Applying these Insights with Protea

Recall that in the *Protea* example we observe (a) that combinations of structural trait values

explain much of the variance in physiological performance traits (Figure 1), and (b) species

differ markedly in structural traits but largely overlap with respect to physiological performance

traits (Figure 2). Our simple, generalized model suggests that the vector of regression

coefficients summarizing the trait-performance associations in four *Protea* models are not

aligned with the major axes of structural trait variation. We explore this possibility here.

Using the princomp function in R (R Core Team 2020; version 4.0.2) we performed a

principal components analysis on the centered and scaled structural trait data in our *Protea*

dataset, yielding eight principal components. We evaluated the alignment of each of these PC

axes with the vector of regression coefficients estimated from each multiple linear regression

models (i.e., there is a regression vector for each of the six physiological performance traits in

our dataset). As the generalized model presented above excludes any species or site

distinctions, we re-ran the structural trait-performance multiple regressions as before, but

excluded the site and species random effects.

We quantified the degree of alignment for each performance trait model by estimating

the angle between the vector of regression coefficients for each response and each of the eight

principal components summarizing the variation in structural traits (Figure 4). As we fit the

multiple regression models using 'brms', we can present a posterior distribution for each of the

angles, where an angle closer to 90 degrees reflects a case in which the PC axis is more

orthogonal to the regression vector, and an angle closer to zero degrees indicates the PC axis

and the regression vector are more parallel (Figure 4). While there is some heterogeneity in the

degree of alignment of the PC axes with the regression vectors representing each physiological

performance trait, it is striking that for each performance trait evaluated, the major axes of trait

variation (i.e., PC axes 1 and 2) are generally the axes that are closest to orthogonal with the

regression vector. Instead, the minor axes of trait variation (often the axes excluded from

analyses that use dimension reduction to estimate functional associations from trait matrices

and performance) are the axes most parallel with the regression vector.

We see little differentiation in performance in our data despite considerable trait

differences because the axis along which trait differences are associated with differences in

physiological performance is largely orthogonal to the axes that account for most structural trait

variation. An important implication of this result is that reducing trait variation to the first few PCs

and attempting to associate those with functional differences may fail to find real associations of

traits with functional differences.

Figure 4: Alignment of each principal component with the regression vector describing the functional

relationship between structural traits and each of six performance traits. (A) Sapwood-specific

conductivity, (B) Leaf-specific conductivity, (C) Photosynthetic rate, (D) Leaf-specific photosynthetic rate,

(E) Stomatal conductance, and (F) Instantaneous water-use efficiency. The horizontal axis is the angle

(0-90°) between each of the eight principal components for the structural trait data and the regression

vector associated with each physiological performance trait. Angles closer to 90° indicate that the

functional axis (i.e., the vector of regression coefficients) and the PC axis are largely orthogonal (angles

closer to 0° indicate they are more aligned). The gray histograms reflect the posterior distribution of

angles, the black circles represent the mean of the posterior, the thick black lines reflect the 50 percent

credible intervals, and the thin black line reflects the 95 percent credible intervals.

DISCUSSION

Given the well-established functional relationships between plant traits and aspects of plant

performance, it may be surprising that the magnitude of species differences in performance is

much less than might be expected given the often large differences among species in suites of

structural traits. Here we unraveled this paradox using a recently published dataset of plant

structural traits and physiological performance traits measured in several natural populations of

Protea species from the Cape Floristic Region, South Africa (Nolting et al. 2021). We (a)

demonstrated that combinations of structural traits predict variation in plant performance well

(Figure 1), and (b) we showed that there are clear species differences in eight leaf and wood

structural traits, but that there is extensive species overlap with respect to six physiological

performance traits measured (Figure 2).

Using a linear regression to model the association between structural traits and

physiological performance, we show that the misalignment between major axes of variation in

structural traits and regression vectors leads to a smaller amount of performance variation than

might be expected. This suggests that in future studies it will be important not only to evaluate

axes of trait variation and covariation in commonly measured functional traits, but also to

determine whether major axes of trait variation align with the axis associated with differences in

performance.

Different Structural Trait Combinations Lead to Similar Performance within and across Species

Our model shows that very different trait combinations lead to similar measures of performance

when the axis of performance variation does not align with the major axes of trait variation.

Depending on the focal study system and the scale of trait measurement, these 'alternative

designs' (Marks and Lechowicz 2006; Marks 2007) might not only exist across distantly related

taxa but might also exist *within* species that demonstrate significant trait variation.

To illustrate how combinations of structural traits can lead to similar performance both

within and across species in our dataset, we plot our sampled individuals in structural trait space

(Figure 5a,b,c: using the first two PC axes), but color the individuals based on their relative

performance values. Each panel in the top row of Figure 5 reflects the position of each

individual in the first two dimensions of structural trait space, i.e., the two most important axes of

structural trait variation. In each panel the points are in the same position, but their coloring

reflects the relative performance of each individual (using a subset of three physiological

performance traits). For example, in Figure 5A the ***position*** of each sampled individual is

determined by its position in the structural trait PCA, but the ***coloring*** of each individual is

based on the quintile of K_s (sapwood-specific hydraulic conductivity) into which it falls.

Confidence ellipses in this figure correspond to species groupings based on the first two PC

axes for structural traits. In the bottom panel of Figure 5 (Figure 5d,e,f), we color the points in

the same way but plot individuals in structural trait space using the PCs that are most aligned

with performance (i.e., the two PCs that were most parallel to the regression vector).

Figure 5: Visualizing variation in relative performance arising from different combinations of structural

traits, for a subset of performance traits (Ks, or sapwood-specific hydraulic conductivity: A and D;

Photosynthetic rate: B and E; Stomatal conductance: C and F). For each panel, individuals from our

empirical dataset are plotted with respect to their position in multivariate structural trait space. For A, B,

and C individuals are plotted using the first two principal components (PC1 and PC2). For D, E, and F

individuals are plotted in the structural trait space that is most aligned with performance (i.e., the PC axes

that are most parallel with the regression vector for each performance trait). The color of the points

reflects the relative performance of each individual, corresponding to one of five ordered categories (Low,

Medium Low, Medium, Medium High, and High). The confidence ellipses correspond to species

groupings. PRLA: *P. laurifolia*, PRNI: *P. nitida*, PREX: *P. eximia*, PRPU: *P. punctata*, and PRRE: *P.*

repens.

Consistent with the univariate examination of among species structural trait variation

(Figure 2), we see that when plotted with respect to PC1 and PC2 (the top two axes of trait

variation), species are well differentiated in structural trait space (Figure 5, top row). In contrast,

when we plot individuals in structural trait space that most closely reflects structural trait

variation associated with the functional axis describing performance (Figure 5, bottom row), we

see that species are weakly differentiated, if at all.

Figure 5 also illustrates that individuals with different combinations of structural traits

(different positions in PCA space) can have similar performance (the same color) and that there

are many different combinations of traits that can lead to performance in the highest (or lowest)

quintile. Returning to the example of sapwood-specific hydraulic conductivity (K_s), we see that

all species (other than *P. nitida*) have individuals with K_s in the highest quintile (e.g., high

relative performance) and that all five of the species have individuals with K_s in the lowest

quintile (e.g., low relative performance). The same pattern is repeated across all six

physiological performance traits measured.

Implications for Trait-Mediated Species Coexistence and Community Ecology

Basic principles of functional ecology hold that species subject to similar selection pressures

should have similar phenotypes, but basic principles of community ecology hold that species

should exhibit trait differences along at least one niche axis in order to coexist. Thus, principles

of functional ecology suggest that species should share traits that allow them to survive in a

particular environment (Keddy 1992), while principles of community ecology suggest that

species should differ in ways that allow them to coexist (Chesson 2000; HilleRisLambers et al.

2012). Environmental filtering may explain the oft-observed association between community-

weighted mean trait values and environments (Ackerly and Cornwell 2007) and evidence of

phylogenetic underdispersion in studies evaluating the relative phylogenetic diversity in local

community assemblages (Webb 2000; Webb et al. 2002; Mayfield and Levine 2010). On the

other hand, there is also substantial evidence of local trait differences within communities that

may promote coexistence (Cavender-Bares et al. 2004*a*, 2004*b*; Kraft et al. 2008; Swenson and

Enquist 2009).

The Marks and Lechowicz (2006) 'alternative designs' model may allow us to resolve

this apparent contradiction. Traits may covary in many different ways to achieve similar

performance, permitting the persistence of species that display high trait diversity within a

community and shared environment that are functionally very similar. These trait differences

promote coexistence through niche differentiation while minimizing fitness differences (Adler et

al. 2007; Umaña et al. 2016). Given the complex nature of how traits interact with one another

to affect performance (Worthy et al. 2020; Funk et al. 2021), the trait space where performance

is roughly equivalent may be extremely large (Marks and Lechowicz 2006; Laughlin 2014;

Laughlin and Messier 2015).

CONCLUSION

In this paper we show that when the major axes of trait variation are largely orthogonal to the

functional axis there may be relatively little variation in functional performance. A paucity of

variation in functional performance arises even if there is both a strong relationship between the

traits of individual organisms and their performance, and substantial variation among individuals

in those traits. To the extent that long-term co-existence among species requires similar levels

of performance within a given habitat, we therefore expect the axis of functional performance to

be orthogonal to major axes of trait variation. Interestingly, Engen and Saether (2021) recently

demonstrated a similar phenomenon in the evolution of multivariate phenotypic traits subject to

natural selection. Specifically, they showed that the G matrix (the additive genetic covariance)

evolves such that at evolutionary equilibrium the eigenvector associated with the smallest

eigenvalue is parallel to the selection gradient. In other words, within a single population natural

selection acts to eliminate genetic variation along axes aligned with fitness. Similarly, we

anticipate that competitive interactions among species will act to eliminate phenotypic variation

along axes aligned with functional performance.

REFERENCES

Ackerly, D. D., and W. K. Cornwell. 2007. A trait-based approach to community assembly:

partitioning of species trait values into within- and among-community components.

Ecology letters 10:135–145.

Adler, P. B., J. Hillerislambers, and J. M. Levine. 2007. A niche for neutrality. Ecology letters

10:95–104.

Aiello-Lammens, M. E., J. A. Slingsby, C. Merow, H. K. Mollmann, D. Euston-Brown, C. S.

Jones, and J. A. Silander Jr. 2017. Processes of community assembly in an

environmentally heterogeneous, high biodiversity region. Ecography 40:561–576.

Allsopp, N. Colville, J.F. Verboom, G.A. 2014. Fynbos: Ecology, evolution, and conservation of

a megadiverse region. Oxford University Press, New York.

Baraloto, C., C. E. Timothy Paine, S. Patiño, D. Bonal, B. Hérault, and J. Chave. 2010.

Functional trait variation and sampling strategies in species-rich plant communities.

Functional ecology 24:208–216.

Becker, P., M. T. Tyree, and M. Tsuda. 1999. Hydraulic conductances of angiosperms versus

conifers: similar transport sufficiency at the whole-plant level. Tree physiology 19:445–

452.

Brown, J. H., and G. A. Lieberman. 1973. Resource Utilization and Coexistence of Seed-Eating

Desert Rodents in Sand Dune Habitats. Ecology 54:788–797.

Bürkner, P.-C. 2017. brms: An R Package for Bayesian Multilevel Models Using Stan. Journal of

statistical software 80:1–28.

———. 2018. Advanced Bayesian Multilevel Modeling with the R Package brms. The R journal

10:395.

———. 2021. Bayesian Item Response Modeling in R with brms and Stan. Journal of statistical

software 100:1–54.

Carlson, J. E., and K. E. Holsinger. 2013. Direct and indirect selection on floral pigmentation by

pollinators and seed predators in a color polymorphic South African shrub. Oecologia

171:905–919.

Carlson, J. E., K. E. Holsinger, and R. Prunier. 2011. Plant responses to climate in the Cape

Floristic Region of South Africa: evidence for adaptive differentiation in the Proteaceae.

Evolution; international journal of organic evolution 65:108–124.

Cavender-Bares, J., D. D. Ackerly, D. A. Baum, and F. A. Bazzaz. 2004*a*. Phylogenetic

overdispersion in Floridian oak communities. The American naturalist 163:823–843.

Cavender-Bares, J., K. Kitajima, and F. A. Bazzaz. 2004*b*. MULTIPLE TRAIT ASSOCIATIONS

IN RELATION TO HABITAT DIFFERENTIATION AMONG 17 FLORIDIAN OAK

SPECIES. Ecological Monographs.

Chesson, P. 2000. Mechanisms of Maintenance of Species Diversity. Annual review of ecology

and systematics 31:343–366.

Clark, J. S., M. Dietze, S. Chakraborty, P. K. Agarwal, I. Ibanez, S. LaDeau, and M. Wolosin.

2007. Resolving the biodiversity paradox. Ecology letters 10:647–59; discussion 659-62.

Cody, M. L. 1986. Structural niches in plant communities. Pages 381–405 *in*Community

ecology. Harper & Row, New York, NY.

Cowling, R. M., P. W. Rundel, B. B. Lamont, M. Kalin Arroyo, and M. Arianoutsou. 1996. Plant

diversity in mediterranean-climate regions. Trends in ecology & evolution 11:362–366.

Díaz, S., J. Kattge, J. H. C. Cornelissen, I. J. Wright, S. Lavorel, S. Dray, B. Reu, et al. 2016.

The global spectrum of plant form and function. Nature 529:167–171.

Engen, S., and B.-E. Sæther. 2021. Structure of the G-matrix in relation to phenotypic

contributions to fitness. Theoretical population biology 138:43–56.

Espino, S., and H. J. Schenk. 2011. Mind the bubbles: achieving stable measurements of

maximum hydraulic conductivity through woody plant samples. Journal of experimental

botany 62:1119–1132.

Felix, R., Gantmacher. 1984. The Theory of Matrices.

Funk, J. L., J. E. Larson, and J. Ricks-Oddie. 2021. Plant traits are differentially linked to

performance in a semiarid ecosystem. Ecology 102:e03318.

Gelman A., Goodrich B., Gabry J., and I. Ali. 2017. R-squared for Bayesian regression models.

http://www.stat.columbia.edu/~gelman/research/unpub- lished/bayes_R2.pdf (4

November 2018).

Gillespie, R. 2004. Community Assembly Through Adaptive Radiationin Hawaiian Spiders.

Science 303.

HilleRisLambers, J., P. B. Adler, W. S. Harpole, J. M. Levine, and M. M. Mayfield. 2012.

Rethinking Community Assembly through the Lens of Coexistence Theory. Annual

review of ecology, evolution, and systematics 43:227–248.

Hubbell, S. P. 2001. The Unified Neutral Theory of Biodiversity and Biogeography. Princeton

University Press, Princeton, NJ.

Hutchinson, G. E. 1959. Homage to Santa Rosalia or Why Are There So Many Kinds of

Animals? The American naturalist 93:145–159.

Keddy, P. A. 1992. Assembly and Response Rules: Two Goals for Predictive Community

Ecology. Journal of vegetation science: official organ of the International Association for

Vegetation Science 3:157–164.

Kraft, N. J. B., R. Valencia, and D. D. Ackerly. 2008. Functional Traits and Niche-Based Tree

Community Assembly in an Amazonian Forest. Science 322:580–582.

Laughlin, D. C. 2014. The intrinsic dimensionality of plant traits and its relevance to community

assembly. (S. Wilson, ed.)The Journal of ecology 102:186–193.

Laughlin, D. C., and J. Messier. 2015. Fitness of multidimensional phenotypes in dynamic

adaptive landscapes. Trends in ecology & evolution 30:487–496.

Linder, H. P. 2003. The radiation of the Cape flora, southern Africa. Biological reviews of the

Cambridge Philosophical Society 78:597–638.

Marks, C. O. 2007. The causes of variation in tree seedling traits: the roles of environmental

selection versus chance. Evolution; international journal of organic evolution 61:455–

469.

Marks, C. O., and M. J. Lechowicz. 2006. Alternative designs and the evolution of functional

diversity. The American naturalist 167:55–66.

Mason, C. M., and L. A. Donovan. 2015. Evolution of the leaf economics spectrum in herbs:

Evidence from environmental divergences in leaf physiology across Helianthus

(Asteraceae). Evolution; international journal of organic evolution 69:2705–2720.

Mayfield, M. M., and J. M. Levine. 2010. Opposing effects of competitive exclusion on the

phylogenetic structure of communities. Ecology letters 13:1085–1093.

Mencuccini, M., and L. Bonosi. 2001. Leaf/sapwood area ratios in Scots pine show acclimation

across Europe. Canadian journal of forest research. Journal canadien de la recherche

forestiere 31:442–456.

Messier, J., B. J. McGill, and M. J. Lechowicz. 2010. How do traits vary across ecological

scales? A case for trait-based ecology. Ecology letters 13:838–848.

Mitchell, N., T. E. Moore, H. K. Mollmann, J. E. Carlson, K. Mocko, H. Martinez-Cabrera, C.

Adams, et al. 2015. Functional Traits in Parallel Evolutionary Radiations and Trait-

Environment Associations in the Cape Floristic Region of South Africa. The American

naturalist 185:525–537.

Nolting, K. M., R. Prunier, G. F. Midgley, and K. E. Holsinger. 2021. Intraspecific trait variation

influences physiological performance and fitness in the South Africa shrub genus Protea

(Proteaceae). Annals of botany 127:519–531.

Peter Linder, H. 2005. Evolution of diversity: the Cape flora. Trends in plant science 10:536–

541.

Prunier, R., K. E. Holsinger, and J. E. Carlson. 2012. The effect of historical legacy on

adaptation: do closely related species respond to the environment in the same way?

Journal of evolutionary biology 25:1636–1649.

Reich, P. B., M. B. Walters, and D. S. Ellsworth. 1997. From tropics to tundra: global

convergence in plant functioning. Proceedings of the National Academy of Sciences of

the United States of America 94:13730–13734.

Schnitzler, J., T. G. Barraclough, J. S. Boatwright, P. Goldblatt, J. C. Manning, M. P. Powell, T.

Rebelo, et al. 2011. Causes of plant diversification in the Cape biodiversity hotspot of

South Africa. Systematic biology 60:343–357.

Schoener, T. W. 1968. The Anolis Lizards of Bimini: Resource Partitioning in a Complex Fauna.

Ecology 49:704–726.

Swenson, N. G., and B. J. Enquist. 2009. Opposing assembly mechanisms in a neotropical dry

forest: implications for phylogenetic and functional community ecology. Ecology

90:2161–2170.

Szabo, F.E. 2015. The Linear Algebra Survival Guide. New York: Academic Press.

Taseski, G. M., D. A. Keith, R. L. Dalrymple, and W. K. Cornwell. 2021. Shifts in fine root traits

within and among species along a fine-scale hydrological gradient. Annals of botany

127:473–481.

Treurnicht, M., J. Pagel, J. Tonnabel, K. J. Esler, J. A. Slingsby, and F. M. Schurr. 2020.

Functional traits explain the Hutchinsonian niches of plant species. Global ecology and

biogeography: a journal of macroecology 29:534–545.

Tyree, M.T., Zimmermann, M.H. 2002. Conducting Units: Tracheids and Vessels. In: Xylem

Structure and the Ascent of Sap. Springer Series in Wood Science. Springer, Berlin,

Heidelberg.

Umaña, M. N., J. Forero-Montaña, R. Muscarella, C. J. Nytch, J. Thompson, M. Uriarte, J.

Zimmerman, et al. 2016. Interspecific Functional Convergence and Divergence and

602   Intraspecific Negative Density Dependence Underlie the Seed-to-Seedling Transition in

603   Tropical Trees. The American naturalist 187:99–109.

604 Umaña, M. N., and N. G. Swenson. 2019. Does trait variation within broadly distributed species

605   mirror patterns across species? A case study in Puerto Rico. Ecology 100:e02745.

606 Van Wilgen, B. W., G. G. Forsyth, H. De Klerk, S. Das, S. Khuluse, and P. Schmitz. 2010. Fire

607   management in Mediterranean-climate shrublands: a case study from the Cape fynbos,

608   South Africa: Fire regimes in Cape fynbos. The Journal of applied ecology 47:631–638.

609 Webb, C. O. 2000. Exploring the Phylogenetic Structure of Ecological Communities: An

610   Example for Rain Forest Trees. The American naturalist 156:145–155.

611 Webb, C. O., D. D. Ackerly, M. A. McPeek, and M. J. Donoghue. 2002. Phylogenies and

612   Community Ecology. Annual review of ecology and systematics 33:475–505.

613 Worthy, S. J., D. C. Laughlin, J. Zambrano, M. N. Umaña, C. Zhang, L. Lin, M. Cao, et al. 2020.

614   Alternative designs and tropical tree seedling growth performance landscapes. Ecology

615   101:e03007.

616 Wright, I. J., P. B. Reich, M. Westoby, D. D. Ackerly, Z. Baruch, F. Bongers, J. Cavender-Bares,

617   et al. 2004. The worldwide leaf economics spectrum. Nature 428:821–827.