1 **Environmental transcriptomics under heat stress: Can environmental RNA reveal changes**

2 **in gene expression of aquatic organisms?**

3 Short title: Using eRNA transcriptomics to assess biotic heat stress responses
4

5 Robert M. Hechler*, [a, b], Matthew C. Yates[c, d], Frédéric J. J. Chaine[e] and Melania E. Cristescu[a]
6
7 Author affiliations:
8 [a] Department of Biology, McGill University, Montréal, QC, H3A 1B1, Canada
9
10 [b] Current affiliation: Department of Ecology and Evolutionary Biology, University of Toronto,
11 Toronto, ON, M5S 3B2, Canada
12
13 [c] Département des Sciences Biologiques, Université du Québec à Montréal, Montréal, QC, H2X
14 1Y4, Canada
15
16 [d] Current affiliation: Great Lakes Institute for Environmental Research, University of Windsor,
17 Windsor, ON, N9C 1A2, Canada
18
19 [e] Department of Biological Sciences, University of Massachusetts Lowell, Lowell, MA, 01854,
20 USA
21
22 *Corresponding author
23 Email: robert.hechler@mail.utoronto.ca
24

Keywords:
environmental transcriptomics; environmental RNA (eRNA); environmental DNA (eDNA);
extra-organismal RNA; gene expression; biological monitoring; biomonitoring

Abstract: To safeguard biodiversity in a changing climate, we require taxonomic information

about species turnover and insights into the health of organisms. Environmental DNA

approaches are increasingly used for species identification, but cannot provide functional

insights. Transcriptomic methods reveal the physiological states of macroorganisms, but are

currently species specific and require tissue sampling or animal sacrifice, making community-

wide assessments challenging. Here, we test if broad functional information (expression level of

the transcribed genes) can be harnessed from environmental RNA (eRNA), which includes extra-

organismal RNA from macroorganisms along with whole microorganisms. We exposed *Daphnia*

pulex as well as phytoplankton prey and microorganism colonizers to control (20 °C) and heat

stress (28 °C) conditions for seven days. We sequenced eRNA from tank water (after complete

removal of *Daphnia*) as well as RNA from *Daphnia* tissue, enabling comparisons of extra-

organismal and organismal RNA based gene expression profiles. Both RNA types detected

similar heat stress responses of *Daphnia.* Using eRNA, we identified 32 *Daphnia* genes to be

differentially expressed following heat stress. Of these, 17 were also differentially expressed and

exhibited similar levels of relative expression in organismal RNA. In addition to the extra-

organismal *Daphnia* response, eRNA detected community-wide heat stress responses consisting

of distinct functional profiles and 121 differentially expressed genes across 8 taxa. Our study

demonstrates that environmental transcriptomics based on eRNA can non-invasively reveal gene

expression responses of macroorganisms following environmental changes, with broad potential

implications for the biomonitoring of ecological health across the trophic chain.

Introduction

In an era of ecological crises, monitoring the presence and health of organisms within an ecosystem is essential for the preservation of biodiversity. Environmental DNA is routinely used for biomonitoring, but such applications are limited to species detection (Cristescu & Hebert, 2018; Deiner et al., 2017). The RNA molecule provides an additional layer of functional information, as transcribed genes reflect an organism's physiological status (Huang et al., 2002). Thus, transcriptomics is used to infer the ecological health of organisms, but such surveys are species-specific and dependent on sampling organismal RNA (oRNA) directly, which is labor-intensive and requires tissue collection or animal sacrifice (Alvarez et al., 2015; Baillon et al., 2015; Gleason & Burton, 2015; Houde et al., 2019; Jeffries et al., 2021; Miller et al., 2017). Non-invasive environmental RNA (eRNA) approaches could overcome these limitations by providing insights into the health of populations and communities (Amarasiri et al., 2021; Cristescu, 2019; Veilleux et al., 2021; Yates et al., 2021). However, using eRNA to inform on ecological health has only been theorized (see Yates et al., 2021) and remains yet to be empirically tested.

We broadly define eRNA as RNA extracted from the environment, including extra-organismal RNA released by macroorganisms as well as whole microorganisms. Microbiologists have long extracted RNA from bulk samples to profile gene expression in microbial communities, referred to as metatranscriptomics (Frias-Lopez et al., 2008; Gilbert et al., 2008; Poretsky et al., 2005). Despite more than two decades of research, metatranscriptomics remains limited to studying the gene expression of microorganisms captured in bulk samples rich in oRNA. The general assumption has been that the eRNA released by macroorganisms into the environment is labile and degrades too rapidly to be reliably detected or quantified (but see

Cristescu, 2019). However, the RNA shed by macroorganisms into their surrounding

environment can be successfully extracted and quantified (T. Jo et al., 2022; Kagzi et al., 2022;

Littlefair et al., 2022; Marshall et al., 2021; Miyata et al., 2021; Tsuri et al., 2021; von Ammon et

al., 2019; Wood et al., 2020). Despite this evidence supporting the extractability of eRNA,

environmental transcriptomics, defined here as eRNA based gene expression profiling, remained

untested.

In aquatic systems, environmental conditions modulate biodiversity and are reflected in

changes of metatranscriptomes (Frias-Lopez et al., 2008; Salazar et al., 2019; Sunagawa et al.,

2015; Sunday et al., 2011). Temperature is a key abiotic factor that influences the physiology and

fitness of ectotherms due to the relationship between body and environmental temperatures

(Huey & Berrigan, 2001; Schulte, 2015; Rodgers, 2021). Rapid increases in temperature, such as

heatwaves, can exceed the thermal limits of aquatic ectotherms and exponentially increase

standard metabolic rates, resulting in decreases in fitness and survival (Harley et al., 2006; Reid

et al., 2019; Rodgers, 2021; Schulte, 2015). Transcriptomics is widely used to study the heat

stress response of many taxa, including fish (Akbarzadeh et al., 2018; Houde et al., 2019; Narum

& Campbell, 2015; Veilleux et al., 2015, 2018), coral holobionts (Savary et al., 2021; Voolstra et

al., 2021), molluscs (Chen et al., 2019; Gleason & Burton, 2015), and copepods (Kelly et al.,

2017; Semmouri et al., 2019). *Daphnia pulex*, a key bioindicator species, exhibits widespread

downregulation of genes involved in metabolic processes under heat stress (Becker et al., 2018;

Yampolsky et al., 2014). Clearly, gene expression surveys can provide species-specific

functional insights important for conserving aquatic biodiversity (Bozinovic & Pörtner, 2015;

Evans & Hofmann, 2012; Wikelski & Cooke, 2006). However, a species by species analysis

based on oRNA cannot capture community-wide changes without extensive efforts. Mapping

eRNA reads to the reference genomes of multiple target species and using existing

metatranscriptomic pipelines, could potentially enable environmental transcriptomics to reveal

gene expression responses across the trophic chain.

We constructed simple mock freshwater communities containing *Daphnia pulex,* as well

as three phytoplankton species and opportunistic microorganisms that colonized the artificial

lake water. The communities were exposed to control (20 °C) and near lethal *Daphnia* heat stress

(28 °C) conditions for seven days. To enable comparisons between eRNA and oRNA based gene

expression profiles, we sequenced eRNA from tank water (after complete *Daphnia* removal) and

oRNA from *Daphnia* tissue. We conducted differential gene expression analyses between the

control and heat stress conditions and compared the *Daphnia* functional stress response

identified in eRNA and oRNA. In addition to the *Daphnia* analysis, we used eRNA to investigate

the community-wide heat stress response. We hypothesized that 1) eRNA captures

transcriptional responses to heat stress and 2) eRNA contains a subset of the differentially

expressed *Daphnia* genes observed in oRNA after exposure to heat stress. Based on our findings,

we discuss the potential of eRNA based transcriptomics for the biological monitoring of

ecological communities and also provide recommendations to overcome potential limitations.

Materials and Methods

Experimental design

We constructed simple mock freshwater communities composed of *Daphnia pulex,* three

algae species (*Ankistrodesmus falcatus, Scenedesmus quadricauda,* and *Pseudokirchneriella*

subcapitata) and microorganisms that colonized the artificial lake water. The communities were

exposed to 20 °C (control) and 28 °C (neat-lethal stress for Daphnia) temperatures for seven days

to mimic natural heatwave conditions and allow for eRNA accumulation in tank water.

Communities were reared in 8 L tanks containing artificial lake water (Celis-Salgado et al.,

2008) and were seeded with a starting population of 50 parthenogenetic females of *Daphnia*

pulex (clone from Illinois, USA; 40.24, -87.78). *Daphnia* populations were kept in controlled

chambers for 45 days at control growing conditions: 20 °C, relative humidity of 50% and a

photoperiod of 16:8 hour light:dark. *Daphnia* were fed 1.5 mL algae (1:1:1 ratio of *A. falcatus, S.*

quadricauda, and *P. subcapitata*) twice per week. After 45 days, the *Daphnia* populations in all

tanks reached a minimum of 500 individuals and we began the experiment by transferring tanks

to controlled experimental chambers pre-set to the respective temperatures (20 °C and 28 °C).

Sample collection

Each temperature treatment had four tanks (biological replicates), and we collected four

eRNA and oRNA technical replicates per tank. We stirred the tanks for 60 seconds and filtered

500 mL of water through a 60 μm mesh to remove all *Daphnia* and ephippia. To capture eRNA,

the *Daphnia* free water was then filtered through a 0.7 μm glass microfiber filter. Each filter was

cut in half and placed in individual 1.5 mL microcentrifuge tubes containing 370 μL RLT buffer

(Qiagen) and 3.7 μL β-mercaptoethanol. Tubes were immediately stored at -80 °C until eRNA

extraction. Negative eRNA filtration control samples (500 mL of distilled water) were collected

at the beginning, middle and end of the sampling day, using the same methods. For each oRNA

sample, six live organisms (3 adults and 3 juveniles) per tank were haphazardly collected using a

sterile pipette and transferred to a 1.5 mL microcentrifuge tube containing 370 μL RLT buffer

(Qiagen) and 3.7 μL β-mercaptoethanol. To maintain the *Daphnia* heat stress transcriptional

signal, tubes were flash frozen in liquid nitrogen for 2 seconds, and immediately stored at -80 °C

until RNA extraction. The following system was used for sample identification:

RNAtype_Temperature&TankID (e.g. sample "eRNA_20T1" refers to eRNA collected at 20 °C

from tank 1).

Extractions of eRNA and oRNA samples

Samples of eRNA and oRNA were extracted using the RNeasy Mini Kit (Qiagen),

following the manufacturer's protocol with the following modifications. eRNA samples were

thawed on ice, gently vortexed for 5 seconds and then centrifuged for 3 minutes at 13,300 rpm to

separate the liquid from filter. To increase the eRNA yield, liquid from two technical replicates

was transferred into a 5 mL microcentrifuge tube containing 800 μL 70% EtOh and pipette

mixed. This mixture was transferred to an RNeasy spin column (700 μL at a time) and

centrifuged at 10,000 rpm for 15 seconds. We repeated this four times, until all remaining liquid

from the 5 mL microcentrifuge tube was transferred to the spin column. For oRNA, *Daphnia*

samples were thawed on ice and homogenized via sterile pestle and mortar mixer for 60 seconds.

We then added 400 μL of 70% EtOh to the homogenized tissue. The resulting mixture was

pipette mixed and transferred to an RNeasy spin column, where it was centrifuged at 10,000 rpm

for 15 seconds. The remainder of the eRNA and oRNA extraction procedures followed the

manufacturer's instructions, with RNA being eluted in 40 μL of DNase and RNase free

molecular grade water.

DNA digestion

Immediately following RNA extraction, an 18 μL aliquot of both RNA types underwent

two rounds of DNA digestion using the DNA-*free*™ DNA Removal Kit (Invitrogen) which

provided rDNase. We completed the first round of DNA digestion following the manufacturer's

instructions, but skipped the DNase inaction step. To ensure no DNA carryover, we completed a

second round of DNA digestion as follows: 2.7 μL of DNase I Buffer and 1 μL rDNase were

added and pipette mixed into each sample and the plate was gently vortexed, centrifuged, and

then incubated at 37 °C for 20 minutes. Following incubation, 2 μL of DNase Inactivation

Reagent was added to each well. The plate was incubated for 2 minutes at room temperature, and

centrifuged at 3700 rpm for 5 minutes to pellet the DNase Inactivation Reagent. The supernatant

containing RNA was transferred to individual sterile microcentrifuge tubes and stored at -80 °C

until library preparation.

RNA-seq library preparation and sequencing

We prepared libraries for whole transcriptome shotgun sequencing (RNA-seq) using the

Illumina Stranded Total RNA Prep with Ribo-Zero Plus kit (rRNA depletion) and IDT® for

Illumina® RNA UD Indexes Set A, following the manufacturer's instructions. Equal volumes of

RNA (post 2 rounds of DNA digestion) from each technical replicate were pooled and used as

input material (250 ng and 26.4-65.56 ng for oRNA and eRNA, respectively) to prepare one

biological replicate library per tank. During library amplification, we used 12 and 17 PCR cycles

for oRNA and eRNA, respectively, which followed the instructions outlined by the manufacturer

for the respective RNA input amounts. Library quantification, quality control and equimolar pool

sequencing of the 16 libraries was conducted at the McGill Genome Centre on one Illumina

NovaSeq 6000 S4 lane using paired-end 100 bp reads.

Contamination prevention

We submerged all experimental and sampling equipment in a 20% bleach solution for 15

minutes and rinsed thoroughly five times with distilled water prior to use. All eRNA samples

were processed in a pre-PCR clean laboratory dedicated to low quality environmental and

ancient DNA/RNA samples. Prior to entering the pre-PCR sterile laboratory, researchers entered

a decontamination room to change into the dedicated clean lab coats and put on face masks,

hairnets and lab clogs with shoe covers. The laboratory workbench was soaked in a 20% bleach

solution for 10 minutes and RNase WiPER™ (iNtRON Biotechnology) and 20% bleach was

used to thoroughly wipe pipettes, vortex mixers and centrifuges. Aerosol filter barrier pipette tips

were used to prevent cross-contamination of samples. Filtration negative control samples and

molecular negative control samples (consisting of reagents or DNA/RNA free molecular grade

water) were processed at each step alongside eRNA samples, and were free of contaminating

nucleic acids as identified by failed PCR amplification using universal COI primers known to

amplify *Daphnia* DNA (Leray et al., 2013). Successful DNA digestion of each eRNA and oRNA

sample was verified by failed PCR amplification of RNA post-DNA digestion x2 using universal

COI primers known to amplify *Daphnia* DNA (Leray et al., 2013). RNA-seq libraries were

prepared from filtration and molecular negative control samples and were free of contamination

as indicated by quality control checks and failed amplification conducted at the McGill Genome

Centre.

Bioinformatic pipeline and statistical approaches

Raw FASTQ files underwent initial sequencing quality inspection using FastQC

(Andrews, 2010). Low quality sequences and adapters were removed using Trimmomatic

(Bolger et al., 2014) (ILLUMINACLIP: lluminaStrandedTotalRNA_adapter.fa:2:30:15

TRAILING:30 HEADCROP:1 MINLEN:90). All statistical analyses were conducted using R (R

Core Team, 2021, R version 4.1.2). With default parameters, STAR (Dobin et al., 2013) was

used to map all eRNA and oRNA reads that passed quality control and trimming to the *Daphnia*

pulex reference genome (Z. Ye et al., 2017). FeatureCounts (Liao et al., 2014) was used on the

BAM file output from STAR to quantify gene counts. Only those genes with a sum of ten or

more counts in either 20 or 28 °C were considered detected and used in the subsequent

differential expression (DE) analyses, for both eRNA and oRNA.

DE analysis was conducted with DESeq2 (Love et al., 2014), comparing all 20 °C

samples versus all 28 °C samples, for both eRNA and oRNA samples. To determine genes as

significantly differentially expressed, we used a false discovery rate (FDR) adjusted p-value <

0.05. A Pearson's chi-squared test with Yates' continuity correction was conducted to test for an

association between the significantly differentially expressed genes identified in eRNA and

oRNA. WEGO 2.0 (J. Ye et al., 2018) was used to classify GO terms (acquired from the

reference genome) to all annotated *D. pulex* DEGs. Gene ontology enrichment analysis was

conducted using topGO (Alexa & Rahnenfuhrer, 2022) by comparing GO terms of the relevant

genes to the genomic background. A FDR corrected p-value < 0.05 indicated significant

enrichment. A principal component analysis (PCA) of the regularized log (rlog)-normalized

eRNA and oRNA gene expression profiles (300 most variant genes) at both optimal and heat

stressed conditions was conducted using DESeq2's plotPCA function (Love et al., 2014).

Heatmaps showing the relative expression (Z-score calculated for each gene) of all significantly

differentially expressed genes between 20 and 28 °C, in both *D. pulex* oRNA and eRNA were

generated using pheatmap (Kolde, 2019).

We used the SqueezeMeta metatranscriptomics pipeline for the eRNA community-wide

analysis (Tamames & Puente-Sánchez, 2019). Briefly, metatranscriptomes were assembled in

sequential mode and contigs were assembled using Megahit (Li et al., 2015). Diamond was used

to align all contigs to the GenBank nr and KEGG databases for taxonomic assignment and

KEGG ID annotation, respectively (Buchfink et al., 2015; Kanehisa & Goto, 2000). We

performed differential gene expression analyses as described above for *Daphnia*, after we

mapped, using Kallisto (Bray et al., 2016), eRNA reads to the reference genomes and

transcriptomes of species known a priori to persist in the communities (*A. falcatus, S.*

quadricauda, and *P. subcapitata*) and highly abundant species identified by SqueezeMeta

(*Brachionus plicatilis*, *Volvox cateri*, *Stylonychia lemnae*, *Chlamydomonas eustigma*)

(Aeschlimann et al., 2014; Han et al., 2019; Hirooka et al., 2017; Nag Dasgupta et al., 2018;

Prochnik et al., 2010; Schomaker & Dudycha, 2021; Suzuki et al., 2018). We used the SQMtools

R package to further analyze the SqueezeMeta data and created stacked-bar plots to represent the

most relatively abundant taxa (Puente-Sánchez et al., 2020). We conducted non-metric

multidimensional scaling analysis of all KEGG IDs to visualize the community-wide functional

profiles.

Results

Daphnia pulex *gene detection and expression profiles*

All *Daphnia* reads within environmental RNA (eRNA) samples are of extra-organismal

origin as we pre-filtered tank water at 60 μm to remove *Daphnia*. Sequencing eRNA from tank

water and organismal RNA (oRNA) from *Daphnia* tissue detected 3,919 (21%) and 17,244

(93%) genes, respectively, of the 18,449 genes that compose the *Daphnia pulex* reference

genome (Z. Ye et al., 2017). All *Daphnia* genes detected in eRNA were also detected in oRNA.

A principal component analysis of the regularized log normalized expression for the 300 most

variable genes showed separation between experimental groups (Fig. 1). The samples were

separated across the first principal component (PC1) by RNA type and across the second

principal component (PC2) by temperature, clustering eRNA, oRNA, 20 °C and 28 °C within

their respective groups. The only exception was 20 °C eRNA sample T1, which unexpectedly

clustered closely with the 28 °C eRNA group. PC1 and PC2 accounted for 64% of explained

variance of all principal components (Fig. S1).

Daphnia pulex *differential gene expression*

Comparison of gene expression profiles at 20 °C and 28 °C identified 32 significantly

differentially expressed genes (DEG; false discovery rate adjusted p-value <0.05) from eRNA (4

upregulated and 28 downregulated) and 2351 DEGs from oRNA samples (1105 upregulated and

1246 downregulated). All eRNA significant DEGs were substantially up and down regulated

(exceeding Log_2 2 fold change), as well as 41 and 202 oRNA DEGs, respectively. The same

directional change (up/down regulated) was observed in 31/32 eRNA DEGs as in oRNA (Table

S1). Of all eRNA DEGs, 17 (2 upregulated and 15 downregulated) were also significantly

differentially expressed in oRNA (Fig. S2). There was a significant association between DEGs

identified in oRNA and DEGs identified in eRNA ($\chi2 = 18.26$, df = 1, p-value = 1.92e-05). A

heatmap comparing the relative expression (Z-score) of the 17 *D. pulex* DEGs common to eRNA

and oRNA revealed four distinct clusters based on hierarchical clustering analyses, separating

samples and genes by temperature and directional change, respectively (Fig. 2). Across both

temperature conditions, eRNA and oRNA clustered together and were not separated into RNA

types, indicating that eRNA and oRNA exhibit similar levels of relative gene expression for

these DEGs. The only exception was eRNA 20 °C T1 sample which, similar to the PCA,

unexpectedly clustered with the 28 °C samples.

Daphnia pulex *functional annotation*

Gene ontology (GO) enrichment analysis of eRNA detected genes identified 50 GO terms

as enriched (FDR-corrected p-value <0.05), with 26, 6, and 18 belonging to Biological Processes

(BP), Cellular Component (CC) and Molecular Function (MF) domains, respectively (Figs. 3 and

S3). Of the 32 significant *D. pulex* DEGs identified from eRNA, 21 had GO annotations from the

reference genome (Z. Ye et al., 2017) (Fig. S4-5). GO enrichment analysis of highly DE

(exceeding Log_2 2 fold change) eRNA genes, and DEGs common to both eRNA and oRNA,

identified structural constituent of cuticle and chitin metabolic process, as enriched, respectively

(FDR-corrected p-value <0.05; Fig. 3).

GO enrichment analysis of oRNA detected genes identified 114 GO terms as enriched

(FDR-corrected p-value <0.05), with 100, 1, and 13 belonging to BP, CC and MF domains,

respectively (Figs. 3 and S3). Of the 2351 DEGs identified from oRNA, 1686 had GO

annotations from the reference genome (Z. Ye et al., 2017) (Figs. S4-5). GO enrichment analysis

of highly DE (exceeding Log_2 2 fold change) oRNA genes identified 16 GO terms as enriched

(FDR-corrected p-value <0.05), including structural constituent of cuticle, chitin metabolic

process, proteolysis, negative regulation of hydrolase activity, negative regulation of proteolysis

and response to stimulus (Fig S3).

Community composition, differential gene expression and functional annotation

eRNA reads originated from a simple mock freshwater community that we constructed,

consisting of *Daphnia pulex* as well as three phytoplankton species and opportunistic

microorganisms that colonized the artificial lake water. All non-*Daphnia* eRNA reads likely

originate from whole microorganisms that were captured on the filter. We used the SqueezeMeta

pipeline (Tamames & Puente-Sánchez, 2019) to align eRNA reads to the GenBank nr database

using its LCA algorithm, which identified 434 taxa. The community composition and relative

abundances remained similar between 20 °C and 28 °C conditions (Fig. S6). Across eRNA

samples, an average of 36% of all reads were aligned to eight species including four eukaryotes

that were a priori known to persist in the communities and four highly abundant taxa identified

by the SqueezeMeta analysis (Table S2). We detected 10,026 genes from these eight species,

with the majority originating from *D. pulex, Scenedesmus quadricauda and Ankistrodesmus*

falcatus, with 3919, 3180 and 1982 genes, respectively (Table S3). Differential gene expression

analysis between 20 °C and 28 °C eRNA samples was performed for each of the eight species

and identified a total of 121 significantly differentially expressed genes (FDR-adjusted p-value <

0.05). The highest numbers of DEGs were identified for *A. falcatus*, *D. pulex* and *S. quadricauda*

with 44, 32 and 20 DEGs respectively.

Within all eRNA reads, 12,211 KEGG IDs were identified and a non-metric

multidimensional scaling plot of these KEGG IDs revealed distinct functional profiles between

20 °C and 28 °C communities (Fig. S7). Of all KEGG IDs, 93 were identified as significantly

differentially expressed (DE; false discovery rate-adjusted p-value <0.05) between 20 °C and 28

°C (Table S4). This included KEGG IDs corresponding to proteins and pathways involved in

stress responses, such as the p53 signaling pathway, ubiquitin mediated proteolysis, metabolic

pathway and cyclin-dependent kinase 4 (Table S4).

Discussion

The use of environmental transcriptomics based on eRNA remained previously untested

despite recent evidence demonstrating that extra-organismal eRNA persists in the environment

long enough to be extracted (Kagzi et al., 2022; Littlefair et al., 2022; Marshall et al., 2021; Tsuri

et al., 2021; Wood et al., 2020). Our findings revealed that environmental transcriptomics is

sensitive in detecting transcriptional heat stress responses even without sampling source

organisms directly. From eRNA released by *Daphnia pulex* into the tank water, we detected

thousands of *D. pulex* genes and identified a subset of heat stress relevant genes to be

differentially expressed between the two temperature conditions, with levels of relative

expression similar to organismal RNA (oRNA). We also detected community wide changes in

functional profiles. Our study demonstrates the ability of environmental transcriptomics to reveal

gene expression responses of macroorganisms, and potentially complex biotic communities,

following environmental changes.

Recovery of Daphnia pulex *extra-organismal eRNA*

Several studies have recently demonstrated robust detection and persistence of extra-

organismal eRNA in the environment (Kagzi et al., 2022; Marshall et al., 2021; Wood et al.,

2020). We similarly were able to recover *Daphnia* eRNA in all samples, and had an average of

0.51% of all eRNA reads mapping to the *D. pulex* reference genome (Table S2). Our recovery of

extra-organismal eRNA reads was similar to metagenomic studies of natural aquatic ecosystems

that found between 0.27% and 1.25% of environmental DNA (eDNA) reads to be likely of extra-

organismal origin (Cowart et al., 2018; Monchamp et al., 2022; Stat et al., 2017; Székely et al.,

2021). We pre-processed our libraries with rRNA depletion, but could have potentially increased

our recovery of *Daphnia* eRNA reads by targeting eukaryotic RNA through poly(A) enrichment.

Nevertheless, our recovery of eRNA was particularly notable, given the general expectation that

within environmental samples, potentially low-abundance eRNA is 'competing' to be sequenced

with larger quantities of oRNA from whole microorganisms. This encouraging finding suggests

that most of the eRNA captured is in a protected state (e.g., cellular, vesicular, or in stable

molecular complexes) while free RNA forms, prone to degradation, are very short lived and

potentially undetectable.

Daphnia pulex *extra-organismal eRNA differential gene expression analysis*

Our broad environmental transcriptomics approach was able to detect 3,919 *D. pulex*

genes from eRNA, building upon targeted RT-PCR of eRNA which detected 6 fish genes (Tsuri

et al., 2021). As expected, we identified many more (2,351) differentially expressed genes

(DEGs) under temperature stress from *Daphnia* oRNA than eRNA (32). This discrepancy in

DEG identification can be easily attributed to large differences in the amount of RNA captured

that is specific to *Daphnia* among other organisms in the community. This is reflected in the

number of reads, 1 and 200 million, that mapped to the *D. pulex* reference genome using eRNA

and oRNA, respectively (Table S2). Having less than 1% of eRNA reads matching *D. pulex*

limited our power to detect DEGs, but is expected given that the libraries were generated from

diverse environmental samples which included RNA transcripts from over 400 taxa, whereas the

oRNA libraries were prepared from concentrated *D. pulex* tissue samples. On the other hand, we

believe this diversity of eRNA within water samples could be beneficial as it opens up exciting

possibilities of conducting community-wide surveys. For our proof-of-concept study, we

performed rRNA depletion and deep sequencing (average of 158 million reads per eRNA

sample, Table S2) in anticipation of such diverse eRNA samples to enable sufficient recovery of

D. pulex eRNA for gene expression analysis. Despite the lower abundance of *D. pulex* reads

from eRNA than oRNA, 97% of eRNA DEGs exhibited the same directional change (up/down

regulated) as in oRNA (Table S1). We found a significant association between the DEGs

identified in oRNA and those identified in eRNA, and similar levels of relative expression in

commonly shared DEGs (Fig 2). These results demonstrate that similar to conventional oRNA

tissue samples, our environmental transcriptomics approach yielded adequate sensitivity in

detecting changes in gene expression of a source organism in response to heat stress.

Changes in gene expression are often observed before phenotypic and demographic

effects occur and may therefore act as an early indication of ecological stress (Fedorenkova et

al., 2010; Jovic et al., 2017; Snell et al., 2003). We found that eRNA and oRNA recovered

similar *D. pulex* chronic heat stress responses which consisted of extensive downregulation of

genes associated with metabolic and cellular processes, membrane, binding, extracellular region,

collagen and cuticle structure (Figs. 3 and S4). These results are consistent with studies that also

found chronically heat stressed *Daphnia* to exhibit widespread downregulation of genes with

similar functions (Becker et al., 2018; Yampolsky et al., 2014). This widespread downregulation

of metabolic genes has been hypothesized to be a molecular compensatory mechanism for

Daphnia at near-lethal temperatures to sacrifice their long-term fitness for immediate survival

(Yampolsky et al., 2014).

After seven days of heat exposure, only few (4/62) heat shock genes were slightly

upregulated (log$_2$ fold change =1.1-1.3; Table S5) in *Daphnia* oRNA. This aligns with findings

suggesting heat shock genes to be less important for the chronic (days) *Daphnia* heat stress

response, but central for mitigating cell damage following acute (hours) heat exposure (Becker et

al., 2018; Yampolsky et al., 2014). Multiple tropical marine fish species similarly do not

differentially express heat shock genes during chronic heat stress (Veilleux et al., 2015),

suggesting that heat shock genes might not be optimal for assessments of communities under

chronic heat stress. Furthermore, we detected chitin related gene ontology (GO) terms to be

commonly enriched in both eRNA and oRNA DEGs (Fig. 3), corroborating *Daphnia* studies that

linked chitin genes to environmental stressors (Becker et al., 2018; Chain et al., 2019; Connon et

al., 2008; Shaw et al., 2007). Hydrolytic and proteolytic related GO terms were enriched in

eRNA (Fig. 3), and are commonly involved in *Daphnia* and copepod heat stress response by

stabilizing and cleaning up misfolded proteins (Becker et al., 2018; Kelly et al., 2017). It is

promising that using eRNA, we were able to detect the widespread downregulation of genes

associated with chronic heat stress with similar levels of relative expression as oRNA. We also

found that many GO terms were commonly shared in both RNA types, and that stress related

terms were enriched in eRNA. Collectively, our results indicate that environmental

transcriptomics based on eRNA is sensitive in detecting heat stress responses of source

organisms through changes in gene expression and functional GO terms. Future empirical work

is necessary to determine how our results extend to more complex natural ecosystems.

Community composition and function

We found the community composition to remain largely unchanged for 20 °C and 28 °C

conditions, after mapping all eRNA reads to GenBank nr (Fig. S6). Since we pre-filtered the tank

water at 60 μm to remove *Daphnia*, all non-*Daphnia* RNA recovered in our eRNA samples

likely originates from whole microorganisms that passed the filter. Within our eRNA samples,

gene expression analysis of eight species detected 10,026 genes with 121 of these being

significantly differentially expressed (DE) between 20 °C and 28 °C (Table S3). Although

community compositions remained similar, analysis of KEGG orthologs revealed distinct

functional profiles for communities in the two temperature treatments (Fig. S7). We found stress-

related KEGG orthologs to be DE, including proteins involved in binding, kinase and

oxidoreductase (Table S4). Our community wide results reflect similar patterns of differential

gene expression and shifts in functional terms observed in conventional aquatic

metatranscriptomic studies based on bulk microorganism samples (Aylward et al., 2015; Frias-

Lopez et al., 2008; Gilbert et al., 2008; Marchetti et al., 2012; Moniruzzaman et al., 2017;

Poretsky et al., 2005; Salazar et al., 2019; Vorobev et al., 2020). For example, a freshwater

metatranscriptome likely under heat stress exhibited similar stress responses which included

oxidoreductase, binding and kinase proteins (Trench-Fiol & Fink, 2020). We build upon this rich

metatranscriptomics literature by demonstrating that in addition to yielding microorganism

insights, environmental transcriptomics can also non-invasively detect the gene expression

response of macroorganisms to environmental stress.

Recommendations and future directions

Interest in non-lethal transcriptomics is growing, as such an approach increases animal

welfare, allows for repeated survey of individuals, and linking gene expression to fitness

following exposure to stress (Czypionka et al., 2015; Jeffries et al., 2021; Veldhoen et al., 2014).

However, conventional non-lethal methods are species specific and involve animal handling and

sampling for blood or tissue, which may still result in organism stress and reduced survival upon

release into the environment (Martins et al., 2018; Portz et al., 2006; Young et al., 2019).

Inferring the physiological state of organisms from non-invasive eRNA samples would

circumvent these stressful sampling procedures and represent a substantial improvement for

animal welfare. We demonstrated that similar to the conventional oRNA approach,

environmental transcriptomics based on eRNA can non-invasively detect heat stress responses of

progenitor organisms.

An exciting prospect of environmental transcriptomics is the potential to monitor the

ecological health of complex communities, including multiple macroorganisms. Biomonitoring

traditionally relies upon identifying bioindicator taxa that are correlated with environmental

conditions, but transcriptomic surveys can provide valuable early warnings since gene expression

changes are likely to occur before compositional shifts could be detected (Veilleux et al., 2021).

Conventional metatranscriptomics has demonstrated that gene expression profiling of

microorganisms reflects environmental conditions (Marchetti et al., 2012; Moran et al., 2013;

Salazar et al., 2019; Shi et al., 2011). Environmental transcriptomics extends beyond

microorganisms and has the potential to provide functional information of species across the

trophic chain as eRNA includes extra-organismal RNA from a diversity of macroorganisms.

However, further empirical work is required before such community-wide surveys should be

applied to biomonitoring efforts; population-level assessments that couple eRNA with targeted

approaches (e.g. qPCR, ddPCR) and stress biomarkers are more realistically applicable in the

short-term. Nevertheless, gaining insights into the ecological health of populations and

communities would enable conservation initiatives to be proactive in their efforts and respond

before local extinction events occur. Environmental transcriptomics is in its infancy and

refinement through mesocosm experiments and natural environment studies must first be

conducted before this approach can be applied further.

We anticipate multiple challenges for environmental transcriptomic biomonitoring in

natural environments, such as achieving adequate sequencing depth of eRNA in a cost-efficient

manner (Table 1). However, sequencing technologies are rapidly advancing and the costs

continue to decrease (Wetterstrand, 2021). The continuation of such technological advancement

will enable future studies to generate more reads less expensively, thereby increasing the

sensitivity of environmental transcriptomics to detect extra-organismal eRNA transcripts.

Additionally, the continued development of low input mRNA-seq library preparation kits will

enable researchers to prepare libraries for environmental transcriptomics with minimal template

eRNA. Molecular technological advancements that enhance extraction efficiency, enrich

eukaryotic RNA and remove ribosomal RNA from diverse samples may increase the yield of

coding eRNA and thus the power of environmental transcriptomics.

While sequencing and molecular advancements will resolve challenges associated with

eRNA depth, perhaps the largest remaining challenge for environmental transcriptomics to

overcome is the lack of a baseline to compare against. Differential gene expression analyses are

conducted by comparing a treatment against a control 'baseline' to identify genes involved with

the stress response. However, in natural ecosystems, it is difficult to obtain a control baseline due

to population demographics and multiple stressors acting simultaneously on a system. This is

similarly shared by metatranscriptomics, and multiple statistical and mathematical modelling

solutions have been developed to overcome this problem (Klingenberg & Meinicke, 2017;

Mallick et al., 2021; Zhang et al., 2021).

By providing functional insights in a non-invasive manner, eRNA based approaches

could push beyond the limitations of eDNA species detection and conventional tissue-based

transcriptomic surveys. We demonstrated that environmental transcriptomics can detect

thousands of genes, and that a subset of functionally relevant genes can be identified as

differentially expressed under heat stress, with levels of relative expression similar to the

conventional oRNA approach. Within all eRNA reads, we were able to reveal changes in

functional profiles for the community in response to heat stress. Collectively, these findings

demonstrate the ability of eRNA to detect changes in gene expression following environmental

stress and the potential of environmental transcriptomics to provide functional information

across the trophic chain. Future empirical work is necessary to determine if this approach can be

extended to natural ecosystems.

Data accessibility:

Sequences for all RNA-seq libraries have been uploaded to the NCBI SRA database and will be

made public upon acceptance of this manuscript. Reviewers can preview the uploaded files:

dataview.ncbi.nlm.nih.gov/object/PRJNA830892?reviewer=mcvpem4l5cupljoaj9jn53kg1e

Author contributions:

RMH and MEC designed the study. RMH and MCY conducted the experiment. RMH conducted

all molecular work. RMH and FJJC performed the bioinformatic analysis and produced the

figures. RMH wrote the first draft of the manuscript and all authors contributed to editing the

manuscript.

Acknowledgements

We thank Elif Irmak Bektas, Cathy Shen and Gabrielle Pellegrin for assisting with animal

husbandry and sample collection. We are grateful to Lars Iversen and Caren Helbing for

providing comments on an early version of this manuscript and Takahiro Maruki for input

regarding the bioinformatic analysis. This study was supported by the Natural Sciences and

Engineering Research Council (NSERC) of Canada through the Canada Research Chair program

and a Discovery Grant to Melania Cristescu.

References

Aeschlimann, S. H., Jönsson, F., Postberg, J., Stover, N. A., Petera, R. L., Lipps, H.-J., Nowacki, M., & Swart, E. C. (2014). The draft assembly of the radically organized stylonychia lemnae macronuclear genome. *Genome Biology and Evolution*, *6*(7), 1707–1723. https://doi.org/10.1093/gbe/evu139

Akbarzadeh, A., Günther, O. P., Houde, A. L., Li, S., Ming, T. J., Jeffries, K. M., Hinch, S. G., & Miller, K. M. (2018). Developing specific molecular biomarkers for thermal stress in salmonids. *BMC Genomics*, *19*(1), 749. https://doi.org/10.1186/s12864-018-5108-9

Alexa, A., & Rahnenfuhrer, J. (2022). *Topgo: Enrichment analysis for gene ontology* (2.46.0). Bioconductor version: Release (3.14). https://doi.org/10.18129/B9.bioc.topGO

Alvarez, M., Schrey, A. W., & Richards, C. L. (2015). Ten years of transcriptomics in wild populations: What have we learned about their ecology and evolution? *Molecular Ecology*, *24*(4), 710–725. https://doi.org/10.1111/mec.13055

Amarasiri, M., Furukawa, T., Nakajima, F., & Sei, K. (2021). Pathogens and disease vectors/hosts monitoring in aquatic environments: Potential of using eDNA/eRNA based approach. *Science of The Total Environment*, *796*, 148810. https://doi.org/10.1016/j.scitotenv.2021.148810

Andrews, S. (2010). *FastQC: a quality control tool for high throughput sequence data*. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Aylward, F. O., Eppley, J. M., Smith, J. M., Chavez, F. P., Scholin, C. A., & DeLong, E. F. (2015). Microbial community transcriptional networks are conserved in three domains at ocean basin scales. *Proceedings of the National Academy of Sciences*, *112*(17), 5443–5448. https://doi.org/10.1073/pnas.1502883112

Baillon, L., Pierron, F., Coudret, R., Normendeau, E., Caron, A., Peluhet, L., Labadie, P., Budzinski, H., Durrieu, G., Sarraco, J., Elie, P., Couture, P., Baudrimont, M., & Bernatchez, L. (2015). Transcriptome profile analysis reveals specific signatures of pollutants in Atlantic eels. *Ecotoxicology*, *24*(1), 71–84. https://doi.org/10.1007/s10646-014-1356-x

Becker, D., Reydelet, Y., Lopez, J. A., Jackson, C., Colbourne, J. K., Hawat, S., Hippler, M., Zeis, B., & Paul, R. J. (2018). The transcriptomic and proteomic responses of Daphnia pulex to changes in temperature and food supply comprise environment-specific and clone-specific elements. *BMC Genomics*, *19*. https://doi.org/10.1186/s12864-018-4742-6

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics (Oxford, England)*, *30*(15), 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Bozinovic, F., & Pörtner, H.-O. (2015). Physiological ecology meets climate change. *Ecology and Evolution*, *5*(5), 1025–1030. https://doi.org/10.1002/ece3.1403

Bray, N. L., Pimentel, H., Melsted, P., & Pachter, L. (2016). Near-optimal probabilistic RNA-seq quantification. *Nature Biotechnology*, *34*(5), Article 5. https://doi.org/10.1038/nbt.3519

Buchfink, B., Xie, C., & Huson, D. H. (2015). Fast and sensitive protein alignment using DIAMOND. *Nature Methods*, *12*(1), Article 1. https://doi.org/10.1038/nmeth.3176

Celis-Salgado, M. P., Cairns, A., Kim, N., & Yan, N. D. (2008). The FLAMES medium: A new, soft-water culture and bioassay medium for Cladocera. *SIL Proceedings, 1922-2010*, *30*(2), 265–271. https://doi.org/10.1080/03680770.2008.11902123

Chain, F. J. J., Finlayson, S., Crease, T., & Cristescu, M. (2019). Variation in transcriptional
responses to copper exposure across Daphnia pulex lineages. *Aquatic Toxicology*, *210*,
85–97. https://doi.org/10.1016/j.aquatox.2019.02.016

Chen, N., Huang, Z., Lu, C., Shen, Y., Luo, X., Ke, C., & You, W. (2019). Different
transcriptomic responses to thermal stress in heat-tolerant and heat-sensitive pacific
abalones indicated by cardiac performance. *Frontiers in Physiology*, *9*.
https://doi.org/10.3389/fphys.2018.01895

Connon, R., Hooper, H. L., Sibly, R. M., Lim, F.-L., Heckmann, L.-H., Moore, D. J., Watanabe,
H., Soetaert, A., Cook, K., Maund, S. J., Hutchinson, T. H., Moggs, J., Coen, W. D.,
Iguchi, T., & Callaghan, A. (2008). Linking molecular and population stress responses in
daphnia magna exposed to cadmium. *Environmental Science & Technology*, *42*(6), 2181–
2188. https://doi.org/10.1021/es702469b

Cordier, T., Lanzén, A., Apothéloz-Perret-Gentil, L., Stoeck, T., & Pawlowski, J. (2019).
Embracing environmental genomics and machine learning for routine biomonitoring.
Trends in Microbiology, *27*(5), 387–397. https://doi.org/10.1016/j.tim.2018.10.012

Cowart, D. A., Murphy, K. R., & Cheng, C.-H. C. (2018). Metagenomic sequencing of
environmental DNA reveals marine faunal assemblages from the West Antarctic
Peninsula. *Marine Genomics*, *37*, 148–160. https://doi.org/10.1016/j.margen.2017.11.003

Cristescu, M. E. (2019). Can environmental rna revolutionize biodiversity science? *Trends in*
Ecology & Evolution, *34*(8), 694–697. https://doi.org/10.1016/j.tree.2019.05.003

Cristescu, M. E., & Hebert, P. D. N. (2018). Uses and misuses of environmental dna in
biodiversity science and conservation. *Annual Review of Ecology, Evolution, and*
Systematics, *49*(1), 209–230. https://doi.org/10.1146/annurev-ecolsys-110617-062306

Czypionka, T., Krugman, T., Altmüller, J., Blaustein, L., Steinfartz, S., Templeton, A. R., &
Nolte, A. W. (2015). Ecological transcriptomics – a non-lethal sampling approach for
endangered fire salamanders. *Methods in Ecology and Evolution*, *6*(12), 1417–1425.
https://doi.org/10.1111/2041-210X.12431

Deiner, K., Bik, H. M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., Creer,
S., Bista, I., Lodge, D. M., Vere, N. de, Pfrender, M. E., & Bernatchez, L. (2017).
Environmental DNA metabarcoding: Transforming how we survey animal and plant
communities. *Molecular Ecology*, *26*(21), 5872–5895. https://doi.org/10.1111/mec.14350

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,
M., & Gingeras, T. R. (2013). STAR: Ultrafast universal RNA-seq aligner.
Bioinformatics, *29*(1), 15–21. https://doi.org/10.1093/bioinformatics/bts635

Evans, T. G., & Hofmann, G. E. (2012). Defining the limits of physiological plasticity: How
gene expression can assess and predict the consequences of ocean change. *Philosophical*
Transactions of the Royal Society B: Biological Sciences, *367*(1596), 1733–1745.
https://doi.org/10.1098/rstb.2012.0019

Fedorenkova, A., Vonk, J. A., Lenders, H. J. R., Ouborg, N. J., Breure, A. M., & Hendriks, A. J.
(2010). Ecotoxicogenomics: Bridging the gap between genes and populations.
Environmental Science & Technology, *44*(11), 4328–4333.
https://doi.org/10.1021/es9037287

Frias-Lopez, J., Shi, Y., Tyson, G. W., Coleman, M. L., Schuster, S. C., Chisholm, S. W., &
DeLong, E. F. (2008). Microbial community gene expression in ocean surface waters.
Proceedings of the National Academy of Sciences of the United States of America,
105(10), 3805–3810. https://doi.org/10.1073/pnas.0708897105

Fulda, S., Gorman, A. M., Hori, O., & Samali, A. (2010). Cellular stress responses: Cell survival
606   and cell death. *International Journal of Cell Biology*, *2010*, 214074.
607   https://doi.org/10.1155/2010/214074

608   Gilbert, J. A., Field, D., Huang, Y., Edwards, R., Li, W., Gilna, P., & Joint, I. (2008). Detection
609   of large numbers of novel sequences in the metatranscriptomes of complex marine
610   microbial communities. *PLoS ONE*, *3*(8). https://doi.org/10.1371/journal.pone.0003042

611   Gleason, L. U., & Burton, R. S. (2015). RNA-seq reveals regional differences in transcriptome
612   response to heat stress in the marine snail Chlorostoma funebralis. *Molecular Ecology*,
613   *24*(3), 610–627. https://doi.org/10.1111/mec.13047

614   Han, J., Park, J. C., Choi, B.-S., Kim, M.-S., Kim, H.-S., Hagiwara, A., Park, H. G., Lee, B.-Y.,
615   & Lee, J.-S. (2019). The genome of the marine monogonont rotifer Brachionus plicatilis:
616   Genome-wide expression profiles of 28 cytochrome P450 genes in response to
617   chlorpyrifos and 2-ethyl-phenanthrene. *Aquatic Toxicology*, *214*, 105230.
618   https://doi.org/10.1016/j.aquatox.2019.105230

619   Harley, C. D. G., Randall Hughes, A., Hultgren, K. M., Miner, B. G., Sorte, C. J. B., Thornber,
620   C. S., Rodriguez, L. F., Tomanek, L., & Williams, S. L. (2006). The impacts of climate
621   change in coastal marine systems. *Ecology Letters*, *9*(2), 228–241.
622   https://doi.org/10.1111/j.1461-0248.2005.00871.x

623   He, S., Wurtzel, O., Singh, K., Froula, J. L., Yilmaz, S., Tringe, S. G., Wang, Z., Chen, F.,
624   Lindquist, E. A., Sorek, R., & Hugenholtz, P. (2010). Validation of two ribosomal RNA
625   removal methods for microbial metatranscriptomics. *Nature Methods*, *7*(10), Article 10.
626   https://doi.org/10.1038/nmeth.1507

627   Hempel, C. A., Wright, N., Harvie, J., Hleap, J. S., Adamowicz, S. J., & Steinke, D. (2022).
628   Metagenomics versus total RNA sequencing: Most accurate data-processing tools,
629   microbial identification accuracy and perspectives for ecological assessments. *Nucleic
630   Acids Research*, *50*(16), 9279–9293. https://doi.org/10.1093/nar/gkac689

631   Hirooka, S., Hirose, Y., Kanesaki, Y., Higuchi, S., Fujiwara, T., Onuma, R., Era, A., Ohbayashi,
632   R., Uzuka, A., Nozaki, H., Yoshikawa, H., & Miyagishima, S. (2017). Acidophilic green
633   algal genome provides insights into adaptation to an acidic environment. *Proceedings of
634   the National Academy of Sciences*, *114*(39), E8304–E8313.
635   https://doi.org/10.1073/pnas.1707072114

636   Houde, A. L. S., Akbarzadeh, A., Günther, O. P., Li, S., Patterson, D. A., Farrell, A. P., Hinch, S.
637   G., & Miller, K. M. (2019). Salmonid gene expression biomarkers indicative of
638   physiological responses to changes in salinity and temperature, but not dissolved oxygen.
639   *Journal of Experimental Biology*, *222*(13). https://doi.org/10.1242/jeb.198036

640   Huang, W., Sher, Y.-P., Peck, K., & Fung, Y. C. B. (2002). Matching gene activity with
641   physiological functions. *Proceedings of the National Academy of Sciences*, *99*(5), 2603–
642   2608. https://doi.org/10.1073/pnas.042684399

643   Huey, R. B., & Berrigan, D. (2001). Temperature, demography, and ectotherm fitness. *The
644   American Naturalist*, *158*(2), 204–210. https://doi.org/10.1086/321314

645   Jeffries, K. M., Teffer, A., Michaleski, S., Bernier, N. J., Heath, D. D., & Miller, K. M. (2021).
646   The use of non-lethal sampling for transcriptomics to assess the physiological status of
647   wild fishes. *Comparative Biochemistry and Physiology Part B: Biochemistry and
648   Molecular Biology*, *256*, 110629. https://doi.org/10.1016/j.cbpb.2021.110629

649 Jo, T. S., Matsuda, N., Hirohara, T., & Yamanaka, H. (2022). *Simple and efficient preservation*
650   *of fish environmental RNA in filtered water samples via RNAlater* [Preprint]. Research
651   Square. https://doi.org/doi.org/10.21203/rs.3.rs-2170577/v1

652 Jo, T., Tsuri, K., Hirohara, T., & Yamanaka, H. (2022). Warm temperature and alkaline
653   conditions accelerate environmental RNA degradation. *Environmental DNA*, *00*, 1–13.
654   https://doi.org/10.1002/edn3.334

655 Jovic, K., Sterken, M. G., Grilli, J., Bevers, R. P. J., Rodriguez, M., Riksen, J. A. G., Allesina, S.,
656   Kammenga, J. E., & Snoek, L. B. (2017). Temporal dynamics of gene expression in heat-
657   stressed Caenorhabditis elegans. *PLOS ONE*, *12*(12), e0189445.
658   https://doi.org/10.1371/journal.pone.0189445

659 Kagzi, K., Hechler, R. M., Fussmann, G. F., & Cristescu, M. E. (2022). Environmental RNA
660   degrades more rapidly than environmental DNA across a broad range of pH conditions.
661   *Molecular Ecology Resources*, *22*(7), 2640–2650. https://doi.org/10.1111/1755-
662   0998.13655

663 Kanehisa, M., & Goto, S. (2000). Kegg: Kyoto encyclopedia of genes and genomes. *Nucleic*
664   *Acids Research*, *28*(1), 27–30. https://doi.org/10.1093/nar/28.1.27

665 Kelly, M. W., Pankey, M. S., DeBiasse, M. B., & Plachetzki, D. C. (2017). Adaptation to heat
666   stress reduces phenotypic and transcriptional plasticity in a marine copepod. *Functional*
667   *Ecology*, *31*(2), 398–406. https://doi.org/10.1111/1365-2435.12725

668 Klingenberg, H., & Meinicke, P. (2017). How to normalize metatranscriptomic count data for
669   differential expression analysis. *PeerJ*, *5*, e3859. https://doi.org/10.7717/peerj.3859

670 Kolde, R. (2019). *pheatmap: Pretty Heatmaps* (1.0.12). https://CRAN.R-
671   project.org/package=pheatmap

672 Leray, M., Yang, J. Y., Meyer, C. P., Mills, S. C., Agudelo, N., Ranwez, V., Boehm, J. T., &
673   Machida, R. J. (2013). A new versatile primer set targeting a short fragment of the
674   mitochondrial COI region for metabarcoding metazoan diversity: Application for
675   characterizing coral reef fish gut contents. *Frontiers in Zoology*, *10*(1), 34.
676   https://doi.org/10.1186/1742-9994-10-34

677 Lewin, H. A., Robinson, G. E., Kress, W. J., Baker, W. J., Coddington, J., Crandall, K. A.,
678   Durbin, R., Edwards, S. V., Forest, F., Gilbert, M. T. P., Goldstein, M. M., Grigoriev, I.
679   V., Hackett, K. J., Haussler, D., Jarvis, E. D., Johnson, W. E., Patrinos, A., Richards, S.,
680   Castilla-Rubio, J. C., … Zhang, G. (2018). Earth BioGenome Project: Sequencing life for
681   the future of life. *Proceedings of the National Academy of Sciences*, *115*(17), 4325–4333.
682   https://doi.org/10.1073/pnas.1720115115

683 Li, D., Liu, C.-M., Luo, R., Sadakane, K., & Lam, T.-W. (2015). MEGAHIT: An ultra-fast
684   single-node solution for large and complex metagenomics assembly via succinct de
685   Bruijn graph. *Bioinformatics*, *31*(10), 1674–1676.
686   https://doi.org/10.1093/bioinformatics/btv033

687 Liao, Y., Smyth, G. K., & Shi, W. (2014). featureCounts: An efficient general purpose program
688   for assigning sequence reads to genomic features. *Bioinformatics*, *30*(7), 923–930.
689   https://doi.org/10.1093/bioinformatics/btt656

690 Littlefair, J. E., Rennie, M. D., & Cristescu, M. E. (2022). Environmental nucleic acids: A field-
691   based comparison for monitoring freshwater habitats using eDNA and eRNA. *Molecular*
692   *Ecology Resources*, *0*, 1–13. https://doi.org/10.1111/1755-0998.13671

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550.
https://doi.org/10.1186/s13059-014-0550-8
Mallick, H., Rahnavard, A., McIver, L. J., Ma, S., Zhang, Y., Nguyen, L. H., Tickle, T. L.,
Weingart, G., Ren, B., Schwager, E. H., Chatterjee, S., Thompson, K. N., Wilkinson, J.
E., Subramanian, A., Lu, Y., Waldron, L., Paulson, J. N., Franzosa, E. A., Bravo, H. C.,
& Huttenhower, C. (2021). Multivariable association discovery in population-scale meta-
omics studies. *PLOS Computational Biology*, *17*(11), e1009442.
https://doi.org/10.1371/journal.pcbi.1009442
Marchetti, A., Schruth, D. M., Durkin, C. A., Parker, M. S., Kodner, R. B., Berthiaume, C. T.,
Morales, R., Allen, A. E., & Armbrust, E. V. (2012). Comparative metatranscriptomics
identifies molecular bases for the physiological responses of phytoplankton to varying
iron availability. *Proceedings of the National Academy of Sciences*, *109*(6), E317–E325.
https://doi.org/10.1073/pnas.1118408109
Marshall, N. T., Vanderploeg, H. A., & Chaganti, S. R. (2021). Environmental (e)RNA advances
the reliability of eDNA by predicting its age. *Scientific Reports*, *11*(1), Article 1.
https://doi.org/10.1038/s41598-021-82205-4
Martins, C. L., Walker, T. I., & Reina, R. D. (2018). Stress-related physiological changes and
post-release survival of elephant fish (Callorhinchus milii) after longlining, gillnetting,
angling and handling in a controlled setting. *Fisheries Research*, *204*, 116–124.
https://doi.org/10.1016/j.fishres.2018.01.016
Miller, K. M., Günther, O. P., Li, S., Kaukinen, K. H., & Ming, T. J. (2017). Molecular indices
of viral disease development in wild migrating salmon. *Conservation Physiology*, *5*(1),
cox036. https://doi.org/10.1093/conphys/cox036
Miyata, K., Inoue, Y., Amano, Y., Nishioka, T., Yamane, M., Kawaguchi, T., Morita, O., &
Honda, H. (2021). Fish environmental RNA enables precise ecological surveys with high
positive predictivity. *Ecological Indicators*, *128*, 107796.
https://doi.org/10.1016/j.ecolind.2021.107796
Monchamp, M.-È., Walsh, D. A., Garner, R. E., Kraemer, S. A., Beisner, B. E., Cristescu, M. E.,
& Gregory-Eaves, I. (2022). Comparative analysis of zooplankton diversity in
freshwaters: What can we gain from metagenomic analysis? *Environmental DNA*, *0*, 1–
15. https://doi.org/10.1002/edn3.335
Moniruzzaman, M., Wurch, L. L., Alexander, H., Dyhrman, S. T., Gobler, C. J., & Wilhelm, S.
W. (2017). Virus-host relationships of marine single-celled eukaryotes resolved from
metatranscriptomics. *Nature Communications*, *8*(1), 16054.
https://doi.org/10.1038/ncomms16054
Moran, M. A., Satinsky, B., Gifford, S. M., Luo, H., Rivers, A., Chan, L.-K., Meng, J., Durham,
B. P., Shen, C., Varaljay, V. A., Smith, C. B., Yager, P. L., & Hopkinson, B. M. (2013).
Sizing up metatranscriptomics. *The ISME Journal*, *7*(2), Article 2.
https://doi.org/10.1038/ismej.2012.94
Nag Dasgupta, C., Nayaka, S., Toppo, K., Singh, A. K., Deshpande, U., & Mohapatra, A. (2018).
Draft genome sequence and detailed characterization of biofuel production by oleaginous
microalga Scenedesmus quadricauda LWG002611. *Biotechnology for Biofuels*, *11*(1),
308. https://doi.org/10.1186/s13068-018-1308-4

Narum, S. R., & Campbell, N. R. (2015). Transcriptomic response to heat stress among
ecologically divergent populations of redband trout. *BMC Genomics*, *16*(1), 103.
https://doi.org/10.1186/s12864-015-1246-5
Poretsky, R. S., Bano, N., Buchan, A., LeCleir, G., Kleikemper, J., Pickering, M., Pate, W. M.,
Moran, M. A., & Hollibaugh, J. T. (2005). Analysis of microbial gene transcripts in
environmental samples. *Applied and Environmental Microbiology*, *71*(7), 4121–4126.
https://doi.org/10.1128/AEM.71.7.4121-4126.2005
Portz, D. E., Woodley, C. M., & Cech, J. J. (2006). Stress-associated impacts of short-term
holding on fishes. *Reviews in Fish Biology and Fisheries*, *16*(2), 125–170.
https://doi.org/10.1007/s11160-006-9012-z
Prochnik, S. E., Umen, J., Nedelcu, A. M., Hallmann, A., Miller, S. M., Nishii, I., Ferris, P.,
Kuo, A., Mitros, T., Fritz-Laylin, L. K., Hellsten, U., Chapman, J., Simakov, O., Rensing,
S. A., Terry, A., Pangilinan, J., Kapitonov, V., Jurka, J., Salamov, A., … Rokhsar, D. S.
(2010). Genomic analysis of organismal complexity in the multicellular green alga
volvox carteri. *Science*. https://doi.org/10.1126/science.1188800
Puente-Sánchez, F., García-García, N., & Tamames, J. (2020). SQMtools: Automated processing
and visual analysis of 'omics data with R and anvi'o. *BMC Bioinformatics*, *21*(1), 358.
https://doi.org/10.1186/s12859-020-03703-2
Reid, A. J., Carlson, A. K., Creed, I. F., Eliason, E. J., Gell, P. A., Johnson, P. T. J., Kidd, K. A.,
MacCormack, T. J., Olden, J. D., Ormerod, S. J., Smol, J. P., Taylor, W. W., Tockner, K.,
Vermaire, J. C., Dudgeon, D., & Cooke, S. J. (2019). Emerging threats and persistent
conservation challenges for freshwater biodiversity. *Biological Reviews of the Cambridge
Philosophical Society*, *94*(3), 849–873. https://doi.org/10.1111/brv.12480
Rodgers, E. M. (2021). Adding climate change to the mix: Responses of aquatic ectotherms to
the combined effects of eutrophication and warming. *Biology Letters*, *17*(10), 20210442.
https://doi.org/10.1098/rsbl.2021.0442
Salazar, G., Paoli, L., Alberti, A., Huerta-Cepas, J., Ruscheweyh, H.-J., Cuenca, M., Field, C.
M., Coelho, L. P., Cruaud, C., Engelen, S., Gregory, A. C., Labadie, K., Marec, C.,
Pelletier, E., Royo-Llonch, M., Roux, S., Sánchez, P., Uehara, H., Zayed, A. A., …
Sunagawa, S. (2019). Gene expression changes and community turnover differentially
shape the global ocean metatranscriptome. *Cell*, *179*(5), 1068-1083.e21.
https://doi.org/10.1016/j.cell.2019.10.014
Savary, R., Barshis, D. J., Voolstra, C. R., Cárdenas, A., Evensen, N. R., Banc-Prandi, G., Fine,
M., & Meibom, A. (2021). Fast and pervasive transcriptomic resilience and acclimation
of extremely heat-tolerant coral holobionts from the northern Red Sea. *Proceedings of the
National Academy of Sciences*, *118*(19). https://doi.org/10.1073/pnas.2023298118
Schomaker, R. A., & Dudycha, J. L. (2021). De novo transcriptome assembly of the green alga
Ankistrodesmus falcatus. *PLOS ONE*, *16*(5), e0251668.
https://doi.org/10.1371/journal.pone.0251668
Schulte, P. M. (2015). The effects of temperature on aerobic metabolism: Towards a mechanistic
understanding of the responses of ectotherms to a changing environment. *Journal of
Experimental Biology*, *218*(12), 1856–1866. https://doi.org/10.1242/jeb.118851
Semmouri, I., Asselman, J., Van Nieuwerburgh, F., Deforce, D., Janssen, C. R., & De
Schamphelaere, K. A. C. (2019). The transcriptome of the marine calanoid copepod
Temora longicornis under heat stress and recovery. *Marine Environmental Research*,
143, 10–23. https://doi.org/10.1016/j.marenvres.2018.10.017

Shakya, M., Lo, C.-C., & Chain, P. S. G. (2019). Advances and challenges in metatranscriptomic analysis. *Frontiers in Genetics*, *10*. https://doi.org/10.3389/fgene.2019.00904

Shaw, J. R., Colbourne, J. K., Davey, J. C., Glaholt, S. P., Hampton, T. H., Chen, C. Y., Folt, C. L., & Hamilton, J. W. (2007). Gene response profiles for Daphnia pulex exposed to the environmental stressor cadmium reveals novel crustacean metallothioneins. *BMC Genomics*, *8*, 477. https://doi.org/10.1186/1471-2164-8-477

Shi, Y., Tyson, G. W., Eppley, J. M., & DeLong, E. F. (2011). Integrated metatranscriptomic and metagenomic analyses of stratified microbial assemblages in the open ocean. *The ISME Journal*, *5*(6), Article 6. https://doi.org/10.1038/ismej.2010.189

Snell, T. W., Brogdon, S. E., & Morgan, M. B. (2003). Gene expression profiling in ecotoxicology. *Ecotoxicology*, *12*(6), 475–483. https://doi.org/10.1023/B:ECTX.0000003033.09923.a8

Stat, M., Huggett, M. J., Bernasconi, R., DiBattista, J. D., Berry, T. E., Newman, S. J., Harvey, E. S., & Bunce, M. (2017). Ecosystem biomonitoring with eDNA: Metabarcoding across the tree of life in a tropical marine environment. *Scientific Reports*, *7*(1), 12240. https://doi.org/10.1038/s41598-017-12501-5

Sunagawa, S., Coelho, L. P., Chaffron, S., Kultima, J. R., Labadie, K., Salazar, G., Djahanschiri, B., Zeller, G., Mende, D. R., Alberti, A., Cornejo-Castillo, F. M., Costea, P. I., Cruaud, C., d'Ovidio, F., Engelen, S., Ferrera, I., Gasol, J. M., Guidi, L., Hildebrand, F., … Bork, P. (2015). Structure and function of the global ocean microbiome. *Science*, *348*(6237), 1261359. https://doi.org/10.1126/science.1261359

Sunday, J. M., Bates, A. E., & Dulvy, N. K. (2011). Global analysis of thermal tolerance and latitude in ectotherms. *Proceedings of the Royal Society B: Biological Sciences*, *278*(1713), 1823–1830. https://doi.org/10.1098/rspb.2010.1295

Suzuki, S., Yamaguchi, H., Nakajima, N., & Kawachi, M. (2018). Raphidocelis subcapitata (=Pseudokirchneriella subcapitata) provides an insight into genome evolution and environmental adaptations in the Sphaeropleales. *Scientific Reports*, *8*(1), 8058. https://doi.org/10.1038/s41598-018-26331-6

Székely, D., Corfixen, N. L., Mørch, L. L., Knudsen, S. W., McCarthy, M. L., Teilmann, J., Heide-Jørgensen, M. P., & Olsen, M. T. (2021). Environmental DNA captures the genetic diversity of bowhead whales (Balaena mysticetus) in West Greenland. *Environmental DNA*, *3*(1), 248–260. https://doi.org/10.1002/edn3.176

Tamames, J., & Puente-Sánchez, F. (2019). Squeezemeta, a highly portable, fully automatic metagenomic analysis pipeline. *Frontiers in Microbiology*, *9*. https://doi.org/doi.org/10.3389/fmicb.2018.03349

Trench-Fiol, S., & Fink, P. (2020). Metatranscriptomics from a small aquatic system: Microeukaryotic community functions through the diurnal cycle. *Frontiers in Microbiology*, *11*. https://doi.org/doi.org/10.3389/fmicb.2020.01006

Tsuri, K., Ikeda, S., Hirohara, T., Shimada, Y., Minamoto, T., & Yamanaka, H. (2021). Messenger RNA typing of environmental RNA (eRNA): A case study on zebrafish tank water with perspectives for the future development of eRNA analysis on aquatic vertebrates. *Environmental DNA*, *3*(1), 14–21. https://doi.org/10.1002/edn3.169

Veilleux, H. D., Misutka, M. D., & Glover, C. N. (2021). Environmental DNA and environmental RNA: Current and prospective applications for biological monitoring. *Science of The Total Environment*, *782*, 146891. https://doi.org/10.1016/j.scitotenv.2021.146891

Veilleux, H. D., Ryu, T., Donelson, J. M., Ravasi, T., & Munday, P. L. (2018). Molecular
response to extreme summer temperatures differs between two genetically differentiated
populations of a coral reef fish. *Frontiers in Marine Science*, 5.
https://doi.org/doi.org/10.3389/fmars.2018.00349
Veilleux, H. D., Ryu, T., Donelson, J. M., van Herwerden, L., Seridi, L., Ghosheh, Y., Berumen,
M. L., Leggat, W., Ravasi, T., & Munday, P. L. (2015). Molecular processes of
transgenerational acclimation to a warming ocean. *Nature Climate Change*, *5*(12), Article
12. https://doi.org/10.1038/nclimate2724
Veldhoen, N., Beckerton, J. E., Mackenzie-Grieve, J., Stevenson, M. R., Truelson, R. L., &
Helbing, C. C. (2014). Development of a non-lethal method for evaluating transcriptomic
endpoints in Arctic grayling (Thymallus arcticus). *Ecotoxicology and Environmental
Safety*, *105*, 43–50. https://doi.org/10.1016/j.ecoenv.2014.03.030
von Ammon, U., Wood, S. A., Laroche, O., Zaiko, A., Lavery, S. D., Inglis, G. J., & Pochon, X.
(2019). Linking Environmental DNA and RNA for Improved Detection of the Marine
Invasive Fanworm Sabella spallanzanii. *Frontiers in Marine Science*, 6.
https://doi.org/10.3389/fmars.2019.00621
Voolstra, C. R., Valenzuela, J. J., Turkarslan, S., Cárdenas, A., Hume, B. C. C., Perna, G.,
Buitrago-López, C., Rowe, K., Orellana, M. V., Baliga, N. S., Paranjape, S., Banc-Prandi,
G., Bellworthy, J., Fine, M., Frias-Torres, S., & Barshis, D. J. (2021). Contrasting heat
stress response patterns of coral holobionts across the Red Sea suggest distinct
mechanisms of thermal tolerance. *Molecular Ecology*, *30*(18), 4466–4480.
https://doi.org/10.1111/mec.16064
Vorobev, A., Dupouy, M., Carradec, Q., Delmont, T. O., Annamalé, A., Wincker, P., & Pelletier,
E. (2020). Transcriptome reconstruction and functional analysis of eukaryotic marine
plankton communities via high-throughput metagenomics and metatranscriptomics.
Genome Research, *30*(4), 647–659. https://doi.org/10.1101/gr.253070.119
Wetterstrand, K. (2021). *DNA Sequencing Costs: Data*. Genome.Gov.
https://www.genome.gov/about-genomics/fact-sheets/DNA-Sequencing-Costs-Data
Wikelski, M., & Cooke, S. J. (2006). Conservation physiology. *Trends in Ecology & Evolution*,
21(1), 38–46. https://doi.org/10.1016/j.tree.2005.10.018
Wood, S. A., Biessy, L., Latchford, J. L., Zaiko, A., von Ammon, U., Audrezet, F., Cristescu, M.
E., & Pochon, X. (2020). Release and degradation of environmental DNA and RNA in a
marine system. *Science of The Total Environment*, *704*, 135314.
https://doi.org/10.1016/j.scitotenv.2019.135314
Yampolsky, L. Y., Zeng, E., Lopez, J., Williams, P. J., Dick, K. B., Colbourne, J. K., & Pfrender,
M. E. (2014). Functional genomics of acclimation and adaptation in response to thermal
stress in Daphnia. *BMC Genomics*, *15*(1), 859. https://doi.org/10.1186/1471-2164-15-859
Yates, M. C., Derry, A. M., & Cristescu, M. E. (2021). Environmental rna: A revolution in
ecological resolution? *Trends in Ecology & Evolution*, *36*(7), 601–609.
https://doi.org/10.1016/j.tree.2021.03.001
Ye, J., Zhang, Y., Cui, H., Liu, J., Wu, Y., Cheng, Y., Xu, H., Huang, X., Li, S., Zhou, A.,
Zhang, X., Bolund, L., Chen, Q., Wang, J., Yang, H., Fang, L., & Shi, C. (2018). WEGO
2.0: A web tool for analyzing and plotting GO annotations, 2018 update. *Nucleic Acids
Research*, *46*(W1), W71–W75. https://doi.org/10.1093/nar/gky400
Ye, Z., Xu, S., Spitze, K., Asselman, J., Jiang, X., Ackerman, M. S., Lopez, J., Harker, B.,
Raborn, R. T., Thomas, W. K., Ramsdell, J., Pfrender, M. E., & Lynch, M. (2017). A new

reference genome assembly for the microcrustacean daphnia pulex. *G3:*
Genes|Genomes|Genetics, *7*(5), 1405–1416. https://doi.org/10.1534/g3.116.038638

Young, T., Walker, S. P., Alfaro, A. C., Fletcher, L. M., Murray, J. S., Lulijwa, R., & Symonds,
J. (2019). Impact of acute handling stress, anaesthesia, and euthanasia on fish plasma
biochemistry: Implications for veterinary screening and metabolomic sampling. *Fish*
Physiology and Biochemistry, *45*(4), 1485–1494. https://doi.org/10.1007/s10695-019-
00669-8

Zhang, Y., Thompson, K. N., Huttenhower, C., & Franzosa, E. A. (2021). Statistical approaches
for differential expression analysis in metatranscriptomics. *Bioinformatics*, *37*, i34–i41.
https://doi.org/10.1093/bioinformatics/btab327

Zhao, S., Zhang, Y., Gamini, R., Zhang, B., & von Schack, D. (2018). Evaluation of two main
RNA-seq approaches for gene quantification in clinical RNA sequencing: PolyA+
selection versus rRNA depletion. *Scientific Reports*, *8*(1), Article 1.
https://doi.org/10.1038/s41598-018-23226-4

Zhao, W., He, X., Hoadley, K. A., Parker, J. S., Hayes, D. N., & Perou, C. M. (2014).
Comparison of RNA-Seq by poly (A) capture, ribosomal RNA depletion, and DNA
microarray for expression profiling. *BMC Genomics*, *15*(1), 419.
https://doi.org/10.1186/1471-2164-15-419

Figures and Tables

Fig. 1. Principal component analysis (PCA) of the top 300 *Daphnia pulex* genes, as ranked by
variance. eRNA refers to extra-organismal RNA released by *D. pulex* into the environment,
whereas oRNA refers to organismal RNA extracted directly from *D. pulex* tissue. Prior to the
PCA, genes were filtered to include only those that had a sum $\geq$ 10 counts in either 20 °C or 28
°C. PCA revealed separation by RNA type and temperature across the PC1 and PC2,
respectively (except eRNA_20T1 clusters more closely with 28 °C samples than with other 20
°C samples).

904
905
906

907
908
Fig. 2. Heatmap showing the relative expression (Z-score calculated for each gene) of all
commonly significantly differentially expressed (DE) *Daphnia pulex* genes (false discovery rate
adjusted p-value <0.05) between 20 °C and 28 °C, in both environmental RNA (eRNA) and
organismal RNA (oRNA). DE analysis was conducted only for those genes that had ≥ 10 counts
in either 20 °C or 28 °C samples. Hierarchical clustering analysis revealed four groups: 20 °C
and 28 °C samples, and up and down regulated genes (except sample eRNA_20T1 clusters with
28 °C samples). Colors indicate levels of relative expression, with blue and red indicating low
and high, respectively. DE statistics and functional description of the genes are provided in Table
S1.
918

919
920
Fig. 3. Bubble plot of the Gene Ontology (GO) enriched terms associated with *Daphnia pulex*
eRNA and oRNA detected genes, and eRNA, oRNA and commonly identified differentially
expressed genes (DEG). The eRNA genes originate from extra-organismal *Daphnia* RNA. We
used a false discovery rate (FDR) corrected p-value <0.05 to identify GO terms as significantly
enriched compared to the genomic background. The bubble color indicates the FDR corrected p-
value for the weight test statistic, while the size is proportional to the fold change between
expected and observed genes annotated with GO term. The biological processes, cellular
components and molecular functions GO domains are represented by BP, CC, and MF,
respectively. Only those GO terms enriched for eRNA genes are shown. Full list of GO terms
enriched for oRNA genes is provided in Fig. S3.
931
932
933
934

Table 1. Problems associated with using environmental transcriptomics based on eRNA for biomonitoring and potential experimental (E), molecular (M), sequencing (S) and bioinformatic (B) solutions.

Problems	Description of problems	Possible solutions	References
Poor understanding of the ecology of eRNA	Little is known about the origin, state, persistence and degradation of eRNA in the environment.	E: Perform experiments that address fundamental questions of the ecology of eRNA (i.e. What are major sources of eRNA? What state is eRNA in? What factors influence eRNA persistence? How do abiotic factors influence eRNA decay rates?)	(Cristescu, 2019)
No understanding of environmental transcriptomics in natural ecosystems	No information available about how our results using environmental transcriptomics will translate to natural ecosystems	E: Perform mesocosm experiments using more complex, multi-trophic communities E: Perform studies using ecosystem wide manipulations E: Perform studies in natural ecosystems	
Rapid degradation of eRNA	eRNA is thought to degrade too rapidly to extract and quantify, but studies indicate that eRNA is detectable for up to 57 hours post organism release	E: Filter water as soon as possible (ideally < 4 hours) E: Freeze samples at -80 °C directly after filtration. Alternatively could potentially use preservative solution if freezing immediately is not possible. M: Keep samples on ice during extraction M: Minimize free-thaw cycles	(T. S. Jo et al., 2022; Kagzi et al., 2022; Marshall et al., 2021; Wood et al., 2020)
Low concentration of eRNA	eRNA samples have much lower RNA concentrations than samples collected from an organisms tissue (i.e. 1 ng/μL vs 200 ng/μL)	E: Increase RNA yield by filtering large volumes of water (> 500 mL); EM: Collect additional technical replicates and pool them during the extraction step; M: Use RNA-seq library prep kits designed for low input starting material	
Low proportion of mRNA	mRNAs are of interest for gene expression analyses as they encode for proteins, but they account for < 5% of total RNA. Non-coding ribosomal and small RNAs account for > 85%	M: Perform rRNA depletion to remove rRNA. However, many commercial kits are optimized to remove human, mouse, rat and bacteria rRNA, and may not deplete rRNA from all species; M: Perform poly(A) enrichment to retain only eukaryotic mRNA. This will eliminate prokaryotic RNA, ribosomal and small RNAs, which make up the majority of total RNA B: If rRNA persists, use it to perform taxonomic identification analysis	(He et al., 2010; Hempel et al., 2022; S. Zhao et al., 2018; W. Zhao et al., 2014)
Low eRNA recovery	eRNA samples are composed of extra-organismal RNA from macro-organisms and whole microorganism. Thus, eRNA reads will be in the minority (estimated <1%), resulting in low power of bioinformatic analyses to detect eRNA based changes in gene expression	E: Pre-filter water with fine mesh to size-selectively remove organisms; E: Use large filter pore size to reduce capture of microorganisms (i.e. > 0.2 μM); E: Sample eRNA from ecosystems that are densely populated with target species M: Perform rRNA depletion on total RNA samples; M: Perform poly(A) enrichment to eliminate prokaryotic RNA; S: Perform deep sequencing to increase eRNA reads (i.e. > 150 million reads per sample)	
Low eRNA read depth	eRNA samples are extremely diverse, resulting in reads from taxa across the trophic chain. Transcript abundance in a sample will influence read depth, and if many micro-organisms are present, eukaryotic genes will have low read depth. Current molecular methods for enrichment and depletion are	M: Create low input poly(A) enrichment methods. Current enrichment methods are designed for organismal RNA from tissue samples and require high concentration of input RNA; M: Increase eukaryotic gene read depth through poly(A) enrichment; MB: Create rRNA depletion methods that remove rRNA from all taxa as current depletion methods are optimized to remove single taxa and bacterial rRNA. If unable to remove all rRNA from eRNA samples, researchers could use rRNA reads for taxonomic information; M: Perform rRNA depletion;	(Yates et al., 2021)

	optimized for organismal, not diverse environmental samples.	M: Amplify biomarker genes via targeted (qPCR, ddPCR) approaches to analyze specific biomarker genes	
Difficulty characterizing read function due to lack of eukaryotic reference genomes	Reference genomes are used to functionally and taxonomically characterize reads. eRNA samples will capture eRNA from eukaryotes across the trophic chain, but genomes are available for only 0.4% of eukaryotic species.	ES: Sequence all eukaryotic species (e.g. Earth BioGenome Project) B: *De novo* assembly of reads into longer scaffolds, which enables the identification of homologs.	(Lewin et al., 2018; Shakya et al., 2019)
Lack of baseline in natural ecosystems to compare against	Differential gene expression analyses are conducted by comparing treatment samples (i.e. stressed community) against a control 'baseline' (i.e. health community). In natural ecosystems it is difficult or often not possible to obtain a baseline control as many stressors act simultaneously on a system	E: Conduct time series eRNA sampling of the same environment and compare time points; E: Compare environments with similar abiotic conditions. If healthy, one could serve as a baseline to compare against; EM: Couple eRNA with targeted PCR approaches (qRT-PCR, ddPCR) and biomarker genes to infer physiological states of specific species; EB: Reconstruct mock communities under neutral conditions as a baseline; B: The cellular stress response is triggered by general stress and as such genes and pathways involved could be broadly indicative of community stress; B: Use housekeeping genes with consistent expression levels as baselines to compare against; B: Supervised machine learning models could be trained with existing metatranscriptomic, transcriptomic and metagenomic data that have identified functional terms, metabolic pathways and taxa associated with stressors. Environmental transcriptomic data could be used as input and the model would predict the stressors involved and health of the community	(Cordier et al., 2019; Fulda et al., 2010; Mallick et al., 2021; Yates et al., 2021; Zhang et al., 2021)
False detection of differential transcript expression due to changes in population densities	Between environments, species across the trophic chain exhibit differences in abundances. RNA transcript abundance varies with DNA abundance. Thus, changes in RNA transcript abundance, independent of differential expression, could be recovered.	EB: Collect and sequence eDNA in parallel. eDNA can acts as a species abundance estimator, independent of transcript abundance, that we can normalize eRNA reads to. Allows us to track whether the proportion of eRNA reads follows species abundance, or is up or down regulated; B: 'Taxon-specific scaling' accounts for variation in abundances by separating and then normalizing metatranscriptomic data into organism-specific bins. A differential expression analysis could then be performed after the normalized matrices are recombined into community wide data	(Klingenberg & Meinicke, 2017; Salazar et al., 2019; Yates et al., 2021; Zhang et al., 2021)