Adhesion of *Crithidia fasciculata* promotes a rapid change in developmental fate driven

by cAMP signaling

Shane Denecke[1,§], Madeline F. Malfara[2,§], Kelly R. Hodges[2], Nikki A. Holmes[2], Andre R. Williams[2], Julia H.

Gallagher-Teske[3,4], Julia M. Pascarella[3], Abigail M. Daniels[1], Geert Jan Sterk[5], Rob Leurs[5], Gordon Ruthel[1],

Rachel Hoang[3], Megan L. Povelones[2,*], Michael Povelones[1,*]

[1]Department of Pathobiology, School of Veterinary Medicine, University of Pennsylvania, Philadelphia, PA

19104, USA

[2]Department of Biology, Villanova University, Villanova, PA 19085, USA

[3]Department of Biology, Haverford College, Haverford, PA 19041, USA

[4]Cancer Clinical Trials Office, Department of Urology, Stanford University, Palo Alto, CA 94304, USA

[5]Amsterdam Institute for Molecules, Medicines and Systems, Division of Medicinal Chemistry, Faculty of

Science, Vrije Universiteit Amsterdam, De Boelelaan 1108, 1081 HZ, Amsterdam, The Netherlands

Running title: Adhesion promotes rapid developmental change

§ Authors contributed equally

* To whom correspondence should be addressed: Michael Povelones: Department of Pathobiology, School of

Veterinary Medicine, University of Pennsylvania, Philadelphia, PA, 19104; Megan Povelones: Department of

Biology, Villanova University, Villanova, PA, 19085

mpove@vet.upenn.edu; Tel (215) 898-4144

megan.povelones@villanova.edu; Tel (610) 519-4863

Keywords: *Crithidia fasciculata*, adhesion, cyclic AMP, signaling, differentiation, kinetoplastid

Abstract

Kinetoplastids are single-celled parasites responsible for human and animal disease. For the vast majority of kinetoplastids, colonization of an insect host is required for transmission. Stable attachment to insect tissues via the single flagellum coincides with differentiation and morphological changes. Although this process is essential for the generation of infectious forms, the molecular mechanism driving differentiation following adherence is not well understood. To study this process, we elaborate upon an in vitro model in which flagellated swimming cells of the kinetoplastid *Crithidia fasciculata* rapidly differentiate following adhesion to artificial substrates. Manipulation of culture parameters revealed that growth phase and time had a strong influence on the proportion of adherent cells. Live imaging of cells transitioning from swimming to an attached cell fate show parasites undergoing a defined sequence of events including an initial adhesion near the base of the flagellum, immediately followed by flagellar shortening, cell rounding, and the formation of a hemidesmosome-like structure between the tip of the shortened flagellum and the substrate. We have also assayed the role of the cyclic AMP (cAMP) signaling pathway in differentiation of *C. fasciculata*. Pharmacological inhibition of cAMP phosphodiesterases eliminated the ability of swimming cells to attach without affecting their growth rate. Further, treatment with inhibitor did not affect the growth rate of established attached cells, indicating its effect is limited to a critical window of time during the early stages of adhesion. Finally, in swimming parasites we have shown that a receptor adenylate cyclase localizes to the distal portion of the flagellum. In attached cells it is absent from the shortened flagellum and instead localizes to the cell body. Similarly, a putative phosphodiesterase, found along the length of the flagellum, also relocalizes to the cell body in attached parasites. These data suggest that in *C. fasciculata* cAMP signaling is required for adherence, that cAMP flux in the flagellum of swimming cells is spatially restricted, and that signaling domains may be reorganized during differentiation and attachment. These studies contribute to our understanding of the flagellum as a multi-functional organelle integrating processes related to motility, signaling, attachment, and differentiation and further develop *C. fasciculata* as a model kinetoplastid.

Author Summary

Parasites from the order Kinetoplastida are transmitted by insects and cause diseases such as Leishmaniasis,

Chagas disease, and Human African Trypanosomiasis. A key aspect of the life cycle of these parasites is their

ability to adhere to surfaces within the insect and subsequently differentiate into forms that are transmissable to

their next host. Here, we explore the molecular mechanisms underpinning both adherence and differentiation

using the mosquito parasite *Crithidia fasciculata*. This organism is closely related to pathogenic species but grows

to high densities in culture and adheres robustly in vitro. We identified factors affecting the rate of adherence and

defined morphological stages of this process including rapid shortening of the cell's single flagellum. Using

electron microscopy, we confirmed that adhesion to artificial substrates results in the formation of a filament-rich

adhesive plaque at the tip of the flagellum that resembles the attachment to insect tissue. We also demonstrate the

key role played by the cyclic AMP signaling pathway during adherence and differentiation. Specifically,

pharmacological inhibition of enzymes that degrade cyclic AMP completely blocks adherence, and tagging of

several proteins in this pathway show that their localization along the flagellum changes following differentiation

to the attached form. These data provide insights into processes critical for all kinetoplastid life cycles.

Introduction

Organisms from the protozoan order Kinetoplastida impose a significant disease burden especially in low-income countries [1]. *Trypanosoma brucei*, *Trypanosoma cruzi*, and *Leishmania* spp. are the etiological agents for African trypanosomiasis, Chagas disease, and Leishmaniasis, respectively. These parasites also cause important veterinary diseases, while still others impact insects critical for pollination. A deeper understanding of the biology of these organisms can provide the basis for rational interventions to limit disease burden. Additionally, kinetoplastid parasites are a compelling model for basic research, representing early-branching eukaryotes that are divergent from more well-studied organisms such as insects, worms, and vertebrates.

Kinetoplastids primarily parasitize insects with only a handful of species using insects as a vector to infect other organisms. In insects, kinetoplastids spend part of their life cycle adhered to the apical surface of an epithelial lining. Adherence, also referred to as adhesion or attachment in the literature, often precedes a developmental transition with distinct morphological and molecular changes. For example, in the kissing bug *Rhodnius prolixus*, *T. cruzi* parasites bind to the perimicrovillar membrane in the posterior midgut, followed by the cuticular lining of the hindgut, before being excreted by the insect as infectious metacyclics [2, 3]. Adherence is necessary not only to maintain the parasites in the gut of their host, but to trigger differentiation to the infectious form. Similarly, *T. brucei* adheres to salivary gland epithelium and *Leishmania* spp. adhere to the midgut and the stomodeal valve. In each case, adherence precedes or is concurrent with differentiation to the final developmental form in the insect. Despite the importance of adherence in the kinetoplastid insect life cycle, the relative inaccessibility of these parasite stages in the insect, and the lack of a robust in vitro model to generate them, means that little is known about adherence mechanisms and the signal transduction pathways that may connect adherence to developmental progression.

Crithidia fasciculata is a monoxenous parasite of mosquitoes that developmentally transitions from a swimming (nectomonad) cell with a long flagellum to an attached (haptomonad) cell with a dramatically shortened flagellum. The tip of this flagellum is attached to the chitinous lining of the hindgut epithelium via a hemidesmosome-like adhesive plaque. A morphologically similar plaque is found in the adhered forms of other kinetoplastid species [4]. Remarkably, for *C. fasciculata* this developmental switch from a swimming to an attached cell is readily produced in vitro by adhesion to a variety of substrates, including Millipore filters [5],

culture debris [6], and plastic dishes [7, 8]. Taking advantage of the ability to produce large numbers of

swimming and attached cells in culture, our recent transcriptomic analysis identified genes encoding putative

cyclic AMP (cAMP) signaling components that were differentially regulated in the different forms of *C.*

fasciculata. This led to the hypothesis that cAMP signaling may be involved in cell fate transitions in these

organisms [7].

Despite its importance, much remains unknown about cAMP signaling in kinetoplastids [9]. The cAMP second

messenger is created from ATP by adenylate cyclases (ACs), many of which are transmembrane receptor

adenylate cyclases (RACs) that probably act as receptors for unidentified ligands. The RAC family of ACs is

dramatically expanded in *T. brucei*, in which they localize to the flagellum, either along the length or at the distal

tip [10]. In many systems protein kinase A (PKA) acts downstream of cAMP, but in kinetoplastids, [11, 12], PKA

is independent cAMP [13]. Instead, several cAMP response proteins (CARPs) are thought to mediate

downstream events. Phosphodiesterases (PDEs) terminate the signal by degrading cAMP to 5'-AMP. In *T. brucei*,

some of these PDEs are localized to the flagellum, while others are found in the cell body [14]. It has been

proposed that PDEs act to limit the diffusion of cAMP and create spatially restricted signaling microdomains [15].

This paradigm has also been demonstrated in other eukaryotic systems [16].

Using a robust and quantitative adherence assay and imaging of live and fixed cells, we have defined the process

of adherence and the conditions that promote it in vitro. In addition, we have used pharmacological inhibitors to

probe the role of cAMP signaling during the transition to an attached state. Finally, we show the localization of

putative components of the *C. fasciculata* cAMP signaling pathway to the flagellum in swimming cells and show

striking changes in the distribution of these proteins upon flagellar shortening. These findings support the use of

C. fasciculata as a model to understand the spatial regulation of signal transduction in the context of adherence

and differentiation of kinetoplastid parasites.

Results

Culture conditions impact in vitro adherence of *C. fasciculata*

To probe the mechanism of *C. fasciculata* adherence, we used an in vitro assay to determine which conditions

impact the adherence rate. Swimming cells, grown on a shaker, were transferred to fresh culture plates and

incubated without shaking to allow cells to adhere. Plates were washed to remove non-adherent cells and then the

number of single adhered cells in representative fields were quantitated (Fig 1A). Over the next 24 hours, these

single cells divide to form attached rosettes. Cells in a rosette can undergo division to produce either two attached

cells or two swimming cells that can leave the rosette. Some proportion of these newly-produced swimming cells

can also encounter the dish and become attached. In our standard assay, swimming cells were again washed away

at 24 hours and rosettes with four or more cells in representative fields were quantitated.

By varying the length of plating time prior to washing, we found that the length of this incubation step was

directly proportional to the number of attached parasites, with the longest time we tested, 240 minutes, producing

the greatest number of rosettes (Fig 1 B,C). Consistent with a previous report [17], we also found that the growth

phase of the cultured cells had a strong effect on the rate of adherence. Cells from an early log phase culture (10^7

cells/mL) adhered and differentiated at a higher frequency compared with the same number of cells taken from a

stationary phase culture (10^8 cells/mL; Fig 1D). In addition to having a reduced rate of cell division, stationary

phase cells are morphologically distinct from cells in exponentially growing cultures, with the latter being rounder

and shorter (Figure S1). Increasing the total number of cells used in the adherence assay, provided those cells

were taken from a culture in log-phase, proportionally increased the total number of attached cells but had

relatively little impact on the percentage of cells that adhered during the 2 hour incubation (Fig 1E). These data

suggest that in vitro adherence of *C. fasciculata* happens at a constant rate and is inversely correlated with the

culture density before the assay begins but not while adherence occurs.

Mechanism of attachment of *C. fasciculata* in vitro

To examine the stages of adherence and differentiation of *C. fasciculata* more precisely, we used time-lapse

microscopy to follow parasites in environmentally controlled conditions. Using flagellar shortening and cell

rounding as morphological markers for differentiation, we observed that these events were always preceded by an

initial adhesion event (Videos S1,S2). During this event, cells first adhere to the plate via their anterior end, near

the flagellar pocket, and then begin to retract their flagellum. As this happens, the cells become rounder as they

undergo their first cellular division as adherent cells. In certain cases, cell division occurs prior to the full

retraction of the flagellum, but this was rare. Parasites sometimes adhered in pairs, a phenomenon that has been

reported previously [18]. In this case, one cell of the pair will typically detach and swim away (Video S2). From

these experiments, it was also possible to directly measure the rate of flagellar shortening, revealing that the

flagellum retracts at a relatively constant rate until it is no longer visible at around 2 hours (Fig 2).

While parasites initially adhere near the base of their flagellum, stably attached cells are connected to the substrate

via the distal tip of the shortened flagellum. Previous reports on *C. fasciculata* and other kinetoplastids have

described plaques at the site of attachment that ultrastructurally resemble hemidesmosomes [6]. Since most

imaging of the adhesive structure was performed on parasites adhered to the insect, we used transmission electron

microscopy (TEM) to determine if hemidesmosomes were present in attached cells where they contact the culture

plate. Consistent with prior findings [8], a cross-section of rosettes showed an osmophilic, filamentous plaque at

the distal tip of the shortened flagellum adjacent to the substrate (Fig 3 A-B). When we detached adherent cells

from culture plates by scraping prior to fixation, the hemidesmosome appeared intact, indicating that attachment

mediated by this structure can be physically disrupted (Fig 3C). Fibrous, desmosome-like structures appear to

connect the proximal part of the flagellum to the cell body at the flagellar pocket, features which have also been

noted in prior studies and which are probably components of the flagellar attachment zone (FAZ) (Fig 3A-C) [6,

19, 20]. When cells were sectioned *en face* near the attachment site with the culture plate, we observed that the

distal portion of the flagellum containing the hemidesmosome was somewhat enlarged and extended beyond the

termination point of flagellar axoneme (Fig 3D). In these sections, hemidesmosomes are circular and the degree to

which they are stained by uranyl acetate is likely a function of how close to the culture plate they were sectioned

(Fig 3D,E). By contrast, in swimming cells, we only found FAZ-like structures connecting the flagellum to the

pocket but did not observe hemidesmosomes at the distal end of the flagellum (Fig 3F). A notable feature of both

attached cells and scraped attached cells was an extracellular network of filamentous material attached to the

plasma membrane of the cell body (Fig 3G and 3A-D). In swimming cells, similar material was sometimes

observed in the flagellar pocket itself, but not on the cell surface (Fig 3F).

Pharmacological Inhibition of PDEs blocks adherence without affecting growth

Our prior work showed that cAMP signaling components are upregulated in swimming *C. fasciculata* relative to

attached parasites [7]. To test whether manipulating the cAMP pathway could influence the ability of *C.*

fasciculata to adhere to plastic in vitro*,* we used previously validated compounds that are known to inhibit cAMP

PDEs in the related parasite, *T. brucei* [21]. As PDEs terminate the signal by degrading the cAMP second

messenger, inhibition of these enzymes has been shown to increase the amount of cAMP in the parasite [22]. The

first compound we tested is known as NPD-001 or Compound A (CpdA) (hereafter NPD-001). The addition of 10

µM of NPD-001 to the adherence assay completely blocked parasite attachment to the plate, resulting in no single

adhered cells after 2 hours of incubation and no visible rosettes after 24 hours of growth (Fig 4A). Interestingly, a

related compound, NPD-008, had only a minor effect on adhesion, slightly reducing the number of single adhered

cells after 2 hours of incubation (Fig 4A). To determine if the effect of NPD-001 on attachment was dose-

dependent, we performed an adherence assay with both 1 µM and 0.1 µM of the compound. Although the effect

of 1 µM was comparable to that of 10 µM, 0.1 µM had a moderate effect on adhesion, suggesting a dose response.

The same concentrations of a related compound, NPD-226, also significantly reduced the number of adhered cells

both at 2 hours and at 24 hours at all three concentrations compared to vehicle-treated controls (Fig 4B). A fourth

compound, NPD-055, had a minor effect on adherence at the highest concentration (10 µM, Fig S3).

To confirm that the adherence-blocking activity of the PDE inhibitors was not due to cell toxicity, we measured

growth rates of swimming cultures of *C. fasciculata* (maintained on a shaker) in the presence of NPD-001.

Addition of this compound to the media did not affect the growth rate of swimming cells compared to vehicle- or

untreated control cells (Fig 5A). Furthermore, addition of NPD-001 to cells that were already differentiated to the

attached state did not affect their attachment or their doubling time over the course of 16 hours (Fig 5B). In this

experiment, we did observe new adherence of swimming cells produced from the rosettes in the vehicle-treated

control. Consistent with our previous finding (Fig 4A), although new swimming cells were produced from the

rosettes, in NPD-001 treated cultures, they failed to adhere.

cAMP signaling proteins are localized to the flagellum

In other kinetoplastids, components of cAMP signaling pathways are localized to flagellar subdomains, indicating

that this organelle may be involved in spatially restricting the distribution of this second messenger [10, 14]. To

see if this feature is also present in *C. fasciculata*, we localized a putative receptor adenylate cyclase (*Cf*RAC1)

and a putative cAMP phosphodiesterase (*Cf*PDEA) by fusing a GFP protein to the C-terminus of each open

reading frame and introducing these constructs into parasites as episomes (Fig S4). In swimming *C. fasciculata*,

*Cf*PDEA::GFP was found along the length of the flagellum, often appearing as two parallel lines of puncta (Fig

6). The extent of *Cf*PDEA::GFP is somewhat diminished along the portion of the flagellum in the flagellar pocket.

*Cf*RAC1::GFP was found in similar puncta, but was restricted to the distal third of the flagellum in swimming

parasites (Fig 6, Fig S5). We compared both of these localization patterns to another cell line in which a YFP

sequence was fused to one of the endogenous copies of the *Cf*PF16 gene (Fig S4), a component of the central pair

of the flagellar axoneme [23]. *Cf*PF16::YFP was uniformly distributed along the length of the flagellum in a

single continuous line without obvious puncta. No fluorescent signal was detected in parental CfC1 cells (Fig S6).

We then followed these localization patterns as cells transitioned from the swimming to the attached form. As

expected, *Cf*PF16::YFP could still be detected in the shortened flagellum which, as supported by our EM studies,

retains the central pair. However, *Cf*RAC1::GFP was no longer localized to the flagellum and was instead

distributed across the entire cell body. A similar redistribution was also observed in attached cells expressing

*Cf*PDEA::GFP (Fig 6). To compare expression levels of these proteins in attached and swimming *C. fasciculata*,

we performed western blotting using an anti-GFP antibody. Consistent with our microscopy results,

*Cf*PDEA::GFP and *Cf*RAC1::GFP are readily detected in both swimming and attached parasites despite their

different subcellular localizations (Fig 6). No fluorescent or anti-GFP western signal was observed in the parental

cells (Fig S6).

Discussion

Adherence and differentiation are both critical for the successful completion of kinetoplastid life cycles. While

signal transduction pathways driving differentiation have been described in multiple species, virtually nothing is

known about mechanisms of attachment and how this process is related to critical developmental events in the

insect vector. Numerous studies have found important roles for cAMP-based signaling in differentiation and

response to host environments, yet many questions remain unanswered in this area as well, including why

kinetoplastids have expanded their repertoire of RACs, what signals are sensed, and how downstream signals are

transduced. In addition, the significance of signaling subdomains in the flagellum is not understood. *C. fasciculata*

provides an opportunity to dissect the relationship between flagellar signaling and adherence in the context of

differentiation. We expect that, with some variations, this connection exists in all kinetoplastids. Our ability to

follow adherence and differentiation in live cells, coupled with a quantitative assay for adherence and expanding

tools for molecular genetics, opens a wide range of avenues for future exploration.

The mechanism of *C. fasciculata* attachment in vitro

The adherence of *C. fasciculata* and other kinetoplastids to substrates in vitro has been noted previously [6, 8, 17,

24-27]. Having developed a reproducible, quantitative assay for this phenomenon, we decided to explore factors

that would increase or decrease adherence. Consistent with previous work, we found that adherence of *C.*

fasciculata to artificial surfaces results in the formation of a hemidesmosome-like structure at the point of

attachment at the tip of the shortened flagellum. Due to the similarity of this attachment structure to that formed in

vivo [28, 29], we can use the in vitro system to determine the conditions required for its formation. As was

reported previously [17], we have also found that cultures in stationary phase are less likely to adhere compared to

log phase cultures, with a large drop off in adherence for parasites growing between 4 to $8x10^7$ cells/mL. *C.*

fasciculata parasites in stationary phase were shown to have a more elongated cell morphology compared to log

phase cells, and previous studies have demonstrated differential protein abundance between these cell states

possibly linked to their nutritional status [30]. Such nutrient deprivation has been shown to trigger adherence of *T.*

cruzi epimastigotes after which the parasites differentiate to the infectious metacyclic form [31]. In our assays,

this effect seems to result from an inherent difference between parasites in log- versus stationary-phase, as plates

seeded with large numbers of log phase cells can form nearly confluent layers, suggesting that there is no upper

limit to density in this assay. As adherence is linearly related to the number of plated log-phase cells, we also

conclude that it is not cooperative, and that each parasite adheres independently to the substrate. This is supported

by our TEM, live-cell, and fluorescence imaging studies which confirm that rosettes are formed by cell division of

individually adhered cells, which are each attached to the surface through their flagellum. In this case, the number

of attached parasites should be limited by substrate surface area. Consistently, previous studies on *Leishmania*

spp. have noted that scratching the surface, thereby increasing the available surface area, increases parasite

adherence [27], although this did not appear to trigger differentiation. It should be noted that the term "rosette"

has also been used to describe an observed tendency of some *Leishmania* spp. parasites to adhere to each other via

their flagella [32-35]. These rosettes are not adhered and are not formed through cell division, but instead result

from clustering of parasites. Thus far we have not seen this phenomenon in our *C. fasciculata* cultures.

We have also used this assay to probe the mechanism of adherence. *C. fasciculata* parasites initially contact the

dish via the anterior end of the cell body. We call this initial event "adhesion" which is rapidly followed by

flagellar shortening, cell rounding, and formation of the hemidesmosome. We call this fully differentiated state

"attached" reflecting the stable attachment between the tip of the flagellum and the substrate presumably

facilitated by the hemidesmosome-like structure and accompanied by changes in gene expression. In this study,

we have found that even at log phase the percentage of *C. fasciculata* parasites that stably attach to the dish is low

and increases linearly with time. Furthermore, our live-cell imaging studies show that not every cell that

encounters the plate undergoes differentiation to the attached form. These aborted adhesion events suggest that

only a subset of cells can make the transition at any one time, which may reflect stochastic variation, timing

within the cell cycle, or other factors. For example, it was shown previously that a subset of stationary-phase *C.*

fasciculata fail to agglutinate after treatment with peanut lectin, indicating possible variations in cell surface

molecules [30]. These glycoproteins and glycoconjugates may connect to the membrane by

glycosylphosphatidylinositol (GPI) anchors, which are found commonly among kinetoplastids. For example, it

has been suggested that differences in cell surface lipophosphoglycan (LPG) between promastigotes and

metacyclics in *L. major* may underpin the ability of the former to adhere to the midgut of their sandfly vector [36,

37]. These findings are particularly interesting in the context of the TEM data reported here, which show a lattice-

like filamentous molecule at the cell surface only in adherent cells. Similar material was detected in the flagellar

pocket of swimming cells, suggesting that it may be synthesized and secreted by this form, but not retained on the

surface. Determining the identity of this molecule will allow for exploration of possible roles in adhesion or

protection in the insect gut, as well as comparisons with other kinetoplastid species.

The predictable regularity of *C. fasciculata* attachment suggests a developmental program resulting in a distinct

cell state. Flagellar shortening occurs at a constant rate and seems triggered by the initial adhesion event. Our

TEM images show that the shortened flagellum of attached *C. fasciculata* are distinct from that of *Leishmania*

amastigotes in that the *C. fasciculata* axoneme retains the central pair of microtubules even at the tip. In addition

to the formation of hemidesmosome filaments, the *C. fasciculata* flagellar membrane at the point of attachment is

also expanded, as has been noted for haptomonads of other species [24, 26, 38]. From these data and our

experiments with NPD-001/CpdA, it appears that the final attached state, once established, is relatively stable. As

in other studies, we have found that the hemidesmosome is extremely resistant to detergents and enzymatic

digestion [39] but can be physically disrupted by scraping [40].

The role of cyclic AMP in *C. fasciculata* adhesion and differentiation

Strikingly, we found that inhibition of PDEs with NPD-001 completely blocked attachment. As treatment of

stably attached *C. fasciculata* parasites with NPD-001 did not cause the cells to detach nor did it prevent division

of attached cells, we hypothesize that elevated cAMP resulting from PDE inhibition disrupts the dynamic

transition from initial adhesion to stably attached. An alternative interpretation is that elevated cAMP impacts the

swimming cell population by dramatically reducing the subset of cells that are competent to adhere, meaning that

these cells would not even form initial adhesions. Our findings contribute to a large body of work showing that

signaling via the cAMP pathway plays a critical role in developmental transitions in kinetoplastid parasites [9,

41], including the epimastigote to metacyclic transition in *T. cruzi* [42] and the transitions to stumpy [42] and

procyclic form *T. brucei* [43]. However, in *T. brucei* there is evidence that differentiation may be triggered by

degradation products of cAMP, rather than cAMP itself [44]. In addition to differentiation, *T. brucei* requires

cAMP signaling for social motility, a process by which parasites generate and sense pH gradients and migrate

collectively through the tissues of their insect host, eventually attaching to the salivary gland epithelium and

differentiating to infectious metacyclics [12, 45, 46]. Treatment with CpdA/NPD-001 blocks social motility in

vitro [45], and genetic knockout of the *T. brucei* PDEB1 gene or other signaling components impairs or blocks

developmental progression through the fly [12, 46]. Due to the impact on social motility, the effect of these

manipulations on *T. brucei* attachment cannot be readily assessed.

Variations in flagellar length are a common feature of kinetoplastid life cycles [47], which may impact the

distribution of flagellar cAMP signaling proteins. We have found that GFP fusions of a putative RAC and PDE

(*Cf*RAC1::GFP and *Cf*PDEA::GFP) both localize to the flagellum in swimming *C. fasciculata*, with

*Cf*RAC1::GFP being enriched in the distal tip. Both *Cf*PDEA::GFP and *Cf*RAC1::GFP display punctate

localization patterns compared to *Cf*PF16::YFP. While *Cf*RAC1 has a predicted transmembrane domain, *Cf*PDEA

does not. However, in *T. brucei*, the phosphodiesterase TbPDEB1 is an integral component of the flagellum that is

resistant to detergent extraction [15]. Lipid analysis of the flagellar membrane has suggested lipid-raft like

membrane heterogeneity, which could also lead to uneven localization of our tagged proteins [48]. Therefore,

anchoring of *Cf*RAC1 and *Cf*PDEA to the flagellar membrane or structures within the flagellum could explain the

puncta that we observe. In addition, the parallel lines of signal found in both tagged cell lines is reminiscent of

intraflagellar transport trains, which associate with two sets of microtubule doublets on either side of the axoneme

[49]. The different distributions of these flagellar proteins, *Cf*RAC1 to the tip and *Cf*PDEA to the entire length, is

also consistent with observations in other organisms, in which specific amino acid sequences are required for

correct localization to flagellar subdomains [10]. However, how these domains are established and the

mechanisms for protein targeting are not yet clear.

Both tagged proteins are no longer enriched in the shortened flagellum of attached parasites, indicating

differential localization in the two developmental forms of *C. fasciculata*. As expected, *Cf*PF16::YFP was

retained in the shortened flagellum, as this structure still has its central pair of microtubules. It is unclear from

these studies if resident signaling proteins are relocalized from the flagellum to the cell body during transition

from swimming to attached, or whether newly synthesized signaling proteins in attached cells fail to enter the

flagellum, perhaps due to alterations in the machinery for flagellar protein sorting. Fluorescence time-course

imaging of differentiating cells should help reveal the dynamic localization patterns of cAMP signaling

components, which would convey critical information about this process.

Our data suggest that cAMP signaling acts during a relatively narrow window to facilitate the transition between

initial adhesion and stable attachment. As illustrated in Figure 7, our model is that the physical contact with the

surface during adhesion initiates flagellar shortening leading to local alterations in cAMP flux which then trigger

a developmental program including changes in transcript abundance and formation of the attachment plaque. This

would mean that the spatial restriction of RACs, PDEs, and other signaling proteins conveys important

statiotemporal information that is vital for successful developmental transitions. *C. fasciculata* is an attractive

system in which to tease out these processes, and future studies will explore the connection between flagellar

dynamics, signaling, adherence, and differentiation in this important group of parasites.

Materials and Methods

Crithidia maintenance

The Cf-C1 strain of *C. fasciculata* was used for all experiments in this study. This strain was originally obtained

from Dr. Stephen Beverley and was grown in complete medium (BHI) consisting of Brain Heart Infusion medium

(Sigma) supplemented with 20 µg/mL hemin (Sigma). Cells were maintained at 28 ºC in non-vented culture flasks

on a rocker and were maintained at a density between 10^5 and 10^8 cells/mL. Swimming cells were counted on a

hemocytometer following fixation in 0.3% formalin.

Plasmids and cell lines

The construct for C-terminal YFP tagging of the *C. fasciculata* putative axoneme central apparatus protein

CFAC1_170028000, named *Cf*PF16 for its shared homology to *L. mexicana* PF16 [23]*,* was created using a

fusion PCR approach described previously [50]. The YFP and neomycin resistance genes were amplified from

pLENTv2eYFPNeo by PCR with forward primer pLENTeYFP_f and reverse primer pLENTNeo_r (Table 1). 500

bp of homology to the C-terminus of *Cf*PF16 was amplified from *C. fasciculata* genomic DNA using forward

primer PF16cterm500_f and reverse primer PF16cterm_r. 500 bp of homology to the 3' UTR of *Cf*PF16 was

amplified using PF16utr_f and PF16utr500_r. All three fragments were purified by PCR clean-up or gel

extraction and 20 ng of each fragment was combined in a fusion PCR reaction using PF16_500bpnest_f and

PF16_500bpnest_r. The resulting construct, PF16eYFPNeo, was gel extracted and ethanol precipitated. To create

a *C. fasciculata* CfPF16::YFP cell line, transfection was performed with 3.5 µg of PF16eYFPNeo introduced into

10^8 cells using a Human T-cell kit and IIb Nucleofector set to program X-001 (Lonza). Other transfections

described in this work used the same conditions except with *Tb*-BSF buffer (90 mM sodium phosphate, 5 mM

potassium chloride, 0.15 mM calcium chloride, 50 mM HEPES, pH 7.3) [51]. Immediately following transfection,

cells were transferred to 30 mL of BHI without selecting drug and diluted 10-fold. For each dilution, 1 mL was

plated into wells of a 24-well plate and cultured overnight without shaking. After approximately 18 hours of

recovery, 1 mL of BHI containing neomycin (G418) was added to each well, for a final selecting drug

concentration of 25 µg/mL. Resistant clones came up in 7-14 days and were fixed and screened for eYFP

fluorescence using wide-field microscopy. The resulting clonal cell line CfPF16YFP_A6 was used for these

studies and was validated by linking PCR across the gene with primer PF16ORF_f binding inside the gene and

reverse primer PF16UTR_r binding in the 3'UTR (outside of the area used for integration). To create the *C.*

*fasciculata Cf*RAC1::GFP and CfPDEA::GFP cell lines, the entire open reading frame of *Cf*RAC1

(CFAC1_090006500) or CfPDEA (CFAC1_140020300) were amplified from *C. fasciculata* genomic DNA and

cloned into the pNUSGFPcH vector [52] using NdeI and KpnI or NdeI and BglII restriction enzyme sites,

respectively. Following confirmation by sequencing (Eurofins Genomics), 10 µg of the final plasmid was

introduced into *C. fasciculata* as described above. Transfected cells were selected using BHI containing 80 µg/mL

hygromycin. After 7 days, selecting drug concentration was increased to 200 µg/mL and cells were screened by

fluorescence microscopy.

Adherence assay

Establishment of cultures of swimming and adherent *C. fasciculata* in vitro were adapted from our previous study

[7]. Briefly, cells from a rocking culture of swimmers were added to 60 mm non-treated tissue culture plates

(VWR) in 2 mL of BHI. For some experiments, cells were spun down in tubes at 800 rcf for 5 minutes and

resuspended in BHI. The cell suspension was allowed to sit in the 60 mm plates undisturbed at 28 ºC for the

indicated amounts of time during which adherence takes place (2 hours for the standard assay). Following this

incubation, each plate was washed 5 times with PBS before addition of fresh BHI media. When assays were

performed in 24-well plates, the same protocol was followed but the wash volume was 2 mL. Other assay

parameters were tested systematically and described in the results.

Quantitation of adherence was performed through imaging on a Zeiss Primovert Digital Microscope. Following

wash steps, plates were imaged by taking either 25 frames at 40x magnification or 10-25 frames at 20x

magnification. Freshly adhered cells (singlets) were counted either manually or, when large numbers of adhered

cells were present, by Fiji script [23]. Rosettes were counted manually as clusters of 4 or more cells at 24 hours.

Transmission electron microscopy and Giemsa staining

Swimming and attached *C. fasciculata* cells were processed and imaged using transmission electron microscopy

(TEM) performed in the Electron Microscopy Resource Lab (Perelman School of Medicine, University of

Pennsylvania). For swimming cells, 10^8 mid-log-phase cells were concentrated by centrifugation and resuspended

in TEM fix (0.1 M sodium cacodylate buffer, pH 7.4, containing 2.5% glutaraldehyde, and 2.0%

paraformaldehyde). For attached cells, rosette cultures in 60 mm dishes were established for 24 hours as described

above. 5 mL of TEM fix was added to the plate at room temperature. A second 24-hour attached sample was

generated by scraping cells off the plate with a disposable plastic scraper, concentrating by centrifugation, and

placing in TEM fix. All samples were fixed initially at room temperature for 2 hours and then placed at 4 ºC

overnight. After subsequent buffer washes, samples were post-fixed in 2.0% osmium tetroxide with 1.5%

$K_3Fe(CN)_6$ for 1 hour at room temperature and rinsed in water prior to *en bloc* staining with 2% uranyl acetate.

After dehydration through a graded ethanol series, the tissue was infiltrated and embedded in EMbed-812

(Electron Microscopy Sciences). Thin sections were stained with uranyl acetate and SATO lead and examined

with a JEOL 1010 electron microscope fitted with a Hamamatsu digital camera and AMT Advantage

NanoSprint500 software. Images of cells cut in cross section and *en face* were captured (Fig S2). Morphological

changes between mid-log phase and stationary phase cells were obtained by performing a cytologic cytocentrifuge

preparation and a standard Wright-Giemsa stain and imaging at 100x.

High content imaging

High content imaging was performed using an ImageXpress Micro 4-High Content imaging device (Molecular

Devices) at 40x and 60x using standard 24 (CellTreat) and black welled (Vision 4tituder) 24 well plates

respectively. To measure rosette growth, cells were allowed to adhere to standard plates for 2 hours followed by 5

washes with 2 mL of PBS and resuspension in fresh BHI media. Adhesion videos were obtained using black

welled plates by adding 4 x 10^5 cells to 500 μL of BHI media and allowing adhesion for 2 hours prior to imaging

at 1.25 min intervals.

Confocal microscopy and image processing

For fluorescence and confocal microscopy, 5 x 10^6 swimming cells from mid-log cultures were concentrated by

centrifugation at 800 rcf for 5 minutes, washed once in PBS, then resuspended in PBS at a final concentration of

10^7 cells/mL. 500 μL of this suspension was then pipetted onto coverslips pretreated with 0.01% poly-L-lysine.

After allowing cells to attach for 20 minutes at room temperature, coverslips were washed twice with PBS,

followed by fixation in 4% paraformaldehyde for 15 minutes. Coverslips were then washed again followed by

permeabilization with 0.1% Triton X-100 in PBS for 5 minutes. Coverslips were washed in PBS prior to staining

with 0.2 μg/mL DAPI in PBS for 5 minutes. After a final PBS wash, coverslips were mounted on a glass slide

using Vectashield mounting medium (Vector Laboratories). For fluorescence microscopy of attached cells, a

standard adherence assay was conducted, as described above, on 1.5 thickness glass-bottom dishes pre-coated

with poly-L-lysine (MatTek). After 24 hours of adherent growth, the final BHI wash contained 16%

formaldehyde added to a final concentration of 4%. The plate was processed from this point as described above

for the swimmer cell slides. The slides and plates were analyzed by confocal microscopy using a Leica SP5-II

with a 100x objective. A Z-series was acquired for GFP/YFP, DAPI and brightfield. Following image acquisition,

deconvolution analysis was performed on the fluorescent images using Huygens Essential deconvolution software

(version 17.04.1p2 64b; SVI). Brightfield images are single confocal sections with a flatfield correction applied.

All images of parental, *Cf*PDEA::GFP, and *Cf*RAC1::GFP were acquired with the same microscope settings.

Equivalent linear adjustments were made to the brightness of each image using Photoshop Version 23.5.1

(Adobe).

Western blot analysis

For swimming cultures, samples were generated from $2x10^6$ log-phase cells. For adhered cultures, an equivalent

number of cells were grown on 60 mm dishes in stationary culture for 24 hours and then dislodged using a cell

scraper and counted by hemacytometer. Both swimming and dislodged adherent cells were transferred to

microcentrifuge tubes and collected by centrifugation, washed once with PBS, then resuspended in Laemmli

SDS-PAGE sample buffer at a concentration of 1×10^5 cell equivalents/µl. Samples boiled for 5 min at 95 ºC and

centrifuged briefly. Cell equivalents of ~1.5×10^6 cells per lane were separated by 10% SDS-PAGE along with the

Precision Plus Protein Kaleidoscope standard (BioRad). The fractionated proteins were transferred to a PVDF

membrane. The membrane was then incubated in blocking buffer (PBS containing 5% milk and 0.2% Tween-20)

for 1 hour at room temperature. A mouse anti-GFP antibody (Roche) was used as the primary antibody at a 1:500

dilution in blocking buffer overnight at 4 °C. Horseradish peroxidase-conjugated goat anti-mouse IgG in blocking

buffer was used as the secondary antibody (1:5000). After washing, Enhanced Chemiluminescent Substrate

(BioRad) was used to develop the blots. For the loading control, the blot was reprobed with a rabbit anti-*Tb*BiP

antibody (a gift of Jay Bangs) at a 1:5000 dilution in blocking buffer, followed by washing and probing with an

anti-rabbit-HRP secondary antibody (1:5000). Images were taken by an Azure biosystems c600 using

chemiluminescence settings.

Pharmacological inhibition

Pharmacological inhibitors were dissolved in DMSO. For vehicle controls, an amount of DMSO equal to that of

the most concentrated sample was added. The compounds were described previously: NPD-001 [22], NPD-008

[53], NPD-055 [54], and NPD-226 [55].

Statistical analyses and data visualization

Prism Version 9.4.1 (458) (GraphPad Software) was used to generate all plots and for all statistical tests. Plots

were assembled into figures using Illustrator Version 26.5 (Adobe).

Acknowledgments

We thank Biao Zuo for preparing TEM samples and assistance with image acquisition and Koranda Walsh and

the PennVet Clinical Pathology lab for processing the Giemsa staining samples. We thank Emmanuel Tetaud,

Sam Dean, Eva Gluenz, and Tom Beneke for plasmids and advice. Thanks to Stephen Beverley for providing the

Cf-C1 wild-type *C. fasciculata* cell line and who, along with Wesley Warren, Chad Tomlinson, and Peter Myler,

made the unpublished genome sequence available on TriTrypDB. We thank Jay Bangs for providing the anti-BiP

antibody and Lindsay Bair, Laura Anastor-Walters, and Nancy Peltier for their technical assistance. We thank

Jesse Quatse for assistance with Photoshop. Finally, we acknowledge the PennVet Imaging Core for providing

access to instrumentation and analysis software used in live-cell and confocal imaging and image processing.

References

1. Diseases GBD, Injuries C. Global burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019. Lancet. 2020;396(10258):1204-22. doi: 10.1016/S0140-6736(20)30925-9. PubMed PMID: 33069326; PubMed Central PMCID: PMCPMC7567026.

2. Goncalves CS, Avila AR, de Souza W, Motta MCM, Cavalcanti DP. Revisiting the *Trypanosoma cruzi* metacyclogenesis: morphological and ultrastructural analyses during cell differentiation. Parasit Vectors. 2018;11(1):83. Epub 20180206. doi: 10.1186/s13071-018-2664-4. PubMed PMID: 29409544; PubMed Central PMCID: PMCPMC5801705.

3. Kollien AH, Schaub GA. The development of *Trypanosoma cruzi* in triatominae. Parasitol Today. 2000;16(9):381-7. doi: 10.1016/s0169-4758(00)01724-5. PubMed PMID: 10951597.

4. Vickerman K, Tetley L. Flagellar surfaces of parasitic protozoa and their role in attachment. In: Bloodgood RA, editor. Ciliary and Flagellar Membranes. Boston, MA: Springer US; 1990. p. 267-304.

5. Brooker BE. Flagellar Adhesion of *Crithidia fasciculata* to Millipore Filters. Protoplasma. 1971;72(1):19-25.

6. Brooker BE. Desmosomes and hemidesmosomes in the flagellate *Crithidia fasciculata*. Z Zellforsch Mikrosk Anat. 1970;105(2):155-66. doi: 10.1007/BF00335467. PubMed PMID: 4195325.

7. Filosa JN, Berry CT, Ruthel G, Beverley SM, Warren WC, Tomlinson C, et al. Dramatic changes in gene expression in different forms of *Crithidia fasciculata* reveal potential mechanisms for insect-specific adhesion in kinetoplastid parasites. PLoS Negl Trop Dis. 2019;13(7):e0007570. Epub 2019/07/30. doi: 10.1371/journal.pntd.0007570. PubMed PMID: 31356610.

8. Hommel M, Robertson E. In vitro attachment of trypanosomes to plastic. Experientia. 1976;32(4):464-6. doi: 10.1007/BF01920798. PubMed PMID: 1269646.

9. Tagoe DN, Kalejaiye TD, de Koning HP. The ever unfolding story of cAMP signaling in trypanosomatids: vive la difference! Front Pharmacol. 2015;6:185. Epub 20150907. doi: 10.3389/fphar.2015.00185. PubMed PMID: 26441645; PubMed Central PMCID: PMCPMC4561360.

10. Saada EA, Kabututu ZP, Lopez M, Shimogawa MM, Langousis G, Oberholzer M, et al. Insect stage-specific receptor adenylate cyclases are localized to distinct subdomains of the *Trypanosoma brucei* flagellar membrane. Eukaryot Cell. 2014;13(8):1064-76. Epub 20140530. doi: 10.1128/EC.00019-14. PubMed PMID: 24879126; PubMed Central PMCID: PMCPMC4135804.

11. Gould MK, Bachmaier S, Ali JA, Alsford S, Tagoe DN, Munday JC, et al. Cyclic AMP effectors in African trypanosomes revealed by genome-scale RNA interference library screening for resistance to the phosphodiesterase inhibitor CpdA. Antimicrob Agents Chemother. 2013;57(10):4882-93. Epub 20130722. doi: 10.1128/AAC.00508-13. PubMed PMID: 23877697; PubMed Central PMCID: PMCPMC3811416.

12. Shaw S, Knüsel S, Abbühl D, Naguleswaran A, Etzensperger R, Benninger M, et al. Cyclic AMP signalling and glucose metabolism mediate pH taxis by African trypanosomes. Nature Communications. 2022;13(1). doi: 10.1038/s41467-022-28293-w.

13. Bachmaier S, Volpato Santos Y, Kramer S, Githure GB, Klöckner T, Pepperl J, et al. Nucleoside analogue activators of cyclic AMP-independent protein kinase A of *Trypanosoma*. Nature Communications. 2019;10(1). doi: 10.1038/s41467-019-09338-z.

14. Oberholzer M, Marti G, Baresic M, Kunz S, Hemphill A, Seebeck T. The *Trypanosoma brucei* cAMP
phosphodiesterases TbrPDEB1 and TbrPDEB2: flagellar enzymes that are essential for parasite virulence. FASEB
J. 2007;21(3):720-31. Epub 20061213. doi: 10.1096/fj.06-6818com. PubMed PMID: 17167070.

15. Luginbuehl E, Ryter D, Schranz-Zumkehr J, Oberholzer M, Kunz S, Seebeck T. The N terminus of
phosphodiesterase TbrPDEB1 of *Trypanosoma brucei* contains the signal for integration into the flagellar
skeleton. Eukaryot Cell. 2010;9(10):1466-75. Epub 20100806. doi: 10.1128/EC.00112-10. PubMed PMID:
20693305; PubMed Central PMCID: PMCPMC2950416.

16. Baillie GS. Compartmentalized signalling: spatial regulation of cAMP by the action of compartmentalized
phosphodiesterases. FEBS Journal. 2009;276(7):1790-9. doi: 10.1111/j.1742-4658.2009.06926.x.

17. Scolaro EJ, Ames RP, Brittingham A. Growth-phase dependent substrate adhesion in *Crithidia fasciculata*. J
Eukaryot Microbiol. 2005;52(1):17-22. doi: 10.1111/j.1550-7408.2005.3315r.x. PubMed PMID: 15702975.

18. Hughes D, Simpson L, Schneider CA. Flagellar adherence of *Crithidia fasciculata* cells in culture. The
Journal of Protozoology. 1983;30(4):635-41. doi: 10.1111/j.1550-7408.1983.tb05335.x.

19. Sunter JD, Yanase R, Wang Z, Catta-Preta CMC, Moreira-Leite F, Myskova J, et al. *Leishmania* flagellum
attachment zone is critical for flagellar pocket shape, development in the sand fly, and pathogenicity in the host.
Proc Natl Acad Sci U S A. 2019;116(13):6351-60. Epub 20190308. doi: 10.1073/pnas.1812462116. PubMed
PMID: 30850532; PubMed Central PMCID: PMCPMC6442623.

20. Wheeler RJ, Sunter JD, Gull K. Flagellar pocket restructuring through the *Leishmania* life cycle involves a
discrete flagellum attachment zone. J Cell Sci. 2016;129(4):854-67. Epub 20160108. doi: 10.1242/jcs.183152.
PubMed PMID: 26746239; PubMed Central PMCID: PMCPMC4760377.

21. Veerman J, van den Bergh T, Orrling KM, Jansen C, Cos P, Maes L, et al. Synthesis and evaluation of analogs
of the phenylpyridazinone NPD-001 as potent trypanosomal TbrPDEB1 phosphodiesterase inhibitors and in vitro
trypanocidals. Bioorg Med Chem. 2016;24(7):1573-81. Epub 20160226. doi: 10.1016/j.bmc.2016.02.032.
PubMed PMID: 26935942.

22. de Koning HP, Gould MK, Sterk GJ, Tenor H, Kunz S, Luginbuehl E, et al. Pharmacological validation of
Trypanosoma brucei phosphodiesterases as novel drug targets. J Infect Dis. 2012;206(2):229-37. Epub 20120130.
doi: 10.1093/infdis/jir857. PubMed PMID: 22291195; PubMed Central PMCID: PMCPMC3379837.

23. Beneke T, Madden R, Makin L, Valli J, Sunter J, Gluenz E. A CRISPR Cas9 high-throughput genome editing
toolkit for kinetoplastids. R Soc Open Sci. 2017;4(5):170095. Epub 20170503. doi: 10.1098/rsos.170095.
PubMed PMID: 28573017; PubMed Central PMCID: PMCPMC5451818.

24. Beattie P, Gull K. Cytoskeletal architecture and components involved in the attachment of *Trypanosoma
congolense* epimastigotes. Parasitology. 1997;115 (Pt 1):47-55. doi: 10.1017/s0031182097001042. PubMed
PMID: 9280895.

25. Kleffmann T, Schmidt J, Schaub GA. Attachment of Trypanosoma cruzi epimastigotes to hydrophobic
substrates and use of this property to separate stages and promote metacyclogenesis. J Eukaryot Microbiol.
1998;45(5):548-55. doi: 10.1111/j.1550-7408.1998.tb05115.x. PubMed PMID: 9783457.

26. Skalický T, Dobáková E, Wheeler RJ, Tesařová M, Flegontov P, Jirsová D, et al. Extensive flagellar
remodeling during the complex life cycle of *Paratrypanosoma*, an early-branching trypanosomatid. Proceedings
of the National Academy of Sciences. 2017;114(44):11757-62. doi: 10.1073/pnas.1712311114.

27. Wakid MH, Bates PA. Flagellar attachment of Leishmania promastigotes to plastic film in vitro. Exp
Parasitol. 2004;106(3-4):173-8. doi: 10.1016/j.exppara.2004.03.001. PubMed PMID: 15172225.

28. Brooker BE. Flagellar attachment and detachment of *Crithidia fasciculata* to the gut wall of *Anopheles*
gambiae. Protoplasma. 1971;73(2):191-202. doi: 10.1007/BF01275594. PubMed PMID: 5126403.

29. Evans DA, Ellis DS, Stamford S. Ultrastructural studies of certain aspects of the development of
Trypanosoma congolense in Glossina morsitans morsitans. J Protozool. 1979;26(4):557-63. doi: 10.1111/j.1550-
7408.1979.tb04195.x. PubMed PMID: 544799.

30. Alcolea PJ, Alonso A, García-Tabares F, Toraño A, Larraga V. An insight into the proteome of *Crithidia*
fasciculata choanomastigotes as a comparative approach to axenic growth, peanut lectin agglutination and
differentiation of *Leishmania* spp. promastigotes. PloS one. 2014;9(12):e113837. doi:
10.1371/journal.pone.0113837.

31. Figueiredo RC, Rosa DS, Soares MJ. Differentiation of *Trypanosoma cruzi* epimastigotes: metacyclogenesis
and adhesion to substrate are triggered by nutritional stress. The Journal of parasitology. 2000;86(6):1213-8. doi:
10.1645/0022-3395(2000)086[1213:DOTCEM]2.0.CO;2. PubMed PMID: 11191893.

32. Greenblatt CL, Handman E, Mitchell GF, Battye FL, Schnur LF, Snary D. Phenotypic diversity of cloned
lines of Leishmania major promastigotes. Z Parasitenkd. 1985;71(2):141-57. doi: 10.1007/BF00926265. PubMed
PMID: 3922149.

33. Iovannisci DM, Plested CP, Moe GR. Evidence for rosettes as an unrecognized stage in the life cycle of
Leishmania parasites. J Eukaryot Microbiol. 2010;57(5):405-14. Epub 20100805. doi: 10.1111/j.1550-
7408.2010.00496.x. PubMed PMID: 20707828; PubMed Central PMCID: PMCPMC3086735.

34. Leishman WB. Critical Review: Kala Azar and Tropical Sore. QJM: An International Journal of Medicine.
1911;os5(1):109-52. doi: 10.1093/oxfordjournals.qjmed.a069298.

35. Noguchi H, Tilden EB. Comparative Studies of Herpetomonads and Leishmanias : I. Cultivation of
Herpetomonads from Insects and Plants. J Exp Med. 1926;44(3):307-25. doi: 10.1084/jem.44.3.307. PubMed
PMID: 19869186; PubMed Central PMCID: PMCPMC2131765.

36. Sacks D, Kamhawi S. Molecular aspects of parasite-vector and vector-host interactions in leishmaniasis. Annu
Rev Microbiol. 2001;55:453-83. doi: 10.1146/annurev.micro.55.1.453. PubMed PMID: 11544364.

37. Pimenta PF, Turco SJ, McConville MJ, Lawyer PG, Perkins PV, Sacks DL. Stage-specific adhesion of
Leishmania promastigotes to the sandfly midgut. Science. 1992;256(5065):1812-5. doi: 10.1126/science.1615326.
PubMed PMID: 1615326.

38. Warburg A, Hamada GS, Schlein Y, Shire D. Scanning electron microscopy of *Leishmania major* in
Phlebotomus papatasi. Z Parasitenkd. 1986;72(4):423-31. doi: 10.1007/BF00927886. PubMed PMID: 3751230.

39. Skalicky T, Dobakova E, Wheeler RJ, Tesarova M, Flegontov P, Jirsova D, et al. Extensive flagellar
remodeling during the complex life cycle of Paratrypanosoma, an early-branching trypanosomatid. Proc Natl
Acad Sci U S A. 2017;114(44):11757-62. Epub 20171016. doi: 10.1073/pnas.1712311114. PubMed PMID:
29078369; PubMed Central PMCID: PMCPMC5676924.

40. Rojas R, Segovia C, Trombert AN, Santander J, Manque P. The effect of tunicamycin on the glucose uptake,
growth, and cellular adhesion in the protozoan parasite *Crithidia fasciculata*. Current Microbiology.
2014;69(4):541-8. doi: 10.1007/s00284-014-0620-x.

41. Makin L, Gluenz E. cAMP signalling in trypanosomatids: role in pathogenesis and as a drug target. Trends in
parasitology. 2015;31(8):373-9. Epub 20150521. doi: 10.1016/j.pt.2015.04.014. PubMed PMID: 26004537;
PubMed Central PMCID: PMCPMC4534343.

42. Gonzales-Perdomo M, Romero P, Goldenberg S. Cyclic AMP and adenylate cyclase activators stimulate
Trypanosoma cruzi differentiation. Exp Parasitol. 1988;66(2):205-12. doi: 10.1016/0014-4894(88)90092-6.
PubMed PMID: 2840306.

43. Rolin S, Paindavoine P, Hanocq-Quertier J, Hanocq F, Claes Y, Le Ray D, et al. Transient adenylate cyclase
activation accompanies differentiation of *Trypanosoma brucei* from bloodstream to procyclic forms. Mol
Biochem Parasitol. 1993;61(1):115-25. doi: 10.1016/0166-6851(93)90164-s. PubMed PMID: 8259124.

44. Laxman S, Riechers A, Sadilek M, Schwede F, Beavo JA. Hydrolysis products of cAMP analogs cause
transformation of *Trypanosoma brucei* from slender to stumpy-like forms. Proceedings of the National Academy
of Sciences. 2006;103(50):19194-9. doi: 10.1073/pnas.0608971103.

45. Oberholzer M, Saada EA, Hill KL. Cyclic AMP regulates social behavior in African trypanosomes. mBio.
2015;6(3):e01954-14. Epub 20150428. doi: 10.1128/mBio.01954-14. PubMed PMID: 25922395; PubMed Central
PMCID: PMCPMC4436052.

46. Shaw S, Demarco SF, Rehmann R, Wenzler T, Florini F, Roditi I, et al. Flagellar cAMP signaling controls
trypanosome progression through host tissues. Nature Communications. 2019;10(1). doi: 10.1038/s41467-019-
08696-y.

47. Kelly FD, Sanchez MA, Landfear SM. Touching the surface: diverse roles for the flagellar membrane in
kinetoplastid parasites. Microbiol Mol Biol Rev. 2020;84(2). Epub 20200401. doi: 10.1128/MMBR.00079-19.
PubMed PMID: 32238446; PubMed Central PMCID: PMCPMC7117551.

48. Tyler KM, Fridberg A, Toriello KM, Olson CL, Cieslak JA, Hazlett TL, et al. Flagellar membrane localization
via association with lipid rafts. J Cell Sci. 2009;122(Pt 6):859-66. Epub 20090224. doi: 10.1242/jcs.037721.
PubMed PMID: 19240119; PubMed Central PMCID: PMCPMC2714428.

49. Bertiaux E, Mallet A, Fort C, Blisnick T, Bonnefoy S, Jung J, et al. Bidirectional intraflagellar transport is
restricted to two sets of microtubule doublets in the trypanosome flagellum. J Cell Biol. 2018;217(12):4284-97.
Epub 20181001. doi: 10.1083/jcb.201805030. PubMed PMID: 30275108; PubMed Central PMCID:
PMCPMC6279389.

50. Dean S, Sunter J, Wheeler RJ, Hodkinson I, Gluenz E, Gull K. A toolkit enabling efficient, scalable and
reproducible gene tagging in trypanosomatids. Open Biol. 2015;5(1):140197. doi: 10.1098/rsob.140197. PubMed
PMID: 25567099; PubMed Central PMCID: PMCPMC4313374.

51. Schumann Burkard G, Jutzi P, Roditi I. Genome-wide RNAi screens in bloodstream form trypanosomes
identify drug transporters. Mol Biochem Parasitol. 2011;175(1):91-4. Epub 20100922. doi:
10.1016/j.molbiopara.2010.09.002. PubMed PMID: 20851719.

52. Tetaud E, Lecuix I, Sheldrake T, Baltz T, Fairlamb AH. A new expression vector for *Crithidia fasciculata* and
Leishmania. Mol Biochem Parasitol. 2002;120(2):195-204. doi: 10.1016/s0166-6851(02)00002-6. PubMed
PMID: 11897125.

53. Blaazer AR, Singh AK, de Heuvel E, Edink E, Orrling KM, Veerman JJN, et al. Targeting a subpocket in
Trypanosoma brucei Phosphodiesterase B1 (TbrPDEB1) enables the structure-based discovery of selective
inhibitors with trypanocidal activity. J Med Chem. 2018;61(9):3870-88. Epub 20180501. doi:
10.1021/acs.jmedchem.7b01670. PubMed PMID: 29672041; PubMed Central PMCID: PMCPMC5949723.

54. de Heuvel E, Singh AK, Boronat P, Kooistra AJ, van der Meer T, Sadek P, et al. Alkynamide phthalazinones
as a new class of TbrPDEB1 inhibitors (Part 2). Bioorg Med Chem. 2019;27(18):4013-29. Epub 20190705. doi:
10.1016/j.bmc.2019.06.026. PubMed PMID: 31378593.

55. Salado IG, Singh AK, Moreno-Cinos C, Sakaine G, Siderius M, Van der Veken P, et al. Lead optimization of
phthalazinone phosphodiesterase inhibitors as novel antitrypanosomal compounds. J Med Chem.
2020;63(7):3485-507. Epub 20200326. doi: 10.1021/acs.jmedchem.9b00985. PubMed PMID: 32196340.

Figure Legends

Figure 1. An adherence assay reveals features of *C. fasciculata* attachment. **A)** Schematic showing steps of the

assay, including the variables tested. **B)** Representative images showing attached rosettes produced by varying the

initial plating time. The dashed red lines were drawn to indicate what was counted manually as individual

rosettes, clusters containing 4 or more cells. **C)** Quantitation of the experiment shown in B. This experiment was

performed in triplicate. Error bars show standard error. **D)** The effect of initial cell concentration on adherence.

10^7 cells/mL from cultures in different phase of growth, were adhered for each sample. Quantitation of a

representative experiment of three replicates is shown with median values indicated by red lines. **E)** Rosettes

resulting from adherence of different numbers of cells, all derived from a mid-log phase culture (10^7 cells/mL). A

representative experiment of three replicates is shown. Red lines indicate median values, and the blue dotted line

represents the proportion of plated cells that were able to adhere.

Figure 2. Flagellar shortening occurs at a relatively constant rate following an initial adhesion event. **A)**

Following 5 minutes of plating and subsequent washing, cells were observed at the indicated times and the length

of the flagellum was measured in 20 representative cells (different cells were measured at each time point). **B)** An

example of a single cell undergoing flagellar shortening. The numbers indicate time in minutes. The schematic on

the right shows a representation of the length of the flagellum at the beginning (red) and end (blue) of the time

course.

Figure 3. Transmission electron microscopy reveals structural differences between attached and swimming cells.

A) Cross section through *C. fasciculata* parasites adhered to tissue culture plastic (Fig S2, yellow plane).

Asterisks indicate hemidesmosomes at the distal tip of the shortened flagella. Scale bar is 1 µm. The dotted box

indicates the region enlarged in panel A'. **A')** Close up of a hemidesmosome (hd) from one of the cells in panel A.

Desmosome-like FAZ structures, connecting the flagellum to the cell body, are also observed (fz). Scale bar is

200 nm. **B)** and **C)** Additional examples of hemidesmosomes (hd) and desmosome-like FAZ (fz). The cell in C)

was scraped from the dish prior to fixation. Scale bar is 200 nm. **D)** Sectioning *en face* shows a region near the

end of the flagellum in an adherent cell (Fig S2, blue planes). The flagellar membrane is enlarged and no flagellar

axoneme can be observed. Instead, diffuse filaments probably comprising the hemidesmosome are seen. Scale bar

is 200 nm. **E)** Another example of a hemidesmosome sectioned *en face*. Scale bar is 200 nm. **F)** Section of a

swimming cell shows the flagellum extending from the flagellar pocket. Dotted box indicates the region enlarged

in F'. Scale bar is 1 µm. **F')** Higher magnification view of the flagellar pocket of a swimming cell. FAZ structures

link the flagellar membrane to the flagellar pocket membrane, but no hemidesmosome is seen. The slight bulge in

the flagellum where it exits the pocket may be a physical correlate to the site of initial adhesion. Filamentous

material can be observed in the flagellar pocket of the swimming cell but was never observed on the surface. In

contrast, the surface of attached cells, **G)** was covered with this material. Some of the same material is also visible

in panels A, B, C, D, and E. Scale bar F' and G is 400 nm.

Figure 4. Phosphodiesterase inhibitors block *C. fasciculata* adherence. **A)** Adherence assays performed in the

presence of 10 µM of either NPD-001 (NPD1) or NPD-008 (NPD8). The number of single adhered cells at 2

hours (open circles) and the number of rosettes ($\geq$ 4 cells, gray squares) at 24 hours were counted and compared

to those from a vehicle-treated control sample (Veh) and an untreated sample (Con). Treatments were present in

the media both during the 2-hour plating time and the subsequent 24 hours of culture. All comparisons were

made. Asterisks indicate Kruskal-Wallis test with Dunn's correction $P<0.0001$ (****) and $P<0.05$ (*) **B)**

Adhesion assays comparing vehicle and untreated controls to assays performed in the presence of different

concentrations of NPD-001 or **C)** NPD-226. The number of single adhered cells at 2 hours (open circles) and the

number of rosettes ($\geq$ 4 cells, gray squares) at 24 hours were counted. All comparisons were made, but only the

groups that are significantly different from the adjacent treatment are shown. Asterisks indicate Kruskal-Wallis

test with Dunn's correction $P<0.0001$ (****), and $P<0.01$ (**).

Figure 5. NPD-001 does not affect the growth rate of *C. fasciculata*. **A)** Graph of *C. fasciculata* growth in

shaking cultures assayed over approximately 31 h in standard medium (Control; blue circles), medium

supplemented containing 10 µM NPD-001 in vehicle (NPD1; orange diamonds), or medium containing vehicle

only (Vehicle, red squares). **B)** Time encoded movies were used to measure the doubling time of established

attached cultures after the addition of NPD-001 in vehicle, or in vehicle alone. The red line indicates the mean. No

significant difference was noted between the doubling times.

Figure 6. Components of cAMP signaling localize to different domains of the flagellum in *C. fasciculata*

swimmers. Deconvolved confocal images of GFP and YFP fusion proteins and DAPI staining of nuclear and

kDNA of fixed swimmer and attached cell cultures. A Z-stack of the entire cells and rosettes were captured.

Fluorescent images are shown individually and merged. A brightfield image for each is included for reference and

was used to generate the red dashed outline of the flagellum in the merged and brightfield images. An overlay of

fluorescent channels on the corresponding brightfield image also shows the extent of GFP or YFP signal with

respect to the length of the flagellum (Fig S4). The scale bars are 5 µm. Western blots of extracts of swimmers

(Sw) and attached (Ad) cells of each line are included supporting the presence of the correct size GFP- or YFP-

fusion protein (Fig S3) in both swimmer and attached cells regardless of their localization patterns. Equivalent

loading was confirmed by re-probing blots with an anti-BiP antibody. Parental Cf-C1 cells lack GFP signal in

swimmers and adherent cells as well as by western analysis (Fig S5).

Figure 7. Model of the developmental transition from swimmer to adherent cell. The localizations of *Cf*RAC1

(red), and *Cf*PDEA (blue) are indicated. Minutes following adhesion via the region of the flagellum near the cell

body (orange), the process of flagellar shortening begins. Flagellar shortening brings *Cf*RAC1 closer to the cell

body. Eventually the source of cAMP is close enough to the cell and the sink for cAMP (*Cf*PDEA) is reduced

such that cAMP levels rise above a critical threshold in the cell body (shown on the graph by a star). This

promotes differentiation to the adherent cell fate characterized cytoplasmic localization of *Cf*RAC1 and *Cf*PDEA

by robust changes in gene regulation and formation of a hemidesmosome.

Figure S1. Density dependent changes to *C. fasciculata* morphology. **A)** Giemsa-stained samples of cells at log-

phase (10^7 cells/mL) or past log-phase growth (10^8 cells/mL). Scale bar is 10 μm. **B)** Plot of individual measured

cell lengths in micrometers. Cells were measured along the anterior-posterior axis of the cell body excluding the

flagellum. The data are normally distributed (Shapiro-Wilk test). The red line indicates the mean and standard

deviation. An unpaired t test indicates the cell length distributions are significantly different with a P value of

0.005.

Figure S2. Illustration of sectioning planes used for transmission electron microscopy. An adherent cell shown in

cross section (yellow plane; sections in Fig 3A-C, and F-G). Sections cut *en face* at different distances from the

culture plate (blue planes). Depending on the position of the *en face* section relative to the culture plate, the

osmiophilic hemidesmosome-like structure will appear darker (Fig 3E) or lighter (Fig 3D) and other cell features,

such as the cell cytoplasm and mitochondrion may be visible (Fig 3D).

Figure S3. Phosphodiesterase inhibitor NPD-055 partially reduces *C. fasciculata* adherence. Adherence assays

performed in the presence in the presence of different amounts of NPD-055 compared to vehicle-only treated

controls. The number of single adhered cells at 2 hours (open circles) and the number of rosettes (≥ 4 cells, gray

squares) at 24 hours were counted and compared to those from a vehicle-treated control sample (Veh). Treatments

were present in the media both during the 2-hour plating time and the subsequent 24 hours of culture. Data are not

normally distributed. Red lines indicate the median. All comparisons were made, but only the groups that are

significantly different from the adjacent treatment are shown. Asterisks indicate Kruskal-Wallis test with Dunn's

correction P<0.0001 (****), P<0.001 (***), and P<0.05 (*).

Figure S4. Schematic diagram of tagged proteins used for localization studies. Open reading frames encoding

*Cf*PDEA (CFAC1_140020300) and *Cf*RAC1 (CFAC1_090006500) were cloned without their C-terminal stop

codon into a vector resulting in an in-frame fusion with the coding sequence of GFP (green rectangle). The

resulting constructs were transfected into *C. fasciculata* and lines episomally expressing the tagged proteins were

selected using hygromycin. A single allele of *Cf*PF16 (CFAC1_170028000) was fused in-frame at its C-terminus

with YFP (yellow rectangle) by homologous recombination followed by neomycin selection. The number of

amino acids (aa) for each fusion protein is provided. Domains and their relative positions indicated in the different

constructs are 3'5'-cyclic nucleotide phosphodiesterase, catalytic domain (PDEase; IPR002073), Periplasmic

binding protein-like I (PBPL; IPR028082), Nucleotide cyclase (AC; IPR029787), Armadillo-type fold

(Armadillo-fold; IPR016024), signal peptide for secretion (SP), and transmembrane domain (TM). Further details

are provided in the Materials and Methods.

Figure S5. Overlay images show the extent of fluorescent signal along the swimming cell flagellum. Overlay of

the brightfield and fluorescent images of parental Cf-C1, *Cf*PDEA::GFP, *Cf*RAC1::GFP, and *Cf*PF16::YFP from

Figure 6. The fluorescence of *Cf*PDEA::GFP reaches the cell body and weakly inside, similar, but less uniform as

*Cf*PF16::YFP. In contrast, *Cf*RAC1::GFP is more restricted to the distal part of the flagellum. Parental cells lack

GFP signal. The scale bars are 5 μm.

Figure S6. Parental Cf-C1 lack GFP signal by confocal and western blot analyses. Deconvolved confocal images

of GFP and DAPI images of fixed swimmer and attached cell cultures taken with the same settings as shown in

Figure 6. Fluorescent images are shown individually and merged. A brightfield image for each is included for

reference and was used to generate the dashed outline of the flagellum in the merged image. A western blot of

extracts of swimming and attached cells probed with anti-GFP and anti-BiP (loading control) antibodies.

Video S1. 4×10^5 parental Cf-C1 cells were added to wells of a 24-well plate and imaged at 75 second intervals for

16 hours to obtain movies of *C. fasciculata* as they adhere. This video shows the full imaging frame at 60x

magnification, which highlights several adhesion events occurring during the imaging time. Note an oblong out-

of-focus object appears to the left of center. The scale bar is 50 μm.

Video S2. Close up and cropped version of Video S1. For reference, the same oblong out-of-focus object that is

visible near the center of Video S1 appears at the left edge of this video. Note when a dividing pair of cells first

adheres to the culture dish surface (time 5:42:30), one cell swims away. The other remains characteristically

attached at the junction between the cell body and the flagellum. Both the cell body and the flagellum move

independently until the flagellum is retracted and the cell body is round (time 8:40:00). The first division as an

adherent cell occurs at approximately 9:08:45. One of the adherent daughters divides again approximately 4 hours

and 4 minutes later (time 13:12:30). The other adherent daughter divides approximately 5 hours and 25 minutes

later (time 14:33:45). The scale bar is 10 μm.

A

Time
Growth phase
Number

Plate

Wash

side	above
10-30 min	
2 h	
24 h	

Culture | Quantitate

B

30' 60' 90'

120' 150' 180'

210' 240' 240'annotated

C Plating time

D Growth phase

E Log-phase cell number

A

Flagellar Shortening

B

A Swimmer Cell Growth Curve

B Attached Cell Division Time