Inbreeding depression in polyploid species: a meta-analysis

Josselin Clo[□], Filip Kolář [□,□]

[□] Department of Botany, Faculty of Science, Charles University in Prague, Benátská 2, CZ-

128 01 Prague, Czech Republic

[□] Institute of Botany of the Czech Academy of Sciences, Zámek 1, CZ-252 43 Průhonice,

Czech Republic

For correspondence: Josselin Clo, Faculty of Science, Charles University in Prague,

Benátská 2, CZ-128 01 Prague, Czech Republic, ORCID 0000-0002-3295-9481,

josselin.clo@gmail.com

Abstract:

Whole-genome duplication is a common mutation in eukaryotes with far-reaching phenotypic effects, the resulting morphological and fitness consequences and how they affect the survival of polyploid lineages are intensively studied. Another important factor may also determine the probability of establishment and success of polyploid lineages: inbreeding depression. Inbreeding depression is expected to play an important role in the establishment of neopolyploid lineages, their capacity to colonize new environments, and in the simultaneous evolution of ploidy and other life-history traits such as self-fertilization. Both theoretically and empirically, there is no consensus on the consequences of polyploidy on inbreeding depression. In this meta-analysis, we investigated the effect of polyploidy on the evolution of inbreeding depression, by performing a meta-analysis within angiosperm species. The main results of our study are that the consequences of polyploidy on inbreeding depression are complex and depends on the time since polyploidization. We found that young polyploid lineages have a much lower amount of inbreeding depression than their diploid relatives and their established counterparts. Natural polyploid lineages are intermediate, and have a higher amount of inbreeding depression than synthetic neopolyploids, and a smaller amount than diploids, suggesting that the negative effect of polyploidy on inbreeding depression decrease with time since polyploidization.

Keywords: polyploidy, genome doubling, fitness, inbreeding depression, polyploid establishment.

Introduction:

Polyploidization has occurred numerous times during the evolution of eukaryotes [1,2] and has been shown to have a broad range of phenotypic and genomic effects and to be an important mechanism for plant adaptation and speciation [3]. Nevertheless, polyploidization initially arises with several disadvantages in neopolyploids, like mitotic and meiotic dysfunction, genomic instability, decrease in fitness, and negative frequency-dependent selection [4–9].

Another important factor may also determine the probability of survival and success of polyploid lineages: inbreeding depression (ID hereafter) [10,11]. ID can be defined as the reduction of fitness found in selfed progenies compared to outcrossed progenies [12], and is predominantly due to the expression of recessive deleterious alleles at their homozygous state [13]. ID theoretically plays an important role in polyploids establishment. The initially low frequency of polyploid lineages within a diploid population may lead to strong bi-parental inbreeding [14]. In such conditions, it has been shown theoretically that a decrease in ID in polyploids compared to diploids is necessary for them to establish [10,11].

Polyploidy is theoretically expected to have both positive and negative effects on the amount of ID (see [18] for review). In the short term, it has been shown theoretically that in autopolyploid species, the strong bottleneck associated with polyploidization can strongly decrease the amount of ID due to the loss and/or the fixation of recessive deleterious mutations [15]. Even if not tested, we can expect similar consequences in allopolyploid species. In autopolyploid species with polysomic inheritance, even without a loss of genetic diversity, if frequencies of deleterious mutations remain similar in different cytotypes, as homozygosity increases at slower rates in autopolyploids compared to diploids [19], the

expression of ID should be less severe in neoautopolyploids. In the long-term, the better

masking of recessive deleterious mutations in autopolyploids should make them segregate at

higher frequencies than in diploid populations [20]. Depending on the dominance coefficients

of the mutations in the autopolyploid heterozygous genotypes, this increase in frequency can

make ID to be smaller [21] or higher [22] in autopolyploids compared to diploids.

Empirically, both a decrease [17,23] or an increase [24,25] of ID in synthetic and natural

autopolyploids compared to diploids have been observed, and no strong consensus can be

made. Allopolyploidy received much less attention, and there are consequently fewer

expectations than in autopolyploid species. Hedrick [26] nevertheless showed that in

homosporous, allopolyploid ferns, because offspring will be strongly homozygous due to

intragametophytic selfing, ID should be lower in allopolyploids compared to diploids.

In this study, we investigated the effect of polyploidy on the evolution of ID. To do so,

we performed a meta-analysis within angiosperm species. The main results of our study are

that the effect of polyploidy on ID is complex and depends on the time since polyploidization.

We found that synthetic polyploid lineages have a lower amount of ID than their diploid

progenitors and their established counterpart. Natural polyploid lineages are intermediate, and

have a higher amount of ID than synthetic neopolyploids, and a smaller amount than diploids.

Material & Methods:

Dataset compilation

For this study, we were interested in the amount of ID in polyploids (auto- and

allopolyploids, and of synthetic and natural genomic origins) compared to their diploid

progenitors. As the studies comparing diploid and polyploid populations of the same species

in the same article were rare, we extended the research to articles estimating the level of ID in

polyploid populations, even without diploid controls.

We used Google Scholar, Web of Science, PubMed and Agricola databases in order to

perform our literature survey. We used the keywords ("neopolyploid*" or "synthetic

polyploid*" or "polyploid*") and ("inbreeding", "inbreeding depression", "fitness"), and

"plant*". To be incorporated in the data collection, the selected study had to (1) define clearly

the type (allo- or autopolyploidy) and level of ploidy of the population under study, (2) give

the level of ID in the populations under study, or at least give the fitness of outbred and inbred

progenies, such that we were able to infer ID ourselves, and (3) that ID equals $\delta = 1-(W_s/W_o)$

when $W_s < W_o$ and $\delta = (W_o/W_s)-1$ when $W_s > W_o$, where W_s and W_o are respectively the

fitness of selfed and outcrossed progenies [12]. Most of the time, the studies reported the

numerical values in tables, but we sometimes had to extract the data directly from the figures,

by using Plot Digitizer [27]. In the following part of the manuscript, we assumed that

estimates found in synthetic polyploids (called neopolyploids in the following parts of the

manuscript) will be our proxies for the short-term consequences of polyploidization on ID,

while estimates from natural populations will be our proxies for the long-term consequences,

even if the time since polyploidization is generally unknown and can greatly differ between

species. A summary of the sampled species and articles can be found in Table 1.

Considering phylogenetic non-independence

We tested if there was a potential phylogenetic correlation of our estimates in our

analyses. To do so, we used the divergence times found in the TimeTree's database [28], in

order to reconstruct the phylogeny of our selected species. We then used the MEGA-X

software [29] to transform the obtained matrix of distance into NEWICK format, with an

UPGMA method. During this process, seven species were not found. We used the TimeTree

database to find closely related species used for replacement of the missing ones (see Figure

S1 and associated text to have the list of species). To test for a potential phylogenetic non-

independence, we used the obtained phylogeny and ran linear mixed effect models by using

the 'metafor' R package [30], and more precisely the rma.mv function that allows integrating

phylogenetic correlation matrix into linear models. We tested if the amount of ID differed

among diploid, neo- and natural polyploids using nested models. A first one in which a

random-effect 'Species' is specified, and a second in which we included the phylogenetic

correlation matrix as an additional random effect. The significance of the phylogenetic matrix

was tested by performing a Likelihood Ratio Test (LRT) between the two abovementioned

models.

Inbreeding depression

We chose to perform a Bayesian meta-analysis, by using the MCMCglmm package

[31]. We wrote the following model:

$$\delta_{ij} = \mu + ploidy_i \, \text{species}_j + \varepsilon_{ij} \, (1)$$

where δ_{ij} is the level of ID, μ is the mean value, and *ploidy $_i$* is the effect of the ploidy i

(diploid, neopolyploid [synthetic polyploid], and established tetraploid). We only included a

single random effect: species $_j$ is the effect of the *j*th species, and ε_{ij} is the residual error. We

assumed that the residual error followed a Gaussian distribution. We performed these models

with two different datasets. The first one in which restricted to studies in which ID is

estimated in diploid and/or neo- and natural polyploids simultaneously (called relatedness-

controlled analysis after). In this dataset, we subtracted the ID level of diploids from the

values of polyploids ($\Delta\delta = \delta_{\text{poly}} - \delta_{\text{diplo}}$). If $\Delta\delta < 0$ (respectively > 0), it means that δ is smaller

(respectively higher) in polyploids compared to diploids. In a second analysis, we included all

studies and compared untransformed values of δ for diploids and polyploids (called complete

analysis after). This analysis is potentially less robust, because we are comparing unrelated

diploid and polyploid species, that can differ for other life-history traits potentially affecting

ID levels.

For all analyses, we used the weakly informative, default priors proposed in

MCMCglmm [31]. For fixed effects, the prior is a normal distribution with mean being equal

to zero and a variance of 10^{10}. For random effects, inverse-Wishart priors are implemented,

with the degree of belief parameter being equal to zero and the expected variance being equal

to 1. For all models, we used a burn period of 1.000.000 iterations, with a thinning interval of

50, and the MCMC chains were run for 6.000.000 iterations in total. The parameter models

and associated 95% credible intervals were thus inferred from the sampling of the posterior

distribution 100.000 times. We did a visual examination of the convergence, posterior traces

and autocorrelation values of our models, as suggested in [31]. The trace of the sampled

posterior, and posterior distributions for both models are available in supplementary materials

(Figures S2 and S3).

Results:

The dataset was composed of 33 articles published between 1940 and 2020. These

articles covered 25 species divided into 15 families of angiosperms. The relatedness-

controlled analysis was based on 99 diploid-polyploids estimates of ID (70 in natural

polyploid populations, 29 in synthetic polyploids). The complete analysis compiled 225

estimates of ID (195 in natural polyploid populations, 30 in synthetic polyploids). Most of the

estimates have been estimated for autopolyploid species (187 in auto- and 23 in allopolyploid

species). A summary can be found in Table 2.

We found that the phylogenetic matrix did not improve the model ($\chi^2 = 0.171$, d.f. = 1,

$p = 0.680$), so we decided to only keep the random effect 'Species'. We found no differences

between allo- and autopolyploid species (Tables S1, S2 & S3).

On average, polyploidy tended to decrease the amount of ID (Figure 1), with synthetic

polyploids having the smallest mean ID level, and natural polyploids being intermediate

between synthetic polyploids and diploids (Figure 1). In the relatedness-controlled analysis,

synthetic polyploids had a significantly smaller level of ID than their diploid relatives (Figure

1A), but natural polyploids had an intermediate level, not significantly different from

synthetic polyploids or diploids (Figure 1A). In the complete analysis, the amount of ID found

in our restricted set of diploid estimates was in line with what was found in bigger studies

(0.42 in [32], 0.38 [95% credible interval 0.27-0.60] in this study), confirming that we can use

this value for comparisons. We found that all cytotypes had significantly different amounts of

ID (Figure 1B). The synthetic polyploids had the lowest amount (Figure 1B, with a decrease

of 69.0% compared to diploids), while natural polyploids were intermediate between the two

other cytotypes (Figure 1B, with a decrease of 30.7% compared to diploids and an increase of

126.8% compared to synthetic polyploids).

Discussion:

Inbreeding depression and establishment of new polyploid lineages

In this study, we found that ID decreases in polyploid populations compared to

diploids. This result increases our understanding of how neopolyploid lineages can establish.

Our meta-analysis confirmed theoretical expectations that due to an initial bottleneck [15], the

masking of deleterious mutations that are in comparable frequencies as in diploids [20,33],

and/or a slower increase in homozygosity during selfing events [19], new polyploid lineages

beneficiate from a strong decrease in ID. Indeed, the small average amount of ID ($\delta = 0.119$)

found in synthetic polyploids suggests that potential bi-parental inbreeding should have a

minor effect on the establishment probability of neopolyploids [10,11]. In the long-term,

however, the amount of ID increases again in natural polyploids, as expected due to the

increase in frequency of recessive deleterious mutations because of their better masking in

polyploids compared to diploids[15,22]. Nevertheless, our results cannot conclude if natural

polyploids have an intermediate or similar amount of ID to their diploid progenitors.

The joint evolution of polyploidy, range expansion and mating system

Our results also give insights into why polyploidization can also lead to the evolution

of other life-history traits. It has been observed that polyploid lineages can lead to the

expansion of geographic [34] and climatic niches [35,36]. A theoretical argument is that a

reduction in ID may favor such expansions, as biparental inbreeding, expected during the

process due to bottleneck events [14], should have a minor effect in (neo)polyploids

compared to diploids [11]. Our results support the theoretical prediction.

Finally, our results indicate that polyploidization could favor the transition from

predominantly outcrossing to predominantly selfing mating systems. If an association

between polyploidy and higher selfing rates has been found [39], it has been shown that this

effect depends on the kind of ploidy. Husband *et al*. [17] showed that autopolyploid species

generally self-fertilize less than diploid ones, while allopolyploids showed the opposite

pattern. Since our dataset is mainly composed of autopolyploid species (Table 1), our finding

seems to be counterintuitive. Nevertheless, recent theoretical advances showed selfing only

promotes autopolyploidization when neopolyploid lineages are at least as fit as their diploid

counterparts [16], which is generally not the case [8,40].

Our results however suggest that geographic expansion and/or the evolution of selfing

are more likely to occur in the very first generations following genome doubling, as ID is

smaller in synthetic polyploids than in natural ones.

Potential limitations

Even if informative, our study could suffer from potential bias. The first one is that

synthetically produced polyploids could not be representative of natural young polyploid

lineages. Nevertheless, the chemical treatments used the generate synthetic polyploids

generally lead to a high to moderate death rate of treated diploids (see for example [41]),

mimicking the expected bottleneck of genetic diversity that occurs during polyploidization

events [15]. A second bias could be that the sampled diploid and natural polyploid lineages

could differ in their selfing rate, which could be problematic if polyploids are more often

predominantly selfers compared to diploids, as the observed decrease in ID could be due to an

increased selfing rate that purges more efficiently deleterious mutations in natural polyploids

compared to diploids [22]. Nevertheless, quick comparisons of the selfing rate found in our

diploid and natural polyploid species suggest no differences (Table S4), and our dataset is

mainly composed of autopolyploid species, that tend to have a smaller selfing rate than their

diploid progenitors on average [17]. These results suggest that our analysis is conservative,

and that the observed decrease in ID is due to the consequences of polyploidization *per se*.

Finally, it is generally assumed that ID will affect the survival probability of polyploid

populations, but such an association remains to be tested empirically.

Conclusions:

Our results are of primary importance for the understanding of polyploid

establishment, and for the joint evolution of polyploidy with other life-history traits such as

mating systems. However, our results remain preliminary, and further studies are needed to

confirm the patterns described. Especially, theoretical and empirical studies in allopolyploid

species would be of primary importance since there are few.

Acknowledgement: We thank the associate editor and anonymous reviewers for helpful

comments.

Funding statement: We thank the European Research Council (project 850852 DOUBLEADAPT), the Czech Science Foundation (project 20-22783S), and the Czech Academy of Sciences (project RVO 67985939).

Data: Data and Rcode are available on GitHub (https://github.com/JosselinCLO/Meta_analysis_Inbreeding_depression).

References:

1. Gregory TR, Mable BK. 2005 Polyploidy in animals. In *The evolution of the genome*, pp. 427–517. Elsevier.

2. Leebens-Mack JH *et al.* 2019 One thousand plant transcriptomes and the phylogenomics of green plants.

3. Van de Peer Y, Mizrachi E, Marchal K. 2017 The evolutionary significance of polyploidy. *Nature Reviews Genetics* **18**, 411.

4. Levin DA. 1975 Minority cytotype exclusion in local plant populations. *Taxon* **24**, 35–43.

5. Comai L. 2005 The advantages and disadvantages of being polyploid. *Nature Reviews Genetics* **6**, 836–846. (doi:10.1038/nrg1711)

6. Otto SP. 2007 The evolutionary consequences of polyploidy. *Cell* **131**, 452–462.

7. Doyle JJ, Coate JE. 2019 Polyploidy, the nucleotype, and novelty: the impact of genome doubling on the biology of the cell. *International Journal of Plant Sciences* **180**, 1–52.

8. Clo J, Kolář F. 2021 Short- and long-term consequences of genome doubling: a meta-analysis. *American Journal of Botany* **108**, 2315–2322. (doi:10.1002/ajb2.1759)

9. Gemble S *et al.* 2022 Genetic instability from a single S phase after whole-genome duplication. *Nature* **604**, 146–151.

10.   Rausch JH, Morgan MT. 2005 THE EFFECT OF SELF-FERTILIZATION, INBREEDING DEPRESSION, AND POPULATION SIZE ON AUTOPOLYPLOID ESTABLISHMEN. *Evolution* **59**, 1867–1875.

11.   Griswold CK. 2021 The effects of migration load, selfing, inbreeding depression, and the genetics of adaptation on autotetraploid versus diploid establishment in peripheral habitats. *Evolution* **75**, 39–55.

12.   Charlesworth D, Charlesworth B. 1987 Inbreeding depression and its evolutionary
288   consequences. *Annual review of ecology and systematics* , 237–268.

289   13.   Charlesworth D, Willis JH. 2009 The genetics of inbreeding depression. *Nature reviews*
290   *genetics* **10**, 783–796.

291   14.   Rosche C, Hensen I, Mraz P, Durka W, Hartmann M, Lachmuth S. 2017 Invasion success in
292   polyploids: the role of inbreeding in the contrasting colonization abilities of diploid versus
293   tetraploid populations of Centaurea stoebe sl. *journal of Ecology* **105**, 425–435.

294   15.   Layman NC, Busch JW. 2018 Bottlenecks and inbreeding depression in autotetraploids.
295   *Evolution* **72**, 2025–2037. (doi:10.1111/evo.13587)

296   16.   Clo J, Padilla-García N, Kolář F. 2022 Polyploidization as an opportunistic mutation: The role
297   of unreduced gametes formation and genetic drift in polyploid establishment. *Journal of*
298   *Evolutionary Biology* **35**, 1099–1109.

299   17.   Husband BC, Ozimec B, Martin SL, Pollock L. 2008 Mating consequences of polyploid
300   evolution in flowering plants: current trends and insights from synthetic polyploids. *International*
301   *Journal of Plant Sciences* **169**, 195–206.

302   18.   Soltis PS, Soltis DE. 2000 The role of genetic and genomic attributes in the success of
303   polyploids. *Proceedings of the National Academy of Sciences* **97**, 7051–7057.

304   19.   Bever JD, Felber F. 1992 The theoretical population genetics of autopolyploidy. *Oxford*
305   *surveys in evolutionary biology* **8**, 185–185.

306   20.   Clo J. 2022 The evolution of the additive variance of a trait under stabilizing selection after
307   autopolyploidization. *Journal of Evolutionary Biology* **35**, 891–897. (doi:10.1111/jeb.14010)

308   21.   Lande R, Schemske DW. 1985 The evolution of self-fertilization and inbreeding depression in
309   plants. I. Genetic models. *Evolution* **39**, 24–40.

310   22.   Ronfort J. 1999 The mutation load under tetrasomic inheritance and its consequences for the
311   evolution of the selfing rate in autotetraploid species. *Genetics Research* **74**, 31–42.

312   23.   Siopa C, Dias MC, Castro M, Loureiro J, Castro S. 2020 Is selfing a reproductive assurance
313   promoting polyploid establishment? Reduced fitness, leaky self-incompatibility and lower
314   inbreeding depression in neotetraploids. *American Journal of Botany* **107**, 526–538.
315   (doi:10.1002/ajb2.1441)

316   24.   Busbice TH, Wilsie CP. 1966 Inbreeding depression and heterosis in autotetraploids with
317   application to Medicago sativa L. *Euphytica* **15**, 52–67.

318   25.   Dewey DR. 1966 Inbreeding Depression in Diploid, Tetraploid, and Hexaploid Crested
319   Wheatgrass 1. *Crop Science* **6**, 144–147.

320   26.   Hedrick PW. 1987 Genetic load and the mating system in homosporous ferns. *Evolution* **41**,
321   1282–1289.

322   27.   Huwaldt JA, Steinhorst S. 2015 Plot Digitizer, version 2.6. 8. *Computer Software]. Retrieved*
323   *from https://sourceforge. net/projects/plotdigitizer*

324   28.   Kumar S, Stecher G, Suleski M, Hedges SB. 2017 TimeTree: a resource for timelines,
325   timetrees, and divergence times. *Molecular biology and evolution* **34**, 1812–1819.

326   29.   Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018 MEGA X: molecular evolutionary genetics
327   analysis across computing platforms. *Molecular biology and evolution* **35**, 1547.

328   30.   Viechtbauer W. 2010 Conducting meta-analyses in R with the metafor package. *Journal of*
329   *statistical software* **36**, 1–48.

330   31.   Hadfield JD. 2010 MCMC methods for multi-response generalized linear mixed models: the
331   MCMCglmm R package. *Journal of statistical software* **33**, 1–22.

332   32.   Winn AA *et al.* 2011 Analysis of inbreeding depression in mixed-mating plants provides
333   evidence for selective interference and stable mixed mating. *Evolution: International Journal of*
334   *Organic Evolution* **65**, 3339–3359.

335   33.   Otto SP, Whitton J. 2000 Polyploid incidence and evolution. *Annual review of genetics* **34**,
336   401–437.

337   34.   Te Beest M, Le Roux JJ, Richardson DM, Brysting AK, Suda J, Kubešová M, Pyšek P. 2012 The
338   more the better? The role of polyploidy in facilitating plant invasions. *Annals of botany* **109**, 19–
339   45.

340   35.   Padilla-García N *et al.* 2022 Niche differentiation following whole-genome duplication? The
341   importance of considering the intricate evolutionary history of polyploids when assessing climatic
342   niche evolution. *bioRxiv* , 2022.02.16.480351. (doi:10.1101/2022.02.16.480351)

343   36.   Glennon KL, Ritchie ME, Segraves KA. 2014 Evidence for shared broad-scale climatic niches of
344   diploid and polyploid plants. *Ecology Letters* **17**, 574–582. (doi:10.1111/ele.12259)

345   37.   Dubcovsky J, Dvorak J. 2007 Genome plasticity a key factor in the success of polyploid wheat
346   under domestication. *Science* **316**, 1862–1866.

347   38.   Fawcett JA, Maere S, Van De Peer Y. 2009 Plants with double genomes might have had a
348   better chance to survive the Cretaceous–Tertiary extinction event. *Proceedings of the National*
349   *Academy of Sciences* **106**, 5737–5742.

350   39.   Barringer BC. 2007 Polyploidy and self-fertilization in flowering plants. *American Journal of*
351   *Botany* **94**, 1527–1533.

352   40.   Porturas LD, Anneberg TJ, Curé AE, Wang S, Althoff DM, Segraves KA. 2019 A meta-analysis of
353   whole genome duplication and the effects on flowering traits in plants. *American journal of*
354   *botany* **106**, 469–476.

355   41.   Julião SA *et al.* 2020 Induction of synthetic polyploids and assessment of genomic stability in
356   Lippia alba. *Frontiers in plant science* **11**, 292.

357   42.   Le S, Griffin RA, Harwood CE, Vaillancourt RE, Harbard JL, Price A, Nghiem CQ, Koutoulis A,
358   Nguyen KD. 2021 Breeding Polyploid Varieties of Acacia: Reproductive and Early Growth
359   Characteristics of the Allotetraploid Hybrid (Acacia mangium$\times$ A. auriculiformis) in
360   Comparison with Diploid Progenitors. *Forests* **12**, 778.

43.   Dewey DR. 1969 Inbreeding Depression in Diploid and Induced-Autotetraploid Crested
Wheatgrass 1. *Crop Science* **9**, 592–595.

44.   Johnston MO, Schoen DJ. 1996 Correlated evolution of self-fertilization and inbreeding
depression: an experimental study of nine populations of Amsinckia (Boraginaceae). *Evolution* **50**,
1478–1491.

45.   Rosquist G. 2001 Reproductive biology in diploid Anthericum ramosum and tetraploid A.
liliago (Anthericaceae). *Oikos* **92**, 143–152.

46.   Inoue K, Masuda M, Maki M. 1998 Brief communication. Inbreeding depression and
outcrossing rate in the endangered autotetraploid plant Aster kantoensis (Asteraceae). *Journal of*
Heredity **89**, 559–562.

47.   Hecker RJ. 1972 Inbreeding depression in diploid and autotetraploid sugarbeet, Beta vulgaris
L. *Euphytica* **21**, 106–111.

48.   Galloway LF, Etterson JR, Hamrick JL. 2003 Outcrossing rate and inbreeding depression in the
herbaceous autotetraploid, Campanula americana. *Heredity* **90**, 308–315.

49.   Galloway LF, Etterson JR. 2007 Inbreeding depression in an autotetraploid herb: a three
cohort field study. *New Phytologist* **173**, 383–392.

50.   Husband BC. 2016 Effect of inbreeding on pollen tube growth in diploid and tetraploid
Chamerion angustifolium: Do polyploids mask mutational load in pollen? *American journal of*
botany **103**, 532–540.

51.   Husband BC, Schemske DW. 1996 Evolution of the magnitude and timing of inbreeding
depression in plants. *Evolution* **50**, 54–70.

52.   Ozimec BC. 2006 Inbreeding depression and mating system evolution in the autotetraploid
Chamerion angustifolium. PhD Thesis, University of Guelph.

53.   Barringer BC, Geber MA. 2008 Mating system and ploidy influence levels of inbreeding
depression in Clarkia (Onagraceae). *Evolution: International Journal of Organic Evolution* **62**, 1040–
1051.

54.   Grindeland JM. 2008 Inbreeding depression and outbreeding depression in Digitalis
purpurea: optimal outcrossing distance in a tetraploid. *Journal of Evolutionary Biology* **21**, 716–
726.

55.   Kwok A. 2013 The role of polyploidy in the evolution of gender dimorphism: an experimental
approach using Fragaria vesca. PhD Thesis, University of Guelph.

56.   Wheelwright NT, Begin E, Ellwanger C, Taylor SH, Stone JL. 2016 Minimal loss of genetic
diversity and no inbreeding depression in blueflag iris (Iris versicolor) on islands in the Bay of
Fundy. *Botany* **94**, 543–554.

57.   Vange V. 2002 Breeding system and inbreeding depression in the clonal plant species Knautia
arvensis (Dipsacaceae): implications for survival in abandoned grassland. *Biological conservation*
108, 59–67.

58.   Wilsie CP. 1958 Effect of Inbreeding on Fertility and Vigor of Alfalfa 1. *Agronomy Journal* **50**,
182–185.

59.   Posler GL, Wilsie CP, Atkins RE. 1972 Inbreeding Medicago sativa L. by Selfing, Sib-mating,
and Intergenerational Crossing 1. *Crop Science* **12**, 49–52.

60.   Pujol B, Zhou S-R, Sanchez Vilas J, Pannell JR. 2009 Reduced inbreeding depression after
species range expansion. *Proceedings of the National Academy of Sciences* **106**, 15379–15383.

61.   Eppley SM, Pannell JR. 2009 Inbreeding depression in dioecious populations of the plant
Mercurialis annua: comparisons between outcrossed progeny and the progeny of self-fertilized
feminized males. *Heredity* **102**, 600–608.

62.   Crane MB, Lewis D. 1942 Genetical studies in pears. *Journal of Genetics* **43**, 31–43.

63.   Nielsen LR, Siegismund HR, Hansen T. 2007 Inbreeding depression in the partially self-
incompatible endemic plant species Scalesia affinis (Asteraceae) from Galápagos islands.
Evolutionary Ecology **21**, 1–12.

64.   Dudash MR, Fenster CB. 2001 The role of breeding system and inbreeding depression in the
maintenance of an outcrossing mating strategy in Silene virginica (Caryophyllaceae). *American*
Journal of Botany **88**, 1953–1959.

65.   Daehler CC. 1998 Variation in self-fertility and the reproductive advantage of self-fertility for
an invading plant (Spartina alterniflora). *Evolutionary Ecology* **12**, 553–568.

66.   Armstrong JM, Robertson RW. 1956 Studies of Colchicine-Induced Tetraploids of Trifolium
Hybridum L.: I. Cross and Self-Fertility and Cytological Observations. *Canadian Journal of*
Agricultural Science **36**, 255–266.

67.   Hokanson K, Hancock J. 2000 Early-acting inbreeding depression in three species of
Vaccinium (Ericaceae). *Sexual Plant Reproduction* **13**, 145–150.

429 **Table 1.** Summary of the sampled species, their ploidy levels, their genomic origin (auto- or

430 allopolyploid), if they are natural or synthetical polyploids.

Species	Ploidy levels	Genomic origin	Natural / synthetic	References
Acacia auriculiformis	2x, 4x	autopolyploid	natural & synthetic	[40]
Agropyron cristatum	2x, 4x	autopolyploid	natural & synthetic	[26,41]
Amsinckia gloriosa	2x, 4x	autopolyploid	Natural	[42]
Anthericum liliago	2x, 4x	allopolyploid	natural	[43]
Aster kantoensis	4x	autopolyploid	natural	[44]
Beta vulgaris	4x	autopolyploid	natural	[45]
Campanula americana	4x	autopolyploid	natural	[46,47]
Centaurea stoebe	2x, 4x	autopolyploid	natural	[13]
Chamerion angustifolium	2x, 4x	autopolyploid	natural & synthetic	[17,48–50]
Clarkia davyi	2x, 4x	allopolyploid	natural	[51]
Clarkia gracilis	2x, 4x	allopolyploid	natural	[51]
Digitalis purpurea	4x	autopolyploid	natural	[52]
Fragaria vesca	2x, 4x	autopolyploid	synthetic	[53]
Iris versicolor	4x	allopolyploid	natural	[54]
Jasione maritima	2x, 4x	autopolyploid	synthetic	[24]
Knautia arvensis	4x	autopolyploid	natural	[55]
Medicago sativa	4x	autopolyploid	natural	[25,56,57]
Mercurialis annua	6x	autopolyploid	natural	[58,59]
Pyrus communis	4x	autopolyploid	natural	[60]
Scalesia affinis	4x	autopolyploid	natural	[61]
Silene virginica	4x	autopolyploid	natural	[62]
Spartina alterniflora	4x	allopolyploid	natural	[63]
Trifolium hybridum	4x	autopolyploid	synthetic	[64]
Vaccinium corymbosum	2x, 4x	autopolyploid	natural	[65]

432   **Table 2.** Summary of the number of estimates for the different categories of ploidy, for the

433   relatedness-controlled and complete analyses.

Dataset	Diploids	Neo-polyploids	Natural polyploids
Relatedness-controlled	84	29	70
Complete	84	30	195

444   **Figure 1.** The evolution of inbreeding depression in polyploids (neo- and natural polyploids)

445   compared to their diploid progenitors. **A.** Relatedness-controlled analysis, with $\Delta\delta = \delta_{poly} -$

446   δ_{diplo}, a negative value showing that inbreeding depression is smaller in polyploids compared

447   to diploids. **B.** Complete analysis. Different letters indicate significantly different mean

448   values. Error bars stand for the 95% credibility intervals.