# The lectin receptor Clec4F participates in the capture of circulating parasite mucins by Kupffer cells in cystic echinococcosis.

**Anabella A. Barrios[1], Camila Mouhape[1], Leonard Schreiber[2], Juliane Nell[2], Mariana Suárez-Martins[1], Geraldine Schlapp[3], María Noel Meikle[3], Ana Paula Mulet[3], Tsui-Ling Hsu[4], Shie-Liang Hsieh[4], Gustavo Mourglia-Ettlin[1], Carlos González[5], Martina Crispo[3], Thomas F. E. Barth[2], Cecilia Casaravilla[1], Stephen J. Jenkins[6] & Álvaro Díaz[1*]**

1. Área Inmunología, Departamento de Biociencias (Facultad de Química) and Cátedra de Inmunología, Instituto de Química Biológica (Facultad de Ciencias), Universidad de la República, Montevideo, Uruguay.

2. Institute of Pathology, University Ulm, Ulm, Germany

3. Unidad de Biotecnología en Animales de Laboratorio, Institut Pasteur de Montevideo, Montevideo, Uruguay.

4. Genomics Research Center, Academia Sinica, Taipei, Taiwan.

5. Retired from: CIMMYT (Centro Internacional de Mejoramiento de Maíz y Trigo), El Batán, México.

6. University of Edinburgh Centre for Inflammation Research, Queen's Medical Research Institute, University of Edinburgh, Edinburgh, UK.

(*) Corresponding author. E-mail: adiaz@fq.edu.uy

**Keywords:** Clec4F, *Echinococcus*, laminated layer, lectin, mucin, Kupffer cells

**Short title:** Kupffer cells capture parasite mucins through Clec4F

## Author contributions

AB: conceptualization, investigation. CM, LS, JN, MS: investigation. GS, MNM, APM, MC: methodology. TLH, SLH, GME: resources. CG: formal analysis (statistics). TFEB, CC: conceptualization, supervision. SJJ: conceptualization, supervision. writing – review & editing. AD. conceptualization, funding acquisition, project administration, supervision, writing – original draft preparation.

**Abstract**

Cystic echinococcosis is caused by the larval stages (hydatids) of cestode parasites

belonging to the species cluster *Echinococcus granulosus sensu lato*. Hydatids are

bladder-like structures that attain large sizes within various internal organs of livestock

ungulates and humans. Hydatids are protected by the massive acellular laminated layer

(LL), composed mainly by mucins. Parasite growth requires LL turnover, and abundant

LL-derived particles are found at infection sites and draining lymph nodes in infected

humans. These observations raise the question of whether LL materials circulate

systemically, and if so how they are dealt with by the hosts. We previously reported that

LL mucins are bound by a recombinant version of the lectin receptor Clec4F. In

rodents, Clec4F is expressed only in Kupffer cells, the liver macrophages exposed to the

vascular space. In this article, we show that Kupffer cells take up LL mucins *in vivo* in a

manner dependent on the mucin glycans and Clec4F. In mice harbouring intraperitoneal

infections, LL mucins were found essentially only at the infection site and in the liver,

where they were taken up by Kupffer cells via Clec4F. It seems likely that the

composition of the LL was adjusted through evolution so that LL debris were taken up

by the major phagocytes of the rodent liver, which is the ancestral infection site for

*Echinococcus* larvae. However, our data reveal that Kupffer cells act as a sink for LL

materials even when the parasite grows in sites other than the liver, as it happens in

natural *E. granulosus* infections.

**Author summary**

Cystic echinococcosis (hydatid disease) is a major neglected tropical disease affecting

both humans and livestock. It is caused certain tapeworm larvae, which develop into

massive bladder-like structures within the liver, lungs or other organs. These larvae are

protected by a thick, sugar-rich coat called the laminated layer (LL). As the parasite

grows, the LL releases its outermost (older) sub-layers as particles, which can

accumulate in the tissues of the infected person or animal. We have addressed the issue

of what is the destiny within the body of these abundant parasite debris. Using mice as a

model, we determined that LL debris are selectively taken up by liver phagocytic cells

known as Kupffer cells, known to clear unwanted components from the blood. More

specifically, we show that Kupffer cells take up the LL debris via a sugar-binding

protein called Clec4F. This protein had previously been implicated in the clearance of

aged platelets from the blood. It makes sense from an evolutionary point of view that a

parasite directs its debris to a host system that normally clears "self" products, as

opposed to host systems dedicated to detecting pathogens and alerting the immune

system of their presence.

**Introduction**

The larval stages of the cestode parasites belonging to the species cluster known as

*Echinococcus granulosus sensu lato* (distinguishable only by molecular means) cause

cystic echinococcosis in livestock animals and humans (1,2). This disease is

characterized by the growth within organ parenchymae of bladder-like parasite

structures called hydatids. Each hydatid is bounded by a thin cellular layer, in turn

protected by a thick acellular coat called the laminated layer (LL) (3–6). The main

component of the LL is a meshwork of mucins, heavily decorated by O-glycans that are

rich in galactose (7,8). Hydatids persist in internal organs of immunocompetent hosts

for up to decades, and they usually elicit weak local inflammation (9). The growth of

the hydatid (potentially through over 3 orders of magnitude in lineal dimension)

requires the shedding of materials from the LL (3–6). In human infections, the abundant

shed LL particles (termed SPEGs, for shed particles of *E. granulosus*) are found in the

tissue surrounding the parasite and in the draining lymph nodes (10).

The persistence of larval *E. granulosus* infection, the massive nature of the LL, and

sheer abundance of shed particles as detected in human infection call the question of

how the host's body disposes of this foreign debris and avoids the potential

inflammation that may result from its build-up. As inflammation is detrimental to the

parasite (9), it makes biological sense that LL constituents may be evolutionarily

perfected for engaging cellular receptors that will ensure non-inflammatory disposal of

the shed materials.

Studies on the responses elicited in mouse dendritic cells (DC) and macrophages by *in*

*vitro*-produced LL particles (pLL, from particulate LL) have suggested an interaction

mode independent of conventional receptors and inefficient phagocytosis, at least in the

absence of opsonization (6,11,12). In broad agreement, none out of 35 human innate

carbohydrate-recognition proteins (recombinantly expressed as fusions with IgG Fc)

bound the LL mucins in an *in vitro* screening (13). However, in the same experiment,

strong binding was observed for mouse Clec4F. This type II, trimeric, cell-surface C-

type lectin has affinity for glycans terminated in N-acetylgalactosamine or galactose

(Gal), especially when they occur in tandem along the same backbone, as it happens in

mucins (14–18). The binding of recombinant Clec4F to the LL mucins was as expected

calcium-dependent and resistant to digestion of non-mucin proteins in the parasite

preparation. Binding was also observed to a number of synthetic glycans representing

some of the main carbohydrate motifs in the *E. granulosus* LL and/or in the LL of the

related species *Echinococcus multilocularis* (13).

In rodents, Clec4F is exclusively expressed in Kupffer cells (KC) (14–16), the sessile

liver macrophages exposed to the vascular space. This restricted expression means that

Clec4F is not expressed in the mouse DC and macrophage sub-types used in the

previously mentioned experiments that suggested a receptor-independent interaction

mode with LL particles (6,11,12). In the ungulate natural hosts of *E. granulosus*, Clec4F

appears to be expressed in lymphoid tissues in addition to the liver (15).

The only demonstrated function of Clec4F (in mice) is the removal from circulation of

de-sialylated platelets by KC (19). Although the macrophage galactose-type lectins

(MGL) and the asialoglycoprotein receptor (ASGR; reported to be expressed in KC in

addition to in hepatocytes) collaborate in this function, Clec4F appears to fulfil an indispensable role (19–21).

KC under basal conditions have a tolerogenic profile as antigen-presenting cells (22). Also, a role of Clec4F in removing de-sialylated platelets (19), conceptually akin to the role of the receptors involved in phagocytosis of apoptotic cells (23), suggests that Clec4F is not wired to inflammatory responses. Thus, Clec4F/KC appears to be an evolutionary option from the parasite's standpoint for LL debris disposal. However, *in vitro* binding by the soluble Clec4F lectin domain expressed in dimeric form (13) does not predict binding by the native cell-surface trimeric receptor, or internalization by KC. In this article, we analyse *in vivo* uptake of soluble and particulate LL materials by KC and the role of Clec4F in this process, in the context of artificial injection of materials as well as experimental infection.

**Results**

*Soluble LL mucins are taken up by Kupffer cells* in vivo *in a manner dependent on the mucin glycans*

We first sought to determine whether KC take up soluble LL mucins (sLL) *in vivo* and whether this is dependent on the mucin glycans. We thus treated the LL mucins either with sodium periodate to oxidize the glycans or applied a mock treatment, tagged both preparations with equivalent amounts of biotin (Figure S1), and injected both preparations i.v. in mice. Twenty minutes later, we quantitated the biotin tag in permeabilized non-parenchymal liver cells using streptavidin conjugated to PE and flow

cytometry. KC were identified using our previously validated gating strategy (Figure

S2) (24). A significant signal was detected in KC, and the signal was markedly higher

for mock-treated than for periodate-treated sLL (Figure 1). No significant biotin signal

was detected in either LSECs or in non-KC CD11b$^+$ cells, a broad definition that

encompasses any mononuclear phagocytes other than KC present in the liver (Figures 1

a, b, and S2).

*The plant lectin PNA and the E492 monoclonal antibody allow detection of laminated*

*layer mucins captured by liver cells* in vivo.

Except in experiments using periodate-treated LL materials, the detection of LL

materials by appropriate carbohydrate-binding probes was preferable to prior labelling

with biotin. Among other advantages, detection with such probes would allow the

detection of endogenous LL mucins in infected animals. One candidate probe was the

plant lectin PNA, which binds Galβ1-3-GalNAcα1-peptide, i.e. the non-decorated

mucin O-glycan core 1 or T-antigen (25). This motif is highly represented in the *E.*

*granulosus* LL, and accordingly, PNA binds the LL mucins in histological sections,

particle lectin fluorescence, and Western blotting (12,26). The second candidate probe

was the monoclonal antibody E492, raised against *E. granulosus* protoscoleces (27).

This antibody binds to LL mucins from *E. multilocularis* (28) and to synthetic glycans

that include the P$_1$ blood antigen motif (Galα1-4Galβ1-4GlcNAc), highly represented

among *E. granulosus* LL mucin glycans (7,8). A third potential probe was recombinant

Clec4F (13). We verified that these potential probes indeed bind to sLL coated on

ELISA plates as well as to LL particles in suspension (Figure 2 a, b).

We took advantage of the defined specificities of PNA and the E492 antibody to gain

insight into the carbohydrate motifs bound by Clec4F in the LL mucins. The binding of

recombinant Clec4F to the solubilized and plate-bound LL mucins was not competed by

PNA, but it was strongly competed by the E492 antibody (Figure 2 c). Therefore, most

of the binding of Clec4F to the native LL mucins appears to rely on recognition of the

$P_1$ motif, whereas recognition of the T-antigen is not significant.

*In vivo* probes should show minimal reactivity with host tissues; for our purposes, lack

of reactivity with the populations of mouse non-parenchymal liver cells under study was

the most important consideration. Neither PNA nor the E492 antibody showed

significant binding in flow cytometry to KC or LSEC from control mice (Figure S3).

PNA bound to a very minor fraction of non-KC CD11b$^+$ cells, whereas the antibody did

not bind. PNA, but not the E492 antibody, bound to a major sub-population of CD45$^+$

CD11b$^-$ F4/80$^-$cells, which were not of particular interest in our study. Recombinant

Clec4F bound to cells in all 4 cell gates defined, most strongly to LSEC and KC (Figure

S3). This probably reflects the fairly broad specificity of Clec4F for glycans terminated

in Gal/GalNAc (15–17), and ruled out this lectin as a probe for our purposes.

Finally, to determine whether uptake of sLL could be measured *in vivo* using PNA or

the E492 antibody, we injected wild-type (WT) C57BL/6J mice i.v. with sLL and

subsequently stained isolated hepatic non-parenchymal cells intracellularly with either

probe. As expected, no staining was observed in cells from mice injected with PBS

only, whereas KC, and to a small extent also LSEC, from mice given sLL stained

intracellularly with either PNA or E492 antibody (Figure 2 d). Furthermore, both probes

appeared to allow more sensitive detection than prior labelling with biotin (Figure 1), so

that captured sLL was detected in almost all KC and also in some LSEC.

In summary both PNA and the E492 antibodies are useful probes for LL mucins

captured by mouse non-parenchymal liver cells.

*Soluble LL mucins are taken up by Kupffer cells* in vivo *in a manner dependent on*

*Clec4F*

We first used Crispr/Cas9 to generate a new Clec4f $^{-/-}$ line on the C57BL/6J background.

These Clec4f $^{-/-}$ mice have 8-bp deletion expected to result in a polypeptide that is

truncated near the N-terminus (Figure 3, a, b), and effectively lack Clec4F protein

expression by flow cytometry (Figure 3 c). Hence, our line appears similar to Clec4f

gene-deficient mice that have already been described (16,19).

We next compared WT and Clec4f $^{-/-}$ mice in terms of the *in vivo* uptake of sLL by liver

non-parenchymal cells. For this purpose, we injected untagged sLL, either untreated or

subjected to the mock treatment used in the previous experiments that involved

periodate oxidation. Uptake of sLL by KC was strongly dependent on Clec4F (Figure 4

a - c), particularly when assessed in terms of mean fluorescence intensity. The low-level

uptake by LSEC observed was as expected independent of Clec4F; in fact, for untreated

sLL, uptake by LSEC was slightly enhanced in the absence of Clec4F, possibly as a

result of the decreased capture by KC. No uptake by non-KC CD11b[+] cells was detected

(Figure 4 a, b and Figure S4). Through a calculation that takes into account the numbers

of each cell type capturing sLL and the intensity of the signal indicating intracellular

sLL, KC were estimated to be responsible for the overwhelming majority of sLL uptake

by non-parenchymal liver cells in WT mice (Figure 4 d). However, in the absence of

Clec4F, LSEC played a significantly, if slightly, more important part in the uptake of

sLL than in the presence of the receptor. Differences in the uptake of untreated *vs* mock-

treated material were not observed, except in the low-level uptake by LSEC in Clec4f$^{-/-}$

mice (Figure 4 a). For consistency with the results shown in Figure 1, we chose to use

mock-treated material in the following experiments.

It can be expected that in the absence of Clec4F, other mechanisms of cellular uptake of

sLL may become apparent, e.g. macropinocytosis or uptake dependent on natural

antibodies or complement (6). Such Clec4F-independent uptake of sLL was hinted by

the previous observations in relation to LSEC, as well as by the 25 – 60 % KC in

Clec4f$^{-/-}$ mice positive for PNA at low levels (Figure 4 a - c). As the Clec4F-

independent mechanisms might become dominant at longer analysis times, we carried

out a similar experiment with a 22 h analysis time. Uptake of sLL was still dominated

by KC among the cell types studied, and this uptake was again strongly dependent on

Clec4F (Figure 5 a –d). In the absence of Clec4F, LSEC made a significantly larger

contribution to sLL uptake than in the presence of the receptor, but this contribution was

still minimal (Figure 5 d). Non-KC CD11b$^{+}$ cells did not take up material detectably

(Figure 5 a – d and Figure S4). Cell surface expression of Clec4F in KC was not altered

by sLL injection (Figure S5).

In summary, soluble circulating LL mucins in naïve mice are taken up mainly by KC

among liver non-parenchymal cells and mostly in Clec4F-dependent fashion.

*LL mucins injected in particulate form are taken up* by *Kupffer cells* in vivo *in Clec4F-*

*dependent manner*

Although the release of soluble mucins from the LL during cystic echinococcosis is a

possibility (6), only particles have been documented to be shed from the LL in human

infections (6,10). It is not known if these particles circulate systemically and/or give rise

to soluble materials that circulate. It was therefore important to determine whether

Clec4F participates in the uptake by KC of LL mucins released in a particulate

presentation into the internal milieu and at an anatomical site distant from the liver. We

thus injected i.p. particulate LL mucins (pLL) (12), and analysed the liver non-

parenchymal fraction 22 h later. LL material was indeed detected by flow cytometry in

KC (Figures 6 a – c and S6 a - c). Uptake by KC was dependent on Clec4F, although

there was a considerable degree of Clec4F-independent uptake as well. In addition to

low-level uptake by LSEC, in this experimental system, non-KC CD11b$^+$ cells were

observed to take up the LL material. Although non-KC CD11b$^+$ cells may include

recruited monocytes/macrophages, the frequency of this population (within all non-

parenchymal cells) did not increase as a result of pLL injection (Figure S7). As

expected, uptake by LSEC and by non-KC CD11b$^+$ cells was independent of Clec4F

and in fact tended to be enhanced in the absence of the receptor (Figures 6 a – c and S6

a - c). Indeed, whereas in WT mice the overwhelming majority of the LL material

uptake by liver non-parenchymal cells was carried out by KC, in Clec4f$^{-/-}$ mice the two

non-KC cell types studied taken together played a significantly larger role than in WT

mice (Figures 6 d and S6 d). The cell surface expression of Clec4F in KC was not

altered by injection of pLL (Figure S8).

272 The mostly Clec4F-dependent uptake of the LL material was confirmed by histology,

273 using the E492 antibody as a probe. Strong staining was observed in KC-like cells of

274 WT animals, whereas only minimal staining was observed in the KC-like cells of

275 Clec4f$^{-/-}$ animals (Figure 6 e). Animals injected with pLL also showed weak staining in

276 hepatocytes (a cell type not analysed by flow cytometry), and this was independent of

277 Clec4F. As hepatocytes are unlikely to phagocytose large particles, this observation

278 suggests the particles injected release small particles and/or soluble mucin materials *in*

279 *vivo* that preferentially circulate systemically.

281 In summary, LL mucins present within tissues in particulate form can generate materials

282 (particulate and/or soluble) that circulate systemically and that are captured by KC via

283 Clec4F.

285 *In experimental* E. granulosus *infection, LL mucins circulate and are captured*

286 *selectively by Kupffer cells, with the participation of Clec4F*

288 We next wished to determine if Clec4F was relevant in the infection setting. The most

289 widely used experimental infection model for *E. granulosus* uses intraperitoneal

290 injection with protoscoleces, parasite stages that usually develop into adult worms but

291 are capable of reverse development into hydatids (1,29). In this model, hydatids develop

292 free or within host collagen capsules in the peritoneal cavity (30). KC uptake of LL

293 materials in this model would require that such materials reach systemic circulation, a

294 possibility for which there was no evidence, in experimental or natural infections. We

295 first analysed infected mice of the BALB/c strain, in which experimental infection

296 reaches higher parasite burdens than in the C57BL/6 strain (31). Indeed, the BALB/c

mice analysed had a median of 25 mL of total parasite volume (Figure S9 a), an

extremely high parasite burden. In these mice, using the E492 antibody as probe, we

detected LL materials by histology in peritoneal cavity cells (i.e. the site of infection)

and the liver, but not in kidneys, intestine or lungs (Figure 7 a). In addition, in some of

the infected mice, occasional positive cells were observed in the spleen, and less

frequently in the mesenteric lymph nodes (Figure 7 a). The use of PNA as probe yielded

similar results, but it was complicated by the expected signal in the lymphoid organs

(32) unrelated to LL material, i.e. detectable in non-infected mice (data not shown).

The above results show that in infected animals LL mucins circulate systemically and

become selectively concentrated in the liver. Within the liver, the signal for LL mucins

was specifically concentrated in KC-like cells (Figure 7 a). The LL signal in KC-like

cells was observed in 12 out of 15 infected mice analysed. Contrary to previous

experiments injecting pLL, no signal was detected in hepatocytes. This may be related

to lower loads of LL material, as suggested by the lower intensity in the staining in KC

in the infection setting in comparison to the pLL injection experiments (compare

Figures 6 e and 7 a).

The selective LL material uptake by KC among the liver non-parenchymal cells was

confirmed by flow cytometry (Figures 7 b – d and S10). Although the data did not reach

statistical significance, a sub-population of non-KC CD11b$^+$ cells appeared to take up

LL materials (Figure 7 b – d). Also, the frequency of non-KC CD11b$^+$ cells within all

liver non parenchymal cells was significantly increased by infection, suggesting cell

recruitment (Figure S11 a). The cell surface expression of Clec4F in KC was not altered

by infection (Figure S12 a).

We next analysed infected C57BL/6 WT and Clec4f$^{-/-}$ mice. In these mice, the parasite

burdens were low: the median burden in the WT C57BL/6 mice was 0.5 mL (this figure

corresponds to the two infection experiments fully analysed, as an additional

experiment with very low parasite burdens was excluded from analysis). Although the

median parasite burden was 3-fold lower in Clec4f$^{-/-}$ mice than in WT mice, the trend

did not reach statistical significance (Figure S8 b). By flow cytometry, uptake of LL

material by KC was detected, in WT but not in Clec4F$^{-/-}$ mice (Figure 7 e - g). No

significant uptake was detected in LSECs (Figures 7 e, f and S10 b). In non-KC CD11b$^{+}$

cells, there was a trend towards low-level, Clec4F-independent uptake (Figure 7 e - g),

which did not reach significance. The frequency of non-KC CD11b$^{+}$ cells was not

significantly altered by infection (Fig S 11 b). Again, Clec4F surface expression in the

KC of WT mice was not altered by infection (Figure S12 b).

**Discussion**

In this article we show that native Clec4F is necessary for the internalization of *E.*

*granulosus* LL mucins by mouse KC *in vivo*, after injection of exogenous mucins as

well as in experimental infection. We also show that LL mucins initially present in

particulate format in the peritoneal cavity reach the liver, because they circulate as

particles and/or because they partly solubilize *in vivo*. The results also show that

parasites growing in the peritoneal cavity release LL materials that circulate

systemically and are preferentially captured by the liver, and within the liver, mostly by

KC.

In addition to KC, both LSEC and certain non-KC CD11b$^+$ cells were observed to take

up LL materials. This was in both cases independent of Clec4F, and in some

experimental settings it was enhanced in the absence of Clec4F (Figures 4 d, 6 d and S6

d). Uptake of materials by non-KC CD11b$^+$ cells was apparent after injection of LL

mucins in particulate presentation (pLL; Figures 6 a – d and S6) but not in a soluble

presentation (sLL; Figures 1, 4 and 5). This contrast suggests that the species released *in*

*vivo* from the large LL particles are (or at least include) small particles, as opposed to

fully soluble mucins like those making up sLL. It is unlikely that the non-KC CD11b$^+$

cell population taking up LL materials derives from peritoneal cavity macrophages, as it

is composed of F4/80$^-$ cells (Figures 6 c and S6 c), whereas most resident peritoneal

macrophages as well as peritoneal macrophages reported to enter the liver upon injury

to this organ are F4/80$^{hi}$ (33). We instead favour the possibility that the population

being discussed corresponds to monocytic cells present in the liver or recruited in

response to LL particles physically trapped in the liver sinusoids.

Hepatocytes could be an additional target cell for LL materials in the liver, as suggested

by the weak but significant staining observed in this cell type after pLL injection

(Figure 6 e); the obvious candidate receptor to mediate this apparent uptake is the

ASGR (34).

Leaving the experiments with sLL aside, the contribution of liver cell types other than

KC to take up the probably insoluble LL mucins appeared to depend on the load of

parasite material. Uptake by non-KC CD11b$^+$ cells was clear after injection of a high-

dose bolus of pLL (Figures 6 a – d and S6 a – d) and it was merely hinted in infection

(Figure 7 b - g). The uptake by hepatocytes and LSEC was observed after pLL injection

(Figures 6 and S6) but not at all in experimental infection, even high-burden (Figure 7).

Thus, a gradation appears to exist in which when the capacity of KC to take up LL

mucins is overwhelmed, non-KC CD11b$^+$ cells become important, followed by

hepatocytes and/or LSEC.

Part of the uptake of LL materials by KC was observed to be Clec4F-independent

(Figures 5 a, 6 a, b, and 7 e). In addition, there was as previously mentioned Clec4F-

independent uptake by non-KC CD11b$^+$ liver cells and by LSEC (Figures 4 a –c, 5 a - c,

6  a – c, and S6 a – c). Clec4F-independent uptake may be mediated by natural anti-

carbohydrate antibodies that bind the LL mucins, either directly or via complement

activation (35). In the context of chronic experimental infection, the possibilities for

Clec4F-independent uptake are even broader than after LL mucin injection. In the first

place, the mucins will be opsonized by specific (homologous) antibodies (6,35). In

second place, contrary to the artificially prepared (soluble and insoluble) mucin

preparations, LL materials shed from live parasites must contain the major non-mucin

structural component of the LL, i.e. calcium inositol hexakisphosphate (calcium Ins$P_6$)

(36–38). Calcium Ins$P_6$ binds host C1q and initiates limited activation of complement

(35,39). In fact, the observed uptake of LL materials by phagocytes at the experimental

infection site (Figure 7 a) must be Clec4F-independent, as peritoneal macrophages are

not expected to express Clec4F (16). Given the ample possibilities of Clec4F-

independent capture, it is noteworthy the Clec4F nonetheless has a clearly detectable

role in LL material uptake (by KC), especially in the infection context.

As mentioned, in human infections affecting solid organs, LL debris accumulate in the

parasite's vicinity (10). Our results show the presence of LL materials at the

experimental infection site but also as mentioned the capture of these materials by the

liver (Figure 7 a). This implies that the accumulation of LL debris in the parasite's

vicinity is moderated by the circulation and capture by the liver of materials derived

from this debris. This is important because, although moderate amounts of LL particles

probably have immune modulatory effects that favour the parasite, large amounts of LL

particles accumulating at the infection site would probably foster inflammation

(11,12,40,41). Inflammation might also be brought about the LL materials remaining in

circulation and possibly being deposited in the glomeruli or other sites. Indeed, if as we

speculated above, monocytes are recruited to the liver in response to trapped LL

particles, then Clec4F-dependent KC capture may be important to avoid liver

inflammation. This aspect may not have been fully captured in our flow cytometry and

histology experiments because of liver perfusion prior to analysis.

The events downstream of LL material capture by KC are certainly worth studying. KC

can present antigens to T cells present in the vascular space, and this normally leads to

tolerogenic responses that can have a systemic impact (22,42,43). The analysis of

immune responses in WT and Clec4f$^{-/-}$ mice infected with *E. granulosus* will be the

subject of a future communication.

Ancestrally, the larval stages of the parasites belonging to the taeniid family (including

those of the genus *Echinococcus*) probably infected the liver of rodent hosts (5,44). In

this context, the interaction of materials shed by the parasites with a receptor (Clec4F)

specifically expressed by phagocytic cells present in the same organ (KC) may have

been evolutionarily advantageous. This may have been so because LL material uptake

by KC would have avoided particle build-up and consequent inflammation in the liver

sinusoids and/or other sites, and/or because KC are poised for tolerogenic antigen

presentation. The extant species *E. multilocularis* dwells almost exclusively in the liver

of rodents (besides accidentally infecting humans). The LL of this species also interacts

with Clec4F, as suggested by experiments using recombinant Clec4F (13) and by the

presence in this species' LL of the blood antigen $P_1$ motif (45). Thus, host Clec4F is

likely an important player in *E. multilocularis* infection.

The hosts of larval *E. granulosus* are mostly ungulates, a taxon in which Clec4F also

appears to be expressed in the liver (15). Therefore, the uptake of LL materials by KC

via Clec4F probably takes place also in the ungulate natural hosts. However, in

ungulates Clec4F appears to be additionally expressed in secondary lymphoid organs

(15), opening the possibility that non-KC, anatomically more widely distributed,

phagocytes may act as additional sinks for LL materials.

Humans do not appear to express Clec4F, or at least not Clec4F with a functional lectin

domain (15). Humans are normally a dead-end for this parasite's life-cycle, meaning

that human *Echinococcus* infection has not been shaped through evolution. Although *E.*

*multilocularis* infection in humans is usually accompanied by strong local inflammation

(and tissue destruction), this is not usually the case with *E. granulosus* (9,46). It is a

possibility that for *E. granulosus*, which grows more slowly than *E. multilocularis*,

Clec4F-independent clearance mechanisms are in most infection settings sufficient to

avoid excessive build-up of LL debris. It is conceivable that such Clec4F-independent

mechanisms in humans include uptake via lectins expressed in KC with specificities

related to Clec4F. As mentioned, the uptake of de-sialylated platelets by mouse KC

requires Clec4F as well as MGLs or ASGR (19,20). It seems likely that a mechanism

for de-sialylated platelet uptake by KC based on a set of collaborating lectins exists

throughout mammals, even if in humans such a system does not include Clec4F. Our

previous screening using recombinant receptors yielded inconclusive results with

respect to the capacity of human MGL and ASGR to interact with the LL mucins (13).

When probing synthetic sugars, human ASGR and MGL appeared to bind less well than

Clec4F to the $P_1$ motif (after normalizing by binding to a known target of all 3

receptors). However, given the sheer abundance and significant breadth of Gal-based

motifs in the LL, it would not be surprising if ASGR and/or MGL are able to internalize

LL materials in the absence of Clec4F. It is important to determine if KC in liver

samples from humans infected with *E. granulosus*, as well as with *E. multilocularis*,

stain for LL materials.

**Materials and Methods**

*Parasite materials*

The starting material for LL mucin preparations, and the source of protoscoleces for

experimental infections, were macroscopically healthy, fertile *E. granulosus sensu lato*

hydatids from natural infections in cattle from Uruguay. For LL mucin preparations,

hydatid walls, which have the LL as the overwhelmingly dominant component (3,4,8),

were extensively washed with phosphate-buffered saline (PBS), then with 2 M NaCl

and finally with water. The walls were then dehydrated by sequential steps with ethanol

(95% v/v) followed by acetone, and finely ground using a mortar and pestle, followed

by grinding between two polished glass plates (12). The resulting fine powders were

rehydrated in endotoxin-free PBS containing 30 mM EDTA (1 ml for every 2 mg dry

mass) to extract the calcium inositol hexakisphosphate deposits, thus leaving the mucin

meshwork as single major structural component (3). The material was then extensively

washed into endotoxin-free PBS. To obtain a fine particulate preparation (pLL), the

suspension was filtered through 85- and 23-µm gauze (12). To obtain the soluble

preparation (sLL), the suspension was subjected to sonication followed by

centrifugation for 1 h at 100,000 g at 4°C. To obtain sLL in which the terminal

monosaccharide residues were oxidized, prior to the solubilization step, the material

was treated with sodium periodate followed by sodium borohydride (or with sodium

borohydride only as mock treatment) (12). For obtaining biotin-tagged sLL, prior to the

solubilization step, the material in PBS was incubated overnight with N-

hydroxysuccinimido-biotin (NHS-biotin; 10 ng per mg of LL dry mass). Sodium

periodate, sodium borohydride and/or NHS-biotin were removed by extensive washing

of the particles into endotoxin-free PBS. Destruction of terminal monosaccharides by

periodate was verified in terms of abrogation of the binding biotinylated PNA (26),

revealed with the help of streptavidin-perodixase, in a dot-blot format. Biotin

incorporation by the sLL was checked using streptavidin-peroxidase also in dot-blot

format. The dry mass content of LL mucin preparations was estimated as in (12). LL

mucin preparations were stored at 4°C in PBS with penicillin-streptomycin during no

more than 6 months (pLL) or 12 months (sLL).

*Clec4f^{-/-} mouse line generation*

All animal procedures were performed at the Specific Pathogen Free animal facility of

the Laboratory Animal Biotechnology Unit of Institut Pasteur de Montevideo.

Experimental protocols were approved by the Institutional Animal Ethics Committee

(protocol number 007-18), in agreement with National law 18.611 and international

animal care guidelines (Guide for the Care and Use of Laboratory Animals) (47)

regarding laboratory animal's protocols. Mice were housed in individually ventilated

cages (Tecniplast, Milan, Italy) containing chip bedding (Toplit 6, SAFE, Augy,

France), in a controlled environment at $20 \pm 1$ºC with a relative humidity of 40-60%, in

a 14/10 h light-dark cycle. Autoclaved food (Labdiet 5K67, PMI Nutrition, IN, US) and

autoclaved filtered water were administered *ad libitum*. The *Clec4f*[−/−] line was produced

using CRISPR-Cas9. Cytoplasmic microinjection was performed in C57BL/6J zygotes

using a mix of 25 ng/µl Cas9 mRNA (Invitrogen, Carlsbad, CA, US), 15 ng/µl of each

sgRNA (2 guides were used) (Synthego, Menlo Park, CA, US), diluted in standard

microinjection buffer (5 mM Tris-HCl, pH 7.4, 0.1 mM EDTA). Viable embryos were

cultured overnight in M16 (Sigma, St Louis, MO, US) microdrops under embryo tested

mineral oil (Sigma), with 5% $CO_2$ in air at 37 ºC. The next day, 2-cell embryos were

transferred into the oviduct of B6D2F1 0.5 days postcoitum pseudopregnant females

(20 embryos/female in average). Standard surgical procedures established in our animal

facility were followed (48). For surgery, recipient females were anesthetized with a

mixture of ketamine (110 mg/kg, Ketonal50, Richmond Veterinaria S.A., Buenos Aires,

Argentina) and xylazine (13 mg/kg, Seton 2%; Calier, Montevideo, Uruguay).

Tolfenamic acid was administered subcutaneously (1 mg/kg, Tolfedine, Vetoquinol,

Madrid, Spain) in order to provide analgesia and anti-inflammatory effects (49).

Pregnancy diagnosis was determined by visual inspection by an experienced animal

caretaker two weeks after embryo transfer, and litter size was recorded on day 7 after

birth. Pups were tail-biopsied and genotyped 21 days after birth, and mutant animals

were maintained as founders. Gene editing efficiency was 53.8% (7 KO/13 born pups).

Genetic modification was confirmed by Sanger sequencing. The Clec4f[−/−] line was

established after producing F1, F2, and finally homozygous F3 animals that were used

for experiments. Primers used for amplification of the genomic locus were 5'-

TCTTTATGATCGCACCCACA-3', 5'-TCCATTCTCGAGAGCCATCT-3'. F1 and F2

heterozygous mice were selected using a heteroduplex mobility assay on a 10%

polyacrylamide gel. In F2, WT animals were differentiated from homozygous KO

animals by Sanger sequencing.

*Mouse experiments*

WT BALB/c mice were obtained from DILAVE (MGAP, Uruguay) and kept at the

Instituto de Higiene animal house. Female WT (C57BL/6J) or Clec4f$^{-/-}$ mice were bred

at the Institut Pasteur de Montevideo, co-housed (so as to equilibrate their microbiota)

at the Instituto de Higiene starting at 5 weeks of age for at least 3 weeks, and used

between 8 and 14 weeks of age. sLL (100 or 200 µg dry mass per mouse as indicated in

each case, dissolved in 100 µl of PBS) or vehicle only was injected into the tail vein.

pLL (450 µg dry mass per mouse, suspended in 200 µl of PBS) or vehicle only, were

injected i.p. Mice were infected by i.p. injection of 2000 (BALB/c) or 2500 (C57BL/6)

protoscoleces (viability ≥ 95%) suspended in 200 µl of PBS; control mice were injected

with vehicle only. Mice were euthanized using isoflurane. The procedures used were

approved by the Comisión Honoraria de Experimentación Animal (CHEA, Universidad

de la República, Uruguay: protocols 101900-001017-16, 101900-000994-16, 101900-

000649-17, 101900-000441-16).

*Preparation of liver non-parenchymal fraction for flow cytometry*

Following perfusion through the inferior vena cava with 10 mL of PBS, livers were

placed in RPMI and finely chopped using a razor blade for approximately 2 min. The

tissue was then digested in 5 ml per g of enzyme mix (RPMI containing 0.4 U/ml

Liberase [Sigma] and 80 U/ml DNase [Sigma]), for 25 min at 37◦C, in an orbital shaker

at 250 rpm with additional manual shaking every 5 min. Digests were poured through a

100 µm strainer and centrifuged at 300 g for 5 min; the cell pellets were re-suspended in

25 ml of RPMI and again centrifuged at 300 g. The pellets were suspended in 2 ml of

RBC lysis buffer (Sigma), and after 2 min, 2 ml of FACS buffer (PBS supplemented

with 0.5% BSA and 2 mM EDTA) was added. After another centrifugation step (300 g,

5 min), cells were re-suspended in 3 ml of FACS buffer, passed through a 40 µm

strainer, and counted using the Nexcelom Cellometer K2 with the AOPI stain for cell

viability. Cells and buffer used with the cells were kept at 4ºC, except for the digestion

step.

*Flow cytometry*

Five hundred thousand live cells where incubated with the viability probe for 10 min at

room temperature, then with 0.025 µg of anti-CD16/32 (Biolegend) in 10% v/v normal

rat serum (Unidad de Reactivos y Biomodelos de Experimentación, Facultad de

Medicina, Universidad de la República). Cells were then surface-stained, fixed in 1%

w/v PFA for 20 min at 4°C, washed, and permeabilized overnight with permeabilisation

buffer (Thermo) for intracellular staining. The details of the viability probes, antibodies

and lectin probes are given in Supplementary Table 1. Finally, cells were washed, spun

at 350 g for 5 min and re-suspended in FACs buffer. The data were acquired in a FACS

Canto II Cytometer (BD), at the Instituto de Higiene, Universidad de la República, and

then analysed using the FlowJo software package. The relative contribution of each

liver non-parenchymal cell type under study to the uptake of LL materials was

estimated by multiplying the frequency of cells positive for the LL probe and belonging

to the particular cell type (within all live single cells) by their arithmetic mean

fluorescence intensity for the probe and then dividing this figure over the summation of

the analogous products for the 3 cell types analysed.

*Histology*

Tissue pieces (approximately 0.5 mm$^3$) were fixed in buffered formalin. Liver samples

corresponded to animals subjected to liver perfusion with PBS (as the materials were

also analyzed by flow cytometry). Paraffin-embedded tissue sections (2 µm) were

probed with biotinylated monoclonal antibody E492 (3 µg/mL; steamer pH 6.1) or

biotinylated PNA (Vector Laboratories, Newark, California, United States, B-1075;

dilution 1:100; steamer pH 9.0) using the avidin-biotin-complex method (50), and

counterstained using haemalum.

*ELISA*

ELISA wells were coated with sLL at 10 µg/mL (total dry mass) in Tris-buffered saline

(TBS) overnight at 4 ºC, and blocked using TBS containing 0.05% w/v Tween 20, 1%

w/v bovine serum albumin and 0.5% v/v Carbo-Free$^{TM}$ blocking solution (Vector

Laboratories). Wells were then probed with biotinylated E492 antibody (51),

biotinylated peanut agglutinin (PNA; Vector laboratories) or recombinant soluble

Clec4F (expressed as a fusion with the $F_c$ fragment of human IgG$_1$ (52)). The

biotinylated PNA or E492 antibody were detected using streptavidin coupled to

peroxidase (ThermoFisher Scientific) and the recombinant Clec4F was detected using a

goat anti-human IgG coupled to peroxidase (AbCam, Cambridge, UK). For competition

ELISA, recombinant Clec4F at 80 ng/mL was mixed with variable concentrations of

either (biotinylated) PNA or E492 antibody as competitors, and bound Clec4F detected

as previously described.

*Particle immunofluorescence*

pLL particles were incubated with biotinylated E492 antibody (0.6 µg/mL), biotinylated

PNA (1:1500) or recombinant soluble Clec4F (7.5 µg/mL), followed by streptavidin-

Cyanine-3 (1:500; Amersham Bioscience) or by goat anti-human IgG coupled to FITC

(1:100, Sigma), as appropriate. Control stains were included using human $IgG_1$ instead

of recombinant Clec4F as well omitting the primary staining reagents.

*Statistical analyses*

As the underlying data do not satisfy the criteria for parametric statistics, rank-based

nonparametric statistical methods, were used throughout. When the numbers of mice in

each treatment-repeat experiment combination were equal, the Mack-Skillings exact test

for a two-way layout was used (53). When the data did not fulfil the above condition,

data from the different repeat experiments were pooled and the modified Wilcoxon-

Mann-Whitney test (only two experimental groups) (53) or the Kruskal-Wallis test

(more than two experimental groups) were used. When the Mack-Skillings or the

Kruskal-Wallis tests resulted in a statistic with a p-value less than 0.05, the test was

followed by the post-hoc multiple comparison test described by Conover (55) and the

Benjamini-Hochberg correction for controlling the false discovery rate (56).

Throughout the paper, the symbols *, ** and *** represent p-values less than 0.05, 0.01

and 0.001 respectively.

**References**

1. Thompson RCA. Biology and Systematics of Echinococcus. Adv Parasitol. 2017;95:65-109.

2. Díaz Á. Immunology of cystic echinococcosis (hydatid disease). British Medical Bulletin. 7 de octubre de 2017;124(1):121-33.

3. Díaz A, Casaravilla C, Irigoín F, Lin G, Previato JO, Ferreira F. Understanding the laminated layer of larval Echinococcus I: structure. Trends in Parasitology. mayo de 2011;27(5):204-13.

4. Díaz A, Casaravilla C, Allen JE, Sim RB, Ferreira AM. Understanding the laminated layer of larval Echinococcus II: immunology. Trends in Parasitology. junio de 2011;27(6):264-73.

5. Díaz Á, Fernández C, Pittini Á, Seoane PI, Allen JE, Casaravilla C. The laminated layer: Recent advances and insights into Echinococcus biology and evolution. Experimental Parasitology. noviembre de 2015;158:23-30.

6. Díaz A, Barrios AA, Grezzi L, Mouhape C, Jenkins SJ, Allen JE, et al. Immunology of a unique biological structure: the Echinococcus laminated layer. Protein & Cell. 2022;

7. Lin G, Todeschini AR, Koizumi A, Neves JL, González H, Dematteis S, et al. Further structural characterization of the Echinococcus granulosus laminated layer carbohydrates: The blood-antigen P1-motif gives rise to branches at different points of the O-glycan chains. Glycobiology. abril de 2013;23(4):438-52.

8. Díaz A, Fontana EC, Todeschini AR, Soulé S, González H, Casaravilla C, et al. The Major Surface Carbohydrates of the *Echinococcus granulosus* Cyst: Mucin-Type *O*-Glycans Decorated by Novel Galactose-Based Structures. Biochemistry. 15 de diciembre de 2009;48(49):11678-91.

9. Díaz Á, Sagasti C, Casaravilla C. Granulomatous responses in larval taeniid infections. Parasite Immunol. mayo de 2018;40(5):e12523.

10. Grimm J, Nell J, Hillenbrand A, Henne-Bruns D, Schmidberger J, Kratzer W, et al. Immunohistological detection of small particles of Echinococcus multilocularis and Echinococcus granulosus in lymph nodes is associated with enlarged lymph nodes in alveolar and cystic echinococcosis. PLoS Negl Trop Dis. diciembre de 2020;14(12):e0008921.

11. Pittini Á, Martínez-Acosta YE, Casaravilla C, Seoane PI, Rückerl D, Quijano C, et al. Particles from the Echinococcus granulosus Laminated Layer Inhibit CD40 Upregulation in Dendritic Cells by Interfering with Akt Activation. Infect Immun. diciembre de 2019;87(12):e00641-19.

12. Casaravilla C, Pittini A, Rückerl D, Seoane PI, Jenkins SJ, MacDonald AS, et al. Unconventional maturation of dendritic cells induced by particles from the

laminated layer of larval Echinococcus granulosus. Infect Immun. agosto de 2014;82(8):3164-76.

13. Hsu TL, Lin G, Koizumi A, Brehm K, Hada N, Chuang PK, et al. The surface carbohydrates of the Echinococcus granulosus larva interact selectively with the rodent Kupffer cell receptor. Molecular and Biochemical Parasitology. noviembre de 2013;192(1-2):55-9.

14. Fadden AJ, Holt OJ, Drickamer K. Molecular characterization of the rat Kupffer cell glycoprotein receptor. Glycobiology. 1 de julio de 2003;13(7):529-37.

15. Taylor ME, Snelling T, Smith DF, Drickamer K. Absence of a human ortholog of rodent Kupffer cell galactose-binding receptor encoded by the CLEC4f gene. Glycobiology. 1 de abril de 2019;29(4):332-45.

16. Yang CY, Chen JB, Tsai TF, Tsai YC, Tsai CY, Liang PH, et al. CLEC4F Is an Inducible C-Type Lectin in F4/80-Positive Cells and Is Involved in Alpha-Galactosylceramide Presentation in Liver. Schulz C, editor. PLoS ONE. 6 de junio de 2013;8(6):e65070.

17. Coombs PJ, Taylor ME, Drickamer K. Two categories of mammalian galactose-binding receptors distinguished by glycan array profiling. Glycobiology. 1 de agosto de 2006;16(8):1C-7C.

18. Ouyang Z, Felix J, Zhou J, Pei Y, Ma B, Hwang PM, et al. Trimeric structure of the mouse Kupffer cell C-type lectin receptor Clec4f. FEBS Lett. enero de 2020;594(1):189-98.

19. Jiang Y, Tang Y, Hoover C, Kondo Y, Huang D, Restagno D, et al. Kupffer cell receptor CLEC4F is important for the destruction of desialylated platelets in mice. Cell Death Differ. 15 de mayo de 2021;28(11):3009-21.

20. Deppermann C, Kratofil RM, Peiseler M, David BA, Zindel J, Castanheira FVES, et al. Macrophage galactose lectin is critical for Kupffer cells to clear aged platelets. J Exp Med. 6 de abril de 2020;217(4):e20190723.

21. Li Y, Fu J, Ling Y, Yago T, McDaniel JM, Song J, et al. Sialylation on O-glycans protects platelets from clearance by liver Kupffer cells. Proc Natl Acad Sci USA. 1 de agosto de 2017;114(31):8360-5.

22. Heymann F, Peusquens J, Ludwig-Portugall I, Kohlhepp M, Ergen C, Niemietz P, et al. Liver inflammation abrogates immunological tolerance induced by Kupffer cells. Hepatology. julio de 2015;62(1):279-91.

23. Rothlin CV, Carrera-Silva EA, Bosurgi L, Ghosh S. TAM receptor signaling in immune homeostasis. Annu Rev Immunol. 2015;33:355-91.

24. Lynch RW, Hawley CA, Pellicoro A, Bain CC, Iredale JP, Jenkins SJ. An efficient method to isolate Kupffer cells eliminating endothelial cell contamination and selective bias. J Leukoc Biol. septiembre de 2018;104(3):579-86.

25. Cummings RD, Etzler, ME. Antibodies and lectins in glycan analysis. En:
Essentials of Glycobiology. Second Edition. Cold Spring Harbor Laboratory Press;
p. 633-47.

26. Casaravilla C, Díaz A. Studies on the structural mucins of the Echinococcus
granulosus laminated layer. Molecular and Biochemical Parasitology. diciembre de
2010;174(2):132-6.

27. Dematteis S, Pirotto F, Marqués J, Nieto A, Orn A, Baz A. Modulation of the
cellular immune response by a carbohydrate rich fraction from Echinococcus
granulosus protoscoleces in infected or immunized Balb/c mice. Parasite Immunol.
enero de 2001;23(1):1-9.

28. Walker M, Baz A, Dematteis S, Stettler M, Gottstein B, Schaller J, et al. Isolation
and characterization of a secretory component of Echinococcus multilocularis
metacestodes potentially involved in modulating the host-parasite interface. Infect
Immun. enero de 2004;72(1):527-36.

29. Araj GF, Matossian RM, Frayha GJ. The host response in secondary hydatidosis of
mice. I. Circulating antibodies. Z Parasitenkd. 3 de junio de 1977;52(1):23-30.

30. Richards KS, Arme C, Bridges JF. Echinococcus granulosus equinus: an
ultrastructural study of murine tissue response to hydatid cysts. Parasitology. junio
de 1983;86 (Pt 3):407-17.

31. Mourglia-Ettlin G, Cucher M, Arbildi P, Rosenzvit M, Dematteis S. Natural and
induced antibodies contribute to differential susceptibility to secondary cystic
echinococcosis of Balb/c and C57Bl/6 mice. Immunobiology. enero de
2016;221(1):103-15.

32. Raedler A, Raedler E, Arndt R, Thiele HG. Terminal galactosyl residues of cell-
surface glycoconjugates exposed on both human and murine immature T- and B-
cells. Cell Tissue Res. 1981;218(1):219-26.

33. Wang J, Kubes P. A Reservoir of Mature Cavity Macrophages that Can Rapidly
Invade Visceral Organs to Affect Tissue Repair. Cell. 21 de abril de
2016;165(3):668-78.

34. Grewal PK. The Ashwell–Morell Receptor. En: Methods in Enzymology [Internet].
Elsevier; 2010 [citado 17 de noviembre de 2021]. p. 223-41. Disponible en:
https://linkinghub.elsevier.com/retrieve/pii/S0076687910790133

35. Barrios AA, Grezzi L, Miles S, Mariconti M, Mourglia-Ettlin G, Seoane PI, et al.
Inefficient and abortive classical complement pathway activation by the calcium
inositol hexakisphosphate component of the Echinococcus granulosus laminated
layer. Immunobiology. septiembre de 2019;224(5):710-9.

36. Irigoín F, Ferreira F, Fernández C, Sim RB, Díaz A. myo-Inositol hexakisphosphate
is a major component of an extracellular structure in the parasitic cestode
Echinococcus granulosus. 2002;8.

37. Irigoín F, Casaravilla C, Iborra F, Sim RB, Ferreira F, Díaz A. Unique precipitation and exocytosis of a calcium salt of *myo* -inositol hexakisphosphate in larval *Echinococcus granulosus*. J Cell Biochem. 15 de diciembre de 2004;93(6):1272-81.

38. Casaravilla C, Brearley C, Soule S, Fontana C, Veiga N, Bessio MI, et al. Characterization of myo-inositol hexakisphosphate deposits from larval Echinococcus granulosus. FEBS Journal. julio de 2006;273(14):3192-203.

39. Irigoín F, Laich A, Ferreira AM, Fernández C, Sim RB, Díaz A. Resistance of the Echinococcus granulosus cyst wall to complement activation: analysis of the role of InsP $_6$ deposits. Parasite Immunol. junio de 2008;30(6-7):354-64.

40. Seoane PI, Rückerl D, Casaravilla C, Barrios AA, Pittini Á, MacDonald AS, et al. Particles from the Echinococcus granulosus laminated layer inhibit IL-4 and growth factor-driven Akt phosphorylation and proliferative responses in macrophages. Sci Rep. diciembre de 2016;6(1):39204.

41. Casaravilla C, Pittini Á, Rückerl D, Allen JE, Díaz Á. Activation of the NLRP3 Inflammasome by Particles from the Echinococcus granulosus Laminated Layer. Herbert DR, editor. Infect Immun [Internet]. 19 de agosto de 2020 [citado 23 de septiembre de 2021];88(9). Disponible en: https://journals.asm.org/doi/10.1128/IAI.00190-20

42. Lüth S, Huber S, Schramm C, Buch T, Zander S, Stadelmann C, et al. Ectopic expression of neural autoantigen in mouse liver suppresses experimental autoimmune neuroinflammation by inducing antigen-specific Tregs. J Clin Invest. octubre de 2008;118(10):3403-10.

43. You Q, Cheng L, Kedl RM, Ju C. Mechanism of T cell tolerance induction by murine hepatic Kupffer cells. Hepatology. septiembre de 2008;48(3):978-90.

44. Hoberg EP, Jones A, Rausch RL, Eom KS, Gardner SL. A phylogenetic hypothesis for species of the genus Taenia (Eucestoda☐: Taeniidae). J Parasitol. febrero de 2000;86(1):89-98.

45. del Puerto L, Rovetta R, Navatta M, Fontana C, Lin G, Moyna G, et al. Negligible elongation of mucin glycans with Gal β1-3 units distinguishes the laminated layer of Echinococcus multilocularis from that of Echinococcus granulosus. International Journal for Parasitology. mayo de 2016;46(5-6):311-21.

46. Basika T, Muñoz N, Casaravilla C, Irigoín F, Batthyány C, Bonilla M, et al. Phagocyte-specific S100 proteins in the local response to the *Echinococcus granulosus* larva. Parasitology. febrero de 2012;139(2):271-83.

47. (1) Committee for the update of the guide for the care and use of laboratory animals. Guide for the care and use of laboratory animals. 8th ed. National Academies Press; 2011.

48. Meikle MN, Schlapp G, Menchaca A, Crispo M. Minimum volume Spatula MVD vitrification method improves embryo survival compared to traditional slow freezing, both for in vivo and in vitro produced mice embryos. Cryobiology. octubre de 2018;84:77-81.

49. Schlapp G, Goyeneche L, Fernández G, Menchaca A, Crispo M. Administration of
the nonsteroidal anti-inflammatory drug tolfenamic acid at embryo transfer
improves maintenance of pregnancy and embryo survival in recipient mice. J Assist
Reprod Genet. febrero de 2015;32(2):271-5.

50. Barth TFE, Herrmann TS, Tappe D, Stark L, Grüner B, Buttenschoen K, et al.
Sensitive and Specific Immunohistochemical Diagnosis of Human Alveolar
Echinococcosis with the Monoclonal Antibody Em2G11. Garcia HH, editor. PLoS
Negl Trop Dis. 25 de octubre de 2012;6(10):e1877.

51. Baz A, Richieri A, Puglia A, Nieto A, Dematteis S. Antibody response in CD4-
depleted mice after immunization or during early infection with Echinococcus
granulosus. Parasite Immunol. marzo de 1999;21(3):141-50.

52. Hsu TL, Cheng SC, Yang WB, Chin SW, Chen BH, Huang MT, et al. Profiling
carbohydrate-receptor interaction with recombinant innate immunity receptor-Fc
fusion proteins. J Biol Chem. 11 de diciembre de 2009;284(50):34479-89.

53. Hollander M, Wolfe DE, Chicken E. Nonparametric Statistical Methods. 3rd ed.
New York: John Wiley & Sons; 2015. (Wiley Series in Probability and Statistics).

**Acknowledgements**

This work was funded by ANII (Government of Uruguay) grant FCE 136130 (to AD),

by a Travel Grant (to SJJ) from the Science and Innovation Fund of the Foreign and

Commonwealth Office (UK Government), by MRC grant MR/L008076/1 (to SJJ), and

by PEDECIBA (Government of Uruguay). AB and CM were supported by PhD

scholarships from Comisión Académica de Posgrado (Universidad de la República,

Uruguay) and ANII respectively.

**Figure legends**

**Figure 1. Soluble LL mucins are taken up selectively by KC, dependent on mucin**

**glycans.** C57BL/6 mice were injected i.v. with 100 µg dry mass of solubilized, biotin-

derivatized LL mucins (sLL), either treated with periodate to oxidize their terminal

monosaccharide residues or mock-treated. Twenty minutes later, mice were sacrificed and

sLL uptake was quantitated in liver non parenchymal cells using streptavidin-PE. Data are

shown for the cell types that are relevant in the context of the article, namely KC, LSEC

and non-KC CD11b$^+$ cells, in terms of percentages of positive cells **(a)** and geometric

means of fluorescence intensity **(b)**. Representative dot-plots for KC are shown in **(c)**. The

signal threshold for each cell type was established using cells from mice injected with

vehicle only and similarly stained; the fluorescence intensities data had the median values

corresponding to vehicle-injected mice subtracted for plotting. Results are pooled from 3

independent experiments (represented in different colours); individual mice and their

overall median values are shown. The statistical analyses were carried out by the Mack-

Skillings test.

**Figure 2. PNA and the E492 antibody are suitable probes for LL mucins captured**

**by mouse KC.** The binding of PNA, the E492 antibody and soluble recombinant

Clec4F to LL mucins coated onto ELISA plate wells **(a)** or present in LL particles in

suspension **(b)** was studied. The capacity of PNA and the E492 antibody to compete

with the binding of recombinant Clec4F to LL mucins was analysed in a competition

ELISA format **(c)**; results are expressed in relation to the response in the absence of

competitor. In (a) and (c), analytical duplicates are shown. WT C57BL/6 mice were

injected i.v. either with PBS only or with solubilized LL mucins (200 µg/mouse) and 22

h later KC and LSEC were analysed by flow cytometry using either PNA or the E492 antibody as probes, after cell permeabilization (**d**). Staining controls shown in (b) and (d) correspond to samples stained in the same way as the positive ones except that the primary probe for LL material was omitted, and the analogous controls in (a) gave no signal.

**Figure 3. Generation of Clec4f gene-deficient mice. (a)** Schematic structure of the mouse *Clec4f* gene. Coding exons appear as solid black bars and non-coding regions as white bars. Introns are represented as lines. **(b)** The target sequences for CRISPR/Cas9, 5'-TGGACGTGCCGCAGGGTCCT-3' and 5'-GCTTCTCCCCTGCAGGACTG-3', are located in the second exon and first intron/second exon junction of the gene respectively. An 8 bp deletion corresponding to nucleotides 5'- CAGGACCC-3' was identified by tail biopsy. The frameshift mutation produces a protein 480 aa shorter. **(c)** The liver non-parenchymal fraction was prepared from Clec4f$^{-/-}$ and WT mice and analyzed for Clec4F expression by flow cytometry without or with cellular permeabilization. KC were defined as shown in Figure S2. The staining controls contain all the fluorophore-coupled antibodies but lack the primary anti-Clec4F antibody.

**Figure 4. Fast uptake of soluble LL mucins by Kupffer cells depends largely on Clec4F.** WT and Clec4f$^{-/-}$ mice were injected i.v. with 200 ug dry mass of sLL, either untreated or mock-treated as in Figure 1, or vehicle only (PBS). Twenty min later, mice were sacrificed and sLL uptake was quantitated in non-parenchymal liver cell populations by flow cytometry in permeabilized cells using PNA. Results are shown for KC, LSEC and non-KC CD11b$^{+}$ cells, in terms of percentages of positive cells **(a)** and mean fluorescence intensities **(b)**, and representative flow cytometry plots are shown for KC and LSEC **(c)**. An

estimation of the fraction of sLL taken up by each of the cell populations under study,

calculated as explained in the Materials and Methods section, is shown in **(d)**; the statistics

shown compare the fractions taken up by KC. The positive cell threshold was established

using cells from the mice injected with PBS only. The data arise from two independent

experiments, represented in blue and red in (a-c) and indicated as 1, 2 in (d). In (a-c),

individual mice and their median values, and in (d) only medians, are shown. The statistical

analyses were carried out by the Mack-Skillings test.

**Figure 5. Clec4F is important for the capture of soluble LL mucins by Kupffer cells**

**even after allowing a long uptake time.** WT and Clec4f$^{-/-}$ C57BL/6 mice were injected i.v.

with 200 µg dry mass per mouse of sLL (mock-treated as in Fig. 1) or PBS only as a

control. Twenty-two h later, mice were sacrificed and sLL uptake was quantitated in liver

non-parenchymal cells by flow cytometry using PNA after cell permeabilization. Results

are shown for KC, LSEC and non-KC CD11b$^+$ cells, in terms of percentages of positive

cells **(a)** and mean fluorescence intensities **(b)**, and representative flow plots for KC and

LSEC are shown in **(c)**. An estimation of the fraction of sLL taken up by each of the cell

populations under study, calculated as explained in the Materials and Methods section, is

shown in **(d);** the statistics shown compare the fractions taken up by KC. The positive cell

threshold was established using staining controls (in which only biotinylated PNA was

omitted). The PNA signal in non-permeabilized cells was negligible, as expected since

EDTA was included in the stain buffer. The data arise from two independent experiments,

represented in blue and red in (a-c) and indicated as 1, 2 in (d). In (a-c), individual mice and

their median values, and in (d) only medians, are shown. The statistical analyses were

carried out by the Mack-Skillings test (a-c) and the modified Wilcoxon-Mann-Whitney tests

(d).

**Figure 6. The uptake by Kupffer cells of LL mucins injected in particulate presentation depends largely on Clec4F.** WT and Clec4f$^{-/-}$ C57BL/6 mice were injected i.p. with 450 µg dry mass of pLL or PBS only as a control. Twenty-two h later, mice were sacrificed and pLL uptake was quantitated in liver non-parenchymal cells by flow cytometry using PNA after cell permeabilisation. Results are shown for KC, LSEC and non-KC CD11b$^+$ cells, in terms of percentages of positive cells **(a)** and mean fluorescence intensities **(b)**, and representative dot-plots for the three cell types are shown in **(c)**. An estimation of the fraction of sLL taken up by each of the cell populations under study, calculated as explained in the Materials and Methods section, is shown in **(d)**; the statistics shown compare the fractions taken up by KC.. The positive cell threshold was established using staining controls. The data arise from two independent experiments, represented in blue and red in (a-c) and indicated as 1, 2 in (d). In (a-c), individual mice and their median values, and in (d) only medians, are shown. The statistical analyses were carried out by the Mack-Skillings test. The analogous flow cytometry results obtained using the E492 antibody for detection are shown in Fig. S6. In addition, pLL was detected in liver sections, using the E492 antibody (**e**). Representative images are shown, at two different magnifications. Black arrows indicate KC-like cells staining strongly for LL material; empty arrowheads indicate hepatocytes staining weakly for LL material.

**Figure 7. LL materials circulate systemically in infected mice and are captured selectively by Kupffer cells via Clec4F.** BALB/c mice (WT) were intraperitoneally infected with *E. granulosus* protoscoleces or injected with the corresponding volume of buffer only. Seven months later, the presence of LL-derived materials in various organs was analysed by histology, using the E492 antibody as a probe (**a**). Images from representative

mice are shown, out of 15 infected and 5 control mice; the bars represent 50 µm. Black

arrows indicate KC-like cells staining for LL material; empty arrowheads indicate

occasional staining for LL material in spleen and lymph node. The staining observed in

adipocytes (upper left-hand corner of the infected mouse mesenteric lymph node section)

was also observed in adipocytes present in kidney and lung sections, irrespective of

infection, and must therefore represent a cross-reaction of the antibody. No staining was

observed in sections probed with the secondary reagent only. The uptake of LL-derived

materials by non-parenchymal liver cells of infected BALB/c mice was also analysed by

flow cytometry, using PNA as probe. Results are shown for KC, LSEC and non-KC

CD11b$^+$ cells, in terms of percentages of positive cells **(a)** and mean fluorescence intensities

**(b)**, and representative dot-plots for the three cell types are shown in **(c)**. The positive cell

threshold was established using staining controls. In addition, WT and Clec4f$^{-/-}$ C57BL/6

mice were similarly infected and the uptake of LL-derived materials analysed by flow

cytometry and presented **(e – g)** as for the BALB/c mice. Results in (a – d) are from a single

experiment and results in (e - g) are pooled from two independent experiments, shown in

blue and red. In (b - g), data shown correspond to individual mice and their median values

are indicated. The statistical analyses were carried out by the modified Wilcoxon-Mann-

Whitney (b, c) and Kruskal-Wallis tests (e, g).

**Supplementary Figures.**

**Figure S1. Summary of preparation of LL materials.** The preparation of pLL and sLL

(periodate- or mock-treated) is summarized in **(a)**. Periodate oxidation of terminal

monosaccharides and further monosaccharides carrying *cis*-diols generates reactive

aldehyde groups, which are reduced to alcohols using sodium borohydride. The mock

treatment consists of the borohydride step only. The effectiveness of the periodate treatment

was verified in terms of abrogation of PNA binding by to the mucins in a dot-blot format

**(b)**. Similar incorporation of the biotin tag to periodate- and mock-treated sLL was verified

using streptavidin-peroxidase in dot-blot format **(c)**.

**Figure S2. Gating strategy for liver cells.** Gating strategies are shown for KC, LSEC, and

non-KC CD11b$^{+}$ cells. For non-KC CD11b$^{+}$ cells, which tend to be substantially smaller

than KC, it was necessary to include an additional doublet exclusion step at the end of the

gating strategy.

**Figure S3 (related to Fig. 3). Reactivity of PNA, E492 antibody and recombinant**

**Clec4F with liver non-parenchymal cells of control mice.** Non-parenchymal cells from

WT C57BL/6 mice were stained for flow cytometry with PNA, E492 antibody or

recombinant mouse Clec4F, followed by appropriate secondary probes. The staining

controls correspond to samples stained in the same way as the positive ones except that the

primary probe was omitted.

**Figure S4 (related to Figs. 4 and 5). Representative flow plots for liver non-KC**

**CD11b$^{+}$ cells.** sLL was injected in WT and Clec4f$^{-/-}$ C57BL/6 mice, and liver non

parenchymal cells analysed after 20 min **(a)** or 22 h **(b)**, as described in Figs. 4 and 5

respectively. Representative flow plots are shown for non-KC CD11b$^{+}$ cells, which were

not found to take up sLL significantly.

**Figure S5 (related to Fig. 5). Surface expression of Clec4F does not change upon**

**injection of sLL.** WT C57BL/6 mice were injected with sLL or vehicle only (PBS) as

described in Fig. 5, and 22 h later KC were analysed for surface expression of Clec4F by

flow cytometry. Data shown, in terms of geometric means of fluorescence intensity (MFI),

arise from 3 independent experiments, shown in different colors. The data were normalized

over the median value of mice injected with PBS only in each experiment.

**Figure S6 (related to Fig. 6). Uptake of pLL by liver non-parenchymal cells measured**

**by flow cytometry using antibody E492.** WT and Clec4f$^{-/-}$ C57BL/6 mice were injected

with pLL and material uptake was analysed as in Figure 6 except that the E492 antibody

was used as probe. Results are shown in terms of percentages of positive cells **(a)** and mean

fluorescence intensities **(b)**, and representative dot-plots are shown in **(c)**. An estimation of

the fraction of sLL taken up by each of the cell populations under study, calculated as

explained in the Materials and Methods section, is shown in **(d)**; the statistics shown

compare the fractions taken up by KC. The positive cell threshold was established using

staining controls. The data arise from two independent experiments, represented in blue and

red in (a-c) and indicated as 1, 2 in (d). In (a-c), individual mice and their median values,

and in (d) only medians, are shown. The statistical analyses were carried out by the Mack-

Skillings test.

**Figure S7 (related to Fig. 6). Frequency of non-KC CD11b$^{+}$ cells in mice injected with**

**pLL.** WT and Clec4f$^{-/-}$ C57BL/6 mice were injected with pLL or vehicle only (PBS) as

described in Fig. 6, and 22 h later the frequency of non-KC CD11b$^{+}$ cells within all live

cells was measured by flow cytometry. Data arise from 2 independent experiments, shown

in different colors.

**Figure S8 (related to Fig. 6). Surface expression of Clec4F does not change upon**

**injection of pLL.** WT C57BL/6 mice were injected with pLL or vehicle only (PBS) as

described in Fig. 6, and 22 h later KC were analysed for surface expression of Clec4F by

flow cytometry. Data shown, in terms of geometric means of fluorescence intensity (MFI),

arise from 2 independent experiments, shown in different colors. The data were normalized

over the median value of mice injected with PBS only in each experiment.

**Figure S9 (related to Fig. 7). Parasite burdens of mice infected with *E. granulosus*.** WT

BALB/c mice **(a)** or WT and Clec4f[-/-] C57BL/6 mice **(b)** were infected with *E. granulosus*

as described in Fig. 7. Parasite burdens are expressed in terms of total volumes of

metacestodes, measured by water displacement. Results for BALB/c mice correspond to a

single experiment and results for C57BL/6 are pooled from 2 experiments, represented in

different colours. Individual mice and their overall median values are shown. Parasite

burdens were not found to be significantly different between WT and Clec4f[-/-] C57BL/6

mice by the Kruskall-Wallis test.

1000

1001    **Figure S10 (related to Fig. 7). Representative flow plots for LSEC in infections.** WT

1002    BALB/c mice **(a)** or WT and Clec4f[-/-] C57BL/6 mice **(b)** were infected with *E. granulosus*

1003    and liver non-parenchymal cells analysed by flow cytometry as described in Fig. 7.

1004    Representative flow plots are shown for LSEC, which were not found to take up LL

1005    materials.

1006

1007    **Figure S11. Frequencies of non-KC CD11b[+] cells in *E. granulosus* experimental**

1008    **infections.** WT BALB/c mice **(a)** or WT and Clec4f[-/-] C57BL/6 mice **(b)** were infected

1009    with *E. granulosus* and liver non-parenchymal cells analysed by flow cytometry as

1010    described in Fig. 7. Frequencies of non-KC CD11b$^+$ cells within total live cells are plotted.

1011    The statistical analysis was carried out by the modified Wilcoxon-Mann-Whitney method.

1012

**Figure S12. Experimental *E. granulosus* infection does not alter the expression of**

**Clec4F in KC.** WT BALB/c mice **(a)** or WT C57BL/6 mice **(b)** were infected with *E.*

*granulosus* as described in Fig. 7. KC were analysed for surface expression of Clec4F by

flow cytometry. Data shown, in terms of geometric means of fluorescence intensity (MFI),

arise from 1 experiment for each of the mouse strains shown. The data were normalized

over the median value of the control mice in each experiment.

1019

# FIGURE 1

# FIGURE 2

## (a) ELISA format

## (b) Immuno/lectin fluorescence on particles

## (c) Competition ELISA format

## (d) Flow cytometry gated on KC and LSEC

# FIGURE 3

## (a)

*Mus musculus clec4f (Chr. 6)*
ENSMUST00000014686.3

## (b)

| WT | C T C A T G G C T C C C A G G A C C C T G C G G C A C G T C C A G G T C A T T C |
|---|---|
| | Leu Met Ala Pro Arg Thr Leu Arg His Val Gln Val Ile |

| *clec4f mut* | C T C A T G G C T C C        T G C G G C A C G T C C A G G T C A T T C |
|---|---|
| | Leu Met Ala Pro     Ala Ala Arg Pro Gly His Ser |

WT   MKEAELNRDMARYCTDNQCVSLQPQGLGPKSAALMAPRTLRHVQVILALMVVTVIFSLLALFVVASQPWRPEWNKEPPSLLLR
*clec4f mut*   MKEAELNRDMARYCTDNQCVSLQPQGLGPKSAALMAPAARPGHSGINGGDCDLFSPCSFCGGFTALET*

## (c)

# FIGURE 4

# Figure 5

# Figure 6

# Figure 7

**(a)**

| | Peritoneal cavity cells | Liver | Spleen | Mesenteric lymph node |
|---|---|---|---|---|
| Control | | | | |
| Infected | | | | |

| | Small intestine | Kidney | Lung |
|---|---|---|---|
| Control | | | |
| Infected | | | |

**(b)** KC, LSEC, Non-KC CD11b⁺ cells — % positive cells (ctrl, inf.)

**(c)** KC, LSEC, Non-KC CD11b⁺ cells — MFI (ctrl, inf.)

**(d)** staining control, control, infected — KC, Non-KC CD11b⁺ cells; PNA binding vs F4/80

**(e)** KC, LSEC, Non-KC CD11b⁺ cells — % positive cells (WT: ctrl, inf.; Clec4f⁻/⁻: ctrl, inf.)

**(f)** KC, LSEC, Non-KC CD11b⁺ cells — MFI (WT: ctrl, inf.; Clec4f⁻/⁻: ctrl, inf.)

**(g)** staining control, WT (control, infected), Clec4f⁻/⁻ (control, infected) — KC, Non-KC CD11b⁺ cells; PNA binding vs F4/80