Mucins shed from the laminated layer in cystic echinococcosis are

captured by Kupffer cells via the lectin receptor Clec4F

Anabella A. Barrios[a], Camila Mouhape[a], Leonard Schreiber[b], Juliane Nell[b], Mariana

Suárez-Martins[a], Geraldine Schlapp[c], María Noel Meikle[c], Ana Paula Mulet[c], Tsui-Ling

Hsu[d], Shie-Liang Hsieh[d], Gustavo Mourglia-Ettlin[a], Carlos González[e], Martina Crispo[c],

Thomas F. E. Barth[b], Cecilia Casaravilla[a], Stephen J. Jenkins[f] & Álvaro Díaz[a]

a. Área Inmunología, Departamento de Biociencias (Facultad de Química) and Cátedra

de Inmunología, Instituto de Química Biológica (Facultad de Ciencias), Universidad

de la República, Montevideo, Uruguay.

b. Institute of Pathology, University Ulm, Ulm, Germany

c. Unidad de Biotecnología en Animales de Laboratorio, Institut Pasteur de

Montevideo, Montevideo, Uruguay.

d. Genomics Research Center, Academia Sinica, Taipei, Taiwan.

e. Retired from: CIMMYT (Centro Internacional de Mejoramiento de Maíz y Trigo),

El Batán, México.

f. University of Edinburgh Centre for Inflammation Research, Queen's Medical

Research Institute, University of Edinburgh, Edinburgh, UK.

Running title: Kupffer cells capture *Echinococcus* mucins via Clec4F

Corresponding author:

Full name: Álvaro Díaz

Department: Área Inmunología, Departamento de Biociencias (Facultad de Química) and

Cátedra de Inmunología, Instituto de Química Biológica (Facultad de Ciencias),

Universidad de la República, Montevideo, Uruguay.

Street address, city, postal code and country: Instituto de Higiene. Avenida Alfredo

Navarro 3051. Montevideo CP 11600. Uruguay.

Tel: + 598 24874320

E-mail: adiaz@fq.edu.uy, alvarojdy@gmail.com

Keywords: *Echinococcus*, laminated layer, lectin, mucin, Kupffer cells, Clec4F

Abstract

Cystic echinococcosis is caused by the larval stages (hydatids) of cestode parasites

belonging to the species cluster *Echinococcus granulosus sensu lato*. Hydatids are

bladder-like structures that attain large sizes within various internal organs of livestock

ungulates and humans. Hydatids are protected by the massive acellular laminated layer

(LL), composed mainly by mucins. Parasite growth requires LL turnover, and abundant

LL-derived particles are found at infection sites in infected humans, raising the question

of how LL materials are dealt with by the hosts. In this article, we show that LL mucins

injected into mice are taken up by Kupffer cells, the liver macrophages exposed to the

vascular space. This uptake is largely dependent on the intact mucin glycans and on

Clec4F, a C-type lectin receptor which in rodents is selectively expressed in Kupffer cells.

This uptake mechanism operates on mucins injected both in soluble form i.v. and in

particulate form i.p. In mice harbouring intraperitoneal infections, LL mucins were found

essentially only at the infection site and in the liver, where they were taken up by Kupffer

cells via Clec4F. Therefore, shed LL materials circulate in the host and Kupffer cells can

act as a sink for these materials even when the parasite grows in sites other than the liver.

Introduction

The larval stages of the cestode parasites belonging to the species cluster known as *Echinococcus granulosus sensu lato* (distinguishable only by molecular means) cause cystic echinococcosis in livestock animals including sheep and cattle, as well as in humans (1, 2). This disease is characterized by the growth within organ parenchymae of bladder-like parasite structures called hydatids. Each hydatid is bounded by a thin cellular layer, in turn protected by a thick acellular coat called the laminated layer (LL) (3–6). The main component of the LL is a meshwork of mucins, heavily decorated by O-glycans that are rich in galactose (7, 8). Hydatids persist in internal organs of immunocompetent hosts for up to decades, and they usually elicit weak local inflammation (9). The growth of the hydatid (potentially through over 3 orders of magnitude in lineal dimension) requires the shedding of materials from the LL (3–6). In human infections, the abundant shed LL particles (termed SPEGs, for shed particles of *E. granulosus*) are found in the tissue surrounding the parasite and in the draining lymph nodes (10).

The persistence of larval *E. granulosus* infection, the massive nature of the LL, and sheer abundance of shed particles as detected in human infection call the question of how the host's body disposes of this foreign debris and avoids the potential inflammation that may result from its build-up. As inflammation is detrimental to the parasite (9), it makes biological sense that LL constituents may be evolutionarily perfected for engaging cellular receptors that will ensure non-inflammatory disposal of the shed materials.

Studies on the responses elicited in mouse dendritic cells (DC) and macrophages by *in vitro*-produced LL particles (pLL, from particulate LL) have suggested an interaction

mode independent of conventional receptors and inefficient phagocytosis, at least in the

absence of opsonization (6, 11, 12). In broad agreement, none out of 35 human innate

carbohydrate-recognition proteins (recombinantly expressed as fusions with IgG Fc)

bound the LL mucins in an *in vitro* screening (13). However, in the same experiment,

strong binding was observed for mouse Clec4F. This type II, trimeric, cell-surface C-type

lectin has affinity for glycans terminated in N-acetylgalactosamine or galactose (Gal),

especially when they occur in tandem along the same backbone, as it happens in mucins

(14–18). The binding of recombinant Clec4F to the LL mucins was as expected calcium-

dependent and resistant to digestion of non-mucin proteins in the parasite preparation.

Binding was also observed to a number of synthetic glycans representing some of the

main carbohydrate motifs in the *E. granulosus* LL and/or in the LL of the related species

Echinococcus multilocularis (13).

In rodents, Clec4F is exclusively expressed in Kupffer cells (KC) (14–16), the sessile

liver macrophages exposed to the vascular space. This restricted expression means that

Clec4F is not expressed in the mouse DC and macrophage sub-types used in the

previously mentioned experiments that suggested a receptor-independent interaction

mode with LL particles (6, 11, 12). In the ungulate natural hosts of *E. granulosus*, Clec4F

appears to be expressed in lymphoid tissues in addition to the liver (15). In humans,

Clec4F is not functional as a lectin and is poorly or not expressed (15).

The only demonstrated function of Clec4F (in mice) is the removal from circulation of

de-sialylated platelets by KC (19). Although the macrophage galactose-type lectins

(MGL) and the asialoglycoprotein receptor (ASGR; reported to be expressed in KC in

addition to in hepatocytes) collaborate in this function, Clec4F appears to fulfil an

indispensable role (19–21).

KC under basal conditions have a tolerogenic profile as antigen-presenting cells (22).

Also, a role of Clec4F in removing de-sialylated platelets (19), conceptually akin to the

role of the receptors involved in phagocytosis of apoptotic cells (23), suggests that Clec4F

is not wired to inflammatory responses. Thus, Clec4F/KC appears to be an evolutionary

option from the parasite's standpoint for LL debris disposal. However, *in vitro* binding

by the soluble Clec4F lectin domain expressed in dimeric form (13) does not predict

binding by the native cell-surface trimeric receptor, or internalization by KC. In this

article, we analyse *in vivo* uptake of soluble and particulate LL materials by KC and the

role of Clec4F in this process, in the context of artificial injection of materials as well as

experimental infection.

Materials and Methods

Parasite materials

The starting material for LL mucin preparations, and the source of protoscoleces for

experimental infections, were macroscopically healthy, fertile *E. granulosus sensu lato*

hydatids from natural infections in cattle from Uruguay. For LL mucin preparations,

hydatid walls, which have the LL as the overwhelmingly dominant component (3, 4, 8),

were extensively washed with phosphate-buffered saline (PBS), then with 2 M NaCl and

finally with water. The walls were then dehydrated by sequential steps with ethanol (95%

v/v) followed by acetone, and finely ground using a mortar and pestle, followed by

grinding between two polished glass plates (12). The resulting fine powders were

rehydrated in endotoxin-free PBS containing 30 mM EDTA (1 ml for every 2 mg dry

mass) to extract the calcium inositol hexakisphosphate deposits, thus leaving the mucin

meshwork as single major structural component (3). The material was then extensively

washed into endotoxin-free PBS. To obtain a fine particulate preparation (pLL), the

suspension was filtered through 85- and 23-µm gauze (12). To obtain the soluble

preparation (sLL), the suspension was subjected to sonication followed by centrifugation

for 1 h at 100,000 g at 4°C. To obtain sLL in which the terminal monosaccharide residues

were oxidized, prior to the solubilization step, the material was treated with sodium

periodate followed by sodium borohydride (or with sodium borohydride only as mock

treatment) (12). For obtaining biotin-tagged sLL, prior to the solubilization step, the

material in PBS was incubated overnight with N-hydroxysuccinimido-biotin (NHS-

biotin; 10 ng per mg of LL dry mass). Sodium periodate, sodium borohydride and/or

NHS-biotin were removed by extensive washing of the particles into endotoxin-free PBS.

Destruction of terminal monosaccharides by periodate was verified in terms of abrogation

of the binding biotinylated PNA (24), revealed with the help of streptavidin-perodixase,

in a dot-blot format. Biotin incorporation by the sLL was checked using streptavidin-

peroxidase also in dot-blot format. The dry mass content of LL mucin preparations was

estimated as in (12). LL mucin preparations were stored at 4°C in PBS with penicillin-

streptomycin during no more than 6 months (pLL) or 12 months (sLL).

Clec4f⁻ᐟ⁻ mouse line generation

Although Clec4f⁻ᐟ⁻ mice had been previously generated by one of the authors' group

(16), the cost of importing the mice from Taiwan to Uruguay and re-deriving them was

high enough that made *de novo* generation of a Clec4f⁻ᐟ⁻ line by CRISPR-Cas9

technology economically convenient. All animal procedures were performed at the

Specific Pathogen Free animal facility of the Laboratory Animal Biotechnology Unit of

Institut Pasteur de Montevideo. Experimental protocols were approved by the

Institutional Animal Ethics Committee (protocol number 007-18), in agreement with

National law 18.611 and international animal care guidelines (Guide for the Care and

Use of Laboratory Animals) (25) regarding laboratory animal's protocols. Mice were

housed in individually ventilated cages (Tecniplast, Milan, Italy) containing chip

bedding (Toplit 6, SAFE, Augy, France), in a controlled environment at $20 \pm 1°C$ with a

relative humidity of 40-60%, in a 14/10 h light-dark cycle. Autoclaved food (Labdiet

5K67, PMI Nutrition, IN, US) and autoclaved filtered water were administered *ad*

libitum. Cytoplasmic microinjection was performed in C57BL/6J zygotes using a mix of

25 ng/µl Cas9 mRNA (Invitrogen, Carlsbad, CA, US), 15 ng/µl of each sgRNA (2

guides were used) (Synthego, Menlo Park, CA, US), diluted in standard microinjection

buffer (5 mM Tris-HCl, pH 7.4, 0.1 mM EDTA). Viable embryos were cultured

overnight in M16 (Sigma, St Louis, MO, US) microdrops under embryo tested mineral

oil (Sigma), with 5% CO_2 in air at 37 ºC. The next day, 2-cell embryos were transferred

into the oviduct of B6D2F1 0.5 days postcoitum pseudopregnant females (20

embryos/female in average). Standard surgical procedures established in our animal

facility were followed (26). For surgery, recipient females were anesthetized with a

mixture of ketamine (110 mg/kg, Ketonal50, Richmond Veterinaria S.A., Buenos Aires,

Argentina) and xylazine (13 mg/kg, Seton 2%; Calier, Montevideo, Uruguay).

Tolfenamic acid was administered subcutaneously (1 mg/kg, Tolfedine, Vetoquinol,

Madrid, Spain) in order to provide analgesia and anti-inflammatory effects (27).

Pregnancy diagnosis was determined by visual inspection by an experienced animal

caretaker two weeks after embryo transfer, and litter size was recorded on day 7 after

birth. Pups were tail-biopsied and genotyped 21 days after birth, and mutant animals

were maintained as founders. Gene editing efficiency was 53.8% (7 KO/13 born pups).

Genetic modification was confirmed by Sanger sequencing. The Clec4f$^{-/-}$ line was

established after producing F1, F2, and finally homozygous F3 animals that were used

for experiments. Primers used for amplification of the genomic locus were 5'-

TCTTTATGATCGCACCCACA-3', 5'-TCCATTCTCGAGAGCCATCT-3'. F1 and F2

heterozygous mice were selected using a heteroduplex mobility assay on a 10%

polyacrylamide gel. In F2, WT animals were differentiated from homozygous KO

animals by Sanger sequencing.

Mouse experiments

WT BALB/c mice were obtained from DILAVE (MGAP, Uruguay) and kept at the

Instituto de Higiene animal house. Female WT (C57BL/6J) or Clec4f$^{-/-}$ mice were bred

at the Institut Pasteur de Montevideo, co-housed (so as to equilibrate their microbiota) at

the Instituto de Higiene starting at 5 weeks of age for at least 3 weeks, and used between

8 and 14 weeks of age. sLL (100 or 200 μg dry mass per mouse as indicated in each case,

dissolved in 100 μl of PBS) or vehicle only was injected into the tail vein. pLL (450 μg

dry mass per mouse, suspended in 200 μl of PBS) or vehicle only, were injected i.p. Mice

were infected by i.p. injection of 2000 (BALB/c) or 2500 (C57BL/6) protoscoleces

(viability ≥ 95%) suspended in 200 μl of PBS; control mice were injected with vehicle

only. Mice were euthanized using isoflurane. The procedures used were approved by the

Comisión Honoraria de Experimentación Animal (CHEA, Universidad de la República,

Uruguay: protocols 101900-001017-16, 101900-000994-16, 101900-000649-17,

101900-000441-16).

Preparation of liver non-parenchymal fraction for flow cytometry

Following perfusion through the inferior vena cava with 10 mL of PBS, livers were placed

in RPMI and finely chopped using a razor blade for approximately 2 min. The tissue was

then digested in 5 ml per g of enzyme mix (RPMI containing 0.4 U/ml Liberase [Sigma]

and 80 U/ml DNase [Sigma]), for 25 min at 37∘C, in an orbital shaker at 250 rpm with

additional manual shaking every 5 min. Digests were poured through a 100 μm strainer

and centrifuged at 300 g for 5 min; the cell pellets were re-suspended in 25 ml of RPMI

and again centrifuged at 300 g. The pellets were suspended in 2 ml of RBC lysis buffer

(Sigma), and after 2 min, 2 ml of FACS buffer (PBS supplemented with 0.5% BSA and

2 mM EDTA) was added. After another centrifugation step (300 g, 5 min), cells were re-

suspended in 3 ml of FACS buffer, passed through a 40 μm strainer, and counted using

the Nexcelom Cellometer K2 with the AOPI stain for cell viability. Cells and buffer used

with the cells were kept at 4ºC, except for the digestion step.

Flow cytometry

Five hundred thousand live cells where incubated with the viability probe for 10 min at

room temperature, then with 0.025 μg of anti-CD16/32 (Biolegend) in 10% v/v normal

rat serum (Unidad de Reactivos y Biomodelos de Experimentación, Facultad de

Medicina, Universidad de la República). Cells were then surface-stained, fixed in 1% w/v

PFA for 20 min at 4°C, washed, and permeabilized overnight with permeabilisation

buffer (Thermo) for intracellular staining. The details of the viability probes, antibodies

and lectin probes are given in Supplementary Table 1. Finally, cells were washed, spun

231 at 350 g for 5 min and re-suspended in FACs buffer. The data were acquired in a FACS

232 Canto II Cytometer (BD), at the Instituto de Higiene, Universidad de la República, and

233 then analysed using the FlowJo software package. The relative contribution of each liver

234 non-parenchymal cell type under study to the uptake of LL materials was estimated by

235 multiplying the frequency of cells positive for the LL probe and belonging to the

236 particular cell type (within all live single cells) by their arithmetic mean fluorescence

237 intensity for the probe and then dividing this figure over the summation of the analogous

238 products for the 3 cell types analysed.

240 *Histology*

242 Tissue pieces (approximately 0.5 mm^3) were fixed in buffered formalin. Liver samples

243 corresponded to animals subjected to liver perfusion with PBS (as the materials were also

244 analyzed by flow cytometry). Paraffin-embedded tissue sections (2 μm) were probed with

245 biotinylated monoclonal antibody E492 (3 μg/mL; steamer pH 6.1) or biotinylated PNA

246 (Vector Laboratories, Newark, California, United States, B-1075; dilution 1:100; steamer

247 pH 9.0) using the avidin-biotin-complex method (28), and counterstained using

248 haemalum.

250 *ELISA*

252 ELISA wells were coated with sLL at 10 μg/mL (total dry mass) in Tris-buffered saline

253 (TBS) overnight at 4 ºC, and blocked using TBS containing 0.05% w/v Tween 20, 1%

254 w/v bovine serum albumin and 0.5% v/v Carbo-Free^{TM} blocking solution (Vector

255 Laboratories). Wells were then probed with biotinylated E492 antibody (29), biotinylated

peanut agglutinin (PNA; Vector laboratories) or recombinant soluble Clec4F (expressed

as a fusion with the F_c fragment of human IgG_1 (30)). The biotinylated PNA or E492

antibody were detected using streptavidin coupled to peroxidase (ThermoFisher

Scientific) and the recombinant Clec4F was detected using a goat anti-human IgG

coupled to peroxidase (AbCam, Cambridge, UK). For competition ELISA, recombinant

Clec4F at 80 ng/mL was mixed with variable concentrations of either (biotinylated) PNA

or E492 antibody as competitors, and bound Clec4F detected as previously described.

Particle immunofluorescence

pLL particles were incubated with biotinylated E492 antibody (0.6 µg/mL), biotinylated

PNA (1:1500) or recombinant soluble Clec4F (7.5 µg/mL), followed by streptavidin-

Cyanine-3 (1:500; Amersham Bioscience) or by goat anti-human IgG coupled to FITC

(1:100, Sigma), as appropriate. Control stains were included using human IgG_1 instead

of recombinant Clec4F as well omitting the primary staining reagents.

Statistical analyses

As the underlying data do not satisfy the criteria for parametric statistics, rank-based

nonparametric statistical methods, were used throughout. When the numbers of mice in

each treatment-repeat experiment combination were equal, the Mack-Skillings exact test

for a two-way layout was used (31). When the data did not fulfil the above condition, data

from the different repeat experiments were pooled and the modified Wilcoxon-Mann-

Whitney test (only two experimental groups) (53) or the Kruskal-Wallis test (more than

two experimental groups) were used. When the Mack-Skillings or the Kruskal-Wallis

tests resulted in a statistic with a p-value less than 0.05, the test was followed by the post-

hoc multiple comparison test described by Conover (55) and the Benjamini-Hochberg

correction for controlling the false discovery rate (56). Throughout the paper, the symbols

*, ** and *** represent p-values less than 0.05, 0.01 and 0.001 respectively.

Results

Soluble LL mucins are taken up by Kupffer cells in vivo *in a manner dependent on the*

mucin glycans

We first sought to determine whether KC take up soluble LL mucins (sLL) *in vivo* and

whether this is dependent on the mucin glycans. We thus treated the LL mucins either

with sodium periodate to oxidize the glycans or applied a mock treatment, tagged both

preparations with equivalent amounts of biotin (Fig. S1), and injected both preparations

i.v. in mice. Twenty minutes later, we quantitated the biotin tag in permeabilized non-

parenchymal liver cells using streptavidin conjugated to PE and flow cytometry. KC were

identified using our previously validated gating strategy (Fig. S2) (32). A significant

signal was detected in KC, and the signal was markedly higher for mock-treated than for

periodate-treated sLL (Fig. 1). No significant biotin signal was detected in either LSECs

or in non-KC CD11b$^+$ cells, a broad definition that encompasses any mononuclear

phagocytes other than KC present in the liver (Fig. 1 a, b, and S2).

The plant lectin PNA and the E492 monoclonal antibody allow detection of laminated

layer mucins captured by liver cells in vivo.

Except in experiments using periodate-treated LL materials, the detection of LL materials

by appropriate carbohydrate-binding probes was preferable to prior labelling with biotin.

Among other advantages, such probes would allow the detection of endogenous LL

mucins in infected animals. One candidate probe was the plant lectin PNA, which binds

Galβ1-3-GalNAcα1-peptide, i.e. the non-decorated mucin O-glycan core 1 or T-antigen

(33). This motif is highly represented in the *E. granulosus* LL, and accordingly, PNA

binds the LL mucins in histological sections, particle lectin fluorescence, and Western

blotting (12, 24). The second candidate probe was the monoclonal antibody E492, raised

against *E. granulosus* protoscoleces (34). This antibody binds to LL mucins from *E.*

multilocularis (35) and to synthetic glycans that include the P_1 blood antigen motif

(Galα1-4Galβ1-4GlcNAc), highly represented among *E. granulosus* LL mucin glycans

(7, 8). A third potential probe was recombinant Clec4F (13). We verified that these

potential probes indeed bind to sLL coated on ELISA plates as well as to LL particles in

suspension (Fig. 2 a, b).

We took advantage of the defined specificities of PNA and the E492 antibody to gain

insight into the carbohydrate motifs bound by Clec4F in the LL mucins. The binding of

recombinant Clec4F to the solubilized and plate-bound LL mucins was not competed by

PNA, but it was strongly competed by the E492 antibody (Fig. 2 c). Therefore, most of

the binding of Clec4F to the native LL mucins appears to rely on recognition of the P_1

motif, whereas recognition of the T-antigen is not significant. This in turn suggests that

in the LL mucins the T-antigen may not be exposed and/or be spaced inadequately for

Clec4F binding, as recombinant Clec4F does bind the T-antigen in glycan arrays (15–17).

In vivo probes should show minimal reactivity with host tissues; for our purposes, lack of

reactivity with the populations of mouse non-parenchymal liver cells under study was the

most important consideration. Neither PNA nor the E492 antibody showed significant

binding in flow cytometry to KC or LSEC from control mice (Fig. S3). PNA bound to a

very minor fraction of non-KC CD11b$^+$ cells, whereas the antibody did not bind. PNA,

but not the E492 antibody, bound to a major sub-population of CD45$^+$ CD11b$^-$ F4/80$^-$

cells, which were not of particular interest in our study. Recombinant Clec4F bound to

cells in all 4 cell gates defined, most strongly to LSEC and KC (Fig. S3). This probably

reflects the fairly broad specificity of Clec4F for glycans terminated in Gal/GalNAc (15–

17), and ruled out this lectin as a probe for our purposes.

Finally, to determine whether uptake of sLL could be measured *in vivo* using PNA or the

E492 antibody, we injected wild-type (WT) C57BL/6J mice i.v. with sLL and

subsequently stained isolated hepatic non-parenchymal cells intracellularly with either

probe. As expected, no staining was observed in cells from mice injected with PBS only,

whereas KC, and to a small extent also LSEC, from mice given sLL stained intracellularly

with either PNA or E492 antibody (Fig. 2 d). Omitting the cell permeabilization step

caused the PNA signal in KC and LSEC of sLL-injected mice to be reduced to negligible

levels (data not shown), ruling out the remote possibility that injection of sLL induces the

expression of ligands for the probe in the glycocalix of the cells under study. Both PNA

and the E492 antibody appeared to allow more sensitive detection than prior labelling

with biotin (Fig. 1), so that captured sLL was detected in almost all KC and also in some

LSEC.

In summary both PNA and the E492 antibody are useful probes for LL mucins captured

by mouse non-parenchymal liver cells.

Soluble LL mucins are taken up by Kupffer cells in vivo *in a manner dependent on Clec4F*

We first used CRISPR/Cas9 to generate a new Clec4f $^{-/-}$ line on the C57BL/6J

background. These Clec4f $^{-/-}$ mice have 8-bp deletion expected to result in a polypeptide

that is truncated near the N-terminus (Fig. 3, a, b), and effectively lack Clec4F protein

expression by flow cytometry (Fig. 3 c). Hence, our line appears similar to Clec4f gene-

deficient mice that have already been described (16, 19).

We next compared WT and Clec4f $^{-/-}$ mice in terms of the *in vivo* uptake of sLL by liver

non-parenchymal cells. For this purpose, we injected untagged sLL, either untreated or

subjected to the mock treatment used in the previous experiments that involved periodate

oxidation. Uptake of sLL by KC was strongly dependent on Clec4F (Fig. 4 a - c),

particularly when assessed in terms of mean fluorescence intensity. The low-level uptake

by LSEC observed was as expected independent of Clec4F; in fact, for untreated sLL,

uptake by LSEC was slightly enhanced in the absence of Clec4F, possibly as a result of

the decreased capture by KC. No uptake by non-KC CD11b^{+} cells was detected (Fig. 4 a,

b and Fig. S4). Through a calculation that takes into account the numbers of each cell

type capturing sLL and the intensity of the signal indicating intracellular sLL, KC were

estimated to be responsible for the overwhelming majority of sLL uptake by non-

parenchymal liver cells in WT mice (Fig. 4 d). However, in the absence of Clec4F, LSEC

played a significantly, if slightly, more important part in the uptake of sLL than in the

presence of the receptor. Differences in the uptake of untreated *vs* mock-treated material

were not observed, except in the low-level uptake by LSEC in Clec4f$^{-/-}$ mice (Fig. 4 a).

For consistency with the results shown in Figure 1, we chose to use mock-treated material

in the following experiments.

It can be expected that in the absence of Clec4F, other mechanisms of cellular uptake of

sLL may become apparent, e.g. macropinocytosis or uptake dependent on natural

antibodies or complement (6). Such Clec4F-independent uptake of sLL was hinted by the

previous observations in relation to LSEC, as well as by the 25 – 60 % KC in Clec4f$^{-/-}$

mice positive for PNA at low levels (Fig. 4 a - c). As the Clec4F-independent mechanisms

might become dominant at longer analysis times, we carried out a similar experiment with

a 22 h analysis time. Uptake of sLL was still dominated by KC among the cell types

studied, and this uptake was again strongly dependent on Clec4F (Fig. 5 a –d). In the

absence of Clec4F, LSEC made a significantly larger contribution to sLL uptake than in

the presence of the receptor, but this contribution was still minimal (Fig. 5 d). Non-KC

CD11b^{+} cells did not take up material detectably (Fig. 5 a – d and S4). Cell surface

expression of Clec4F in KC was not altered by sLL injection (Fig. S5).

In summary, soluble circulating LL mucins in naïve mice are taken up mainly by KC

among liver non-parenchymal cells and mostly in Clec4F-dependent fashion.

LL mucins injected in particulate form are taken up by *Kupffer cells* in vivo *in Clec4F-*

dependent manner

Although the release of soluble mucins from the LL during cystic echinococcosis is a

possibility (6), only particles have been documented to be shed from the LL in human

infections (6, 10). It is not known if these particles circulate systemically and/or give rise

to soluble materials that circulate. It was therefore important to determine whether Clec4F

participates in the uptake by KC of LL mucins released in a particulate presentation into

the internal milieu and at an anatomical site distant from the liver. We thus injected i.p.

particulate LL mucins (pLL) (12), and analysed the liver non-parenchymal fraction 22 h

later. LL material was indeed detected by flow cytometry in KC (Fig. 6 a – c and S6 a -

c). Uptake by KC was dependent on Clec4F, although there was a considerable degree of

Clec4F-independent uptake as well. In addition to low-level uptake by LSEC, in this

experimental system, non-KC CD11b$^+$ cells were observed to take up the LL material.

Although non-KC CD11b$^+$ cells may include recruited monocytes/macrophages, the

frequency of this population (within all non-parenchymal cells) did not increase as a result

of pLL injection (Fig. S7). As expected, uptake by LSEC and by non-KC CD11b$^+$ cells

was independent of Clec4F and in fact tended to be enhanced in the absence of the

receptor (Fig. 6 a – c and S6 a - c). Indeed, whereas in WT mice the overwhelming

majority of the LL material uptake by liver non-parenchymal cells was carried out by KC,

in Clec4f$^{-/-}$ mice the two non-KC cell types studied taken together played a significantly

larger role than in WT mice (Fig. 6 d and S6 d). The cell surface expression of Clec4F in

KC was not altered by injection of pLL (Fig. S8).

The mostly Clec4F-dependent uptake of the LL material was confirmed by histology,

using the E492 antibody as a probe. Strong staining was observed in KC-like cells of WT

animals, whereas only minimal staining was observed in the KC-like cells of Clec4f$^{-/-}$

animals (Fig. 6 e). Animals injected with pLL also showed weak staining in hepatocytes

(a cell type not analysed by flow cytometry), and this was independent of Clec4F. As

hepatocytes are unlikely to phagocytose large particles, this observation suggests the

particles injected release small particles and/or soluble mucin materials *in vivo* that

preferentially circulate systemically.

In summary, LL mucins present within tissues in particulate form can generate materials

(particulate and/or soluble) that circulate systemically and that are captured by KC via

Clec4F.

In experimental E. granulosus *infection, LL mucins circulate and are captured selectively*

by Kupffer cells, with the participation of Clec4F

We next wished to determine if Clec4F was relevant in the infection setting. The most

widely used experimental infection model for *E. granulosus* uses intraperitoneal injection

with protoscoleces, parasite stages that usually develop into adult worms but are capable

of reverse development into hydatids (1, 36). In this model, hydatids develop free or

within host collagen capsules in the peritoneal cavity (37). KC uptake of LL materials in

this model would require that such materials reach systemic circulation, a possibility for

which there was no evidence, in experimental or natural infections. We first analysed

infected mice of the BALB/c strain, in which experimental infection reaches higher

parasite burdens than in the C57BL/6 strain (38). Indeed, the BALB/c mice analysed had

a median of 25 mL of total parasite volume (Fig. S9 a), an extremely high parasite burden.

In these mice, using the E492 antibody as probe, we detected LL materials by histology

in peritoneal cavity cells (i.e. the site of infection) and the liver, but not in kidneys,

intestine or lungs (Fig. 7 a). In addition, in some of the infected mice, occasional positive

cells were observed in the spleen, and less frequently in the mesenteric lymph nodes (Fig.

7 a). The use of PNA as probe yielded similar results, but it was complicated by the

expected signal in the lymphoid organs (39) unrelated to LL material, i.e. detectable in

non-infected mice (data not shown).

The above results show that in infected animals LL mucins circulate systemically and

become selectively concentrated in the liver. Within the liver, the signal for LL mucins

was specifically concentrated in KC-like cells (Fig. 7 a). The LL signal in KC-like cells

was observed in 12 out of 15 infected mice analysed. Contrary to previous experiments

injecting pLL, no signal was detected in hepatocytes. This may be related to lower loads

of LL material, as suggested by the lower intensity in the staining in KC in the infection

setting in comparison to the pLL injection experiments (compare Fig. 6 e and 7 a).

The selective LL material uptake by KC among the liver non-parenchymal cells was

confirmed by flow cytometry (Fig. 7 b – d and S10). Although the data did not reach

statistical significance, a sub-population of non-KC CD11b^{+} cells appeared to take up LL

materials (Fig. 7 b – d). Also, the frequency of non-KC CD11b^{+} cells within all liver non

parenchymal cells was significantly increased by infection, suggesting cell recruitment

(Fig. S11 a). The cell surface expression of Clec4F in KC was not altered by infection

(Fig. S12 a).

We next analysed infected C57BL/6 WT and Clec4f$^{-/-}$ mice. In these mice, the parasite

burdens were low: the median burden in the WT C57BL/6 mice was 0.5 mL (this figure

corresponds to the two infection experiments fully analysed, as an additional experiment

with very low parasite burdens was excluded from analysis). Although the median

parasite burden was 3-fold lower in Clec4f$^{-/-}$ mice than in WT mice, the trend did not

reach statistical significance (Fig. S8 b). By flow cytometry, uptake of LL material by

KC was detected, in WT but not in Clec4F$^{-/-}$ mice (Fig. 7 e - g). No significant uptake

was detected in LSECs (Fig. 7 e, f and S10 b). In non-KC CD11b$^+$ cells, there was a trend

towards low-level, Clec4F-independent uptake (Fig. 7 e - g), which did not reach

significance. The frequency of non-KC CD11b$^+$ cells was not significantly altered by

infection (Fig. S 11 b). Again, Clec4F surface expression in the KC of WT mice was not

altered by infection (Fig. S12 b).

Discussion

In this article we show that native LL materials shed from *E. granulosus sensu lato*

hydatids are taken up selectively by KC, in a fashion mostly dependent on Clec4F. It is

thought that pathogens capable of immune evasion engage innate receptors that normally

bind altered self, as opposed to receptors specialized in detecting pathogens and triggering

inflammatory responses (40). *E. granulosus* is well adapted for immune evasion (2), and

the fact that its most abundant surface molecules engage a receptor involved in the

clearing of aged host platelets (19, 20) seems to be in line with the above principle.

KC are specialized in removing unwanted materials from the blood. Our results imply

that the LL mucins (which even after their artificial solubilisation behave as having very

high molecular sizes (24)) have the capacity to circulate systemically, thus reaching KC

from a distant anatomical site. This applies to exogenous LL mucins initially present in

the peritoneal cavity in particulate format, which must therefore either circulate as

particles and/or partly solubilize *in vivo*. It also applies to the LL materials released by

live parasites growing in the peritoneal cavity.

In addition to KC, both LSEC and certain non-KC CD11b$^+$ cells were observed to take

up LL materials. This was in both cases independent of Clec4F, and in some experimental

settings it was enhanced in the absence of Clec4F (Fig. 4 d, 6 d and S6 d). Uptake of

materials by non-KC CD11b$^+$ cells was apparent after injection of LL mucins in

particulate presentation (pLL; Fig. 6 a – d and S6) but not in a soluble presentation (sLL;

Fig. 1, 4 and 5). This contrast suggests that the species released *in vivo* from the large LL

particles and capable of circulating are (or at least include) small particles, as opposed to

fully soluble mucins like those making up sLL. The fact that uptake by the non-KC

CD11b$^+$ population was clear after injection of a high-dose bolus of pLL whereas it was

merely hinted in experimental infection (Fig. 7 b - g) is most probably explained by this

uptake mechanism becoming relevant only when the uptake capacity of KC is

overwhelmed. It is unlikely that the non-KC CD11b$^+$ cell population taking up LL

materials derives from peritoneal cavity macrophages, as it is composed of F4/80$^-$ cells

(Fig. 6 c and S6 c), whereas most resident peritoneal macrophages as well as peritoneal

macrophages reported to enter the liver upon injury to this organ are F4/80^{hi} (41). We

instead favour the possibility that the population being discussed corresponds to

monocytic cells present in the liver or recruited in response to LL particles physically

trapped in the liver sinusoids.

Hepatocytes could be an additional target cell for LL materials in the liver, as suggested

by the weak but significant staining observed in this cell type after pLL injection (Fig. 6

e). The obvious candidate receptor to mediate this apparent uptake is the ASGR (42).

Ligand uptake by hepatocytes via the ASGR is reported to have an upper size limit of 70

nm (43). Thus the signal in hepatocytes observed in our experiments with pLL (which

has particle sizes in the micrometer range (12)) may correspond to uptake of soluble

mucins released from the particles *in vivo*. In the same experiments as well as in the

infection experiments (Fig. 7), the preferential uptake by KC in relation to hepatocytes

surely reflects the differential capacity of KC to phagocytose particles (even if

additionally, Clec4F may bind LL mucins better than the ASGR, as suggested by *in vitro*

evidence (13)). Our results do not allow to weigh the importance of hepatocytes in the

capture of fully soluble injected LL materials (sLL).

Part of the uptake of LL materials by KC was observed to be Clec4F-independent (Fig. 5

a, 6 a, b, and 7 e). Clec4F-independent uptake by KC, as well as by non-KC CD11b[+] cells

and LSEC as previously mentioned, may be mediated by natural anti-carbohydrate

antibodies that bind the LL mucins, either directly or via complement activation (44). In

the context of chronic experimental infection, the possibilities for Clec4F-independent

uptake are even broader than after LL mucin injection. In the first place, the mucins will

be opsonized by specific (homologous) antibodies (6, 44). In second place, contrary to

the artificially prepared (soluble and insoluble) mucin preparations, LL materials shed

from live parasites must contain the major non-mucin structural component of the LL,

i.e. calcium inositol hexakisphosphate (calcium InsP_6) (45–47). Calcium InsP_6 binds host

C1q and initiates limited activation of complement (44, 48). In fact, the observed uptake

of LL materials by phagocytes at the experimental infection site (Fig. 7 a) must also be

Clec4F-independent, as peritoneal macrophages are not expected to express Clec4F (16).

Given the ample possibilities of Clec4F-independent capture, it is noteworthy the Clec4F

nonetheless has a clearly detectable role in LL material uptake (by KC), especially in the

infection context.

As mentioned, in human infections affecting solid organs, LL debris accumulate in the

parasite's vicinity (10). Our results show the presence of LL materials at the experimental

infection site but also as mentioned the capture of these materials by the liver (Fig. 7 a).

This implies that the accumulation of LL debris in the parasite's vicinity is moderated by

the circulation and capture by the liver of materials derived from this debris. This is

important because, although moderate amounts of LL particles probably have immune

modulatory effects that favour the parasite, large amounts of LL particles accumulating

at the infection site would probably foster inflammation (11, 12, 49, 50). Inflammation

might also be brought about the LL materials remaining in circulation and possibly being

deposited in the glomeruli or other sites. Indeed, if as we speculated above, monocytes

are recruited to the liver in response to trapped LL particles, then Clec4F-dependent KC

capture may be important to avoid liver inflammation. This aspect may not have been

fully captured in our flow cytometry and histology experiments because of liver perfusion

prior to analysis.

The events downstream of LL material capture by KC are certainly worth studying. KC

can present antigens to T cells present in the vascular space, and this normally leads to

tolerogenic responses that can have a systemic impact (22, 51, 52). The analysis of

immune responses in WT and Clec4f$^{-/-}$ mice infected with *E. granulosus* will be the

subject of a future communication.

Ancestrally, the larval stages of the parasites belonging to the taeniid family (including

those of the genus *Echinococcus*) probably infected the liver of rodent hosts (5, 53). In

this context, the interaction of materials shed by the parasites with a receptor (Clec4F)

specifically expressed by phagocytic cells present in the same organ (KC) may have been

evolutionarily advantageous. This may have been so because LL material uptake by KC

would have avoided particle build-up and consequent inflammation in the liver sinusoids

and/or other sites, and/or because KC are poised for tolerogenic antigen presentation. The

extant species *E. multilocularis* dwells almost exclusively in the liver of rodents (besides

accidentally infecting humans). The LL of this species also interacts with Clec4F, as

suggested by experiments using recombinant Clec4F (13) and by the presence in this

species' LL of the blood antigen P_1 motif (54). Thus, host Clec4F is likely an important

player in *E. multilocularis* infection.

The natural hosts of larval *E. granulosus* are mostly ungulates, including sheep, cattle,

pigs, goats, camels and horses. Like in rodents, in ungulates Clec4F also appears to be

expressed in the liver (15). Therefore, the uptake of LL materials by KC via Clec4F

probably takes place also in the ungulate natural hosts. However, in ungulates Clec4F

appears to be additionally expressed in secondary lymphoid organs (15), opening the

possibility that non-KC, anatomically more widely distributed, phagocytes may act as

additional sinks for LL materials.

As mentioned, humans do not express Clec4F with a functional lectin domain (15).

Humans are normally a dead-end for this parasite's life-cycle, meaning that human

Echinococcus infection has not been shaped through evolution. Although *E.*

multilocularis infection in humans is usually accompanied by strong local inflammation

(and tissue destruction), this is not usually the case with *E. granulosus* (9, 55). It is a

possibility that for *E. granulosus*, which grows more slowly than *E. multilocularis*,

Clec4F-independent clearance mechanisms are in most infection settings sufficient to

avoid excessive build-up of LL debris. It is conceivable that such Clec4F-independent

mechanisms in humans include uptake via lectins expressed in KC with specificities

related to Clec4F. As mentioned, the uptake of de-sialylated platelets by mouse KC

requires Clec4F as well as MGLs or ASGR (19, 20). It seems likely that a mechanism for

de-sialylated platelet uptake by KC based on a set of collaborating lectins exists

throughout mammals, even if in humans such a system does not include Clec4F. Our

previous screening using recombinant receptors yielded inconclusive results with respect

to the capacity of human MGL and ASGR to interact with the LL mucins (13). When

probing synthetic sugars, human ASGR and MGL appeared to bind less well than Clec4F

to the P_1 motif (after normalizing by binding to a known target of all 3 receptors) (13).

However, given the sheer abundance and significant breadth of Gal-based motifs in the

LL, it would not be surprising if ASGR and/or MGL were able to internalize LL materials

in the absence of Clec4F. It is important to determine if KC in liver samples from humans

infected with *E. granulosus*, as well as with *E. multilocularis*, stain for LL materials.

Statement of Ethics

Mouse studies followed the ARRIVE guidelines. Experimental protocols were

individually reviewed and approved by the Institutional Animal Ethics Committees of the

Institut Pasteur Montevideo or the Universidad de la República, Uruguay, as detailed in

the Materials and Methods section.

Conflict of Interest Statement

The authors declare no conflict of interest.

Funding sources

This work was funded by Agencia Nacional de Investigación e Innovación (ANII;

Government of Uruguay) grant FCE 136130 (to AD), by a Travel Grant (to SJJ) from the

Science and Innovation Fund of the Foreign and Commonwealth Office (UK

Government), and by PEDECIBA (Government of Uruguay). AB and CM were

supported by PhD scholarships from Comisión Académica de Posgrado (Universidad de

la República, Uruguay) and ANII respectively.

Author contributions

AB: conceptualization, investigation. CM, LS, JN, MS: investigation. GS, MNM, APM,

MC: methodology. TLH, SLH, GME: resources. CG: formal analysis (statistics). TFEB,

CC: conceptualization, supervision. SJJ: conceptualization, funding acquisition,

supervision, writing – review & editing. AD: conceptualization, funding acquisition,

project administration, supervision, writing – original draft preparation.

References

References

1. Thompson RCA. 2017. Biology and Systematics of Echinococcus. Adv Parasitol
95:65–109.

2. Díaz Á. 2017. Immunology of cystic echinococcosis (hydatid disease). British
Medical Bulletin 124:121–133.

3. Díaz A, Casaravilla C, Irigoín F, Lin G, Previato JO, Ferreira F. 2011.
Understanding the laminated layer of larval Echinococcus I: structure. Trends in
Parasitology 27:204–213.

4. Díaz A, Casaravilla C, Allen JE, Sim RB, Ferreira AM. 2011. Understanding the
laminated layer of larval Echinococcus II: immunology. Trends in Parasitology
27:264–273.

5. Díaz Á, Fernández C, Pittini Á, Seoane PI, Allen JE, Casaravilla C. 2015. The
laminated layer: Recent advances and insights into Echinococcus biology and
evolution. Experimental Parasitology 158:23–30.

6. Díaz A, Barrios AA, Grezzi L, Mouhape C, Jenkins SJ, Allen JE, Casaravilla C.
2022. Immunology of a unique biological structure: the Echinococcus laminated
layer. Protein & Cell https://doi.org/10.1093/procel/pwac023.

7. Lin G, Todeschini AR, Koizumi A, Neves JL, González H, Dematteis S, Hada N,
Previato JO, Ferreira F, Mendonça-Previato L, Díaz A. 2013. Further structural
characterization of the Echinococcus granulosus laminated layer carbohydrates:
The blood-antigen P1-motif gives rise to branches at different points of the O-
glycan chains. Glycobiology 23:438–452.

8.   Díaz A, Fontana EC, Todeschini AR, Soulé S, González H, Casaravilla C, Portela

M, Mohana-Borges R, Mendonça-Previato L, Previato JO, Ferreira F. 2009. The

Major Surface Carbohydrates of the *Echinococcus granulosus* Cyst: Mucin-Type

O -Glycans Decorated by Novel Galactose-Based Structures. Biochemistry

48:11678–11691.

9.   Díaz Á, Sagasti C, Casaravilla C. 2018. Granulomatous responses in larval taeniid

infections. Parasite Immunol 40:e12523.

10.   Grimm J, Nell J, Hillenbrand A, Henne-Bruns D, Schmidberger J, Kratzer W,

Gruener B, Graeter T, Reinehr M, Weber A, Deplazes P, Möller P, Beck A, Barth

TFE. 2020. Immunohistological detection of small particles of Echinococcus

multilocularis and Echinococcus granulosus in lymph nodes is associated with

enlarged lymph nodes in alveolar and cystic echinococcosis. PLoS Negl Trop Dis

14:e0008921.

11.   Pittini Á, Martínez-Acosta YE, Casaravilla C, Seoane PI, Rückerl D, Quijano C,

Allen JE, Díaz Á. 2019. Particles from the Echinococcus granulosus Laminated

Layer Inhibit CD40 Upregulation in Dendritic Cells by Interfering with Akt

Activation. Infect Immun 87:e00641-19.

12.   Casaravilla C, Pittini A, Rückerl D, Seoane PI, Jenkins SJ, MacDonald AS,

Ferreira AM, Allen JE, Díaz A. 2014. Unconventional maturation of dendritic cells

induced by particles from the laminated layer of larval Echinococcus granulosus.

Infect Immun 82:3164–3176.

13.   Hsu T-L, Lin G, Koizumi A, Brehm K, Hada N, Chuang P-K, Wong C-H, Hsieh

S-L, Díaz A. 2013. The surface carbohydrates of the Echinococcus granulosus

larva interact selectively with the rodent Kupffer cell receptor. Molecular and

Biochemical Parasitology 192:55–59.

14. Fadden AJ, Holt OJ, Drickamer K. 2003. Molecular characterization of the rat

Kupffer cell glycoprotein receptor. Glycobiology 13:529–537.

15. Taylor ME, Snelling T, Smith DF, Drickamer K. 2019. Absence of a human

ortholog of rodent Kupffer cell galactose-binding receptor encoded by the CLEC4f

gene. Glycobiology 29:332–345.

16. Yang C-Y, Chen J-B, Tsai T-F, Tsai Y-C, Tsai C-Y, Liang P-H, Hsu T-L, Wu C-

Y, Netea MG, Wong C-H, Hsieh S-L. 2013. CLEC4F Is an Inducible C-Type

Lectin in F4/80-Positive Cells and Is Involved in Alpha-Galactosylceramide

Presentation in Liver. PLoS ONE 8:e65070.

17. Coombs PJ, Taylor ME, Drickamer K. 2006. Two categories of mammalian

galactose-binding receptors distinguished by glycan array profiling. Glycobiology

16:1C-7C.

18. Ouyang Z, Felix J, Zhou J, Pei Y, Ma B, Hwang PM, Lemieux MJ, Gutsche I,

Zheng F, Wen Y. 2020. Trimeric structure of the mouse Kupffer cell C-type lectin

receptor Clec4f. FEBS Lett 594:189–198.

19. Jiang Y, Tang Y, Hoover C, Kondo Y, Huang D, Restagno D, Shao B, Gao L,

Michael McDaniel J, Zhou M, Silasi-Mansat R, McGee S, Jiang M, Bai X, Lupu

F, Ruan C, Marth JD, Wu D, Han Y, Xia L. 2021. Kupffer cell receptor CLEC4F

is important for the destruction of desialylated platelets in mice. Cell Death Differ

28:3009–3021.

20.   Deppermann C, Kratofil RM, Peiseler M, David BA, Zindel J, Castanheira FVES,

715   van der Wal F, Carestia A, Jenne CN, Marth JD, Kubes P. 2020. Macrophage

716   galactose lectin is critical for Kupffer cells to clear aged platelets. J Exp Med

717   217:e20190723.

718   21.   Li Y, Fu J, Ling Y, Yago T, McDaniel JM, Song J, Bai X, Kondo Y, Qin Y,

719   Hoover C, McGee S, Shao B, Liu Z, Sonon R, Azadi P, Marth JD, McEver RP,

720   Ruan C, Xia L. 2017. Sialylation on O-glycans protects platelets from clearance by

721   liver Kupffer cells. Proc Natl Acad Sci USA 114:8360–8365.

722   22.   Heymann F, Peusquens J, Ludwig-Portugall I, Kohlhepp M, Ergen C, Niemietz P,

723   Martin C, van Rooijen N, Ochando JC, Randolph GJ, Luedde T, Ginhoux F, Kurts

724   C, Trautwein C, Tacke F. 2015. Liver inflammation abrogates immunological

725   tolerance induced by Kupffer cells. Hepatology 62:279–291.

726   23.   Rothlin CV, Carrera-Silva EA, Bosurgi L, Ghosh S. 2015. TAM receptor signaling

727   in immune homeostasis. Annu Rev Immunol 33:355–391.

728   24.   Casaravilla C, Díaz A. 2010. Studies on the structural mucins of the Echinococcus

729   granulosus laminated layer. Molecular and Biochemical Parasitology 174:132–

730   136.

731   25.   (1)   Committee for the update of the guide for the care and use of laboratory

732   animals. 2011. Guide for the care and use of laboratory animals, 8th ed. National

733   Academies Press.

734   26.   Meikle MN, Schlapp G, Menchaca A, Crispo M. 2018. Minimum volume Spatula

735   MVD vitrification method improves embryo survival compared to traditional slow

freezing, both for in vivo and in vitro produced mice embryos. Cryobiology

84:77–81.

27.   Schlapp G, Goyeneche L, Fernández G, Menchaca A, Crispo M. 2015.

Administration of the nonsteroidal anti-inflammatory drug tolfenamic acid at

embryo transfer improves maintenance of pregnancy and embryo survival in

recipient mice. J Assist Reprod Genet 32:271–275.

28.   Barth TFE, Herrmann TS, Tappe D, Stark L, Grüner B, Buttenschoen K,

Hillenbrand A, Juchems M, Henne-Bruns D, Kern P, Seitz HM, Möller P, Rausch

RL, Kern P, Deplazes P. 2012. Sensitive and Specific Immunohistochemical

Diagnosis of Human Alveolar Echinococcosis with the Monoclonal Antibody

Em2G11. PLoS Negl Trop Dis 6:e1877.

29.   Baz A, Richieri A, Puglia A, Nieto A, Dematteis S. 1999. Antibody response in

CD4-depleted mice after immunization or during early infection with

Echinococcus granulosus. Parasite Immunol 21:141–150.

30.   Hsu T-L, Cheng S-C, Yang W-B, Chin S-W, Chen B-H, Huang M-T, Hsieh S-L,

Wong C-H. 2009. Profiling carbohydrate-receptor interaction with recombinant

innate immunity receptor-Fc fusion proteins. J Biol Chem 284:34479–34489.

31.   Hollander M, Wolfe DE, Chicken E. 2015. Nonparametric Statistical Methods, 3rd

ed. John Wiley & Sons, New York.

32.   Lynch RW, Hawley CA, Pellicoro A, Bain CC, Iredale JP, Jenkins SJ. 2018. An

efficient method to isolate Kupffer cells eliminating endothelial cell contamination

and selective bias. J Leukoc Biol 104:579–586.

33. Cummings RD, Etzler, ME. Antibodies and lectins in glycan analysis, p. 633–647.

In Essentials of GlycobiologySecond Edition. Cold Spring Harbor Laboratory

Press.

34. Dematteis S, Pirotto F, Marqués J, Nieto A, Orn A, Baz A. 2001. Modulation of

the cellular immune response by a carbohydrate rich fraction from Echinococcus

granulosus protoscoleces in infected or immunized Balb/c mice. Parasite Immunol

23:1–9.

35. Walker M, Baz A, Dematteis S, Stettler M, Gottstein B, Schaller J, Hemphill A.

2004. Isolation and characterization of a secretory component of Echinococcus

multilocularis metacestodes potentially involved in modulating the host-parasite

interface. Infect Immun 72:527–536.

36. Araj GF, Matossian RM, Frayha GJ. 1977. The host response in secondary

hydatidosis of mice. I. Circulating antibodies. Z Parasitenkd 52:23–30.

37. Richards KS, Arme C, Bridges JF. 1983. Echinococcus granulosus equinus: an

ultrastructural study of murine tissue response to hydatid cysts. Parasitology 86 (Pt

3):407–417.

38. Mourglia-Ettlin G, Cucher M, Arbildi P, Rosenzvit M, Dematteis S. 2016. Natural

and induced antibodies contribute to differential susceptibility to secondary cystic

echinococcosis of Balb/c and C57Bl/6 mice. Immunobiology 221:103–115.

39. Raedler A, Raedler E, Arndt R, Thiele HG. 1981. Terminal galactosyl residues of

cell-surface glycoconjugates exposed on both human and murine immature T- and

B-cells. Cell Tissue Res 218:219–226.

40. Díaz A, Allen JE. 2007. Mapping immune response profiles: The emerging

scenario from helminth immunology. Eur J Immunol 37:3319–3326.

41. Wang J, Kubes P. 2016. A Reservoir of Mature Cavity Macrophages that Can

Rapidly Invade Visceral Organs to Affect Tissue Repair. Cell 165:668–678.

42. Grewal PK. 2010. The Ashwell–Morell Receptor, p. 223–241. *In* Methods in

Enzymology. Elsevier.

43. Rensen PC, Sliedregt LA, Ferns M, Kieviet E, van Rossenberg SM, van Leeuwen

SH, van Berkel TJ, Biessen EA. 2001. Determination of the upper size limit for

uptake and processing of ligands by the asialoglycoprotein receptor on hepatocytes

in vitro and in vivo. J Biol Chem 276:37577–37584.

44. Barrios AA, Grezzi L, Miles S, Mariconti M, Mourglia-Ettlin G, Seoane PI, Díaz

A. 2019. Inefficient and abortive classical complement pathway activation by the

calcium inositol hexakisphosphate component of the Echinococcus granulosus

laminated layer. Immunobiology 224:710–719.

45. Irigoín F, Ferreira F, Fernández C, Sim RB, Díaz A. 2002. myo-Inositol

hexakisphosphate is a major component of an extracellular structure in the

parasitic cestode Echinococcus granulosus 8.

46. Irigoín F, Casaravilla C, Iborra F, Sim RB, Ferreira F, Díaz A. 2004. Unique

precipitation and exocytosis of a calcium salt of *myo* -inositol hexakisphosphate in

larval *Echinococcus granulosus*. J Cell Biochem 93:1272–1281.

47.   Casaravilla C, Brearley C, Soule S, Fontana C, Veiga N, Bessio MI, Ferreira F,

Kremer C, Diaz A. 2006. Characterization of myo-inositol hexakisphosphate

deposits from larval Echinococcus granulosus. FEBS Journal 273:3192–3203.

48.   Irigoín F, Laich A, Ferreira AM, Fernández C, Sim RB, Díaz A. 2008. Resistance

of the Echinococcus granulosus cyst wall to complement activation: analysis of the

role of InsP$_6$ deposits. Parasite Immunol 30:354–364.

49.   Seoane PI, Rückerl D, Casaravilla C, Barrios AA, Pittini Á, MacDonald AS, Allen

JE, Díaz A. 2016. Particles from the Echinococcus granulosus laminated layer

inhibit IL-4 and growth factor-driven Akt phosphorylation and proliferative

responses in macrophages. Sci Rep 6:39204.

50.   Casaravilla C, Pittini Á, Rückerl D, Allen JE, Díaz Á. 2020. Activation of the

NLRP3 Inflammasome by Particles from the Echinococcus granulosus Laminated

Layer. Infect Immun 88.

51.   Lüth S, Huber S, Schramm C, Buch T, Zander S, Stadelmann C, Brück W, Wraith

DC, Herkel J, Lohse AW. 2008. Ectopic expression of neural autoantigen in

mouse liver suppresses experimental autoimmune neuroinflammation by inducing

antigen-specific Tregs. J Clin Invest 118:3403–3410.

52.   You Q, Cheng L, Kedl RM, Ju C. 2008. Mechanism of T cell tolerance induction

by murine hepatic Kupffer cells. Hepatology 48:978–990.

53.   Hoberg EP, Jones A, Rausch RL, Eom KS, Gardner SL. 2000. A phylogenetic

hypothesis for species of the genus Taenia (Eucestoda : Taeniidae). J Parasitol

86:89–98.

54.   del Puerto L, Rovetta R, Navatta M, Fontana C, Lin G, Moyna G, Dematteis S,

823   Brehm K, Koziol U, Ferreira F, Díaz A. 2016. Negligible elongation of mucin

824   glycans with Gal β1-3 units distinguishes the laminated layer of Echinococcus

825   multilocularis from that of Echinococcus granulosus. International Journal for

826   Parasitology 46:311–321.

827   55.   Basika T, Muñoz N, Casaravilla C, Irigoín F, Batthyány C, Bonilla M, Salinas G,

828   Pacheco JP, Roth J, Durán R, Díaz A. 2012. Phagocyte-specific S100 proteins in

829   the local response to the *Echinococcus granulosus* larva. Parasitology 139:271–

830   283.

 Figure legends

Fig. 1. Soluble LL mucins are taken up selectively by KC, dependent on mucin glycans.

C57BL/6 mice were injected i.v. with 100 μg dry mass of solubilized, biotin-derivatized LL

mucins (sLL), either treated with periodate to oxidize their terminal monosaccharide residues

or mock-treated. Twenty minutes later, mice were sacrificed and sLL uptake was quantitated

in permeabilized liver non parenchymal cells using streptavidin-PE. Data are shown for the

cell types that are relevant in the context of the article, namely KC, LSEC and non-KC

CD11b$^+$ cells, in terms of percentages of positive cells **(a)** and geometric means of

fluorescence intensity **(b)**. Representative dot-plots for KC are shown in **(c)**. The signal

threshold for each cell type was established using cells from mice injected with vehicle only

and similarly stained; the fluorescence intensities data had the median values corresponding

to vehicle-injected mice subtracted for plotting. Results are pooled from 3 independent

experiments (represented in different colours); individual mice and their overall median

values are shown. The statistical analyses were carried out on the crude data, by the Mack-

Skillings test.

Fig. 2. PNA and the E492 antibody are suitable probes for LL mucins captured by

mouse KC. The binding of PNA, the E492 antibody and soluble recombinant Clec4F to

LL mucins coated onto ELISA plate wells **(a)** or present in LL particles in suspension **(b)**

was studied. The capacity of PNA and the E492 antibody to compete with the binding of

recombinant Clec4F to LL mucins was analysed in a competition ELISA format **(c)**;

results are expressed in relation to the response in the absence of competitor. In (a) and

(c), analytical duplicates are shown. WT C57BL/6 mice were injected i.v. either with PBS

only or with solubilized LL mucins (200 μg/mouse) and 22 h later KC and LSEC were

analysed by flow cytometry using either PNA or the E492 antibody as probes, after cell permeabilization (**d**). Staining controls shown in (b) and (d) correspond to samples stained in the same way as the positive ones except that the primary probe for LL material was omitted, and the analogous controls in (a) gave no signal.

Fig. 3. Generation of Clec4f gene-deficient mice. (a) Schematic structure of the mouse *Clec4f* gene. Coding exons appear as solid black bars and non-coding regions as white bars. Introns are represented as lines. **(b)** The target sequences for CRISPR/Cas9, 5'-TGGACGTGCCGCAGGGTCCT-3' and 5'-GCTTCTCCCCTGCAGGACTG-3', are located in the second exon and first intron/second exon junction of the gene respectively. An 8 bp deletion corresponding to nucleotides 5'- CAGGACCC-3' was identified by tail biopsy. The frameshift mutation produces a protein 480 aa shorter. **(c)** The liver non-parenchymal fraction was prepared from Clec4f$^{-/-}$ and WT mice and analyzed for Clec4F expression by flow cytometry without or with cellular permeabilization. KC were defined as shown in Figure S2. The staining controls contain all the fluorophore-coupled antibodies but lack the primary anti-Clec4F antibody.

Fig. 4. Fast uptake of soluble LL mucins by Kupffer cells depends largely on Clec4F. WT and Clec4f$^{-/-}$ mice were injected i.v. with 200 ug dry mass of sLL, either untreated or mock-treated as in Figure 1, or vehicle only (PBS). Twenty min later, mice were sacrificed and sLL uptake was quantitated in non-parenchymal liver cell populations by flow cytometry in permeabilized cells using PNA. Results are shown for KC, LSEC and non-KC CD11b$^+$ cells, in terms of percentages of positive cells **(a)** and mean fluorescence intensities **(b)**, and representative flow cytometry plots are shown for KC and LSEC **(c)**. An estimation of the fraction of sLL taken up by each of the cell populations under study, calculated as explained

in the Materials and Methods section, is shown in **(d)**; the statistics shown compare the

fractions taken up by KC. The positive cell threshold was established using cells from the

mice injected with PBS only. The data arise from two independent experiments, represented

in blue and red in (a-c) and indicated as 1, 2 in (d). In (a-c), individual mice and their median

values, and in (d) only medians, are shown. The statistical analyses were carried out by the

Mack-Skillings test.

Fig. 5. Clec4F is important for the capture of soluble LL mucins by Kupffer cells even

after allowing a long uptake time. WT and Clec4f$^{-/-}$ C57BL/6 mice were injected i.v. with

200 μg dry mass per mouse of sLL (mock-treated as in Fig. 1) or PBS only as a control.

Twenty-two h later, mice were sacrificed and sLL uptake was quantitated in liver non-

parenchymal cells by flow cytometry using PNA after cell permeabilization. Results are

shown for KC, LSEC and non-KC CD11b^{+} cells, in terms of percentages of positive cells **(a)**

and mean fluorescence intensities **(b)**, and representative flow plots for KC and LSEC are

shown in **(c)**. An estimation of the fraction of sLL taken up by each of the cell populations

under study, calculated as explained in the Materials and Methods section, is shown in **(d);**

the statistics shown compare the fractions taken up by KC. The positive cell threshold was

established using staining controls (in which only biotinylated PNA was omitted). The PNA

signal in non-permeabilized cells was negligible. The data arise from two independent

experiments, represented in blue and red in (a-c) and indicated as 1, 2 in (d). In (a-c),

individual mice and their median values, and in (d) only medians, are shown. The statistical

analyses were carried out by the Mack-Skillings test (a-c) and the modified Wilcoxon-Mann-

Whitney tests (d).

Fig. 6. The uptake by Kupffer cells of LL mucins injected in particulate presentation

depends largely on Clec4F. WT and Clec4f$^{-/-}$ C57BL/6 mice were injected i.p. with 450 μg

dry mass of pLL or PBS only as a control. Twenty-two h later, mice were sacrificed and pLL

uptake was quantitated in liver non-parenchymal cells by flow cytometry using PNA after

cell permeabilisation. Results are shown for KC, LSEC and non-KC CD11b^{+} cells, in terms

of percentages of positive cells **(a)** and mean fluorescence intensities **(b)**, and representative

dot-plots for the three cell types are shown in **(c)**. An estimation of the fraction of sLL taken

up by each of the cell populations under study, calculated as explained in the Materials and

Methods section, is shown in **(d)**; the statistics shown compare the fractions taken up by KC..

The positive cell threshold was established using staining controls. The data arise from two

independent experiments, represented in blue and red in (a-c) and indicated as 1, 2 in (d). In

(a-c), individual mice and their median values, and in (d) only medians, are shown. The

statistical analyses were carried out by the Mack-Skillings test. The analogous flow

cytometry results obtained using the E492 antibody for detection are shown in Fig. S6. In

addition, pLL was detected in liver sections, using the E492 antibody (**e**). Representative

images are shown, at two different magnifications. Black arrows indicate KC-like cells

staining strongly for LL material; empty arrowheads indicate hepatocytes staining weakly for

LL material.

Fig. 7. LL materials circulate systemically in infected mice and are captured selectively

by Kupffer cells via Clec4F. BALB/c mice (WT) were intraperitoneally infected with *E.*

granulosus protoscoleces or injected with the corresponding volume of buffer only. Seven

months later, the presence of LL-derived materials in various organs was analysed by

histology, using the E492 antibody as a probe **(a)**. Images from representative mice are

shown, out of 15 infected and 5 control mice; the bars represent 50 μm. Black arrows indicate

KC-like cells staining for LL material; empty arrowheads indicate occasional staining for LL

material in spleen and lymph node. The staining observed in adipocytes (upper left-hand

corner of the infected mouse mesenteric lymph node section) was also observed in adipocytes

present in kidney and lung sections, irrespective of infection, and must therefore represent a

cross-reaction of the antibody. No staining was observed in sections probed with the

secondary reagent only. The uptake of LL-derived materials by non-parenchymal liver cells

of infected BALB/c mice was also analysed by flow cytometry, using PNA as probe. Results

are shown for KC, LSEC and non-KC $CD11b^{+}$ cells, in terms of percentages of positive cells

(b) and mean fluorescence intensities **(c)**, and representative dot-plots for the three cell types

are shown in **(d)**. The positive cell threshold was established using staining controls. In

addition, WT and $Clec4f^{-/-}$ C57BL/6 mice were similarly infected and the uptake of LL-

derived materials analysed by flow cytometry and presented **(e – g)** as for the BALB/c mice.

Results in (a – d) are from a single experiment and results in (e - g) are pooled from two

independent experiments, shown in blue and red. In (b - g), data shown correspond to

individual mice and their median values are indicated. The statistical analyses were carried

out by the modified Wilcoxon-Mann-Whitney (b, c) and Kruskal-Wallis tests (e, g).

Supplementary Figures.

Fig. S1. Summary of preparation of LL materials. The preparation of pLL and sLL

(periodate- or mock-treated) is summarized in **(a)**. Periodate oxidation of terminal

monosaccharides and further monosaccharides carrying *cis*-diols generates reactive aldehyde

groups, which are reduced to alcohols using sodium borohydride. The mock treatment

consists of the borohydride step only. The effectiveness of the periodate treatment was

verified in terms of abrogation of PNA binding by to the mucins in a dot-blot format **(b)**.

Similar incorporation of the biotin tag to periodate- and mock-treated sLL was verified using

streptavidin-peroxidase in dot-blot format **(c)**.

Fig. S2. Gating strategy for liver cells. Gating strategies are shown for KC, LSEC, and non-

KC CD11b$^+$ cells. For non-KC CD11b$^+$ cells, which tend to be substantially smaller than KC,

it was necessary to include an additional doublet exclusion step at the end of the gating

strategy.

Fig. S3 (related to Fig. 3). Reactivity of PNA, E492 antibody and recombinant Clec4F

with liver non-parenchymal cells of control mice. Non-parenchymal cells from WT

C57BL/6 mice were stained for flow cytometry with PNA, E492 antibody or recombinant

mouse Clec4F, followed by appropriate secondary probes. The staining controls correspond

to samples stained in the same way as the positive ones except that the primary probe was

omitted.

Fig. S4 (related to Figs. 4 and 5). Representative flow plots for liver non-KC CD11b$^+$

cells. sLL was injected in WT and Clec4f$^{-/-}$ C57BL/6 mice, and liver non parenchymal cells

analysed after 20 min **(a)** or 22 h **(b)**, as described in Figs. 4 and 5 respectively. Representative

flow plots are shown for non-KC CD11b$^+$ cells, which were not found to take up sLL

significantly.

Fig. S5 (related to Fig. 5). Surface expression of Clec4F does not change upon injection

of sLL. WT C57BL/6 mice were injected with sLL or vehicle only (PBS) as described in Fig.

5, and 22 h later KC were analysed for surface expression of Clec4F by flow cytometry. Data

shown, in terms of geometric means of fluorescence intensity (MFI), arise from 3

independent experiments, shown in different colors. The data were normalized over the

median value of mice injected with PBS only in each experiment.

Fig. S6 (related to Fig. 6). Uptake of pLL by liver non-parenchymal cells measured by

flow cytometry using antibody E492. WT and Clec4f$^{-/-}$ C57BL/6 mice were injected with

pLL and material uptake was analysed as in Figure 6 except that the E492 antibody was used

as probe. Results are shown in terms of percentages of positive cells **(a)** and mean

fluorescence intensities **(b)**, and representative dot-plots are shown in **(c)**. An estimation of

the fraction of sLL taken up by each of the cell populations under study, calculated as

explained in the Materials and Methods section, is shown in **(d)**; the statistics shown compare

the fractions taken up by KC. The positive cell threshold was established using staining

controls. The data arise from two independent experiments, represented in blue and red in (a-

c) and indicated as 1, 2 in (d). In (a-c), individual mice and their median values, and in (d)

only medians, are shown. The statistical analyses were carried out by the Mack-Skillings test.

Fig. S7 (related to Fig. 6). Frequency of non-KC CD11b^{+} cells in mice injected with pLL.

WT and Clec4f$^{-/-}$ C57BL/6 mice were injected with pLL or vehicle only (PBS) as described

in Fig. 6, and 22 h later the frequency of non-KC CD11b^{+} cells within all live cells was

measured by flow cytometry. Data arise from 2 independent experiments, shown in different

colors.

1002

Fig. S8 (related to Fig. 6). Surface expression of Clec4F does not change upon injection

of pLL. WT C57BL/6 mice were injected with pLL or vehicle only (PBS) as described in

Fig. 6, and 22 h later KC were analysed for surface expression of Clec4F by flow cytometry.

Data shown, in terms of geometric means of fluorescence intensity (MFI), arise from 2

independent experiments, shown in different colors. The data were normalized over the

median value of mice injected with PBS only in each experiment.

1009

Fig. S9 (related to Fig. 7). Parasite burdens of mice infected with *E. granulosus*. WT

BALB/c mice **(a)** or WT and Clec4f$^{-/-}$ C57BL/6 mice **(b)** were infected with *E. granulosus*

as described in Fig. 7. Parasite burdens are expressed in terms of total volumes of

metacestodes, measured by water displacement. Results for BALB/c mice correspond to a

single experiment and results for C57BL/6 are pooled from 2 experiments, represented in

different colours. Individual mice and their overall median values are shown. Parasite burdens

were not found to be significantly different between WT and Clec4f$^{-/-}$ C57BL/6 mice by the

Kruskall-Wallis test.

1018

1019 **Fig. S10 (related to Fig. 7). Representative flow plots for LSEC in infections.** WT

1020 BALB/c mice **(a)** or WT and Clec4f$^{-/-}$ C57BL/6 mice **(b)** were infected with *E. granulosus*

1021 and liver non-parenchymal cells analysed by flow cytometry as described in Fig. 7.

1022 Representative flow plots are shown for LSEC, which were not found to take up LL materials.

1023

1024 **Fig. S11. Frequencies of non-KC CD11b$^+$ cells in *E. granulosus* experimental infections.**

1025 WT BALB/c mice **(a)** or WT and Clec4f$^{-/-}$ C57BL/6 mice **(b)** were infected with *E.*

1026 *granulosus* and liver non-parenchymal cells analysed by flow cytometry as described in Fig.

1027 7. Frequencies of non-KC CD11b$^+$ cells within total live cells are plotted. The statistical

1028 analysis was carried out by the modified Wilcoxon-Mann-Whitney method.

1029

Fig. S12. Experimental *E. granulosus* infection does not alter the expression of Clec4F

in KC. WT BALB/c mice **(a)** or WT C57BL/6 mice **(b)** were infected with *E. granulosus* as

described in Fig. 7. KC were analysed for surface expression of Clec4F by flow cytometry.

Data shown, in terms of geometric means of fluorescence intensity (MFI), arise from 1

experiment for each of the mouse strains shown. The data were normalized over the median

value of the control mice in each experiment.

1036

FIGURE 1

FIGURE 2

(a) ELISA format

PNA

E492 Ab

Clec4F

(b) Immuno/lectin fluorescence on particles

PNA   E492   Rec. Clec4F

Probe

Staining control

Bright field

(c) Competition ELISA format

Clec4F

Competitor:
● PNA
○ E492 Ab

(d) Flow cytometry gated on KC and LSEC

staining control   PNA   staining control   E492

KC
PBS
sLL

LSEC
PBS
sLL

FIGURE 3

(a)

Mus musculus clec4f (Chr. 6)
ENSMUST00000014686.3

(b)

WT	C T C	A T G	G C T	C C C	A G G	A C C	C T G	C G G	C A C	G T C	C A G	G T C	A T T C
	Leu	Met	Ala	Pro	Arg	Thr	Leu	Arg	His	Val	Gln	Val	Ile

clec4f mut	C T C	A T G	G C T	C C		T G C	G G C	A C G	T C C	A G G	T C A	T T C
	Leu	Met	Ala	Pro		Ala	Ala	Arg	Pro	Gly	His	Ser

WT MKEAELNRDMARYCTDNQCVSLQPQGLGPKSAALMAPRTLRHVQVILALMVVTVIFSLLALFVVASQPWRPEWNKEPPSLLLR
clec4f mut MKEAELNRDMARYCTDNQCVSLQPQGLGPKSAALMAPAARPGHSGINGGDCDLFSPCSFCGGFTALET*

(c)

staining control | Clec4f⁻/⁻ | WT

Non-permeabilized cells   0%   35%

Permeabilized cells   0%   70%

Cle4F

F4/80

FIGURE 4

Figure 5

Figure 6

Figure 7

(a)

	Peritoneal cavity cells	Liver	Spleen	Mesenteric lymph node
Control				
Infected				

	Small intestine	Kidney	Lung
Control			
Infected			

(b) KC — LSEC — Non-KC CD11b⁺ cells
% positive cells vs ctrl / inf.

(c) KC — LSEC — Non-KC CD11b⁺ cells
MFI vs ctrl / inf.

(d) staining control / control / infected
KC
Non-KC CD11b⁺ cells
PNA binding vs F4/80

(e) KC — LSEC — Non-KC CD11b⁺ cells
% positive cells vs ctrl / inf. WT / Clec4f⁻/⁻

(f) KC — LSEC — Non-KC CD11b⁺ cells
MFI vs ctrl / inf. WT / Clec4f⁻/⁻

(g) staining control / WT control infected / Clec4f⁻/⁻ control infected
KC
Non-KC CD11b⁺ cells
PNA binding vs F4/80