Multiple Postsynaptic Protein Levels in Adult Superior Colliculus Are Unaffected by Dark Rearing from Birth

Abbreviated title: Postsynaptic proteins stable in dark rearing

Parag S. Juvale (ORCID ID #0009-0005-2614-9828)[1§], David B. Mudd[2#], Nitheyaa Shree (ORCID ID #0000-0002-9681-6308)[2¶], and Sarah L. Pallas (ORCID ID #0000-0002-1760-7437)[1*]

[1] Department of Biology, University of Massachusetts Amherst, Amherst, MA 01003, USA

[2] Neuroscience Institute, Georgia State University, Atlanta, GA 30303, USA

§Current address: Aston Pharmacy School, Aston University, Birmingham, UK

Current address: Office of Technology Transfer, Emory University, Atlanta, GA

¶Current address: Graduate Program in Biochemistry, University of Bristol, UK

Conflict of interest: *The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.*

Author Contributions: Juvale, Mudd, and Pallas contributed to the conception and design of the study. Juvale, Mudd, and Shree collected data, organized the database, and performed the statistical analyses. Mudd wrote the first draft of the manuscript. Juvale wrote subsequent versions of the manuscript and Pallas edited the manuscript. All authors read the manuscript, contributed to manuscript revision, and approved the submitted version.

Funding: Support for this work was provided by a Sigma Xi GIAR grant and a UMass Amherst predissertation grant to P.S.J.; a GSU Brains & Behavior Fellowship, and a GSU Center for Neuromics student grant awarded to D.B.M.; and a GSU Brains & Behavior Seed grant, UMass startup funds, a National Science Foundation grant (IOS-1656838), and a DARPA grant (HR0011-18-2-0019, TA2) awarded to S.L.P.

*Correspondence:
Sarah L. Pallas
Department of Biology
University of Massachusetts-Amherst
Amherst, MA 01003 USA
spallas@umass.edu

Keywords: Visual deprivation, superior colliculus, GABA, adult plasticity, visual refinement, GABA$_A$ receptor, gephyrin, chloride co-transporter

Number of pages: 25
Number of words: 5899
Number of figures: 7
Number of tables: 0
Number of words in Abstract: 255

Acknowledgments: We thank Profs. Angela Mabb, Larry Schwartz, Thomas Maresca, and Lillian Fritz-Laylin for instrument access and guidance with Western Blotting, Profs. Gerald Downes, Margaret Stratton, and Stephen Moss for providing technical guidance and support on P.S.J.'s thesis committee, Pallas lab members for technical support and manuscript review, and the animal care staff at GSU and UMass. We thank Dr. Andrew Stephens for providing the Histone H3 antibody.

Abstract

Visual deprivation by dark rearing in kittens and monkeys delays visual pathway development and prolongs the critical period. In contrast, receptive fields (RFs) in superior colliculus (SC) of Syrian hamsters *(Mesocricetus auratus)* refine normally with spontaneous activity alone, requiring only brief juvenile visual experience to maintain refined RFs in adulthood (Carrasco et al., 2005). Extending dark rearing past puberty leads to lower GAD and GABA levels due to reduced BDNF-TrkB signaling, resulting in RF re-enlargement (Carrasco et al., 2011; Mudd et al., 2019). Previous studies in kittens and monkeys have reported that dark rearing is associated with changes in both GABA ligand and $GABA_A$ receptor levels. Given the reduced GABA levels in SC of dark reared adult hamsters, we asked if dark rearing also causes changes in $GABA_A$ receptor levels. We examined expression of $GABA_A$ receptor subunits, their anchoring protein gephyrin, and the cation-chloride co-transporters KCC2 and NKCC1 in dark reared hamsters. Surprisingly, we found that dark rearing from birth until puberty had no effect on the levels of any of these postsynaptic elements, revealing a new form of maladaptive, presynaptic only inhibitory plasticity in which, rather than extending the critical period as seen in kittens and monkeys, hamster receptive fields refine normally and then lose refinement in adulthood. These results suggest that attempts to increase plasticity in adulthood for rehabilitation or recovery from injury should consider the possibility of unintended negative consequences. In addition, our results demonstrate the interesting finding that changes in neurotransmitter levels are not necessarily coordinated with changes in postsynaptic components.

Introduction

During brain development, synaptic connections are elaborated and then refined to a mature state under the influence of neural activity. Activity due to sensory experience during an early "critical period" is important for shaping some aspects of neural circuit development. In the visual pathway, although spontaneous retinal activity is important for initial axon pruning (Kutsarova et al., 2017), it has long been thought that early visual experience is essential to attain refined connectivity patterns during development (Wiesel and Hubel, 1965; Maffei and Galli-Resta, 1990; Meister et al., 1991; Wong et al., 1993; Katz and Shatz, 1996; Firth et al., 2005). Once critical periods have closed, plasticity is often limited or even prevented, thus protecting refined circuits from destabilization (Hubel and Wiesel, 1970; Takesian and Hensch, 2013; Hübener and Bonhoeffer, 2014; Pallas, 2017; Hensch and Quinlan, 2018; Reh et al., 2020; Ribic, 2020; Mitchell and Maurer, 2022).

In some mammals, dark rearing is reported to delay or prevent refinement, prolonging critical period plasticity (Cynader and Mitchell, 1980; Mower et al., 1985; Mower, 1991; Lee and Nedivi, 2002; Nakadate et al., 2012). In apparent contradiction to this common view, we have reported that spontaneous activity is sufficient for refinement of receptive fields in both visual cortex (V1) and superior colliculus (SC) of dark reared (DR) Syrian hamsters *(Mesocricetus auratus)*, and early light exposure for 3-7 days is necessary only to maintain the refinement into adulthood (Carrasco et al., 2005; Carrasco and Pallas, 2006; Balmer and Pallas, 2015; Mudd et al., 2019). Syrian hamster pups spend the first 3-4 weeks after birth underground in the wild (Adler, 1948; Nowosielski-Slepowron and Park, 1987), so it would not be beneficial to have the development of their visual function depend on light exposure. If early visual experience continues to be unavailable, GABAergic lateral inhibition in SC and V1 declines and RFs expand, but not until approximately two months of age (~puberty) (Carrasco et al., 2005; Balmer and Pallas, 2015; Mudd et al., 2019). Pharmacological activation of TrkB receptors can mimic the effects of early light exposure in DR hamsters, resulting in long-term maintenance of refined receptive fields and visual acuity (Mudd et al., 2019), perhaps through promoting GABA synthesis (Zhang et al., 2018).

Thus, hamsters, contrary to what has been found in cats and monkeys, need visual experience to maintain

refined receptive fields in adulthood, but not to refine them during development **(Figure 1)**.

$GABA_A$ receptors are pentameric, ionotrophic receptors consisting of five subunits grouped around a

central chloride ion pore. The functional characteristics of the receptor largely depend upon the subunit

composition (Sigel et al., 1990) and organization (Minier and Sigel, 2004). Of the many subunit

arrangements, alpha1 and alpha2 subunits have been linked to synaptic localization of $GABA_A$ receptors.

However, these two subunit types have different kinetics and are expressed at different points in

development. At birth, receptors containing the alpha2 subunit are widely expressed throughout the brain,

whereas alpha1 expression is initially low in major areas of the brain like the neocortex, the hippocampus,

and the cerebellum (Laurie et al., 1992; Fritschy et al., 1994; Dunning et al., 1999; Chen et al., 2001).

During the first several postnatal weeks, alpha1 expression increases, coinciding with a reduction in

alpha2 expression (Fritschy et al., 1994). This alpha2 to alpha1 switch in subunit expression underlies a

developmental decrease in inhibitory postsynaptic current (ipsc) decay time and an increase in ipsc

amplitude (Fritschy et al., 1994; Okada et al., 2000; Yu et al., 2006).

Our lab demonstrated previously that the expansion of RFs in SC of adult, DR hamsters is associated

with a loss of GABA immunoreactivity (Carrasco et al., 2011). Iontophoretic application of $GABA_A$

agonists *in vivo* restored RFs to a normal adult size. In addition to a loss of GABA-immunoreactivity,

$GABA_A$ agonists and antagonists were less effective in SC and V1 neurons of DR hamsters than in

normally reared (NR) hamsters (Carrasco et al., 2011).

The incomplete development of RF properties in V1 of visually deprived cats has been associated

with a failure in developmental maturation of NMDA and $GABA_A$ receptors (Carmignoto and Vicini,

1992; Chen et al., 2000; Chen et al., 2001; Erisir and Harris, 2003). Although a failure to maintain refined

RFs is a different phenomenon than a failure to refine them during a critical period, the mechanism(s)

could be similar or convergent. We thus tested the hypothesis that maintenance of refined RFs in

adulthood depends on the stability of mature receptors and other postsynaptic signaling components. This

hypothesis predicts that the detrimental, post-critical period receptive field plasticity observed in DR adult

hamsters results from a deprivation-induced failure to maintain these postsynaptic proteins in their mature

state. Contrary to this hypothesis, we find, using Western blot assays of synaptosomes, that the quantity,

subunit composition, and localization of $GABA_A$ receptors in SC of adult dark reared hamsters with re-

expanded RFs resemble those seen in normally reared subjects. Furthermore, levels of the synaptic

scaffolding proteins gephyrin and PSD-95 are normal, as are the adult expression levels of cation-chloride

co-transporters (KCC2/NKCC) in DR subjects. These findings suggest that, although a change in

effectiveness of $GABA_A$ receptors was reported previously using pharmacological agents (Carrasco et al.,

2011; Balmer and Pallas, 2015), the loss of RF refinement in adulthood is mediated primarily by

reductions in GABA expression in the presynaptic terminals rather than by significant postsynaptic

alterations. This result is at odds with the common view that presynaptic changes in the ligand must

occur together with corresponding postsynaptic changes in receptor levels (Fisher-Lavie and Ziv, 2013;

Sudhof, 2018 ; Sanderson et al., 2020). Taken together, our results rule out several hypotheses about the

mechanistic basis of refined RF maintenance throughout adulthood and provide insights into regulation of

critical period plasticity that could help to understand the regulation of GABAergic signaling at the

synaptic level. Similar research in diurnal animals that have photopic vision as do humans could help to

provide insight on treatment and therapeutic modalities in adults facing issues with plasticity in

adulthood.

Materials and Methods

Subjects

A total of 42 adult Syrian hamsters (Cricetidae: *Mesocricetus auratus*) (aged P90-P100) of both sexes

were bred within our animal facility and used as subjects in this study. Syrian hamsters are an altricial

rodent species that is ideal for studying the developing visual system due to their robust and well-

characterized visual responses, short gestation time, and large litters (Chalupa, 1981; Huck et al., 1988;

Pratt and Lisk, 1989; Razak et al., 2003; Carrasco et al., 2005). Sexual maturity in this species occurs

between postnatal days (P)56 and P60 (Diamond and Yanagimachi, 1970; Fitzgerald and Zucker, 1976).

Breeding females were singly housed. Male breeders were introduced and supervised until intromission

was observed, after which they were removed. Weanlings and adult males were housed in single sex

social groups of up to 5 adults per cage and adult females were housed with female siblings or

individually in standard rodent cages. Running wheels were not provided because they have been shown

to alter the timing of visual cortical plasticity (Baroncelli et al., 2010; Tognini et al., 2012; Kalogeraki et

al., 2014) but a variety of other enrichment items were available. All subjects were provided with *ad*

libitum access to rodent chow and water.

Treatment groups

Animals used in this study were bred in-house to control sensory experiences throughout development.

Dams of DR subjects were transferred into total darkness 1-3 days before parturition. An antechamber

and light-impenetrable black curtain separated the dark housing room from the hallway, ensuring that any

accidental openings of the hallway doors did not expose the animals to light. Dark reared animals were

housed inside light-tight stackable cages with a standard HVAC filtration system consistent with the other

animal rooms in the facility. During general animal husbandry purposes, the hamsters were exposed to

dim red light at a wavelength not visible to Syrian hamsters (Huhman and Albers, 1994). NR hamsters

were maintained in a standard 12/12 light cycle room from before birth into adulthood.

Western blotting

Animals were euthanized with Euthasol at >150 mg/kg IP prior to tissue collection. Brains were

immediately extracted and frozen in 2-methylbutane on dry ice, then stored at -80°C or immediately

dissected for preparation of lysates. In order to analyze differences in protein levels between NR and DR

hamsters, we used immunoblotting (Western blots). Western blots can detect protein levels at a 1-3 ng

resolution (Coorssen et al., 2002; Ghosh et al., 2014), allowing high resolution quantification of proteins.

Note that normal levels of synaptic GABA concentration have been estimated to be between 1.5-3 mM

(Coorssen et al., 2002; Tretter et al., 2008; Ghosh et al., 2014). GAPDH or β-actin were used as loading

controls to normalize for any differences in the amount of lysate pipetted into each gel lane.

Protein extraction was done as described by Shi et al. (1997) with a few modifications. Briefly,

individual left and right SC brain areas were excised and homogenized in a lysis buffer (10 mM

phosphate buffer, pH 7.0, 5 mM EGTA, 5 mM EDTA, 1 mM DTT) containing Halt protease inhibitor

(ThermoFisher Scientific). The lysate was centrifuged at 13,000 rpm at 4°C for 10 min, and the

supernatant was saved for the analysis of cytosolic proteins (cytosolic fraction). The resulting pellet was

resuspended in 2 mM HEPES buffer, pH 7.2. It was then centrifuged at 70,000 rpm at 4°C for 30 min.

The pellet thus obtained was resuspended in 0.5 mM HEPES, pH 7.3, containing 0.32 M sucrose and

centrifuged at 2,000 rpm for 8 min. Synaptosomes are present in the supernatant with this method. The

success of the synaptosome isolation protocol was confirmed by assessing the presence of Histone H3,

which should only be present in the cytosolic fractions and not in the synaptosome fractions **(Figure 2)**.

Synaptic proteins were then quantified using the Pierce BCA Protein Assay Kit (ThermoFisher Scientific)

mixed with 2X sample buffer and heated for 15 min at 60 °C. Twenty µg of the synaptosome proteins

were loaded per well in pre-cast Bio-Rad gels and electrophoresis was carried out at 110 V for 90 min in a

Bio-Rad electrophoresis tank. Proteins were then transferred onto nitrocellulose membranes at 70 V for

75 min, blocked in BSA for 1 h, and probed with primary antibodies overnight. Primary antibodies used

in this study included: rabbit anti-GABA$_A$Rα1 (1:1000, Cat#: AGA-001, Alomone Labs); rabbit anti-

GABA$_A$Rα2 (1:1000, Cat#: ab72445, Abcam); rabbit anti- GABA$_A$Rα5 (1:1000, Cat#: ab10098, Abcam);

rabbit anti-Gephyrin (1:1000, Cat#: ab32206, Abcam); mouse anti-PSD-95 (1:1000, Cat#: ab2723,

Abcam); mouse anti-KCC2 (1:1000, Cat#: 75-013, NeuroMab); rabbit anti-NKCC1 (1:1000, Cat#:

ab59791, Abcam); mouse anti-β-actin (1:1000, Cat#: A2228, Sigma-Aldrich) and mouse anti-GAPDH

(1:1000, Cat#: 600-GAPDH, PhosphoSolutions). Protein bands were labeled using either appropriate

fluorescent secondaries or appropriate HRP-conjugated secondary antibodies, then imaged on an Odyssey

CLx fluorescent imaging system (Li-Cor) or developed with enhanced chemiluminescent (ECL) substrate

in a Bio-Rad ChemiDoc Imager. All of the proteins studied here were analyzed and quantified as a ratio

of the optical density of the protein of interest compared to the density of the loading control (either

GAPDH or β-actin). Note that GAPDH is found in the pre-and post-synaptic sites (Frederikse et al., 2016)

along with β-actin, and thus makes an effective control protein to measure relative densities.

Statistical Analysis

A Student's *t*-test was used to compare parametric data with equal variance between treatment groups and

a normally distributed control data set. In the case of non-parametric data (data that were not normally

distributed and/or exhibited unequal variance), a Mann-Whitney Rank Sum (U) test was employed.

Descriptive statistics for these analyses are provided as mean ± standard error of the mean (SEM) in the

text. Whiskers represent the 5th (lower) and 95th (upper) percentage of the data.

Results

The failure to maintain RF refinement in adult DR animals involves deficits in overall GABA expression

and $GABA_A$ receptor function (Carrasco et al., 2011), leading to a loss of lateral inhibition and thus

expansion of RFs after P60 (Carrasco et al., 2005; Balmer and Pallas, 2015; Mudd et al., 2019). Using

adult hamsters (postnatal day (P)90-P100), we explored several possible ways that dark rearing during a

critical period for RF refinement could affect levels of $GABA_A$ receptors and other postsynaptic proteins

associated with inhibitory synaptic function in adult SC. We used Western blotting on synaptosomes in

normal and dark reared animals to study postsynaptic proteins that might regulate synaptic plasticity.

Dark rearing does not affect the subunit composition of $GABA_A$ receptors in adult SC

Deprivation-induced decreases in both $GABA_A$ and NMDA receptor levels in cat visual cortex have been

reported previously and were proposed to be involved in functional deficits (Carmignoto and Vicini,

1992; Chen et al., 2000; Chen et al., 2001). In DR hamsters, GAD and GABA immunoreactivity declines

(Carrasco et al., 2011; Otfinoski & Pallas, in prep.) and $GABA_A$ receptors are less efficient when tested

pharmacologically (Carrasco et al., 2011), thus we expected to see changes in levels of postsynaptic

$GABA_A$ proteins. However, in our previous study using Western blots on synaptosomes, no significant

changes in the level of the $GABA_A$ receptor alpha1 subunit were observed (Mudd et al., 2019). These

results raised the question of whether subunit composition of $GABA_A Rs$ might be altered by dark rearing

in a way that reduced their effectiveness without affecting alpha1 levels.

We reasoned that the developmental alpha2 to alpha1 switch, if reversed in adulthood, could underlie

the reduction in $GABA_A$ receptor function that was previously observed in studies of RF enlargement in

adult SC (Carrasco et al., 2011). We explored this possibility by examining the expression of each subunit

in synaptosomes of SC in normally reared and visually deprived adults. Hamsters in the experimental

group were dark reared from before birth. We used Western blotting for a high resolution, quantitative

assay of synaptic membrane-bound alpha1 and alpha2 $GABA_A$ receptor expression in the synaptosome

fractions obtained from adult SC. We found that there were no significant differences in either the overall

levels of alpha2 protein, observed as a ratio of alpha2 to GAPDH (NR: 1.21 ± 0.052, n=8; DR: $1.33 \pm$

0.185, n=8; U=26, n=8, p=0.574; Mann-Whitney Rank Sum Test) **(Figure 3A)**, or in the ratio of

alpha1/alpha2 expression, observed as a ratio of the normalized alpha 1 density (alpha 1/GAPDH ratio) to

normalized alpha 2 density (alpha 2/GAPDH ratio) in the SC of adult DR (1.651 ± 0.277, n=5) compared

to adult NR hamsters (1.19 ± 0.084, n=4) (T(7)= -1.436, n=4, p=0.194 Student's t-test) **(Figure 3B)**. This

was a surprising result considering our previous finding that dark rearing reduces the response to

pharmacological application of GABA agonists and antagonists (Carrasco et al., 2011). These findings

argue against a reversal of the normal developmental transition from alpha2 to alpha1-dominant

expression as a cause of the deprivation-induced RF enlargement in adult SC, and they support the

interpretation that visual experience is not needed to maintain mature $GABA_A$ receptor alpha1/alpha2

subunit composition.

$GABA_A$ receptors can also be expressed extrasynaptically, where they can be activated by GABA

derived from synaptic spillover or non-neuronal sources. This low concentration GABA source generates

"tonic" inhibition (Farrant and Nusser, 2005). Alpha5 subunit-containing receptors are primarily

expressed extrasynaptically and have been implicated in regulating the induction of synaptic plasticity for

LTP in hippocampus (Saab et al., 2010; Zurek et al., 2012; Zurek et al., 2014; Jacob, 2019). However,

alpha5 GABA$_A$Rs can relocate to the synapse and colocalize with gephyrin (Brady and Jacob, 2015). To

investigate the possible role of synaptic alpha5 levels in adult RF maintenance we quantified and

compared the alpha5/GAPDH ratios between NR (0.544 ± 0.0520, n=8) and DR (0.471 ± 0.0935, n=8)

adult hamsters **(Figure 4A)** using Western blotting. We found no significant differences in alpha5 protein

levels between groups (U (20) = 0.308, p = 0.878, Mann-Whitney Rank Sum test). We compared the ratio

of alpha1/alpha5 between adult NR (1.027 ± 0.0815, n=10) and DR (0.995 ± 0.0926, n =9) hamsters and

found no differences between these groups (U (18) =35, p=0.438, Mann-Whitney Rank Sum test) **(Figure**

4B). Because we were only studying proteins from synaptosome preparations (i.e., GABA$_A$ receptor

subunits localized in the synapses), these results suggest that the localization of alpha1 and alpha5

subunit-containing GABA$_A$ receptors in SC is not being altered by early visual experience.

Dark rearing does not affect the normal location of GABA$_A$ receptors in adult SC

The regulation of GABA$_A$ receptors at the synapse is pivotal for maintaining correct levels of inhibitory

synaptic transmission (Jacob et al., 2008). Impaired trafficking of GABA$_A$ receptors into and out of

synaptic membranes could affect their synaptic localization in SC and thus their overall response to

presynaptically released GABA. GABA$_A$ receptor trafficking is partially regulated by endocytosis – the

controlled removal of receptors from the postsynaptic membrane into the cytoplasm (see Lévi and Triller,

2006, for review). Endocytosed receptors are subsequently reinserted into the postsynaptic membrane or

undergo lysosomal degradation (Kittler et al., 2000). We reasoned that if internalization was dysregulated,

either by decreased receptor reinsertion or increased receptor degradation, it could negatively impact the

efficacy of GABA$_A$ receptors at the synapse. We examined this possibility by comparing the ratio of

postsynaptic membrane-bound to cytosolic alpha1 subunit- containing receptors between our treatment

groups. No differences were observed in the postsynaptic membrane/cytosol ratio of alpha1 expressing

receptors between DR (0.768 ± 0.044, n=7) and NR (0.778 ± 0.1, n=6) adult hamsters (U(11)= 20.00,

p=0.945, Mann-Whitney Rank Sum test) **(Figure 5A)**. These results indicate that the overall trafficking of

alpha1 subunit-expressing synaptic $GABA_A$ receptors is not affected by dark rearing.

We also examined the possibility that extrasynaptic alpha5 receptor internalization may be

dysregulated and thus responsible for changes in tonic $GABA_A$ inhibition (Davenport et al., 2021). There

were no significant differences in cytosolic membrane localization of alpha5 subunits between adult DR

$(1.320 \pm 0.198, n=8)$ and NR groups, however $(1.753 \pm 0.449, n=6)$ (U(12)=19.00, p=0.573, Mann-

Whitney Rank Sum test) **(Figure 5B)**. These results indicate that the internalization of extrasynaptic

alpha5 subunit-expressing $GABA_A$ receptors is not responsible for the decreased efficacy of $GABA_A$

receptors observed in RFs that fail to maintain refinement in adulthood after dark rearing.

Inhibitory and excitatory scaffolding proteins in SC are not affected by dark rearing

One factor influencing the accumulation and retention of $GABA_A$ receptors at postsynaptic sites is the

membrane scaffolding protein gephyrin (Kneussel et al., 1999; Sun et al., 2004; Jacob et al., 2005; Tretter

et al., 2008). Decreased expression of gephyrin results in less clustering (Essrich et al., 1998) and more

mobility of $GABA_A$ receptors at the synapse (Jacob et al., 2005). We surmised that decreased gephyrin

expression could be responsible for the weaker $GABA_A$ receptor signaling observed in neurons with RFs

that failed to maintain refinement in adulthood. We quantified and compared postsynaptic membrane-

bound gephyrin expression between DR and NR adults using Western blotting. Gephyrin levels in DR

adults $(0.786 \pm 0.124, n=17)$ were similar to those in NR adults $(0.736 \pm 0.158, n=16)$ (U(111)=-0.247,

p=0.377, Mann-Whitney Rank Sum test) **(Figure 6A)**. This indicates that maintenance of adult gephyrin

expression levels is not affected by dark rearing and suggests that if $GABA_A$ receptor accumulation and

trafficking is being affected, then it is occurring independently of gephyrin levels.

PSD-95 is the primary glutamate (AMPA and NMDA) receptor scaffolding protein in CNS neurons

(Chen et al., 2015), and it functions like gephyrin does for $GABA_A$ receptors. Although it does not

directly impact $GABA_A$ receptor function, PSD-95 has an influence on visual circuit plasticity. For

example, mice lacking PSD-95 have lifelong ocular dominance plasticity in primary visual cortex that

results from an increase in the overall proportion of silent synapses, despite having normal inhibitory tone

(Funahashi et al., 2013; Huang et al., 2015). Thus, we examined whether the dark rearing-induced re-

enlargement of RFs could be mediated by a reduction in adult PSD-95 expression. We found that PSD-95

protein levels were not significantly different in DR (0.613± 0.96, n=10) compared to NR adult hamsters

(0.486 ± 0.868, n=9) (U(31)=-0.978, p=0.270, Mann-Whitney Rank Sum test) **(Figure 6B)**. These results

suggest that differences in PSD-95 levels do not underlie the re-enlargement of RFs in SC following dark

rearing from birth.

Cation-chloride co-transporters undergo their normal developmental switch in adult dark reared

subjects

Levels of inhibitory GABAergic signaling in neurons are dependent on the intracellular chloride (Cl^-)

concentration. The K^+ Cl^- co-transporter (KCC2) is responsible for regulating intracellular Cl^- in mature

adult neurons with an outward K^+ current (Rivera et al., 1999) and also regulates the formation, function,

and plasticity of glutamatergic synapses (Li et al., 2007; Gauvain et al., 2011; Chevy et al., 2015). At the

beginning of postnatal life, $GABA_A$ receptor effects are excitatory because the Na^+-K^+-$2Cl^-$ co-

transporter 1 (NKCC1) that mediates Cl^- uptake is dominant (Cherubini et al., 1991; Lee et al., 2005;

Cancedda et al., 2007). By the end of the second postnatal week in rats and mice NKCC1 is replaced by

KCC2 as the dominant cation-chloride co-transporter in the brain, shifting the resting membrane potential

and thus causing $GABA_A$ receptors to produce inhibitory PSPs (Rivera et al., 1999; Pfeffer et al., 2009;

Moore et al., 2019). In V1, the developmental switch from NKCC1 dominance to KCC2 dominance

occurs at the same time as a period of BDNF/TrkB mediated synaptic imbalance (Zhang et al., 2018). We

surmised that a shift in the ratio of KCC2:NKCC1 could underlie the reinstatement of RF size plasticity in

dark reared adults, leading to re-enlargement of RFs in SC. Examination of the expression of KCC2 and

NKCC1 in adult SC neurons revealed no significant differences between our treatment groups, however.

KCC2 levels were not significantly different between NR (0.979 ± 0.115, n=8) and DR groups of adult

hamsters (0.963 ± 0.154, n=8) (T(14)=0.082, p=0.936, t-test) **(Figure 7A)**. The same was true of NKCC1

levels (NR 1.050 ± 0.0419, n=8; DR 1.081 ± 0.0814, n=8) ($T(14)=-0.339$, p=0.740, t-test) **(Figure 7B)**,

and of the ratio of the two cation-chloride co-transporters within groups ($T(8)=1.096$, p=0.305, t-test)

(Figure 7C).

Discussion

The goal of this study was to examine potential postsynaptic mechanisms through which light exposure

during an early critical period ensures the long-term stability of visual receptive fields in the hamster

superior colliculus. Our previous results established a correlation between the maintenance of RF

refinement and levels of GABA immunoreactivity in SC (Carrasco et al., 2011; Mudd et al., 2019) and

V1 (Balmer and Pallas, 2015), but potential postsynaptic changes in $GABA_A$ receptor and related protein

levels had not been examined. We have reported here that at the high-resolution level of Western blot

protein quantification, visual deprivation-induced failure to maintain refined RFs in SC does not appear to

involve changes in $GABA_A$R subunit composition, inhibitory or excitatory scaffolding protein expression,

or cation-Chloride co-transporter ratios. These results exclude several possible mechanisms that could

explain the reduced activation of $GABA_A$Rs with GABA agonists reported in DR adult SC (Carrasco et

al., 2011), and support activity-dependent regulation of GABA expression as the primary mechanism

underlying TrkB-mediated maintenance of RF refinement (Mudd et al., 2019). The finding that a change

in GABA levels could affect RF refinement in adulthood has important implications for the treatment of

memory impairments or brain injury.

This study supports our previous research that provided substantial evidence of a novel, maladaptive

adult plasticity in which visually deprived hamsters refine SC RFs normally but fail to maintain them in

adulthood. Our research differs from these previous studies in suggesting that dark reared hamsters lose

visual refinement in adulthood and not, as in the case of monkeys, ferrets, and cats, during development

(Mower and Christen, 1985; Mower et al., 1986; Mower, 1991; Carmignoto and Vicini, 1992; Fagiolini et

al., 1994; Chen et al., 2000; Chen et al., 2001; Lee and Nedivi, 2002; Erisir and Harris, 2003). Some of

these previous studies looked only at early and/or adult ages in the animals, thereby missing the RF

refinement that happens in between the two ages. Because diminished GABA release, contrary to what

we expected, did not elicit measurable changes in the levels of postsynaptic GABA$_A$ receptors, scaffold

proteins, or chloride co-transporters, this study provides a valuable demonstration that changes in

neurotransmitter availability do not necessarily result in coordinated changes in postsynaptic receptors.

Maturation of GABAergic signaling in visual cortex, particularly of the fast-spiking, parvalbumin-

containing basket cells, is thought to open and then close the critical period for plasticity (Fagiolini et al.,

2004; Sale et al., 2010; Toyoizumi et al., 2013; Capogna et al., 2021). Combined pre- and postsynaptic

alteration of synaptic strength has been seen in other sensory deprivation paradigms, including in dark

reared and monocularly deprived visual cortex, although with an earlier time course (Carmignoto and

Vicini, 1992; Chen et al., 2000; Chen et al., 2001). However, the retinorecipient layers of the superior

colliculus have no basket cells and contain very few GABAergic parvalbumin neurons, and the plasticity

described here occurs after the critical period has closed, suggesting that SC may accomplish plasticity

through a different mechanism than visual cortex. On the other hand, previous studies found that, as in

visual cortex (Hanover et al., 1999; Huang et al., 1999; Viegi et al., 2002), deprivation-induced receptive

field plasticity in adult SC is mediated by the BDNF receptor TrkB (Mudd et al., 2019). Furthermore,

reduced GABA and GABA$_A$ receptor efficacy in response to iontophoretically-applied GABA$_A$R agonists

and antagonists is observed in both SC and V1 of dark reared hamsters (Carrasco et al., 2011; Balmer and

Pallas, 2015), arguing for mechanistic elements in common.

Early visual experience is not necessary for maturation or maintenance of GABA$_A$ receptor subunit

composition at the synapse

GABA$_A$ receptors contain fast acting chloride (Cl$^-$) channels (Pfeiffer et al., 1982; Sigel and

Steinmann, 2012). The subunit composition of GABA$_A$ receptors changes during development from an

alpha 2 to alpha 1 dominant condition (Laurie et al., 1992; Fritschy et al., 1994; Chen et al., 2001) and

also changes in some disease states (Levitt, 2005; Tyson and Anderson, 2014; Deidda et al., 2015;

Kimoto et al., 2015; Schmidt and Mirnics, 2015; Tang et al., 2021) in a way that affects receptor

functional properties (Farrant and Nusser, 2005) and localization (Jacob et al., 2005). We studied synaptic levels of the GABA$_A$ receptor alpha1 and alpha2 subunits to quantify their expression levels under normal and DR conditions. The normalized expression levels of GABA$_A$R alpha1 relative to GABA$_A$R alpha2 levels were not altered in DR hamsters when compared to those of NR hamsters, arguing that the altered inhibitory synaptic efficacy that we previously observed was not caused by an immature GABA$_A$R subunit composition at the synapse.

An increase or decrease in the level of any receptor subunit is best understood in context, because different conclusions would be drawn if subunits changed independently or in concert. Thus, we also analyzed the alpha1/alpha2 ratios in individual animals. We did not find any change in alpha1/alpha2 ratios in NR compared to DR adult hamsters. These results suggest that early visual experience is not necessary for maturation or maintenance of mature synaptic GABA$_A$R subunit composition in adulthood. Thus, the failure to maintain refined RFs in adult DR hamsters cannot be explained by a return to a juvenile type of GABA$_A$R subunit composition.

Level and localization of the extrasynaptic GABA$_A$R subunit alpha 5 does not change with dark rearing

GABA$_A$R subunit alpha 5 is predominantly an extrasynaptic membrane receptor subunit that regulates tonic inhibition. It is important in neuronal circuit development, learning, and memory (Brady and Jacob, 2015), has been implicated in regulating the induction of synaptic plasticity in hippocampus (Saab et al., 2010; Zurek et al., 2012; Zurek et al., 2014), and can relocate to the synaptic region in learning and memory deficits (Brady and Jacob, 2015). Because the excitation/inhibition (E/I) balance could be affected if alpha5 subunit levels changed or if they moved into the synapse, we compared its expression between NR and DR cases. We did not see any significant changes in the overall levels of GABA$_A$R alpha5 subunits, or in the alpha1/alpha5 ratio, suggesting that dark rearing-induced RF enlargement is not caused by changes in the GABA$_A$R subunit alpha 5 levels or localization in the synapse.

Dark rearing does not affect the rate of internalization of GABA_AR subunits at the synapse

Because clathrin-dependent endocytosis is likely important for regulating inhibitory signaling and

synaptic plasticity (Kittler et al., 2000), we explored the internalization of $GABA_A$ receptor alpha1 and

alpha5 subunits by comparing their synaptic vs. extrasynaptic location in normally reared and dark reared

subjects. We did not observe a significant change in location of either subunit type as assayed by the ratio

of synaptosome-bound to cytosolic fractions. This implies that a lack of visual experience does not affect

the trafficking of the $GABA_AR$ subunits between the synaptic membrane and the cytosol or the

phosphorylation events that maintain the balance between internalization and postsynaptic membrane

insertion of the receptor subunits.

Early visual experience is not necessary to maintain scaffolding protein levels at the synapse

Another finding of this work is that the expression levels of the postsynaptic scaffold proteins PSD-

95 and gephyrin were not altered in adulthood following lifelong lack of light exposure, suggesting that

any changes in inhibitory function are probably not caused by a significant change in scaffolding protein

expression. At any rate, the clustering of $GABA_ARs$ at inhibitory synapses in SC may happen in a

gephyrin-independent manner (Kneussel et al., 2001), or total gephyrin expression may not be as

important as the formation of gephyrin nanodomains within inhibitory synapses (Pennacchietti et al.,

2017). Future studies with additional techniques would be required to determine if changes in receptor

clustering may be occurring and what role gephyrin or PSD-95 may have in mediating any such effects.

Early light exposure is not necessary for maturation of the cation-chloride co-transporters

We investigated the status of the chloride transporters KCC2 and NKCC1 due to their role in maintaining

chloride balance inside of the neurons and thus in setting the reversal potential. The cation-chloride co-

transporters could have reverted to their early developmental state, leading to a lower threshold for

excitation in dark reared animals, possibly explaining the RF expansion we observed. However, we did

not observe any changes in the cation-chloride co-transporters in dark-reared compared to normally

reared adult hamsters, suggesting that the RF enlargement was not caused by alterations in the cation-

chloride co-transporters.

A GABA-BDNF feedback loop maintains inhibitory networks, thereby maintaining RF refinement in

adulthood

GABA-GABA$_A$R interaction is known to regulate various downstream signaling pathways, and a

major regulator of GABA itself is BDNF-TrkB signaling triggered by NMDA receptor activity (Marini et

al., 1998). Our data are consistent with previous studies suggesting a positive feedback loop between the

BDNF-TrkB pathway and GABA expression, in which GABA facilitates BDNF expression, and BDNF

facilitates the production of GABA by GAD (Sánchez-Huertas and Rico, 2010) and its synaptic release

(Huang et al., 1999; Morales et al., 2002; Gianfranceschi et al., 2003; Jovanovic et al., 2004; Kuczewski

et al., 2008; Porcher et al., 2011; Hanno-Iijima et al., 2015), maintaining RF size and visual acuity

through GABAergic lateral inhibition (Mudd et al., 2019). Signaling via the MAPK cascade and the

transcription factor cAMP-responsive element-binding protein (CREB) appears to play a substantial role

in this process (Obrietan et al., 2002; Sánchez-Huertas and Rico, 2010). BDNF-TrkB interaction leads to

dimerization and auto-phosphorylation of TrkB, thereby triggering MAPK, PLC gamma, and PI3K

pathways (Yoshii and Constantine-Paton, 2007). These pathways in turn lead to the activation of

downstream effectors and mediators to initiate a CREB-dependent transcription process that can lead to

an increase in GABA$_A$R levels as well as more BDNF production (Huang and Reichardt, 2003; Yoshii

and Constantine-Paton, 2007; Porcher et al., 2011; Esvald et al., 2020). In addition, an increase in the

transmembrane localization of GABA$_A$Rs is mediated by BDNF-dependent inhibition of receptor

internalization in addition to ongoing reinsertion of the receptor into the postsynaptic membrane (Porcher

et al., 2011). This positive feedback regulation is critical in developing neurons and hence constituted a

major motivation for the work reported here. In this study however, neither the GABA$_A$R subunit

composition at the synapse nor subunit composition in the extrasynaptic regions was affected by dark

rearing. Chloride transport proteins (KCC2, NKCC2) also remained at normal levels. One possibility is

that GABA expression levels alone are the key factor in RF re-enlargement in hamster SC (Carrasco et

al., 2011; Mudd et al., 2019). If so, it would suggest that this type of delayed plasticity resulting from a

lack of early visual experience occurs through a different mechanism than described in other types of

plasticity resulting from dark rearing (Mower et al., 1985; Chen et al., 2000; Chen et al., 2001).

It is possible that changes in $GABA_AR$ signaling occurred that are not reflected here in the expression

levels of postsynaptic receptor composition, scaffolding molecules, or ion transporters, or that we missed

some transient changes in $GABA_AR$ signaling-associated proteins that cause $GABA_AR$ functional

changes. In the future, it would be interesting to study protein localization and interactions with

immunohistochemistry, or to study the properties of synaptic and extrasynaptic responses in the SC with

patch clamp experiments, and the subcellular dynamics of associated proteins involved. This might

improve the understanding of the molecular processes active in this deprivation-induced, maladaptive

plasticity in the SC.

Suggested alternative explanations include that the changes leading to RF expansion and thus the

visual acuity deficits take place earlier than the time point that we studied and return to normal by P90.

The width of the synaptic cleft decreases during development (Li and Cline, 2010), thus, another

interesting possibility is that dark rearing from birth gradually increases the width of the synaptic cleft in

adulthood while keeping the postsynaptic signaling components in place. Alternatively, the synapses may

be present but silent due to the presynaptic loss of GABA (Carrasco et al., 2011). It is also possible that

other scaffolding proteins and their partners are involved. Potential candidates include gephyrin binding

partners such as $GABA_AR$ beta2 and beta3 subunits (Kowalczyk et al., 2013), the scaffold protein radixin

that binds $GABA_AR$ alpha5 subunit to the actin cytoskeleton (Loebrich et al., 2006) and regulates

synaptic $GABA_AR$ density (Hausrat et al., 2015), or gephyrin post-translational modification events that

influence inhibitory synaptic plasticity by affecting postsynaptic scaffolding (Zacchi et al., 2014). More

recently, distinct spectrin isoforms have been shown to affect synaptic inhibition by selectively targeting

specific $GABA_AR$ subunits, including α1 and α2, to particular regions of the neuron (Smalley et al.,

2023).

In summary, our results argue that visual experience is not necessary to maintain mature levels and

composition of several postsynaptic proteins that are essential for retino-SC synaptic communication.

Unlike many previous dark rearing studies in which both GABA and $GABA_A$ receptors were found to be

downregulated (Carmignoto and Vicini, 1992; Chen et al., 2000; Chen et al., 2001; Kilman et al., 2002;

Nahmani and Turrigiano, 2014), we report here that in the SC, $GABA_A R$ levels, subunit composition, and

localization in adulthood are unaffected by dark rearing. The scaffold proteins gephyrin and PSD-95, and

the chloride transporters KCC2 and NKCC2 are also not affected by dark rearing. This novel,

experience-dependent form of adult plasticity may involve an as yet unidentified postsynaptic

mechanism, or only a reduction in GABA release (Carrasco et al., 2005; Carrasco and Pallas, 2006;

Carrasco et al., 2011), thereby challenging the common view that presynaptic changes in ligand

availability are always associated with matching postsynaptic changes in their receptors. Either possibility

is encouraging with respect to understanding this form of adult plasticity and might help adults with

memory impairments, traumatic brain injury, or inhibition-associated neurological disorders.

REFERENCES

Adler S. 1948. Origin of the Golden Hamster Cricetus auratus as a laboratory animal. Nature 162:256-
257.
Balmer TS, Pallas SL. 2015. Refinement but not maintenance of visual receptive fields is independent of
visual experience. Cerebral Cortex 25:904-917.
Balmer TS, Pallas SL. 2015. Visual experience prevents dysregulation of GABA$_B$ receptor-dependent
short-term depression in adult superior colliculus. J Neurophysiol 113:2049-2061.
Baroncelli L, Sale A, Viegi A, Maya Vetencourt JF, De Pasquale R, Baldini S, Maffei L. 2010. Experience-
dependent reactivation of ocular dominance plasticity in the adult visual cortex. Exp Neurol
226:100-109.
Brady ML, Jacob TC. 2015. Synaptic localization of α5 GABA (A) receptors via gephyrin interaction
regulates dendritic outgrowth and spine maturation. Devel Neurobiol 75:1241-1251.
Cancedda L, Fiumelli H, Chen K, Poo M-m. 2007. Excitatory GABA action is essential for morphological
maturation of cortical neurons *in vivo*. J Neurosci 27:5224-5235.
Capogna M, Castillo PE, Maffei A. 2021. The ins and outs of inhibitory synaptic plasticity: Neuron types,
molecular mechanisms and functional roles. Eur J Neurosci 54:6882-6901.
Carmignoto G, Vicini S. 1992. Activity-dependent decrease in NMDA receptor responses during
development of the visual cortex. Science 258:1007-1011.
Carrasco MM, Mao YT, Balmer TS, Pallas SL. 2011. Inhibitory plasticity underlies visual deprivation-
induced loss of receptive field refinement in the adult superior colliculus. Eur J Neurosci 33:58-
68.
Carrasco MM, Pallas SL. 2006. Early visual experience prevents but cannot reverse deprivation-induced
loss of refinement in adult superior colliculus. Vis Neurosci 23:845-852.
Carrasco MM, Razak KA, Pallas SL. 2005. Visual experience is necessary for maintenance but not
development of receptive fields in superior colliculus. J Neurophysiol 94:1962-1970.
Chalupa LM. 1981. Some observations on the functional organization of the golden hamster's visual
system. Behav Brain Res 3:189-200.
Chen L, Cooper NGF, Mower GD. 2000. Developmental changes in the expression of NMDA receptor
subunits (NR1, NR2A, NR2B) in the cat visual cortex and the effects of dark rearing. Molec Brain
Res 78:196-200.
Chen L, Yang C, Mower GD. 2001. Developmental changes in the expression of GABA(A) receptor
subunits (alpha(1), alpha(2), alpha(3)) in the cat visual cortex and the effects of dark rearing.
Molec Brain Res 88:135-143.
Chen X, Levy JM, Hou A, Winters C, Azzam R, Sousa AA, Leapman RD, Nicoll RA, Reese TS. 2015. PSD-95
family MAGUKs are essential for anchoring AMPA and NMDA receptor complexes at the
postsynaptic density. Proc Natl Acad Sci U S A 112:E6983-E6992.
Cherubini E, Gaiarsa JL, Ben-Ari Y. 1991. GABA: An excitatory transmitter in early postnatal life. Trends
in Neurosciences 14:515-519.
Chevy Q, Heubl M, Goutierre M, Backer S, Moutkine I, Eugène E, Bloch-Gallego E, Lévi S, Poncer JC. 2015.
KCC2 gates activity-driven AMPA receptor traffic through cofilin phosphorylation. J Neurosci
35:15772.
Coorssen JR, Blank PS, Albertorio F, Bezrukov L, Kolosova I, Backlund PS, Zimmerberg J. 2002.
Quantitative femto- to attomole immunodetection of regulated secretory vesicle proteins
critical to exocytosis. Analytical Biochemistry 307:54-62.
Cynader M, Mitchell DE. 1980. Prolonged sensitivity to monocular deprivation in dark-reared cats. J
Neurophysiol 43:1026-1040.

Davenport CM, Rajappa R, Katchan L, Taylor CR, Tsai MC, Smith CM, de Jong JW, Arnold DB, Lammel S,
Kramer RH. 2021. Relocation of an extrasynaptic GABA(A) receptor to inhibitory synapses
freezes excitatory synaptic strength and preserves memory. Neuron 109:123-134.
Deidda G, Allegra M, Cerri C, Naskar S, Bony G, Zunino G, Bozzi Y, Caleo M, Cancedda L. 2015. Early
depolarizing GABA controls critical-period plasticity in the rat visual cortex. Nat Neurosci 18:87-
96.
Diamond M, Yanagimachi R. 1970. Reproductive development in the female Golden Hamster in relation
to spontaneous estrus. Biol Reprod 2:223-229.
Dunning DD, Hoover CL, Soltesz I, Smith MA, O'Dowd DK. 1999. GABAA Receptor–mediated miniature
postsynaptic currents and α-subunit expression in developing cortical neurons. J Neurophysiol
82:3286-3297.
Erisir A, Harris JL. 2003. Decline of the critical period of visual plasticity is concurrent with the reduction
of NR2B subunit of the synaptic NMDA receptor in layer 4. J Neurosci 23:5208-5218.
Esvald EE, Tuvikene J, Sirp A, Patil S, Bramham CR, Timmusk T. 2020. CREB family transcription factors
are major mediators of BDNF transcriptional autoregulation in cortical neurons. J Neurosci
40:1405-1426.
Fagiolini M, Fritschy JM, Low K, Mohler H, Rudolph U, Hensch TK. 2004. Specific GABAA circuits for visual
cortical plasticity. Science 303:1681-1683.
Fagiolini M, Pizzorusso T, Berardi N, Domenici L, Maffei L. 1994. Functional postnatal development of
the rat primary visual cortex and the role of visual experience: Dark rearing and monocular
deprivation. Vision Res 34:709-720.
Farrant M, Nusser Z. 2005. Variations on an inhibitory theme: phasic and tonic activation of GABAA
receptors. Nat Rev Neurosci 6:215-229.
Firth SI, Wang CT, Feller MB. 2005. Retinal waves: mechanisms and function in visual system
development. Cell Calcium 37:425-432.
Fisher-Lavie A, Ziv NE. 2013. Matching dynamics of presynaptic and postsynaptic scaffolds. J Neurosci
33:13094-13100.
Fitzgerald K, Zucker I. 1976. Circadian organization of the estrous cycle of the golden hamster. Proc Natl
Acad Sci USA 73:2923-2927.
Frederikse PH, Nandanoor A, Kasinathan C. 2016. "Moonlighting" GAPDH Protein Localizes with AMPA
Receptor GluA2 and L1 Axonal Cell Adhesion Molecule at Fiber Cell Borders in the Lens. Curr Eye
Res 41:41-49.
Fritschy J-M, Paysan J, Enna A, Mohler H. 1994. Switch in the expression of rat GABAA-receptor subtypes
during postnatal development: An immunohistochemical study. J Neurosci 14:5302-5324.
Funahashi R, Maruyama T, Yoshimura Y, Komatsu Y. 2013. Silent synapses persist into adulthood in layer
2/3 pyramidal neurons of visual cortex in dark-reared mice. J Neurophysiol 109:2064-2076.
Gauvain G, Chamma I, Chevy Q, Cabezas C, Irinopoulou T, Bodrug N, Carnaud M, Lévi S, Poncer JC. 2011.
The neuronal K-Cl cotransporter KCC2 influences postsynaptic AMPA receptor content and
lateral diffusion in dendritic spines. Proc Natl Acad Sci U S A 108:15474-15479.
Ghosh R, Gilda JE, Gomes AV. 2014. The necessity of and strategies for improving confidence in the
accuracy of western blots. Expert Review of Proteomics 11:549-560.
Gianfranceschi L, Siciliano R, Walls J, Morales B, Kirkwood A, Huang ZJ, Tonegawa S, Maffei L. 2003.
Visual cortex is rescued from the effects of dark rearing by overexpression of BDNF. Proc Natl
Acad Sci U S A 100:12486-12491.
Hanno-Iijima Y, Tanaka M, Iijima T. 2015. Activity-dependent bidirectional regulation of GAD expression
in a homeostatic fashion Is mediated by BDNF-dependent and independent pathways. PloS one
10:e0134296.

Hanover JL, Huang ZJ, Tonegawa S, Stryker MP. 1999. Brain-derived neurotrophic factor overexpression induces precocious critical period in mouse visual cortex. J Neurosci 19:RC40 (41-45).

Hausrat TJ, Muhia M, Gerrow K, Thomas P, Hirdes W, Tsukita S, Heisler FF, Herich L, Dubroqua S, Breiden P, Feldon J, Schwarz JR, Yee BK, Smart TG, Triller A, Kneussel M. 2015. Radixin regulates synaptic GABAA receptor density and is essential for reversal learning and short-term memory. Nat Commun 6:6872.

Hensch TK, Quinlan EM. 2018. Critical periods in amblyopia. Vis Neurosci 35:E014.

Huang EJ, Reichardt LF. 2003. Trk receptors: roles in neuronal signal transduction. Annu Rev Biochem 72:609-642.

Huang X, Stodieck SK, Goetze B, Cui L, Wong MH, Wenzel C, Hosang L, Dong Y, Lowel S, Schluter OM. 2015. Progressive maturation of silent synapses governs the duration of a critical period. Proc Natl Acad Sci USA 112:E3131-E3140.

Huang ZJ, Kirkwood A, Pizzorusso T, Porciatti V, Morales B, Bear MF, Maffei L, Tonegawa S. 1999. BDNF regulates the maturation of inhibition and the critical period of plasticity in mouse visual cortex. Cell 98:739-755.

Hubel DH, Wiesel TN. 1970. The period of susceptibility of the physiological effects of unilateral eye closure in kittens. J Physiol (Lond) 206:419-436.

Hübener M, Bonhoeffer T. 2014. Neuronal Plasticity: Beyond the Critical Period. Cell 159:727-737.

Huck UW, Lisk RD, McKay MV. 1988. Social dominance and reproductive success in pregnant and lactating golden hamsters (Mesocricetus auratus) under seminatural conditions. Physiol Behav 44:313-319.

Huhman KL, Albers HE. 1994. Neuropeptide Y microinjected into the suprachiasmatic region phase shifts circadian rhythms in constant darkness. Peptides 15:1475-1478.

Jacob TC. 2019. Neurobiology and therapeutic potential of α5-GABA type A receptors. Front Molec Neurosci 12:179.

Jacob TC, Bogdanov YD, Magnus C, Saliba RS, Kittler JT, Haydon PG, Moss SJ. 2005. Gephyrin regulates the cell surface dynamics of synaptic GABAA receptors. J Neurosci 25:10469-10478.

Jacob TC, Moss SJ, Jurd R. 2008. GABAA receptor trafficking and its role in the dynamic modulation of neuronal inhibition. Nat Rev Neurosci 9:331-343.

Jovanovic JN, Thomas P, Kittler JT, Smart TG, Moss SJ. 2004. Brain-Derived Neurotrophic Factor modulates fast synaptic inhibition by regulating GABAA receptor phosphorylation, activity, and cell-surface stability. J Neurosci 24:522-530.

Kalogeraki E, Greifzu F, Haack F, Lowel S. 2014. Voluntary physical exercise promotes ocular dominance plasticity in adult mouse primary visual cortex. J Neurosci 34:15476-15481.

Katz LC, Shatz CJ. 1996. Synaptic activity and the construction of cortical circuits. Science 274:1133-1138.

Kilman V, van Rossum MC, Turrigiano GG. 2002. Activity deprivation reduces miniature IPSC amplitude by decreasing the number of postsynaptic GABA(A) receptors clustered at neocortical synapses. J Neurosci 22:1328-1337.

Kimoto S, Zaki MM, Bazmi HH, Lewis DA. 2015. Altered markers of cortical gamma-Aminobutyric Acid neuronal activity in schizophrenia: Role of the NARP gene. JAMA psychiatry 72:747-756.

Kittler JT, Delmas P, Jovanovic JN, Brown DA, Smart TG, Moss SJ. 2000. Constitutive endocytosis of GABAA receptors by an association with the Adaptin AP2 complex modulates inhibitory synaptic currents in hippocampal neurons. J Neurosci 20:7972-7977.

Kneussel M, Brandstätter JH, Laube B, Stahl S, Müller U, Betz H. 1999. Loss of postsynaptic GABAA receptor clustering in gephyrin-deficient mice. J Neurosci 19:9289-9297.

Kneussel M, Helmut Brandstätter J, Gasnier B, Feng G, Sanes JR, Betz H. 2001. Gephyrin-independent clustering of postsynaptic GABAA receptor subtypes. Molec Cell Neurosci 17:973-982.

Kowalczyk S, Winkelmann A, Smolinsky B, Förstera B, Neundorf I, Schwarz G, Meier JC. 2013. Direct
binding of GABAA receptor β2 and β3 subunits to gephyrin. Eur J Neurosci 37:544-554.
Kuczewski N, Langlois A, Fiorentino H, Bonnet S, Marissal T, Diabira D, Ferrand N, Porcher C, Gaiarsa JL.
2008. Spontaneous glutamatergic activity induces a BDNF-dependent potentiation of GABAergic
synapses in the newborn rat hippocampus. J Physiol 586:5119-5128.
Kutsarova E, Munz M, Ruthazer ES. 2017. Rules for Shaping Neural Connections in the Developing Brain.
Frontiers in neural circuits 10:111.
Laurie DJ, Wisden W, Seeburg PH. 1992. The distribution of thirteen GABAA receptor subunit mRNAs in
the rat brain. III. Embryonic and postnatal development. J Neurosci 12:4151-4172.
Lee H, Chen CX, Liu YJ, Aizenman E, Kandler K. 2005. KCC2 expression in immature rat cortical neurons is
sufficient to switch the polarity of GABA responses. Eur J Neurosci 21:2593-2599.
Lee WC, Nedivi E. 2002. Extended plasticity of visual cortex in dark-reared animals may result from
prolonged expression of cpg15-like genes. J Neurosci 22:1807-1815.
Lévi S, Triller A. 2006. Neurotransmitter Dynamics. In: Kittler JT, Moss SJ, Kittler JT, Moss SJs. The
Dynamic Synapse: Molecular Methods in Ionotropic Receptor Biology. Boca Raton, FL: CRC
Press/Taylor & Francis.
Levitt P. 2005. Disruption of interneuron development. Epilepsia 46:22-28.
Li H, Khirug S, Cai C, Ludwig A, Blaesse P, Kolikova J, Afzalov R, Coleman SK, Lauri S, Airaksinen MS,
Keinänen K, Khiroug L, Saarma M, Kaila K, Rivera C. 2007. KCC2 interacts with the dendritic
cytoskeleton to promote spine development. Neuron 56:1019-1033.
Li J, Cline HT. 2010. Visual deprivation increases accumulation of dense core vesicles in developing optic
tectal synapses in Xenopus laevis. J Comp Neurol 518:2365-2381.
Loebrich S, Bähring R, Katsuno T, Tsukita S, Kneussel M. 2006. Activated radixin is essential for GABAA
receptor α5 subunit anchoring at the actin cytoskeleton. EMBO J 25:987-999.
Maffei L, Galli-Resta L. 1990. Correlation in the discharge of neighboring rat retinal ganglion cells during
prenatal life. Proc Natl Acad Sci, USA 87:2861-2864.
Marini AM, Rabin SJ, Lipsky RH, Mocchetti I. 1998. Activity-dependent release of Brain-derived
Neurotrophic Factor underlies the neuroprotective effect of N-Methyl-D-aspartate. J Biol Chem
273:29394-29399.
Meister M, Wong RO, Baylor DA, Shatz CJ. 1991. Synchronous bursts of action potentials in ganglion cells
of the developing mammalian retina. Science 252:939-943.
Minier F, Sigel E. 2004. Positioning of the alpha-subunit isoforms confers a functional signature to
gamma-aminobutyric acid type A receptors. Proc Natl Acad Sci U S A 101:7769-7774.
Mitchell DE, Maurer D. 2022. Critical Periods in Vision Revisited. Ann Rev Vision Sci 8:291-321.
Moore YE, Conway LC, Wobst HJ, Brandon NJ, Deeb TZ, Moss SJ. 2019. Developmental Regulation of
KCC2 Phosphorylation Has Long-Term Impacts on Cognitive Function. Front Mol Neurosci
12:173.
Morales B, Choi S-Y, Kirkwood A. 2002. Dark rearing alters the development of GABAergic transmission
in visual cortex. J Neurosci 22:8084-8090.
Mower GD. 1991. The effect of dark rearing on the time course of the critical period in cat visual cortex.
Devel Brain Res 58:151-158.
Mower GD, Caplan CJ, Christen WG, Duffy FH. 1985. Dark rearing prolongs physiological but not
anatomical plasticity of the cat visual cortex. J Comp Neurol 235:448-466.
Mower GD, Christen WG. 1985. Role of visual experience in activating critical period in cat visual cortex.
J Neurophysiol 53:572-589.
Mower GD, White WF, Rustad R. 1986. [3H]muscimol binding of GABA receptors in the visual cortex of
normal and monocularly deprived cats. Brain Res 380:253-260.

Mudd DB, Balmer TS, Kim SY, Machhour N, Pallas SL. 2019. TrkB activation during a critical period mimics
the protective effects of early visual experience on perception and the stability of receptive
fields in adult superior colliculus. J Neurosci 39:4475-4488.
Nahmani M, Turrigiano GG. 2014. Deprivation-induced strengthening of presynaptic and postsynaptic
inhibitory transmission in layer 4 of visual cortex during the critical period. J Neurosci 34:2571-
2582.
Nakadate K, Imamura K, Watanabe Y. 2012. Effects of monocular deprivation on the spatial pattern of
visually induced expression of c-Fos protein. Neuroscience 202:17-28.
Nowosielski-Slepowron B, Park A. 1987. Variation in the growth of the preweaning Syrian hamster
(Cricetus auratus). Acta Morphol Neerl Scand 25:187-199.
Obrietan K, Gao X-B, van den Pol AN. 2002. Excitatory actions of GABA Increase BDNF expression via a
MAPK-CREB–dependent mechanism—A positive feedback circuit in developing neurons. J
Neurophysiol 88:1005-1015.
Okada M, Onodera K, Van Renterghem C, Sieghart W, Takahashi T. 2000. Functional correlation of
GABAA receptor α subunits expression with the properties of IPSCs in the developing thalamus. J
Neurosci 20:2202-2208.
Pallas SL. 2017. The impact of ecological niche on adaptive flexibility of sensory circuitry. Frontiers in
neuroscience 11:344.
Pennacchietti F, Vascon S, Nieus T, Rosillo C, Das S, Tyagarajan SK, Diaspro A, Del Bue A, Petrini EM,
Barberis A, Cella Zanacchi F. 2017. Nanoscale molecular reorganization of the inhibitory
postsynaptic density is a determinant of GABAergic synaptic potentiation. J Neurosci 37:1747-
1756.
Pfeffer CK, Stein V, Keating DJ, Maier H, Rinke I, Rudhard Y, Hentschke M, Rune GM, Jentsch TJ, Hübner
CA. 2009. NKCC1-dependent GABAergic excitation drives synaptic network maturation during
early hippocampal development. J Neurosci 29:3419-3430.
Pfeiffer F, Graham D, Betz H. 1982. Purification by affinity chromatography of the glycine receptor of rat
spinal cord. J Biol Chem 257:9389-9393.
Porcher C, Hatchett C, Longbottom RE, McAinch K, Sihra TS, Moss SJ, Thomson AM, Jovanovic JN. 2011.
Positive feedback regulation between {gamma}-Aminobutyric Acid Type A (GABAA) receptor
signaling and Brain-derived Neurotrophic Factor (BDNF) release in developing neurons. J Biol
Chem 286:21667-21677.
Pratt NC, Lisk RD. 1989. Effects of social stress during early pregnancy on litter size and sex ratio in the
golden hamster (Mesocricetus auratus). J Reprod Fertil 87:763-769,.
Razak KA, Huang L, Pallas SL. 2003. NMDA receptor blockade in the superior colliculus increases
receptive field size without altering velocity and size tuning. J Neurophysiol 90:110-119.
Reh RK, Dias BG, Nelson CA, 3rd, Kaufer D, Werker JF, Kolb B, Levine JD, Hensch TK. 2020. Critical period
regulation across multiple timescales. Proc Natl Acad Sci U S A.
Ribic A. 2020. Stability in the Face of Change: Lifelong Experience-Dependent Plasticity in the Sensory
Cortex. Frontiers in Cellular Neuroscience 14.
Rivera C, Voipio J, Payne JA, Ruusuvuori E, Lahtinen H, Lamsa K, Pirvola U, Saarma M, Kaila K. 1999. The
K+/Cl− co-transporter KCC2 renders GABA hyperpolarizing during neuronal maturation. Nature
397:251-255.
Saab BJ, MacLean AJ, Kanisek MM, Zurek AA, Martin LJ, Roder JC, Orser BA. 2010. Short-term memory
impairment after isoflurane in mice is prevented by the α5 γ-aminobutyric acid Type A receptor
inverse agonist L-655,708. Anesthesiol 113:1061-1071.
Sale A, Berardi N, Spolidoro M, Baroncelli L, Maffei L. 2010. GABAergic inhibition in visual cortical
plasticity. Frontiers in Cellular Neuroscience 4.

Sánchez-Huertas C, Rico B. 2010. CREB-dependent regulation of GAD65 transcription by BDNF/TrkB in
cortical interneurons. Cereb Cortex 21:777-788.
Sanderson TM, Georgiou J, Collingridge GL. 2020. Illuminating Relationships Between the Pre- and Post-
synapse. Frontiers in neural circuits 14:9.
Schmidt MJ, Mirnics K. 2015. Neurodevelopment, GABA system dysfunction, and schizophrenia.
Neuropsychopharmacology : official publication of the American College of
Neuropsychopharmacology 40:190-206.
Shi J, Aamodt SM, Constantine-Paton M. 1997. Temporal correlations between functional and molecular
changes in NMDA receptors and GABA neurotransmission in the superior colliculus. J Neurosci
17:6264-6276.
Sigel E, Baur R, Trube G, Möhler H, Malherbe P. 1990. The effect of subunit composition of rat brain
GABAA receptors on channel function. Neuron 5:703-711.
Sigel E, Steinmann ME. 2012. Structure, function, and modulation of GABA(A) receptors. J Biol Chem
287:40224-40231.
Smalley JL, Cho N, Ng SFJ, Choi C, Lemons AHS, Chaudry S, Bope CE, Dengler JS, Zhang C, Rasband MN,
Davies PA, Moss SJ. 2023. Spectrin-beta 2 facilitates the selective accumulation of GABAA
receptors at somatodendritic synapses. Communications Biology 6:11.
Sudhof TC. 2018 Towards an Understanding of Synapse Formation. Neuron 100:276-293.
Sun C, Sieghart W, Kapur J. 2004. Distribution of alpha1, alpha4, gamma2, and delta subunits of GABAA
receptors in hippocampal granule cells. Brain research 1029:207-216.
Takesian AE, Hensch TK. 2013. Balancing plasticity/stability across brain development. Prog Brain Res
207:3-34.
Tang X, Jaenisch R, Sur M. 2021. The role of GABAergic signalling in neurodevelopmental disorders.
Nature Rev Neurosci 22:290-307.
Tognini P, Manno I, Bonaccorsi J, Cenni MC, Sale A, Maffei L. 2012. Environmental enrichment promotes
plasticity and visual acuity recovery in adult monocular amblyopic rats. PloS one 7:e34815.
Toyoizumi T, Miyamoto H, Yazaki-Sugiyama Y, Atapour N, Hensch Takao K, Miller Kenneth D. 2013. A
theory of the transition to critical period plasticity: Inhibition selectively suppresses spontaneous
activity. Neuron 80:51-63.
Tretter V, Jacob TC, Mukherjee J, Fritschy JM, Pangalos MN, Moss SJ. 2008. The clustering of GABA(A)
receptor subtypes at inhibitory synapses is facilitated via the direct binding of receptor alpha 2
subunits to gephyrin. J Neurosci 28:1356-1365.
Tyson JA, Anderson SA. 2014. GABAergic interneuron transplants to study development and treat
disease. Trends Neurosci 37:169-177.
Viegi A, Cotrufo T, Berardi N, Mascia L, Maffei L. 2002. Effects of dark rearing on phosphorylation of
neurotrophin Trk receptors. Eur J Neurosci 16:1925-1930.
Wiesel TN, Hubel DH. 1965. Comparison of the effects of unilateral and bilateral eye closure on cortical
unit responses in kittens. J Neurophysiol 28:1029-1040.
Wong ROL, Meister M, Shatz CJ. 1993. Transient period of correlated bursting activity during
development of the mammalian retina. Neuron 11:923-938.
Yoshii A, Constantine-Paton M. 2007. BDNF induces transport of PSD-95 to dendrites through PI3K-AKT
signaling after NMDA receptor activation. Nature Neurosci 10:702-711.
Yu Z-Y, Wang W, Fritschy J-M, Witte OW, Redecker C. 2006. Changes in neocortical and hippocampal
GABAA receptor subunit distribution during brain maturation and aging. Brain Research
1099:73-81.
Zacchi P, Antonelli R, Cherubini E. 2014. Gephyrin phosphorylation in the functional organization and
plasticity of GABAergic synapses. Front Cell Neurosci 8:103.

Zhang H, Mu L, Wang D, Xia D, Salmon A, Liu Q, Wong-Riley MTT. 2018. Uncovering a critical period of
synaptic imbalance during postnatal development of the rat visual cortex: role of brain-derived
neurotrophic factor. J Physiol 596:4511-4536.
Zurek AA, Bridgwater EM, Orser BA. 2012. Inhibition of alpha5 gamma-Aminobutyric acid type A
receptors restores recognition memory after general anesthesia. Anesth Analg 114:845-855.
Zurek AA, Yu J, Wang D-S, Haffey SC, Bridgwater EM, Penna A, Lecker I, Lei G, Chang T, Salter EWR, Orser
BA. 2014. Sustained increase in α5 GABAA receptor function impairs memory after anesthesia. J
Clin Invest 124:5437-5441.

Figure 1: Differences in the visual refinement of hamster visual pathways depending on exposure to light during the critical period of heightened neural plasticity. Animals exposed to light during postnatal development gradually improve RF refinement in SC and V1 and maintain it throughout life (as indicated by the blue line). Animals that do not experience postnatal light also show RF refinement by P60, but the refinement progressively declines in adulthood (as indicated by the orange line).

Figure 2. Histone H3 expression in the synaptosomal and cytosolic lysates. Western blot showing histone H3 (15 kDa) bands in the synaptosomal and cytosolic fractions of SC. GAPDH was used as a loading control. The presence of histone H3 in the cytosolic fraction and its absence in the synaptosomal fraction shows an effective synaptosomal separation occurred in these experiments.

Figure 3. GABA$_A$α2 receptor subunit levels in SC are not affected by early dark rearing. (A) Image: Individual Western blots of normally reared (NR) and dark-reared (DR) treatment groups generated using 20 µg of SC protein per lane. GAPDH was used as a loading control. **Plot:** Boxplot showing the normalized GABA$_A$Rα2 expression level in normally reared vs. dark reared hamsters. **(B) Image:** Individual Western blots of SC tissue from NR and DR animals comparing GABA$_A$Rα1 and GABA$_A$Rα2 expression with corresponding GAPDH expression. **Plot:** Boxplot showing the ratio of normalized values of GABA$_A$Rα1/GAPDH to the normalized values of GABA$_A$Rα2/GAPDH. Boxes in each individual boxplot show the median and 25th and 75th percentiles of the data (whiskers show 5% and 95% levels). Individual data points obtained from each animal within a group are shown as dots. For Western blots in this and the following figures, lanes presented together are from the same gel, and each measured protein was normalized against GAPDH (unless stated otherwise) as a loading control. Taken together, these results reveal that the levels and ratio of synaptic GABA$_A$Rα2 receptor subunits and their ratio with GABA$_A$Rα1 subunits are similar in normal and dark reared adult SC.

Figure 4. GABA$_A$α5 receptor subunit levels in SC are not affected by early dark rearing. A and B

Images: Representative Western blots of NR and DR treatment groups as in Figure 3. All lanes presented

together are from the same gel(s), and each gel was run with GAPDH as a loading control. **(A) Plot:**

Boxplot showing normalized GABA$_A$Rα5 expression levels compared between normal and dark reared

hamsters. **(B) Plot:** Boxplot showing the ratio of normalized (against GAPDH levels) GABA$_A$Rα1:

GABA$_A$Rα5 expression ratios. Boxes in each individual boxplot show the median and 25th and 75th

percentiles of the data, with whiskers at 5 and 95%. Individual data points in each group are shown as

dots. These results show that the levels of synaptic GABA$_A$α5 receptor subunits and their ratio with

GABA$_A$Rα1 subunits are similar in normal and dark reared adult SC.

Figure 5. Internalization of GABA$_A$ receptors in SC is not affected by early dark rearing. **(A)** Adult

levels of the cytosolic vs. the synaptic membrane-attached ratio of the synaptically-targeted GABA$_A$Rα1

and **(B)** the synaptically-targeted GABA$_A$Rα5 subunits were not affected by early light deprivation.

Images: Representative Western blots represent bands of cytoplasmic and membrane bound receptor

subunit proteins, each from the same animal, measured as a ratio against GAPDH and compared between

NR and DR groups. **Plots:** Boxplot showing the ratio of normalized (against GAPDH) values of cytosol:

membrane ratios of each subunit. Boxes in each individual boxplot show the median and 25th and 75th

percentile of the data, with whiskers indicating 5 and 95%. Individual data points obtained in each group

are shown as dots. These results show that the internalization of synaptic GABA$_A$Rα1 and GABA$_A$Rα5

subunits is similar in normal and dark reared adult SC.

Figure 6. Gephyrin and PSD-95 expression in SC are not affected by dark rearing. **Images: (A)**

Gephyrin and **(B)** PSD-95 expression was similar between adult NR and DR groups (upper panels).

Plots: Boxplot showing the ratio of normalized values of gephyrin vs. GAPDH (A) or β-actin (B) . Boxes

in each individual boxplot show the median and 25th and 75th percentiles of the data, with whiskers at 5

and 95%. Individual data points obtained in each group are shown as dots. These results show that the

levels and ratio of the scaffold proteins are similar in normal and dark reared adult SC.

Figure 7. Cation-chloride co-transporter expression in SC is not affected by early dark rearing.

Images: Example Western blots of NR and DR samples labeled for cation-chloride co-transporters **(A)**

KCC2 (140 kDa) and **(B)** NKCC1 (150 kDa) compared to the GAPDH loading control and **(C)** a

comparison of the within subject ratio of KCC2:NKCC1 in NR and DR adult hamsters. **Plots:** Boxplots

showing the levels of KCC2 and NKCC1 proteins, normalized against GAPDH **(A and B, respectively)**

and comparison of normalized values of KCC2/GAPDH to that of NKCC1/GAPDH **(C)**. Boxes in each

boxplot show the median, 25th and 75th percentiles of the data. Whiskers are at 5 and 95% percentiles.

Individual data points obtained in each group are shown as dots. These results show that the number and

ratio of cation chloride co-transporters are similar in normal and dark reared adult SC.