Multiple Postsynaptic Protein Levels in Adult Superior Colliculus Are Unaffected by Dark Rearing from Birth

Abbreviated title: **Postsynaptic proteins unaffected by dark rearing**

Parag S. Juvale[1], David B. Mudd[2#], Nitheyaa Shree[2¶], and Sarah L. Pallas[1]*

[1] Department of Biology, University of Massachusetts Amherst, Amherst, MA 01003, USA

[2] Neuroscience Institute, Georgia State University, Atlanta, GA 30303, USA

\# Current address: Office of Technology Transfer, Emory University, Atlanta, GA

¶Current address: Graduate Program in Biochemistry, University of Bristol, UK

Conflict of interest: *The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.*

Author Contributions: Juvale, Mudd, and Pallas contributed to the conception and design of the study. Juvale, Mudd, and Shree organized the database and performed the statistical analysis. Mudd wrote the first draft of the manuscript. Juvale wrote subsequent versions of the manuscript. Pallas edited the manuscript. All authors read the manuscript, contributed to manuscript revision, and approved the submitted version.

Funding: Support for this work was provided by a Sigma Xi GIAR grant and a UMass Amherst predissertation grant to P.S.J.; a GSU Brains and Behavior Fellowship and a GSU Center for Neuromics student grant awarded to D.B.M.; and a GSU Brains and Behavior Seed grant, UMass startup funds, a National Science Foundation grant (IOS-1656838), and a DARPA grant (HR0011-18-2-0019, TA2) awarded to S.L.P.

*Correspondence:

Sarah L. Pallas

Department of Biology

University of Massachusetts-Amherst

Amherst, MA 01003 USA

spallas@umass.edu

Keywords: Visual deprivation, superior colliculus, GABA signaling, adult plasticity, visual refinement, $GABA_A$ receptor, scaffold protein, chloride co-transporter

Number of pages: 25

Number of words: 9222

Number of figures: 6

Number of tables: 0

Number of words in Abstract: 205

Acknowledgments: We would like to thank Drs. Angela Mabb, Larry Schwartz, Thomas Maresca, and Lillian Fritz-Laylin for providing instrument access and guidance with Western Blot imaging and analysis, members of the Pallas lab for technical support and manuscript review, and the animal care staff at GSU and UMass.

Abstract

Dark rearing in kittens and monkeys delays visual pathway development and prolongs the critical period. In contrast, receptive fields (RFs) in superior colliculus (SC) and visual cortex (V1) of Syrian hamsters refine normally with spontaneous activity alone, requiring only brief juvenile visual experience to maintain refined RFs in adulthood. Continued visual deprivation into adulthood reduces GAD and GABA levels due to reduced BDNF-TrkB signaling, resulting in RF re-enlargement. To explore possible postsynaptic changes caused by dark rearing, we examined expression of $GABA_A$ receptors and their anchoring protein gephyrin and cation-chloride co-transporters KCC2 and NKCC1 in dark reared hamsters. We found that dark rearing from birth until puberty had no effect on levels of postsynaptic $GABA_A$ receptors or their subunit composition, anchoring proteins, or chloride transporters. Thus, visual deprivation from birth to puberty reveals a new form of maladaptive, inhibitory plasticity in which, rather than extending the critical period, hamster receptive fields refine normally and then lose refinement in adulthood. These results suggest that attempts to increase plasticity in adulthood for rehabilitation or recovery from injury should consider the possibility of unintended negative consequences. In addition, our results demonstrate the interesting finding that changes in neurotransmitter levels are not necessarily coordinated with changes in postsynaptic components.

Contribution to the field statement

Raising animals in darkness from birth prevents development of the visual system in some mammals. In Syrian hamsters, however, the visual system develops normally in the dark, but continued visual deprivation into adulthood leads to a gradual re-expansion of receptive field size. We showed previously that a loss of presynaptic inhibition contributes to the reversal of visual refinement in adulthood. However, the postsynaptic mechanisms that contribute to the reversal of visual refinement are not yet known. Here we show that despite the reduction in GABAergic lateral inhibition there are no detectable changes in postsynaptic $GABA_A$ receptor levels or in several other postsynaptic proteins. These results

show that changes in presynaptic ligand levels do not always correspond with changes in their

postsynaptic receptors. They suggest that in some species dark rearing extended into adulthood can re-

establish plasticity, potentially producing negative effects on visual acuity.

Introduction

During brain development, synaptic connections are elaborated and then refined to a mature state

under the influence of neural activity. Activity due to sensory experience during an early "critical period"

is important for shaping some aspects of neural circuit development. In the visual pathway, although

spontaneous retinal activity is important for initial axon pruning, it has historically been thought that early

visual experience is essential to attain refined connectivity patterns during development (Maffei and

Galli-Resta, 1990; Meister et al., 1991; Wong et al., 1993; Katz and Shatz, 1996; Firth et al., 2005). Once

critical periods have closed, plasticity is often limited or even prevented, thus protecting refined circuits

from destabilization (Hubel and Wiesel, 1970; Takesian and Hensch, 2013; Hübener and Bonhoeffer,

2014; Pallas, 2017; Hensch and Quinlan, 2018; Reh et al., 2020; Ribic, 2020).

In some mammals, dark rearing is reported to delay or prevent refinement, prolonging critical period

plasticity (Cynader and Mitchell, 1980; Mower et al., 1985; Mower, 1991; Lee and Nedivi, 2002;

Nakadate et al., 2012). In apparent contradiction to this common view, we have reported that

spontaneous activity is sufficient for refinement of receptive fields in both visual cortex (V1) and superior

colliculus (SC) of dark reared (DR) Syrian hamsters, and early light exposure for 3-7 days is necessary

only to maintain the refinement in adulthood (Carrasco et al., 2005; Carrasco and Pallas, 2006; Balmer

and Pallas, 2015b; Mudd et al., 2019). Syrian hamster pups spend the first 3-4 weeks after birth

underground in the wild (Adler, 1948; Nowosielski-Slepowron and Park, 1987), so it would not be

beneficial to have the development of visual function depend on light exposure. If early visual experience

is unavailable, GABAergic lateral inhibition in SC and V1 declines and RFs expand, but not until about

two months of age (puberty) (Carrasco et al., 2005; Balmer and Pallas, 2015b; Mudd et al., 2019).

Pharmacological activation of TrkB receptors can mimic the effects of early light exposure in DR

hamsters, resulting in long-term maintenance of refined receptive fields and visual acuity (Mudd et al.,

2019). Thus, hamsters, unlike cats and monkeys, need visual experience to maintain refined receptive

fields in adulthood, but not to refine them during development **(Figure 1)**.

$GABA_A$ receptors are pentameric, ionotrophic receptors consisting of five subunits grouped around a

central chloride ion pore. The functional characteristics of the receptor largely depend upon the subunit

composition (Sigel et al., 1990) and organization (Minier and Sigel, 2004). Of the many subunit

arrangements, alpha1 and alpha2 subunits have been linked to synaptic localization of $GABA_A$ receptors.

However, these two subunit types have different kinetics and are expressed at different points in

development. At birth, receptors containing the alpha2 subunit are widely expressed throughout the brain,

whereas alpha1 expression is low in major areas of the brain like the neocortex, the hippocampus, and the

cerebellum (Laurie et al., 1992; Fritschy et al., 1994; Dunning et al., 1999; Chen et al., 2001). During the

first several postnatal weeks, alpha1 expression increases, coinciding with a reduction in alpha2

expression (Fritschy et al., 1994). This alpha2 to alpha1 switch in subunit expression underlies a

developmental decrease in inhibitory postsynaptic current (ipsc) decay time and an increase in ipsc

amplitude (Fritschy et al., 1994; Okada et al., 2000; Yu et al., 2006).

It was demonstrated previously that the expansion of RFs in SC of adult, DR hamsters is associated

with a loss of GABA immunoreactivity. Iontophoretic application of $GABA_A$ agonists *in vivo* restored

RFs to a normal adult size (Carrasco et al., 2011). In addition to a loss of GABA-immunoreactivity,

$GABA_A$ agonists and antagonists were less effective in SC and V1 neurons of DR hamsters than in

normally reared (NR) hamsters (Carrasco et al., 2011).

The incomplete development of RF properties in V1 of visually deprived cats has been associated

with a failure in developmental maturation of NMDA and $GABA_A$ receptors (Carmignoto and Vicini,

1992; Chen et al., 2000; Chen et al., 2001; Erisir and Harris, 2003). Although a failure to maintain

refined RFs is a different phenomenon than a failure to refine them during a critical period, the

mechanism(s) could be similar or convergent. Here, we provide molecular evidence complementing our

previous work (Carrasco et al., 2005; Carrasco and Pallas, 2006; Balmer and Pallas, 2015b) that suggests

the presence of divergent synaptic development mechanisms between hamsters and some other mammals

(Regal et al., 1976; Timney et al., 1978; Leventhal and Hirsch, 1980; Mower et al., 1981; Morales et al.,

2002; Viegi et al., 2002) in regulating RF development and maintenance under dark rearing conditions.

We thus tested the hypothesis that maintenance of refined RFs in adulthood depends on the stability of

mature receptors and other postsynaptic signaling components. This hypothesis predicts that the

detrimental, post-critical period receptive field plasticity observed in DR adults results from a

deprivation-induced failure to maintain these postsynaptic proteins in their mature state. Contrary to this

prediction, we find using Western blot assays of synaptosomes that the quantity, subunit composition, and

localization of $GABA_A$ receptors in SC of adult dark reared hamsters with re-expanded RFs resemble

those seen in normally reared subjects. Furthermore, levels of the synaptic scaffolding proteins gephyrin

and PSD-95 are normal, as are the adult expression levels of cation-chloride co-transporters

(KCC2/NKCC) in DR subjects. These findings suggest that, although a change in effectiveness of

$GABA_A$ receptors was reported previously using pharmacological agents (Carrasco et al., 2011; Balmer

and Pallas, 2015a), the loss of RF refinement in adulthood is mediated primarily by reductions in GABA

expression in the presynaptic terminals rather than by substantial postsynaptic alterations. This result is at

odds with the common view that presynaptic changes in the ligand must occur together with

corresponding postsynaptic changes in receptor levels (Fisher-Lavie and Ziv, 2013; Sudhof, 2018 ;

Sanderson et al., 2020). Taken together, our results rule out several hypotheses about the mechanistic

basis of maintenance of refined RFs throughout adulthood, and provide insights into regulation of critical

period plasticity that could lead to novel treatment targets for memory disorders and traumatic brain

injury.

Materials and Methods

Subjects

A total of 42 adult Syrian hamsters (Cricetidae: *Mesocricetus auratus*) of both sexes were bred within

our animal facility and used as subjects in this study. Syrian hamsters are an altricial rodent species that

is ideal for studying the developing visual system due to their robust and well-characterized visual

responses, short gestation time, and large litters (Chalupa, 1981; Huck et al., 1988; Pratt and Lisk, 1989;

Razak et al., 2003; Carrasco et al., 2005). Sexual maturity in this species occurs between postnatal day

(P) 56 and P60 (Diamond and Yanagimachi, 1970; Fitzgerald and Zucker, 1976). Breeding females were

singly housed. Male breeders were introduced and supervised until intromission was observed, after

which they were removed. Weanlings and adult males were housed in single sex social groups of up to 5

adults per cage and adult females were housed with female siblings or individually in standard rodent

cages. Running wheels were not provided because they have been shown to alter the timing of visual

cortical plasticity (Baroncelli et al., 2010; Tognini et al., 2012; Kalogeraki et al., 2014) but a variety of

other enrichment items were available. All subjects were provided with *ad libitum* access to rodent chow

and water.

Treatment groups

Animals used in this study were bred in-house to control sensory experiences throughout development.

Dams of DR subjects were transferred into total darkness 1-3 days before parturition. An antechamber

and light-impenetrable black curtain separated the dark housing room from the hallway, ensuring that any

accidental openings of the hallway doors did not expose the animals to light. Dark reared animals were

housed inside light-tight stackable cages with a standard HVAC filtration system consistent with the other

animal rooms in the facility. During general animal husbandry purposes, the hamsters were exposed to

dim red light at a wavelength not visible to Syrian hamsters (Huhman and Albers, 1994). NR hamsters

were maintained in a standard 12/12 light cycle room from before birth into adulthood.

Western blotting

Animals were euthanized with Euthasol at >150 mg/kg IP. Brains were immediately extracted and frozen

in 2-methylbutane on dry ice, then stored at -80°C or immediately dissected for preparation of lysates.

Protein extraction was done as described by Shi et al. (1997) with a few modifications. Briefly, individual

left and right superficial SC brain areas were excised and homogenized in a lysis buffer (10 mM

phosphate buffer, pH 7.0, 5 mM EGTA, 5 mM EDTA, 1 mM DTT) containing Halt protease inhibitor

(ThermoFisher Scientific). The lysate was centrifuged at 16,000 × g at 4°C for 10 min, and the pellet was

resuspended in 2 mM HEPES buffer, pH 7.2. It was then centrifuged at 70,000 rpm at 4°C for 30 min.

The pellet thus obtained was resuspended in 0.5 mM HEPES, pH 7.3, containing 0.32 M sucrose and

centrifuged at 450 × g for 8 min. Synaptic proteins were quantified using the Pierce BCA Protein Assay

Kit (ThermoFisher Scientific), mixed with 2X sample buffer, and boiled for 15 min at 60 °C. Twenty μg

of protein was loaded per well in pre-cast Bio-Rad gels and electrophoresis was carried out at 110 V for

90 min in a Bio-Rad electrophoresis tank. Proteins were then transferred onto nitrocellulose membranes at

70 V for 75 min, blocked in BSA for 1 h, and probed with primary antibodies overnight. Primary

antibodies used in this study included: rabbit anti-$GABA_AR\alpha1$ (1:1000, Cat#: AGA-001, Alomone Labs);

rabbit anti- $GABA_AR\alpha2$ (1:1000, Cat#: ab72445, Abcam); rabbit anti- $GABA_AR\alpha5$ (1:1000, Cat#:

ab10098, Abcam); rabbit anti-Gephyrin (1:1000, Cat#: ab32206, Abcam); mouse anti-PSD-95 (1:1000,

Cat#: ab2723, Abcam); mouse anti-KCC2 (1:1000, Cat#: 75-013, NeuroMab); rabbit anti-NKCC1

(1:1000, Cat#: ab59791, Abcam); and mouse anti-GAPDH (1:1000, Cat#: 600-GAPDH,

PhosphoSolutions). Protein bands were labeled using either appropriate IR dye fluorescent secondaries or

appropriate HRP-conjugated secondary antibodies and imaged on an Odyssey CLx fluorescent imaging

system (Li-Cor) or developed with enhanced chemiluminescent (ECL) substrate in a Bio-Rad ChemiDoc

Imager. Protein expression levels in the bands were quantified as the optical density of the labeled protein

normalized against the optical density of the housekeeping protein GAPDH using ImageJ (NIH).

Statistical Analysis

A Student's *t*-test was used to compare parametric data with equal variance between treatment groups and

a normally distributed control data set. In the case of non-parametric data that were not normally

distributed and/or exhibited unequal variance, a Mann-Whitney Rank Sum (U) test was employed.

Descriptive statistics for these analyses are provided as mean ± standard error of the mean (SEM) in the

text.

208 Results

The failure to maintain RF refinement in adult DR animals involves deficits in overall GABA

expression and $GABA_A$ receptor function (Carrasco et al., 2011), leading to a loss of lateral inhibition and

thus expansion of RFs after P60 (Carrasco et al., 2005; Balmer and Pallas, 2015b; Mudd et al., 2019). In

this study we explored several possible ways that visual deprivation during a critical period for RF

refinement could affect $GABA_A$ receptor efficacy and other postsynaptic proteins associated with

inhibitory synaptic function in adult SC. We used Western blotting on synaptosomes to study synaptic

proteins that can regulate synaptic plasticity but observed no significant changes in their relative

abundance under dark rearing conditions.

Dark rearing does not affect the subunit composition of $GABA_A$ receptors in adult SC

Deprivation-induced decreases in both $GABA_A$ and NMDA receptor levels in cat visual cortex have

been reported previously and were proposed to be involved in functional deficits (Carmignoto and Vicini,

1992; Chen et al., 2000; Chen et al., 2001). In DR hamsters, GAD and GABA immunoreactivity declines

and $GABA_A$ receptors are less efficient when tested pharmacologically (Carrasco et al., 2011), thus we

expected to see changes in levels of $GABA_A$ proteins postsynaptically. However, in our previous study

using Western blots on synaptosomes, no significant changes in the level of the $GABA_A$ receptor alpha1

subunit were observed (Mudd et al., 2019). These results raised the question of whether subunit

composition of GABA$_A$Rs might be altered by visual deprivation in a way that reduced their effectiveness

without affecting alpha1 levels.

We reasoned that the developmental alpha2 to alpha1 switch, if reversed in adulthood, could underlie

the reduction in GABA$_A$ receptor function that was previously observed in studies of RF enlargement in

adult SC (Carrasco et al., 2011). We explored this possibility by examining the expression of each subunit

in synaptosomes of SC in normally reared and visually deprived adults. Hamsters in the visual

deprivation group were dark reared from before birth. We used Western blotting to quantify the amount

of synaptic membrane-bound alpha1 and alpha2 GABA$_A$ receptor expression in the synaptosome

fractions obtained from adult SC. We found that there were no significant differences in either the overall

levels of alpha2 protein, observed as a ratio of alpha2 to GAPDH (NR: 1.21 ± 0.052, n=8; DR: $1.33 \pm$

0.185, n=8; U=26, n=8, p=0.574; Mann-Whitney Rank Sum Test) **(Figure 2A)**, or the ratio of

alpha1/alpha2 expression in the SC between adult DR (1.651 ± 0.277, n=5) and adult NR hamsters ($1.19\pm$

0.084, n=4) (T(7)= -1.436, n=4, p=0.194 Student's t-test) **(Figure 2B)**, a surprising result considering our

previous finding that dark rearing reduces the response to pharmacological application of GABA agonists

and antagonists (Carrasco et al., 2011). These findings argue against a reversion of the normal

developmental transition from alpha2 to alpha1-dominant expression as a cause of the deprivation-

induced RF enlargement in adult SC and support the interpretation that visual experience is not needed to

maintain mature GABA$_A$ receptor alpha subunit composition.

GABA$_A$ receptors can also be expressed extrasynaptically, where they can be activated by ambient

GABA derived from synaptic spillover or non-neuronal sources. This low concentration GABA source

generates "tonic" inhibition (Farrant and Nusser, 2005), and constitutes a significant (~75%) portion of

the total inhibitory forces acting on neurons in hippocampus (Mody and Pearce, 2004; Magnin et al.,

2019). Alpha5 subunit-containing receptors are primarily expressed extrasynaptically and have been

implicated in regulating the induction of synaptic plasticity for LTP in hippocampus (Saab et al., 2010;

Zurek et al., 2012; Zurek et al., 2014; Jacob, 2019). However, alpha5 GABA$_A$Rs have also been shown to

relocate to the synapse, colocalize with gephyrin (Brady and Jacob, 2015), and are critical for memory

consolidation (Davenport et al., 2021). To investigate the possible role of alpha5 levels in adult RF

maintenance we performed Western blotting on synaptosomes and compared between NR (0.544 units $\pm$

0.0520, n=8) and DR (0.471 $\pm$ 0.0935, n=8) adult hamsters **(Figure 3A)**. We found no significant

differences in alpha5 protein levels between groups (t (20) = 0.308, p = 0.878, Mann-Whitney Rank Sum

test. We also compared the ratio of alpha1/alpha5 between NR (1.027 $\pm$ 0.0815, n=10) and DR (0.995 $\pm$

0.0926, n =9) adult hamsters and found no differences between these groups (U (18) =35, p=0.438, Mann-

Whitney Rank Sum test) **(Figure 3B)**. Because we were only studying proteins from synaptosome

preparations (i.e., GABA$_A$ receptor subunits localized in the synapses), these results suggest that the

localization of alpha1 and alpha5 subunit-containing GABA$_A$ receptors in SC is not being altered by early

visual experience and argue against a deprivation-induced shift in tonic vs. synaptic inhibition (Davenport

et al., 2021).

Dark rearing does not affect the normal endocytosis of GABA$_A$ receptors in adult SC

The regulation of GABA$_A$ receptors at the synapse is pivotal for maintaining correct levels of

inhibitory synaptic transmission (Jacob et al., 2008). Impaired trafficking of GABA$_A$ receptors into and

out of synaptic membranes could affect their synaptic localization in SC and thus their overall response to

presynaptically released GABA. GABA$_A$ receptor trafficking is partially regulated by endocytosis- the

controlled removal of receptors from the postsynaptic membrane into the cytoplasm. Endocytosed

receptors are subsequently reinserted into the postsynaptic membrane or undergo lysosomal degradation

after longer periods (Kittler et al., 2000). We reasoned that if endocytosis was dysregulated, either by

decreased receptor reinsertion or increased receptor degradation, it could negatively impact the efficacy of

GABA$_A$ receptors at the synapse. We examined this possibility by comparing the ratio of postsynaptic

membrane-bound to cytosolic alpha1 subunit containing receptors between our treatment groups. No

differences were observed in the postsynaptic membrane/cytosol ratio of alpha1 expressing receptors

between DR (0.768 units $\pm$ 0.044, n=7) and NR (0.778 $\pm$ 0.1, n=6) adult hamsters (U(11)= 20.00,

p=0.945, Mann-Whitney Rank Sum test) **(Figure 4A)**. These results indicate that the overall trafficking of

alpha1 (synaptic) subunit- expressing GABA$_A$ receptors is not affected by dark rearing.

We also examined the possibility that extrasynaptic alpha5 receptor endocytosis-mediated trafficking

may be dysregulated and thus responsible for changes in tonic GABA$_A$ inhibition (Davenport et al.,

2021). There were no significant differences in cytosolic membrane localization of alpha5 subunits

between adult DR (1.320 ± 0.198, n=8) and NR groups (1.753 ± 0.449, n=6) (U(12)=19.00, p=0.573,

Mann-Whitney Rank Sum test) **(Figure 4B)**. These results indicate that the overall exo- and endocytosis-

mediated trafficking of alpha5 (extrasynaptic) subunit-expressing GABA$_A$ receptors is not responsible for

the decreased efficacy of GABA$_A$ receptors observed in RFs that fail to maintain refinement in adulthood

after dark rearing.

Inhibitory and excitatory scaffolding proteins in SC are not affected by dark rearing

One factor influencing the accumulation and retention of GABA$_A$ receptors at postsynaptic sites is the

membrane scaffolding protein gephyrin (Kneussel et al., 1999; Sun et al., 2004; Jacob et al., 2005; Tretter

et al., 2008). Decreased expression of gephyrin results in less clustering (Essrich et al., 1998) and more

overall mobility of GABA$_A$ receptors at the synapse (Jacob et al., 2005). We surmised that decreased

gephyrin expression could be responsible for the weaker GABA$_A$ receptor signaling we observed in

neurons with RFs that fail to maintain refinement in adulthood. We quantified and compared postsynaptic

membrane-bound gephyrin expression between DR and NR adults using Western blotting. Gephyrin

levels in DR adults (0.786 ± 0.124, n=17) were similar to those in NR adults (0.736 ± 0.158, n=16)

(U(111)=-0.247, p=0.377, Mann-Whitney Rank Sum test) **(Figure 5A)**. This indicates that maintenance

of adult gephyrin expression levels is not affected by dark rearing and suggests that if GABA$_A$ receptor

accumulation and trafficking is being affected, then it is occurring in a gephyrin-independent manner.

PSD-95 is the primary glutamate (AMPA and NMDA) receptor scaffolding protein in CNS neurons

(Chen et al., 2015), and it functions in a similar way as gephyrin does for GABA$_A$ receptors. Although it

does not have a direct impact on GABA$_A$ receptor function, PSD-95 has been shown to have an influence

on visual plasticity. For example, mice lacking PSD-95 have lifelong ocular dominance plasticity in V1

that results from an increase in the overall proportion of silent synapses, despite having normal inhibitory

tone (Funahashi et al., 2013; Huang et al., 2015). The possibility that the plasticity involved with the dark

rearing-induced re-enlargement of RFs could be mediated by a reduction in adult PSD-95 expression was

examined. We found that PSD-95 protein levels were not significantly different in DR (0.613± 0.96,

n=10) compared to NR adult hamsters (0.486 ± 0.868, n=9) (U(31)=-0.978, p=0.270, Mann-Whitney

Rank Sum test) **(Figure 5B)**. These results suggest that differences in PSD-95 levels do not underlie the

re-enlargement of RFs in SC following dark rearing from birth.

Cation-chloride co-transporters undergo their normal developmental switch in adult dark reared

subjects

Levels of inhibitory GABAergic signaling in neurons are dependent on the intracellular chloride (Cl^-)

concentration. The K^+ Cl^- co-transporter (KCC2) is responsible for regulating intracellular Cl^- in mature

adult neurons by an outward K^+ current (Rivera et al., 1999) and also regulates the formation, function,

and plasticity of glutamatergic synapses (Li et al., 2007; Gauvain et al., 2011; Chevy et al., 2015). At the

beginning of postnatal life $GABA_A$ receptor effects are excitatory because the Na^+-K^+-$2Cl^-$ co-transporter

1 (NKCC1) that mediates Cl^- uptake is dominant (Cherubini et al., 1991; Lee et al., 2005; Cancedda et

al., 2007). By the end of the second postnatal week after birth in rats and mice NKCC1 is replaced by

KCC2 as the dominant cation-chloride co-transporter in the brain, shifting $GABA_A$ receptors to produce

inhibitory PSPs (Rivera et al., 1999; Pfeffer et al., 2009; Moore et al., 2019). In V1, the developmental

switch from NKCC1 dominance to KCC2 dominance occurs at the same time as a period of BDNF/TrkB

mediated synaptic imbalance (Zhang et al., 2018). We surmised that a shift in the ratio of KCC2:NKCC1

could underlie the reinstating/restoring of plasticity for RF size refinement in dark reared adults, leading

to re-enlargement of RFs in SC. Examination of the expression of KCC2 and NKCC1 in adult SC

neurons revealed no significant differences between our treatment groups, however. KCC2 levels were

not significantly different between NR (0.979 ± 0.115, n=8) and DR groups of adult hamsters (0.963 ±

0.154, n=8) (T(14)=0.082, p=0.936, t-test) **(Figure 6A)**. The same was true of NKCC1 levels (NR 1.050

$\pm$ 0.0419, n=8; DR 1.081 $\pm$ 0.0814, n=8) (T(14)=-0.339, p=0.740, t-test) **(Figure 6B)**, and of the ratio of

the two cation-chloride co-transporters within groups (T(8)=1.096, p=0.305, t-test) **(Figure 6C)**.

Discussion

The goal of this study was to examine potential postsynaptic mechanisms through which light

exposure during an early critical period ensures the long-term stability of visual receptive fields in the

hamster superior colliculus. Although previous results established a correlation between the maintenance

of RF refinement and GABA immunoreactivity levels in SC (Carrasco et al., 2011; Mudd et al., 2019)

and V1 (Balmer and Pallas, 2015b), potential postsynaptic changes in protein levels had not been

examined. We have reported here that at the high resolution level of Western blots, visual deprivation-

induced failure to maintain refined RFs in SC does not appear to involve changes in GABA$_A$R subunit

composition, inhibitory or excitatory scaffolding protein expression, or cation-Chloride co-transporter

ratios. These results exclude several possible mechanisms that could explain the reduced activation of

GABA$_A$Rs with GABA agonists reported in DR adult SC (Carrasco et al., 2011), and support activity-

dependent regulation of GABA expression as the primary mechanism underlying TrkB-mediated

maintenance of RF refinement (Mudd et al., 2019). The finding that a change in GABA levels could

affect neural plasticity in adulthood has important implications for the treatment of memory impairments

or brain injury in adults.

This study supports our previous research that provided substantial evidence of a novel,

maladaptive adult plasticity in hamsters, in which visually deprived hamsters develop refined RFs

normally but fail to maintain them in adulthood. Interestingly, this contrasts with previous literature on

other mammals including cats (Cynader and Mitchell, 1980; Carmignoto and Vicini, 1992; Chen et al.,

2000; Chen et al., 2001; Mower, 1991; Mower et al., 1985; Mower et al., 1986; Erisir and Harris, 2003),

and rats (Fagiolini et al., 1994; Lee and Nedivi, 2002), in which the visual system does not develop

normally in response to dark rearing. Some of these previous studies looked only at early and/or adult

ages in the animals, thereby missing the RF refinement that happens in between the two ages. Because

diminished GABA release, contrary to what we expected, did not elicit changes in the levels of major

postsynaptic GABA$_A$ receptors, scaffold proteins, or chloride co-transporters, this study provides a

valuable demonstration that changes in neurotransmitter availability do not necessarily result in

coordinated changes in postsynaptic receptors.

Maturation of GABAergic signaling in visual cortex, particularly of the fast-spiking, parvalbumin-

containing basket cells, is thought to open and then close the critical period for plasticity (Fagiolini et al.,

2004; Sale et al., 2010; Toyoizumi et al., 2013; Capogna et al., 2021). Combined pre- and postsynaptic

alteration of synaptic strength has been seen in other sensory deprivation paradigms, including in dark

reared and monocularly deprived visual cortex, although with an earlier time course (Carmignoto and

Vicini, 1992; Chen et al., 2000; Chen et al., 2001). However, the retinorecipient layers of the superior

colliculus have no basket cells and contain very few GABAergic parvalbumin neurons, and the plasticity

described here occurs after the critical period should have closed, suggesting that SC may accomplish

plasticity through a different mechanism than visual cortex. On the other hand, previous studies found

that, as in visual cortex (Hanover et al., 1999; Huang et al., 1999; Viegi et al., 2002), deprivation-induced

receptive field plasticity in adult SC is mediated by the BDNF receptor TrkB (Mudd et al., 2019).

Furthermore, reduced GABA and GABA$_A$ receptor efficacy in response to iontophoretically-applied

GABA$_A$R agonists and antagonists is observed in both SC and V1 of visually deprived hamsters

(Carrasco et al., 2011; Balmer and Pallas, 2015b), arguing for mechanistic elements in common.

Early visual experience is not necessary for maturation or maintenance of GABA$_A$ receptor subunit

composition at the synapse

GABA$_A$ receptors contain fast acting chloride (Cl$^-$) channels (Pfeiffer et al., 1982; Sigel and

Steinmann, 2012). The subunit composition of GABA$_A$ receptors changes during development from an

alpha 2 to alpha 1 dominant condition (Laurie et al., 1992; Fritschy et al., 1994; Chen et al., 2001) and

also changes in some disease states (Levitt, 2005; Tyson and Anderson, 2014; Deidda et al., 2015;

Kimoto et al., 2015; Schmidt and Mirnics, 2015; Tang et al., 2021) in a way that affects receptor

functional properties (Farrant and Nusser, 2005) and localization (Jacob et al., 2005).   We studied

synaptic levels of the $GABA_A$ receptor alpha1 and alpha2 subunits to quantify their expression levels

under normal and DR conditions. The overall levels of $GABA_A$R alpha1 and $GABA_A$R alpha2 were not

altered in DR hamsters when compared to those of NR hamsters, arguing that the altered inhibitory

synaptic efficacy that we previously observed was not caused by an immature $GABA_A$R subunit

composition at the synapse.

An increase or decrease in the level of any receptor subunit is best understood in context, because

different conclusions would be drawn if subunits changed independently or in concert. Thus, we also

analyzed the alpha1/alpha2 ratios in individual animals. We did not find any change in alpha1/alpha2

ratios in NR compared to DR adult hamsters. These results suggest that early visual experience is not

necessary for maturation or maintenance of mature synaptic $GABA_A$R subunit composition in adulthood.

Thus, the failure to maintain refined RFs in adult DR hamsters cannot be explained by a return to a

juvenile type of $GABA_A$R subunit composition.

Level and localization of the extrasynaptic $GABA_A$R subunit alpha 5 does not change with visual

deprivation

$GABA_A$R subunit alpha 5 is predominantly an extrasynaptic membrane receptor subunit that regulates

tonic inhibition. It is important in neuronal circuit development, learning, and memory (Brady and Jacob,

2015), has been implicated in regulating the induction of synaptic plasticity in hippocampus (Saab et al.,

2010; Zurek et al., 2012; Zurek et al., 2014), and can relocate to the synaptic region in learning and

memory deficits (Brady and Jacob, 2015). Because the excitation/inhibition (E/I) balance could be

affected if alpha5 subunit levels changed or if they moved into the synapse, we compared its expression

between NR and DR cases. We did not see any significant changes in the overall levels of $GABA_A$R

alpha5 subunits, or in the alpha1/alpha5 ratio, suggesting that dark rearing-induced RF enlargement is not

caused by E/I imbalance via extrasynaptic receptor subunit changes.

Visual deprivation does not affect the rate of endocytosis of GABA$_A$R subunits at the synapse

Because clathrin-dependent endocytosis is likely important for regulating inhibitory signaling and

synaptic plasticity (Kittler et al., 2000), we examined the endocytosis of GABA$_A$ receptor alpha1 and

alpha5 subunits in normally reared and visually deprived subjects. We did not observe a significant

change in endocytosis of either subunit type as assayed by the ratio of synaptosome-bound to cytosolic

fractions. This implies that visual deprivation does not affect the trafficking of the GABA$_A$R subunits

between the synaptic membrane and the cytosol or the phosphorylation events that maintain the balance

between endocytosis and postsynaptic membrane insertion of the receptor subunits.

Early visual experience is not necessary to maintain scaffolding protein levels at the synapse

Another finding of this work is that the expression levels of the postsynaptic scaffold proteins PSD-

95 and gephyrin were not altered in adulthood following early visual deprivation, suggesting that any

changes in inhibitory function are probably not caused by a significant change in scaffolding protein

expression. Alternatively, the clustering of GABA$_A$Rs at inhibitory synapses in SC may happen in a

gephyrin-independent manner (Zita et al., 2007), or total gephyrin expression may not be as important as

the formation of gephyrin nanodomains within inhibitory synapses (Pennacchietti et al., 2017). Future

studies with different techniques would be required to determine if changes in receptor clustering may be

occurring and to what extent gephyrin or PSD-95 may have in mediating any such effects.

Early light exposure is not necessary for maturation of the cation-Chloride co-transporters

We investigated the status of the chloride transporters KCC2 and NKCC1 due to their role in

maintaining chloride balance inside the neurons and thus in maintaining the E/I balance. The cation-

chloride co-transporters could have reverted to their early developmental state, leading to disturbed E/I

balance and more excitation in dark reared animals, possibly explaining the RF expansion we observed.

However, we did not observe any changes in the cation-chloride co-transporters in dark-reared compared

to normally reared adult hamsters, suggesting that the changes in E/I balance due to lack of early visual

experience were not caused by alterations in the cation-chloride co-transporters.

A GABA-BDNF feedback loop maintains inhibitory networks, thereby maintaining RF refinement in

adulthood

GABA-GABA$_A$R interaction is known to regulate various downstream signaling pathways, and a

major regulator of GABA itself is BDNF-TrkB signaling triggered by NMDA receptor activity (Marini et

al., 1998). Our data are consistent with previous studies suggesting a positive feedback loop between the

BDNF-TrkB pathway and GABA expression, in which GABA facilitates BDNF expression, and BDNF

facilitates the synaptic release of GABA (Huang et al., 1999; Morales et al., 2002; Gianfranceschi et al.,

2003; Jovanovic et al., 2004; Kuczewski et al., 2008; Porcher et al., 2011; Hanno-Iijima et al., 2015),

maintaining RF size and visual acuity through GABAergic lateral inhibition (Mudd et al., 2019).

Signaling via the MAPK cascade and the transcription factor cAMP-responsive element-binding protein

(CREB) appears to play a substantial role in this process (Obrietan et al., 2002). BDNF-TrkB interaction

leads to dimerization and auto-phosphorylation of TrkB, thereby triggering MAPK, PLC gamma, and

PI3K pathways (Yoshii and Constantine-Paton, 2007). These pathways in turn lead to the activation of

downstream effectors and mediators to initiate a CREB-dependent transcription process that can lead to

an increase in GABA$_A$R levels as well as more BDNF production (Huang and Reichardt, 2003; Yoshii

and Constantine-Paton, 2007; Porcher et al., 2011; Esvald et al., 2020). In addition, an increase in the

transmembrane localization of GABA$_A$Rs is mediated by BDNF-dependent inhibition of receptor

endocytosis in addition to ongoing reinsertion of the receptor into the postsynaptic membrane (Porcher et

al., 2011). This positive feedback regulation is critical in developing neurons and hence constituted a

major motivation for the work reported here. In this study however, neither the GABA$_A$R subunit

composition at the synapse nor subunit composition in the extrasynaptic regions was affected by dark

rearing. Chloride transport proteins (KCC2, NKCC2) also remained normal in levels. One possibility is

that GABA expression levels alone are the key factor in RF re-enlargement (Carrasco et al., 2011; Mudd

et al., 2019), suggesting that this type of delayed plasticity resulting from a lack of early visual experience

occurs through a different mechanism than described in other types of plasticity resulting from visual

deprivation.

It is possible that changes in GABA$_A$R signaling occurred that are not reflected here in the expression

levels of postsynaptic receptors, scaffolding molecules, or ion transporters, or that we missed some

transient changes in GABA$_A$R signaling-associated proteins that cause GABA$_A$R functional changes. In

the future, it would be interesting if investigators used immunolabelling techniques to study protein

localization and interactions, or patch clamp experiments to study the properties of synaptic and

extrasynaptic responses in the SC as well as the subcellular dynamics of associated proteins involved.

This might improve the understanding of the molecular processes active in this deprivation-induced

maladaptive plasticity in the SC.

Suggested alternative explanations to understand this form of adult neural plasticity include that

the changes leading to RF expansion and thus the visual acuity deficits take place earlier than the time

point that we studied and return to normal by P90. Previous research has shown that the width of the

synaptic cleft decreases during development (Li and Cline, 2010). Thus, another interesting possibility is

that dark rearing from birth gradually increases the width of the synaptic cleft again in adulthood, while

maintaining the postsynaptic signaling components in place. It is also possible that other scaffolding

proteins and their partners are involved, for example gephyrin binding partners such as GABA$_A$R beta2

and beta3 subunits (Kowalczyk et al., 2013), the scaffold protein radixin that binds GABA$_A$R alpha5

subunit to the actin cytoskeleton (Loebrich et al., 2006) and regulates synaptic GABA$_A$R density (Hausrat

et al., 2015), or gephyrin post-translational modification events that influence inhibitory synaptic

plasticity by affecting postsynaptic scaffolding (Zacchi et al., 2014). More recently, distinct spectrin

isoforms have been shown to affect synaptic inhibition by selectively targeting specific GABA$_A$Rs

subtypes to dendritic and axon initial segment synapses (Smalley et al., 2023).

In summary, our results argue that visual experience is not necessary to maintain mature levels and

composition of several postsynaptic proteins that are essential for retino-SC synaptic communication.

Unlike many previous visual deprivation studies in which both GABA and GABA$_A$ receptors are

downregulated (Carmignoto and Vicini, 1992; Chen et al., 2000; Chen et al., 2001; Kilman et al., 2002;

Nahmani and Turrigiano, 2014), we report here that in SC GABA$_A$R levels, subunit composition, and

localization in adulthood are unaffected by dark rearing. The scaffold proteins gephyrin and PSD-95, and

the chloride transporters KCC2 and NKCC2 are also not affected by a lack of light exposure. The novel,

experience-dependent form of adult plasticity that we observed previously may involve an as yet

unidentified postsynaptic mechanism or solely a reduction in GABA release (Carrasco et al., 2005;

Carrasco and Pallas, 2006; Carrasco et al., 2011), thereby challenging the common argument that

presynaptic changes in ligand availability are always associated with matching postsynaptic changes in

their receptors. Either possibility is encouraging with respect to understanding this form of adult plasticity

and could help to improve treatment and therapeutic modalities in adults with memory impairments,

traumatic brain injury, or inhibition-associated neurological disorders.

REFERENCES

Adler S (1948) Origin of the Golden Hamster Cricetus auratus as a laboratory animal. Nature 162:256-
257.
Balmer TS, Pallas SL (2015a) Visual experience prevents dysregulation of GABA$_B$ receptor-dependent
short-term depression in adult superior colliculus. J Neurophysiol 113:2049-2061.
Balmer TS, Pallas SL (2015b) Refinement but not maintenance of visual receptive fields is independent
of visual experience. Cerebral Cortex 25:904-917.
Baroncelli L, Sale A, Viegi A, Maya Vetencourt JF, De Pasquale R, Baldini S, Maffei L (2010)
Experience-dependent reactivation of ocular dominance plasticity in the adult visual cortex. Exp
Neurol 226:100-109.
Brady ML, Jacob TC (2015) Synaptic localization of α5 GABA (A) receptors via gephyrin interaction
regulates dendritic outgrowth and spine maturation. Devel Neurobiol 75:1241-1251.
Cancedda L, Fiumelli H, Chen K, Poo M-m (2007) Excitatory GABA action is essential for
morphological maturation of cortical neurons *in vivo*. J Neurosci 27:5224-5235.
Capogna M, Castillo PE, Maffei A (2021) The ins and outs of inhibitory synaptic plasticity: Neuron
types, molecular mechanisms and functional roles. Eur J Neurosci 54:6882-6901.
Carmignoto G, Vicini S (1992) Activity-dependent decrease in NMDA receptor responses during
development of the visual cortex. Science 258:1007-1011.
Carrasco MM, Pallas SL (2006) Early visual experience prevents but cannot reverse deprivation-induced
loss of refinement in adult superior colliculus. Vis Neurosci 23:845-852.
Carrasco MM, Razak KA, Pallas SL (2005) Visual experience is necessary for maintenance but not
development of receptive fields in superior colliculus. J Neurophysiol 94:1962-1970.

Carrasco MM, Mao YT, Balmer TS, Pallas SL (2011) Inhibitory plasticity underlies visual deprivation-
induced loss of receptive field refinement in the adult superior colliculus. Eur J Neurosci 33:58-
68.
Chalupa LM (1981) Some observations on the functional organization of the golden hamster's visual
system. Behav Brain Res 3:189-200.
Chen L, Cooper NGF, Mower GD (2000) Developmental changes in the expression of NMDA receptor
subunits (NR1, NR2A, NR2B) in the cat visual cortex and the effects of dark rearing. Molec
Brain Res 78:196-200.
Chen L, Yang C, Mower GD (2001) Developmental changes in the expression of GABA(A) receptor
subunits (alpha(1), alpha(2), alpha(3)) in the cat visual cortex and the effects of dark rearing.
Molec Brain Res 88:135-143.
Chen X, Levy JM, Hou A, Winters C, Azzam R, Sousa AA, Leapman RD, Nicoll RA, Reese TS (2015)
PSD-95 family MAGUKs are essential for anchoring AMPA and NMDA receptor complexes at
the postsynaptic density. Proc Natl Acad Sci U S A 112:E6983-E6992.
Cherubini E, Gaiarsa JL, Ben-Ari Y (1991) GABA: An excitatory transmitter in early postnatal life.
Trends in Neurosciences 14:515-519.
Chevy Q, Heubl M, Goutierre M, Backer S, Moutkine I, Eugène E, Bloch-Gallego E, Lévi S, Poncer JC
(2015) KCC2 gates activity-driven AMPA receptor traffic through cofilin phosphorylation. J
Neurosci 35:15772.
Cynader M, Mitchell DE (1980) Prolonged sensitivity to monocular deprivation in dark-reared cats.
Journal of Neurophysiology 43:1026-1040.
Davenport CM, Rajappa R, Katchan L, Taylor CR, Tsai MC, Smith CM, de Jong JW, Arnold DB,
Lammel S, Kramer RH (2021) Relocation of an extrasynaptic GABA(A) receptor to inhibitory
synapses freezes excitatory synaptic strength and preserves memory. Neuron 109:123-134.
Deidda G, Allegra M, Cerri C, Naskar S, Bony G, Zunino G, Bozzi Y, Caleo M, Cancedda L (2015) Early
depolarizing GABA controls critical-period plasticity in the rat visual cortex. Nat Neurosci 18:87-
96.
Diamond M, Yanagimachi R (1970) Reproductive Development in the Female Golden Hamster in
Relation to Spontaneous Estrus. Biology of Reproduction 2:223-229.
Dunning DD, Hoover CL, Soltesz I, Smith MA, O'Dowd DK (1999) GABAA Receptor–mediated
miniature postsynaptic currents and α-subunit expression in developing cortical neurons. J
Neurophysiol 82:3286-3297.
Erisir A, Harris JL (2003) Decline of the critical period of visual plasticity is concurrent with the
reduction of NR2B subunit of the synaptic NMDA receptor in layer 4. J Neurosci 23:5208-5218.
Esvald EE, Tuvikene J, Sirp A, Patil S, Bramham CR, Timmusk T (2020) CREB family transcription
factors are major mediators of BDNF transcriptional autoregulation in cortical neurons. J
Neurosci 40:1405-1426.
Fagiolini M, Pizzorusso T, Berardi N, Domenici L, Maffei L (1994) Functional postnatal development of
the rat primary visual cortex and the role of visual experience: Dark rearing and monocular
deprivation. Vision Res 34:709-720.
Fagiolini M, Fritschy JM, Low K, Mohler H, Rudolph U, Hensch TK (2004) Specific GABA$_A$ circuits for
visual cortical plasticity. Science 303:1681-1683.
Farrant M, Nusser Z (2005) Variations on an inhibitory theme: phasic and tonic activation of GABAA
receptors. Nat Rev Neurosci 6:215-229.
Firth SI, Wang CT, Feller MB (2005) Retinal waves: mechanisms and function in visual system
development. Cell Calcium 37:425-432.
Fisher-Lavie A, Ziv NE (2013) Matching dynamics of presynaptic and postsynaptic scaffolds. J Neurosci
33:13094-13100.
Fitzgerald K, Zucker I (1976) Circadian organization of the estrous cycle of the golden hamster. Proc Natl
Acad Sci USA 73:2923-2927.

Fritschy J-M, Paysan J, Enna A, Mohler H (1994) Switch in the expression of rat GABA$_A$-receptor
571   subtypes during postnatal development: An immunohistochemical study. J Neurosci 14:5302-
572   5324.
573   Funahashi R, Maruyama T, Yoshimura Y, Komatsu Y (2013) Silent synapses persist into adulthood in
574   layer 2/3 pyramidal neurons of visual cortex in dark-reared mice. J Neurophysiol 109:2064-2076.
575   Gauvain G, Chamma I, Chevy Q, Cabezas C, Irinopoulou T, Bodrug N, Carnaud M, Lévi S, Poncer JC
576   (2011) The neuronal K-Cl cotransporter KCC2 influences postsynaptic AMPA receptor content
577   and lateral diffusion in dendritic spines. Proc Natl Acad Sci U S A 108:15474-15479.
578   Gianfranceschi L, Siciliano R, Walls J, Morales B, Kirkwood A, Huang ZJ, Tonegawa S, Maffei L (2003)
579   Visual cortex is rescued from the effects of dark rearing by overexpression of BDNF. Proc Natl
580   Acad Sci U S A 100:12486-12491.
581   Hanno-Iijima Y, Tanaka M, Iijima T (2015) Activity-dependent bidirectional regulation of GAD
582   expression in a homeostatic fashion Is mediated by BDNF-dependent and independent pathways.
583   PloS one 10:e0134296.
584   Hanover JL, Huang ZJ, Tonegawa S, Stryker MP (1999) Brain-derived neurotrophic factor
585   overexpression induces precocious critical period in mouse visual cortex. J Neurosci 19:RC40
586   (41-45).
587   Hausrat TJ, Muhia M, Gerrow K, Thomas P, Hirdes W, Tsukita S, Heisler FF, Herich L, Dubroqua S,
588   Breiden P, Feldon J, Schwarz JR, Yee BK, Smart TG, Triller A, Kneussel M (2015) Radixin
589   regulates synaptic GABAA receptor density and is essential for reversal learning and short-term
590   memory. Nat Commun 6:6872.
591   Hensch TK, Quinlan EM (2018) Critical periods in amblyopia. Vis Neurosci 35:E014.
592   Huang EJ, Reichardt LF (2003) Trk receptors: roles in neuronal signal transduction. Annu Rev Biochem
593   72:609-642.
594   Huang X, Stodieck SK, Goetze B, Cui L, Wong MH, Wenzel C, Hosang L, Dong Y, Lowel S, Schluter
595   OM (2015) Progressive maturation of silent synapses governs the duration of a critical period.
596   Proc Natl Acad Sci USA 112:E3131-E3140.
597   Huang ZJ, Kirkwood A, Pizzorusso T, Porciatti V, Morales B, Bear MF, Maffei L, Tonegawa S (1999)
598   BDNF regulates the maturation of inhibition and the critical period of plasticity in mouse visual
599   cortex. Cell 98:739-755.
600   Hubel DH, Wiesel TN (1970) The period of susceptibility of the physiological effects of unilateral eye
601   closure in kittens. J Physiol (Lond) 206:419-436.
602   Hübener M, Bonhoeffer T (2014) Neuronal Plasticity: Beyond the Critical Period. Cell 159:727-737.
603   Huck UW, Lisk RD, McKay MV (1988) Social dominance and reproductive success in pregnant and
604   lactating golden hamsters (Mesocricetus auratus) under seminatural conditions. Physiol Behav
605   44:313-319.
606   Huhman KL, Albers HE (1994) Neuropeptide Y microinjected into the suprachiasmatic region phase
607   shifts circadian rhythms in constant darkness. Peptides 15:1475-1478.
608   Jacob TC (2019) Neurobiology and Therapeutic Potential of α5-GABA Type A Receptors. Front Molec
609   Neurosci 12:179.
610   Jacob TC, Moss SJ, Jurd R (2008) GABAA receptor trafficking and its role in the dynamic modulation of
611   neuronal inhibition. Nat Rev Neurosci 9:331-343.
612   Jacob TC, Bogdanov YD, Magnus C, Saliba RS, Kittler JT, Haydon PG, Moss SJ (2005) Gephyrin
613   regulates the cell surface dynamics of synaptic GABAA receptors. J Neurosci 25:10469-10478.
614   Jovanovic JN, Thomas P, Kittler JT, Smart TG, Moss SJ (2004) Brain-Derived Neurotrophic Factor
615   modulates fast synaptic inhibition by regulating GABAA receptor phosphorylation, activity, and
616   cell-surface stability. J Neurosci 24:522-530.
617   Kalogeraki E, Greifzu F, Haack F, Lowel S (2014) Voluntary physical exercise promotes ocular
618   dominance plasticity in adult mouse primary visual cortex. J Neurosci 34:15476-15481.
619   Katz LC, Shatz CJ (1996) Synaptic activity and the construction of cortical circuits. Science 274:1133-
620   1138.

Kilman V, van Rossum MC, Turrigiano GG (2002) Activity deprivation reduces miniature IPSC amplitude by decreasing the number of postsynaptic GABA(A) receptors clustered at neocortical synapses. J Neurosci 22:1328-1337.

Kimoto S, Zaki MM, Bazmi HH, Lewis DA (2015) Altered markers of cortical gamma-Aminobutyric Acid neuronal activity in schizophrenia: Role of the NARP gene. JAMA psychiatry 72:747-756.

Kittler JT, Delmas P, Jovanovic JN, Brown DA, Smart TG, Moss SJ (2000) Constitutive endocytosis of GABAA receptors by an association with the Adaptin AP2 complex modulates inhibitory synaptic currents in hippocampal neurons. J Neurosci 20:7972-7977.

Kneussel M, Brandstätter JH, Laube B, Stahl S, Müller U, Betz H (1999) Loss of postsynaptic GABAA receptor clustering in gephyrin-deficient mice. J Neurosci 19:9289-9297.

Kowalczyk S, Winkelmann A, Smolinsky B, Förstera B, Neundorf I, Schwarz G, Meier JC (2013) Direct binding of GABAA receptor β2 and β3 subunits to gephyrin. Eur J Neurosci 37:544-554.

Kuczewski N, Langlois A, Fiorentino H, Bonnet S, Marissal T, Diabira D, Ferrand N, Porcher C, Gaiarsa JL (2008) Spontaneous glutamatergic activity induces a BDNF-dependent potentiation of GABAergic synapses in the newborn rat hippocampus. J Physiol 586:5119-5128.

Laurie DJ, Wisden W, Seeburg PH (1992) The distribution of thirteen GABA$_A$ receptor subunit mRNAs in the rat brain. III. Embryonic and postnatal development. J Neurosci 12:4151-4172.

Lee H, Chen CX, Liu YJ, Aizenman E, Kandler K (2005) KCC2 expression in immature rat cortical neurons is sufficient to switch the polarity of GABA responses. Eur J Neurosci 21:2593-2599.

Lee WC, Nedivi E (2002) Extended plasticity of visual cortex in dark-reared animals may result from prolonged expression of cpg15-like genes. J Neurosci 22:1807-1815.

Leventhal AG, Hirsch HV (1980) Receptive-field properties of different classes of neurons in visual cortex of normal and dark-reared cats. J Neurophysiol 43:1111-1132.

Levitt P (2005) Disruption of interneuron development. Epilepsia 46:22-28.

Li H, Khirug S, Cai C, Ludwig A, Blaesse P, Kolikova J, Afzalov R, Coleman SK, Lauri S, Airaksinen MS, Keinänen K, Khiroug L, Saarma M, Kaila K, Rivera C (2007) KCC2 interacts with the dendritic cytoskeleton to promote spine development. Neuron 56:1019-1033.

Li J, Cline HT (2010) Visual deprivation increases accumulation of dense core vesicles in developing optic tectal synapses in Xenopus laevis. J Comp Neurol 518:2365-2381.

Loebrich S, Bähring R, Katsuno T, Tsukita S, Kneussel M (2006) Activated radixin is essential for GABAA receptor α5 subunit anchoring at the actin cytoskeleton. EMBO J 25:987-999.

Maffei L, Galli-Resta L (1990) Correlation in the discharge of neighboring rat retinal ganglion cells during prenatal life. Proc Natl Acad Sci USA 87:2861-2864.

Magnin E, Francavilla R, Amalyan S, Gervais E, David LS, Luo X, Topolnik L (2019) Input-specific synaptic location and function of the alpha5 GABAA receptor subunit in the mouse CA1 hippocampal neurons. J Neurosci 39:788-801.

Marini AM, Rabin SJ, Lipsky RH, Mocchetti I (1998) Activity-dependent release of Brain-derived Neurotrophic Factor underlies the neuroprotective effect of N-Methyl-D-aspartate. J Biol Chem 273:29394-29399.

Meister M, Wong RO, Baylor DA, Shatz CJ (1991) Synchronous bursts of action potentials in ganglion cells of the developing mammalian retina. Science 252:939-943.

Minier F, Sigel E (2004) Positioning of the alpha-subunit isoforms confers a functional signature to gamma-aminobutyric acid type A receptors. Proc Natl Acad Sci U S A 101:7769-7774.

Mody I, Pearce RA (2004) Diversity of inhibitory neurotransmission through GABAA receptors. Trends in Neurosciences 27:569-575.

Moore YE, Conway LC, Wobst HJ, Brandon NJ, Deeb TZ, Moss SJ (2019) Developmental Regulation of KCC2 Phosphorylation Has Long-Term Impacts on Cognitive Function. Front Mol Neurosci 12:173.

Morales B, Choi S-Y, Kirkwood A (2002) Dark rearing alters the development of GABAergic transmission in visual cortex. J Neurosci 22:8084-8090.

Mower GD (1991) The effect of dark rearing on the time course of the critical period in cat visual cortex. Dev Brain Res 58:151-158.

Mower GD, Berry D, Burchfiel JL, Duffy FH (1981) Comparison of the effects of dark rearing and binocular suture on development and plasticity of cat visual cortex. Brain Res 220:255-267.

Mower GD, Caplan CJ, Christen WG, Duffy FH (1985) Dark rearing prolongs physiological but not anatomical plasticity of the cat visual cortex. J Comp Neurol 235:448-466.

Mudd DB, Balmer TS, Kim SY, Machhour N, Pallas SL (2019) TrkB activation during a critical period mimics the protective effects of early visual experience on perception and the stability of receptive fields in adult superior colliculus. J Neurosci 39:4475-4488.

Nahmani M, Turrigiano GG (2014) Deprivation-induced strengthening of presynaptic and postsynaptic inhibitory transmission in layer 4 of visual cortex during the critical period. J Neurosci 34:2571-2582.

Nakadate K, Imamura K, Watanabe Y (2012) Effects of monocular deprivation on the spatial pattern of visually induced expression of c-Fos protein. Neuroscience 202:17-28.

Nowosielski-Slepowron B, Park A (1987) Variation in the growth of the preweaning Syrian hamster (Cricetus auratus). Acta Morphol Neerl Scand 25:187-199.

Obrietan K, Gao X-B, van den Pol AN (2002) Excitatory actions of GABA Increase BDNF expression via a MAPK-CREB–dependent mechanism—A positive feedback circuit in developing neurons. J Neurophysiol 88:1005-1015.

Okada M, Onodera K, Van Renterghem C, Sieghart W, Takahashi T (2000) Functional correlation of GABAA receptor α subunits expression with the properties of IPSCs in the developing thalamus. J Neurosci 20:2202-2208.

Pallas SL (2017) The impact of ecological niche on adaptive flexibility of sensory circuitry. Frontiers in neuroscience 11:344.

Pennacchietti F, Vascon S, Nieus T, Rosillo C, Das S, Tyagarajan SK, Diaspro A, Del Bue A, Petrini EM, Barberis A, Cella Zanacchi F (2017) Nanoscale molecular reorganization of the inhibitory postsynaptic density Is a determinant of GABAergic synaptic potentiation. J Neurosci 37:1747-1756.

Pfeffer CK, Stein V, Keating DJ, Maier H, Rinke I, Rudhard Y, Hentschke M, Rune GM, Jentsch TJ, Hübner CA (2009) NKCC1-Dependent GABAergic Excitation Drives Synaptic Network Maturation during Early Hippocampal Development. J Neurosci 29:3419-3430.

Pfeiffer F, Graham D, Betz H (1982) Purification by affinity chromatography of the glycine receptor of rat spinal cord. J Biol Chem 257:9389-9393.

Porcher C, Hatchett C, Longbottom RE, McAinch K, Sihra TS, Moss SJ, Thomson AM, Jovanovic JN (2011) Positive feedback regulation between {gamma}-Aminobutyric Acid Type A (GABAA) receptor signaling and Brain-derived Neurotrophic Factor (BDNF) release in developing neurons. J Biol Chem 286:21667-21677.

Pratt NC, Lisk RD (1989) Effects of social stress during early pregnancy on litter size and sex ratio in the golden hamster (Mesocricetus auratus). J Reprod Fertil 87:763-769,.

Razak KA, Huang L, Pallas SL (2003) NMDA receptor blockade in the superior colliculus increases receptive field size without altering velocity and size tuning. J Neurophysiol 90:110-119.

Regal DM, Boothe R, Teller DY, Sackett GP (1976) Visual acuity and visual responsiveness in dark-reared monkeys (Macaca nemestrina). Vision Res 16:523-530.

Reh RK, Dias BG, Nelson CA, 3rd, Kaufer D, Werker JF, Kolb B, Levine JD, Hensch TK (2020) Critical period regulation across multiple timescales. Proc Natl Acad Sci U S A.

Ribic A (2020) Stability in the Face of Change: Lifelong Experience-Dependent Plasticity in the Sensory Cortex. Frontiers in Cellular Neuroscience 14.

Rivera C, Voipio J, Payne JA, Ruusuvuori E, Lahtinen H, Lamsa K, Pirvola U, Saarma M, Kaila K (1999) The K+/Cl− co-transporter KCC2 renders GABA hyperpolarizing during neuronal maturation. Nature 397:251-255.

Saab BJ, MacLean AJ, Kanisek MM, Zurek AA, Martin LJ, Roder JC, Orser BA (2010) Short-term memory impairment after isoflurane in mice is prevented by the α5 γ-aminobutyric acid Type A receptor inverse agonist L-655,708. Anesthesiol 113:1061-1071.

Sale A, Berardi N, Spolidoro M, Baroncelli L, Maffei L (2010) GABAergic inhibition in visual cortical plasticity. Frontiers in Cellular Neuroscience 4.

Sanderson TM, Georgiou J, Collingridge GL (2020) Illuminating Relationships Between the Pre- and Post-synapse. Frontiers in neural circuits 14:9.

Schmidt MJ, Mirnics K (2015) Neurodevelopment, GABA system dysfunction, and schizophrenia. Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology 40:190-206.

Shi J, Aamodt SM, Constantine-Paton M (1997) Temporal correlations between functional and molecular changes in NMDA receptors and GABA neurotransmission in the superior colliculus. J Neurosci 17:6264-6276.

Sigel E, Steinmann ME (2012) Structure, function, and modulation of GABA(A) receptors. J Biol Chem 287:40224-40231.

Sigel E, Baur R, Trube G, Möhler H, Malherbe P (1990) The effect of subunit composition of rat brain GABAA receptors on channel function. Neuron 5:703-711.

Smalley JL, Cho N, Ng SFJ, Choi C, Lemons AHS, Chaudry S, Bope CE, Dengler JS, Zhang C, Rasband MN, Davies PA, Moss SJ (2023) Spectrin-beta 2 facilitates the selective accumulation of GABAA receptors at somatodendritic synapses. Communications Biology 6:11.

Sudhof TC (2018) Towards an Understanding of Synapse Formation. Neuron 100:276-293.

Sun C, Sieghart W, Kapur J (2004) Distribution of alpha1, alpha4, gamma2, and delta subunits of GABAA receptors in hippocampal granule cells. Brain research 1029:207-216.

Takesian AE, Hensch TK (2013) Balancing plasticity/stability across brain development. Prog Brain Res 207:3-34.

Tang X, Jaenisch R, Sur M (2021) The role of GABAergic signalling in neurodevelopmental disorders. Nature Rev Neurosci 22:290-307.

Timney B, Mitchell DE, Giffin F (1978) The development of vision in cats after extended periods of dark-rearing. Experimental Brain Research 31:547-560.

Tognini P, Manno I, Bonaccorsi J, Cenni MC, Sale A, Maffei L (2012) Environmental enrichment promotes plasticity and visual acuity recovery in adult monocular amblyopic rats. PloS one 7:e34815.

Toyoizumi T, Miyamoto H, Yazaki-Sugiyama Y, Atapour N, Hensch Takao K, Miller Kenneth D (2013) A theory of the transition to critical period plasticity: Inhibition selectively suppresses spontaneous activity. Neuron 80:51-63.

Tretter V, Jacob TC, Mukherjee J, Fritschy JM, Pangalos MN, Moss SJ (2008) The clustering of GABA(A) receptor subtypes at inhibitory synapses is facilitated via the direct binding of receptor alpha 2 subunits to gephyrin. J Neurosci 28:1356-1365.

Tyson JA, Anderson SA (2014) GABAergic interneuron transplants to study development and treat disease. Trends Neurosci 37:169-177.

Viegi A, Cotrufo T, Berardi N, Mascia L, Maffei L (2002) Effects of dark rearing on phosphorylation of neurotrophin Trk receptors. Eur J Neurosci 16:1925-1930.

Wong ROL, Meister M, Shatz CJ (1993) Transient period of correlated bursting activity during development of the mammalian retina. Neuron 11:923-938.

Yoshii A, Constantine-Paton M (2007) BDNF induces transport of PSD-95 to dendrites through PI3K-AKT signaling after NMDA receptor activation. Nature Neurosci 10:702-711.

Yu Z-Y, Wang W, Fritschy J-M, Witte OW, Redecker C (2006) Changes in neocortical and hippocampal GABAA receptor subunit distribution during brain maturation and aging. Brain Research 1099:73-81.

Zacchi P, Antonelli R, Cherubini E (2014) Gephyrin phosphorylation in the functional organization and plasticity of GABAergic synapses. Front Cell Neurosci 8:103.

Zhang H, Mu L, Wang D, Xia D, Salmon A, Liu Q, Wong-Riley MTT (2018) Uncovering a critical
period of synaptic imbalance during postnatal development of the rat visual cortex: role of brain-
derived neurotrophic factor. J Physiol 596:4511-4536.
Zita MM, Marchionni I, Bottos E, Righi M, Del Sal G, Cherubini E, Zacchi P (2007) Post-
phosphorylation prolyl isomerisation of gephyrin represents a mechanism to modulate glycine
receptors function. EMBO J 26:1761-1771.
Zurek AA, Bridgwater EM, Orser BA (2012) Inhibition of alpha5 gamma-Aminobutyric acid type A
receptors restores recognition memory after general anesthesia. Anesth Analg 114:845-855.
Zurek AA, Yu J, Wang D-S, Haffey SC, Bridgwater EM, Penna A, Lecker I, Lei G, Chang T, Salter
EWR, Orser BA (2014) Sustained increase in $\alpha 5$ GABAA receptor function impairs memory
after anesthesia. J Clin Invest 124:5437-5441.

Figure 1: Differences in the visual refinement in the hamster SC depending on exposure to light

during the critical period of heightened neural plasticity. Hamsters exposed to light during their critical

period gradually improve RF refinement in the SC and maintain it throughout life (as indicated by the blue

line). Hamsters that do not experience light during the critical period also show RF refinement in the SC by

P60, but the refinement progressively declines in adulthood (as indicated by the orange line)

(Carrasco et al., 2005; Carrasco and Pallas, 2006; Balmer and Pallas, 2015b).

A.

B.

Figure 2. Adult GABA$_A$α2 receptor subunit composition is not affected by early dark rearing. (A)

image: Individual Western blots of normally reared (NR) and dark-reared (DR) treatment groups

generated using 20 μg of SC protein per lane. GAPDH was used as a loading control. graph:

Densitometric analyses of GABA$_A$Rα2 in normally vs. dark reared hamsters. **(B)** Comparison of

GABA$_A$Rα2 expression with GABAA$_A$Rα1 expression, shown as individual Western blots in the upper

panel and as a ratio in the lower panel. Data presented as mean ± SEM. In this and the following figures,

lanes presented together are from the same gel(s), and each measured protein was normalized against

GAPDH as a loading control. These results suggest that the number and ratio of synaptic GABA$_A$α2

receptors are similar in normal and dark reared adult SC.

Figure 3. Adult GABA$_A$α5 receptor subunit composition is not affected by early dark rearing. (A

and B top: Example Western blots of NR and DR treatment groups as in Figure 2. All presented lanes are

from the same gel(s), and each gel was run with GAPDH as a loading control. **(A graph)** GABA$_A$Rα5

comparison. **(B graph)** GABA$_A$Rα1: GABA$_A$Rα5 ratio. Data presented as mean ± SEM normalized

against GAPDH levels. These results suggest that the number and ratio of synaptic GABA$_A$α5 receptors

are similar in normal and dark reared adult SC.

A.

B.

Figure 4. Endocytosis of GABA$_A$ receptors is not affected by early dark rearing. **(A)** Adult levels of

the cytosolic vs. synaptic membrane-attached ratio of the extrasynaptically-targeted GABA$_A$Rα1 and **(B)**

the synaptically-targeted GABA$_A$Rα5 subunits were not affected by early light deprivation. Example

Western blots (upper panels) represent bands of labeled intracellular and membrane bound receptor

subunit proteins (upper bands) from the same animal, measured as a ratio against GAPDH and compared

between NR and DR groups. GAPDH was used as a loading control. Graph data presented as normalized

mean optical density ± SEM.

831 **Figure 5. Adult gephyrin and PSD-95 expression are not affected by dark rearing.** (A) Gephyrin

832 and (B) PSD-95 expression was similar between adult (>P90) NR and DR groups. GAPDH and β-actin

833 were used as loading controls. Graph data are presented as normalized mean relative optical density ±

834 SEM.

Figure 6. Adult cation-chloride co-transporters expression is not affected by early dark rearing.

Western blot examples of NR and DR samples labeled for cation-chloride co-transporters **(A)** KCC2 (140

840   kDa) and **(B)** NKCC1 (150 kDa), **(C)** and a comparison of the within subject ratio of KCC2:NKCC1 in

841   NR and DR adult (>P90) hamsters. Graph data presented as mean relative optical density ± SEM.