Dopamine signaling regulates predator-driven changes in *Caenorhabditis elegans'*

egg laying behavior

Amy Pribadi[1,2ϒ], Michael A. Rieger[2ϒ], Kaila Rosales[2], Kirthi C. Reddy[2] and Sreekanth H.

Chalasani[1,2,*]

[1]Biological Sciences Graduate Program, University of California, San Diego, La Jolla,

CA, USA.

[2]Molecular Neurobiology Laboratory, The Salk Institute for Biological Studies, La Jolla,

CA, USA.

[ϒ]Equal author contribution.

* Address for correspondence: schalasani@salk.edu (S.H.C.).

Abstract

Prey respond to predators by altering their behavior to optimize their own fitness and

survival. Specifically, prey are known to avoid predator-occupied territories to reduce

their risk of harm or injury to themselves and their progeny. We probe the interactions

between *Caenorhabditis elegans* and its naturally cohabiting predator *Pristionchus*

uniformis to reveal the pathways driving changes in prey behavior. While *C. elegans*

prefers to lay its eggs on a bacteria food lawn, the presence of a predator inside a lawn

induces *C. elegans* to lays more eggs away from that lawn. We confirm that this change

in egg laying is in response to bites from predators, rather than to predatory secretions.

Moreover, predator-exposed prey continue to lay their eggs away from the dense lawn

even after the predator is removed, indicating a form of learning. Next, we find that

mutants in dopamine synthesis significantly reduce egg laying behavior off the lawn in

both predator-free and predator-inhabited lawns, which we can rescue by transgenic

complementation or supplementation with exogenous dopamine. Moreover, we find that

dopamine is likely released from multiple dopaminergic neurons and requires

combinations of both D1- (DOP-1), and D2-like (DOP-2 and DOP-3) dopamine

receptors to alter predator-induced egg laying behavior, whereas other combinations

modify baseline levels of egg laying behavior. Together, we show that dopamine

signaling can alter both predator-free and predator-induced foraging strategies, implying

a role for this pathway in defensive behaviors.

Keywords

Predator-prey, *C. elegans*, *P. uniformis*, egg laying, dopamine.

Introduction

Predator-prey systems offer a rich variety of prey behaviors to explore, from

innate to learned responses. Prey responses to predators also vary depending on the

predation strategy [1,2], as well as the prey's abilities and the environmental context of

both species [3]. Additionally, prey can evaluate the cost/benefit of engaging in these

antipredator behaviors, since they might impose additional costs by reducing access to

food or mates [4]. While predators kill and consume prey, they can also influence prey

behavior without necessarily inflicting direct harm, in both wild and laboratory contexts

[5,6]. However, these changes in prey behavior often involve costs like reduced access

to food or mates [6]. For example, reintroducing wolves into Yellowstone National Park

resulted in changes to the grazing patterns of female elks with calves, with more time

devoted to vigilance behaviors [7,8]. In the laboratory setting, rats presented with cat

odor spent more time in shelter than exploring, feeding, or mating [9,10]. Laboratory

experiments in model organisms can lack the natural context of predator-prey dynamics,

but observation in the wild lacks the ability to link predator-prey behaviors to molecules

and neural pathways. To bridge the gap between ecological relevance and mechanistic

insight, we explored a predator-prey system in nematodes that brings a naturalistic

predator-prey interaction into the laboratory, making it more amenable to controlled

experimentation.

Caenorhabditis elegans is a nematode that lives in rotting vegetation and eats

the bacteria found there [11]. With 302 neurons and a mapped connectome [12], it is a

model well-suited to study behavior with the manipulation of genes and circuits often at

the resolution of a single cell. Much research in predator-prey relationships involve

organisms that possess vision, but less is understood about analogous interspecies

interactions where prey response is entirely dependent on olfaction and

mechanosensation, such as *C. elegans*. While much research in predator-prey

relationships involve organisms that have vision, little is known about defensive

behaviors in olfactory/mechanosensory-dependent organisms like *C. elegans*. With

different dependencies on sensory modalities, *C. elegans*-specific behaviors may not

necessarily mimic defensive behavior traditionally associated with sighted prey, such as

freezing [13,14]. *C. elegans* spends most of its time searching for food or eating it, as

well as laying eggs, so predator threat may influence these activities. The motor

sequences required for changes to navigation when searching for food, such as the

frequency of turns and reversals, are subject to the integration of input from several

sensory neurons, and their modulation by biogenic amine neurotransmitter signaling

[15,16]. Although non-predative, there are numerous examples of *C. elegans* altering

this system of navigational decision making in response to encounters with potentially

aversive stimuli. For example, *C. elegans* will learn to avoid pathogenic bacteria such *S.*

marcescens, a behavior mediated by serotonin signaling [17]. *C. elegans* will also sense

and navigate away from certain metal ions such as Cu^{2+}, and neurons mediating Cu^{2+}

response are modulated in turn by octopamine and serotonin [18]. This response is also

enhanced by the presence of food which is mediated by dopaminergic signaling [19].

Dopaminergic signaling also impacts how an animal locomotes in response to

mechanically aversive stimuli such as the touch response, which itself is again modified

by the presence of food [20]. These same neurotransmitters also impact the decision of

when and where to lay eggs. Exogenous serotonin is known to promote the rate of egg

laying off food, meanwhile exogenous octopamine and tyramine can inhibit this behavior

[21]. Dopaminergic signaling couples locomotor behavior and egg laying, promoting the

rate of egg laying when animals are roaming [22]. Like other potentially aversive stimuli,

predator responses may be expected to modify how an animal navigates its

environment. Like these stimuli, predator-evoked changes to exploration would likely

intersect with the availability of food, potentially impacting activities like egg laying, all of

which is expected to be modulated by biogenic amine neurotransmitter and receptor

signaling pathways.

Previous studies have shown that nematodes of the *Pristionchus* genus can

predate on other nematodes like *C. elegans* [23] and are found in necromenic

association with beetles [24,25] as well as in rotting vegetation along with

Caenorhabditis [26,27]. Members of the *Pristionchus* genus exhibit mouth polyphenism,

with either two-toothed "eurystomatous" (Eu) or one-toothed "stenostomatous"(St)

mouthforms [28,29]. The Eu mouthform in *P. pacificus* has been shown to enable more

successful killing of nematode prey like *C. elegans* [30,31]. While *P. pacificus* is a

relatively well-studied species within *Pristionchus,* it is uncertain whether *C. elegans*

actually interacts with *P. pacificus* in nature. In contrast, the gonochoristic species *P.*

uniformis has been found in the same sample with wild *C. elegans* isolates [26], thus *P.*

uniformis may represent a likelier candidate for naturalistic predative antagonism to *C.*

elegans. Although *P. uniformis* was first characterized as a St-only species [32], it has

recently been re-assessed and found to possess both a bacterivorous St and the

predatory Eu mouthform [33], and we too find that in standard growth conditions most *P.*

uniformis strain JU1051 individuals have an Eu mouthform (**Fig. 1-figure supplement**

1).

To test the hypothesis that, like other aversive stimuli, predators were able to

exert an influence on patterns of *C. elegans* exploration, we wondered whether we

could observe changes to *C. elegans* position and egg laying relative to food when

animals experienced predator threat, and how factors like predator presence and

bacterial topology intersect. As navigation and egg laying are influenced by biogenic

amine signaling, we also wondered whether we could then use the molecular tools

developed in *C. elegans* to discover the mechanisms underlying any observed changes

to behavior. In this study, we show that *C. elegans* avoids a bacterial lawn that is

occupied by its naturally cohabiting predator *P. uniformis* [26], and lays its eggs away

from that lawn. We find that predator-exposed *C. elegans* potentiates the probability of

egg laying off of the lawn, and this effect is sustained for many hours even after the

predator is removed. This potentiation is further exaggerated when food is present

outside the main bacterial lawn. Furthermore, we find that *C. elegans* egg laying

locations are regulated by biogenic amine signaling in both baseline and predator-

exposed conditions. Complete loss of dopamine synthesis resulted in overall reductions

to egg laying at off lawn locations, which was restored by supplementation with

exogenous dopamine. However, loss of signaling through combinations of D1 and D2

receptor homologs was able to perturb predator-induced off lawn egg laying behavior

while maintaining baseline levels. Taken together we present a framework for

interrogating prey behavior in nematodes, define some of the dynamics of this behavior,

and identify potential molecular regulators of egg laying under predator threat.

Results

C. elegans **avoids bacterial lawns inhabited by** *Pristionchus* **predators.**

We recently showed that *P. pacificus* bites *C. elegans* adults even though it is difficult to

consume them. This biting of adult *C. elegans* prey forces these animals to leave the

bacterial lawn, resulting in more exclusive access to the lawn by the predator [34].

Using a modified version of the protocol in our previous study [34], we placed three

predators and three *C. elegans* on an assay plate containing a small, dense bacterial

lawn. Animals were restricted to an arena that included the lawn and a small area of

empty agar (see **Methods**). Control plates ("Mock") had six *C. elegans* to maintain a

consistent number of animals between plates with and without predators. These

behavioral arenas were imaged under various experimental conditions, and coordinates

of the eggs in arenas were determined. These coordinates were used to compute the

distances of individual eggs from lawn center as well as their position relative to the

lawn edge. Since *Pristionchus* also lay eggs, we used a *C. elegans* strain that

expresses the GFP fluorophore in all of its eggs (*Pelt-2*::GFP) (**Fig. 1a**).

To observe whether predator biting affects *C. elegans* prey behavior, we chose

several different types of predators: *P. pacificus* strains PS312 and RS5194, a

stenostomatous-only *P. pacificus* mutant TU445 *eud-1(tu445)* [35], and an isolate of *P.*

uniformis, JU1051. *P. pacificus* strain RS5194 is more aggressive than PS312 as

characterized by an increased probability of bite per encounter [34] so both strains were

included in this analysis. The stenostomatous-only (non-predative) mutant was included

to demonstrate whether biting was required for predators to alter *C. elegans* behavior.

We also included the cohabiting predator *P. uniformis*. As a more naturalistic predator

which has coevolved with *C. elegans*, we wondered how prey response to this predator

may differ from *P. pacificus. P. uniformis* males and females were considered

separately, while only hermaphrodite *P. pacificus* were used. We first tested if short-

term predator exposure could alter where eggs were laid by determining the numbers of

eggs on and off bacterial lawns in our experimental arenas. These tabulations allowed

us to fit a logistic regression model (**Methods**, **Equations 1-2**) that estimated the

probability of off lawn egg laying ("P(off)") as a function of time and in interaction with

various predators or other conditions. To prevent eggs hatching into L1s, which secrete

pheromones that promote lawn-leaving [36] this assay only ran for six hours. *C. elegans*

in general showed an increase to P(off) over time regardless of predator condition

although animals exposed to the aggressive strain *P. pacificus* RS5194 showed slightly

higher P(off) at 6 hours compared to Mock (*C. elegans* only 0.22, RS5194 exposed

animals 0.29, p=0.043) (**Figure 1-figure supplement 2**). We also observed an increase

to P(off) between 3-5 hours when exposed to *P. uniformis* females but by 6 hours P(off)

in both Mock control and *P. uniformis* female exposed conditions appeared comparable.

We hypothesized that increasing predator exposure time would more greatly

increase the probability of off lawn egg laying in predator-exposed animals. We

conducted a long-term assay with L4 *C. elegans* and J4 *Pristionchus* instead of adults

and stopped the assay after 20 hours of exposure. Juveniles develop into adulthood

over the course of the assay, (*C. elegans* starts laying eggs approximately 8-10 hours

after the L4 stage [37]). Thus, as eggs were laid primarily in the latter portion of the 20-

hour time period, this limited L1 hatching during the assay. Arenas with *P. pacificus*

eud-1 mutants showed similar P(off) compared to Mock (*C. elegans* only) exposed

animals, while all other *Pristionchus* predators showed pronounced increases to the

probability of off lawn egg laying (**Figure 1b-c**). These data indicate that interactions

between *eud-1* mutants and prey (secretions, contacts, and others) are unable to alter

the locations of *C. elegans* eggs. We confirmed that this change was primarily due to

altered egg laying location and not overall changes to the number of eggs (no significant

change in egg numbers after predator exposure, **Figure 1-figure supplement 3**). While

P. uniformis males triggered a similar proportion of *C. elegans* eggs to be laid off lawn

(P(off)=0.72) compared to both strains of *P. pacificus* (RS5194 0.74, PS312 0.73), *P.*

uniformis females had an intermediate effect (P(off) = 0.62). Taken together, these

experiments show that *C. elegans* change their location of egg laying away from a lawn

occupied by primarily eurystomatous *Pristionchus* capable of biting.

Next, we tested whether *Pristionchus* biting-induced injury was required for the

change in *C. elegans* egg location. We used a *C. elegans* reporter strain expressing

GFP (green fluorescent protein) under the control of a *nlp-29* promoter. This strain

(*Pnlp-29*::GFP) has been shown to increase GFP expression upon wounding the cuticle

with a microinjection needle, a laser beam, or fungal infection [38,39]. We paired each

of the predators tested in our egg location assay with this reporter strain and monitored

GFP fluorescence relative to the co-injection marker (*Pcol-12*::*dsRED*) (**Fig. 1c**). We

found that both isolates of *P. pacificus* (PS312 and RS5194) were able to increase

reporter fluorescence in this reporter strain within four hours (**Figure 1-figure**

supplement 4). In the 20-hour assay, *C. elegans* exposed to *P. pacificus* RS5194 were

killed and could not be measured, but animals exposed to *P. pacificus* PS312 adults

showed increased reporter fluorescence (**Figure 1d**). In contrast, the stenostomatus

eud-1 mutant was unable to increase GFP fluorescence even after 20 hours. Notably,

neither *P. uniformis* males nor females were able to increase GFP fluorescence in this

Pnlp-29::GFP reporter strain. However, while it is difficult to confirm biting when the

bites are relatively ineffective, *C. elegans* do appear to sense putative bites from *P.*

uniformis by exhibiting escape response typical of other aversive stimuli [34,40]

(**Supplementary Video 1**). It is possible that these bites are causing low level of harm

without damaging the cuticle sufficiently to increase expression from the *Pnlp-29*::GFP

reporter. We planned to use the *C. elegans* egg location assay for the remainder of our

studies in non-fluorescent wildtype *C. elegans* and so chose a predator that does not

lay eggs (*P. uniformis* males) in our assays (**Figure 1e**). Furthermore, failure to elicit a

change in *Pnlp-29*::GFP fluorescence also indicated that changes to P(off) when

exposed to *P. uniformis* animals in our egg location assay was not the result of

extensive injury.

We next tested how the ratio of predators (*P. uniformis* males) to prey (*C.*

elegans) affected the location of prey eggs and the expected value of P(off). While

maintaining the same arena size and total number of animals (six), we altered the ratio

of predators and prey. We found that the presence of even a single predator was able to

increase the P(off) and adding additional predators resulted in greater increase to P(off),

appearing to asymptote after ≥2 predators in the arena (**Figure 1-figure supplement**

5a). These changes to predator:prey ratio did not alter the overall abundance of *C.*

elegans eggs (**Figure 1-figure supplement 5b**). These data are consistent with results

in our previous study using *P. pacificus* [34].

As exposure to *P. uniformis* males did not result in strong injury to *C. elegans* but

nevertheless was associated with changes to off lawn egg laying, we wondered whether,

rather than biting itself, this phenomenon was due to compounds secreted by the

predator. We have previously shown that *P. pacificus* secretions are aversive to *C.*

elegans [41]. We tested whether *P. uniformis* was secreting an aversive chemical that

drives *C. elegans* away from the bacterial lawn. We conditioned lawns with *P. uniformis*

males or sterile *C. elegans* as a control (to simulate changes in lawn caused by animal

movement) and tested whether naïve *C. elegans* would alter their egg location on these

lawns. We were unable to detect a shift in P(off) as the result of exposure to *P.*

uniformis-conditioned lawns (**Figure 1-figure supplement 6a**). We did detect, curiously,

an increase to the overall number of *C. elegans* eggs, though this was likely driven by

an outlier effect (**Figure 1-figure supplement 6b**). These data suggest that *P. uniformis*

males likely do not secrete a *C. elegans* aversive signal that can account for the

observed predator-induced change to egg location.

Predator-induced changes to off lawn laying are associated with sustained

avoidance of the lawn by prey.

While *C. elegans* exhibits increased P(off) when occupying a lawn with predators, it may

be that *C. elegans* is not truly avoiding the lawn in general, but simply altering its

decision about where to lay its eggs. To determine where the prey themselves were

located throughout the course of a predator exposure experiment, we used an imaging

setup (WormWatcher) to monitor the locations of mScarlet-expressing *C. elegans* over

20 hours with images of acquired every 4 minutes (**Figure 2a**). We found that, when

exposed to *P. uniformis*, *C. elegans* exhibited a shift in location to just outside the lawn

boundary, starting at approximately 5-6 hours (**Figure 2b**). This shift in location was

sustained in predator-containing arenas through the remainder of the 20-hour assay,

while Mock controls exposed only to other *C. elegans* remained mainly within the lawn.

Thus, we infer that changes to P(off) observed in our egg location assays is likely a

consequence of this sustained avoidance.

Change in bacterial topography alone contributes to, but does not account for

extent of egg location change

We observed that arenas containing *C. elegans* hermaphrodites and *P. uniformis* males,

but not controls, had streaks of bacteria outside the main lawn (**Figure 1e**). Given the

duration of our assay, these streaks might represent bacteria that sticks on the *C.*

elegans body and gets deposited onto the agar as it exits the lawn. Over the duration of

the assay, these streaks grow and are visible to the naked eye by the end of the 20-

hour period. We tested whether the presence of streaks outside the main lawn alone

could account for the change in egg location. We artificially streaked bacterial lawns at

the beginning of the assay and monitored the location of the eggs in these predator-free

arenas (**Figure 3b**). Indeed, artificial streaking was able to induce an increase in P(off)

nearly three-fold, however this response was greater in arenas containing *P. uniformis*

(**Figure 3c**). These data showed that the presence of bacteria outside the main lawn

can drive egg location change but may not be the only contributor to the decision of

where to lay eggs when exposed to predator.

Egg location change lasts many hours even in the absence of predator.

Next, we tested whether changes to egg location persist even in the absence of

predators. We "trained" *C. elegans* prey in our egg location assay setup with *P.*

uniformis males for 20 hours and transferred only the prey to a test arena. The position

of eggs laid in the test arena was quantified over 6 hours and subjected to the same

analyses as our other egg location assays, allowing us to test more nuanced

hypotheses about the effect of recent exposure to predator.

We tested animals recently exposed to *P. uniformis* or Mock (*C. elegans* only)

controls in three environments: completely filled arenas, normal small (~1.5 mm radius)

lawn arenas, and arenas with artificial streaks as in Figure 3 (**Figure 4a**). In a

completely filled arena, there is no region which can be labeled as "off lawn", and so

P(off) cannot be quantified. Instead, we were able to use this arena to estimate

predator-induced changes to overall distributional properties of eggs. We looked at the

average distance from center eggs were laid over 6 hours (**Figure 4b**) as an estimate of

the prey's central tendencies, as well as the coefficient of variation of the egg

distribution (**Figure 4c**) which estimates changes to the width of the egg distribution that

may have been brought about by recent predator exposure. We were unable to detect

significant differences due to predator exposure, though we did detect a significant main

effect of time on each metric. The average distance of eggs from center decreased over

the course of 6 hours, while the coefficient of variation of these distributions tended to

increase (**Figure 4c**). The estimated slopes for these effects over time are shown in

Figure 4-figure supplement 1.

When we tested artificial streaking (**Figure 3**), results suggested both effects of

changes to bacterial topology and an interaction with the presence of *P. uniformis* males.

We were curious about dynamics of this interaction in the absence of predator. We

tested *C. elegans* recently exposed to predator or non-exposed controls in our learning

paradigm in arenas containing either a small main lawn or a lawn with artificial streaks,

and determined the number of eggs laid at 1-6 hours in independent arenas. We found

a significant three-way interaction between time, recent predator exposure, and

bacterial topology (**Figure 4d**). Animals exposed only to other *C. elegans* and then

subsequently laying eggs in test arenas with small unperturbed lawns tended to have

very low values of P(off) in general, which decayed negatively over time. By contrast,

when tested in arenas with artificial streaks, not only was P(off) increased generally, but

showed a positive relationship with off lawn laying increasing over time. When exposed

to *P. uniformis* males and tested in arenas with unperturbed lawns, as expected animals

did show a potentiation of P(off) and this potentiation to P(off) was comparable to that

exhibited by *C. elegans* in the artificially streaked arenas at the early time points.

However, in contrast to the temporal dynamics shown by changes to bacterial topology,

P(off) was flatter with recent predator exposure across all time points. Finally, combining

recent predator exposure and testing on lawns with artificial streaks showed the

greatest potentiation to P(off) in general, with a similarly flat response over time. These

results suggest that for at least 6 hours there are two separate phenomena: egg laying

off the lawn driven by the presence of low concentrations of bacteria at a distance from

the main lawn, and egg laying off the lawn driven by recent predator exposure. Predator

exposure and artificial streaks together exhibit a combined effect on potentiating P(off)

overall which is more than additive. With respect to the time evolution over 6 hours,

recent predator exposure appears to trump the effects of bacterial topology, indicated

by the relatively flat slopes in predator-exposed animals in either bacterial topological

condition. The estimated slopes for these effects over time are shown in **Figure 4-**

figure supplement 2.

We wondered whether this elevation to P(off) would persist at even greater

periods of time away from predator exposure. We transferred 20-hour exposed *C.*

elegans to a rest plate completely filled with food for 1, 2, or 24 hours (**Figure 5a**). We

then quantified eggs laid on a test plate containing artificial streaks, as it appeared that

artificial streaking of the bacteria was likely to bring about the greatest potentiation of

predator-induced changes to P(off). Consistent with the positive slope conditions

observed over 6 hours in artificially streaked test arenas (**Figure 4d**), we saw a

significant elevation of the baseline level of P(off) at 24 hours in the Mock control

condition where animals were not exposed to predator (**Figure 5b**). Predator-exposed

animals showed elevation to P(off) at all time points including 24 hours, with a flatter

relationship over time. This indicates that *C. elegans* are able to "remember" their

recent predator experience for at least 24 hours. However, it is also clear that the

baseline probability of off lawn egg laying increases by 24 hours regardless of predator

exposure, as exhibited in the Mock condition. Thus, we computed changes to the fold

change between Predator and Mock observed at each time point, and defined this as

the Predator Response (see **Methods**, **Equation 3**). This difference of differences

captures the overall magnitude of observed shifts in egg laying behavior associated with

the presence of predator . Although our data are not paired, the generalized linear

modeling frameworks allows us to compute estimated confidence intervals on this fold

change for performing statistical inference. We found that this response was

significantly lower at 24 hours than at 1 or 2 hours, as a result of the increase in the

baseline P(off) in the Mock condition (**Figure 5c**). This indicates that while P(off)

remains elevated, *C. elegans* may be beginning to extinguish its "memory" of recent

exposure by 24 hours.

Biogenic amine signaling regulates off lawn laying behavior.

Biogenic amines already are well established as modulators of egg laying behavior in general as well as egg laying during different locomotor modes [21,22,42]. Additionally, biogenic amines are known to modulate behaviors over long time scales [43], and the change in egg location behavior upon predator exposure appears to last for many hours. We hypothesized that egg laying behavior in response to predator might be subject to modulation by biogenic amines, and therefore tested mutants in genes required for their synthesis. We observed variable changes both to the baseline P(off) probabilities in animals not exposed to predator, as well as to the magnitude of predator exposure response (**Figure 6a-b**). This is consistent with previous studies showing that dopamine and serotonin signaling is required for overall locomotion [44,45]. In general, all mutants were able to show potentiation in P(off) when exposed to predator (**Figure 6a**). Mutants in the *C. elegans* homolog of the mammalian vesicular monoamine transporter (VMAT), *cat-1* [46] exhibited a lower P(off) in the absence of predators. Although *cat-1* mutant animals showed an increase to P(off) with predator exposure, the magnitude of predator response (as in **Figure 5c**) was lower than WT animals (**Figure 6b**). We also found that mutants in *cat-2* (which encodes tyrosine hydroxylase for dopamine synthesis [47,48]) and *tph-1* (tryptophan hydroxylase for serotonin synthesis [49]) had similar changes to baseline off lawn egg laying, but nevertheless increase P(off) in the presence of predator. This magnitude of increase was greater than WT in *tph-1* mutants given the very low baseline P(off) in these animals in Mock conditions, and the increase in *cat-2*(*e1112*) was similar in fold change magnitude compared to WT, again given their low baseline of P(off) in non-exposed conditions. Mutants in *tbh-1* (which encodes tyramine beta-hydroxylase which converts tyramine into octopamine [21]) showed a similar baseline of P(off) to WT animals and a greater potentiation with predator. Mutants in *tdc-1* (tyrosine decarboxylase, which converts tyrosine into tyramine), showed an elevation of P(off) in Mock controls and a slight decrease in fold potentiation in the presence of predator compared to WT animals. Tyramine is known to inhibit egg laying [21], however we did not detect significant changes to the number of eggs laid per *C. elegans* animals in *tdc-1* mutants (not shown). Taken together, these data show loss of

biogenic amine neurotransmitters can modify off lawn egg laying behavior, attenuating

or even increasing the observed response to predator, though these two phenomena

were not so clearly separable. Loss of both dopamine and serotonin neurotransmitters

in *cat-1* mutants, however, not only reduced the general probability of off lawn laying but

also contributed to the largest blunting of the predator response. We focused our

remaining studies on dopaminergic signaling, but future work will investigate the role of

serotonin signaling as serotonin has been previously shown to modify egg laying

behavior [50,51].

We continued to investigate the consequence of loss of dopamine synthesis by

testing a second mutant allele of *cat-2*, *n4547*. Both *cat-2* mutants showed a similar

reduction to baseline P(off) in Mock conditions (**Figure 6c**), and a similar magnitude of

Predator Response (**Figure 6d**). In *C. elegans* adult hermaphrodites, CAT-2 protein is

expressed by eight neurons (four CEPs, two ADEs and PDEs), and dopamine signaling

has been previously shown to affect modulation of locomotion as well as learning [43].

Additionally, analysis of the dopamine transporter promoter has identified specific

elements that drive expression of transgenes in subsets of these dopaminergic neurons

[52]. Using these cell-selective promoter elements, we expressed full-length coding

sequence of the *cat-2* cDNA under either CEP- or ADE/PDE- specific promoters.

Transgenic rescue in ADE/PDE partially restored baseline P(off) in Mock controls

(**Figure 6e**), with rescue in CEP neurons resulting in the greatest increase to baseline

P(off), even greater than WT levels. In this particular experiment, *cat-2*(*e1112*) mutants

did in fact show a blunted Predator Response even though this metric accounts for the

reduced levels of baseline P(off) in the Mock condition, and both rescues also show

significantly lower Predator Response compared to WT (**Figure 6f**). This indicates some

variability in absolute loss of dopamine synthesis on modulating predator response vs.

modulating off lawn egg laying in general. The cohorts of *cat-2* mutants used in **Figure**

6a-b, **Figure 6c-d,** as well as the results shown in **Figure 7** described below, indicate

that changes to the underlying probability of laying eggs off the lawn is likely driving any

observed effects to predator response. Additionally, differences in promoter strength

used to drive expression of *cat-2* may explain why dopaminergic cell types show

differing ability to restore baseline P(off). Nevertheless, it is clear that re-expression of

CAT-2 protein in either ADE/PDE or CEP only is sufficient to at least partially restore

baseline off lawn egg laying behavior.

We also monitored locomotor activity of *cat-2*(*e1112*) animals over the course of

20 hours using the WormWatcher imager. Mutants were still capable of elevating

distance from center upon predator exposure. However, there were approximately 40%

fewer timepoints at which mutants differed between Mock and predator-exposed

conditions as compared to controls (**Figure 6-figure supplement 1a-b**). When

computing confidence intervals for the fold increase (change between Predator and

Mock conditions), both *cat-2*(*e1112*) mutants and WT exhibited similar response,

though *cat-2* mutants did show lower magnitudes of change at a few time points (**Figure**

6-figure supplement 1c). A mutant in the dopamine reuptake transporter gene *dat-1*,

which has increased amounts of dopamine at synapses [53,54], showed a nearly

identical response to WT animals (**Figure 6-figure supplement 1d-f**). Towards the end

of the 20-hour observation period, however, *dat-1* mutants in the mock condition began

to move away from the lawn, consistent with the role of excess dopamine in altering

locomotion [16,43,55]. These results suggest dopamine signaling is required for off lawn

exploration and changes in this pathway likely affects both animal position and egg

laying distribution.

Next, we hypothesized that adding exogenous dopamine would restore normal

egg laying to *cat-2* dopamine deficient mutants. To test our hypothesis, we first exposed

wildtype and *cat-2* mutant *C. elegans* to *P. uniformis* males for 20 hours (training) and

then transferred them to a plate with a lawn and artificial streaks (as in **Figure 5-6**) with

and without exogenous dopamine (**Figure 7a**). This assay setup avoids exogenous

dopamine from altering *P. uniformis* behavior, and leverages our data that prey

responses persist for 24 hours even without predators. Previously 2 mM exogenous

dopamine has been shown to rescue basal slowing upon encountering food [44] and

density pattern discrimination of PDMS pillars [56] in *cat-2* mutants. Consistent with our

previous results, *cat-2* mutants exhibited reduced off lawn egg laying in both control and

predator-exposed conditions (**Figure 7b**). We found that adding 2 mM dopamine

restored normal off lawn egg laying in both of these conditions. In the case of *cat-2*

mutants, addition of exogenous dopamine restored baseline P(off) to significantly

greater levels than in WT, and thus exhibited a net reduction the predator response

(**Figure 7c**). Together, these data suggest that dopamine signaling is required for off

lawn egg laying in both control and predator-exposed conditions.

Dopamine receptor signaling alters both baseline and predator-evoked egg laying

behavior.

Complete loss of dopamine synthesis appeared to primarily affect baseline levels of egg

laying activity off the bacterial lawn, so we next explored the roles of the cognate

dopamine receptors in modifying this behavior. The *C. elegans* genome encodes at

least four dopamine receptors (*dop-1*, *-2*, *-3*, and *-4*) with viable mutants in each [43].

These receptors each have multiple protein isoforms whose sequence alignments are

depicted in **Figure 8a**. *C. elegans* DOP-1 is a homolog of the mammalian D1-like

receptors and DOP-2/3 are homologs of mammalian D2-like receptors [43]. DOP-4 is

also D1-like, however this receptor belongs to a unique invertebrate family of D1-like

including receptors found in *Drosophila melanogaster* and *Apis mellifera* [57]. We

tested single mutants in each of these four receptors in our egg location assay along

with a quadruple mutant that lacked all four receptors. P(off) was increased with

exposure to predator in all cases (**Figure 8b**). Complete loss of all four receptors was

associated with a trend to reduce the baseline of P(off) in Mock controls (p=0.08 after

multiple testing correction) and did not show a significant change to the predator

response compared to WT (**Figure 8c**), which were similar effects observed when

removing dopamine synthesis. Loss of individual receptors had varying results. Loss of

dop-1, *dop-2*, *dop-3* all elevated baseline P(off) to varying degrees (**Figure 8b**) and

showed concomitant reductions to the magnitude of predator-induced fold increases

(**Figure 8c**). Thus, loss of single receptors, though able to modulate overall fold change

in P(off) when predator was present, still appeared to do so as a consequence of

changes to background. Only *dop-4* single mutants show Mock condition P(off) not

significantly distinct from WT and also showed comparable predator-evoked response.

Since dopamine receptors are known to exist as heteromers [58], we analyzed

mutants in pairwise combinations. Again, all combinations of two *dop-* mutants showed

an elevation of P(off) when predator was present (**Figure 9a**). These combinations also
had differing effects on baseline P(off) in Mock controls. *dop-1;dop-4* mutants were the
most similar to WT. *dop-1;dop-2*, *dop-2;dop-4* and *dop-2;dop-3* all showed elevation of
baseline off lawn egg laying activity relative to WT, and *dop-1;dop-3* and *dop-3;dop-4*
showed reductions to baseline P(off). The magnitude of Predator Response in these
mutant combinations is shown ordered from highest to lowest in **Figure 9b**. WT and
dop-3;dop-4 double mutants show the highest fold change increase in P(off) relative to
their respective Mock controls. All combinations containing *dop-4* rank intermediate with
dop-2;dop-3 and *dop-1;dop-3* ranking lowest. Other than *dop-3;dop-4*, all other
combinations showed reduction to predator response relative to WT.

The ranked magnitudes of fold change in predator exposed conditions suggested
combinations with *dop-4* mutants were intermediate or closer to WT response level,
regardless of changes to baseline P(off). To test the hypothesis of the presence or
absence of just *dop-4* influencing predator-evoked behavior, we performed an
experiment comparing triple mutant animals in *dop-1;dop-2;dop-3* to quadruple mutants
of all four receptors (**Figure 9c**). Once again, the quadruple mutant showed reduction to
the baseline P(off) in the Mock control as in **Figure 8**, however, the triple mutant
showed a comparable level of off lawn laying in the mock condition relative to WT. This
enabled us to more easily interpret the significant reduction to Predator Response
observed when comparing the triple mutant to the quadruple mutant or WT (**Figure 9f**).
Taken together, dopaminergic receptor signaling can affect both baseline off lawn laying
activity as well as predator response, and the specific exclusion of *dop-1*, *dop-2*, and
dop-3 from the assembly of available receptors modulates predator response while
maintaining otherwise normal levels of egg laying activity.

Discussion

In this study, we show that *C. elegans* responds to its predator *P. uniformis* by changing
egg laying location relative to a shared food patch. When given the option to find lower
density bacterial streaks off of the main lawn, *C. elegans* shift to laying more eggs off
the lawn basally consistent with a boost in exploratory behavior when alternate food
sources are present (**Figure 3,4**). When exposed to predator, *C. elegans* is even more

likely to lay its eggs off the lawn (**Figure 3**) when these new food options are available
and this effect is greater than either exposure to predator or the presence of these
bacterial streaks alone, and persists even in the absence of predator for many hours
(**Figure 4,5**).

We show that basally and in predator-exposed contexts, a shift to laying eggs off
the lawn is modulated by biogenic amine signaling. Biogenic amines like dopamine and
serotonin have been previously shown to play a role in driving responses to predator
threat in honey bees [59], ants [60], and fruit flies [61]. Consistently, we find that loss of
dopamine synthesis modulates baseline *C. elegans* egg laying which is consistent with
changes to locomotion observed in these mutants [43]. This behavior is rescued by
transgenic re-expression of *cat-2* in CEP or ADE/PDE neurons or with the application
exogenous dopamine (**Figure 6,7**). Finally, we show that loss of specific combinations
of dopaminergic receptors can exhibit effects to the basal rate of off lawn egg laying, but
importantly also appear to modulate the magnitude of predator response (**Figure 8,9**).
Other biogenic amines such as serotonin also appear to exert effects on off lawn egg
laying and their contributions to predator-evoked response merits future investigation.

CEP neurons have been previously implicated in learning the size of bacterial
lawns. We previously showed that *C. elegans* learns the size of the lawn by using high
threshold sensory neurons that detect lawn edges, which in turn signal to CEP neurons
to release dopamine. In this paradigm, we speculate that information about lawn size
was stored in amount of dopamine released from CEP neurons [55]. PDE neurons are
involved increasing egg laying during roaming, and dopamine release can increase the
probability of egg laying in the absence of food [53], so dopamine release from PDE in
this predator-prey assay could also encourage egg laying off lawn. While it is the case
that effects observed in our transgenic rescue experiments could be due to artifacts of
promoter usage, this known division of labor between CEP and PDE could also explain
the intermediate levels of rescue to off lawn laying we observe.

We observe a role for multiple dopamine receptors in this prey response to
predator threat. The *C. elegans* genome encodes at least four dopamine receptors [62].
While DOP-1 and DOP-2/3 are the *C. elegans* homologs of the mammalian D1-like and
D2-like receptors respectively, DOP-4 is a D1-like receptor unique to invertebrates

[57,63]. We find that the *dop-1*; *dop-2*; *dop-3* triple mutant animals have a reduced

response to predator threat while maintaining normal off lawn egg laying behavior.

Complete loss of all four receptors, or the double loss of *dop-3* and *dop-4*, results in

greatest reductions to baseline off lawn laying. Studies in mammals where

pharmacology and receptor knockouts have shown that knockouts in D1- and D2-like

receptors can have opposing effects on behavior [57,64,65,66]. Here we show that

specific combinations of receptors can exert varying effects. While we did not identify

the site of action of these receptors, we suggest that the combined action of DOP-1, -2,

and -3 receptors act downstream of dopamine release to alter prey egg location in

predator-exposed animals.

Ideas and speculation

We speculate that responses by *C. elegans* to predator exposure fit within the

broader context of prey refuge, wherein a prey adopts a strategy to reduce predation

risk. The prey refuge brings with it the potential cost of decreased feeding opportunities,

which is weighed against the benefit of minimal harm induced by the predator [67]. This

theoretical framework is consistent with the interactions we find between predator and

changes to bacterial topology. Predator-exposed *C. elegans* shift egg to streaks away

from a central lawn, and this strategy may lower the encounter probability with

Pristionchus thus minimizing risk to the prey. This is especially so given the

observations we have previously made that *Pristionchus* predators prefer to patrol a

main lawn when it is available, thus leaving refuges open for exploitation by *C. elegans*

[34]. However, such refuges afforded by these streaks may have detriment to *C.*

elegans fitness due to their lower density of available food, and longer term monitoring

of health and fitness of prey in these conditions has yet to be tested. Variations to the

number of available refuges for fleeing prey, as well as their local food density and

quality, can be modified in the future to gain a better appreciation for this intriguing

model of prey risk minimization strategy in *C. elegans*.

Dopamine has been shown to affect multiple *C. elegans* behaviors including

locomotion, foraging, and learning [16,55]. For example, we previously showed that this

pathway is required for learning the size of a bacterial lawn and then driving a search

strategy when removed from that lawn [55]. Furthermore, dopamine has been shown to

promote egg laying when animals roam [22]. This may explain the interaction effects we

observe between predator exposure and artificially streaking bacteria. The combination

of these inputs may motivate a roaming program, which continues to promote egg

laying at a distance, explaining the large boost in P(off) observed in Figures 3c and 4d.

Our video tracking data (Figure 6-figure supplement 1) suggests *cat-2*(*e1112*) mutants

are able to avoid the predator at least some of the time at perhaps an attenuated

magnitude of response. However, despite this, they very rarely lay eggs off the lawn at

all with P(off) values as low as 0.004 and as high as 0.13 across all experiments. Given

that loss of dopamine synthesis appears to suppress P(off) and addition of exogenous

dopamine restores this baseline (Figure 6-7), this is consistent with the hypothesis that

dopamine is important in modulating egg laying while roaming. Even when *cat-2*

mutants are straying from the lawn, they are still by and large laying eggs on the lawn.

It is curious that in combination, loss of signaling via the DOP-1;DOP-2;DOP-3

receptors modulates predator response without modulating baseline P(off), while

additional loss of DOP-4 modulates the baseline. This suggests that potentially the

route through which egg laying while roaming is altered requires DOP-4, while predator

response proceeds through signaling via the other receptors. Double mutant

combinations in our data however are complex with both effects to magnitude of

predator response and baseline. These data nevertheless stratify combinations with

DOP-3 and either DOP-1 or DOP-2 as showing the most attenuated predator responses

(**Figure 9b**). In Chase and colleagues' work identifying DOP-3, triple mutants in *dop-*

1;dop-2;dop-3 show attenuated basal slowing response in the presence of food, but

nevertheless show normal dopamine-dependent paralysis, and this is also exhibited by

dop-1;dop-3 double mutants [63]. It may be that predator-evoked changes to shifting

egg locations is linked to lawn edge detection. As *dop-1;dop-2;dop-3* mutants in our

work here show similar background levels of P(off) in the mock control condition, this

may indicate that lawn edge detection is not the driving force in basal off lawn egg

laying. However, when *C. elegans* learns to associate the main lawn with an aversive

stimulus such as predator threat, then proper detection of the lawn edge would be

crucial to avoiding it. However, when *dop-4* is also mutated, it may be that this mimics

loss of dopaminergic signaling observed in dopamine synthesis mutants, which in turn

serves to modulate the baseline off lawn laying rate.

In the future, the role of serotonin should be further investigated. Serotonin has

been shown to modulate dopamine-dependent behaviors. For example, while dopamine

signaling is required for basal slowing when encountering a lawn of food, serotonin can

enhance the slowing response if the animal is starved [44]. Thus, dopamine modulates

basal behavior while serotonin modulates it in an experience-dependent manner.

Whether serotonin acts in a similar manner in this assay is yet to be investigated.

In summary, after predator exposure, *C. elegans* lays eggs in areas of high food

variability that still have some food, rather than laying eggs in a dense food patch

inhabited and preferred by predators. Loss of dopamine synthesis alters baseline egg

laying activity restored by exogenous dopamine, while nuanced combinations of

dopaminergic receptors exert effects on specific predator-evoked response. This study

lays the foundation for studying prey behavior in *C. elegans*. Future studies can use this

system to interrogate the impact of various neurotransmitter signaling pathways on *C.*

elegans feeding, reproductive, and general exploration strategies modified by

experience.

Materials and Methods

C. elegans **and** *Pristionchus spp.* **Strains**

Nematode strains used in this study are shown in the following table:

Strain Name	Genotype	Figure	Notes
N2	Wild-type	Figure 1-figure supplement 5-6, Figure 2-9	
CX7389	*kyIs392 [Pstr-2::GFP::rab-3; Pttx-3::lin-10::dsRed; Pelt-2::GFP]*	Figure 1b-c, Figure 1-figure supplement 2-3	Fluorescent eggs
CZ6326	*frIs7 [nlp-29p::GFP + col-12p::DsRed] IV*	Figure 1d-e, Figure 1-figure	Injury reporter

		supplement 4	
ARM112	*wamSi112 [eft-3p::mScarlet::unc-54 3'UTR + Cbr-unc-119(+)] II; unc-119(ed3) III*	Figure 2, Figure 6-figure supplement 1	WT whole-body fluorescent strain
IV983	*cat-2(e1112) wamSi112[eftp-3::mScarlet::unc-54 3'UTR + Cbr-unc-119(+)] II*	Figure 6-figure supplement 1	Created for this study
IV988	*wamSi112[eftp-3::mScarlet::unc-54 3'UTR + Cbr-unc-119(+)] II; dat-1(ok157) III*	Figure 6-figure supplement 1	Created for this study
CB1111	*cat-1(e1111) X*	Figure 6a-b	
CB1112	*cat-2(e1112) II*	Figure 6, 7	
MT13113	*tdc-1(n3419) II*	Figure 6a-b	
MT15434	*tph-1(mg280) II*	Figure 6a-b	
MT9455	*tbh-1(n3247) X*	Figure 6a-b	
MT15620	*cat-2(n4547) II*	Figure 6c-d	
IV111	*cat-2(e1112) II; ueEx51 [p27::cat-2-sl2-GFP; Pelt-2::GFP]*	Figure 6e-f	
IV552	*cat-2(e1112) II; ueEx355 [Pdat-1p19::cat-2-sl2-GFP; Pelt-2::GFP]*	Figure 6e-f	
LX645	*dop-1(vs100) X*	Figure 8b-c	
LX702	*dop-2(vs105) V*	Figure 8b-c	
LX703	*dop-3(vs106) X*	Figure 8b-c	
RB1254	*C52B11.3(ok1321) X*	Figure 8b-c	
LX705	*dop-1(vs100);dop-3(vs106) X*	Figure 9a-b	
LX706	*dop-2(vs105) V; dop-1(vs100)*	Figure 9a-b	

	X		
IV984	*dop-4(ok1321);dop-1(vs100) X*	Figure 9a-b	Created for this study
IV985	*dop-2(vs105) V;dop-4(ok1321) X*	Figure 9a-b	Created for this study
IV986	*dop-4(ok1321) dop-3(vs106) X*	Figure 9a-b	Created for this study
LX734	*dop-2(vs105) V; dop-1(vs100); dop-3(vs106) X.*	Figure 9c-d	
CF2805	*dop-2(vs105) V; dop-4(ok1321) dop-1(vs100) dop-3(vs106) X*	Figure 8b-c, Figure 9c-d	
CF1903	*glp-1(e2144) III*		
JU1051	*P. uniformis* wild isolate	Figure 1-9 and figure supplements	From Marie-Anne Félix
PS312	*P. pacificus* California isolate	Figure1b-e, Figure 1-figure supplement 1, Figure 1 supplement 2-4	From Ralf J. Sommer
RS5194	*P. pacificus* Japanese isolate	Figure1b-e, Figure 1 supplement 2-4	From Ralf J. Sommer
TU445	*P. pacificus eud-1(tu445) X*	Figure1b-e, Figure 1-figure supplement 2-4	From Ralf J. Sommer

Nematode growth

Nematode strains were maintained at 20°C on 6 cm Petri plates containing Nematode
Growth Medium (NGM) seeded with *Escherichia coli* OP50 bacteria as food [37].

Egg location assay

Assay plates are created by spotting 0.5ul of OP50 liquid culture (OD600=0.5) on 35mm
standard NGM plates [37]. The bacterial lawns are allowed to grow at 20°C for 30 hours,
then stored for up to one month at 4°C. Whatman filter paper with ¼" punch forms the
"corral" and encircles the lawn, allowing approximately 1.5mm of clean agar in between
the lawn edge and the corral edge. All animals are allowed to crawl on a clean section
of agar to clean them of bacteria and picked to the assay plate using a sanitized
eyelash, placed next to the lawn on a clean area of agar. Three predators are picked
first, staged by overall size and pigment development as J4s. Then three *C. elegans*
L4s are picked to the assay plate. The animals are allowed to interact for a determined
amount of time, 20 hours for an overnight assay, at 20°C. For short-term exposure (6
hours and under), gravid *C. elegans* adults and adult predators are used by picking L4s
or J4s the day before to plates with plenty of food. The juveniles are allowed to grow
overnight into adulthood and then used in the same assay setup. After their interaction,
corrals and all adults are removed from the plate and the area inside the corral is
imaged using an AxioZoom V16 (ZEISS).

For the streaked lawn variation, streaks are formed by gently dragging a sanitized
eyelash through the center of the lawn in radial streaks ten times, followed by two
concentric circular streaks halfway between the lawn and the corral edge. The streaked
lawn is then used immediately.

Injury assay

Injury assays are set up in the same way as the egg location assays, using a *C. elegans*
strain containing the array *frIs7* [*Pnlp-29*::GFP + *Pcol-12*::*DsRed*]. After the set
interaction time, worms are immobilized by placing the plates on ice and imaged on an
AxioZoom V16 (ZEISS) within one hour, with exposure times kept constant for
fluorescence imaging (25ms).

Learning assay

C. elegans are trained using the 20-hour egg location assay. At the same time as the

animals used for training are transferred to their assay plates, test plates are set up.

Three types of test plates are used: a filled lawn (10ul of OP50 (OD600=0.5), a streaked

lawn (same as the streaked lawn variant of the egg location assay), and a small lawn

(same as the original assay plate). The training plates with animals on them and the test

plates are incubated at 20°C for 20 hours, during which the *C. elegans* is exposed to

JU1051 males and the smears on the test plates are allowed to grow. (The bacteria on

the other test plates are also allowed to grow at this time so that the bacteria are at a

similar metabolic state and density across test plates, and streaks are already present).

Filter paper corrals like those used in the egg location assay are centered over the test

plate lawns.

　　　　After the *C. elegans* are incubated in their training conditions for 20 hours, they

are carefully removed with an eyelash pick from their training plates to a clean section

of an NGM plate. The animal is allowed to crawl for a few seconds to remove bacteria

and then picked to a test plate halfway between the central lawn and the corral edge.

For the filled test lawns, the animals are placed in an equivalent position relative to the

corral edge. The test plates are then imaged every hour on an AxioZoom V16 for 6

hours. For the variant including rest plates, the *C. elegans* are picked from their training

plate to rest plates ("filled lawn" plates) for the set rest time. They are then transferred to

a streaked lawn test plate and egg locations are observed after two hours. In learning

experiments, all three *C. elegans* in predator exposed conditions are transferred to a

rest plate or test arena. In Mock controls, where there are six *C. elegans* present, three

C. elegans are selected randomly for transfer.

Exogenous dopamine assay

When adding exogenous dopamine to the learning assay, a 200mM stock of dopamine

hydrochloride (Code 122000100 Lot: A0427132, CAS: 62-31-7, Acros Organics) in

water was prepared. Two hours before the trained worms needed to be transferred to

the test plates, 50μl of the dopamine stock or water as a control was gently applied onto

the streaked lawn test plate. The plates were allowed to diffuse and dry with the lids off

for two hours, at which time the trained worms were transferred to the test plates. The

trained worms were allowed to lay eggs for two hours before their plates were imaged.

Egg location image quantification

Egg location images are quantified in FIJI with the experimenter blinded to the condition

by randomizing the file order and obscuring the filenames (using the

Filename_Randomizer macro found at

https://imagej.nih.gov/ij/macros/Filename_Randomizer.txt). Eggs are manually selected

with the multipoint tool and lawns are selected as circles. Distances from each egg from

to the lawn edge are calculated in Python. All assays are performed with their relevant

controls over at least three separate days.

WormWatcher assays

Assays conducted in the WormWatcher (Tau Scientific Instruments, West Berlin, New

Jersey, USA) were performed on a single 6cm 2.5% agar NGM plate in a 12-arena

setup. The 12-well corral was created by cutting a 3x4 array of ¼" circles into a plastic

sheet using a Cricut machine. OP50 was spotted in the 3x4 pattern using the same

concentration and allowed to grow for the same amount of time as in the egg location

assay. The increased agar percentage on the WormWatcher plates helped prevent

worms from escaping under the plastic edges of the corral.

The assays were set up like the egg location assays, with three L4 *C. elegans* and three

J4 JU1051 males or six L4 *C. elegans* in control arenas. The positions of predator-

containing and/or mutant-containing arenas was alternated on different assay days. The

WormWatcher was set to acquire fluorescent frames with a green LED excitation light

every four minutes. A reference darkfield image was acquired before and after every

experiment to reference the positions of the arenas and the size and positions of the

lawns. After the experiment was completed, each arena was inspected and imaged to

determine whether any worms escaped away or into it. Custom code was written to

segment the *C. elegans* and arenas in each position and the median distance from the

lawn edge to the mid-point of each worm body per well was recorded. Data from arenas

were discarded if two worms had escaped from an arena, or if a *P. uniformis* was seen

in a control arena.

Pristionchus mouthform analysis

Pristionchus mouthform analysis was performed as reported in [68]. Briefly, *Pristionchus*

were egg-prepped via bleaching and eggs were either cultured on standard solid NGM

plates or in liquid culture. After eggs reached adulthood, they were immobilized on

agarose slides with sodium azide. The slides of different strains from different culture

conditions were mixed and their labels obscured while they were observed. The slides

were scored as either Eu (wide mouth, two teeth) or St (narrow mouth, one tooth) while

the experimenter was blinded to strain.

Statistical methods

Egg location data. For egg location assays the number of eggs on and off the bacterial

lawn were quantified from images. Tabulated egg data as numbers of eggs off and on

the lawn were analyzed via binomial generalized linear models (logistic regression) in R

using the glm function to fit one, two, or three-way interactions between independent

variables [69]. These models are fit using the logit link function (**Equation 1**):

$$log\left(\frac{P(off)}{1-P(off)}\right) \sim X \cdot \beta \qquad (1)$$

where P(off) is the expected probability of off lawn egg laying, X is the design matrix of

categorical or continuous predictors and β is the vector of fitted coefficients. The

quantity $\frac{P(off)}{1-P(off)}$ is the "odds ratio" of laying eggs off the lawn, and thus the logit is the

logarithmic scale odds ratio. Changes to the log odds ratio can be interpreted as

changes to odds of laying eggs off lawn vs. on lawn. The expected probability P(off)

under different conditions and associated confidence intervals can be determined from

exponentiation of logit scale quantities using the inverse logit function (**Equation 2**):

$$P(off) \sim \frac{e^{X \cdot \beta}}{1+e^{X \cdot \beta}} \qquad (2)$$

These estimates for the expected value of P(off) with its associated 95% confidence

interval were used for overlaying on plots. Omnibus effects in the data were determined

by likelihood ratio tests/Analysis of Deviance using the Anova function in the car

package in R [70]. Where significant main effects or interactions were detected, post

hoc linear hypotheses included both comparisons between groups as well as higher

order comparisons of magnitudes of change were computed (as in Figures 5 – 9, e.g.

the change to the magnitude of change between Predator and Mock conditions between

genotypes). The "Predator Response" in Figures 5 – 9 is specifically defined as the

change to the expected value of the log odds ratio (Equation 1) between Mock and

Predator conditions (**Equation 3**):

$$\log Predator\ Response = \log\left(\frac{P(off)}{1-P(off)}\right)_{Predator} - \log\left(\frac{P(off)}{1-P(off)}\right)_{Mock}$$

739   (3)

which can straightforwardly be computed for a particular experimental condition or

genotype from logistic models by linear combination of the coefficients in β (**Equations**

1-2, with associated standard error and confidence intervals used for inference). In plain

language this represents the change in odds of off lawn egg laying observed in the

Predator condition relative to the Mock control. As natural logarithms are cumbersome

for easy interpretation on plots, Figures 5-9 use base 2 logarithms where each unit

change corresponds to a two-fold change in the Predator Response as defined in

Equation 3 above.

All linear hypotheses were computed using the glht function in the multcomp

package in R with associated correction for multiple testing performed using the

multivariate normal distribution (Z tests with the "single step" method for generalized

linear models, according to the simultaneous p-value estimation method of

Hothorn, Brez, and Westfall [71].) All statistical inference for differences between groups

is performed on the logit scale but linear scale P(off) values are shown on plots for ease

of interpretation.

WormWatcher positional tracking data. Distance from body to center of arena over

20 hours of observation in WormWatcher assays was subjected to non-parametric

bootstrap resampling with replacement for 10^5 iterations with empirical 95% intervals

determined using the quantile function in R. Significant changes to position with respect

to time between conditions were inferred at p<0.05 where empirical bootstrapped

intervals failed to overlap.

Egg count data and P*nlp-29*::GFP fluorescence data. Average number of eggs per

individual *C. elegans* in assays as well as logarithmic scale normalized fluorescence in

Figure 1 and Figure 1-figure supplement 1 were tested for main effects and interactions

between independent variables using general linear models using the lm function and

the Anova function from the car package. Omnibus effects in the data were determined

by ANOVA. Where significant main effects or interactions were detected, post hoc linear

hypothesis tests for differences between conditions were determined using the glht

function in the multcomp package in R with associated correction for multiple testing

performed using the multivariate t distribution (the "single step" method for

ANOVA/linear models according to the simultaneous p-value estimation method of

Hothorn, Brez, and Westfall [71]).

Mouthform analysis. Changes to abundance of stenostomatous or eurystomatous

Pristionchus was determined by Fisher's exact test.

CLUSTAL alignment of DOP receptors. Alignment of receptors shown in Figure 8 was

performed using Clustal Omega on the EMBL-EBI server at

https://www.ebi.ac.uk/Tools/msa/clustalo/ [72].

Data Availability

Raw data for experiments in Figures 1 – 9 and figure supplements are provided as

source data files in MS Excel format. Analysis code for computation of associated

effects can be found on the Shrek Lab github: https://github.com/shreklab.

Acknowledgments

We would like to thank Nadia Haghani and Adeline Sov for their help in collecting the

Pristionchus mouthform data. Anthony Fouad wrote custom code to help run the

WormWatcher imaging setup as well as analyze the resulting images. We would also

like to thank Kathleen Quach, Kirthi Reddy, Jess Haley, Wen Mai Wong, and Callum

Walsh for contributing comments on this manuscript. This work was funded by a

Graduate Research Fellowship from the National Science Foundation (A.P.), an

Innovation grant from Kavli Institute of Brain and Mind (A.P.), and a NIH R01 MH113905

(S.H.C.).

References

1. Belgrad BA, Griffen BD (2016) Predator–prey interactions mediated by prey personality and predator hunting mode. Proceedings of the Royal Society B: Biological Sciences 283: 20160408.
2. Palmer MS, Packer C (2021) Reactive anti-predator behavioral strategy shaped by predator characteristics. PloS one 16: e0256147.
3. Garcia J, Koelling RA (1966) Relation of cue to consequence in avoidance learning. Psychonomic science 4: 123-124.
4. Kavaliers M, Choleris E (2001) Antipredator responses and defensive behavior: ecological and ethological approaches for the neurosciences. Neurosci Biobehav Rev 25: 577-586.
5. Lima SL (1998) Nonlethal effects in the ecology of predator-prey interactions. Bioscience 48: 25-34.
6. Sih A (1980) Optimal behavior: can foragers balance two conflicting demands? Science 210: 1041-1043.
7. Childress MJ, Lung MA (2003) Predation risk, gender and the group size effect: does elk vigilance depend upon the behaviour of conspecifics? Animal behaviour 66: 389-398.
8. Laundré JW, Hernández L, Altendorf KB (2001) Wolves, elk, and bison: reestablishing the" landscape of fear" in Yellowstone National Park, USA. Canadian Journal of Zoology 79: 1401-1409.
9. Choi J-S, Kim JJ (2010) Amygdala regulates risk of predation in rats foraging in a dynamic fear environment. Proceedings of the National Academy of Sciences 107: 21773-21777.
10. Kim EJ, Kong M-S, Park SG, Mizumori SJ, Cho J, Kim JJ (2018) Dynamic coding of predatory information between the prelimbic cortex and lateral amygdala in foraging rats. Science advances 4: eaar7328.
11. Schulenburg H, Félix M-A (2017) The natural biotic environment of Caenorhabditis elegans. Genetics 206: 55-86.
12. White JG, Southgate E, Thomson JN, Brenner S (1986) The structure of the nervous system of the nematode Caenorhabditis elegans. Philos Trans R Soc Lond B Biol Sci 314: 1-340.
13. Yilmaz M, Meister M (2013) Rapid innate defensive responses of mice to looming visual stimuli. Current Biology 23: 2011-2015.
14. De Franceschi G, Vivattanasarn T, Saleem AB, Solomon SG (2016) Vision guides selection of freeze or flight defense strategies in mice. Current biology 26: 2150-2154.
15. Gray JM, Hill JJ, Bargmann CI (2005) A circuit for navigation in Caenorhabditis elegans. Proc Natl Acad Sci U S A 102: 3184-3191.
16. Hills T, Brockie PJ, Maricq AV (2004) Dopamine and glutamate control area-restricted search behavior in Caenorhabditis elegans. J Neurosci 24: 1217-1225.
17. Zhang Y, Lu H, Bargmann CI (2005) Pathogenic bacteria induce aversive olfactory learning in Caenorhabditis elegans. Nature 438: 179-184.
18. Guo M, Wu T-H, Song Y-X, Ge M-H, Su C-M, Niu W-P, Li L-L, Xu Z-J, Ge C-L, Al-Mhanawi MTH, Wu S-P, Wu Z-X (2015) Reciprocal inhibition between sensory ASH and ASI neurons modulates nociception and avoidance in Caenorhabditis elegans. Nature Communications 6: 5655.
19. Ezcurra M, Tanizawa Y, Swoboda P, Schafer WR (2011) Food sensitizes C. elegans avoidance behaviours through acute dopamine signalling. Embo J 30: 1110-1122.

20. Kindt KS, Quast KB, Giles AC, De S, Hendrey D, Nicastro I, Rankin CH, Schafer WR (2007) Dopamine mediates context-dependent modulation of sensory plasticity in C. elegans. Neuron 55: 662-676.

21. Alkema MJ, Hunter-Ensor M, Ringstad N, Horvitz HR (2005) Tyramine Functions independently of octopamine in the Caenorhabditis elegans nervous system. Neuron 46: 247-260.

22. Cermak N, Stephanie KY, Clark R, Huang Y-C, Baskoylu SN, Flavell SW (2020) Whole-organism behavioral profiling reveals a role for dopamine in state-dependent motor program coupling in C. elegans. Elife 9: e57093.

23. Sommer RJ (2006) Pristionchus pacificus. WormBook: 1-8.

24. Hong RL, Sommer RJ (2006) Chemoattraction in Pristionchus nematodes and implications for insect recognition. Curr Biol 16: 2359-2365.

25. Herrmann M, Mayer WE, Sommer RJ (2006) Nematodes of the genus Pristionchus are closely associated with scarab beetles and the Colorado potato beetle in Western Europe. Zoology (Jena) 109: 96-108.

26. Félix M-A, Ailion M, Hsu J-C, Richaud A, Wang J (2018) Pristionchus nematodes occur frequently in diverse rotting vegetal substrates and are not exclusively necromenic, while Panagrellus redivivoides is found specifically in rotting fruits. PloS one 13: e0200851.

27. Felix MA, Jovelin R, Ferrari C, Han S, Cho YR, Andersen EC, Cutter AD, Braendle C (2013) Species richness, distribution and genetic diversity of Caenorhabditis nematodes in a remote tropical rainforest. BMC Evol Biol 13: 10.

28. Von Lieven AF, Sudhaus W (2000) Comparative and functional morphology of the buccal cavity of Diplogastrina (Nematoda) and a first outline of the phylogeny of this taxon. Journal of Zoological Systematics and Evolutionary Research 38: 37-63.

29. Sudhaus W, Fürst von Lieven A (2003) A phylogenetic classification and catalogue of the Diplogastridae (Secernentea, Nematoda). Journal of Nematode Morphology and Systematics 6: 43-90.

30. Serobyan V, Ragsdale EJ, Sommer RJ (2014) Adaptive value of a predatory mouth-form in a dimorphic nematode. Proc Biol Sci 281: 20141334.

31. Wilecki M, Lightfoot JW, Susoy V, Sommer RJ (2015) Predatory feeding behaviour in Pristionchus nematodes is dependent on phenotypic plasticity and induced by serotonin. Journal of Experimental Biology 218: 1306-1313.

32. Fedorko A, Stanuszek S (1971) Pristionchus uniformis sp. n.(Nematoda, Rhabditida, Diplogasteridae), a facultative parasite of Leptinotarsa decemlineata Say and Melolontha melolontha L. in Poland. Morphology and biology. Acta Parasitologica Polonica 19: 95-112.

33. Kanzaki N, Ragsdale EJ, Herrmann M, Sommer RJ (2014) Two new and two recharacterized species from a radiation of Pristionchus (Nematoda: Diplogastridae) in Europe. Journal of Nematology 46: 60.

34. Quach KT, Chalasani SH (2022) Flexible reprogramming of Pristionchus pacificus motivation for attacking Caenorhabditis elegans in predator-prey competition. Current Biology 32: 1675-1688. e1677.

35. Ragsdale EJ, Muller MR, Rodelsperger C, Sommer RJ (2013) A developmental switch coupled to the evolution of plasticity acts through a sulfatase. Cell 155: 922-933.

36. Scott E, Hudson A, Feist E, Calahorro F, Dillon J, De Freitas R, Wand M, Schoofs L,
O'Connor V, Holden-Dye L (2017) An oxytocin-dependent social interaction between
larvae and adult C. elegans. Scientific Reports 7: 1-13.
37. Brenner S (1974) The genetics of Caenorhabditis elegans. Genetics 77: 71-94.
38. Pujol N, Cypowyj S, Ziegler K, Millet A, Astrain A, Goncharov A, Jin Y, Chisholm AD,
Ewbank JJ (2008) Distinct innate immune responses to infection and wounding in the C.
elegans epidermis. Curr Biol 18: 481-489.
39. Pujol N, Zugasti O, Wong D, Couillault C, Kurz CL, Schulenburg H, Ewbank JJ (2008)
Anti-fungal innate immunity in C. elegans is enhanced by evolutionary diversification of
antimicrobial peptides. PLoS pathogens 4: e1000105.
40. Hilliard MA, Bargmann CI, Bazzicalupo P (2002) C. elegans responds to chemical repellents
by integrating sensory inputs from the head and the tail. Curr Biol 12: 730-734.
41. Liu Z, Kariya MJ, Chute CD, Pribadi AK, Leinwand SG, Tong A, Curran KP, Bose N,
Schroeder FC, Srinivasan J, Chalasani SH (2018) Predator-secreted sulfolipids induce
defensive responses in C. elegans. Nat Commun 9: 1128.
42. Horvitz HR, Chalfie M, Trent C, Sulston JE, Evans PD (1982) Serotonin and octopamine in
the nematode Caenorhabditis elegans. Science 216: 1012-1014.
43. Chase DL, Koelle MR (2007) Biogenic amine neurotransmitters in C. elegans. WormBook:
1-15.
44. Sawin ER, Ranganathan R, Horvitz HR (2000) C. elegans locomotory rate is modulated by
the environment through a dopaminergic pathway and by experience through a
serotonergic pathway. Neuron 26: 619-631.
45. Flavell SW, Pokala N, Macosko EZ, Albrecht DR, Larsch J, Bargmann CI (2013) Serotonin
and the neuropeptide PDF initiate and extend opposing behavioral states in C. elegans.
Cell 154: 1023-1035.
46. Duerr JS, Frisby DL, Gaskin J, Duke A, Asermely K, Huddleston D, Eiden LE, Rand JB
(1999) The cat-1 gene of Caenorhabditis elegans encodes a vesicular monoamine
transporter required for specific monoamine-dependent behaviors. J Neurosci 19: 72-84.
47. Sulston J, Dew M, Brenner S (1975) Dopaminergic neurons in the nematode Caenorhabditis
elegans. J Comp Neurol 163: 215-226.
48. Lints R, Emmons SW (1999) Patterning of dopaminergic neurotransmitter identity among
Caenorhabditis elegans ray sensory neurons by a TGFbeta family signaling pathway and
a Hox gene. Development 126: 5819-5831.
49. Sze JY, Victor M, Loer C, Shi Y, Ruvkun G (2000) Food and metabolic signalling defects in
a Caenorhabditis elegans serotonin-synthesis mutant. Nature 403: 560-564.
50. Schafer WR (2005) Egg-laying. WormBook: 1-7.
51. Schafer WF (2006) Genetics of egg-laying in worms. Annu Rev Genet 40: 487-509.
52. Flames N, Hobert O (2009) Gene regulatory logic of dopamine neuron differentiation. Nature
458: 885-889.
53. Nass R, Hahn MK, Jessen T, McDonald PW, Carvelli L, Blakely RD (2005) A genetic screen
in Caenorhabditis elegans for dopamine neuron insensitivity to 6-hydroxydopamine
identifies dopamine transporter mutants impacting transporter biosynthesis and
trafficking. J Neurochem 94: 774-785.
54. Carvelli L, McDonald PW, Blakely RD, Defelice LJ (2004) Dopamine transporters
depolarize neurons by a channel mechanism. Proc Natl Acad Sci U S A 101: 16046-
16051.

55. Calhoun AJ, Tong A, Pokala N, Fitzpatrick JA, Sharpee TO, Chalasani SH (2015) Neural
Mechanisms for Evaluating Environmental Variability in Caenorhabditis elegans. Neuron
86: 428-441.
56. Han B, Dong Y, Zhang L, Liu Y, Rabinowitch I, Bai J (2017) Dopamine signaling tunes
spatial pattern selectivity in C. elegans. Elife 6.
57. Sugiura M, Fuke S, Suo S, Sasagawa N, Van Tol HHM, Ishiura S (2005) Characterization of
a novel D2-like dopamine receptor with a truncated splice variant and a D1-like
dopamine receptor unique to invertebrates from Caenorhabditis elegans. Journal of
Neurochemistry 94: 1146-1157.
58. Perreault ML, Hasbi A, O'Dowd BF, George SR (2014) Heteromeric Dopamine Receptor
Signaling Complexes: Emerging Neurobiology and Disease Relevance.
Neuropsychopharmacology 39: 156-168.
59. Nouvian M, Mandal S, Jamme C, Claudianos C, d'Ettorre P, Reinhard J, Barron AB, Giurfa
M (2018) Cooperative defence operates by social modulation of biogenic amine levels in
the honey bee brain. Proceedings of the Royal Society B: Biological Sciences 285:
20172653.
60. Aonuma H (2020) Serotonergic control in initiating defensive responses to unexpected tactile
stimuli in the trap-jaw ant Odontomachus kuroiwae. Journal of experimental biology 223:
jeb228874.
61. Gibson WT, Gonzalez CR, Fernandez C, Ramasamy L, Tabachnik T, Du RR, Felsen PD,
Maire MR, Perona P, Anderson DJ (2015) Behavioral responses to a repetitive visual
threat stimulus express a persistent state of defensive arousal in Drosophila. Current
Biology 25: 1401-1415.
62. Hobert O (2013) The neuronal genome of Caenorhabditis elegans. WormBook: 1-106.
63. Chase DL, Pepper JS, Koelle MR (2004) Mechanism of extrasynaptic dopamine signaling in
Caenorhabditis elegans. Nat Neurosci 7: 1096-1103.
64. Kelly MA, Rubinstein M, Phillips TJ, Lessov CN, Burkhart-Kasch S, Zhang G, Bunzow JR,
Fang Y, Gerhardt GA, Grandy DK (1998) Locomotor activity in D2 dopamine receptor-
deficient mice is determined by gene dosage, genetic background, and developmental
adaptations. Journal of Neuroscience 18: 3470-3479.
65. Gong W, Neill D, Lynn M, Justice Jr J (1999) Dopamine D1/D2 agonists injected into
nucleus accumbens and ventral pallidum differentially affect locomotor activity
depending on site. Neuroscience 93: 1349-1358.
66. McNamara FN, Clifford JJ, Tighe O, Kinsella A, Drago J, Croke DT, Waddington JL (2003)
Congenic D1A dopamine receptor mutants: ethologically based resolution of behavioural
topography indicates genetic background as a determinant of knockout phenotype.
Neuropsychopharmacology 28: 86-99.
67. Sih A (1987) Prey refuges and predator-prey stability. Theoretical Population Biology 31: 1-
12.
68. Werner MS, Sieriebriennikov B, Loschko T, Namdeo S, Lenuzzi M, Dardiry M, Renahan T,
Sharma DR, Sommer RJ (2017) Environmental influence on Pristionchus pacificus
mouth form through different culture methods. Scientific Reports 7: 1-12.
69. Team RDC (2009) A language and environment for statistical computing. http://www R-
project org.
70. Fox J, Weisberg S (2018) An R companion to applied regression: Sage publications.

71. Hothorn T, Bretz F, Westfall P (2008) Simultaneous inference in general parametric models.
Biometrical Journal: Journal of Mathematical Methods in Biosciences 50: 346-363.
72. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,
Remmert M, Söding J (2011) Fast, scalable generation of high-quality protein multiple
sequence alignments using Clustal Omega. Molecular systems biology 7: 539.
978
979

Figure legends

Figure 1. Predators influence prey egg location. (a) Schematic showing egg location assay setup. Small lawns (approx. 3 mm) in diameter are enclosed in a filter paper corralled arena. Six animals are placed into the arena, 3 GFP+ *C. elegans* strain CX7389 and 3 *Pristionchus spp.* (or 6 *C. elegans* in Mock controls). After 20 hours, eggs are visualized and <x,y> positions in the arena are determined. (b) Histograms of egg distributions in Mock (N=10 arenas) or 5 predator conditions: *P. pacificus eud-1*(*tu445*) mutants (N=11 arenas), *P. pacificus* strain PS312 (California isolate) (N=11 arenas), *P. pacificus* strain RS5194 (Japan isolate) (N=9 arenas), *P. uniformis* strain JU1051 males (N=11 arenas), and *P. uniformis* JU1051 females (N=10 arenas). Bolded bars show average distribution of egg distance from center (in mm) with faint bars indicating the individual arena distributions. Lawn edge is marked at radial distance approximately 1.5 mm from center. (c) Distributions of eggs are quantified as <# eggs off lawn, # eggs on lawn> in each arena and the observed probability of off lawn egg laying (P(off)) is plotted in each condition (# eggs off lawn / total # of eggs, average of ~90 eggs per arena). Statistical analysis was performed by logistic regression in R modeling the [# off, # on] egg counts as a function of predator condition, with significant effects determined by likelihood ratio Analysis of Deviance in R. Model estimates are overlaid on plots as expected values of P(off) from the logistic model ± 95% confidence intervals. We detected a significant main effect of predator condition ($p < 2.2 \times 10^{-16}$). Post hoc comparisons with correction for multiple testing were computed using the single step multivariate normal procedure in the *multcomp* package in R according to simultaneous method of Hothorn, Brez, and Westfall [71]. (d) *C. elegans* expressing P*nlp-29*::GFP and a P*col-12*::dsRed co-injection marker paired with various predators after 20 hours. *P. pacificus* RS5194 animals all died following 20 hours of predator exposure. GFP signal was quantified and normalized to dsRed signal for each animal. (e) $\log_2$ (GFP/dsRed) signal is shown relative to the Mock mean (=0). N= 79 Mock, 47 *P. pacificus eud-1*(*tu445*), 34 *P. pacificus* PS312, 44 *P. uniformis* JU1051 males, 49 *P. uniformis* JU1051 females. Statistical analysis was performed with ANOVA and we detected a significant main effect of predator condition ($p < 2.2 \times 10^{-16}$). Model estimates are overlaid on plots as mean $\log_2$ normalized fluorescence ± 95% confidence intervals.

Post hoc comparisons with correction for multiple testing were performed using the

single step multivariate t procedure in the *multcomp* package in R [71]. (f)

Representative images of egg location assay plates after 20 hours of Mock (upper) or

exposure to *P. uniformis* males (lower). Red arrows indicate example eggs laid off lawn.

n.s.=p>0.1, †=p<0.1, * p<0.05, ** p<0.01, ***p<0.001, ****<p<0.0001.

Figure 1-source data 1. **Egg position data in various predator conditions**. For each

test arena, data tabulate the arena, strain, condition, <x,y> coordinates in pixels, lawn

radius in pixels, the egg distance from arena center in pixels, the egg distance from

lawn edge in pixels, position as 1 (off) or 0 (on) lawn, the conversion factor for pixel data

in mm-per-pixel, the calculated distance from center in mm, the calculated distance from

the lawn edge in mm.

Figure 1-source data 2. **P*nlp-29*::GFP and *Pcol-12*::dsRed data in various predator**

conditions. Individual animal fluorescence data is tabulated with strain, predator

condition, GFP intensity, dsRed intensity, the calculated GFP/dsRed ratio, the

calculated $\log_2$ GFP/dsRed ratio, and the normalized $\log_2$ GFP/dsRed set relative to the

average of the Mock (Control) condition.

Figure 1-figure supplement 1. Eurystomatous and stenostomatous animals in *P.*

***uniformis* and *P. pacificus*.** Proportion of stenostomatous and eurystomatous animals

in *P. uniformis* females (N=40), *P. uniformis* males (N=33), and *P. pacificus* strain

PS312 (N=69). We did not detect significant differences in the steno-/eurystomatous

ratio between *P. uniformis* males and females (Fisher's Exact Test, p = 0.77). We did

detect a significantly lower proportion of stenostomatous animals in *P. pacificus* PS312

compared to *P. uniformis* males (Fisher's Exact Test, p = 0.036) and a trend to lower

proportion of stenostomatous animals compared to *P. uniformis* females (Fisher's Exact

Test, p = 0.095).

Figure 1-figure supplement 2-source data 1. Egg position data in various

predator conditions from 1 – 6 hours. For each test arena, data tabulate the arena,

strain, condition, <x,y> coordinates in pixels, lawn radius in pixels, the egg distance

from arena center in pixels, the egg distance from lawn edge in pixels, position as 1 (off)

or 0 (on) lawn, the conversion factor for pixel data in mm-per-pixel, the calculated

distance from center in mm, the calculated distance from the lawn edge in mm, time in

hours.

Figure 1-figure supplement 3. Number of eggs laid in arenas after 20 hours of

exposure to various predators. Average number of eggs per individual *C. elegans*

animal from assays shown in Figure 1B and 1C are plotted for each predator condition.

We did not detect a significant effect of predator condition on the number of eggs laid

(one-way ANOVA, p = 0.14). Expected values are overlaid on plots as mean number of

eggs ± 95% confidence intervals.

Figure 1-figure supplement 4. 6 hour time course of P*nlp-29*::GFP fluorescence

with various predators. We determined the $\log_2$ (P*nlp-29*::GFP/P*col-12*::dsRed)

relative to the Mock condition mean from 2 - 6 hours. Relative fluorescence data were

analyzed by ANOVA/linear regression as a two-way interaction of time as a continuous

variable and predator condition. We detected a significant two-way interaction of

predator condition and time (ANOVA p < 2.2×10^{-16}). To explore this interaction visually,

observed fluorescence data are only plotted once per condition, but trendlines ± 95%

confidence intervals are shown for each pairwise comparison of predator condition to

Mock control. In Mock conditions, *P. uniformis* male, and *P. uniformis* female conditions,

we did not observe a significant relationship of fluorescence over time, but a positive

relationship was observed for both *P. pacificus* strains. Pairwise comparisons at

individual time points between predator and Mock conditions were computed with

correction for multiple testing using the single step method in the *multcomp* package in

R as in Figure 1. n.s.=p>0.1, †=p<0.1, * p<0.05, ** p<0.01, ***p<0.001, ****<p<0.0001.

Figure 1-figure supplement 4-source data 1. P*nlp-29*::GFP and P*col-12*::dsRed

data in various predator conditions from 2 – 6 hours. Individual animal fluorescence

data is tabulated with strain, predator condition, GFP intensity, dsRed intensity, the

calculated GFP/dsRed ratio, the calculated log2 GFP/dsRed ratio, the normalized $\log_2$

GFP/dsRed set relative to the average of the Mock (Control) condition, and time in

hours.

Figure 1-figure supplement 5. Different ratios of predator:prey alter the probability

of off lawn egg laying. (a) The number eggs laid off and on lawn was determined in

different ratios of predator:prey (0:6 N=9 arenas, 1:5 N=8 arenas, 2:4, 3:3, 4:2, 5:1 all

N=10 arenas). Plotted are observed P(off) in each ratio condition. <# off, #on> data

were analyzed by logistic regression/analysis of deviance. Overlaid are model expected

values of P(off) ± 95% confidence intervals. We detected a significant main effect of

predator:prey ratio (likelihood ratio p < 2.2×10^{-16}). Post hoc comparisons with correction

for multiple testing were computed using the single step method in the *multcomp*

package in R. (b) The average number of eggs per individual *C. elegans* in the same

arenas as (a) was calculated and data analyzed by ANOVA. Overlaid are mean values

± 95% confidence intervals. We did not detect a significant effect of predator:prey ratio

on the number of eggs laid (ANOVA p = 0.92). n.s.=p>0.1, †=p<0.1, * p<0.05, ** p<0.01,

p<0.001, *<p<0.0001.

Figure 1-figure supplement 5-source data 1. Egg position data in various

predator:prey ratios. For each test arena, data tabulate the arena, ratio of

predator:prey, <x,y> coordinates in pixels, lawn radius in pixels, the egg distance from

arena center in pixels, the egg distance from lawn edge in pixels, position as 1 (off) or 0

(on) lawn, the conversion factor for pixel data in mm-per-pixel, the calculated distance

from center in mm, the calculated distance from the lawn edge in mm.

Figure 1-figure supplement 6. Bacteria pre-conditioned with *P. uniformis* males is

not sufficient to alter egg laying behavior. (a) The number of eggs laid off and on

lawn was determined in arenas with lawns pre-conditioned with Mock (sterile *C.*

elegans) (N=7 arenas) or *P. uniformis* males (N=5 arenas). Plotted are observed values

of P(off) with data further analyzed by logistic regression/analysis of deviance as in

Figure 1, and Figure 1-figure supplement 2. Overlaid are logistic model estimates of the

expected values of P(off) ± 95% confidence intervals. We did not detect a significant

effect of lawn condition on P(off) (likelihood ratio p = 0.73). (b) The average number of

eggs per individual *C. elegans* animal in arenas in (a) was determined and is plotted for

each condition, with data analyzed by ANOVA. Overlaid are mean estimates ± 95%

confidence intervals. We did detect a significant increase to the average number of

eggs laid in lawns conditioned by *P. uniformis males* (ANOVA p = 0.03). * p<0.05

Figure 1-figure supplement 6-source data 1. Egg position data in arenas with conditioned lawns. For each test arena, data tabulate the arena, condition, <x,y> coordinates in pixels, lawn radius in pixels, the egg distance from arena center in pixels, the egg distance from lawn edge in pixels, position as 1 (off) or 0 (on) lawn, the conversion factor for pixel data in mm-per-pixel, the calculated distance from center in mm, the calculated distance from the lawn edge in mm.

Figure 2. *C. elegans* shows sustained avoidance of the lawn of the lawn when exposed to predator. (a) Schematic for WormWatcher experiments for location tracking. Distance of midpoint of fluorescent *C. elegans* (strain ARM112, P*eft-3*::mScarlet) to center of the arena is tracked over 20 hours (15 frames per hour, $t_{resolution}$ = 4 min). (b) Worms tracked by WormWatcher included ARM112 strain *C. elegans* in Mock (N=12 arenas), or predator (*P. uniformis* males, N=12 arenas), and are plotted as individual traces (thin lines, average position of all worms in an arena, range 2-6 worms per arena, average = 3), representing average distance from center in mm over time. Data were analyzed by non-parametric bootstrap resampling with replacement with 1×10^5 iterations. Bold lines represent the estimated average distance over time, with shading representing empirically determined 2.5% - 97.5% quantiles (95% interval) of bootstrap samples. $p < 0.05$ significance can be inferred from regions of lack of overlap of bootstrapped intervals between Mock and predator-exposed conditions, identified with lines above traces showing regions of 0% overlap. Regions with 0% overlap account for 71% of all time points, all occurring in the region >5 hours.

Figure 2-source data 1. **WormWatcher tracking data for predator and mock-exposed ARM112 mScarlet expressing *C. elegans*.** For each test arena, 20 hours (15 frames per hour, tresolution = 4 min) of tracking data are tabulated, showing frame number, time (in hours), arena, condition, number of worms tracked per time point, lawn radius (in pixels), distance of body midpoint to center of arena (in pixels), distance of body midpoint to edge of lawn (in pixels), position as 1 (off) or 0 (on) lawn, the conversion factor for pixel data in mm-per-pixel, the calculated distance from center in mm, the calculated distance from the lawn edge in mm.

Figure 3. Bacterial topography alone does not account for predator associated changed to egg laying location. (a) Representative image of an assay plate after 20 hours with artificially streaked lawns. (b) Histograms of egg distributions in Mock (N=14 arenas), artificially streaked (N=14 arenas), and predator-exposed (N=17 arenas) conditions. Bolded bars show average distribution of egg distance from center (in mm) with faint bars indicating the individual arena distributions. Lawn edge is marked at radial distance approximately 1.5 mm from center. (c) Distributions of eggs are quantified as [# off lawn, # on lawn] in each arena as in Figure 1, and the observed probability of off lawn laying (P(off)) is plotted in each condition, with data analyzed by logistic regression/analysis of deviance. Overlaid are logistic model estimates of the expected values of P(off) ± 95% confidence intervals. We detected a significant effect of condition (likelihood ratio $p < 2.2 \times 10^{-16}$). Post hoc comparisons with correction for multiple testing were computed using the single step method in the *multcomp* package in R. n.s.=$p > 0.1$, †=$p < 0.1$, * $p < 0.05$, ** $p < 0.01$, ***$p < 0.001$, ****$< p < 0.0001$.

Figure 3-source data 1. **Egg position data in arenas with and without predator exposure and artificial streaking.** For each test arena, data tabulate the arena, condition, <x,y> coordinates in pixels, lawn radius in pixels, the egg distance from arena center in pixels, the egg distance from lawn edge in pixels, position as 1 (off) or 0 (on) lawn, the conversion factor for pixel data in mm-per-pixel, the calculated distance from center in mm, the calculated distance from the lawn edge in mm.

Figure 4. Sustained changes to egg laying is observed following prior predator exposure. (a) Schematic of egg laying learning assay: after 20 hours of exposure to either Mock (*C. elegans* only) or predator condition (*P. uniformis* males) worms are transferred to arenas either completely filled with bacteria, or arenas with a normal sized small lawn or a lawn with artificial streaks. In the predator exposed condition, all three *C. elegans* are transferred, while in the Mock condition, three *C. elegans* selected at random from amongst the 6 are transferred. (b) Analysis of distributional properties of *C. elegans* egg laying for 1 to 6 hours in arenas completely filled with bacteria after Mock (N=8-9 arenas per time point), or predator exposure (N=9 arenas per time point). Plotted are the mean distance from lawn center in mm for each time point and condition.

(c) Data points represent the coefficient of variation (standard deviation divided by the

mean) for egg distances in (b) for each time point and for each condition. Data in (b)

and (c) were analyzed by linear regression/ANOVA modeling interactions of time as

continuous variable and predator exposure condition. Overlaid on plots are trendlines

for each condition from linear models with shading showing 95% confidence intervals.

We detected a significant main effect of time on both average distance from center

(ANOVA $p = 3.0 \times 10^{-6}$) as well as on the dispersal of the eggs measured by the

coefficient of variation (ANOVA $p = 0.0016$) but no significant main effect of predator

condition (average distance, $p = 0.51$; coefficient of variation, $p = 0.14$) or interaction

effects on either variable (average distance, $p = 0.76$; coefficient of variation, $p = 0.97$).

(d) Analysis of off lawn egg laying in animals exploring small lawns or lawns with

artificial streaks after 20 hours of Mock or predator exposure. Data points in (d)

represent observe P(off) in each time point and condition (N=9-12 arenas per time

point/condition). Off lawn egg laying probability was analyzed by logistic

regression/analysis of deviance modeling a three-way interaction between time as a

continuous variable, lawn type, and predator exposure condition. We detected a

significant three-way interaction between these independent variables (likelihood ratio

$p = 1.5 \times 10^{-7}$). To explore this interaction visually, observed P(off) data points are plotted

only once per time point/lawn type/condition, but for reference, the trend lines ± 95%

confidence intervals from logistic regression are shown for pairwise comparisons

between: artificially streaked and small lawns for Mock exposed animals, Predator vs.

Mock in small lawns, Predator exposure/small lawns compared to the artificially

streaked/Mock exposed animals, and finally artificially streaked lawns compared to

small lawns for Predator exposed animals. Pairwise comparisons at individual time

points between lawn types/conditions were computed with correction for multiple testing

using the single step method in the *multcomp* package in R. n.s.=p>0.1, †=p<0.1, *

p<0.05, ** p<0.01, ***p<0.001, ****<p<0.0001.

Figure 4-figure supplement 1. Table of slopes for temporal changes in the

distributional properties of eggs after predator exposure in filled arenas. Slopes

showing the temporal change per hour in mean distance from center (mm/hour) and

coefficient of variation (units/hour) in Figure 4b and 4c in predator-exposed and control

conditions.

Exposure	Parameter	Slope	Std. Error	95% C.I.	p (slope=0)
Mock	Mean Distance from Center	-0.07 mm/hr	0.02	[-0.11, -0.03]	**0.0016
Predator	Mean Distance from Center	-0.07 mm/hr	0.02	[-0.11, -0.04]	***3×10^{-4}
Mock	Coefficient of Variation	+0.017 /hr	0.008	[+0.002, 0.032]	*0.03
Predator	Coefficient of Variation	+0.017 /hr	0.007	[+0.003, 0.032]	*0.02

1191

Figure 4-figure supplement 2. **Table of slope for temporal changes to the**

probability of off lawn egg laying with and without predator exposure, and in

arenas with differing bacterial topology. Slopes showing the temporal change per

hour in log odds ratio ($\log\frac{P(off)}{1-P(off)}$) from logistic models in Figure 4d in different

conditions. On logit scale, a slope of +0.1=1.1-fold, 0.22=1.25-fold, +0.4 = 1.5-fold,

+0.69=2-fold change in the P(off)/P(on) odds ratio.

Exposure	Lawn Type	Slope	Std. Error	95% C.I.	p (slope=0)
Mock	Small lawn	-0.32	0.09	[-0.49, -0.15]	***2×10^{-4}
Mock	Artificially streaked	+0.24	0.04	[+0.18, +0.30]	****2.2×10^{-16}
Predator	Small lawn	+0.07	0.04	[9×10^{-4}, +0.15]	*0.047

| Predator | Artificially streaked | +0.10 | 0.02 | [0.05, +0.15] | ****3.9x10^{-5} |

1198

1199

Figure 4-source data 1. Egg position data in filled arenas after predator exposure.
For each test arena, data tabulate the arena, condition, <x,y> coordinates in pixels, the
egg distance from arena center in pixels, the conversion factor for pixel data in mm-per-
pixel, the calculated distance from center in mm, time in hours.

**Figure 4-source data 2. Egg position data in small or artificially streaked arenas
after predator exposure.** For each test arena, data tabulate the arena, predator
exposure condition, lawn type, <x,y> coordinates in pixels, lawn radius in pixels, the egg
distance from arena center in pixels, the egg distance from lawn edge in pixels, position
as 1 (off) or 0 (on) lawn, the conversion factor for pixel data in mm-per-pixel, the
calculated distance from center in mm, the calculated distance from the lawn edge in
mm, time in hours.

**Figure 5. Changes to egg laying behavior after predator exposure continue for 24
hours.** (a) Schematic of egg laying learning assay. *C. elegans* are exposed to Mock or
Predator condition (*P. uniformis* males) for 20 hours and transferred to a rest plate for 1,
2, or 24 hours. After rest, animals are transferred to a test arena containing artificial
streaks as in Figure 4 and positions of laid eggs are determined in order to determine
the proportion of eggs laid off and on the lawn. (b) Observed P(off) in test arenas is
plotted by condition and length of rest period (Mock/1HR N=12 arenas, Mock/2HRS
N=15 arenas, Mock/24HRS N=20 arenas, Predator-exposed/1HR N=12 arenas,
Predator-exposed/2HR N=17 arenas, Predator-exposed/24HRS N=19 arenas). Data
were analyzed by logistic regression/analysis of deviance fitting a two-way interaction of
categorical length of rest period (1-24HRS) and Mock or Predator condition, with
expected values of P(off) ± 95% confidence intervals from logistic model overlaid on
plot. We found a significant two-way interaction of rest period length and predator
exposure condition (likelihood ratio p=3.4x10^{-11}). (c) Log$_2$ fold change in computed

Predator Response is plotted for each rest time period, where Predator Response is

defined as the change to the odds ratio of [off lawn/on lawn] egg laying between

Predator and Mock conditions (see **Methods, Equation 1-3**). These are displayed as

point estimates with 95% confidence intervals as derived from logistic regression. Post

hoc comparisons between conditions, as well as changes to Predator Response, with

correction for multiple testing, were computed using the single step method in the

multcomp package in R as in previous figures. n.s.=p>0.1, †=p<0.1, * p<0.05, ** p<0.01,

p<0.001, *<p<0.0001.

Figure 5-source data 1. Egg position data after periods of 1HR, 2HRS, or 24HRS

following predator exposure. For each test arena, data tabulate the arena, predator

exposure condition, rest period, <x,y> coordinates in pixels, lawn radius in pixels, the

egg distance from arena center in pixels, the egg distance from lawn edge in pixels,

position as 1 (off) or 0 (on) lawn, the conversion factor for pixel data in mm-per-pixel,

the calculated distance from center in mm, the calculated distance from the lawn edge

in mm.

Figure 6. Loss of biogenic amine synthesis results in changes to the probability

of laying eggs off the bacterial lawn. (a) Plotted are observed P(off) data for either

Mock or Predator-exposed arenas in various mutants in biogenic amine synthesis

genes (Mock: WT N=17 arenas, *cat-1*(*e1111*) N=18, *cat-2*(*e1112*) N=13, *tph-1*(*mg280*)

N=12, *tbh-1*(*n3247*) N=12, *tdc-1*(*n3419*) N=15, Predator-exposed: WT N=16, , *cat-*

1(*e1111*) N=9, *cat-2*(*e1112*) N=12, *tph-1*(*mg280*) N=12, *tbh-1*(*n3247*) N=12, *tdc-*

1(*n3419*) N=16). Data were analyzed by logistic regression/analysis of deviance fitting a

two-way interaction of genotype and predator exposure, with overlaid expected values

of P(off) from logistic modeling ± 95% confidence intervals. We detected a significant

two-way interaction of genotype and predator exposure condition (likelihood ratio

p<2.2×10^{-16}). (b) Log_2 Predator Response (as in **Figure 5** and **Methods, Equation 3**) is

plotted as point estimates with error bars showing 95% confidence intervals across

genotypes. (c) Observed P(off) data in Mock or Predator-exposed conditions in WT or

two *cat-2* mutant alleles *n4547* and *e1112* (Mock N=9 arenas per genotype, Predator

N=8 arenas per genotype). Data analyzed as in (a) with overlaid expected values for

P(off) from the logistic model ± 95% confidence intervals. We failed to detect a

significant interaction between genotype and predator condition (likelihood ratio p =

0.22) but we were able to detect a main effect of genotype ($p<2.2 \times 10^{-6}$) and a main

effect of predator exposure ($p<2.2 \times 10^{-6}$). (d) Log$_2$ Predator Response across

genotypes as in (b). (e) Observed P(off) in WT or *cat-2*(*e1112*) mutant animals with or

without transgenic rescue of *cat-2* cDNA in either ADE/PDE or CEP neurons (Mock/WT

N=11 arenas, Predator/WT N=10 arenas, Mock/*cat-2*(*e1112*) N=10 arenas,

Predator/*cat-2*(*e1112*) N=11 arenas, *cat-2*(*e1112*); *p27::cat-2-sl2-GFP* (ADE/PDE) N=10

arenas for each condition, *cat-2*(*e1112*); *Pdat-1p19::cat-2-sl2-GFP* (CEP) N=11 arenas

per condition). Data analyzed as in (a),(c) with overlaid expected values for P(off) from

the logistic model ± 95% confidence intervals. We detected a significant two-way

interaction of genotype and predator exposure condition (likelihood ratio p<2.2x10-16).

(f) Log$_2$ Predator Response as described in (b) and (d) across genotype/transgenic

rescue conditions. Post hoc with correction for multiple testing, were computed using

the single step method in the *multcomp* package in R as in previous figures. n.s.=p>0.1,

†=p<0.1, * p<0.05, ** p<0.01, ***p<0.001, ****<p<0.0001.

Figure 6-source data 1. Egg position data in biogenic amine mutants with and

without predator exposure. For each test arena, data tabulate the arena, predator

exposure condition, genotype, <x,y> coordinates in pixels, lawn radius in pixels, the egg

distance from arena center in pixels, the egg distance from lawn edge in pixels, position

as 1 (off) or 0 (on) lawn, the conversion factor for pixel data in mm-per-pixel, the

calculated distance from center in mm, the calculated distance from the lawn edge in

mm.

Figure 6-source data 2. Egg position data in *cat-2* mutant alleles with and without

predator exposure. For each test arena, data tabulate the arena, predator exposure

condition, genotype, <x,y> coordinates in pixels, lawn radius in pixels, the egg distance

from arena center in pixels, the egg distance from lawn edge in pixels, position as 1 (off)

or 0 (on) lawn, the conversion factor for pixel data in mm-per-pixel, the calculated

distance from center in mm, the calculated distance from the lawn edge in mm.

Figure 6-source data 3. Egg position data in *cat-2* mutant alleles with and without

predator exposure and rescue of *cat-2* cDNA. For each test arena, data tabulate the

arena, predator exposure condition, genotype, <x,y> coordinates in pixels, lawn radius

in pixels, the egg distance from arena center in pixels, the egg distance from lawn edge

in pixels, position as 1 (off) or 0 (on) lawn, the conversion factor for pixel data in mm-

per-pixel, the calculated distance from center in mm, the calculated distance from the

lawn edge in mm.

Figure 6-figure supplement 1. **Mutants in dopamine synthesis and re-uptake show**

varying degrees of predator avoidance. (a) WormWatcher data tracking position (15

frames per hour, $t_{resolution}$ = 4 min) in ARM112 mScarlet-expressing *C. elegans* ("WT")

exposed to Mock (N=8 arenas) or Predator (N=8 arenas) over the course of 20 hours.

(b) Position tracking of ARM112 animals crossed into the *cat-2*(*e1112*) mutant

background in Mock (N=8) or Predator (N=9) conditions. Data in (a),(b) are plotted as

individual traces (thin lines) representing average distance from center over time per

arena. Data were analyzed by non-parametric bootstrap resampling with replacement

with 1×10^5 iterations. Bold lines represent the estimated average distance over time,

with shading representing empirically determined 2.5% - 97.5% quantiles (95% interval)

of bootstrap samples. $p < 0.05$ significance can be inferred from regions of lack of

overlap of bootstrapped intervals between Mock and predator-exposed conditions,

identified with lines above traces showing regions of 0% overlap, accounting for 63% of

all time points in (a) and 38% of time points in (b). (c) The bootstrap estimates for the

magnitude of change between Predator - Mock distance from center in both genotypes.

(d),(e) WormWatcher data tracking position as in (a),(b) in either WT ARM112 controls

or ARM112 crossed into the *dat-1*(*ok157*) mutant background. Lack of bootstrap

confidence interval overlap between conditions is evident for 76% of all time points for

WT and 82% of all time points for *dat-1*(*ok157*) mutants. (f) Bootstrap estimates for the

magnitude of change between Predator - Mock distance from center in both genotypes.

Figure 6-figure supplement 1-source data 1. WormWatcher tracking data for

predator and mock-exposed ARM112 mScarlet expressing C. elegans and

ARM112 animals with *cat-2*(*e1112*) mutant allele. For each test arena, 20 hours (15

frames per hour, tresolution = 4 min) of tracking data are tabulated, showing frame

number, time (in hours), arena, condition, genotype, number of worms tracked per time

point, lawn radius (in pixels), distance of body midpoint to center of arena (in pixels),

distance of body midpoint to edge of lawn (in pixels), position as 1 (off) or 0 (on) lawn,

the conversion factor for pixel data in mm-per-pixel, the calculated distance from center

in mm, the calculated distance from the lawn edge in mm.

Figure 6-figure supplement 1-source data 2. WormWatcher tracking data for

predator and mock-exposed ARM112 mScarlet expressing C. elegans and

ARM112 animals with *dat-1*(*ok157*) mutant allele. For each test arena, 20 hours (15

frames per hour, tresolution = 4 min) of tracking data are tabulated, showing frame

number, time (in hours), arena, condition, genotype, number of worms tracked per time

point, lawn radius (in pixels), distance of body midpoint to center of arena (in pixels),

distance of body midpoint to edge of lawn (in pixels), position as 1 (off) or 0 (on) lawn,

the conversion factor for pixel data in mm-per-pixel, the calculated distance from center

in mm, the calculated distance from the lawn edge in mm.

Figure 7. **Addition of exogenous dopamine rescues egg laying behavior in**

dopamine synthesis deficient mutants. (a) Schematic of the egg laying learning

assay. *C. elegans* exposed to either Mock or Predator condition for 20 hours are

transferred to testing arenas containing artificially streaked bacteria with or without the

addition of 2 mM dopamine. (b) Observed P(off) data are plotted for each genotype,

predator exposure, and presence of exogenous dopamine (N=11 arenas all conditions

except *cat-2*(*e1112*)/Mock/+3mM dopamine condition which had N=10 arenas). Data

were analyzed as in previous figures by logistic regression/analysis of deviance fitting a

three-way interaction of genotype, predator exposure, and dopamine, with overlaid

expected values of P(off) from logistic modeling ± 95% confidence intervals. We

detected a significant three-way interaction of genotype, predator exposure, and

dopamine (likelihood ratio p=0.0006). (c) Log_2 Predator Response ± 95% confidence

intervals (as in **Figure 5-6**, see **Methods, Equation 3**) in each genotype with and

without addition of 2 mM dopamine. Post hoc with correction for multiple testing, were

computed using the single step method in the *multcomp* package in R as in previous

figures. n.s.=p>0.1, †=p<0.1, * p<0.05, ** p<0.01, ***p<0.001, ****<p<0.0001.

Figure 7-source data 1. Egg position data in *cat-2*(*e1112*) mutants with and

without predator exposure and addition of 3mM dopamine. For each test arena,

data tabulate the arena, predator exposure condition, genotype, dopamine addition,

<x,y> coordinates in pixels, lawn radius in pixels, the egg distance from arena center in

pixels, the egg distance from lawn edge in pixels, position as 1 (off) or 0 (on) lawn, the

conversion factor for pixel data in mm-per-pixel, the calculated distance from center in

mm, the calculated distance from the lawn edge in mm.

Figure 8. Mutations in DOP family dopaminergic receptors influence egg laying

behavior with predator exposure. (a) CLUSTAL Omega multiple protein sequence

alignment of the three isoforms of dopaminergic receptors DOP-1, the 6 of DOP-2, the 3

of DOP-3, and DOP-4 are shown visually as a colormap where black squares represent

sequence alignment gaps, and amino acids colors are group by type (e.g. uncharged,

charged). (b) Observed P(off) data are shown for the Mock and Predator-exposed

conditions in WT (Mock N=12 arenas, Predator N=11 arenas), a quadruple mutant for

all four receptor genes (N=10/condition), and single receptor mutants *dop-1(vs100*)

(N=12/condition), *dop-2*(*vs105*) (Mock N=12, Predator N=11), *dop-3*(*vs106*) (Mock

N=12, Predator N=11), and *dop-4*(*ok1321*) (Mock N=11, Predator N=12).). Data were

analyzed as in previous figures by logistic regression/analysis of deviance fitting a two-

way interaction of genotype and predator exposure, with overlaid expected values of

P(off) from logistic modeling ± 95% confidence intervals. We detected a significant two-

way interaction of genotype and predator condition (likelihood ratio $p<2.2x10^{-6}$). (c)

Log_2 Predator Response ± 95% confidence intervals as in **Figures 5-7** (see **Methods,**

Equation 3) across genotypes. Post hoc comparisons with correction for multiple

testing, were computed using the single step method in the multcomp package in R as

in previous figures. n.s.=p>0.1, †=p<0.1, * p<0.05, ** p<0.01, ***p<0.001,

****<p<0.0001.

Figure 8-source data 1. CLUSTAL multiple protein sequence alignment of DOP

receptor amino acid sequences. FASTA format for amino acid sequences of DOP-1,

DOP-2, DOP-3, DOP-4 proteins and isoforms obtained from Wormbase.

Figure 8-source data 2. **Egg position data in dopamine receptor mutants with and**

without predator exposure. For each test arena, data tabulate the arena, predator

exposure condition, genotype, <x,y> coordinates in pixels, lawn radius in pixels, the egg

distance from arena center in pixels, the egg distance from lawn edge in pixels, position

as 1 (off) or 0 (on) lawn, the conversion factor for pixel data in mm-per-pixel, the

calculated distance from center in mm, the calculated distance from the lawn edge in

mm.

Figure 9. **Loss of dopaminergic signaling via combinations of DOP receptors is**

associated with changes to both baseline egg laying behavior as well as the

magnitude of predator response. (a) Observed P(off) data are shown for Mock and

Predator-exposed conditions in WT or various pairwise combinations of dopamine

receptor mutants (N=12 arenas per condition except Mock/*dop-2*(*vs105*);*dop-4*(*ok1321*)

N= 9, and Predator/*dop-2*(*vs105*);*dop-4*(*ok1321*) N=10). Data were analyzed as in

previous figures by logistic regression/analysis of deviance fitting a two-way interaction

of genotype and predator exposure, with overlaid expected values of P(off) from logistic

modeling ± 95% confidence intervals. We detected a significant two-way interaction of

genotype and predator condition (likelihood ratio p<2.2x10-6). (b) Log_2 Predator

Response ± 95% confidence intervals as in **Figures 5-8** (see **Methods, Equation 3**)

across receptor mutant combinations. Below (b) is shown a qualitative visualization of

the Predator Response ranked from highest to lowest. (c) Observed P(off) data shown

for Mock and Predator-exposed conditions in WT, a triple mutant *dop-1*(*vs100*);*dop-*

2(*vs105*);*dop-3*(*vs106*), and a quadruple mutant in all four *dop-* genes (N=10 arenas per

condition except Mock/quadruple mutant N=9). Data analyzed as in (a) with overlaid

expected values for P(off) ± 95% confidence intervals. We detected a significant two-

way interaction of genotype and predator condition (likelihood ratio $p=2x10^{-13}$). (d) Log_2

Predator Response ± 95% confidence intervals as in (b). Post hoc comparisons with

correction for multiple testing, were computed using the single step method in the

multcomp package in R as in previous figures. n.s.=p>0.1, †=p<0.1, * p<0.05, ** p<0.01,

p<0.001, *<p<0.0001.

Figure 9-source data 1. Egg position data in pairwise combinations of dopamine

receptor mutants with and without predator exposure. For each test arena, data

tabulate the arena, predator exposure condition, genotype, <x,y> coordinates in pixels,

lawn radius in pixels, the egg distance from arena center in pixels, the egg distance

from lawn edge in pixels, position as 1 (off) or 0 (on) lawn, the conversion factor for pixel

data in mm-per-pixel, the calculated distance from center in mm, the calculated distance

from the lawn edge in mm.

Figure 9-source data 2. Egg position data in triple and quadruple dopamine

receptor mutants with and without predator exposure. For each test arena, data

tabulate the arena, predator exposure condition, genotype, <x,y> coordinates in pixels,

lawn radius in pixels, the egg distance from arena center in pixels, the egg distance

from lawn edge in pixels, position as 1 (off) or 0 (on) lawn, the conversion factor for pixel

data in mm-per-pixel, the calculated distance from center in mm, the calculated distance

from the lawn edge in mm.

1417

a

Pristionchus spp.

GFP+ *C. elegans*

filter paper corral

OP50 lawn

20 hours

b

□ on lawn □ off lawn

Mock

lawn edge ~1.5mm

P. pacificus eud-1 (tu445)

P. pacificus RS5194

P. pacificus PS312

P. uniformis males

P. uniformis females

Egg distance from lawn center (mm)

c

P(off)

Mock | *eud-1 (tu445)* | PS312 | RS5194 | JU1051 males | JU1051 females

P. pacificus | *P. uniformis*

d

P*col-12*::dsRED   P*nlp-29*::GFP   Merge

Alone

P. pacificus

eud-1(tu445)

PS312

P. uniformis

JU1051 M

JU1051 F

e

Δlog₂ (GFP/dsRed)

n/a

Mock | *eud-1 (tu445)* | PS312 | RS5194 | JU1051 males | JU1051 females

P. pacificus | *P. uniformis*

f

C. elegans only

off-lawn eggs

+ *P. uniformis* males

1mm

off-lawn eggs

a

Artificially Streaked Arena

1mm

b

on lawn   off lawn   on streaks

Mock

lawn edge ~1.5mm

Artificially streaked

+*P. uniformis* males

% (y-axis)

Egg distance from lawn center (mm)

c

P(off)

****   ****   ****

Mock   Artificial Streaks   *P. uniformis* males

a

Training

20h exposure
+/- predator

Testing

+/- DA

b

- ● Mock
- ● *P. uniformis* males

c

a

Clustal Omega Protein Multiple Sequence Alignment

alignment
gap G A I L M V P F W Y N C Q S T D E R H K

DOP-1

DOP-2

DOP-3

DOP-4

1   951

b

● Mock ● *P. uniformis* males

P(off)

WT ****
dop-1 (vs100);
dop-2 (vs105);
dop-3 (vs106);
dop-4 (ok1321) ****
dop-1
(vs100) ****
dop-2
(vs105) ****
dop-3
(vs106) ****
dop-4
(ok1321) ****

Changes to Mock
(vs. WT)

"d1234"

↓†   ↑****   ↑****   ↑**   n.s.

c

log₂ Predator Response

n.s.   ****   ****   ****   n.s.

WT   d1234   *dop-1*
(vs100)   *dop-2*
(vs105)   *dop-3*
(vs106)   *dop-4*
(ok1321)