Commitment in the Pink1/Parkin mitophagy decision circuit

Christopher S. Waters[1], Sigurd B. Angenent[2], Steven J. Altschuler[1]*, Lani F. Wu[1]*

[1] Department of Pharmaceutical Chemistry, University of California San Francisco; San Francisco, CA 94158, USA.

[2] Mathematics Department, University of Wisconsin Madison; Madison, WI 53706, USA.

* Corresponding authors: Steven.Altschuler@ucsf.edu, Lani.Wu@ucsf.edu

Contents:

Abstract

Mechanisms that prevent accidental degradation of healthy mitochondria by the Pink1/Parkin mitophagy pathway are poorly understood. On the surface of damaged mitochondria, Pink1 accumulates and acts as the input signal to a positive feedback loop of Parkin recruitment, which in turn promotes mitochondrial degradation via mitophagy. However, Pink1 also transiently associates with healthy mitochondria where it could errantly recruit Parkin and thereby activate this positive feedback loop. Here, we mapped the relationship between Pink1 input levels and Parkin recruitment dynamics using quantitative live-cell microscopy and mathematical modeling. We found that Parkin is recruited to the mitochondria only if Pink1 levels exceed a threshold and only after a delay that is inversely proportional to Pink1 levels. The properties of threshold and delay emerge from the Pink1/Parkin circuit topology and provide a mechanism for cells to assess damage signals before committing to mitophagy.

Introduction

When mitochondria accumulate significant oxidative damage, they cease to function properly and can become depolarized. Cells maintain a healthy pool of mitochondria by degrading depolarized mitochondria via mitophagy (mitochondrial autophagy). A core molecular circuit mediating the mitophagy decision involves Pink1, a mitochondrial kinase, and Parkin, a cytoplasmic E3 ubiquitin ligase (*1*). On healthy mitochondria, Pink1 is constantly recruited to the mitochondrial outer membrane (MOM), whereupon Pink1 undergoes polarization-dependent degradation (*2*, *3*). However, mitochondrial depolarization prevents Pink1 degradation, leading to accumulation of Pink1 on the MOM (*2*, *3*). Upon Pink1 accumulation, positive feedback in the Pink1/Parkin circuit leads to extensive labeling of the mitochondrial surface with phospho-Ubiquitin (pUb) and to pUb-dependent recruitment of downstream mitophagy machinery (*1*, *4*, *5*).

Positive feedback in the Pink1/Parkin circuit proceeds through several steps: Pink1 phosphorylates mitochondrial ubiquitin (Ub), phosphorylated ubiquitin recruits autoinhibited Parkin to the MOM, Pink1 phosphorylates Parkin's ubiquitin-like (UBL) domain to activate Parkin, and activated Parkin deposits new Ub on proteins on the MOM (*1*, *6–14*). Newly deposited Ub can then be phosphorylated by Pink1, starting new rounds of amplification (*1*, *15*). Together, these steps form a Pink1-dependent positive feedback circuit topology, where Pink1 provides an input signal that regulates continued recruitment of Ub and Parkin (Fig. 1). Previous work speculated that the Pink1/Parkin feedback topology plays a role in preventing errant mitophagy decisions (*4*, *6*, *15–17*). This speculation is motivated by the observation that mitochondria tolerate natural transient depolarization events (*18*), by the transient association of Pink1 with healthy mitochondria (*1*, *2*), and by the existence of additional layers of positive feedback in downstream mitophagy steps (*19*, *20*). Emergent properties, such as activation thresholds, have been proposed

as governors for Pink1/Parkin activation (*6, 15–17*). However, in practice, it remains unclear how noise tolerance within the Pink1/Parkin circuit arises.

Here, we investigated how cells avoid accidental degradation of healthy mitochondria via the Pink1/Parkin positive feedback circuit. We used quantitative live-cell microscopy to directly map the relationship between discrete Pink1 levels and Parkin recruitment dynamics. Our input-response mapping revealed inherent regulatory properties that allow the Pink1/Parkin pathway to only activate in the presence of both elevated and sustained mitochondrial Pink1 levels. We found that both of these properties can be modulated through inhibition of the mitochondrial deubiquitinase USP30, an antagonist of the Pink1/Parkin circuit (*4, 21–23*). Mathematical modeling then revealed that the topology of the Pink1/Parkin circuit, alone, is sufficient to generate the observed regulatory properties.

Results

To study the Pink1/Parkin circuit, we examined the relationship between circuit input (mitochondrial Pink1 levels) and activation (mitochondrial Parkin recruitment) on healthy mitochondria. We designed an experimental system that would: avoid confounding side effects from chemically induced mitochondrial depolarization (e.g., dysregulation of ATP, ROS and Calcium homeostasis) (*24–28*), enable precise control of mitochondrial Pink1 input levels, and allow simultaneous monitoring of circuit input and activation at a single-cell level over time.

Building a fluorescent Parkin probe to read out circuit activation.

First, we focused on establishing methods to monitor circuit activation. A long-established readout of Pink1/Parkin circuit activation and of mitophagy initiation is Parkin relocalization from the cytoplasm to the mitochondria (*2*). This relocalization is commonly monitored through N-terminally tagged Parkin fusions. However, N-terminal epitope tags have been reported to disrupt Parkin's autoinhibited state, presumably by weakening interactions involving Parkin's N-terminal UBL domain (*29*) (Fig. 2A). Because Parkin's UBL domain is also required at multiple steps of Parkin's activation (*10, 11, 13*), we hypothesized that an N-terminal tag may additionally limit Parkin's activity. To test this, untagged Parkin (Parkin[Untagged]) and Parkin, N-terminally tagged with monomeric Kusabira-Orange 2, (Parkin-mKO2[Nterm-I1]) were individually expressed in HeLa cells lacking endogenous parkin (*3, 30*), mitochondria were depolarized with 20mM carbonyl cyanide m-chlorophenyl hydrazone (CCCP), and mitophagy activity was assessed using immunofluorescence (IF; Fig. 2B-D). Compared to Parkin[Untagged], Parkin-mKO2[Nterm-I1] had significantly reduced mitochondrial parkin recruitment and significantly reduced downstream mitochondrial degradation (Fig. 2E-J; Fig. S1A-B). We hypothesized that the deficit in Parkin-mKO2[Nterm-I1] activity could be due to an M1I substitution carried over from the mCherry-Parkin source. While correction of this substitution (Parkin-mKO2[Nterm-M1]) partially restored the fusion's recruitment and mitophagy activity, a significant deficit remained (Fig. 2E-J; Fig. S1A-B).

What might cause N-terminal tag-based Parkin dysfunction? The N-terminal surface of Parkin's UBL domain has not been reported to be involved in intra- or inter-protein interactions (*12–14*, *31*). However, protein structure analysis suggests that the positively charged amino group at Parkin's N-terminus is an integral part of the UBL domain fold (Fig. S2A-C). Specifically, the NH_3^+ group of Methionine 1 (M1) appears to form a salt bridge with the negatively charged sidechain of glutamic acid 16 (E16) on an adjacent beta strand (Fig. S2C). Similar salt bridges were observed in published structures for other N-terminal UBL domains and for Ubiquitin itself (Fig. S2D-F). Further, alignment of all known UBL domain sequences revealed that negatively charged residues are conserved at an equivalent position to E16 in N-terminal, but not internal, UBL domains (Fig. S2G). Internal UBL domains, which by definition lack an NH_3^+ group, appear to have distinct stabilizing interactions (Fig. S2H-I). Finally, we found that a Parkin (E16A) mutant (Parkin[Untagged-E16A]) displayed both reduced mitochondrial recruitment and mitochondrial degradation capability (Fig. S2J). Together, these data suggest that the conserved NH_3^+-E16 interaction in Parkin's UBL domain is functionally important.

To engineer a fluorescently tagged mKO2-Parkin with preserved activity, we investigated alternative tagging locations. C-terminal tags were disregarded both due to proximity to Parkin's catalytic Ring2 domain and due to involvement of Parkin's C-terminal surface in intra-protein interactions (*14*) (Fig. S2A). Instead, we identified an internal tag insertion site, between A92 and G93, within a disordered (*12*, *13*) and non-conserved linker region (Fig. 2A-C). Characterization of the mitophagy activity of internally tagged Parkin (Parkin-mKO2[Internal]) revealed a slight but not statistically significant reduction in both mitochondrial recruitment and mitochondrial degradation activities (Fig. 2E-J; Fig. S1A-B). We note that mKO2 fluorescence is correlated with Parkin antibody staining for all constructs (Fig. S1 C-D.) Consistent with the literature, we also observed that Parkin-mKO2[Internal] recruitment to depolarized mitochondria was reversed upon mitochondrial repolarization, a well-established property of the circuit (*2*, *3*, *32*). Taken together, we chose to monitor activation of the Pink1/Parkin pathway using Parkin-mKO2[Internal].

Control of circuit input with simultaneous monitoring of circuit activation

Next, we focused on establishing methods to activate and monitor circuit input. To do so, we developed the ability to directly control the amount of Pink1 on mitochondria in HeLa cells stably expressing Parkin-mKO2[internal] (hereafter, abbreviated to Parkin-mKO2). We adapted existing approaches, allowing chemically induced recruitment of Pink1 to mitochondria (*2*, *33–35*), for quantitative live-cell microscopy (Fig. 3A-C). Specifically, Pink1's mitochondrial targeting transmembrane region (MTS; amino acids 1-110) was replaced with an FKBP domain (half of an FRB-FKBP rapamycin analog (rapalog) induced heterodimerization system) (*22*) and labeled with mNeonGreen (Pink1-mNG; Fig. 3A). Further, an FRB domain tether was fused to a MOM-targeting domain and to a SNAP tag for far-red live-cell imaging via staining with SNAP-Cell 647-SiR (mtTether-SNAP + SNAP-647; Fig. 3A-B). Pink1-mNG was expressed using a doxycycline-inducible promoter, while mtTether-SNAP was stably expressed and localized to the mitochondria (Fig. S3A-C). We verified that addition of 200nM rapalog causes the expected

sequence of: Pink1-mNG recruitment to mitochondria, Parkin-mKO2 recruitment to mitochondria, and mitochondrial aggregation/degradation (Fig. 3D; Fig. S3C-D). We further observed that: Parkin-mKO2 was not recruited in the absence of rapalog (Fig. S3B/D; Fig. 3D), mitochondria did not depolarize in response to Pink1-mNG or Parkin-mKO2 recruitment (data not shown), and MTOR signaling was not affected by rapalog treatment (Fig. S3E). We selected 12 hours post rapalog addition as our final imaging time point to give cells sufficient time to undergo Parkin-mKO2 recruitment. Finally, cell-to-cell heterogeneity of Pink1-mNG expression (Fig. S3A-B) allowed us to simultaneously observe a wide variety of mitochondrial Pink1-mNG levels in each imaging experiment. Thus, our system allowed us to control mitochondrial Pink1-mNG input and to simultaneously monitor mitochondrial Pink1-mNG and Parkin-mKO2 levels via live-cell microscopy.

Quantification of circuit dynamics

Using the rapalog-induced recruitment system, we defined an approach to quantify circuit dynamics at single-cell resolution. We quantified mitochondrial Pink1-mNG levels (*via* mean Pink1-mNG intensity) and Parkin-mKO2 recruitment (*via* Pearson correlation of Parkin-mKO2 and mtTether-SNAP intensities) (Fig. 3D-E; methods). Each cell was assigned: a circuit Pink1 input score (time-averaged mitochondrial Pink1-mNG levels before Parkin-mKO2 recruitment; methods, Fig. 3E, top); a time of circuit activation defined by when Parkin recruitment is first observed (Fig. 2E, bottom; methods); and a circuit Parkin output score (maximum observed level of Parkin-mKO2 recruitment; Fig. 3E, bottom). This triplet of scores summarized circuit behavior for each cell.

Pink1 input threshold determines decision for Parkin recruitment

These scores were first used to investigate the input-to-output relationship in this Pink1-Parkin circuit. We found that cells exhibited a bimodal distribution of Parkin output scores, having either undetectable or high Parkin responses (Fig. 3F). (Bimodality was also observed using other scores; Fig. S4.) Further, a threshold for Pink1 input scores effectively separated these low vs high Parkin output cells (Fig. 3F-G; methods). These data suggest that the Pink1-Parkin circuit displays switch-like behavior, where circuit activation is triggered if input levels surpass an input threshold.

Pink1 levels determine timing of Parkin recruitment.

Next, we examined how circuit input scores influence the detection time of Parkin recruitment. We found that the relationship between single-cell Pink1 input scores and time to detection of Parkin response was well fit by a hyperbolic model (R-squared value of 0.89; Fig. 3H; a linear model is inconsistent with the observed threshold behavior). That is, we observe a "reciprocal delay" behavior: doubling the amount by which Pink1 input exceeds the threshold will halve the time to trigger a Parkin response. Both the input threshold and reciprocal delay required of Pink1 levels for Parkin recruitment were recapitulated by fixed-cell time series (Fig. S5). Taken together, our data suggest two rules for Parkin recruitment: Pink1 input must exceed a threshold,

and Pink1 input must persist above the threshold for a duration which is inversely proportional to the amount that Pink1 input exceeds the threshold. Together, the input threshold and reciprocal delay behaviors describe a dual-factor regulatory paradigm for the Pink1/Parkin pathway.

Dual factor authentication can be perturbed

We next tested how chemical perturbations reported to broadly sensitize or desensitize the Pink1/Parkin circuit, in depolarization-based assays, would affect the activation behaviors identified here (Fig. 4A-B). To reduce pathway sensitivity, we tested the reactive oxygen species (ROS) scavenger N-acetyl-L-cysteine (NAC) (*24*). ROS generated by depolarized mitochondria is thought to promote mitophagy (*25*). We found that treatment with 2mM NAC delayed Parkin recruitment but did not alter the input threshold (Fig. 4C; Fig. S6). The direction of this effect is consistent with the published effect of NAC treatment (*24*) and confirms that NAC treatment can directly decrease Pink1/Parkin circuit sensitivity, even in the absence of overt mitochondrial depolarization, via an unknown mechanism. To increase pathway sensitivity, we tested the USP30 inhibitor CMPD39 (*36*). USP30 is thought to both limit the basal mitochondrial ubiquitin level (*6, 16, 17, 37*) and directly antagonize Pink1/Parkin circuit activation (*4, 21–23*) through removal of mitochondrial ubiquitin. In agreement with these roles, we found that treatment with CMPD39 shortened the Parkin recruitment delay and lowered the input threshold (Fig. 4D; Fig. S7).

Input-coupled amplification is sufficient to produce dual factor authentication.

We next investigated emergent behaviors of the Pink1/Parkin circuit topology using mathematical models (Fig. 5 for simplified model, supplementary text and Fig. S8 for more general model). A key feature of these models is that the input controls the feedback strength, which we refer to as "input-coupled positive feedback" (Fig. 5A-B). Mathematical analysis shows that when Pink1 levels are zero the circuit is unable to become activated and is in a stable "off" state, which makes intuitive sense as both Parkin activation and Parkin recruitment depend on Pink1 phosphorylation steps. If Pink1 levels increase, positive feedback for Parkin recruitment will be initiated above a Pink1 input threshold due to a transcritical bifurcation arising within the system (Fig. 4C-D; supplementary text) (*38–41*). Finally, if an on-state level of Parkin is defined (e.g., high enough for detection, recruitment of downstream mitophagy components, etc.), then mathematical analysis shows that the time to reach this on-state is inversely related to Pink1 levels (Fig. 4E-F; supplementary text). Thus, our conceptual models were sufficient to recapitulate the noise-tolerance properties of the Pink1/Parkin circuit observed experimentally.

Discussion

This study reveals an intrinsic mechanism by which the Pink1/Parkin circuit can avoid errant activation. We developed an approach to modulate and measure mitochondrial Pink1 input levels and simultaneously monitor Parkin localization response using a novel tagging strategy. Our input-to-response mapping confirms previous speculation that a Pink1 threshold buffers against

improper activation of the Pink1/Parkin pathway. However, our analysis also reveals that Parkin activation does not simply follow from the requirement that Pink1 levels exceed a threshold. Rather, appreciable Parkin activation is delayed for a duration of time that is inversely proportional to Pink1 levels. The dependence of circuit activation on persistence of input signal could allow mitochondria to ignore transient depolarizations yet respond to persistent depolarization or to repeated transient depolarizations (*32*). These regulatory properties can be modulated to tune the sensitivity of the pathway. Finally, mathematical modeling shows that input-coupled positive feedback—central to the Pink1/Parkin circuit topology—can intrinsically give rise to input thresholds and reciprocal delay behaviors that enable noise-tolerance.

Our study opens several questions for future investigation. First, previous hypotheses that an activation threshold for the Pink1/Parkin circuit exists and can be controlled by USP30 (*6*, *15–17*) were confirmed by our results. However, our data does not resolve whether USP30-mediated control of this threshold is via removal of Parkin-deposited Ub or of basal mitochondrial Ub. Second, a number of additional proteins have been implicated in controlling mitophagy. These proteins include the putative ubiquitin phosphatase PTEN-L (*42*), deubiquitinases such as USP15 (*43*), and Parkin-binding proteins such as Miro (*27*). It remains unclear how each of these proteins modulates Pink1/Parkin dynamics within the conceptual framework described here. Finally, depolarization-dependent signals including ROS generation, calcium release, and ATP depletion (*24–28*) have been associated with modulating Pink1/Parkin activation sensitivity. However, the mechanisms by which these signals affect the Pink1/Parkin pathway have been difficult to disentangle using traditional depolarization-based assays (*24*). The quantitative framework described in this work provides a starting point for understanding how various proteins and signals affect properties of Pink1/Parkin circuit dynamics.

Input-coupled feedback circuits not only provides control of when amplification can occur, but also where amplification can occur. For example, because the presence of input is required for maintenance of activation, Pink1/Parkin circuit activation is not expected to spread to healthy mitochondria where Pink1 is absent. When amplification is decoupled from input, as in many classic feedback systems, spatial constraint of activation may be lost without additional mechanisms (*38*, *40*). Beyond mitophagy, there are other systems where input-coupled positive feedback may be operating. In EGF receptor (EGFR) signaling, an EGFR-driven amplification loop involving GAB1, PI3K and PIP3 matches the topology of an input coupled positive feedback circuit (*44*, *45*). A similar amplification loop also exists for insulin receptor (IR) signaling via IRS, PI3K, and PIP3 (*45*). Our work motivates future studies into how input-coupled positive feedback circuits enable different cellular-decision making strategies in space and time.

Fig. 1. Pink1 provides continual input to the Pink1/Parkin positive feedback loop. Cartoon schematic of the Pink1/Parkin positive feedback loop. Phosphorylation sites, yellow circles. Parkin's Ubiquitin-like (UBL) domain and catalytic Ring 2 domain labeled to illustrate conformational changes required for Parkin's activation. In each round of amplification, 1) Pink1 phosphorylates Ubiquitin, 2) pUbiquitin recruits cytoplasmic parkin to the mitochondria, 3) Pink1 phosphorylates Parkin's UBL domain to activate Parkin, 4) pParkin deposits additional ubiquitin. Ubiquitin is attached to proteins on the MOM to form ubiquitin chains. Ubiquitin chains not shown.

Fig. 2. Identification of an internal Parkin tagging site that preserved function. (A) Domains and selected residues of Parkin. **(B)** Parkin fusion proteins, assessed for activity. LD, ligase dead. **(C)** Immunofluorescence (IF) approach for assessing Parkin function. **(D)** Conservation of residues surrounding Parkin's internal tagging site. **(E)** Parkin recruitment to mitochondria (Tom20) as a function of Parkin expression. Data are sliding medians of single-cell data shown in Fig. S1A. AFU, arbitrary fluorescence units. **(F)** Mean change in Parkin localization across Parkin expression levels from (E). Data are mean and 95% confidence intervals, calculated via bootstrap analysis. *adjusted p-value<0.05 by bootstrap analysis and Bonferroni multiple comparison adjustment. **(G)** Representative images showing Parkin

recruitment. Scale bars, 20um. **(H)** Total mitochondrial intensity as a function of Parkin expression. Data are sliding medians of single-cell data shown in Fig. S1B. **(I)** Mean mitochondrial loss across Parkin expression levels. Representation and analysis as in (F). **(J)** Representative images showing mitochondrial loss. Scale bars, 20um.

Fig. 3. Input requirements for activation of the Pink1/Parkin circuit. (A) Rapalog-induced Pink1 recruitment system. MTS, mitochondrial targeting sequence. T82L, mutation required for rapalog binding. Expression methods: Doxycycline-inducible (dox.) vs cytomegalovirus (CMV) constitutive promoters. **(B)** Live-cell imaging approach. **(C)** Cartoon of synthetic system function **(D)** Representative images showing Pink1-mNG and Parkin-mKO2 recruitment to mitochondria. Scale bars, 10um. **(E)** Quantification of Pink1/Parkin dynamics for the cell shown in (D). Co-localization, intensity correlation. Black dots, timepoints shown in (D). **(F)** Parkin output scores versus Pink1 input scores for individual cells. Datapoints colored by local point density. **(G)** Pink1 input threshold separates Parkin activated from Parkin non-activated cells.

(H) Parkin recruitment time versus Pink1 input scores for individual cells. Datapoints colored by local point density. Fitted hyperbolic curve is shown.

Fig. 4. Dissecting the effects of mitophagy modifiers on Pink1/Parkin activation. (A) Reported effects of mitophagy perturbations from the literature. Chemical structure of the USP30 inhibitor, CMPD39. **(B)** Potential effects of perturbations on pathway behavior. Combination effects also possible. **(C)** Measured threshold and delay values for live cells in NAC or vehicle conditions. Data are mean and 95% confidence intervals calculated via bootstrap analysis. *adjusted p-value<0.05 by bootstrap analysis and Bonferroni multiple comparison adjustment. Single-cell data shown in Fig. S6. **(D)** Quantified values for USP30 inhibition as in (C). Single-cell data shown in Fig. S7.

Fig. 5. Threshold and delay properties arise within a minimal model of input-coupled positive feedback. (A) Schematic of input-coupled positive feedback. **(B)** Schematic and rate equations for an input-coupled positive feedback model approximating the Pink1/Parkin circuit. X represents free Ubiquitin/Parkin and pX represents phosphorylated mitochondrial (blue) Parkin/Ubiquitin, and E represents Pink1. In the model, pX recruits X to be phosphorylated by E. Newly formed pX binds to the mitochondria. Brackets indicate concentration. Variables: a, input threshold; b, delay scaler; $[pX_{init}]$, initial seed; $[pX_{det}]$, detection point. Generalized analysis in supplement. Assumes, mass action, steady state, $[X_{tot}] = [X] + [pX]$, and $[pX_{det}] \ll [X_{tot}]$. **(C)** Rate analyses for values of [E] below or above the input threshold. Recruitment and removal rates from B are plotted. SSS, stable steady state; USS, unstable steady state. **(D)** Steady state analysis illustrating the transcritical bifurcation, TCB. Examples from (C), marked with empty circles. **(E)** Exponential growth of [pX] from $[pX_{init}]$ to $[pX_{det}]$ for various values of [E]. Equation

describing hyperbolic relationship between [E] and time to reach [pX$_{det}$] is shown. **(F)** Graphed relationship between [E] and time to reach [pX$_{det}$]. Examples from (E), marked with empty circles.

Fig. S1. Comparison of Parkin tagging strategies. (A & B) Single-cell data used to generate Fig. 2E-J. Points, single cells; contours, datapoint density; red lines, sliding medians. Single-cell data from three experimental replicates was pooled. DMSO: n=~60,000 cells/condition; CCCP:

n=~150,000 cells/condition; ~35% of cells had Parkin expression within the range used to calculate the moving medians, 3000-20000 AFU/px. **(C)** mKO2 fluorescence versus anti-Parkin immunostaining intensity. Data are sliding medians of single cell data. **(D)** Top: representative images showing mKO2 fluorescence and anti-Parkin immunostaining. Parkin-mKO2[Internal], shown. Bottom: pixel intensity traces along line overlaid on images. Line is 33um long for scale.

Fig. S2. Structural importance of Parkin's N-terminus. (A-C) Human Parkin structure. **(A)** Full protein, colored by domain. **(B)** UBL domain colored blue to red, N to C terminus. **(C)** N-terminus (N.t.) of UBL domain. Carbons colored as in B; Oxygen, red; Nitrogen, blue; sulfur, yellow. Dotted lines, predicted interactions. # indicates predicted salt bridge between E16 and the N-terminal NH_3^+, in all panels. **(D-F)** N.t. structures of Ubiquitin and other N-terminal UBL domains. **(G)** Sequence logo showing amino acid conservation in UBL domains. Analysis done

across all unique UBL domains from all species in the Uniprot database. Residues used for motif-based alignment are marked. **(H/I)** N.t. structures of internal UBL domains. Carbons of additional N-terminal residues colored gray. Asterisks, potential stabilizing interactions; dashed circle, potential nonpolar interaction. **(J)** Functional quantification of untagged Parkin with an E16A mutation (Parkin$^{Untagged-E16A}$) as in Fig. 2D-J. Data collected and analyzed alongside data in Fig. 2D-J. Data are mean and 95% confidence intervals calculated via bootstrap analysis. *adjusted p-value<0.05 by bootstrap analysis and Bonferroni multiple comparison adjustment.

Fig. S3. Characterization of the inducible Pink1/Parkin circuit. (A) Cumulative distribution of single-cell Pink1 expression levels following 48h doxycycline treatment. **(B)** Representative images showing population-level expression levels. Scale bar, 50um. **(C)** Top: Representative cell showing colocalization between MtTether-SNAP stained with SNAP-647 and mitochondria immunostained for Tom20. Bottom: pixel intensity values along white line in the single-channel images above. Scale bar in merged channel image, 20um. **(D)** Representative images showing mitochondrial loss following prolonged rapalog-driven Pink1-mNG and Parkin-mKO2 recruitment. Asterisk indicates cell of interest. Dotted line in bottom right image indicates cell boundary. Scale bars, 10um. **(E)** Western blot comparison of phosphorylated S6 (top pair) to unphosphorylated S6 (bottom pair) following treatment with either rapamycin or rapalog.

Fig. S4. Alternative Pink1/Parkin quantification methods also reveal output bimodality. (A-B) Alternative quantifications of mitochondrial Pink1 levels or mitochondrial Parkin recruitment for single cell data shown in Fig. 3F. **(A)** Pathway activation estimated as maximum achieved fraction of total Parkin-mKO2 localized on mitochondria. Datapoints colored by local density. Pink1 input threshold from Fig. 3F shown as a vertical black line. **(B)** Histogram showing ability of input threshold in (A) to separate Parkin activation populations. **(C)** Pathway input estimated as Pink1 expression level. An input threshold for this metric was recalculated and is shown as a vertical black line. **(D)** Histogram showing ability of input threshold in (C) to separate Parkin activation populations.

Fig. S5. Fixed-cell data recapitulates the threshold and reciprocal delay behaviors required for activation of the Pink1/Parkin circuit. (A) Approach for quantifying Pink1-mNG and Parkin-mKO2 dynamics via fixed-cell timeseries. **(B)** Mitochondria-Parkin co-localization versus mitochondrial Pink1 levels for individual cells. Single-cell data from three experimental replicates was pooled. A selection of treatment durations are shown. Datapoints colored by local point density. Vertical black lines denote separation values identifying the mitochondrial Pink1-mNG level capable of driving Parkin-mKO2 recruitment for that treatment duration. Relative

datapoint scarcity at 24h due to mitochondrial degradation. **(C)** Calculated separation values versus rapalog treatment durations. Datapoints from (B). Fitted hyperbola, solid red line; hyperbola's limit, dashed red line. Relationship to single cells: cells with mitochondrial Pink1 will undergo detectable Parkin translocation when the separation value drops below their mitochondrial Pink1 level. **(D)** The hyperbola, in (C), rearranged to the form used for live-cell data in Fig. 3H.

Fig. S6. Dissecting the effects of NAC on Pink1/Parkin circuit dynamics. (A) Experimental schematic for measurement of Pink1/Parkin circuit dynamics in live cells following ROS modulation. All treatments were maintained through subsequent steps. **(B/D)** Parkin activation scores versus Pink1 input scores of individual cells. Single-cell data from two experimental replicates was pooled. Calculated threshold is marked. Single-cell datapoints, colored by local point density. **(C/E)** Time of Parkin recruitment vs Pink1 input scores of individual cells. Single-cell datapoints, colored by local point density. Fitted Hyperbolas are shown with R-squared values. Thresholds from (B/D) are also shown.

Fig. S7. Dissecting the effects of USP30 inhibition on Pink1/Parkin circuit dynamics. (A) Experimental schematic for measurement of Pink1/Parkin circuit dynamics in live cells following USP30 inhibition. All treatments were maintained through subsequent steps. **(B/D)** Parkin activation scores versus Pink1 input scores of individual cells. Single-cell data from three experimental replicates was pooled. Calculated threshold is marked. Single-cell datapoints, colored by local point density. **(C/E)** Time of Parkin recruitment vs Pink1 input scores of individual cells. Single-cell datapoints, colored by local point density. Fitted Hyperbolas are shown with R-squared values. Thresholds from (B/D) are also shown.

A

B

$$U_f \underset{k_3}{\overset{k_1}{\rightleftarrows}} U_m \underset{k_5}{\overset{k_4E}{\rightleftarrows}} U_p \underset{k_7}{\overset{k_6A_f}{\rightleftarrows}} A_b \underset{k_9}{\overset{k_8E}{\rightleftarrows}} A_p$$

with k_2U_f arcing from U_m to A_p.

Fig. S8. Parameterization of Pink1/Parkin circuit for mathematical model analysis in supplemental text. (A) Parameterized model of the Pink1/Parkin circuit. Variables are defined: Parkin, A; Ubiquitin, U; Pink1, E. Rate constants "k" are marked for each step. **(B)** Simple schematic of the model. Rate of each step obtained by multiplying the source moiety (e.g. U_f) by the rate shown (e.g. k_1)

Supplementary text

MATHEMATICAL MODEL ANALYSIS

<u>General framework</u>

<u>Assumptions and conclusions</u>

We consider two models of increasing complexity for the dynamics in the system describing Pink1–Parkin–Ubiquitin in a Mitochondrion discussed in the main body of the paper. Simplifying somewhat, we may say that in the experiment the amount $P(t)$ at time t of Parkin in a mitochondrion is tracked as it grows until it reaches a certain threshold level P_{th}. The initial amount of Parkin P_{in} is assumed to be much smaller than the threshold level P_{th}. This yields a measurement of the time T it takes the Parkin level to grow from P_{in} to P_{th}. The level E of Pink1 affects the rate at which Parkin increases and thus the experiment gives us a measurement of the growth time T as a function of the Pink1 level E.

The main assumptions concerning the system that we make are

i. The system contains a positive feedback mechanism

ii. Pink1 acts as a catalyst: some of the reaction rates in the system increase when the amount of Pink1 is increased

iii. The initial amounts of Parkin and Ubiquitin on the Mitochondria are much smaller than the threshold amount

iv. The rate at which free Ubiquitin spontaneously moves to the mitochondrion so small that it may be ignored

v. The amounts of free Ubiquitin and Parkin are large and may be considered constant

These assumptions lead to the following consequences:

<u>Exponential Growth:</u> The amount of Parkin grows (or decays) exponentially; if $P(t)$ is the amount of Parkin at time t, then the models predict $P(t) \approx P_{in}e^{\lambda_E t}$, where the growth rate λ_E depends on the many reaction rates in the model, and in particular on the amount E of Pink1 in the system.

<u>Switch-like behavior:</u> The system exhibits switch-like behavior in its dependence on E: there is a critical value E_* such that $\lambda_E < 0$ when $E < E_*$ and such that $\lambda_E > 0$ when $E > E_*$. This means that if the Pink1 amount is below E_* the Parkin levels will decrease exponentially, and if the Pink1 amount exceeds E_* then Parkin grows exponentially.

<u>Time to detection decreases with increasing Pink1 levels:</u> The exponent λ_E is an increasing function of the Pink1 level E. When $\lambda_E > 0$ the Parkin level grows exponentially according to

$$P(t) \approx P(0)e^{\lambda_E t}.$$

Let P_{th} be the threshold level at which Parkin is detected, and assume that the initial amount P_{in} of Parkin is small compared to P_{th}. If the growth rate λ_E is negative then the amount $P(t)$ of Parkin will only decay and thus never reach the threshold level. On the other hand, if the growth rate λ_E is positive, then the threshold time $T_{\text{th}}(E)$ follows from

$$(1) \quad P(T_{\text{th}}) = P_{\text{th}} \iff P_{\text{in}}e^{\lambda_E T_{\text{th}}} = P_{\text{th}} \iff T_{\text{th}} = \frac{\ln(P_{\text{th}}/P_{\text{in}})}{\lambda_E}.$$

For each of the models we find that the growth rate λ_E increases when E is increased, which therefore implies that the time to detection T_{th} is a decreasing function of E.

<u>General form of the models</u>

In each of the models we have a vector

$$X(t) = \begin{bmatrix} x_1(t) \\ \vdots \\ x_n(t) \end{bmatrix}$$

containing the total amounts of the components that the model tracks, such as Parkin, Ubiquitin, (phosphorelated or not), or combinations of these. Using mass action kinetics we arrive at a system of differential equations governing the time dependence of $X(t)$. Assuming that we only consider the system when the amounts of non-free Parkin and Ubiquitin are small, we arrive at a linear system

$$(2) \quad X'(t) = \mathcal{M}_E X(t),$$

where

- E is the amount of Pink1 in the system; this quantity is assumed to be kept constant;

- $\mathcal{M}_E$ is a matrix containing the reaction rates, and which can be further decomposed as

$$\mathcal{M}_E = \mathcal{M}_0 + E\mathcal{M}_1.$$

The matrix $\mathcal{M}_0$ contains the rates at which reactions take place in the absence of Pink1, while $\mathcal{M}_1$ accounts for the change in the reaction rates caused by the presence of Pink1.

<u>Eigenvalue analysis of the models</u>

Using the method of eigenvalues and vectors one shows that the general solution of a linear equation such as (Eq. 2) is a superposition of exponentially growing or decaying terms, i.e.

(3) $\quad X(t) = e^{\lambda_1 t} V_1 + e^{\lambda_2 t} V_2 + \cdots + e^{\lambda_n t} V_n$

in which $\lambda_1, \ldots, \lambda_n$ are the eigenvalues of the matrix $\mathcal{M}_E$, $V_1, \ldots, V_n$ are corresponding eigenvectors. Since we are interested in the time it takes $X(t)$ to grow from a small initial amount to a detectable threshold value, we want to consider the fastest growing term(s) in (Eq. 3), i.e. the terms corresponding to the largest eigenvalues λ_i. In studying the eigenvalues $\lambda_1, \ldots, \lambda_n$ we note that in all versions of our model the matrix $\mathcal{M}_E$ has non-negative off-diagonal entries, and is irreducible. This implies, by the Perron-Frobenius theorem, that $\mathcal{M}_E$ has a unique dominant eigenvalue λ_E, i.e.

i. λ_E is a real eigenvalue of $\mathcal{M}_E$ (not a complex eigenvalue)

ii. every other (possibly complex) eigenvalue μ of $\mathcal{M}_E$ satisfies $\mathrm{Re}\,\mu < \lambda_E$

iii. corresponding to the eigenvalue λ_E, the matrix $\mathcal{M}_E$ has positive left and right eigenvectors W_E, V_E respectively; by definition these satisfy

$$\mathcal{M}_E V_E = \lambda_E V_E, \qquad \mathcal{M}_E^\top W_E = \lambda_E W_E.$$

They can be normalized so that $\langle W_E, V_E \rangle = 1$.

The dominant eigenvalue tells us the largest exponential rate with which solutions to (Eq. 2) can grow. More precisely, the eigenvalue decomposition (Eq. 3) contains one term corresponding to the dominant eigenvalue λ_E. If we denote this term by $x(t)V_E$ and group the remaining terms into a slower growing component $Y(t)$ then we have

(4) $\quad X(t) = x(t)V_E + Y(t), \qquad \langle W_E, Y(t) \rangle = 0.$

The left-eigenvector W_E allows one to find the coefficient $x(t)$ from the vector $X(t)$ via

$$x(t) = \langle W_E, X(t) \rangle.$$

By taking the inner product with W_E on both sides in (Eq. 2) one finds that the V_E component of X satisfies an ordinary differential equation

(5) $\quad x'(t) = \lambda_E x(t),$

whose solution can be written as

(6) $\quad x(t) = e^{\lambda_E t} x(0)$

To compute the time to detection we assume that Parkin is detected when $x(t)$ reaches a specific threshold value x_{th}. Then the time to detection is

$$T_{\text{th}}(E) = \frac{1}{\lambda_E} \ln \frac{x_{\text{th}}}{x(0)}.$$

The minimal Pink-Parkin model

In our simplest model we only keep track of the Parkin in the system, assuming it exists in one of two states: on the mitochondria or free (not on the mitochondria).

Parkin on the mitochondria can bind free Parkin and this process is aided by Pink1 as an enzymatic catalyst. Parkin on the mitochondria also spontaneously leaves the mitochondria.

Our model keeps track of the following quantities:

A_t total amount of Parkin in the system; a constant

A_m amount of Parkin on the mitochondria

A_f amount of free Parkin, $A_f + A_m = A_t$

E amount of Pink1 in the system; constant in time

Since A_t and E are time independent and since A_m and A_f are constrained by $A_m + A_f = A_t$, the time evolution of the system is completely determined by that of A_m . The following differential equation takes both recruitment and degradation into account:

$$\frac{dA_m}{dt} = k_{\text{fb}}E(A_t - A_m)A_m - k_{\text{off}}A_m.$$

Here k_{fb} and k_{off} are reaction constants.

We make one further simplifying assumption, namely, during the observations in the experiment, the total amount A_t of Parkin is much larger than the amount A_m on the mitochondria. We may therefore replace $A_t - A_m$ by A_t, which leads us to the differential equation

(7) $$\frac{dA_m}{dt} = k_{\text{fb}}EA_tA_m - k_{\text{off}}A_m = (k_{\text{fb}}EA_t - k_{\text{off}})A_m.$$

The coefficient $k_{\text{fb}}EA_t - k_{\text{off}}$ is constant in time, so this differential equation is of the type $\frac{dx}{dt} = kx$, and its solution is given by the exponential growth formula $x(t) = e^{kt}x(0)$. In terms of A_m we get

$$A_m(t) = e^{(k_{\text{fb}}EA_t - k_{\text{off}})t}A_m(0).$$

Indeed, equation (Eq. 7) is of the form (Eq. 2), if lets $X(t)$ be the vector with only one component $X(t) = [A_m(t)]$, and if one lets $\mathcal{M}_E$ be the 1×1-matrix $\mathcal{M}_E = [k_{\text{fb}}EA_t - k_{\text{off}}]$. The dominant (and only) eigenvalue of $\mathcal{M}_E$ is $\lambda_E = k_{\text{fb}}EA_t - k_{\text{off}}$. However, since both the vector $X(t)$ and the matrix $\mathcal{M}_E$ only have one component the eigenvalue analysis is not needed to solve the differential equation (Eq. 7).

In the experiment one begins with a given small amount $A_m(0)$ of Parkin and measures how long it takes before the amount $A_m(t)$ of Parkin reaches a fixed detectable level, A_{th}. By solving the equation $A_m(t) = A_{th}$ for t we find

$$T_{th}(E) = \frac{\ln \frac{A_{th}}{A_m(0)}}{k_{fb}A_t E - k_{off}} \ .$$

The critical value of E at which the systems "switches on" is

$$E_* = \frac{k_{off}}{k_{fb}A_t}.$$

The full model

The nonlinear model

The following variables appear in the full model

- U_f, U_m, U_p The free, mitochondrial, and phosphorelated forms of Ubiquitin

- A_f, A_b, A_p The free, bound, and phosphorelated and bound forms of Parkin

- E The amount of Pink1 present in the system; a constant.

Assuming mass-action kinetics, these variables evolve according to the following set of differential equations (see model schematic in Fig. S8B).

$$\frac{dU_f}{dt} = -k_1 U_f - k_2 U_f A_p + k_3 U_m$$
$$\frac{dU_m}{dt} = k_1 U_f + k_2 U_f A_p - k_3 U_m - k_4 E U_m + k_5 U_p$$
$$\frac{dU_p}{dt} = k_4 E U_m - k_5 U_p - k_6 U_p A_f + k_7 A_b$$
$$\frac{dA_f}{dt} = -k_6 U_p A_f + k_7 A_b$$
$$\frac{dA_b}{dt} = k_6 U_p A_f - k_7 A_b - k_8 E A_b + k_9 A_p$$
$$\frac{dA_p}{dt} = k_8 E A_b - k_9 A_p$$

Linearization assuming abundant free Parkin and Ubiquitin

We can simplify the model by assuming that A_f and U_f are nearly constant because free Parkin and Ubiquitin are abundantly present. This leads to a reduced system with four components U_m, U_p, A_b, A_p, which satisfy the following four linear differential equations:

$$
\begin{aligned}
\frac{dU_m}{dt} &= -(k_3 + k_4 E)U_m & &+k_5 U_p & &+k_2 U_f A_p \\
\frac{dU_p}{dt} &= k_4 E U_m - (k_5 + k_6 A_f)U_p & &+k_7 A_b & & \\
\frac{dA_b}{dt} &= & &+k_6 A_f U_p - (k_7 + k_8 E)A_b & &+k_9 A_p \\
\frac{dA_p}{dt} &= & &k_8 E A_b & &-k_9 A_p
\end{aligned}
$$

(8)

This linear system is of the form $X'(t) = \mathcal{M}_E X(t)$ (see (Eq. 2)) where

$$
X(t) = \begin{bmatrix} U_m \\ U_p \\ A_b \\ A_p \end{bmatrix}
$$

The matrix in this linear system is

$$
\mathcal{M}_E = \begin{bmatrix}
-k_3 - k_4 E & k_5 & 0 & k_2 U_f \\
k_4 E & -k_5 - k_6 A_f & k_7 & 0 \\
0 & k_6 A_f & -k_7 - k_8 E & k_9 \\
0 & 0 & k_8 E & -k_9
\end{bmatrix} = \mathcal{M}_0 + E\mathcal{M}_1
$$

where

$$
\mathcal{M}_0 = \begin{bmatrix}
-k_3 & k_5 & 0 & k_2 U_f \\
0 & -k_5 - k_6 A_f & k_7 & 0 \\
0 & k_6 A_f & -k_7 & k_9 \\
0 & 0 & 0 & -k_9
\end{bmatrix}
\qquad
\mathcal{M}_1 = \begin{bmatrix}
-k_4 & 0 & 0 & 0 \\
k_4 & 0 & 0 & 0 \\
0 & 0 & -k_8 & 0 \\
0 & 0 & k_8 & 0
\end{bmatrix}
$$

The dominant eigenvalue of $\mathcal{M}_E$ is the largest real root of the characteristic equation

$$
\begin{vmatrix}
-k_3 - k_4 E - \lambda & k_5 & 0 & k_2 U_f \\
k_4 E & -k_5 - k_6 A_f - \lambda & k_7 & 0 \\
0 & k_6 A_f & -k_7 - k_8 E - \lambda & k_9 \\
0 & 0 & k_8 E & -k_9 - \lambda
\end{vmatrix} = 0.
$$

Theorem. Assume $k_2, k_4, k_5, k_6, k_7, k_8, k_9, U_f$, and A_f are positive constants. Then

(a) The dominant eigenvalue λ_E is a continuous function of E for $E \geq 0$.

(b) When $E = 0$, one has $\lambda_E < 0$

(c) For $E \to \infty$ the dominant eigenvalue converges to a positive limiting value $\lambda_+^\infty > 0$.

(d) There is a unique positive number E_* such that $\lambda_{E_*} = 0$ and for all $E \geq E_*$ one has $\frac{d\lambda_E}{dE} > 0$.

Proof. The dominant eigenvalue is a simple eigenvalue and therefore depends continuously on the parameters in the matrix $\mathcal{M}_E$. Statement (a) follows.

To verify (b), we set $E = 0$ and compute

$$\det(\mathcal{M}_E - \lambda) = \det(\mathcal{M}_0 - \lambda)$$

$$= \begin{vmatrix} -k_3 - \lambda & k_5 & 0 & k_2 U_f \\ 0 & -k_5 - k_6 A_f - \lambda & k_7 & 0 \\ 0 & k_6 A_f & -k_7 - \lambda & k_9 \\ 0 & 0 & 0 & -k_9 - \lambda \end{vmatrix}$$

$$= (\lambda + k_3)(\lambda + k_9)\left(\lambda^2 + (k_5 + k_7 + k_6 A_f)\lambda + k_5 k_7\right)$$

It follows that when $E = 0$ the eigenvalues of the matrix $\mathcal{M}_E$ are

$$\lambda_1 = -k_3, \quad \lambda_2 = -k_9,$$

$$\lambda_3, \lambda_4 = \frac{-k_5 - k_7 - k_6 A_f \pm \sqrt{(k_5 - k_7)^2 + 2(k_5 + k_7)k_6 A_f + (k_6 A_f)^2}}{2}$$

All four eigenvalues are real and negative. The dominant eigenvalue λ_E is the largest of these,

$$\lambda_E = \max\{\lambda_1, \lambda_2, \lambda_3, \lambda_4\} \text{ if } E = 0$$

and it follows that for $E = 0$ one has $\lambda_E < 0$.

We turn to (c). To analyze the eigenvalues for large values of E we write the characteristic polynomial as

(9) $\quad \det(\mathcal{M}_E - \lambda) = D_0(\lambda) + D_1(\lambda)E + D_2(\lambda)E^2$

where D_0, D_1, D_2 are polynomials in λ, which upon computation turn out to satisfy

$$D_0(\lambda) = \det(\mathcal{M}_0 - \lambda)$$
$$= (k_3 + \lambda)(k_9 + \lambda)\left(\lambda^2 + (k_5 + k_7 + k_6 A_f)\lambda + k_5 k_7\right)$$
$$= \lambda^4 + \text{lower order terms},$$

$$D_1(\lambda) = \lambda\left(k_4(\lambda + k_6 A_f + k_7)(\lambda + k_9) + (\lambda + k_3)(k_8\lambda + k_5 k_8 + (k_8 - k_7)k_7 A_f)\right)$$
$$= (k_4 + k_8)\lambda^3 + \left[k_4(k_7 + k_9) + (k_3 + k_5)k_8 + (k_4 + k_8 - k_7)k_6 A_f\right]\lambda^2$$
$$\quad + (k_3 k_5 k_8 + k_3(k_8 - k_7)k_6 A_f)\lambda$$
$$= (k_4 + k_8)\lambda^3 + \text{lower order terms},$$

and

$$D_2(\lambda) = k_4 k_8 (\lambda^2 + k_6 A_f \lambda - k_2 U_f k_6 A_f)$$
$$= k_4 k_8 \lambda^2 + \text{lower order terms}$$

For large values of E the four eigenvalues of $\mathscr{M}_E$ can be separated into those eigenvalues λ for which $\lambda \ll E$, and those for which λ is comparable to E or larger.

If $|\lambda| \sim E$ or $|\lambda| \gg E$ then the dominant terms in the characteristic polynomial (Eq. 9) are those that contain $\lambda^4, \lambda^3 E, \lambda^2 E^2$. Two eigenvalues are therefore approximated by the nonzero roots of

$$\lambda^4 + (k_4 + k_8)\lambda^3 E + k_4 k_8 \lambda^2 E^2 = 0.$$

This yields two very negative eigenvalues

$$\lambda_1 \approx -k_4 E, \quad \lambda_2 \approx -k_8 E.$$

If on the other hand $|\lambda| \ll E$ then $D_2(\lambda)E^2$ is the dominant term in the characteristic polynomial (Eq. 9), and thus two of the eigenvalues are close to the roots of $D_2(\lambda) = 0$, i.e. $\lambda^2 + k_6 A_f \lambda - k_2 U_f k_6 A_f = 0$, which are given by

$$\lambda_{\pm}^{\infty} = \frac{-k_6 A_f \pm \sqrt{(k_6 A_f)^2 + 4 k_2 U_f k_6 A_f}}{2}.$$

Of these, λ_{-}^{∞} is negative and λ_{+}^{∞} is positive. Since λ_{+}^{∞} is the only positive eigenvalue, it is the dominant eigenvalue.

The dominant eigenvalue therefore satisfies

$$\lambda_E \approx \lambda_{+}^{\infty} = \frac{-k_6 A_f + \sqrt{(k_6 A_f)^2 + 4 k_2 U_f k_6 A_f}}{2} > 0.$$

This implies statement (c).

We conclude by verifying (d). Since λ_E is a simple eigenvalue, it is a differentiable function of E and its derivative is given by

$$\frac{d\lambda_E}{dE} = \left\langle W_E, \frac{d\mathscr{M}_E}{dE} V_E \right\rangle = \langle W_E, \mathscr{M}_1 V_E \rangle$$

where W_E and V_E are the left and right eigenvalues normalized by $\langle W_E, V_E \rangle = 1$.

If we write

$$W_E = \begin{bmatrix} w_1 \\ w_2 \\ w_3 \\ w_4 \end{bmatrix} \text{ and } V_E = \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{bmatrix}$$

then

(10) $\langle W_E, \mathcal{M}_1 V_E \rangle = k_4(w_2 - w_1)v_1 + k_8(w_4 - w_3)v_3.$

Since $v_1, v_3 > 0$ it suffices to show that $w_2 > w_1$ and $w_4 > w_3$. Expanding the eigenvalue equation $\mathcal{M}_E^\top W_E = \lambda_E W_E$ and rearranging terms we get

$$-k_3 w_1 + k_4 E(w_2 - w_1) = \lambda_E w_1$$
$$k_5(w_1 - w_2) + k_6 A_f(w_3 - w_2) = \lambda_E w_2$$
$$k_7(w_2 - w_3) + k_8 E(w_4 - w_3) = \lambda_E w_3$$
$$k_2 U_f w_1 + k_9(w_3 - w_4) = \lambda_E w_4$$

Using the assumption that $\lambda_E \geq 0$ we conclude from the first equation

$$k_4 E(w_2 - w_1) = \lambda_E w_1 + k_3 w_1 \geq k_3 w_1 > 0 \implies w_2 - w_1 > 0.$$

The second equation then implies

$$k_6 A_f(w_3 - w_2) = \lambda_E w_2 + k_5(w_2 - w_1) > 0 \implies w_3 - w_2 > 0.$$

Finally the third equation leads to

$$k_8 E(w_4 - w_3) = \lambda_E w_3 + k_7(w_3 - w_2) > 0 \implies w_4 - w_3 > 0.$$

It follows that the terms on the right in (Eq. 10) are positive, and hence $\frac{d\lambda_E}{dE} > 0$.

To complete the proof we note that we have shown that the continuous function $E \mapsto \lambda_E$ satisfies $\lambda_0 < 0$ and $\lambda_E > 0$ for all large enough E. Therefore there is an $E_* > 0$ at which $\lambda_{E_*} = 0$. Since λ_E is strictly increasing when $E > 0$ there cannot be two different values E_*, E_{**} at which $\lambda_{E_*} = \lambda_{E_{**}} = 0$.

Acknowledgements:

We thank all of the member of the Altschuler-Wu lab for their support. We also thank Jay Debnath, Orion Weiner, and Matt Jacobson for helpful feedback and perspectives during this study. We thank the UCSF HDFCCC Laboratory for Cell Analysis Shared Resource Facility for training and assistance with cell sorting on the BD FACS Aria 2. The HDFCCC Laboratory for Cell Analysis Shared Resource Facility is supported through a grant from the National Institute of Health (P30CA082103). Protein structure analyses performed via UCSF ChimeraX, developed by the Resource for Biocomputing, Visualization, and Informatics at the University of California, San Francisco. ChimeraX development supported by National Institutes of Health R01-GM129325 and the Office of Cyber Infrastructure and Computational Biology, National Institute of Allergy and Infectious Diseases.

Funding:

Author contributions:

Conceptualization: CSW, SJA, LFW

Methodology: CSW, SBA, SJA, LFW

Investigation: CSW

Analysis: CSW

Visualization: CSW

Modeling: CSW, SBA

Funding acquisition: CSW, SJA, LFW

Supervision: SJA, LFW

Writing – original draft: CSW, SBA

Writing – review & editing: CSW, SBA, SJA, LFW

Competing interests: SJA and LFW are founders and scientific advisory board members of Nine Squared Therapeutics.

Data and materials availability: Upon scientific review, code will be deposited on Zenodo. Due to their size, raw image datasets will be made available by SJA and LFW upon request. Plasmids containing mNeonGreen will be available through the Allele Biotechnologies depository due to a material transfer agreement limiting distribution of mNeonGreen. All other plasmids generated for this study will be made available by SJA and LFW upon completion of a material transfer agreement.

Materials and Methods:

Plasmid generation

All plasmid for expressing variants of tagged Parkin (pCMV-Parkin, pCMV-Parkin(E16A), pCMV-mKO2-Parkin, pCMV-mKO2-Parkin(C431N), pCMV-mKO2-Parkin(I1M), and pCMV-Parkin-A92mKO2) were derived from the pCMV-mCherry-Parkin (Addgene, #23956). The mKO2 insert was derived from pLL3.7m-Clover-Geminin(1-110)-IRES-mKO2-Cdt(30-120) (Addgene, #83841). Lentiviral Parkin expression vector, pLv-CMV-Parkin(A92mKO2), and Lentiviral tether expression vector pLV-EF1a-SNAP-FRB-FIS1(MTS) were created using the backbone from pLV-EF1a-IRES-Puro (Addgene, #85132). The SNAP tag insert was derived from pSNAPf (NEB, N9183S) and the FRB-Fis1 insert was derived from pC4-RhE-FRB-Fis1. The Lentiviral doxycycline-inducible FKBP-Pink1(d1-110)-mNeonGreen expression vector, pLv-TetOn-FKBP-Pink1(d1-110)-mNG-Neo, was created using the backbone from pLv-TetOn-MCS1-P2A-MCS2 (Neo) (Addgene, #89180). The FKBP insert was derived from pC4M-F2E-GFP-FKBP (Addgene, #68058), the Pink1 insert was derived from pCMVTNT-Pink1-C-Myc (Addgene, #13314) and the mNeonGreen insert was obtained as cDNA (Allele Biotech, ABP-FP-MNEONSB) (*46*). All plasmid assembly was done using standard cloning methods. All plasmids were grown following transformation into NEB STBL competent *E. coli* (NEB, C3040H). All inserts were verified via sanger sequencing (Elim Biopharmaceuticals).

Cell culture

HeLa cells were cultured with a base media consisting of RPMI (Thermo Fisher Scientific, 11875119) with 10% fetal bovine serum (Gemini Bio., #100-106) and antibiotic/antimycotic (Thermo Fisher Scientific, 15240062). HEK293T cells were cultured with a base media consisting of DMEM (Thermo Fisher Scientific, 11965092) with 10% fetal bovine serum and penicillin/streptomycin (Corning, 30-002-CI). Immunofluorescence and live-cell imaging experiments were performed in 96 well glass bottom, tissue culture treated imaging plates (Thermo Scientific Nunc, 164588). Cells were kept at 37C and 5% CO2 during culturing and during live cell microscopy.

Immunofluorescence

Cells were fixed in PBS (Invitrogen, AM9625) with 4% paraformaldehyde (Electron Microscopy Sciences, 15710) for 20 minutes at room temperature and were then permeabilized in PBS with 0.5% Triton X-100 (Sigma-Aldrich, 93443) for 20 minutes. Cells and then washed three times in PBS and blocked in PBS with 0.3% Triton X-100 and 3% Bovine Serum Albumin (BSA) (Fisher bioreagents, BP9703) for 1 hour. Blocked cells were Incubated in staining buffer (PBS with 0.3% Triton X-100 and 1% BSA) with primary antibody overnight at 4C when applicable. Cells were then washed three times with PBS and incubated in staining buffer with secondary antibody for 2 hours at room temperature when applicable. Fluorescently conjugated primary antibody was included in the secondary antibody stain when applicable. Finally, cells were washed

twice in PBS, stained with 1:5000 hoechst 33342 (Invitrogen, H3570) for ten minutes, and washed once more in PBS. Primary antibodies: 1:500 ms-anti-Parkin (Abcam, ab77924); 1:1000 Rabbit anti-Tom20 Alexa Fluor (AF) 647 (Abcam, ab209606). Secondary antibodies: 1:1000 Goat anti-mouse AF plus 488 (Invitrogen, A32731).

Activity assessment of Parkin tag variants

HeLa cells were seeded in 96 well imaging plates and allowed to grow for 36 hours. Individual wells were transfected with 25ng of plasmid expressing one tagged Parkin variant using a Lipofectamine 3000 kit (Invitrogen, L3000-015) via manufacturer's instructions. Media on transfected cells was replaced after 3.5 hours and cells were allowed to recover. Wells were treated with either 20uM carbonyl cyanide 3-chlorophenylhydrazone, (CCCP; Sigma Aldrich, C2759) or 2:5000 DMSO (Sigma Life Science, D2650) control for either 10 hours or 30 minutes with timing such that cells were fixed 48 hours after transfection. Cells were fixed and subjected to immunofluorescence staining for Parkin and Tom20. Cells were imaged on an Operetta automated microscope in confocal mode with a 20x water objective, no pixel binning, and standard filter sets.

Cell line generation

Lentivirus was made using HEK293T cells and tittered for 10% transduction efficiency. HeLa cells were co-transduced with lentivirus delivering MtTether-SNAP: EF1a-SNAP-FRB-Fis1(MTS), CMV-Parkin(A92-mKO2), and TetOn-FKBP-Pink1(110-581)-mNeonGreen using 8ug/mL polybrene in growth media. Cells were selected for the TetOn Pink1 expression cassette via treatment with 400ug/mL Geniticin (Life Tech. Corp., 10131035). Triple positive cells were isolated using fluorescence-activated cell sorting (FACS). Prior to FACS, Pink1-mNG expression was induced with 50ng/mL Doxycycline Hydrochloride (Sigma-Aldrich, D3072) for 48 hours and MtTether-SNAP was stained with 1:2000 SNAP-647 SiR (NEB, s9102) for 30 minutes. Triple positive cells were then allowed to expand and recover in the absence of doxycycline, to eliminate Pink1-mNG expression prior to being frozen for storage.

Live imaging of chemically induced recruitment of Pink1 and Parkin to mitochondria

Triple positive HeLa cells were seeded in 96 well imaging plates with 250 ng/mL doxycycline. Pink1-mNG expression was induced for 48 hours with media and doxycycline. Media with doxycycline was refreshed after 24 hours due to chemical instability of doxycycline in aqueous solution. All stain, wash, and imaging medias used in subsequent steps were growth media containing doxycycline. Prior to imaging, cells were stained with 1:1000 SNAP-AF647-SiR for 30 minutes, stained with 1:1000 hoechst 33342 for 10 minutes and washed 2x20 minutes. Hoechst staining of live HeLa cells was observed to have no effect on cell viability or growth rate. Cells were imaged in growth media once (~30 minutes) to capture the pre-treatment state. Cells were then treated with 200nM rapalog (A/C heterodimerizer, previously AP21967) (Takara Bio., 635055) and immediately subjected to live cell microscopy with images being taken every 30 minutes for 12 hours. For live cell experiments longer than 12 hours, imaging frequency was every

45 minutes. Cells were imaged using an Operetta microscope in confocal mode with a 40x water objective, 2x2 pixel binning, and standard filter sets. The exact imaging time for each was recorded.

Modulation of threshold and rate with ROS and USP30 inhibitor

Triple positive HeLa cells were seeded, treated with Doxycycline for 48 hours, stained, and imaged as with the live cell imaging of the synthetic system described above. A 2-hour pretreatment with 2mM N-acetyl cysteine (NAC; Sigma-Aldrich, A9165) was accomplished by addition to all staining, wash, and imaging medias. Vehicle treatment for NAC was water. A 1-hour pretreatment with 1uM CMPD39 (also called USP30 inhibitor 18; MedChem Express, HY-141659) was accomplished by addition to the wash and imaging medias. Vehicle treatment for CMPD39 was DMSO.

Immunofluorescence-based analysis of synthetic system

Triple positive HeLa cells were seeded and treated with Doxycycline for 48 hours as with the live cell imaging of the synthetic system described above. Individual wells of the 96 well plates were treated with rapalog for treatment durations ranging from 1 hour to 24 hours. Treatments were started at different times such that all cells were fixed at 72 hours following seeding. Cells were fixed, subjected to immunofluorescence staining for Tom20, and imaged on an Operetta microscope in confocal mode with a 40x water objective, no pixel binning, and standard filter sets.

Western blots

Triple positive cells were treated with 100nM Rapamycin or 200nM rapalog for either 6 or 24 hours. Cell pellets were collected and lysed in RIPA buffer (Sigma, R0278) with Halt protease and phosphatase inhibitor cocktail (Thermo Fisher Scientific, 78440). Lysates were run on Mini-Protean TGX 4-20% SDS-PAGE gels (Bio-Rad, 4561093). Western blotting was performed using standard approaches. Odyssey blocking buffer (LI-COR Biosciences, 927-50000) was used for blocking and antibody incubation steps. Western blots were imaged using a LI-COR Odyssey CLx infrared imaging system. Primary antibodies: 1:1000 Rb-anti-pS6 (Ser235/236; Cell Signaling Technologies, 4856), 1:1000 Rabbit anti-S6 (Cell Signaling Technologies, 2217), and 1:1000 Mouse anti-GAPDH (Cell Signaling Technologies, 97166). Secondary antibodies: 1:5000 Goat anti-rabbit IRDye 800CW (LI-COR, 926-32211) and 1:5000 Goat anti-mouse IRDye 680LT (LI-COR, 926-68020).

Microscopy image processing and cell segmentation

All image quantification and analyses were done in MATLAB. Images were first subjected to flatfield illumination correction using pre-generated objective-specific flatfield correction estimations. Uniform background fluorescence was then removed for each image. Finally, mild fluorescence crossover from the TRITC channel (mKO2) into the FITC channel (mNeonGreen or AF488) was corrected in a pixelwise manner using a multiply and subtract compensation approach.

Crossover correction was empirically calibrated for each microscope objective and for each set of exposure times used in this study. Correction multipliers ranged from 0.02 to 0.05. Processed images were then subjected to cell segmentation using an in-house watershed segmentation pipeline. Segmentation was able to identify nucleus and cytoplasm boundaries for each cell, in each field of view, and at each timepoint (when applicable). A perinuclear region was also identified by removing the outer 25% of pixels from the edge of the cytoplasm region in order to remove small segmentation errors due to overlap with adjacent cells.

Single cell quantification for Parkin tag variant comparison

Following single cell segmentation, prior to quantification, single cell data from three experimental technical replicates was pooled. For each cell, foreground images were calculated for both the anti-Parkin and anti-Tom20 channels by applying a tophat filter with a radius of 8 pixels to each cell image. Colocalization of anti-Parkin and anti-Tom20 intensity was calculated as the pixelwise Pearson correlation coefficient of the foreground images for those channels. For each cell, total anti-Tom20 intensity was calculated by identifying a mitochondrial mask (anti-Tom20 foreground image intensity above 2500 AFU) and then calculating the total pixel intensity of the original anti-Tom20 image within that mask.

To compare cells with similar Parkin expression levels, a sliding window median approach was used for each construct and each condition. For each construct, DMSO-treated data was used to normalize the CCCP-treated data. When comparing Parkin-Tom20 staining colocalization, the DMSO sliding median values were subtracted from the corresponding CCCP-treated values to estimate the change in localization across parkin expression levels. When comparing total cellular anti-Tom20 intensity, CCCP-treated values were divided by the corresponding DMSO-treated values to estimate the proportion of anti-Tom20 staining remaining across parkin expression levels. Finally, these normalized sliding medians were summarized by taking the mean value along their length. Hypothesis testing was performed using a bootstrap approach of random resampling with replacement of single cell data, 1000 times. Due to all 6 Parkin tag/mutant variants (Untagged, Internal, Nterm-M1, Nterm-I1, Nterm-LD, and Untagged-E16A) being assessed in parallel, a Bonferroni multiple comparison correction for 5 comparisons was used.

Single cell quantification and tracking for live-cell Pink1/Parkin dynamics

Following single cell segmentation, foreground images and mitochondrial masks were calculated. First, foreground images were calculated for all images by applying a 2D bandpass gaussian filter (sequential high pass gaussian with a wavelength of 50 pixels to remove background intensity and low pass gaussian with wavelength of 3 pixels to reduce pixel noise) using the filt2 MATLAB function (47). Then, a mitochondria mask was calculated using the MtTether-SNAP foreground image and a threshold that was calculated on a per-cell basis. This cell-specific threshold was calculated as the median pixel intensity of the original cell MtTether-SNAP image raised to the exponent of 0.8 and multiplied by 2. This approach was empirically designed to be effective over a range of MtTether-SNAP expression levels.

Next, a wide variety of distinct cellular phenotypes (features) were quantified. Colocalization features were calculated as the pixelwise Pearson correlation coefficients between pairs of foreground images. Mitochondrial Pink1 levels were calculated as the mean pixelwise Pink1-mNG foreground intensity in the mitochondrial mask. Colocalization features and mitochondrial Pink1-mNG features were calculated only in the perinuclear region to avoid intensity artifacts at cell edges. Expression level features were calculated as means from the original channel intensity images.

Following feature extraction, single cells were tracked from one timepoint to the next by finding the closest cell both in distance and phenotypic similarity. This was accomplished using a distance metric defined as the weighted sum of: the distance between nucleus centroids, the ratio of cell areas, the ratio of nucleus area, and the ratio of marker expression levels between candidate cells. Only cells which were successfully tracked across all timepoints were used for downstream analyses. Additionally, cells with high or low Parkin-mKO2 expression or tether expression levels (approximately top and bottom 5%) were discarded. Cells with high Pink1-mNG expression levels above 10000 AFU/px were discarded as well due to masking of visible Pink1-mNG recruitment to mitochondria at such high expression levels. Similarly, cells which did not show an increased Pink1-mitochondria colocalization upon rapalog treatment were discarded as well. Quantification of mitochondrial Pink1 levels was prone to overestimation for cells with no Pink1-mitochondria colocalization, but was effective if Pink1 was colocalized with mitochondria. Finally, because some cells with low initial Pink1-mNG expression levels underwent doxycycline-dependent Pink1-mNG expression induction during the experiment, cells with Pink1-mNG expression levels which increased 1.5-fold or more over the course of the experiment were discarded.

Quantification of Pink1 input, and detection time of Parkin recruitment, and Parkin output scores for live cell data

Each cell was quantified individually. First, the Parkin output score was calculated as the maximum observed colocalization between the Parkin-mKO2 and MtTether-SNAP channels. The maximum value was used because extended Parkin-mKO2 recruitment eventually caused mitochondrial degradation and an associated decrease in measured foreground colocalization. Next, the detection time of Parkin recruitment was determined as the time at which the colocalization of Parkin-mKO2 and MtTether-SNAP reached a value of 0.4. The value 0.4 was chosen because it is half-way between the values measured for cells with and without parkin recruitment. Finally, the Pink1 input score was calculated from the mitochondrial Pink1 level. Due to noise over time in this quantification, the data was smoothed via a moving mean with a window size of 5 timepoints. The Pink1 input score was then calculated by taking the mean value of this smoothed data from when Pink1 was recruited either to when Parkin was recruited (for cells with Parkin response scores > 0.4) or to the end of the experiment (for cells with Parkin response scores < 0.4).

Threshold and delay function quantification from live cell Pink1/Parkin experiments

First, an input threshold value was calculated as the Pink1 input score which caused one half of cells to exhibit Parkin recruitment (Parkin response scores > 0.4). Next a hyperbolic curve was fit to the Pink1 input score vs detection time of Parkin recruitment data. This hyperbola was fit using a geometric distance, sum of squares, approach. Specifically, "delay scaler" in the following function was optimized to fit the curve: *Detection time of Parkin recruitment = (delay scaler) / (Pink1 input score – input threshold)*. When comparing across conditions, hypothesis testing was performed using a bootstrap approach. Random sampling with replacement and re-quantification of the input threshold and delay scaler values was performed 1000 times. A Bonferroni multiple comparison correction was used to correct for 2 comparisons in each experiment.

Single cell quantification for fixed-cell Pink1/Parkin dynamics

Following single cell segmentation, foreground images were calculated as for live cell data above. Gaussian bandpass wavelengths used were 100 pixels and 3 pixels. A Mitochondria mask was calculated using the anti-Tom20 mitochondrial foreground image and a fixed threshold of 1000 AFU. A single intensity threshold was used to calculate the mitochondria mask due to uniform Tom20 staining across cells. Next, features were calculated the same way as described above for live cell data.

Finally, the data quality control was performed. Mis-segmented cells (small cell areas, small nucleus areas, or high levels of hoechst staining in the cytoplasm) were discarded. Cells with high or low Parkin expression (approximately top or bottom 5%) were discarded. Cells with low Tom20 staining (average cellular intensity less than 1200 AFU/pixel) were discarded to remove cells that had undergone mitochondrial degradation at longer timepoints. Cells with a low Pink1-Tom20 intensity colocalization scores (less than 0.6) were discarded to ensure that only cells with mitochondrial Pink1 were analyzed.

Threshold and delay function quantification from fixed cell Pink1/Parkin experiments

Each cell had a quantified mitochondrial Pink1 level (Pink1 intensity on mitochondria) and a Parkin recruitment level (Parkin-Tom20 intensity colocalization). At each timepoint (rapalog treatment length), a variant of Otsu's thresholding method was used to identify a Pink1 level capable of separating cells into two Parkin recruitment level populations. In short, an algorithm searched for a Pink1 level separation value, able to minimize variance in Parkin recruitment level for the two populations of cells: Pink1 levels above or below the separation value. Functionally, this separation value identified the mitochondrial Pink1 level required to trigger Parkin recruitment for each length of rapalog treatment. A hyperbola with the equation *Pink1 separation value = A/(rapalog treatment time) + B* was fit to the data by identifying values of A and B using a nonlinear least squares approach. The hyperbola fitted here is an algebraic rearrangement (swapping the dependent and independent variables) of the hyperbola fit to the live cell data. The hyperbolic relationship extracted from the fixed cell data (required Pink1 level as a function of Rapalog treatment time) is distinct in identity but similar in nature to the hyperbolic relationship

extracted from the live cell data (Parkin recruitment time as a function of Mitochondrial Pink1 levels).

Analysis of existing UBL domain structures

PDB accession numbers for structures shown in Fig. S2 are as follows: hParkin, 5c1z; hUbiquitin, 1f9j; hRAD23B, 1uel; hNedd8, 1ndd; hHoil1, 2lgy; hUbiquilin, 2klc. Visualization and analysis were performed in the UCSF ChimeraX software (*48*). Hydrogen bonds were predicted using the show hydrogen bonds option in UCSF ChimeraX.

Conservation at the N-terminus of N-terminal ubiquitin-like domains

The amino acid sequence and the location of 67594 annotated UBL domains in the Uniprot database were downloaded. Analysis performed in MATLAB. Entries with annotation errors were identified and discarded as follows. UBL domain entries with non-standard residues (including "J", "O", "U", "X" or "Z") or with lengths of less than 65 amino acids were discarded. Domains annotated to start at the protein's N-terminus, but which did not start with methionine were discarded. Entries corresponding to Ubiquitin polymers rather than to UBL domains were identified as starting or ending with tandem glycines. These Ubiquitin entries were discarded to prevent over-representation in the dataset. Finally, any additional UBL domains originating from proteins containing at least one discarded domain were also discarded. This cleanup left 28447 UBL domain entries.

Alignment using traditional methods could not be used successfully due to the large number of sequences and high observed sequence variability. Instead, the conserved positioning of specific residues in the stereotyped ubiquitin fold were used to align the sequences. Specifically, the following schema was used: hydrophobic residues in the first beta sheet at positions 1, 3 and, 5; a variable-length linker with length "x" ranging from 7 to 17 residues; hydrophobic residues in the second beta sheet at positions 5+x+1 and 5+x+3; hydrophobic residues in the first alpha helix at positions 5+x+9, 5+x+12, and 5+x+16. These positions correspond to M1, V3, V5, V15, V17, I23, L26 and V30 of Parkin's UBL domain. Potential alignments were evaluated using scoring system where non-standard hydrophobic residues were allowed but were assigned a penalty score. Furthermore, extreme linker lengths, "x", were penalized for being too short or too long. Finally, a small tie-breaker bonus was awarded to alignments which contained a positively charged residue at the equivalent position of K27 of Parkin's UBL. This positively charged residue was used in the alignment because it was present in nearly all of the published UBL domain structures which were used as reference when designing the alignment schema. Using this approach, 19623 sequences out of 28447 were successfully aligned for residues surrounding the region of interest, M1 and E16 of Parkin's UBL.

Finally, aligned sequences were separated into those belonging to N-terminal or internal UBL domains and were submitted to WebLogo3 (*49*) to create the sequence logos used to visualize residue conservation.

Alignment of mammalian Parkin ortholog protein sequences

Full protein sequences for Parkin were downloaded from Uniprot for human (O60260) pig (Q2L7G3), dog (A0A8C0PPD2), rat (Q9JK66), mouse (Q9WVS6), and guinea pig (H0V739). These sequences were aligned using the Clustal Omega multiple sequence alignment tool by EMBL-EBI (*50*).

References and Notes:

1. A. M. Pickrell, R. J. Youle, The Roles of PINK1, Parkin, and Mitochondrial Fidelity in Parkinson's Disease. *Neuron*. 85, 257–273 (2015).

2. D. P. Narendra, S. M. Jin, A. Tanaka, D.-F. Suen, C. A. Gautier, J. Shen, M. R. Cookson, R. J. Youle, PINK1 Is Selectively Stabilized on Impaired Mitochondria to Activate Parkin. *Plos Biol*. 8, e1000298 (2010).

3. N. Matsuda, S. Sato, K. Shiba, K. Okatsu, K. Saisho, C. A. Gautier, Y. Sou, S. Saiki, S. Kawajiri, F. Sato, M. Kimura, M. Komatsu, N. Hattori, K. Tanaka, PINK1 stabilized by mitochondrial depolarization recruits Parkin to damaged mitochondria and activates latent Parkin for mitophagy. *J Cell Biol*. 189, 211–221 (2010).

4. A. Ordureau, S. A. Sarraf, D. M. Duda, J.-M. Heo, M. P. Jedrychowski, V. O. Sviderskiy, J. L. Olszewski, J. T. Koerber, T. Xie, S. A. Beausoleil, J. A. Wells, S. P. Gygi, B. A. Schulman, J. W. Harper, Quantitative Proteomics Reveal a Feedforward Mechanism for Mitochondrial PARKIN Translocation and Ubiquitin Chain Synthesis. *Mol Cell*. 56, 360–375 (2014).

5. M. Lazarou, D. A. Sliter, L. A. Kane, S. A. Sarraf, C. Wang, J. L. Burman, D. P. Sideris, A. I. Fogel, R. J. Youle, The ubiquitin kinase PINK1 recruits autophagy receptors to induce mitophagy. *Nature*. 524, 309–314 (2015).

6. A. Ordureau, J. A. Paulo, J. Zhang, H. An, K. N. Swatek, J. R. Cannon, Q. Wan, D. Komander, J. W. Harper, Global Landscape and Dynamics of Parkin and USP30-Dependent Ubiquitylomes in iNeurons during Mitophagic Signaling. *Mol Cell*. 77, 1124-1142.e10 (2020).

7. L. A. Kane, M. Lazarou, A. I. Fogel, Y. Li, K. Yamano, S. A. Sarraf, S. Banerjee, R. J. Youle, PINK1 phosphorylates ubiquitin to activate Parkin E3 ubiquitin ligase activity. *J Cell Biol*. 205, 143–153 (2014).

8. A. Kazlauskaite, C. Kondapalli, R. Gourlay, D. G. Campbell, M. S. Ritorto, K. Hofmann, D. R. Alessi, A. Knebel, M. Trost, M. M. K. Muqit, Parkin is activated by PINK1-dependent phosphorylation of ubiquitin at Ser65. *Biochem J*. 460, 127–139 (2014).

9. F. Koyano, K. Okatsu, H. Kosako, Y. Tamura, E. Go, M. Kimura, Y. Kimura, H. Tsuchiya, H. Yoshihara, T. Hirokawa, T. Endo, E. A. Fon, J.-F. Trempe, Y. Saeki, K. Tanaka, N. Matsuda, Ubiquitin is phosphorylated by PINK1 to activate parkin. *Nature*. 510, 162–166 (2014).

10. C. Kondapalli, A. Kazlauskaite, N. Zhang, H. I. Woodroof, D. G. Campbell, R. Gourlay, L. Burchell, H. Walden, T. J. Macartney, M. Deak, A. Knebel, D. R. Alessi, M. M. K. Muqit, PINK1 is activated by mitochondrial membrane potential depolarization and stimulates Parkin E3 ligase activity by phosphorylating Serine 65. *Open Biol*. 2, 120080 (2012).

11. K. Shiba-Fukushima, Y. Imai, S. Yoshida, Y. Ishihama, T. Kanao, S. Sato, N. Hattori, PINK1-mediated phosphorylation of the Parkin ubiquitin-like domain primes mitochondrial translocation of Parkin and regulates mitophagy. *Sci Rep-uk*. 2, 1002 (2012).

12. V. Sauvé, G. Sung, N. Soya, G. Kozlov, N. Blaimschein, L. S. Miotto, J.-F. Trempe, G. L. Lukacs, K. Gehring, Mechanism of parkin activation by phosphorylation. *Nat Struct Mol Biol*. 25, 623–630 (2018).

13. C. Gladkova, S. L. Maslen, J. M. Skehel, D. Komander, Mechanism of parkin activation by PINK1. *Nature*. 559, 410–414 (2018).

14. J.-F. Trempe, V. Sauvé, K. Grenier, M. Seirafi, M. Y. Tang, M. Ménade, S. Al-Abdul-Wahid, J. Krett, K. Wong, G. Kozlov, B. Nagar, E. A. Fon, K. Gehring, Structure of Parkin Reveals Mechanisms for Ubiquitin Ligase Activation. *Science*. 340, 1451–1455 (2013).

15. B. E. Riley, J. A. Olzmann, A Polyubiquitin Chain Reaction: Parkin Recruitment to Damaged Mitochondria. *Plos Genet*. 11, e1004952 (2015).

16. E. V. Rusilowicz-Jones, J. Jardine, A. Kallinos, A. Pinto-Fernandez, F. Guenther, M. Giurrandino, F. G. Barone, K. McCarron, C. J. Burke, A. Murad, A. Martinez, E. Marcassa, M. Gersch, A. J. Buckmelter, K. J. Kayser-Bricker, F. Lamoliatte, A. Gajbhiye, S. Davis, H. C. Scott, E. Murphy, K. England, H. Mortiboys, D. Komander, M. Trost, B. M. Kessler, S. Ioannidis, M. K. Ahlijanian, S. Urbé, M. J. Clague, USP30 sets a trigger threshold for PINK1–PARKIN amplification of mitochondrial ubiquitylation. *Life Sci Alliance*. 3, e202000768 (2020).

17. M. Gersch, C. Gladkova, A. F. Schubert, M. A. Michel, S. Maslen, D. Komander, Mechanism and regulation of the Lys6-selective deubiquitinase USP30. *Nat Struct Mol Biol*. 24, 920–930 (2017).

18. G. Twig, A. Elorza, A. J. A. Molina, H. Mohamed, J. D. Wikstrom, G. Walzer, L. Stiles, S. E. Haigh, S. Katz, G. Las, J. Alroy, M. Wu, B. F. Py, J. Yuan, J. T. Deeney, B. E. Corkey, O. S. Shirihai, Fission and selective fusion govern mitochondrial segregation and elimination by autophagy. *Embo J*. 27, 433–446 (2008).

19. B. Richter, D. A. Sliter, L. Herhaus, A. Stolz, C. Wang, P. Beli, G. Zaffagnini, P. Wild, S. Martens, S. A. Wagner, R. J. Youle, I. Dikic, Phosphorylation of OPTN by TBK1 enhances its binding to Ub chains and promotes selective autophagy of damaged mitochondria. *Proc National Acad Sci*. 113, 4039–4044 (2016).

20. B. S. Padman, T. N. Nguyen, L. Uoselis, M. Skulsuppaisarn, L. K. Nguyen, M. Lazarou, LC3/GABARAPs drive ubiquitin-independent recruitment of Optineurin and NDP52 to amplify mitophagy. *Nat Commun*. 10, 408 (2019).

21. J. Liang, A. Martinez, J. D. Lane, U. Mayor, M. J. Clague, S. Urbé, USP30 deubiquitylates mitochondrial Parkin substrates and restricts apoptotic cell death. *Embo Rep*. 16, 618–627 (2015).

22. C. N. Cunningham, J. M. Baughman, L. Phu, J. S. Tea, C. Yu, M. Coons, D. S. Kirkpatrick, B. Bingol, J. E. Corn, USP30 and parkin homeostatically regulate atypical ubiquitin chains on mitochondria. *Nat Cell Biol*. 17, 160–169 (2015).

23. B. Bingol, J. S. Tea, L. Phu, M. Reichelt, C. E. Bakalarski, Q. Song, O. Foreman, D. S. Kirkpatrick, M. Sheng, The mitochondrial deubiquitinase USP30 opposes parkin-mediated mitophagy. *Nature*. 510, 370–375 (2014).

24. B. Xiao, J.-Y. Goh, L. Xiao, H. Xian, K.-L. Lim, Y.-C. Liou, Reactive oxygen species trigger Parkin/PINK1 pathway–dependent mitophagy by inducing mitochondrial recruitment of Parkin. *J Biol Chem*. 292, 16697–16708 (2017).

25. B. Xiao, X. Deng, G. G. Y. Lim, S. Xie, Z. D. Zhou, K.-L. Lim, E.-K. Tan, Superoxide drives progression of Parkin/PINK1-dependent mitophagy following translocation of Parkin to mitochondria. *Cell Death Dis*. 8, e3097–e3097 (2017).

26. C.-M. Hung, P. S. Lombardo, N. Malik, S. N. Brun, K. Hellberg, J. L. V. Nostrand, D. Garcia, J. Baumgart, K. Diffenderfer, J. M. Asara, R. J. Shaw, AMPK/ULK1-mediated phosphorylation of Parkin ACT domain mediates an early step in mitophagy. *Sci Adv*. 7, eabg4544 (2021).

27. D. Safiulina, M. Kuum, V. Choubey, N. Gogichaishvili, J. Liiv, M. A. Hickey, M. Cagalinec, M. Mandel, A. Zeb, M. Liiv, A. Kaasik, Miro proteins prime mitochondria for Parkin translocation and mitophagy. *Embo J*. 38, e99384 (2019).

28. R. Gómez-Sánchez, M. E. Gegg, J. M. B.-S. Pedro, M. Niso-Santano, L. Alvarez-Erviti, E. Pizarro-Estrella, Y. Gutiérrez-Martín, A. Alvarez-Barrientos, J. M. Fuentes, R. A. González-Polo, A. H. V. Schapira, Mitochondrial impairment increases FL-PINK1 levels by calcium-dependent gene expression. *Neurobiol Dis*. 62, 426–440 (2014).

29. L. Burchell, V. K. Chaugule, H. Walden, Small, N-Terminal Tags Activate Parkin E3 Ubiquitin Ligase Activity by Disrupting Its Autoinhibited Conformation. *Plos One*. 7, e34748 (2012).

30. S. R. Denison, F. Wang, N. A. Becker, B. Schüle, N. Kock, L. A. Phillips, C. Klein, D. I. Smith, Alterations in the common fragile site gene Parkin in ovarian and other cancers. *Oncogene*. 22, 8370–8378 (2003).

31. A. F. Schubert, C. Gladkova, E. Pardon, J. L. Wagstaff, S. M. V. Freund, J. Steyaert, S. L. Maslen, D. Komander, Structure of PINK1 in complex with its substrate ubiquitin. *Nature*. 552, 51–56 (2017).

32. J. L. Bowling, M. C. Skolfield, W. A. Riley, A. P. Nolin, L. C. Wolf, D. E. Nelson, Temporal integration of mitochondrial stress signals by the PINK1:Parkin pathway. *Bmc Mol Cell Biology*. 20, 33 (2019).

33. P. J. Belshaw, S. N. Ho, G. R. Crabtree, S. L. Schreiber, Controlling protein association and subcellular localization with a synthetic ligand that induces heterodimerization of proteins. *Proc National Acad Sci*. 93, 4604–4607 (1996).

34. M. Lazarou, S. M. Jin, L. A. Kane, R. J. Youle, Role of PINK1 Binding to the TOM Complex and Alternate Intracellular Membranes in Recruitment and Activation of the E3 Ligase Parkin. *Dev Cell*. 22, 320–333 (2012).

35. J. H. Bayle, J. S. Grimley, K. Stankunas, J. E. Gestwicki, T. J. Wandless, G. R. Crabtree, Rapamycin Analogs with Differential Binding Specificity Permit Orthogonal Control of Protein Activity. *Chem Biol*. 13, 99–107 (2006).

36. E. V. Rusilowicz-Jones, F. G. Barone, F. M. Lopes, E. Stephen, H. Mortiboys, S. Urbé, M. J. Clague, Benchmarking a highly selective USP30 inhibitor for enhancement of mitophagy and pexophagy. *Life Sci Alliance*. 5, e202101287 (2022).

37. E. Marcassa, A. Kallinos, J. Jardine, E. V. Rusilowicz-Jones, A. Martinez, S. Kuehl, M. Islinger, M. J. Clague, S. Urbé, Dual role of USP30 in controlling basal pexophagy and mitophagy. *Embo Rep*. 19, e45595 (2018).

38. A. Jilkine, S. B. Angenent, L. F. Wu, S. J. Altschuler, A Density-Dependent Switch Drives Stochastic Clustering and Polarization of Signaling Molecules. *Plos Comput Biol*. 7, e1002271 (2011).

39. M. Swat, A. Kel, H. Herzel, Bifurcation analysis of the regulatory modules of the mammalian G1/S transition. *Bioinformatics*. 20, 1506–1511 (2004).

40. Y. Chen, J. E. Ferrell, C. elegans colony formation as a condensation phenomenon. *Nat Commun*. 12, 4947 (2021).

41. P. S. Dodds, D. J. Watts, A generalized model of social and biological contagion. *J Theor Biol*. 232, 587–604 (2005).

42. L. Wang, Y.-L. Cho, Y. Tang, J. Wang, J.-E. Park, Y. Wu, C. Wang, Y. Tong, R. Chawla, J. Zhang, Y. Shi, S. Deng, G. Lu, Y. Wu, H. W.-S. Tan, P. Pawijit, G. G.-Y. Lim, H.-Y. Chan, J. Zhang, L. Fang, H. Yu, Y.-C. Liou, M. Karthik, B.-H. Bay, K.-L. Lim, S.-K. Sze, C. T. Yap, H.-M. Shen, PTEN-L is a novel protein phosphatase for ubiquitin dephosphorylation to inhibit PINK1–Parkin-mediated mitophagy. *Cell Res*. 28, 787–802 (2018).

43. T. Cornelissen, D. Haddad, F. Wauters, C. V. Humbeeck, W. Mandemakers, B. Koentjoro, C. Sue, K. Gevaert, B. D. Strooper, P. Verstreken, W. Vandenberghe, The deubiquitinase USP15 antagonizes Parkin-mediated mitochondrial ubiquitination and mitophagy. *Hum Mol Genet*. 23, 5227–5242 (2014).

44. A. Kiyatkin, E. Aksamitiene, N. I. Markevich, N. M. Borisov, J. B. Hoek, B. N. Kholodenko, Scaffolding Protein Grb2-associated Binder 1 Sustains Epidermal Growth Factor-induced Mitogenic and Survival Signaling by Multiple Positive Feedback Loops*. *J Biol Chem*. 281, 19925–19938 (2006).

45. N. Borisov, E. Aksamitiene, A. Kiyatkin, S. Legewie, J. Berkhout, T. Maiwald, N. P. Kaimachnikov, J. Timmer, J. B. Hoek, B. N. Kholodenko, Systems-level interactions between insulin–EGF networks amplify mitogenic signaling. *Mol Syst Biol*. 5, 256–256 (2009).

46. N.C. Shaner, G.G. Lambert, A. Chammas, Y.Ni, P.J. Cranfill, M.A. Baird, B.R. Sell, J.R. Allen, R.N. Day, M. Israelsson, M.W. Davidson, J. Wang, A bright monomeric green fluorescent protein derived from branchiostoma lanceolatum, *Nature Methods*, **10**, 407-409 (2013).

47. C. Greene, Filt2 2D geospatial data filter. *MATLAB Central File Exchange*. https://www.mathworks.com/matlabcentral/fileexchange/61003-filt2-2d-geospatial-data-filter (2017, Retrieved September 19, 2022)

48. E.F. Pettersen, T.D. Goddard, C.C. Huang, E.C. Meng, G.S. Couch, T.I. Croll, J.H. Morris, T. E. Ferrin, UCSF ChimeraX: Structure visualization for researchers, educators, and developers. *Protein Sci*. **30(1)**, 70-82 (2021).

49. G.E. Crooks, G. Hon, J.M. Chandonia, S.E. Brenner, WebLogo: A sequence logo generator. *Genome Research*. **14**, 1188-1190 (2004).

50. F. Madeira, M. Pearce, A.R.N. Tivey, P. Basutkar, J. Lee, O. Edbali, N. Madhusoodanan, A. Kolesnikov, R. Lopez, Search and sequence analysis tools services from EMBL-EBI in 2022. Nucleic Acids Research. **50(W1)**, W276-W279 (2022).