Brevicidine, a bacterial non-ribosomally produced cyclic antimicrobial lipopeptide with a unique *modus operandi*

Xinghong Zhao[a,b,c,1,*], Xinyi Zhong[a,1], Hongping Wan[c,1], Lu Liu[a], Xu Song[a], Yuanfeng Zou[a], Lixia Li[a],
Renyong Jia[c], Juchun Lin[a], Huaqiao Tang[a], Shan Zhao[c], Yifei Lang[c], Zhongqiong Yin[a,*], Oscar P.
Kuipers[b]

[a] Centre for Sustainble Antimicrobials, Department of Pharmacy, Sichuan Agricultural University,
   Chengdu, 611130, China.

[b] Department of Molecular Genetics, Groningen Biomolecular Sciences and Biotechnology
   Institute, University of Groningen, Groningen, 9747 AG, The Netherlands.

[c] Key Laboratory of Animal Disease and Human Health of Sichuan Province, Sichuan Agricultural
   University, Chengdu, 611130, China.

[1] These authors contributed equally to this work.

* Correspondence: xinghong.zhao@sicau.edu.cn; yinzhongq@163.com

Abstract

Due to the accelerated appearance of antibiotic-resistant (AMR) pathogens in clinical infections, new first-in-class antibiotics, operating via novel modes of action, are desperately-needed. Brevicidine, a bacterial non-ribosomally produced cyclic lipopeptide, has shown potent and selective antimicrobial activity against Gram-negative pathogens. However, before our investigations, little was known about how brevicidine exerts its potent bactericidal effect against Gram-negative pathogens. In this study, we find that brevicidine has potent antimicrobial activity against AMR *Enterobacteriaceae* pathogens, with a MIC value ranging between 0.5µM and 2µM. In addition, brevicidine showed potent anti-biofilm activity against the *Enterobacteriaceae* pathogens, with same 100% inhibition and 100% eradication concentration of 4µM. Further mechanistic studies showed that brevicidine exerts its potent bactericidal activity via interacting with lipopolysaccharide in the outer membrane, targeting phosphatidylglycerol and cardiolipin in the inner membrane, and dissipating the proton motive force of bacteria. This results in metabolic perturbation, including inhibition of adenosine triphosphate synthesis, inhibits the dehydrogenation of nicotinamide adenine dinucleotides, accumulation of reactive oxygen species in bacteria, and inhibition of protein synthesis. Lastly, brevicidine showed a good therapeutic effect in a mouse peritonitis–sepsis model. Our findings pave the way for further research on clinical applications of brevicidine, to combat the prevalent infections caused by AMR Gram-negative pathogens worldwide.

Keywords: Brevicidine; Antimicrobial resistance; Antimicrobial mechanism; Bacteraemia model; *Escherichia coli*

Author summary

Due to the accelerated appearance of antimicrobial resistance in bacterial pathogens, many bacterial infections have become increasingly difficult, or even impossible, to treat with conventional antimicrobials. To address the problem posed by drug-resistant infections new first-in-class antibiotics, operating via novel modes of action, are desperately needed. Here, we show that the bacterial non-ribosomally produced cyclic antimicrobial lipopeptide brevicidine has potent antimicrobial against multi-drug resistance Gram-negative pathogens, with a MIC value ranging between $0.5\mu M$ and $2\mu M$. Mechanism studies elucidate that brevicidine exerts its potent bactericidal activity via interacting with lipopolysaccharide in the outer membrane, targeting phosphatidylglycerol and cardiolipin in the inner membrane, and dissipating the proton motive force of bacteria. In addition, brevicidine showed a good therapeutic effect in a mouse peritonitis–sepsis model. This study showcases the excellent antibiotic properties of brevicidine in an age where new antibiotics that target Gram-negative bacteria are desperately needed.

Introduction

Due to the accelerated appearance of antimicrobial resistance (AMR) in bacterial pathogens, many bacterial infections have become increasingly difficult, or even impossible, to treat with conventional antimicrobials. The O'Neill report on AMR, published in 2014, predicted that deaths attributable to resistant infections will reach 10 million annually by 2050 [1,2]. In addition, due to the COVID-19 pandemic, the AMR situation is getting worse since increasing evidence has emerged that the COVID-19 pandemic also contributes to AMR [2–6]. Despite the AMR situation being so urgent, the number of newly approved antimicrobials has been steadily decreasing over the past 20 years, especially those for treating Gram-negative pathogen infections [7–9]. Bacteria produce a wealth of non-ribosomally produced peptide (NRP) antimicrobials, including clinically approved antimicrobial cyclic lipopeptides, such as vancomycin, daptomycin, and colistin [10]. This shows that the group of cyclic lipopeptides makes up a rich source of molecules able to control pathogenic bacterial infections.

Brevicidine (Bre), a bacterial non-ribosomally produced cyclic lipopeptide, was first isolated from the *Brevibacillus laterosporus* DSM25 in 2018 [11]. This cyclic lipopeptide contains 12 amino acids (4-Methyl-Hexanoyl-D-Asn-D-Tyr-D-Trp-D-Orn-Orn-Gly-D-Orn-Trp-Thr-Ile-Gly-Ser) with a 4-Methyl-Hexanoyl chain at its N-terminus and a lactone bond between Thr9 and Ser12 (Fig 1) [11,12]. Brevicidine displays potent and selective antimicrobial activity against Gram-negative pathogens, including *Enterobacter cloacae*, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Klebsiella pneumoniae* [11], which are listed as "critical" priority pathogens that need

R&D of new antimicrobials by the World Health Organization (WHO) [13]. Notably,

brevicidine showed no cytotoxicity and hemolytic activity at a relatively high

concentration of 128μg/ml [11], indicating brevicidine is a promising antimicrobial

candidate. However, before our investigations, little was known about how

brevicidine exerts its potent bactericidal effect against Gram-negative pathogens. In

addition, the antimicrobial activity against AMR pathogens and the therapeutic effect

in vivo of brevicidine still needs to be investigated before it is considered for a clinical

trial.

88
Fig 1. The structure of brevicidine and the predicted biosynthetic gene cluster. A, The
non-ribosomal peptide synthetases genes harbored by the *Brevibacillus laterosporus* DSM 25
genome; **B**, the catalytic domains encoded by the gene cluster, and the substrates incorporated
by the respective modules. Domains: A, adenylation; T, thiolation; C, condensation; E,
epimerization; TE, thioesterase; **C**, the structure of brevicidine.

In this study, we aimed to identify the precise mechanism of action by which

brevicidine kills Gram-negative pathogens. Moreover, the anti-biofilm activity against

E. coli, the antimicrobial activity against AMR *Enterobacteriaceae*, and the *in vivo*

therapeutic effect of brevicidine were also investigated. The results showed that

brevicidine exerts its potent bactericidal effect by interacting with lipopolysaccharide (LPS) in the outer membrane, targeting phosphatidylglycerol (PG) and cardiolipin (CL) in the inner membrane, and subsequently disrupting the proton motive force of Gram-negative pathogens. This leads to adenosine triphosphate (ATP) synthesis inhibition, nicotinamide adenine dinucleotide (NADH) and reactive oxygen species (ROS) accumulation, and bacteria death. In addition, brevicidine showed potent antimicrobial activity against AMR Gram-negative pathogens and good anti-biofilm activity against Gram-negative pathogen. Notably, brevicidine showed a good therapeutic effect in an *E. coli*-induced mouse peritonitis–sepsis model. These results will pave the way for further research on the clinical applications of brevicidine.

Results and Discussion

Brevicidine shows potent antimicrobial activity against antibiotic-resistant (AMR) *Enterobacteriaceae* pathogens

The antimicrobial activity of brevicidine against AMR *Enterobacteriaceae* pathogens was measured by a MIC assay. The results show that brevicidine has potent antimicrobial activity, with a MIC value of 0.5μM to 2μM (Table 1), against *Enterobacteriaceae* pathogens, including the tested colistin-resistant strains and multidrug-resistant strains [14]. These results indicate that brevicidine has a different mechanism of action from colistin, ampiciline, ceftriaxone, cefotaxime, aztreonam, gentamicin, ofloxacin, amoxicillin, streptomycin, lincomycin, doxycycline, or ciprofloxacin, which makes it a more attractive antimicrobial candidate for treating AMR Gram-negative pathogen infections.

Table 1 MIC values of brevicidine against *Enterobacteriacea*e bacterial pathogens.

Microorganism	MIC (µM)			
	Brevicidine	**Colistin**	**Amikacin**	**Ampiciline**
Escherichia coli CVCC3749	1	2	2	> 256
Escherichia coli B2[a]	1	8	32	> 256
Escherichia coli 16QD[a]	1	8	8	> 256
Escherichia coli EPZC17-45[b]	2	2	4	> 256
Escherichia coli EPZC17-22[b]	1	1	2	> 256
Escherichia coli E21-5[b]	0.5	2	2	> 256
Escherichia coli E21-71[b]	0.5	4	4	> 256
Escherichia coli E21-75[b]	0.5	2	8	> 256
Escherichia coli E21-107[b]	0.5	1	4	> 256
Enterobacter cloacae LMG02783[c]	1	> 64	8	> 256
Klebsiella pneumoniae LMG20218	2	2	4	> 256

[a]*Escherichia coli* carrying mcr-1, blaNDM-5, mdfA genes, etc., which is resistant against colistin, carbapenems, rifampin, etc. [b] *Escherichia coli* (clinical isolation) carrying multidrug resistance genes, *Escherichia coli* EPZC17-45, resistant to ampiciline, ceftriaxone, cefotaxime, aztreonam, gentamicin, doxycycline, and ciprofloxacin; *Escherichia coli* EPZC17-22, resistant to ampiciline, ceftriaxone, cefotaxime, aztreonam, gentamicin, doxycycline, and ciprofloxacin; *Escherichia coli* E21-5, resistant to ampiciline, amoxicillin, ceftriaxone, gentamicin, cefotaxime, streptomycin, doxycycline; *Escherichia coli* E21-71, resistant to ampiciline, amoxicillin, ceftriaxone, cefotaxime, doxycycline, and ciprofloxacin, and ofloxacin; *Escherichia coli* E21-75, resistant to ampiciline, amoxicillin, ceftriaxone, cefotaxime, doxycycline, ofloxacin, and ciprofloxacin; *Escherichia coli* E21-107, resistant to ampiciline, amoxicillin, ceftriaxone, gentamicin, cefotaxime, streptomycin, doxycycline, and lincomycin. [c]*Enterobacter cloacae* carrying mcr-9 gene, which is a potent colistin-resistant gene.

Subsequently, the safety of brevicidine was assessed by hemolytic activity (Rat and mouse erythrocytes), cytotoxicity (PK-15 cell line), and toxicity (*Galleria mellonella*) assays. The results showed that brevicidine, at a concentration range of 2µM to 128µM, neither caused cell lysis on rat erythrocytes nor on mouse erythrocytes (S1 Fig). Moreover, no significant cytotoxicity on PK-15 cell line was observed after treatment with brevicidine at a concentration range of 0.5µM to 128µM for 48h (S2 Fig). Further, the *in vivo* toxicity of brevicidine was evaluated in *Galleria mellonella*. The results showed that no *Galleria mellonella* died during five days post-treatment with brevicidine at a concentration range of 4mg/kg to 128mg/kg. These findings

demonstrate that brevicidine is a promising candidate that should be further studied

for antimicrobial development.

Fig 2. Brevicidine showed potent anti-biofilm activity against *E. coli* O101. A, Biofilm inhibition
of various antimicrobials (brevicidine, colistin, and amikacin) versus *E. coli* O101. Biofilm
inhibition was assessed by incubating the antimicrobials with bacteria in growth media for 24 h,
discarding the growth media and staining the adhered biomass with CV. The percentage of
biofilm inhibition is reported relative to untreated bacteria (defined as 100%) and sterility control
wells (defined as 0%). All data were presented as means ± s.d. **B**, Biofilm eradication assay results
for various antimicrobials (brevicidine, colistin, and amikacin) versus *E. coli* O101. Biofilm biomass,
quantified based on CV staining, is indicated in pink. Biofilm metabolism, quantified based on TTC
metabolism, is indicated in red. All data were presented as means ± s.d.

154 Brevicidine shows potent anti-biofilm activity against *E. coli*

Bacterial biofilm formation could increase the resistance of pathogens to

conventional antimicrobials, which could decrease the therapeutic effect of

antimicrobials [15,16]. Therefore, the anti-biofilm activity of antimicrobials plays a

vital role in their therapeutic effect on biofilm-formation bacterial pathogen

infections. In this study, the anti-biofilm activity of brevicidine against *E. coli* was

evaluated according to the protocol described previously [17]. Colistin (Col) was used

as an anti-biofilm antimicrobial control, while amikacin (Amk) was used as a control antimicrobial without anti-biofilm activity. The results show that brevicidine has potent anti-biofilm activity against *E. coli*, with the same 100% inhibition and eradication concentration of 4μM, which is better than the anti-biofilm activity of colistin against *E. coli* (Fig 2). These results demonstrate that brevicidine shows high potential for the development of therapeutic antimicrobial.

Brevicidine dissipates the proton motive force of Gram-negative bacteria

In a previous study, brevicidine had shown bacterial lytic activity at a concentration of 10×MIC [11]. However, it did not disrupt the membrane at a concentration of 2× MIC, evidenced by an atomic force microscopy assay. Therefore, the authors suggested that a more in-depth mechanism study is needed to characterize the targets as well as the membrane disruption effect of brevicidine [11]. In this study, to investigate the killing capacity of brevicidine at concentrations of low MIC values, 4× MIC, 2×MIC, and 1×MIC, a time-killing assay was performed as the previously described method [18,19]. Brevicidine quickly decreased the population of *E. coli* at a concentration of 4×MIC, which killed all of the tested bacteria in 1h. Interestingly, the population of *E. coli* did not decrease under the treatment of brevicidine at concentrations of 2×MIC and 1×MIC within 30min (Fig 3A). These results indicate that brevicidine has different modes of action between high and low MIC concentrations; it shows bacterial lytic activity at concentrations of 4-fold to 10-fold MIC, while it does not disrupt the membrane at concentrations of 1-fold to 2-fold MIC [11]. The atomic force microscopy, which was used in the previous study [11], is not a common technique for investigating the membrane effect of antimicrobials

because of its low resolution. Therefore, in this study, we performed several

fluorescent probe assays, SEM assay, and TEM assay to get insight into the effect of

brevicidine on the Gram-negative bacterial membrane.

Fig 3. Brevicidine kills Gram-negative bacteria via dissipating their proton motive force. A, time-killing curves of brevicidine at concentrations of 4×MIC, 2×MIC, and 1×MIC against *E. coli* O101. **B**, *E. coli* O101 cells pretreated with propidium iodide were exposed to antimicrobial peptides brevicidine at concentrations of 2×MIC, 1×MIC, and 0.5×MIC, and the extent of membrane leakage was visualized as an increase in fluorescence. **C**, NPN fluorescence in *E. coli* O101 cells upon exposure to brevicidine at concentrations of 2×MIC, 1×MIC, and 0.5×MIC. Representative examples from three technical replicates are shown. **D**, DiSC$_3$(5) fluorescence in *E. coli* O101 cells upon exposure to brevicidine at concentrations of 2×MIC, 1×MIC, and 0.5×MIC. Representative examples from three technical replicates are shown. **E**, ATP concentration of *E. coli* O101 cells treated with brevicidine at concentrations of 2×MIC, 1×MIC, and 0.5×MIC. **F**, Resorufin fluorescence in *E. coli* O101 cells upon exposure to brevicidine at concentrations of 2×MIC, 1×MIC, and 0.5×MIC. Representative examples from three technical replicates are shown. **G**, Total ROS accumulation in *E. coli* O101 cells treated with brevicidine at concentrations of 2×MIC, 1×MIC, and 0.5×MIC. Representative examples from three technical replicates are shown. **H**, Growth curves of *E. coli* O101 exposed to brevicidine at concentrations of 2×MIC, 1×MIC, and 0.5×MIC. Exogenous addition of NAC (6 µM) diminished the bactericidal activity of brevicidine.

The effect of brevicidine on the membrane integrity of Gram-negative bacteria was

investigated by using the fluorescent probes N-Phenyl-1-naphthylamine (NPN) and

propidium iodide (PI). NPN is an outer-membrane barrier permeant fluorescent

probe [20]. If the outer membrane is disrupted, the fluorescent probe can enter,

reaching the phospholipid layer and resulting in a significant increase of fluorescence.

The fluorescent probe PI is permeant to membrane intact bacteria, and the

disruption of membrane integrity can lead to a prominent increase in fluorescence

[18]. There was no fluorescence increase observed for both brevicidine-treated

NPN-containing cell suspensions and brevicidine-treated PI-containing cell

suspensions (Fig 3 B and C), indicating brevicidine does not disrupt the membrane

integrity at concentrations of 2×MIC, 1×MIC, and 0.5×MIC.

Although brevicidine showed no effect on the bacterial membrane integrity, it could

dissipate the proton motive force of bacteria as some novel antimicrobials do [21,22].

To test the hypothesis that brevicidine dissipated the proton motive force, a $DiSC_3(5)$

(3,3'-dipropylthiadicarbocyanine iodide) assay was performed. $DiSC_3(5)$ accumulates

in the cytoplasmic membrane of bacteria in response to the $\Delta\psi$ component of the

proton motive force [22,23]. When the transmembrane ΔpH potential of bacteria is

disrupted, cells compensate by increasing the $\Delta\psi$, resulting in enhanced $DiSC_3(5)$

uptake into the cytoplasmic membrane and therefore decreased fluorescence [22,23].

After brevicidine treatment, the fluorescence signal of $DiSC_3(5)$ was immediately

reduced (Fig 3D). In addition, the descent rate of $DiSC_3(5)$ fluorescence signal was

displayed in a dose-dependent manner (Fig 3D). These results demonstrate that

brevicidine dissipats the proton motive force of bacteria.

The proton motive force of bacteria is essential for the generation of ATP, which is an

essential bioactive compound for live bacteria [24,25]. If the proton motive force of

bacteria is dissipated, the process of ATP biosynthesis will be inhibited or even

terminated. To confirm that brevicidine can dissipate the proton motive force of

bacteria, the ATP levels of brevicidne, at concentrations of 2×MIC, 1×MIC, and 0.5×

MIC, treated bacteria were measured by using the BacTiter-Glo Microbial Cell

Viability Assay (Promega) kit. The results showed that brevicidine decreased the ATP

level of bacteria as expected (Fig 3E). These results confirmed that brevicidine could

dissipate the proton motive force of bacteria, which is one of the mechanisms by

brevicidine acts as a bactericidal antimicrobial.

Nonfluorescent resazurin can be reduced to resorufin, a highly fluorescent compound,

in an NADH-dependent manner in the presence of NADH dehydrogenase [26–29].

Therefore, the NADH level of cells can be measured by the resazurin/NADH

dehydrogenase/NADH system. The dissipation of the proton motive force of bacteria

can result in an increase in the NADH levels of cells. To investigate the influence of

brevicidine on the NADH level of bacteria, a Resazurin assay was performed. The

results show that the NADH level of bacteria was greatly increased with the

treatment of brevicidine (Fig 3F). These results demonstrate that the

dehydrogenation of NADH can be inhibited by the addition of brevicidine, which is an

antimicrobial that dissipates the proton motive force of bacterial. Considering the

importance of the dehydrogenation of NADH to the electron transport chain of

bacteria, we hypothesized that brevicidine treatment could result in the

accumulation of ROS and thereafter kills bacteria. Therefore, the ROS levels of

bacteria treated with different concentrations of brevicidine were measured by using

a commercial kit (Beyotime, catalogue no. S0033S). The results show that brevicidine

increases the ROS level of bacteria (Fig 3G), which is one of the reasons that

brevicidine shows bactericidal activity. This was further demonstrated by growth

curve assay, which showed that the bactericidal activity of brevicidine was

attenuated by the addition of the antioxidant N-Acetyl-L-cysteine (NAC, 6 mM) (Fig

3H). Taking together, these results demonstrate that brevicidine exerts its bactericidal

activity via dissipating the proton motive force of bacteria and thereafter inhibits ATP

biosynthesis, inhibits the dehydrogenation of NADH, accumulates ROS in bacteria,

and results in bacteria death.

To investigate the effect of brevicidine on the cell membrane in intuitive ways, we

performed fluorescence microscopy, SEM, and TEM assays. The fluorescence

microscopy assay results are consistent with the PI and NPN assays, which

demonstrate that brevicidine is not an antimicrobial that kills pathogens by

disrupting bacteria membrane integrity (Fig 4A). Interestingly, after treatment with

brevicidine (1×MIC) for 15 min, the bacterial surface fold was observed (Fig 4B). In

addition, after treatment with brevicidine (1×MIC) for 120 min, most of the bacteria

were killed, evidenced by SEM and TEM images (Fig 4 B and C). These results are

consistent with the results of time killing assay (Fig 3A), which showed that

brevicidine started to kill bacteria after 30min treatments. Moreover, the

morphology data are also consistent that brevicidine exerts its bactericidal activity

via dissipating the proton motive force of bacteria because this mode of action can

finally cause bacteria lysis, as we observed.

Fig 4. The morphology analysis of *E. coli* O101 under the treatment of brevicidine. A, Fluorescence microscopy images of *E. coli* O101 cells, which were challenged with brevicidine at concentrations of 2× MIC, 1× MIC, and 0.5× MIC for 5min. Green denotes a cell with an intact membrane, whereas red denotes a cell with a compromised membrane. **B**, SEM image of *E. coli* O101 cells after treatment with brevicidine at a concentration of 1× MIC for 15min, 30min, 60min, and 120min. **C**, TEM image of *E. coli* O101 cells after treatment with brevicidine at a concentration of 1× MIC for 120min.

Brevicidine targets the membrane components LPS, PG, and CL

As brevicidine dissipates the proton motive force of bacteria, we hypothesized that brevicidine could target the component of the bacterial inner membrane. To get an insight into the potential target, an agar diffusion assay was performed to investigate if there is an interaction between brevicidine and LPS (outer membrane component), phosphatidylethanolamine (PE, outer and inner membrane component), PG (inner membrane component), or CL (inner membrane component) [30,31]. The main component of mammalian cellular membrane phosphatidylcholine (PC) was used as a control. The results show that the addition of PC or PE did not attenuate the antimicrobial activity of brevicidine (Fig 5A). In contrast, the antimicrobial activity was completely abolished by the addition of LPS, PG, or CL (Fig 5A). Subsequently, to investigate the dose-dependent effect of LPS, PG, or CL on the antimicrobial activity of brevicidine, a MIC assay was performed with different concentrations of LPS, PG, CL, PC, or PE. The MIC value of brevicidine had no changes with the addition of PC and PE, suggesting there is no interaction between brevicidine and PE (or PC) (Fig 5B). This could explain why brevicidine targets the Gram-negative bacteria but has low cytotoxicity to mammalian cells (S2 Fig) [11]. LPS, PG, and CL decreased the antimicrobial activity of brevicidine in a dose-dependent manner, and the MIC value of brevicidine increased 8-fold, 16-fold, and 64-fold by the addition of 128μg/ml of LPS, PG, and CL, respectively (Fig 5B). These results indicate that brevicidine could bind to the cell membrane components LPS, PG, and CL.

Fig 5. Brevicidine targets the membrane components LPS, PG, and CL. A, Exogenous addition of
LPS, PG, or CL abolishes the antibacterial activity of brevicidine against *E. coli* O101. Inhibition
zones of the mixtures of 1 nmol of brevicidine and 4 mmol of PC, LPS, PE, PG and CL, against *E.
coli* O101 for overnight incubation at 37 °C. **B**, Increased MICs of brevicidine against *E. coli* O101
in the presence of LPS, PG and CL, ranging from 0 μg/ml to 128 μg/ml, based on MIC assays. **C**,
ITC analysis of the interaction between brevicidine and LPS. **D**, ITC analysis of the interaction
between brevicidine and PG. **E**, ITC analysis of the interaction between brevicidine and CL. The K_D
values of brevicidine to LPS, PG and CL were calculated using the Nano Analyze Software (Waters
LLC).

To gain an insight into the interaction between brevicidine and LPS, PG, or CL, an ITC

measurement was performed. The results show that CL and PG showed comparable

binding affinity to brevicidine, with K_D values of 0.156 μM and 0.170 Mm (Fig 5 C and

D), respectively. Notably, CL and PG have much stronger (approximately 10-fold

stronger) binding affinity to brevicidine than LPS, which has a K_D value of 1.785 μM

(Fig 5E). These results demonstrate that brevicidine has multiple targets on the

bacteria, which is good for inhibiting the development of bacterial antibiotic

resistance. Together, the observed results demonstrate that brevicidine exerts its

bactericidal activity via interacting with LPS in the outer membrane and targeting CL

and PG in the inner membrane, and thereafter dissipating the proton motive force of

bacteria.

Brevicidine inhibits the oxidative phosphorylation and protein

synthesis processes of *E. coli*

To gain an insight into the antimicrobial mechanism of brevicidine, transcriptome

analysis was performed on *E. coli* treated with or without brevicidine at a

concentration of 0.5μM. The whole distribution of differentially expressed genes

(DEGs) is shown in Fig 6A. When padj value<0.05 and log2(FoldChange)>0, among

4622 genes, 587 were upregulated and 622 were downregulated, and 3413 genes

were not significantly changed (Fig 6A). Subsequently, Kyoto Encyclopedia of Genes

and Genomes (KEGG) enrichment analysis was performed on the genes that had

significantly expression changes. The results showed that ribosome, oxidative

phosphorylation, and aminoacyl-tRNA biosynthesis pathways were significantly (padj

336
Fig 6. Transcriptome analysis of *E. coli* O101 under the treatment of brevicidine (0.5 μM) for 1 h.
A, The volcano map of DEGs. The abscissa represents the multiple changes of gene expression
between brevicidine treated and untreated *E. coli* O101; the ordinate represents the statistical
significance of the change of gene expression (pvalue<0.05 |log2FoldChange|>0). **B**, Scatter plot
of KEGG pathway enrichment analysis of DEGs (Brevicidine vs untreated). The vertical axis
represents the KEGG enrichment analysis pathway name, and the horizontal axis represents the
gene ratio of the genes in the pathway. The size of the dots indicates the number of DEGs in the
pathway, and the color of the dots corresponds to different padj value ranges. **C**, Genes involved
in oxidative phosphorylation, ribosome, and aminoacyl-tRNA biosynthesis were identified as
significantly different between the treatment of brevicidine and untreated, with pvalue values
<0.05 and log2FoldChange <0 in expression level. All genes that had significantly changes were
down-regulated in oxidative phosphorylation, ribosome, and aminoacyl-tRNA biosynthesis
pathways. All data are presented as means (n= 3 biological replicates).

value<0.05) affected by brevicidine treatment (Fig 6B). After *E. coli* exposure to

brevicidine for 1h oxidative phosphorylation related genes including ATP synthase,

NADH-quinone oxidoreductase, and cytochrome oxidase were significantly

downregulated (Fig 6C). This change would contribute to the lower level of ATP,

higher level of NADH and ROS of the brevicidine treated cells. The downregulated

tRNA ligase and ribosome synthesis metabolism indicate that the protein synthesis

process was inhibited by brevicidine treatment.

Fig 7. Scheme of the mechanism of action brevicidine against *E. coli*. Brevicidine exerts its bactericidal activity via interacting with LPS in the outer membrane (OM) and targeting CL and PG in the inner membrane (IM), and thereafter dissipating the proton motive force of the bacteria. This results in metabolic perturbation, including decreases in the intracellular ATP level, inhibits the dehydrogenation of NADH and accumulates ROS in bacteria. In addition, transcriptome analysis results showed that brevicidine inhibits the synthesis of t-RNA ligase (essential for amino acid transport) and ribosome protein synthesis, which can result in protein synthesis inhibition. All these actions together lead to the death of Gram-negative pathogens.

Collectively, our findings demonstrate that brevicidine exerts its bactericidal activity

via interacting with LPS in the outer membrane and targeting CL and PG in the inner

membrane, and thereafter dissipation of the proton motive force of bacteria. This

can result in metabolic perturbation, including inhibits ATP biosynthesis, inhibits the

dehydrogenation of NADH, accumulates ROS in bacteria, and inhibits protein

synthesis (Fig 7).

Brevicidine showed a good therapeutic effect in a mouse

peritonitis–sepsis model

Given the attractive mode of action and potent antimicrobial and anti-biofilm activity

of brevicidine, we investigated its potential as a therapeutic in a mouse

peritonitis–sepsis model (Fig 8A). Mice were infected intraperitoneally with *E. coli* at

a dose that leads to 80% of death. At 1h post-infection, brevicidine was introduced

intraperitoneally at single doses ranging from 10mg/kg to 40mg/kg. All

brevicidine-treated mice were survived (Fig 8B). Consistently, the bacterial load in

different organs of mice significantly reduced under brevicidine treatment in a

dose-dependent manner at either 1 or 7 days post-infection (Fig 8 C-G, and S3 Fig).

These results are consistent with the previous study, which showed that brevicidine

had good antimicrobial activity in a mouse thigh model [11]. Together, these findings

demonstrate the potential of brevicidine as a novel therapeutic antimicrobial in AMR

Gram-negative pathogen infections, in particular, AMR *Enterobacteriaceae* pathogen

infections.

Fig 8. Brevicidine showed a good therapeutic effect in a mouse peritonitis–sepsis model. A, Scheme of the experimental protocol for the mouse peritonitis–sepsis model. **B**, Survival rates of mice in the mouse peritonitis–sepsis model (n= 10). Increased survival rates of mice over 7 d by a dose that leads to 80% of death of *E. coli* O101 (2.0×10^9 c.f.u.), treated with brevicidine (10 mg per kg, 20 mg per kg, or 40 mg per kg) or with colistin (20 mg per kg). **C to G**, Brevicidine significantly reduced the bacterial load of organs of the peritonitis-sepsis mouse. At 24h post-infection, the survived mice (n = 6) were euthanized by cervical dislocation. Bacterial loads (Log10 CFU per gram of *E. coli* O101) of the heart (**C**), liver (**D**), spleen (**E**), lung (**F**) and kidney (**G**) were counted. All data were presented as means ± SD (n = 6). Correlation analyses were evaluated by Pearson r2, ns: ***$p < 0.01$ and ****$p < 0.001$ vs. untreated group.

Conclusions

AMR poses a major threat to human health around the world; many bacterial infections have become increasingly difficult, or even impossible, to treat with conventional antimicrobials. Therefore, new first-in-class antibiotics, operating via novel modes of action, are desperately-needed. Here, we show that brevicidine has potent antimicrobial activity against AMR *Enterobacteriaceae* pathogens, with a MIC value range of 0.5µM to 2µM. In addition, brevicidine showed potent anti-biofilm activity against the *Enterobacteriaceae* pathogens, with the same 100% inhibition and eradication concentration of 4µM. Mechanistic studies showed that brevicidine exerts its potent bactericidal activity via interacting with LPS in the outer membrane, targeting PG and CL in the inner membrane, and dissipating the proton motive force of bacteria. This results in metabolic perturbation, including inhibition of ATP biosynthesis, inhibition of the dehydrogenation of NADH, accumulation of ROS in bacteria, and inhibition of protein synthesis (Fig 7). Lastly, brevicidine showed a good therapeutic effect in a mouse peritonitis–sepsis model. Our findings pave the way for further research on clinical applications of brevicidine, to combat the prevalent infections caused by AMR Gram-negative pathogens worldwide.

Materials and Methods

Ethical approval

The animal experiment protocol was approved by the Animal Ethical and welfare Committee of Sichuan Agricultural University [permission number 20220087].

Bacterial strains used and growth conditions

Bacterial strains used in this study are listed in S1 Table. All bacterial strains were inoculated in LB and incubated at 37°C with aeration at 220 rpm for preparing the overnight cultures.

Purification of brevicidine

Methods for the purification of brevicidine have been described in detail previously [12]. The matrix-assisted laser desorption ionization-time-of-flight mass spectrometry data is shown in S4 Fig, which is showing the correct molecular weight of the purified brevicidine.

Minimum inhibitory concentration (MIC) assay

MIC values of antimicrobials against the tested bacterial pathogens were determined according to the standard guidelines [32]. Briefly, cell concentration was adjusted to approximately 5×10^5 cells per ml in cation-adjusted Mueller-Hinton broth (MHB). Subsequently, antimicrobials were added to the above bacterial cultures at different concentrations. After 24h of incubation at 37°C, the MIC value was defined as the lowest concentration of antimicrobial with no visible growth.

For measuring the MIC values of antimicrobials against indicator strains in the presence of cell membrane components, MHB was replaced with MHB containing different concentrations of cell membrane components, and the effect of these two conditions was compared.

Hemolytic activity assay

Hemolytic activity assay was performed as the method described in previous studies

[11,19]. Briefly, fresh rat or mouse blood was washed with 0.1 M PBS three times, and the erythrocytes were collected. Subsequently, brevicidine was added at final concentrations of 128, 64, 32, 16, 8, 4, 2 and 0 µM in PBS containing 2% (v/v) erythrocytes. The cells were incubated at 37 °C for 1 h and centrifuged for 5 min at 5000 g. The supernatants were transferred to a 96-well plate, and the absorbance was measured at a wavelength of 570 nm with a Thermo Scientific Varioskan™ LUX multimode microplate reader. The absorbance relative to the positive control, which was treated with 10% Triton X-100, was defined as the percentage of hemolysis. Representative examples from three technical replicates are shown.

Mammalian cytotoxicity

The cytotoxicity of brevicidine was evaluated on a porcine kidney cell line (PK-15) by using the cell counting kit-8 (CCK-8; Dojindo Molecular Technologies, Kumamoto, Japan) [33]. PK-15 cells (in Dulbecco's Modified Eagle's medium supplemented with 10% Fetal Bovine Serum) were seeded into 96-well plates and incubated at 37 °C with 5% CO_2. After 24 h, the medium was replaced with a fresh medium (DMEM with 2% FBS, 100 µL per well) containing different concentrations of brevicidine [33]. After 48 h incubation, the CCK-8 reagent was added to the cultures according to the manufacturer's instructions, and the plates were incubated at 37 °C for 2 h with 5% CO_2. Subsequently, the absorbance values were measured by using a Varioskan™ LUX multimode microplate reader (Thermo Fisher Scientific) at 450 nm.

In vivo toxicity in the *Galleria mellonella*

A toxicity analysis was performed on *G. mellonella* according to previously described protocols [34,35]. The worms, 300 to 350 mg per worm, were stored in the dark at

4 °C with wood shavings before use. For each treatment, 10 healthy worms were

placed in Petri dishes with filter paper on the bottom, and 10 μl of the compound

was injected into the worm through the last left proleg. The final dosages injected

into the worms were 4, 8, 16, 32, 64, and 128 mg/kg. NaCl 0.9% was used as a

negative control. The inoculated worms were incubated at 37 °C for 5 d, and record

the mortality daily.

Biofilm inhibition assay

Biofilm inhibition assay was performed according to a previously described protocol

[17]. Briefly, an overnight culture of *E. coli* O101 was diluted 100-fold in tryptone soy

broth (TSB) and incubated at 37 °C with aeration at 220 r.p.m. Bacteria were grown to

an OD_{600} of 0.8, and then the concentration of cells was adjusted to an OD_{600} of 0.04

in TSB. The cell suspension was added to a 96-well plate, and antimicrobials were

added at different final concentrations. After incubation at 37 °C for 24 h, the wells

were washed with MiliQ water three times, and the adhered biomasses were stained

with 105 μl of a 0.1% crystal violet (CV) (wt/vol) solution. The plate was placed on a

low-speed orbital shaker and allowed to stain at room temperature for 30 min.

Subsequently, the plate was washed with MiliQ water three times, and then 70%

(vol/vol) ethanol was added to the wells (110 μl per well) and incubated for 30 min at

room temperature with gentle shaking. The absorbance was measured by using a

Thermo Scientific Varioskan™ LUX multimode microplate reader with a wavelength of

595 nm. The inhibition rates were calculated by the raw CV absorbance values from

each growth condition vs the raw CV absorbance value of normally grown biofilm.

Representative examples from three technical replicates are shown.

Biofilm eradication assay

Biofilm eradication assay was performed according to a previously described protocol [17]. Briefly, an overnight culture of *E. coli* O101 was diluted 100-fold in tryptone soy broth (TSB) and incubated at 37 °C with aeration at 220 r.p.m. Bacteria were grown to an OD_{600} of 0.8, and then the concentration of cells was adjusted to an OD_{600} of 0.04 in TSB. The cell suspension was added to 96-well plates and incubated at 37 °C for 24 h. After washing the wells with TSB three times, antimicrobials were added to the wells at different concentrations, and the plates (one for CV staining, the other one for TTC staining) were incubated at 37 °C for 24 h. For measuring the metabolic activity of biofilm, a final concentration of 0.05% TTC was added to wells with antimicrobials. Subsequently, the adhered biomasses were measured as the method described in the biofilm inhibition assay. After washing all the wells of the TTC plate three times with distilled water, dimethyl sulfoxide (DMSO) was added to the wells at 205 µl per well. After that, the plate was placed on an orbital shaker rotating at low speed and mixed at room temperature for 30 min. The absorbance of the TTC plate was measured by using a Thermo Scientific Varioskan™ LUX multimode microplate reader with a wavelength of 500 nm. The biofilm eradication (Biomass and metabolism) rate was calculated by the method described in the previous study [17]. Representative examples from three technical replicates are shown.

Time-dependent killing assay

The time-dependent killing assay was performed according to the procedure described previously [18,19]. An overnight culture of *E. coli* O101 was diluted 100-fold in MHB and incubated at 37 °C with aeration at 220 r.p.m. Bacteria were

grown to an OD of 0.6, and then the concentration of cells was adjusted to 7.6×10^7

cells per mL. Bacteria were then challenged with antimicrobials at concentrations of

1×MIC, 2×MIC, or 4×MIC in glass culture tubes at 37 °C with aeration at 220 r.p.m.

Bacteria not treated with antimicrobials were used as normal growth control. At

desired time points, two hundred μl aliquots were taken, centrifuged at 5,000 g for 5

min, and resuspended in 200 μl of MHB. Ten-fold serially diluted samples were plated

on MHA plates. After incubation at 37 °C for 24 h, colonies were counted, and the

colony forming units (c.f.u.) per ml were calculated. Each experiment was performed

in triplicate.

Membrane integrity assay

A fresh culture of *E. coli* O101 was pelleted at 5,000 g for 5 min and washed three

times with MHB. The cell density was normalized to an OD_{600} of 0.2, loaded with

propidium iodide (final concentration 2.5 μg per ml), and incubated for 5 min in the

dark for probe fluorescence to stabilize. After the cell suspension was added to a

96-well microplate, brevicidine was added to final concentrations of 2×MIC, 1×MIC,

or 0.5×MIC, with the antimicrobials added after ~30 s, and fluorescence was

monitored for 15 min. The excitation and emission wavelengths on the fluorescence

spectrometer were adjusted to 533 nm and 617 nm, respectively. Representative

examples from three technical replicates are shown.

Outer membrane permeability assay

The integrity of the outer membrane was investigated with the fluorescent probe

N-Phenyl-1-naphthylamine (NPN, Aladdin). A fresh culture of *E. coli* O101 was

pelleted at 5,000 g for 5 min and washed three times with 10 mM HEPES containing

10 mM glucose (GHEPES, pH 7.2). The cell density was normalized to an OD_{600} of 0.2

in GHEPES, loaded with NPN (final concentration 30 μM), and incubated for 30 min in

the dark for probe fluorescence to stabilize. After the cell suspension (190 μL) was

added to a 96-well microplate, brevicidine (10 μL) was added to final concentrations

of 2×MIC, 1×MIC, or 0.5×MIC, with the antimicrobials added after ~30 s, and

fluorescence was monitored for 15 min. The excitation and emission wavelengths on

the fluorescence spectrometer were adjusted to 350 nm and 420 nm, respectively.

Representative examples from three technical replicates are shown.

Fluorescence microscopy assay

This assay was performed according to the procedure described previously [29,36]. *E.*

coli O101 was grown to an OD_{600} of 0.6 in MHB. The culture was pelleted at 5,000 g

for 5 min and washed with MHB three times. After normalization of the cell density

to an OD_{600} of 0.2 in MHB, brevicidine was added to a final concentration of 2×MIC,

1×MIC or 0.5×MIC. After incubation at 37 °C for 5min, cells were collected by

centrifugation. Subsequently, SYTO® 9 and propidium iodide (LIVE/DEAD Baclight

Bacterial Viability Kit, Invitrogen) were added to the above cells. After incubation at

room temperature for 15 min, cells were washed three times with MHB. Then the

cell suspensions were loaded on 1.5 % agarose pads and analyzed by Nikon 80i

(Japan) microscope.

DiSC$_3$(5) assay

DiSC3(5) assay was adapted from previously described procedures [22,29]. Briefly, a

fresh *E. coli* O101 culture was pelleted at 5,000 g for 5 min and washed three times

with MHB. The cell density was normalized to an OD_{600} of 0.2, loaded with 6 μM

DiSC$_3$(5) dye (3,3'-dipropylthiadicarbocyanine iodide), and incubated for 30 min in

the dark for probe fluorescence to stabilize. Subsequently, the cell suspension was

added to a 96-well microplate and incubated for 30 min. After antimicrobials were

added at final concentrations of 2×MIC, 1×MIC, or 0.5×MIC, with the antimicrobials

added after ~30 s, fluorescence was monitored for 100 min. The excitation and

emission wavelengths on the fluorescence spectrometer were adjusted to 622 nm

and 670 nm, respectively. Representative examples from three technical replicates

are shown.

Resazurin assay

Nonfluorescent resazurin can be reduced to resorufin, a highly fluorescent compound,

in an NADH-dependent manner in the presence of NADH dehydrogenase [26–29].

Therefore, the resazurin/NADH dehydrogenase/NADH system can be used to detect

the NADH level of cells. *E. coli* O101 cells were grown in MHB until the OD$_{600}$ reached

0.2. Resazurin was added to the cell culture at a final concentration of 100 μg per ml.

Brevicidine was added to final concentrations of 2×MIC, 1×MIC, or 0.5×MIC. Milli Q

water was used as an untreated control. Fluorescence was recorded by using a

Thermo Scientific Varioskan™ LUX multimode microplate reader with an emission

wavelength of 560 nm. Representative examples from three technical replicates are

shown.

ATP determination

582   The ATP levels were determined using a BacTiter-Glo Microbial Cell Viability Assay kit

583   (Promega). A fresh culture of *E. coli* O101 was pelleted at 5,000 g for 5 min and

584   washed three times with MHB. The cell density was normalized to an OD$_{600}$ of 0.2,

loaded with brevicidine at final concentrations of 2×MIC, 1×MIC, or 0.5×MIC. At desired time points, 100 µl of cell culture were taken out, mixed with 100 µl BacTiter-Glo reagent, and incubated for 5 minutes at room temperature. Luminescence was measured with an Infinite M2000pro Microplate reader (Tecan). CCCP (40 µg/mL) was used as a positive control. The relative ATP levels were calculated using the measured Luminescence values vs the Luminescence value of untreated cells at relative time points.

Reactive oxygen species (ROS) measurement

The levels of ROS in *E. coli* O101 treated with different concentrations of brevicidine were measured by 10 µM of 2',7'-dichlorofluorescein diacetate (DCFH-DA), following the manufacturer's instruction (Beyotime, catalogue no. S0033S). Briefly, a fresh culture of *E. coli* O101 was pelleted at 5,000 g for 5 min and washed three times with MHB. The cell density was normalized to an OD_{600} of 1.0, loaded with DCFH-DA at a final concentration of 10 µM, and the mixture incubated at 37 °C for 30 min. After washing with MHB three times, 190 µl of probe-labeled bacterial cells were added to a 96-well plate, and then 10 µl of brevicidine. Fluorescence was recorded by using a Thermo Scientific Varioskan™ LUX multimode microplate reader with the excitation wavelength at 488 nm and the emission wavelength at 525 nm. The antioxidant N-Acetyl-L-cysteine (NAC, 6 mM) was used as a control to neutralize the production of ROS.

Scanning electron microscopy (SEM) and transmission electron

microscopy (TEM) assays

The morphology of 1μM brevicidne treated *E. coli* O101 was recorded in vacuum on a

ZEISS Gemini SEM300 with an acceleration voltage of 3kV. For TEM measurements,

samples were prepared by delivering 2μl of 1μM brevicidine treated *E. coli* O101

suspension to carbon-coated copper grids and dried in a vacuum system. The

morphology was obtained under a JEM-2100Plus Electron Microscope using an

acceleration voltage of 120kV.

Spot-on-lawn assay

An overnight cultured *E. coli* O101 was added to 0.7% MHA (w/v, temperature 42 °C)

at a final concentration of 0.25% (v/v), and then the mixture was poured onto the

plates, with 20 mL for each 15 cm diameter circular plate. Subsequently, a

spot-on-lawn assay was used to analyze the antimicrobial activity of brevicidine in

the presence of cell membrane components [18]. In short, 20 μL brevicidine (50 μM)

containing 200 μM lipopolysaccharides (LPS, Solarbio, catalogue no. L8880, ≥98%),

L-α-phosphatidylcholine, (PC, Aladdin, catalogue no. L130331, ≥99%),

L-α-phosphatidylglycerol (PG, Aladdin, catalogue no. L130372; ≥99%),

L-α-phosphatidylethanolamine (PE, Aladdin, catalogue no. L130310; ≥99%) or

Cardiolipin (CL, Aladdin, catalogue no. B130227;≥99%) was loaded to the agar plate

that contains indicator strain. After the brevicidiene/cell membrane component

mixture solution drops had dried, the plates were transferred to a 37 °C incubator for

overnight incubation.

Isothermal Titration Calorimetry (ITC) assay

ITC was performed with the Low Volume NanoITC (TA Instruments) at 25°C to

determine the interaction between brevicidine and LPS, PG, or CL. Brevicidine was

diluted in a buffer (20 mM HEPES, pH7.0) to a final concentration of 50 μM. Samples

were degassed before use. The chamber was filled with 300 μL of the brevicidine

solution, and LPS (250μM), PG (500μM), or CL (500μM) was titrated into the

chamber at a rate of 1.96 μL/200 s with a stirring rate of 300 rpm. The control

experiment was performed with 20 mM HEPES titration to 50 μM brevicidine. The K_D

values of brevicidine to LPS, PG, or CL were calculated using the Nano Analyse

Software (Waters LLC).

Transcriptome analysis of E. coli O101 treated with brevicidine

Three replicates of overnight bacteria cultures were diluted at 1:100 in MHB and

grown at 37˚C with aeration at 220rpm. Bacteria were grown to an OD_{600} of 0.4, and

then the concentration of cells was adjusted to an OD_{600} of 0.2 in MHB. Subsequently,

the bacteria were treated with brevicidine at a final concentration of 0.5μM. Bacteria

without brevicidine treatment were used as untreated control. Three replicates were

performed for both brevicidine treated and untreated bacterial cultures, 4ml for each

sample. After incubation at 37˚C with aeration at 220rpm for 1h, the bacteria from

different treatments were collected by centrifuging, and the bacterial pallets were

stored in liquid nitrogen immediately. Subsequently, the samples were sent to

Novogene Co., Ltd (China) for transcriptome analysis. After RNA extraction, mRNA

purification, and cDNA synthesis, the samples were sequenced on the Illumina

NovaSeq6000. The quality of the resulting fastq reads was mapped on the reference

genome using Bowtie2 2.3.4.3 using default settings. Feature Counts 1.5.0-p3 was

used to get the gene counts. DESeq2 1.20 was used to identify all genes, and padj

≤0.05 and |log2 (foldchange) | ≥ 0 of genes between two treatments were regarded as differentially expressed genes (DEGs). The volcano map and KEGG enrichment of DEGs were performed with Magic Novogene package (https://magic.novogene.com). The RNA-Seq data have been deposited in the NCBI Gene Expression Omnibus with the accession number PRJNA872022.

Mouse peritonitis–sepsis model

Brevicidine was tested against *E. coli* O101 in a mouse peritonitis–sepsis model to assess its *in vivo* bioavailability. BALB/c male mice (n=6 per group) were infected with 0.4ml of bacterial suspension (2×10^9 c.f.u. per mouse) via intraperitoneal injection, a concentration that achieves approximately 80% mortality within 48 h post-infection. At 1h post-infection, mice were treated with 0.9% NaCl, brevicidine (40mg/kg), brevicidine (20mg/kg), brevicidine (10mg/kg), or colistin (20mg/kg) via intraperitoneal injection. Mice without *E. coli* O101 infections were used as the normal group control. Once the infected mice died, different organs, including the heart, liver, spleen, lung, and kidney, were collected and homogenized in sterilized PBS for bacterial load quantification. At 24h post-infection, the organs of the survived mice were collected to measure the bacterial load. In addition, the same mouse peritonitis–sepsis model was determined for 7d (n=10), and the bacterial load in the organs of surviving mice was quantified at 7 days post-infection.

Statistical analysis

GraphPad Prism 8.0 was used to fit the data of Fig 2, Fig 3, Fig 5, Fig 8, S1 Fig, S2 Fig, and S3 Fig. The statistical significance of the data was assessed using a two-tailed Student's t-test with GraphPad Prism 8.0. Correlation analyses were evaluated by

677   Pearson r2, ns: p>0.05, *p<0.05, **p<0.01, ***p<0.001, and ****p<0.0001.

Supporting information

S1 Fig. Brevicidine show exceptional low hemolytic activity. Mice (**A**) and rat (**B**) erythrocytes were incubated with brevicidines at concentrations ranging from 2 to 128 µM. Their hemolytic activity was assessed by the release of hemoglobin. Cells treated without a tested compound were used as no-lysis control. Cells treated with 10% Triton X-100 were used as complete lysis control. The data are representative of three independent experiments.

S2 Fig. Brevicidine show exceptional low cytotoxicity. Cytotoxicity of brevicidine against PK-15 cells. The data are representative of three independent experiments.

S3 Fig. Brevicidine significantly reduced the bacterial load of organs of the peritonitis-sepsis mouse. At 7 days post-infection, the survived mice (n = 2 to 6) were euthanized by cervical dislocation. Bacterial loads (Log10 CFU per gram of *E. coli* O101) of the heart (**A**), liver (**B**), spleen (**C**), lung (**D**) and kidney (**E**) were counted. All data were presented as means ± SD (n = 6, or n=2 for the untreated group). Correlation analyses were evaluated by Pearson r2, ns: $p > 0.05$, *$p < 0.05$, **$p < 0.01$, and ***$p < 0.001$ vs. untreated group.

S4 Fig. MALDI-TOF MS data of HPLC purified brevicidine.

S1 Table Strains used in this study.

Data Availability Statement: The transcriptome (RNA sequencing) data that support the findings of this study have been deposited in the NCBI Sequence Read Archive (SRA) with the accession code PRJNA872022.

Finding: Xinghong Zhao was supported by the Science and Technology Project of Sichuan Province (2022YFH0057). Hongping Wan was supported by the Science and Technology Project of Sichuan Province (2022YFH0062).

Competing Interests: The authors have declared that no competing interests exist.

Acknowledgments

705 We thank Prof. Kui Zhu (National Center for Veterinary Drug Safety Evaluation,

706 College of Veterinary Medicine, China Agricultural University, Beijing, China.) for

707 providing the multidrug resistant *Escherichia coli* (B2 and 16QD) strains.

References

1. O'neill JIM. Antimicrobial resistance: tackling a crisis for the health and wealth of nations. Rev Antimicrob Resist. 2014;1: 1–16.

2. Kwon JH, Powderly WG. The post-antibiotic era is here. Science. American Association for the Advancement of Science; 2021. p. 471.

3. Knight GM, Glover RE, McQuaid CF, Olaru ID, Gallandat K, Leclerc QJ, et al. Antimicrobial resistance and COVID-19: Intersections and implications. Elife. 2021;10: e64139.

4. Strathdee SA, Davies SC, Marcelin JR. Confronting antimicrobial resistance beyond the COVID-19 pandemic and the 2020 US election. Lancet. 2020;396: 1050–1053.

5. Lai C-C, Chen S-Y, Ko W-C, Hsueh P-R. Increased antimicrobial resistance during the COVID-19 pandemic. Int J Antimicrob Agents. 2021;57: 106324.

6. Rizvi SG, Ahammad SZ. COVID-19 and antimicrobial resistance: A cross-study. Sci Total Environ. 2022;807: 150873.

7. Butler MS, Blaskovich MAT, Cooper MA. Antibiotics in the clinical pipeline at the end of 2015. J Antibiot (Tokyo). 2017;70: 3.

8. Butler MS, Blaskovich MA, Cooper MA. Antibiotics in the clinical pipeline in 2013. J Antibiot (Tokyo). 2013;66: 571.

9. Roemer T, Boone C. Systems-level antimicrobial drug and drug synergy discovery. Nat Chem Biol. 2013;9: 222.

10. Süssmuth RD, Mainz A. Nonribosomal peptide synthesis—principles and prospects. Angew Chemie Int Ed. 2017;56: 3770–3821.

11. Li Y-X, Zhong Z, Zhang W-P, Qian P-Y. Discovery of cationic nonribosomal peptides as Gram-negative antibiotics through global genome mining. Nat Commun. 2018;9: 3273.

12. Zhao X, Li Z, Kuipers OP. Mimicry of a Non-ribosomally Produced Antimicrobial, Brevicidine, by Ribosomal Synthesis and Post-translational Modification. Cell Chem Biol. 2020;27: 1262-1271.e4. doi:10.1016/j.chembiol.2020.07.005

13. Tacconelli E, Magrini N, Kahlmeter G, Singh N. Global priority list of antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics. World Heal Organ. 2017;27: 318–327.

14. Song M, Liu Y, Huang X, Ding S, Wang Y, Shen J, et al. A broad-spectrum antibiotic adjuvant reverses multidrug-resistant Gram-negative pathogens. Nat Microbiol. 2020;5: 1040–1050.

15. Liu Y, Shi L, Su L, van der Mei HC, Jutte PC, Ren Y, et al. Nanotechnology-based antimicrobials and delivery systems for biofilm-infection control. Chem Soc Rev. 2019;48: 428–446.

16. Sharma G, Sharma S, Sharma P, Chandola D, Dang S, Gupta S, et al. Escherichia coli biofilm: development and therapeutic strategies. J Appl Microbiol. 2016;121: 309–319.

17. Haney EF, Trimble MJ, Hancock REW. Microtiter plate assays to assess antibiofilm activity against bacteria. Nat Protoc. 2021;16: 2615–2632.

752   18.   Zhao X, Yin Z, Breukink E, Moll GN, Kuipers OP. An Engineered Double Lipid II
753   Binding Motifs-Containing Activity against Enterococcus faecium. Antimicrob
754   Agents Chemother. 2020;64: 1–12.
755   19.   Ling LL, Schneider T, Peoples AJ, Spoering AL, Engels I, Conlon BP, et al. A new
756   antibiotic kills pathogens without detectable resistance. Nature. 2015;517:
757   455.
758   20.   Helander IM, Mattila-Sandholm T. Fluorometric assessment of Gram-negative
759   bacterial permeabilization. J Appl Microbiol. 2000;88: 213–219.
760   21.   Cochrane SA, Findlay B, Bakhtiary A, Acedo JZ, Rodriguez-Lopez EM, Mercier P,
761   et al. Antimicrobial lipopeptide tridecaptin A1 selectively binds to
762   Gram-negative lipid II. Proc Natl Acad Sci. 2016;113: 11561–11566.
763   22.   Stokes JM, Yang K, Swanson K, Jin W, Cubillos-Ruiz A, Donghia NM, et al. A
764   deep learning approach to antibiotic discovery. Cell. 2020;180: 688–702.
765   23.   Wu M, Maier E, Benz R, Hancock REW. Mechanism of interaction of different
766   classes of cationic antimicrobial peptides with planar bilayers and with the
767   cytoplasmic membrane of Escherichia coli. Biochemistry. 1999;38: 7235–7242.
768   24.   Ahmed S, Booth IR. The use of valinomycin, nigericin and trichlorocarbanilide
769   in control of the protonmotive force in Escherichia coli cells. Biochem J.
770   1983;212: 105–112.
771   25.   Bakker EP, Mangerich WE. Interconversion of components of the bacterial
772   proton motive force by electrogenic potassium transport. J Bacteriol.
773   1981;147: 820–826.
774   26.   Winartasaputra H, Mallet VN, Kuan SS, Guilbault GG. Fluorometric and
775   colorimetric enzymic determination of triglycerides (triacylglycerols) in serum.
776   Clin Chem. 1980;26: 613–617.
777   27.   McMillian MK, Li L, Parker JB, Patel L, Zhong Z, Gunnett JW, et al. An improved
778   resazurin-based cytotoxicity assay for hepatic cells. Cell Biol Toxicol. 2002;18:
779   157–173.
780   28.   Barnes S, Spenney JG. Stoichiometry of the NADH-oxidoreductase reaction for
781   dehydrogenase determinations. Clin Chim acta. 1980;107: 149–154.
782   29.   Li Z, Chakraborty P, de Vries RH, Song C, Zhao X, Roelfes G, et al.
783   Characterization of two relacidines belonging to a novel class of circular
784   lipopeptides that act against Gram-negative bacterial pathogens. Environ
785   Microbiol. 2020;22: 5125–5136.
786   30.   Sohlenkamp C, Geiger O. Bacterial membrane lipids: diversity in structures
787   and pathways. FEMS Microbiol Rev. 2016;40: 133–159.
788   31.   Santos RS, Figueiredo C, Azevedo NF, Braeckmans K, De Smedt SC.
789   Nanomaterials and molecular transporters to overcome the bacterial
790   envelope barrier: Towards advanced delivery of antibiotics. Adv Drug Deliv
791   Rev. 2018;136: 28–48.
792   32.   Wiegand I, Hilpert K, Hancock REW. Agar and broth dilution methods to
793   determine the minimal inhibitory concentration (MIC) of antimicrobial
794   substances. Nat Protoc. 2008;3: 163.
33.   Zhao X, Cui Q, Fu Q, Song X, Jia R, Yang Y, et al. Antiviral properties of

resveratrol against pseudorabies virus are associated with the inhibition of IκB

kinase activation. Sci Rep. 2017;7. doi:10.1038/s41598-017-09365-0

34.   Ce R, Silva RC, Trentin DS, Marchi JGB De, Paese K, Guterres SS, et al. Galleria

mellonella larvae as an in vivo model to evaluate the toxicity of polymeric

nanocapsules. J Nanosci Nanotechnol. 2020;20: 1486–1494.

35.   Cebrián R, Belmonte-Reche E, Pirota V, de Jong A, Morales JC, Freccero M, et

al. G-quadruplex DNA as a target in pathogenic bacteria: efficacy of an

extended naphthalene diimide ligand and its mode of action. J Med Chem.

2021;65: 4752–4766.

36.   Li Z, de Vries RH, Chakraborty P, Song C, Zhao X, Scheffers D-J, et al. Novel

modifications of nonribosomal peptides from Brevibacillus laterosporus MG64

and investigation of their mode of action. Appl Environ Microbiol. 2020;86.