Brevicidine, a bacterial non-ribosomally produced cyclic antimicrobial lipopeptide with a unique *modus operandi*

Xinghong Zhao[a,b,c,1,*], Xinyi Zhong[a,1], Hongping Wan[c,1], Lu Liu[a], Xu Song[a], Yuanfeng Zou[a], Lixia Li[a],

Renyong Jia[c], Juchun Lin[a], Huaqiao Tang[a], Gang Ye[a], Jianqing Yang[d], Shan Zhao[c], Yifei Lang[c],

Zhongqiong Yin[a,*], Oscar P. Kuipers[b]

[a] Lab for Sustainble Antimicrobials, Department of Pharmacy, Sichuan Agricultural University, Chengdu, 611130, China.

[b] Department of Molecular Genetics, Groningen Biomolecular Sciences and Biotechnology Institute, University of Groningen, Groningen, 9747 AG, The Netherlands.

[c] Key Laboratory of Animal Disease and Human Health of Sichuan Province, Sichuan Agricultural University, Chengdu, 611130, China.

[d] State Key Laboratory of Crop Gene Exploration and Utilization in Southwest China, Sichuan Agricultural University at Wenjiang, Chengdu, Sichuan 611130, China

[1] These authors contributed equally to this work.

* Correspondence: xinghong.zhao@sicau.edu.cn; yinzhongq@163.com

Abstract

Due to the accelerated appearance of antibiotic-resistant (AMR) pathogens in clinical infections, new first-in-class antibiotics, operating via novel modes of action, are desperately-needed. Brevicidine, a bacterial non-ribosomally produced cyclic lipopeptide, has shown potent and selective antimicrobial activity against Gram-negative pathogens. However, before our investigations, little was known about how brevicidine exerts its potent bactericidal effect against Gram-negative pathogens. In this study, we find that brevicidine has potent antimicrobial activity against AMR *Enterobacteriaceae* pathogens, with a MIC value ranging between 0.5μM (0.8mg/L) and 2μM (3.0mg/L). In addition, brevicidine showed potent anti-biofilm activity against the *Enterobacteriaceae* pathogens, with same 100% inhibition and 100% eradication concentration of 4μM (6.1mg/L). Further mechanistic studies showed that brevicidine exerts its potent bactericidal activity via interacting with lipopolysaccharide in the outer membrane, targeting phosphatidylglycerol and cardiolipin in the inner membrane, and dissipating the proton motive force of bacteria. This results in metabolic perturbation, including inhibition of adenosine triphosphate synthesis, inhibits the dehydrogenation of nicotinamide adenine dinucleotides, accumulation of reactive oxygen species in bacteria, and inhibition of protein synthesis. Lastly, brevicidine showed a good therapeutic effect in a mouse peritonitis–sepsis model. Our findings pave the way for further research on clinical applications of brevicidine, to combat the prevalent infections caused by AMR Gram-negative pathogens worldwide.

Keywords: Brevicidine; Antimicrobial resistance; Antimicrobial mechanism; Bacteraemia model; *Escherichia coli*

Introduction

Due to the accelerated appearance of antimicrobial resistance (AMR) in bacterial pathogens, many bacterial infections have become increasingly difficult, or even impossible, to treat with conventional antimicrobials. The O'Neill report on AMR, published in 2014, predicted that deaths attributable to resistant infections will reach 10 million annually by 2050 (1, 2). In addition, due to the COVID-19 pandemic, the AMR situation is getting worse since increasing evidence has emerged that the COVID-19 pandemic also contributes to AMR (2–6). Despite the AMR situation being so urgent, the number of newly approved antimicrobials has been steadily decreasing over the past 20 years, especially those for treating Gram-negative pathogen infections (7–9). Bacteria produce a wealth of non-ribosomally produced peptide (NRP) antimicrobials, including clinically approved antimicrobial cyclic lipopeptides, such as vancomycin, daptomycin, and colistin (10). This shows that the group of cyclic lipopeptides makes up a rich source of molecules able to control pathogenic bacterial infections.

Brevicidine (Bre), a bacterial non-ribosomally produced cyclic lipopeptide, was first isolated from the *Brevibacillus laterosporus* DSM25 in 2018 (11). This cyclic lipopeptide contains 12 amino acids (4-Methyl-Hexanoyl-D-Asn-D-Tyr-D-Trp-D-Orn-Orn-Gly-D-Orn-Trp-Thr-Ile-Gly-Ser) with a 4-Methyl-Hexanoyl chain at its N-terminus and a lactone bond between Thr9 and Ser12 (Fig. 1) (11, 12). In addition, Nathaniel et al reported the total synthetic route of brevicidine in 2022 (13), which makes it more attractive for antibiotic development. Brevicidine displays potent and selective

antimicrobial activity against Gram-negative pathogens, including *Enterobacter*

cloacae, *Escherichia coli*, *Pseudomonas aeruginosa*, and *Klebsiella pneumoniae* (11,

14), which are listed as "critical" priority pathogens that need R&D of new

antimicrobials by the World Health Organization (WHO) (15). Notably, brevicidine

showed no cytotoxicity and hemolytic activity at a relatively high concentration of

128μg/ml (11, 14), indicating brevicidine is a promising antimicrobial candidate. In

our previous study, we found that brevicidine dissipated the proton motive force of

bacteria. However, before the present investigations, little was known about how

brevicidine exerts its potent bactericidal effect against Gram-negative pathogens. In

addition, the antimicrobial activity against AMR pathogens and the therapeutic effect

in vivo of brevicidine still needs to be investigated before it is considered for a clinical

trial.

76
77 **Fig. 1. The structure of brevicidine and the predicted biosynthetic gene cluster** (14). **A**, The
78 non-ribosomal peptide synthetases genes harbored by the *Brevibacillus laterosporus* DSM 25
79 genome; **B**, the catalytic domains encoded by the gene cluster, and the substrates incorporated
80 by the respective modules. Domains: A, adenylation; T, thiolation; C, condensation; E,
81 epimerization; TE, thioesterase; **C**, the structure of brevicidine.

In this study, we aimed to identify the precise molecular mechanism of action by which brevicidine kills Gram-negative pathogens. Moreover, the anti-biofilm activity against *E. coli*, the antimicrobial activity against AMR *Enterobacteriaceae*, and the *in vivo* therapeutic effect of brevicidine were also investigated. The results showed that brevicidine exerts its potent bactericidal effect by interacting with lipopolysaccharide (LPS) in the outer membrane as well as targeting phosphatidylglycerol (PG) and cardiolipin (CL) in the inner membrane, and subsequently disrupting the proton motive force of Gram-negative pathogens. This leads to adenosine triphosphate (ATP) synthesis inhibition, nicotinamide adenine dinucleotide (NADH) and reactive oxygen species (ROS) accumulation, and bacteria death. In addition, brevicidine showed potent antimicrobial activity against AMR Gram-negative pathogens and good anti-biofilm activity against Gram-negative pathogen. Notably, brevicidine showed a good therapeutic effect in an *E. coli*-induced mouse peritonitis–sepsis model. These results will pave the way for further research on the clinical applications of brevicidine.

Results and Discussion

Brevicidine shows potent antimicrobial activity against antibiotic-resistant (AMR) *Enterobacteriaceae* pathogens

The antimicrobial activity of brevicidine against AMR *Enterobacteriaceae* pathogens was measured by a MIC assay. The results show that brevicidine has potent antimicrobial activity, with a MIC value of 0.5μM (0.8mg/L) to 2μM (3.0mg/L) (Table 1), against *Enterobacteriaceae* pathogens, including the tested colistin-resistant

strains and multidrug-resistant strains (16). These results indicate that brevicidine has

no cross-resistance with colistin, ampicilin, ceftriaxone, cefotaxime, aztreonam,

gentamicin, ofloxacin, amoxicillin, streptomycin, lincomycin, doxycycline, or

ciprofloxacin, which makes it a more attractive antimicrobial candidate for treating

AMR Gram-negative pathogen infections.

Table 1 MIC values of brevicidine against *Enterobacteriacea*e bacterial pathogens.

Microorganism	MIC [μM (mg/L)]			
	Brevicidine	**Colistin**	**Amikacin**	**Ampicilin**
Escherichia coli ATCC25922	0.5 (0.8)	2 (2.3)	2 (1.2)	8 (2.8)
Escherichia coli CVCC3749	1 (1.5)	2 (2.3)	2 (1.2)	> 256 (89)
Escherichia coli B2[a]	1 (1.5)	8 (9.2)	32 (19)	> 256 (89)
Escherichia coli 16QD [a]	1 (1.5)	8 (9.2)	8 (4.7)	> 256 (89)
Escherichia coli EPZC17-45[b]	1 (1.5)	2 (2.3)	4 (2.3)	> 256 (89)
Escherichia coli EPZC17-22[b]	1 (1.5)	1 (1.6)	2 (1.2)	> 256 (89)
Escherichia coli E21-5[b]	0.5 (0.8)	2 (2.3)	2 (1.2)	> 256 (89)
Escherichia coli E21-71[b]	0.5 (0.8)	4 (4.6)	4 (2.3)	> 256 (89)
Escherichia coli E21-75[b]	0.5 (0.8)	2 (2.3)	8 (4.7)	> 256 (89)
Escherichia coli E21-107[b]	0.5 (0.8)	1 (1.6)	4 (2.3)	> 256 (89)
Enterobacter cloacae LMG02783[c]	1 (1.5)	> 64 (74)	8 (4.7)	> 256 (89)
Klebsiella pneumoniae LMG20218	2 (3.0)	2 (2.3)	4 (2.3)	> 256 (89)
Staphylococcus aureus ATCC 43300[d]	32 (48.6)	32 (37)	4 (2.3)	16 (5.6)
Enterococcus faecium LMG16003[e]	> 32 (48.6)	> 32 (37)	256 (150)	2 (0.7)

[a]*Escherichia coli* carrying mcr-1, blaNDM-5, mdfA genes, etc., which is resistant against colistin,
carbapenems, rifampin, etc. [b] *Escherichia coli* (clinical isolation) carrying multidrug resistance
genes, *Escherichia coli* EPZC17-45, resistant to ampicilin, ceftriaxone, cefotaxime, aztreonam,
gentamicin, doxycycline, and ciprofloxacin; *Escherichia coli* EPZC17-22, resistant to ampicilin,
ceftriaxone, cefotaxime, aztreonam, gentamicin, doxycycline, and ciprofloxacin; *Escherichia coli*
E21-5, resistant to ampicilin, amoxicillin, ceftriaxone, gentamicin, cefotaxime, streptomycin,
doxycycline; *Escherichia coli* E21-71, resistant to ampicilin, amoxicillin, ceftriaxone, cefotaxime,
doxycycline, and ciprofloxacin, and ofloxacin; *Escherichia coli* E21-75, resistant to ampicilin,
amoxicillin, ceftriaxone, cefotaxime, doxycycline, ofloxacin, and ciprofloxacin; *Escherichia coli*
E21-107, resistant to ampicilin, amoxicillin, ceftriaxone, gentamicin, cefotaxime, streptomycin,
doxycycline, and lincomycin. [c]*Enterobacter cloacae* carrying mcr-9 gene, which is a potent
colistin-resistant gene. [d] *Staphylococcus aureus* resistant to oxacillin and methicillin. [e]
Enterococcus faecalis resistant to vancomycin.

Fig. 2. Brevicidine showed potent anti-biofilm activity against *E. coli* O101. A, Biofilm inhibition

of various antimicrobials (brevicidine, colistin, and amikacin) versus *E. coli* O101. Biofilm

inhibition was assessed by incubating the antimicrobials with bacteria in growth media for 24 h,

discarding the growth media and staining the adhered biomass with CV. The percentage of

biofilm inhibition is reported relative to untreated bacteria (defined as 100%) and sterility control

wells (defined as 0%). All data were presented as means ± Standard Deviation. **B**, Biofilm

eradication assay results for various antimicrobials (brevicidine, colistin, and amikacin) versus *E.*

coli O101. Biofilm biomass, quantified based on CV staining, is indicated in pink. Biofilm

metabolism, quantified based on TTC metabolism, is indicated in red. All data were presented as

means ± Standard Deviation.

Brevicidine shows potent anti-biofilm activity against *E. coli*

Bacterial biofilm formation could increase the resistance of pathogens to

conventional antimicrobials, which could decrease the therapeutic effect of

antimicrobials (17, 18). Therefore, the anti-biofilm activity of antimicrobials plays a

vital role in their therapeutic effect on biofilm-formation bacterial pathogen

infections. In this study, the anti-biofilm activity of brevicidine against *E. coli* was

evaluated according to the protocol described previously (19). Colistin (Col) was used

as an anti-biofilm antimicrobial control, while amikacin (Amk) was used as a control

antimicrobial without anti-biofilm activity. The results show that brevicidine has

potent anti-biofilm activity against *E. coli*, with the same 100% inhibition and

eradication concentration of 4μM (6.1 mg/L), which is better than the anti-biofilm

activity of colistin against *E. coli* (Fig. 2). These results demonstrate that brevicidine

shows high potential for the development of therapeutic antimicrobial.

Brevicidine disrupts the electron transport chain of Gram-negative

bacteria

In previous studies, brevicidine had shown bacterial lytic activity at a concentration

of 10×MIC (11, 14). However, it did not disrupt the membrane at a concentration of

2×MIC, evidenced by an atomic force microscopy assay. Therefore, the authors

suggested that a more in-depth mechanism study is needed to characterize the

targets as well as the membrane disruption effect of brevicidine (11). In our previous

study, we found that brevicidine, at a concentration of 1×MIC, could dissipate the

proton motive force of *E. coli* , evidence by a DiSC$_3$(5) assay (14). However, before our

investigations, little is known about the molecular mechanism of brevicidine on

gram-negative bacteria. In this study, to investigate the killing capacity of brevicidine

at concentrations of low MIC values, 4×MIC, 2×MIC, and 1×MIC, a time-killing assay

was performed as the previously described method (20, 21). Brevicidine quickly

decreased the population of *E. coli* at a concentration of 4×MIC, which killed all of

the tested bacteria in 1h. Interestingly, the population of *E. coli* did not decrease

under the treatment of brevicidine at concentrations of 2×MIC and 1×MIC within

30min (Fig. 3A). These results indicate that brevicidine has different modes of action

between high and low MIC concentrations; it shows bacterial lytic activity at

concentrations of 4-fold to 10-fold MIC, while it does not disrupt the membrane at

concentrations of 1-fold to 2-fold MIC (11, 14). The atomic force microscopy, which

was used in the previous study (11), is not a common technique for investigating the

membrane effect of antimicrobials because of its low resolution. Therefore, in this

study, we performed several fluorescent probe assays, scanning electron microscopy

(SEM) assay, and transmission electron microscopy (TEM) assay to get insight into the

effect of brevicidine, at low MIC level concentration, on the Gram-negative bacterial

membrane.

The effect of brevicidine on the membrane integrity of Gram-negative bacteria was

investigated by using the fluorescent probes N-Phenyl-1-naphthylamine (NPN) and

propidium iodide (PI). NPN is an outer-membrane barrier permeant fluorescent

probe (22). If the outer membrane is disrupted, the fluorescent probe can enter,

reaching the phospholipid layer and resulting in a significant increase of fluorescence.

The fluorescent probe PI is permeant to membrane intact bacteria, and the

disruption of membrane integrity can lead to a prominent increase in fluorescence

(20). There was no fluorescence increase observed for both brevicidine-treated

NPN-containing cell suspensions and brevicidine-treated PI-containing cell

suspensions (Fig. 3 B and C), indicating brevicidine does not disrupt the membrane

integrity at concentrations of 2×MIC, 1×MIC, and 0.5×MIC.

Fig. 3. Brevicidine kills Gram-negative bacteria via dissipating their electron transport chain. A,

time-killing curves of brevicidine at concentrations of 4×MIC, 2×MIC, and 1×MIC against *E. coli*

O101. **B**, *E. coli* O101 cells pretreated with propidium iodide were exposed to antimicrobial

peptides brevicidine at concentrations of 2×MIC, 1×MIC, and 0.5×MIC, and the extent of

membrane leakage was visualized as an increase in fluorescence. **C**, NPN fluorescence in *E. coli*

O101 cells upon exposure to brevicidine at concentrations of 2×MIC, 1×MIC, and 0.5×MIC.

Representative examples from three technical replicates are shown. **D**, ATP concentration of *E.*

coli O101 cells treated with brevicidine at concentrations of 2×MIC, 1×MIC, and 0.5×MIC. **E**, Total

ROS accumulation in *E. coli* O101 cells treated with brevicidine at concentrations of 2×MIC,

1×MIC, and 0.5×MIC. Representative examples from three technical replicates are shown. **F**,

Growth curves of *E. coli* O101 exposed to brevicidine at concentrations of 2×MIC, 1×MIC, and

0.5×MIC. Exogenous addition of NAC (6 μM) diminished the bactericidal activity of brevicidine.

Although brevicidine showed no effect on the bacterial membrane integrity, our

previous study had shown that brevicidine dissipated the proton motive force of

bacteria as some novel antimicrobials do (14, 23, 24). The proton motive force of

bacteria is essential for the generation of ATP, which is an essential bioactive

compound for live bacteria (25, 26). If the proton motive force of bacteria is

dissipated, the process of ATP biosynthesis will be inhibited or even terminated. To

confirm that brevicidine can dissipate the proton motive force of bacteria, the ATP

levels of brevicidne, at concentrations of 2×MIC, 1×MIC, and 0.5×MIC, treated

bacteria were measured by using the BacTiter-Glo Microbial Cell Viability Assay

(Promega) kit. The results showed that brevicidine decreased the ATP level of

bacteria as expected (Fig. 3D). These results confirmed that brevicidine could

dissipate the proton motive force of bacteria, which is one of the mechanisms by

which brevicidine acts as a bactericidal antimicrobial.

The dissipation of the proton motive force of bacteria can result in an increase in the

NADH levels of cells. Indeed, our previous study showed that the dehydrogenation of

NADH can be inhibited by the addition of brevicidine (14). Considering the

importance of the dehydrogenation of NADH to the electron transport chain of

bacteria, we hypothesized that brevicidine treatment could result in the

accumulation of ROS and thereafter kills bacteria. Therefore, the ROS levels of

bacteria treated with different concentrations of brevicidine were measured by using

a commercial kit (Beyotime, catalogue no. S0033S). The results show that brevicidine

increases the ROS level of bacteria (Fig. 3E), which is one of the reasons that

brevicidine shows bactericidal activity. This was further demonstrated by growth

curve assay, which showed that the bactericidal activity of brevicidine was

attenuated by the addition of the antioxidant N-Acetyl-L-cysteine (NAC, 6 mM) (Fig.

3F). Taking together, these results demonstrate that brevicidine exerts its bactericidal

activity via dissipating the proton motive force of bacteria and thereafter inhibits ATP

biosynthesis, inhibits the dehydrogenation of NADH, accumulates ROS in bacteria,

and results in bacteria death.

Fig. 4. The morphology analysis of *E. coli* O101 under the treatment of brevicidine. A,
Fluorescence microscopy images of *E. coli* O101 cells, which were challenged with brevicidine at
concentrations of 2× MIC, 1× MIC, and 0.5× MIC for 5min. Green denotes a cell with an intact
membrane, whereas red denotes a cell with a compromised membrane. **B**, SEM image of *E. coli*
O101 cells after treatment with brevicidine at a concentration of 1× MIC for 15min, 30min, 60min,
and 120min. **C**, TEM image of *E. coli* O101 cells after treatment with brevicidine at a
concentration of 1× MIC for 120min.

To investigate the effect of brevicidine on the cell membrane in intuitive ways, we

performed fluorescence microscopy, SEM, and TEM assays. The fluorescence

microscopy assay results are consistent with the PI and NPN assays, which

demonstrate that brevicidine is not an antimicrobial that kills pathogens by

disrupting bacteria membrane integrity (Fig. 4A). Interestingly, after treatment with

brevicidine (1×MIC) for 15 min, the bacterial surface fold was observed (Fig. 4B). In

addition, after treatment with brevicidine (1×MIC) for 120 min, most of the bacteria

were broken, evidenced by SEM and TEM images (Fig. 4 B and C). These results are

consistent with the results of time killing assay (Fig. 3A), which showed that

brevicidine started to kill bacteria after 30min treatments.

Brevicidine targets the membrane components LPS, PG, and CL

As brevicidine dissipates the proton motive force of bacteria, we hypothesized that

brevicidine could target the component of the bacterial inner membrane. In addition,

previous studies showed that brevicidine could interact with LPS (27, 28). To get an

insight into the potential target, an agar diffusion assay was performed to investigate

if there is an interaction between brevicidine and LPS (outer membrane component),

phosphatidylethanolamine (PE, outer and inner membrane component), PG (inner

membrane component), or CL (inner membrane component) (29, 30). The main

component of mammalian cellular membrane phosphatidylcholine (PC) was used as

a control. The results show that the addition of PC or PE did not attenuate the

antimicrobial activity of brevicidine (Fig. 5A). In contrast, the antimicrobial activity

was completely abolished by the addition of LPS, PG, or CL (Fig. 5A).

Fig. 5. Brevicidine targets the membrane components LPS, PG, and CL. A, Exogenous addition of LPS, PG, or CL abolishes the antibacterial activity of brevicidine against *E. coli* O101. Inhibition zones of the mixtures of 1 nmol of brevicidine and 4 mmol of PC, LPS, PE, PG and CL, against *E. coli* O101 for overnight incubation at 37 °C. Representative examples from three technical replicates are shown. **B**, Increased MICs of brevicidine against *E. coli* O101 in the presence of LPS, PG and CL, ranging from 0 µg/ml to 128 µg/ml, based on MIC assays. Representative examples from three technical replicates are shown. **C**, ITC analysis of the interaction between brevicidine and LPS. **D**, ITC analysis of the interaction between brevicidine and PG. **E**, ITC analysis of the interaction between brevicidine and CL. The K_D values of brevicidine to LPS, PG and CL were calculated using the Nano Analyze Software (Waters LLC).

Subsequently, to investigate the dose-dependent effect of LPS, PG, or CL on the

antimicrobial activity of brevicidine, a MIC assay was performed with different

concentrations of LPS, PG, CL, PC, or PE. The MIC value of brevicidine had no changes

with the addition of PC and PE, suggesting there is no interaction between

brevicidine and PE (or PC) (Fig. 5B). This could explain why brevicidine targets the

Gram-negative bacteria but has low cytotoxicity to mammalian cells (11, 14). LPS, PG,

and CL decreased the antimicrobial activity of brevicidine in a dose-dependent

manner, and the MIC value of brevicidine increased 8-fold, 16-fold, and 64-fold by

the addition of 128μg/ml of LPS, PG, and CL, respectively (Fig. 5B). These results

indicate that brevicidine could bind to the cell membrane components LPS, PG, and

CL.

To gain an insight into the interaction between brevicidine and LPS, PG, or CL, an ITC

measurement was performed. The results show that CL and PG showed comparable

binding affinity to brevicidine, with K_D values of 0.156 μM and 0.170 Mm (Fig. 5 C

and D), respectively. Notably, CL and PG have much stronger (approximately 10-fold

stronger) binding affinity to brevicidine than LPS, which has a K_D value of 1.785 μM

(Fig. 5E). These results demonstrate that brevicidine has multiple targets on the

bacteria, which is good for inhibiting the development of bacterial antibiotic

resistance. Together, the observed results demonstrate that brevicidine exerts its

bactericidal activity via interacting with LPS in the outer membrane and targeting CL

and PG in the inner membrane, and thereafter dissipating the proton motive force of

bacteria.

Brevicidine inhibits the oxidative phosphorylation and protein synthesis processes of *E. coli*

To gain an insight into the antimicrobial mechanism of brevicidine, transcriptome analysis was performed on *E. coli* treated with or without brevicidine at a concentration of 0.5µM (0.8 mg/L). The whole distribution of differentially expressed genes (DEGs) is shown in Fig. 6A. When padj value<0.05 and log2(FoldChange)>0, among 4622 genes, 587 were upregulated and 622 were downregulated, and 3413 genes were not significantly changed (Fig. 6A). Subsequently, Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment analysis was performed on the genes that had significantly expression changes. The results showed that ribosome, oxidative phosphorylation, and aminoacyl-tRNA biosynthesis pathways were significantly (padj value<0.05) affected by brevicidine treatment (Fig. 6B). After *E. coli* exposure to brevicidine for 1h oxidative phosphorylation related genes including ATP synthase, NADH-quinone oxidoreductase, and cytochrome oxidase were significantly downregulated (Fig. 6C). This change would contribute to the lower level of ATP, higher level of NADH and ROS of the brevicidine treated cells. The downregulated tRNA ligase and ribosome synthesis metabolism indicate that the protein synthesis process was inhibited by brevicidine treatment.

Fig. 6. Transcriptome analysis of *E. coli* O101 under the treatment of brevicidine [0.5 μM (0.8 mg/L)] for 1 h. A, The volcano map of DEGs. The abscissa represents the multiple changes of gene expression between brevicidine treated and untreated *E. coli* O101; the ordinate represents the statistical significance of the change of gene expression (pvalue<0.05 |log2FoldChange|>0). **B**, Scatter plot of KEGG pathway enrichment analysis of DEGs (Brevicidine vs untreated). The vertical axis represents the KEGG enrichment analysis pathway name, and the horizontal axis represents the gene ratio of the genes in the pathway. The size of the dots indicates the number of DEGs in the pathway, and the color of the dots corresponds to different padj value ranges. **C**, Genes involved in oxidative phosphorylation, ribosome, and aminoacyl-tRNA biosynthesis were identified as significantly different between the treatment of brevicidine and untreated, with pvalue values <0.05 and log2FoldChange <0 in expression level. All genes that had significantly changes were down-regulated in oxidative phosphorylation, ribosome, and aminoacyl-tRNA biosynthesis pathways. All data are presented as means (n= 3 biological replicates).

Fig. 7. Scheme of the proposed mechanism of action brevicidine against *E. coli*. Brevicidine exerts its bactericidal activity via interacting with LPS in the outer membrane (OM) and targeting CL and PG in the inner membrane (IM), which could make a positive potential surround the outside surface of inner membrane (Note: no evidence shows brevicidine directly interacts with these channel proteins), and thereafter dissipating the proton motive force of the bacteria. This results in metabolic perturbation, including decreases in the intracellular ATP level, inhibits the dehydrogenation of NADH and accumulates ROS in bacteria. In addition, transcriptome analysis results showed that brevicidine inhibits the synthesis of t-RNA ligase (essential for amino acid transport) and ribosome protein synthesis, which can result in protein synthesis inhibition. All these actions together lead to the death of Gram-negative pathogens.

Collectively, our findings demonstrate that brevicidine exerts its bactericidal activity via interacting with LPS in the outer membrane and targeting CL and PG in the inner membrane, and thereafter dissipation of the proton motive force of bacteria. This can result in metabolic perturbation, including inhibits ATP biosynthesis, inhibits the dehydrogenation of NADH, accumulates ROS in bacteria, and inhibits protein synthesis (Fig. 7).

Fig. 8. Brevicidine showed a good therapeutic effect in a mouse peritonitis–sepsis model. A,
Scheme of the experimental protocol for the mouse peritonitis–sepsis model. **B**, Survival rates of
mice in the mouse peritonitis–sepsis model (n= 10). Increased survival rates of mice over 7 d by a
dose that leads to 80% of death of *E. coli* O101 (2.0×10^9 c.f.u.), treated with brevicidine
(10 mg per kg, 20 mg per kg, or 40 mg per kg) or with colistin (20 mg per kg). **C to G**, Brevicidine
significantly reduced the bacterial load of organs of the peritonitis-sepsis mouse. At 24h
post-infection, the survived mice (n = 6) were euthanized by cervical dislocation. Bacterial loads
(Log10 CFU per gram of *E. coli* O101) of the heart (**C**), liver (**D**), spleen (**E**), lung (**F**) and kidney (**G**)
were counted. All data were presented as means ± Standard Deviation (n = 6). Correlation
analyses were evaluated by Pearson r2, ns: ***p < 0.01 and ****p < 0.001 vs. untreated group.

354 **Brevicidine showed a good therapeutic effect in a mouse**

355 **peritonitis–sepsis model**

356 Given the attractive mode of action and potent antimicrobial and anti-biofilm activity

357 of brevicidine, we investigated its potential as a therapeutic in a mouse

358 peritonitis–sepsis model (Fig. 8A). Mice were infected intraperitoneally with *E. coli* at

359 a dose that leads to 80% of death. At 1h post-infection, brevicidine was introduced at

360 single intravenous doses ranging from 10mg/kg to 40mg/kg. All brevicidine-treated

361 mice were survived (Fig. 8B). Consistently, the bacterial load in different organs of

362 mice significantly reduced under brevicidine treatment in a dose-dependent manner

363 at either 1 or 7 days post-infection (Fig. 8 C-G, and Supplementary Fig. S1). These

364 results are consistent with the previous study, which showed that brevicidine had

365 good antimicrobial activity in a mouse thigh model (11). Together, these findings

366 demonstrate the potential of brevicidine as a novel therapeutic antimicrobial in AMR

367 Gram-negative pathogen infections, in particular, AMR *Enterobacteriaceae* pathogen

368 infections.

369 **Conclusions**

370 AMR poses a major threat to human health around the world; many bacterial

371 infections have become increasingly difficult, or even impossible, to treat with

372 conventional antimicrobials. Therefore, new first-in-class antibiotics, operating via

373 novel modes of action, are desperately-needed. Here, we show that brevicidine has

374 potent antimicrobial activity against AMR *Enterobacteriaceae* pathogens, with a MIC

375 value range of 0.5μM (0.8mg/L) to 2μM (3.0mg/L). In addition, brevicidine showed

potent anti-biofilm activity against the *Enterobacteriaceae* pathogens, with the same 100% inhibition and eradication concentration of 4μM (6.1mg/L). Mechanistic studies showed that brevicidine exerts its potent bactericidal activity via interacting with LPS in the outer membrane, targeting PG and CL in the inner membrane, and dissipating the proton motive force of bacteria. This results in metabolic perturbation, including inhibition of ATP biosynthesis, inhibition of the dehydrogenation of NADH, accumulation of ROS in bacteria, and inhibition of protein synthesis (Fig. 7). Lastly, brevicidine showed a good therapeutic effect in a mouse peritonitis–sepsis model. Our findings pave the way for further research on clinical applications of brevicidine, to combat the prevalent infections caused by AMR Gram-negative pathogens worldwide.

Materials and Methods

Ethical approval

The animal experiment protocol was approved by the Animal Ethical and welfare Committee of Sichuan Agricultural University [permission number 20220087].

Bacterial strains used and growth conditions

Bacterial strains used in this study are listed in S1 Table. All bacterial strains were inoculated in LB and incubated at 37°C with aeration at 220 rpm for preparing the overnight cultures.

Purification of brevicidine

Methods for the purification of brevicidine have been described in detail previously (12). The matrix-assisted laser desorption ionization-time-of-flight mass spectrometry

data and high-performance liquid chromatograph trace of purified brevicidine are shown in Supplementary Figs. S2 and S3, which are showing the correct molecular weight and high purity of the purified brevicidine.

Minimum inhibitory concentration (MIC) assay

MIC values of antimicrobials against the tested bacterial pathogens were determined according to the standard guidelines (31). Briefly, cell concentration was adjusted to approximately 5×10^5 cells per ml in cation-adjusted Mueller-Hinton broth (MHB). Subsequently, antimicrobials were added to the above bacterial cultures at different concentrations. After 24h of incubation at 37°C, the MIC value was defined as the lowest concentration of antimicrobial with no visible growth.

For measuring the MIC values of antimicrobials against indicator strains in the presence of cell membrane components, MHB was replaced with MHB containing different concentrations of cell membrane components, and the effect of these two conditions was compared.

Biofilm inhibition assay

Biofilm inhibition assay was performed according to a previously described protocol (19). Briefly, an overnight culture of *E. coli* O101 was diluted 100-fold in tryptone soy broth (TSB) and incubated at 37 °C with aeration at 220 r.p.m. Bacteria were grown to an OD_{600} of 0.8, and then the concentration of cells was adjusted to an OD_{600} of 0.04 in TSB. The cell suspension was added to a 96-well plate, and antimicrobials were added at different final concentrations. After incubation at 37 °C for 24 h, the wells were washed with MiliQ water three times, and the adhered biomasses were stained with 105 μl of a 0.1% crystal violet (CV) (wt/vol) solution. The plate was placed on a

low-speed orbital shaker and allowed to stain at room temperature for 30 min.

Subsequently, the plate was washed with MiliQ water three times, and then 70%

(vol/vol) ethanol was added to the wells (110 μl per well) and incubated for 30 min at

room temperature with gentle shaking. The absorbance was measured by using a

Thermo Scientific Varioskan™ LUX multimode microplate reader with a wavelength of

595 nm. The inhibition rates were calculated by the raw CV absorbance values from

each growth condition vs the raw CV absorbance value of normally grown biofilm.

Representative examples from three technical replicates are shown.

Biofilm eradication assay

Biofilm eradication assay was performed according to a previously described protocol

(19). Briefly, an overnight culture of *E. coli* O101 was diluted 100-fold in tryptone soy

broth (TSB) and incubated at 37 °C with aeration at 220 r.p.m. Bacteria were grown to

an OD_{600} of 0.8, and then the concentration of cells was adjusted to an OD_{600} of 0.04

in TSB. The cell suspension was added to 96-well plates and incubated at 37 °C for 24

h. After washing the wells with TSB three times, antimicrobials were added to the

wells at different concentrations, and the plates (one for CV staining, the other one

for TTC staining) were incubated at 37 °C for 24 h. For measuring the metabolic

activity of biofilm, a final concentration of 0.05% TTC was added to wells with

antimicrobials. Subsequently, the adhered biomasses were measured as the method

described in the biofilm inhibition assay. After washing all the wells of the TTC plate

three times with distilled water, dimethyl sulfoxide (DMSO) was added to the wells at

205 μl per well. After that, the plate was placed on an orbital shaker rotating at low

speed and mixed at room temperature for 30 min. The absorbance of the TTC plate

was measured by using a Thermo Scientific Varioskan™ LUX multimode microplate

reader with a wavelength of 500 nm. The biofilm eradication (Biomass and

metabolism) rate was calculated by the method described in the previous study (19).

Representative examples from three technical replicates are shown.

Time-dependent killing assay

The time-dependent killing assay was performed according to the procedure

described previously (20, 21). An overnight culture of *E. coli* O101 was diluted

100-fold in MHB and incubated at 37 °C with aeration at 220 r.p.m. Bacteria were

grown to an OD of 0.6, and then the concentration of cells was adjusted to 7.6×10^7

cells per mL. Bacteria were then challenged with antimicrobials at concentrations of

1×MIC, 2×MIC, or 4×MIC in glass culture tubes at 37 °C with aeration at 220 r.p.m.

Bacteria not treated with antimicrobials were used as normal growth control. At

desired time points, two hundred μl aliquots were taken, centrifuged at 5,000 g for 5

min, and resuspended in 200 μl of MHB. Ten-fold serially diluted samples were plated

on MHA plates. After incubation at 37 °C for 24 h, colonies were counted, and the

colony forming units (c.f.u.) per ml were calculated. Each experiment was performed

in triplicate.

Membrane integrity assay

A fresh culture of *E. coli* O101 was pelleted at 5,000 g for 5 min and washed three

times with MHB. The cell density was normalized to an OD_{600} of 0.2, loaded with

propidium iodide (final concentration 2.5 μg per ml), and incubated for 5 min in the

dark for probe fluorescence to stabilize. After the cell suspension was added to a

96-well microplate, brevicidine was added to final concentrations of 2×MIC, 1×MIC,

or 0.5×MIC, with the antimicrobials added after ~30 s, and fluorescence was

monitored for 15 min. The excitation and emission wavelengths on the fluorescence

spectrometer were adjusted to 533 nm and 617 nm, respectively. Representative

examples from three technical replicates are shown.

Outer membrane permeability assay

The integrity of the outer membrane was investigated with the fluorescent probe

N-Phenyl-1-naphthylamine (NPN, Aladdin). A fresh culture of *E. coli* O101 was

pelleted at 5,000 g for 5 min and washed three times with 10 mM HEPES containing

10 mM glucose (GHEPES, pH 7.2). The cell density was normalized to an OD_{600} of 0.2

in GHEPES, loaded with NPN (final concentration 30 µM), and incubated for 30 min in

the dark for probe fluorescence to stabilize. After the cell suspension (190 µL) was

added to a 96-well microplate, brevicidine (10 µL) was added to final concentrations

of 2×MIC, 1×MIC, or 0.5×MIC, with the antimicrobials added after ~30 s, and

fluorescence was monitored for 15 min. The excitation and emission wavelengths on

the fluorescence spectrometer were adjusted to 350 nm and 420 nm, respectively.

Representative examples from three technical replicates are shown.

Fluorescence microscopy assay

This assay was performed according to the procedure described previously (32, 33). *E.*

coli O101 was grown to an OD_{600} of 0.6 in MHB. The culture was pelleted at 5,000 g

for 5 min and washed with MHB three times. After normalization of the cell density

to an OD_{600} of 0.2 in MHB, brevicidine was added to a final concentration of 2×MIC,

1×MIC or 0.5×MIC. After incubation at 37 °C for 5min, cells were collected by

centrifugation. Subsequently, SYTO® 9 and propidium iodide (LIVE/DEAD Baclight

Bacterial Viability Kit, Invitrogen) were added to the above cells. After incubation at

room temperature for 15 min, cells were washed three times with MHB. Then the

cell suspensions were loaded on 1.5 % agarose pads and analyzed by Nikon 80i

(Japan) microscope.

ATP determination

The ATP levels were determined using a BacTiter-Glo Microbial Cell Viability Assay kit

(Promega). A fresh culture of *E. coli* O101 was pelleted at 5,000 g for 5 min and

washed three times with MHB. The cell density was normalized to an OD_{600} of 0.2,

loaded with brevicidine at final concentrations of 2×MIC, 1×MIC, or 0.5×MIC. At

desired time points, 100 µl of cell culture were taken out, mixed with 100 µl

BacTiter-Glo reagent, and incubated for 5 minutes at room temperature.

Luminescence was measured with an Infinite M2000pro Microplate reader (Tecan).

CCCP (40 µg/mL) was used as a positive control. The relative ATP levels were

calculated using the measured Luminescence values vs the Luminescence value of

untreated cells at relative time points.

Reactive oxygen species (ROS) measurement

The levels of ROS in *E. coli* O101 treated with different concentrations of brevicidine

were measured by 10 µM of 2',7'-dichlorofluorescein diacetate (DCFH-DA), following

the manufacturer's instruction (Beyotime, catalogue no. S0033S). Briefly, a fresh

culture of *E. coli* O101 was pelleted at 5,000 g for 5 min and washed three times with

MHB. The cell density was normalized to an OD_{600} of 1.0, loaded with DCFH-DA at a

final concentration of 10 µM, and the mixture incubated at 37 °C for 30 min. After

washing with MHB three times, 190 µl of probe-labeled bacterial cells were added to

a 96-well plate, and then 10 µl of brevicidine. Fluorescence was recorded by using a

Thermo Scientific Varioskan™ LUX multimode microplate reader with the excitation

wavelength at 488 nm and the emission wavelength at 525 nm. The antioxidant

N-Acetyl-L-cysteine (NAC, 6 mM) was used as a control to neutralize the production

of ROS.

Scanning electron microscopy (SEM) and transmission electron microscopy (TEM) assays

520   The morphology of 1μM (1.52 mg/L) brevicidne treated *E. coli* O101 was recorded in vacuum on a ZEISS Gemini SEM300 with an acceleration voltage of 3kV. For TEM measurements, samples were prepared by delivering 2μl of 1μM (1.52 mg/L) brevicidine treated *E. coli* O101 suspension to carbon-coated copper grids and dried in a vacuum system. The morphology was obtained under a JEM-2100Plus Electron Microscope using an acceleration voltage of 120kV.

Spot-on-lawn assay

527   An overnight cultured *E. coli* O101 was added to 0.7% MHA (w/v, temperature 42 °C) at a final concentration of 0.25% (v/v), and then the mixture was poured onto the plates, with 20 mL for each 15 cm diameter circular plate. Subsequently, a spot-on-lawn assay was used to analyze the antimicrobial activity of brevicidine in the presence of cell membrane components (20). In short, 20 μL brevicidine (50 μM) containing 200 μM lipopolysaccharides (LPS, from Escherichia coli 055:B5, Solarbio, catalogue no. L8880, ≥98%), L-α-phosphatidylcholine, (PC, Aladdin, catalogue no. L130331, ≥99%), L-α-phosphatidylglycerol (PG, Aladdin, catalogue no. L130372; ≥99%), L-α-phosphatidylethanolamine (PE, Aladdin, catalogue no. L130310; ≥99%) or Cardiolipin (CL, Aladdin, catalogue no. B130227;≥99%) was loaded to the agar plate that contains indicator strain. After the brevicidiene/cell membrane component mixture solution drops had dried, the plates were transferred to a 37 °C incubator for overnight incubation.

Isothermal Titration Calorimetry (ITC) assay

ITC was performed with the Low Volume NanoITC (TA Instruments) at 25°C to determine the interaction between brevicidine and LPS, PG, or CL. Brevicidine was diluted in a buffer (20 mM HEPES, pH7.0) to a final concentration of 50 µM. Samples were degassed before use. The chamber was filled with 300 µL of the brevicidine solution, and LPS (250µM), PG (500µM), or CL (500µM) was titrated into the chamber at a rate of 1.96 µL/200 s with a stirring rate of 300 rpm. The control experiment was performed with 20 mM HEPES titration to 50 µM brevicidine. The K_D values of brevicidine to LPS, PG, or CL were calculated using the Nano Analyse Software (Waters LLC).

Transcriptome analysis of *E. coli* O101 treated with brevicidine

Three replicates of overnight bacteria cultures were diluted at 1:100 in MHB and grown at 37°C with aeration at 220rpm. Bacteria were grown to an OD_{600} of 0.4, and then the concentration of cells was adjusted to an OD_{600} of 0.2 in MHB. Subsequently, the bacteria were treated with brevicidine at a final concentration of 0.5µM (0.8 mg/L). Bacteria without brevicidine treatment were used as untreated control. Three replicates were performed for both brevicidine treated and untreated bacterial cultures, 4ml for each sample. After incubation at 37°C with aeration at 220rpm for 1h, the bacteria from different treatments were collected by centrifuging, and the bacterial pallets were stored in liquid nitrogen immediately. Subsequently, the samples were sent to Novogene Co., Ltd (China) for transcriptome analysis. After RNA extraction, mRNA purification, and cDNA synthesis, the samples were sequenced on the Illumina NovaSeq6000. The quality of the resulting fastq reads was mapped on the reference genome using Bowtie2 2.3.4.3 using default settings. Feature Counts

1.5.0-p3 was used to get the gene counts. DESeq2 1.20 was used to identify all genes,

and padj ≤0.05 and |log2 (foldchange) | ≥ 0 of genes between two treatments were

regarded as differentially expressed genes (DEGs). The volcano map and KEGG

enrichment of DEGs were performed with Magic Novogene package

(https://magic.novogene.com). The RNA-Seq data have been deposited in the NCBI

Gene Expression Omnibus with the accession number PRJNA872022.

Mouse peritonitis–sepsis model

Brevicidine was tested against *E. coli* O101 in a mouse peritonitis–sepsis model to

assess its *in vivo* bioavailability. BALB/c male mice (n=6 per group) were infected with

0.4ml of bacterial suspension ($2×10^9$ c.f.u. per mouse) via intraperitoneal injection, a

concentration that achieves approximately 80% mortality within 48 h post-infection.

At 1h post-infection, mice were treated with 0.9% NaCl, brevicidine (40mg/kg),

brevicidine (20mg/kg), brevicidine (10mg/kg), or colistin (20mg/kg) via intravenous

injection. Mice without *E. coli* O101 infections were used as the normal group control.

Once the infected mice died, different organs, including the heart, liver, spleen, lung,

and kidney, were collected and homogenized in sterilized PBS for bacterial load

quantification. At 24h post-infection, the organs of the survived mice were collected

to measure the bacterial load. In addition, the same mouse peritonitis–sepsis model

was determined for 7d (n=10), and the bacterial load in the organs of surviving mice

was quantified at 7 days post-infection.

Statistical analysis

GraphPad Prism 8.0 was used to fit the data of Fig. 2, Fig. 3, Fig. 5, Fig. 8, and

Supplementary Fig. S1. The statistical significance of the data was assessed using a

two-tailed Student's t-test with GraphPad Prism 8.0. Correlation analyses were

evaluated by Pearson r2, ns: p>0.05, *p<0.05, **p<0.01, ***p<0.001, and ****p<0.0001.

Data Availability Statement: The transcriptome (RNA sequencing) data that support the findings of this study have been deposited in the NCBI Sequence Read Archive (SRA) with the accession code PRJNA872022.

Finding: Xinghong Zhao was supported by the Science and Technology Project of Sichuan Province (2022YFH0057). Hongping Wan was supported by the Science and Technology Project of Sichuan Province (2022YFH0062).

Competing Interests: The authors have declared that no competing interests exist.

Acknowledgments

We thank Prof. Kui Zhu (National Center for Veterinary Drug Safety Evaluation, College of Veterinary Medicine, China Agricultural University, Beijing, China.) for providing the multidrug resistant *Escherichia coli* (B2 and 16QD) strains.

References

1. O'neill, J. I. M. (2014) Antimicrobial resistance: tackling a crisis for the health and wealth of nations. *Rev. Antimicrob. Resist.* **1**, 1–16

2. Kwon, J. H., and Powderly, W. G. (2021) The post-antibiotic era is here. *Science.* **373**, 471

3. Knight, G. M., Glover, R. E., McQuaid, C. F., Olaru, I. D., Gallandat, K., Leclerc, Q. J., Fuller, N. M., Willcocks, S. J., Hasan, R., and van Kleef, E. (2021) Antimicrobial resistance and COVID-19: Intersections and implications. *Elife.* **10**, e64139

4. Strathdee, S. A., Davies, S. C., and Marcelin, J. R. (2020) Confronting antimicrobial resistance beyond the COVID-19 pandemic and the 2020 US election. *Lancet.* **396**, 1050–1053

5. Lai, C.-C., Chen, S.-Y., Ko, W.-C., and Hsueh, P.-R. (2021) Increased antimicrobial resistance during the COVID-19 pandemic. *Int. J. Antimicrob. Agents.* **57**, 106324

6. Rizvi, S. G., and Ahammad, S. Z. (2022) COVID-19 and antimicrobial resistance: A cross-study. *Sci. Total Environ.* **807**, 150873

7. Butler, M. S., Blaskovich, M. A. T., and Cooper, M. A. (2017) Antibiotics in the clinical pipeline at the end of 2015. *J. Antibiot. (Tokyo).* **70**, 3

8. Butler, M. S., Blaskovich, M. A., and Cooper, M. A. (2013) Antibiotics in the clinical pipeline in 2013. *J. Antibiot. (Tokyo).* **66**, 571

9. Roemer, T., and Boone, C. (2013) Systems-level antimicrobial drug and drug synergy discovery. *Nat. Chem. Biol.* **9**, 222

10. Süssmuth, R. D., and Mainz, A. (2017) Nonribosomal peptide synthesis—principles and prospects. *Angew. Chemie Int. Ed.* **56**, 3770–3821

11. Li, Y.-X., Zhong, Z., Zhang, W.-P., and Qian, P.-Y. (2018) Discovery of cationic nonribosomal peptides as Gram-negative antibiotics through global genome mining. *Nat. Commun.* **9**, 3273

12. Zhao, X., Li, Z., and Kuipers, O. P. (2020) Mimicry of a Non-ribosomally Produced Antimicrobial, Brevicidine, by Ribosomal Synthesis and Post-translational Modification. *Cell Chem. Biol.* **27**, 1262-1271.e4

13. Al Ayed, K., Ballantine, R. D., Hoekstra, M., Bann, S. J., Wesseling, C. M. J., Bakker, A. T., Zhong, Z., Li, Y.-X., Brüchle, N. C., and van der Stelt, M. (2022) Synthetic studies with the brevicidine and laterocidine lipopeptide antibiotics including analogues with enhanced properties and in vivo efficacy. *Chem. Sci.* **13**, 3563–3570

14. Zhao, X., and Kuipers, O. P. (2021) BrevicidineB, a new member of the brevicidine family, displays an extended target specificity. *Front. Microbiol.* **12**, 1482

15. Tacconelli, E., Magrini, N., Kahlmeter, G., and Singh, N. (2017) Global priority list of antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics. *World Heal. Organ.* **27**, 318–327

16. Song, M., Liu, Y., Huang, X., Ding, S., Wang, Y., Shen, J., and Zhu, K. (2020) A

broad-spectrum antibiotic adjuvant reverses multidrug-resistant Gram-negative pathogens. *Nat. Microbiol.* **5**, 1040–1050

17. Liu, Y., Shi, L., Su, L., van der Mei, H. C., Jutte, P. C., Ren, Y., and Busscher, H. J. (2019) Nanotechnology-based antimicrobials and delivery systems for biofilm-infection control. *Chem. Soc. Rev.* **48**, 428–446

18. Sharma, G., Sharma, S., Sharma, P., Chandola, D., Dang, S., Gupta, S., and Gabrani, R. (2016) Escherichia coli biofilm: development and therapeutic strategies. *J. Appl. Microbiol.* **121**, 309–319

19. Haney, E. F., Trimble, M. J., and Hancock, R. E. W. (2021) Microtiter plate assays to assess antibiofilm activity against bacteria. *Nat. Protoc.* **16**, 2615–2632

20. Zhao, X., Yin, Z., Breukink, E., Moll, G. N., and Kuipers, O. P. (2020) An Engineered Double Lipid II Binding Motifs-Containing Activity against Enterococcus faecium. *Antimicrob. Agents Chemother.* **64**, 1–12

21. Ling, L. L., Schneider, T., Peoples, A. J., Spoering, A. L., Engels, I., Conlon, B. P., Mueller, A., Schäberle, T. F., Hughes, D. E., and Epstein, S. (2015) A new antibiotic kills pathogens without detectable resistance. *Nature*. **517**, 455

22. Helander, I. M., and Mattila-Sandholm, T. (2000) Fluorometric assessment of Gram-negative bacterial permeabilization. *J. Appl. Microbiol.* **88**, 213–219

23. Cochrane, S. A., Findlay, B., Bakhtiary, A., Acedo, J. Z., Rodriguez-Lopez, E. M., Mercier, P., and Vederas, J. C. (2016) Antimicrobial lipopeptide tridecaptin A1 selectively binds to Gram-negative lipid II. *Proc. Natl. Acad. Sci.* **113**, 11561–11566

24. Stokes, J. M., Yang, K., Swanson, K., Jin, W., Cubillos-Ruiz, A., Donghia, N. M., MacNair, C. R., French, S., Carfrae, L. A., and Bloom-Ackerman, Z. (2020) A deep learning approach to antibiotic discovery. *Cell*. **180**, 688–702

25. Ahmed, S., and Booth, I. R. (1983) The use of valinomycin, nigericin and trichlorocarbanilide in control of the protonmotive force in Escherichia coli cells. *Biochem. J.* **212**, 105–112

26. Bakker, E. P., and Mangerich, W. E. (1981) Interconversion of components of the bacterial proton motive force by electrogenic potassium transport. *J. Bacteriol.* **147**, 820–826

27. Malina, A., and Shai, Y. (2005) Conjugation of fatty acids with different lengths modulates the antibacterial and antifungal activity of a cationic biologically inactive peptide. *Biochem. J.* **390**, 695–702

28. Li, Y. X., Zhong, Z., Zhang, W. P., and Qian, P. Y. (2018) Discovery of cationic nonribosomal peptides as Gram-negative antibiotics through global genome mining. *Nat. Commun.* **9**, 2–10

29. Sohlenkamp, C., and Geiger, O. (2016) Bacterial membrane lipids: diversity in structures and pathways. *FEMS Microbiol. Rev.* **40**, 133–159

30. Santos, R. S., Figueiredo, C., Azevedo, N. F., Braeckmans, K., and De Smedt, S. C. (2018) Nanomaterials and molecular transporters to overcome the bacterial envelope barrier: Towards advanced delivery of antibiotics. *Adv. Drug Deliv. Rev.* **136**, 28–48

31.   Wiegand, I., Hilpert, K., and Hancock, R. E. W. (2008) Agar and broth dilution
methods to determine the minimal inhibitory concentration (MIC) of
antimicrobial substances. *Nat. Protoc.* **3**, 163

32.   Li, Z., Chakraborty, P., de Vries, R. H., Song, C., Zhao, X., Roelfes, G., Scheffers,
D., and Kuipers, O. P. (2020) Characterization of two relacidines belonging to a
novel class of circular lipopeptides that act against Gram-negative bacterial
pathogens. *Environ. Microbiol.* **22**, 5125–5136

33.   Li, Z., de Vries, R. H., Chakraborty, P., Song, C., Zhao, X., Scheffers, D.-J.,
Roelfes, G., and Kuipers, O. P. (2020) Novel modifications of nonribosomal
peptides from Brevibacillus laterosporus MG64 and investigation of their
mode of action. *Appl. Environ. Microbiol.*